MULTIVALUED SECTIONS AND SELF-MAPS OF SPHERE BUNDLES

M. C. CRABB

ABSTRACT. Let G be a finite group and V a finite dimensional (non-zero) orthogonal G-module such that, for each prime p dividing the order of G, the subspace of V fixed by a Sylow p-subgroup of G is non-zero and, if the dimension of V is odd, has dimension greater than 1. Using ideas of Avvakumov, Karasev, Kudrya and Skopenkov and work of Noakes on self-maps of sphere bundles, we show that, for any principal G-bundle $P \to X$ over a compact ENR X, there exists a G-map from P to the unit sphere S(V) in V.

1. INTRODUCTION

Let G be a (non-trivial) finite group and let V be an orthogonal G-module of dimension $n \ge 1$. Given a principal G-bundle $P \to X$ over a compact ENR X we form the n-dimensional real vector bundle $\xi = P \times_G V$ and its associated sphere bundle $S(\xi) = P \times_G S(V)$ (where S(V) is the unit sphere in V).

Theorem 1.1. Suppose that either

(a) for each prime p dividing the order of G the fixed subspace V^H of a Sylow p-subgroup H of G is non-zero and, if n is odd, has dimension greater than 1, or (b) n is odd, the fixed subspace V^H of a Sylow 2-subgroup H is non-zero and dim X < 2(n-1).

Then there exists a G-equivariant map

$$P \to S(V),$$

or, equivalently, a section of the sphere bundle $S(\xi)$.

For a fixed integer q > 1, the symmetric group \mathfrak{S}_q , the group of permutations of the set $I = \{1, \ldots, q\}$, acts on the Euclidean vector space $\mathbb{R}[I]$ with orthonormal basis $e_i, i \in I$. Let L denote the (q-1)-dimensional quotient of $\mathbb{R}[I]$ by the fixed subspace generated by $e_I = e_1 + \ldots + e_q$. We give L the inner product for which the orthogonal complement of $\mathbb{R}e_I$ projects isometrically onto L. Given a prime pwe can write $q = \sum_r a_r p^r$, where $0 \leq a_r < p$, and set $\alpha_p(q) = \sum_r a_r$. Then, for a Sylow p-subgroup H of \mathfrak{S}_q , the dimension of the fixed subspace L^H is equal to $\alpha_p(q) - 1$. (See the discussion in Section 3.)

Thus we deduce from Theorem 1.1(a) the following recent result of Avvakumov, Karasev, Kudrya and Skopenkov [1, 2], which was applied in [3] to the topological Tverberg theorem.

Date: August 2022, revised May2024.

²⁰²⁰ Mathematics Subject Classification. Primary 55M25, 55M35, 55R25, 55P69, 55S40, 57S17; Secondary 55M20, 55R70.

Key words and phrases. Sylow p-subgroup, sphere bundle, multivalued section, Euler class, Tverberg theorem.

Corollary 1.2. ([1, 2, 3].) Let $P \to X$ be a principal \mathfrak{S}_q -bundle over a compact ENR X and let λ denote the real vector bundle $P \times_{\mathfrak{S}_q} L \to X$ of dimension q-1 over X. Suppose that q is not a prime power and that $k \ge 1$ is a positive integer. If k = 1 and q is even, suppose further that $\alpha_p(q) > 2$ for every prime p.

Then there exists an \mathfrak{S}_{q} -equivariant map

$$P \to S(\mathbb{R}^k \otimes L)$$

to the unit sphere in $\mathbb{R}^k \otimes L$, or, equivalently, a section of the sphere bundle $S(\mathbb{R}^k \otimes \lambda)$ of $\mathbb{R}^k \otimes \lambda$.

The traditional approach to such problems, following Sullivan [16], is to localize at a prime p. If $H \leq G$ is a Sylow p-subgroup (trivial if p does not divide the order of G) and V^H is non-zero, then there is an H-equivariant map $P \to S(V)$. Equivalently, the pullback $S(\pi^*\xi)$ to the finite cover $\pi : P/H \to X = P/G$ has a section. Since the order of this cover, the index of H in G, is prime to p, it follows by a standard argument that the p-local obstruction to the existence of a section of $S(\xi)$ is zero in the metastable range: dim X < 2(n-1). If this holds for all primes p, the global obstruction is zero and the desired section, or equivariant map, exists in this range of dimensions. (An expository proof along these lines for the case of the symmetric group \mathfrak{S}_q was included in [8, Proposition 6.5].) Integral cohomology is enough to establish the result by the same reasoning in the range dim $X \leq n$, as in [5, Theorem 5.1] and [4, Theorem 1.2].

The new proof in [1, 2, 3] without the dimensional restriction is surprisingly elementary, especially when linked to a result of Noakes [15] on the degree of selfmaps of sphere bundles. This connection is explained in Section 2, where the main construction in [3] is formulated in the language of multivalued sections of sphere bundles and part (a) of Theorem 1.1 is established. In Section 3 we discuss the extension (no longer a corollary) of Corollary 1.2 that was established in [2] to the case in which k = 1 and q is even but not twice a power of an odd prime and give examples to show that this restriction on q is necessary. The more technical Section 4 describes the stable homotopy-theoretic argument in the metastable range that was outlined above and completes the proof of part (b) of Theorem 1.1.

2. Self-maps of sphere bundles

We consider an orthogonal real vector bundle ξ of dimension *n* over a connected compact ENR *X*.

Definition 2.1. For a (fibrewise) self-map $f : S(\xi) \to S(\xi)$ of the sphere bundle of ξ we define the *degree of* f, $deg(f) \in \mathbb{Z}$, to be the degree of the restriction of fto any fibre, which is independent of the fibre since X is required to be connected.

We review first some results of Noakes [15] from the 1970s, beginning with a basic construction.

Definition 2.2. For a Euclidean vector space V of dimension at least 2 and an integer r, we define a continuous map

$$p_r: S(V) \times S(V) \to S(V)$$

as follows. Given $u, v \in S(V)$ lying in some 2-dimensional subspace $E \subseteq V$, we may choose a complex structure on E (compatible with the inner product) and define $\rho_r(u, v) = z^r u$, where $v = zu, z \in \mathbb{C}, |z| = 1$. Since complex conjugation commutes

with the *r*th power, this is independent of the choice of complex structure, and, since $\rho_r(u, u) = u$ and $\rho_r(u, -u) = (-1)^r u$, it is independent of the choice of *E*.

Remark 2.3. More precisely, writing $O(\mathbb{C}, V)$ for the Stiefel manifold of \mathbb{R} -linear isometric maps $\mathbb{C} \hookrightarrow V$, we have a surjective map

$$S(\mathbb{C}) \times \mathcal{O}(\mathbb{C}, V) \to S(V) \times S(V), \quad (z, a) \mapsto (a(1), a(z)),$$

and ρ_r lifts to the map

$$\tilde{\rho}_r : S(\mathbb{C}) \times \mathcal{O}(\mathbb{C}, V) \to S(V), \quad (z, a) \mapsto (a(1), a(z^r)).$$

This provides a formal proof that ρ_r is continuous.

When $V = \mathbb{C}$ we have the explicit formula

$$S(\mathbb{C}) \times S(\mathbb{C}) \to S(\mathbb{C}) : \rho_r(u, v) = u^{1-r} v^r$$

from which we see that, in general, $\rho_r(v, u) = \rho_{1-r}(u, v)$ and, for a second integer $s, \rho_{rs}(u, v) = \rho_r(u, \rho_s(u, v))$. For any V, when r = 0, the map $v \mapsto \rho_0(u, v) = u$ is constant; and when r = -1, the map $v \mapsto -\rho_{-1}(u, v) = v - 2\langle v, u \rangle u$ is reflection in the hyperplane orthogonal to u.

Lemma 2.4. Suppose that dim V is even. Then, for a fixed $u \in S(V)$, the map $v \mapsto \rho_r(u, v) : S(V) \to S(V)$ has degree r.

The assertion is clear if r = 0 and when dim V = 2 it is evident from the explicit formula.

Proof. We give a proof from a differentiable viewpoint [13].

Suppose that dim V > 2. Let $E \subseteq V$ be the orthogonal complement of $\mathbb{R}u$. Consider the smooth map $\pi : S(\mathbb{C}) \times S(E) \to S(V)$ taking (x + yi, e), where $x, y \in \mathbb{R}$, to xu + ye. The restriction of $\pi : (S(\mathbb{C}) - \{\pm 1\}) \times S(E) \to S(V) - \{\pm u\}$ is a double cover, with covering involution $(x + yi, e) \mapsto (x - yi, -e)$, which is orientation-preserving, for an appropriate choice of orientations of the connected manifolds $S(\mathbb{C}) \times S(E)$ and S(V), because dim V is even.

Now the map $f: v \mapsto \rho_r(u, v)$ lifts to the smooth map $\tilde{f}: (z, e) \mapsto (z^r, e) :$ $S(\mathbb{C}) \times S(E) \to S(\mathbb{C}) \times S(E)$. Its degree is clearly deg $\tilde{f} = r$.

The degree of π , for the chosen orientations, is equal to 2. Because the inverse image of a regular value v in the complement of $\{\pm u\}$ consists of two points at each of which the sign of the determinant of $d\pi$ is +1.

Since $\deg(\pi) \deg(f) = \deg(\pi \circ f) = \deg(f \circ \pi) = \deg(f) \deg(\pi)$, we see that $\deg f = r$.

By the symmetry, the degree of the map $u \mapsto \rho_r(u, v) : S(V) \to S(V)$, for a fixed v, is equal to 1 - r.

Proposition 2.5. (Noakes [15]). Let ξ be an even-dimensional vector bundle. Then there is a non-negative integer $N(\xi)$ such that there is a self-map $S(\xi) \to S(\xi)$ of degree 1 - d if and only if d is divisible by $N(\xi)$.

Proof. Consider two self-maps $f, g: S(\xi) \to S(\xi)$. For each r we may form $h = \rho_r(f,g): S(\xi) \to S(\xi)$. Its degree, by Lemma 2.4, is given by

$$\deg(h) = (1 - r)\deg(f) + r\deg(g),$$

that is, $1 - \deg(h) = (1 - r)(1 - \deg(f)) + r(1 - \deg(g))$. It follows that the set I of integers of the form $1 - \deg(f)$ is an ideal.

(Indeed, $0 \in I$ (because the identity map has degree 1) and $x, y \in I \Rightarrow (1-r)x + ry \in I$ for all $r \in \mathbb{Z}$. If I contains an element $y \neq 0$, then $ry \in I$ for all $r \in \mathbb{Z}$. So I contains a least positive element a. Any element of I can then be written as qa + b with $q \in \mathbb{Z}$, $0 < b \leq a$. So $(1-r)(qa) + r(qa + b) = qa + rb \in I$ for all $r \in \mathbb{Z}$. We must have $qa + sb - b < 0 \leq qa + sb$ for some integer s, and so $0 \leq qa + sb < b \leq a$. Hence qa + sb = 0and $b = qa + (s + 1)b \in I$. By minimality, b = a.)

Corollary 2.6. Let ξ be even-dimensional. If $S(\xi)$ admits a section, then $N(\xi) = 1$ and hence $S(\xi)$ has self-maps of all degrees.

Proof. Given a section s, the self-map taking the constant value s(x) in the fibre at $x \in X$ has degree 0.

Proposition 2.7. (Some properties of $N(\xi)$).

(i). Suppose that ξ and η are two even-dimensional vector bundles over X. If ξ can be embedded as a subbundle of η , then $N(\eta)$ divides $N(\xi)$.

(ii). The cohomology Euler class $e(\xi) \in H^n(X; \mathbb{Z}(\xi))$ with twisted integer coefficients is annihilated by $N(\xi)$, that is, $N(\xi) \cdot e(\xi) = 0$.

(iii). Suppose that ξ is a complex line bundle such that $m \cdot c_1(\xi) = 0 \in H^2(X; \mathbb{Z})$ for some $m \ge 1$. Then $N(\xi)$ divides m.

It follows from (ii) that, if the rational Euler class of ξ is non-zero, then $N(\xi) = 0$.

Proof. (i). We can write $S(\eta) = S(\xi) *_X S(\xi^{\perp})$ as the fibrewise join of the sphere bundles of ξ and its orthogonal complement ξ^{\perp} in η . A fibrewise self-map f of $S(\xi)$ gives a fibrewise self-map $f *_X 1$ (the fibrewise join with the identity on $S(\xi^{\perp})$) of $S(\eta)$ with the same degree.

(ii). The homomorphism $H^n(X; \mathbb{Z}(\xi)) \to H^n(X; \mathbb{Z}(\xi))$ induced by a self-map $f : S(\xi) \to S(\xi)$ of degree d is multiplication by d and takes $e(\xi)$ to itself. So $(1-d)e(\xi) = 0$.

(iii). Choose an isomorphism $\xi^{\otimes (m+1)} \to \xi$ between ξ and its (m+1)-fold complex tensor product. The (m+1)th power map $z \mapsto z^{\otimes (m+1)} : S(\xi) \to S(\xi^{\otimes (m+1)}) \cong S(\xi)$ has degree m+1.

Lemma 2.8. Suppose that there exists a self-map $S(\xi) \to S(\xi)$ of degree 0. Then the sphere bundle $S(\xi)$ admits a section.

Proof. This is a standard nilpotence result. Suppose that $f: E \to E$ is a self-map of a local trivially fibre bundle over X such that the restriction of f to the fibre $f_x: E_x \to E_x$ at $x \in X$ is homotopic to a constant for every x. Then some power of f is homotopic to a map that is constant in each fibre.

(Using the local contractibility of X and compactness, one gets a finite open cover U_i , i = 1, ..., k, of X and maps f_i homotopic to f such that f_i is constant in the fibres over each point of U_i . Then the kth power f^k is homotopic to the composition $f_1 \circ \cdots \circ f_k$, which is constant in every fibre.)

Example 2.9. Suppose that ξ_1 and ξ_2 are complex line bundles over X such that $m_1 \cdot c_1(\xi_1) = 0$ and $m_2 \cdot c_1(\xi_2) = 0$, where m_1 and m_2 are coprime positive integers. Then $\xi = \xi_1 \oplus \xi_2$ is isomorphic to $\mathbb{C} \oplus (\xi_1 \otimes \xi_2)$ as a complex bundle.

Proof. For $N(\xi)$ divides $N(\xi_i)$, which divides m_i , for i = 1, 2. Hence, $N(\xi) = 1$ and so $S(\xi)$ has a section. We conclude that ξ splits as the direct sum of a trivial

line bundle \mathbb{C} and a complex line bundle classified by its first Chern class $c_1(\xi) = c_1(\xi_1) + c_1(\xi_2) = c_1(\xi_1 \otimes \xi_2)$.

We look next at *d*-valued sections of a sphere bundle. (A different class of multivalued sections is considered in [10, Section 5].)

Definition 2.10. Let ξ be an orthogonal real vector bundle over X. For an integer $d \ge 1$, a *d*-valued section of the sphere bundle $S(\xi)$ is a *d*-fold cover $\tilde{X} \to X$ embedded, fibrewise over X, in $S(\xi)$.

Such a d-valued section determines, up to homotopy, a self-map of the sphere bundle. To prepare for the proof, we recall a version of the Pontryagin-Thom Umkehr construction introduced by McDuff in [12].

Let M be a connected, closed smooth manifold with tangent bundle τM . We write $(\tau M)_M^+$ for the fibrewise one-point compactification of τM ; its fibre at $x \in M$ is the one-point compactification $(\tau_x M)^+$ of the tangent space $\tau_x M$ at x. It can be identified by stereographic projection with the sphere bundle $S(\mathbb{R} \oplus \tau M)$ of the direct sum $\mathbb{R} \oplus \tau M$ of the tangent bundle and a trivial line bundle, so that the point at infinity in a fibre corresponds to the 'North Pole' (1,0):

$$v \in \tau_x M \mapsto (\frac{\|v\|^2}{4} + 1)^{-1} (\frac{\|v\|^2}{4} - 1, v) \in S(\mathbb{R} \oplus \tau_x M).$$

Consider a finite subset $N \subseteq M$ with cardinality #N = d. The restriction of τM to N will be denoted by ν ; it is the normal bundle of the inclusion of the 0-dimensional manifold N in M. With a suitably scaled Riemannian metric on M we have a tubular neighbourhood $D(\nu) \hookrightarrow M$ embedding the closed unit disc bundle of the normal bundle ν of N into M. So $D(\nu)$ is the union of d disjoint closed discs $D(\nu_y), y \in N$. Over $D(\nu) \subseteq M$ we can identify the restriction of τM with the pullback of ν .

Construction 2.11. To the subset N, we associate a section s_N of $(\tau M)_M^+$ taking the value ∞ (that is, $s_N(x) = \infty \in (\tau_x M)^+$) outside the open unit disc bundle $B(\nu)$ and given on $D(\nu)$ by

$$s_N(y,v) = [\psi(v)] \in \nu_y^+ = \tau_{(y,v)} M^+ \text{ for } y \in N, \ v \in D(\nu_y),$$

where ψ is the homeomorphism

$$D(\nu_y)/S(\nu_y) \to \nu_y^+ : [v] \mapsto v/(1 - \|v\|^2)^{1/2}.$$

We now look at the special case in which M is the sphere S(V). The tangent space $\tau_x S(V)$ at $x \in S(V) = M$ is the orthogonal complement of $\mathbb{R}x$ in V. The direct sum $\mathbb{R} \oplus \tau S(V)$ is thus identified with the trivial bundle $S(V) \times V$ (with $(1,0) \in \mathbb{R} \oplus \tau_x S(V)$ corresponding to $x \in V$), and so $(\tau M)_M^+ = S(\mathbb{R} \oplus \tau M)$ is identified with the trivial sphere bundle $S(V) \times S(V) \to S(V)$ (projecting onto the first factor) in such a way that $\infty \in (\tau_x S(V))_{S(V)}^+$ corresponds to (x, x).

The space of sections of the trivial bundle $S(V) \times S(V) \to S(V)$ is just the space of self-maps $S(V) \to S(V)$. And so s_N gives a map $f_N : S(V) \to S(V)$. In particular, if N is empty (and d = 0), this gives the identity map $1 : S(V) \to S(V)$.

Lemma 2.12. The degree of f_N is equal to 1 - d.

Proof. For a general connected manifold M of dimension m, the homotopy classes of sections s of the sphere bundle $(\tau M)_M^+$ are classified by an integer deg $(s) \in \mathbb{Z}$. To avoid twisted coefficients, let us assume that M is oriented. The sections s

M. C. CRABB

are determined by the difference class $\deg(s) = \delta(s_{\emptyset}, s) \in H^m(M; \mathbb{Z}) = \mathbb{Z}$, the obstruction to existence of a homotopy from the section s_{\emptyset} taking the value ∞ in each fibre to s. For the section s_N , we have $\delta(s_{\emptyset}, s_N) = d$ (as in [12]).

When we specialize to M = S(V), the section s_{\emptyset} corresponds to the identity map $1 = f_{\emptyset} : S(V) \to S(V)$ and for any map $f : S(V) \to S(V)$ the difference class is equal to $\delta(1, f) = 1 - \deg(f) = \deg(1) - \deg(f) \in \mathbb{Z}$, which is the obstruction to existence of a homotopy from 1 to f. Thus, $d = 1 - \deg(f_N)$ as required.

This argument is described in a more general setting in Proposition 4.7. \Box

The construction that we have described is implicit in the arguments of [1, 2, 3], although not there linked to the, now classical, construction given in [12].

Proposition 2.13. (Compare [3, Lemma 10]). Suppose that the sphere bundle $S(\xi)$ of an orthogonal real vector bundle ξ over X admits a d-valued section. Then there exists a self-map $S(\xi) \to S(\xi)$ of fibre degree 1 - d.

Proof. Since X is compact, for $\epsilon > 0$ sufficiently small, the closed balls $D_{\epsilon}(\tilde{x})$ of radius ϵ (in the Euclidean metric) centred at the points $\tilde{x} \in \tilde{X}$ in any fibre are disjoint. So the construction of f_N for a finite set N can be carried through fibrewise over X to produce a map $f_{\tilde{X}} : S(\xi) \to S(\xi)$ with fibre degree 1 - d. \Box

Remark 2.14. When d = 2 we can write down explicitly a map $f : S(\xi) \to S(\xi)$ of degree 1-2 = -1 specified in each fibre by the (linear) reflection that interchanges the two points of \tilde{X} in the fibre.

Remark 2.15. When ξ is a complex line bundle, the cover $\tilde{X} \to X$ determines a section of the sphere bundle $S(\xi^{\otimes d})$ of the complex *d*-fold tensor power by taking the product of the *d* elements in the fibre of ξ . The section trivializes the line bundle $\xi^{\otimes d}$ and so gives a self-map of degree 1 - d as in Proposition 2.7(iii).

This is already enough to establish Theorem 1.1(a) when n is even.

Proof of Theorem 1.1(a) when n is even. For a prime p dividing #G, choose a point $x \in S(V)$ such that the stabilizer G_x of x contains a Sylow p-subgroup of G. The cardinality d of the orbit Gx is, thus, not divisible by p. Then $\tilde{X} = P \times_G Gx \subseteq P \times_G S(V) = S(\xi)$ gives a d-valued section of $S(\xi)$. By Proposition 2.13, $S(\xi)$ has a self-map of fibre degree 1 - d. Hence $N(\xi)$ divides d, which divides #G. Since p does not divide d, it does not divide $N(\xi)$.

Hence $N(\xi) = 1$, because $N(\xi)$ divides #G and is not divisible by any prime p dividing #G. Thus $S(\xi)$ has a self-map of fibre degree 0 and so has a section, by Lemma 2.8.

Proposition 2.16. Suppose that G is an elementary abelian p-group of order p^s , $s \ge 1$, and V is a G-module such that $V^G = 0$. Then there exists a principal G-bundle $P \to X$ admitting no G-map $P \to S(V)$.

Proof. It is enough to consider a representation $V = \mathbb{C}^k \otimes (\mathbb{R}[G]/\mathbb{R})$, where $k \ge 1$. Take $G = C_p \times \cdots \times C_p$, where C_p is the group of *p*th roots of unity in \mathbb{C} acting on $P = S(\mathbb{C}^m) \times \cdots \times S(\mathbb{C}^m)$ by $(g_i) \cdot (v_i) = (g_i v_i)$. For *m* sufficiently large the \mathbb{F}_p -Euler class $e(\xi) \in H^n(X; \mathbb{F}_p)$ of the complex vector bundle ξ is non-zero. \Box

Remark 2.17. An argument using complex K-theory Euler classes, due to Munkholm [14], shows similarly that, if G is cyclic of prime power order p^s and V is a G-module

 $\mathbf{6}$

such that $V^G = 0$, then there exists a principal *G*-bundle $P \to X$ admitting no G-map $P \to S(V)$. The same result for any *p*-group G^{-1} can be established by using the Euler class in stable cohomotopy and Carlsson's verification of Segal's Burnside ring conjecture [6]. (In the notation of Section 4, the equivariant stable cohomotopy class $\gamma(V) \in \omega_G^0(*; -V)$ of V generates a free summand $\mathbb{Z}\gamma(V)$, because its image under the *G*-fixed-point map $\omega_G^0(*; -V) \to \omega^0(*) = \mathbb{Z}$ is 1. By the Segal conjecture, there is a bundle $P \to X$ such that the Euler class $\gamma(\xi) \in \omega^0(X; -\xi)$, corresponding to the image of $\gamma(V)$ in $\omega_G^0(P; -V)$, is non-zero; the sphere bundle $S(\xi) \to X$ has no section.)

To complete the proof of Theorem 1.1(a), we need a generalization of Proposition 2.13. We start by extending Construction 2.11.

Construction 2.18. Suppose that we are given a family h of pointed maps h_x : $\nu_x^+ \rightarrow \nu_x^+$ for $x \in N$. To the subset N and family h, we associate a section $s_{N,h}$ of $(\tau M)_M^+$ taking the value ∞ outside the open unit disc bundle $B(\nu)$ and given on $D(\nu)$ by

$$\beta_{N,h}(y,v) = [h_y(\psi(v))] \in \nu_y^+ = \tau_{(y,v)}M^+ \text{ for } y \in N, \ v \in D(\nu_y).$$

When M = S(V) is a sphere, the section $s_{N,h}$ corresponds to a self-map $f_{N,h}$: $S(V) \to S(V)$, which (by the argument used to prove Lemma 2.12) has degree $1 - \sum_{y \in N} \deg h_y$.

Suppose that $\pi : \tilde{X} \to X$ is a finite cover defining a *d*-valued section of $S(\xi)$. Then the fibre of $\pi^*\xi$ at $u \in \tilde{X}$ splits as an orthogonal direct sum $\mathbb{R}u \oplus \eta_x$ and $\pi^*\xi$ is, thus, a direct sum $\mathbb{R} \oplus \eta$ of a trivial line bundle and a real vector bundle η of dimension n-1. Suppose that the space \tilde{X} is expressed as a disjoint union of subspaces $\tilde{X}_{\gamma}, \gamma \in \Gamma$ such that the restriction $\tilde{X}_{\gamma} \to X$ of π is a finite cover, of order d_{γ} say. (For example, the \tilde{X}_{γ} might be the connected components of \tilde{X} .)

Proposition 2.19. (Compare [2, Lemma 3.1]). In the notation introduced above, suppose that $h: S(\eta) \to S(\eta)$ is a self-map of the sphere bundle $S(\eta)$ over \tilde{X} such that, for each $\gamma \in \Gamma$, the restriction of h to \tilde{X}_{γ} has constant fibre degree, m_{γ} say.

Then $S(\xi)$ admits a self-map of fibre degree

$$1 - \sum_{\gamma \in \Gamma} m_{\gamma} d_{\gamma} \, .$$

Proof. The bundle map $h: S(\eta) \to S(\eta)$ extends radially to a map $\eta_{\tilde{X}}^+ \to \eta_{\tilde{X}}^+$. Then the construction of $f_{N,h}$ above can be performed fibrewise over X to produce the required self-map $f_{\tilde{X},h}$ of $S(\xi)$ as in Proposition 2.13.

Proof of Theorem 1.1(a) when n is odd. Consider a prime p dividing #G and a Sylow p-subgroup H of G. Since $S(V^H)$ is an infinite set and G has only finitely many subgroups, there exist two points $x, y \in S(V^H)$ such that $G_x = G_y$ and $y \notin \{\pm x\}$. It is possible that G_x contains a Sylow p'-subgroup for some other prime p'.

We can, therefore, choose finitely many points x_{γ} , y_{γ} , indexed by $\gamma \in \Gamma$, such that: (i) $x_{\gamma} \neq \pm y_{\gamma}$ and $G_{x_{\gamma}} = G_{y_{\gamma}}$; (ii) for each prime p dividing #G at least one of the subgroups $G_{x_{\gamma}}$ contains a Sylow p-subgroup of G; (iii) if $\gamma \neq \delta$, then the subgroups $G_{x_{\gamma}}$ and $G_{x_{\delta}}$ of G are not conjugate.

¹I am grateful to Roman Karasev for pointing this out.

M. C. CRABB

Let $\tilde{X}_{\gamma} = P \times_G Gx_{\gamma} \subseteq P \times_G S(V) = S(\xi)$. Then $\tilde{X}_{\gamma} \to X$ is a finite cover of order $d_{\gamma} = \#Gx_{\gamma}$. If $\gamma \neq \delta$, the sets \tilde{X}_{γ} and \tilde{X}_{δ} are disjoint, because the stabilizers $G_{x_{\gamma}}$ and $G_{x_{\delta}}$ are not conjugate. Take $\tilde{X} = \bigsqcup_{\gamma \in \Gamma} \tilde{X}_{\gamma}$. By construction, d_{γ} divides #G and for each prime p dividing #G there is some γ such that p does not divide d_{γ} . So there are integers m_{γ} such that $1 - \sum_{\gamma} m_{\gamma} d_{\gamma} = 0$.

Now y_{γ} gives a section of $S(\pi^*\xi) = S(\mathbb{R} \oplus \eta)$ over \tilde{X}_{γ} : $[a, gx_{\gamma}] \mapsto [a, gy_{\gamma}]$ for $a \in P, g \in G$ (well-defined, because $G_{x_{\gamma}} = G_{y_{\gamma}}$). Then, because $gy_{\gamma} \neq \pm gx_{\gamma}$, the η -component is non-zero and so determines a section of $S(\eta)$ over \tilde{X}_{γ} . Hence, there is a map $S(\eta) \to S(\eta)$ with degree m_{γ} over \tilde{X}_{γ} .

It is easy to write down a counterexample if the assumption that the dimension of V^H is greater than 1 is omitted.

Proposition 2.20. Let G be cyclic of order 6, generated by an element g, and let A and B be the G-modules $A = \mathbb{R}$, $B = \mathbb{C}$ with g acting as $t \mapsto -t$, $z \mapsto e^{2\pi i/3}z$, respectively. Take $V = A \oplus B$ and let P be the 6-dimensional free G-space $S(\mathbb{R}^2 \otimes A) \times S(\mathbb{R}^3 \otimes B)$. Then

(i). There is a G-map $S(V) \to S(V)$ with non-equivariant degree d if and only if $d \equiv \pm 1 \pmod{3}$.

(ii). There is no G-map $P \to S(V)$.

Proof. (i). For the non-equivariant degree of a *G*-map is congruent (mod 3) to the degree of its restriction $S(\mathbb{R}) \to S(\mathbb{R})$ to the fixed subspace $S(V^H)$ of the cyclic subgroup *H* of order 3. And we can realize a map of degree $\pm(3r+1)$ by the join of $t \mapsto \pm t : S(\mathbb{R}) \to S(\mathbb{R})$ and $z \mapsto z^{3r+1} : S(\mathbb{C}) \to S(\mathbb{C})$.

(ii). Let α and β be the real 1- and 2-dimensional vector bundles over $X = P/G = S(\mathbb{R}^2 \otimes A)/\langle g^3 \rangle \times S(\mathbb{R}^3 \otimes B)/\langle g^2 \rangle$ associated with the modules A and B. So $\xi = \alpha \oplus \beta$.

Now, writing $\tilde{\mathbb{Z}}$ for integer coefficients twisted by the Hopf line bundle over S^1 , we have, for any compact ENR Y, an equivalence

$$H^i(S^1 \times Y; \tilde{\mathbb{Z}}) = H^{i-1}(Y; \mathbb{F}_2), \quad i \in \mathbb{Z},$$

given by reduction $(\mod 2)$ and the projection to the second factor

$$H^{i}(S^{1} \times Y; \widetilde{\mathbb{Z}}) \to H^{i}(S^{1}; \mathbb{F}_{2}) = H^{i}(Y; \mathbb{F}_{2}) \oplus H^{i-1}(Y; \mathbb{F}_{2}) \to H^{i-1}(Y; \mathbb{F}_{2}).$$

(This is the classical statement that the real projective plane is a mod 2 Moore space. Given the naturality, it is enough to check when Y is a point.)

There is only one non-orientable 2-dimensional real vector bundle over X, namely $\mathbb{R} \oplus \alpha$, because the group $H^2(X; \mathbb{Z}(\alpha)) = H^1(S(\mathbb{R}^3 \otimes B)/\langle g^2 \rangle; \mathbb{F}_2)$ that classifies such bundles is zero. Since the Pontryagin class $p_1(\beta) \in H^4(X; \mathbb{Z})$ is non-zero, ξ is not isomorphic to $\mathbb{R} \oplus (\mathbb{R} \oplus \alpha)$.

Theorem 2.21. Suppose that for each prime p dividing #G the fixed subspace V^H of a Sylow p-subgroup H is non-zero and, if n is odd, of dimension greater than 1. Then there exists a G-map $S(V) \rightarrow S(V)$ with non-equivariant degree equal to 0.

Proof. Indeed, since the definition of ρ_r , Definition 2.2, is equivariant with respect to the action of the orthogonal group of V, for any even-dimensional G-module V there is (as was already shown in [15]) a non-negative integer $N_G(V)$ such that there is a G-map $S(V) \to S(V)$ with non-equivariant degree 1 - d if and only if d is divisible by $N_G(V)$. And then it follows from the equivariant version of Proposition

2.13 that, if $V^H \neq 0$ for a Sylow *p*-subgroup *H*, the prime *p* does not divide $N_G(V)$. Hence, if this holds for all primes, we must have $N_G(V) = 1$.

The proof when n is odd uses the equivariant version of Proposition 2.19.

Proposition 2.22. Suppose that G is a p-group and that $V^G = 0$. Then any G-map $S(V) \rightarrow S(V)$ has non-equivariant degree congruent to $1 \pmod{p}$. If n is even, then $N_G(V)$ is divisible by p. If n is odd, p = 2.

Proof. In general, for the non-trivial *p*-group *G*, the degree of a *G*-map $f: S(V) \to S(V)$ is congruent (mod *p*) to the degree of its restriction $f^G: S(V^G) \to S(V^G)$ to the fixed subspace. (See, for example, [11]. The map *f* has an equivariant degree in the Burnside ring A(G) of *G*, the Grothendieck group of finite *G*-sets. And so the assertion reduces to the fact that the cardinality of a finite *G*-set is congruent (mod *p*) to the cardinality of the fixed subset.)

If n is odd, the map $-1: S(V) \to S(V)$ has degree -1.

Proposition 2.23. Suppose that G is a group of order $2p^s$, where p is an odd prime, $s \ge 1$, and that $V^G = 0$. Suppose further that for a Sylow p-subgroup H, $\dim V^H = 1$. Then any G-map $S(V) \to S(V)$ has non-equivariant degree congruent to $\pm 1 \pmod{p}$.

If n is even, then $N_G(V)$ is divisible by p.

Proof. Observe that H is a normal subgroup and the quotient group G/H acts on V^H as ± 1 . The degree of a G-map $f: S(V) \to S(V)$ is congruent (mod p) to the degree of its restriction $f^H: S(V^H) \to S(V^H)$. Since f^H commutes with the action of G/H, we must have $f^H = \pm 1$.

When n is even, for each $r \in \mathbb{Z}$ there is a G-map $S(V) \to S(V)$ with degree $1 - rN_G(V)$, and so the product $rN_G(V)(rN_G(V) - 2)$ must be divisible by p. This is only possible if p divides $N_G(V)$.

3. Symmetric groups

For a subset $J \subseteq I = \{1, \ldots, q\}$, as in the Intrduction, we write $e_J = \sum_{i \in J} e_i \in \mathbb{R}[I]$ and $\overline{e}_J \in L$ for its image in the quotient $L = \mathbb{R}[I]/\mathbb{R}e_I$. Consider a prime $p \leq q$ and write $q = \sum_r a_r p^r$, where $0 \leq a_r < p$. A Sylow *p*-subgroup *H* of \mathfrak{S}_q determines a partition

$$I = \bigsqcup_{i=1}^{\alpha_p(q)} J_i$$

of I into $\alpha_p(q) = \sum_r a_r$ subsets each of order a power of p with a_r of order p^r . The fixed subspace $\mathbb{R}[I]^H$ has dimension $\alpha_p(q)$ with the basis e_{J_i} . So L^H has dimension $\alpha_p(q) - 1$ and is spanned by the \overline{e}_{J_i} subject to the single relation $\sum_i \overline{e}_{J_i} = 0$. The order of a Sylow p-subgroup is the highest power, $p^{\nu_p(q)}$, of p that divides $\#\mathfrak{S}_q$, where the p-adic valuation $\nu_p(q!)$ of q! is equal to $(q - \alpha_p(q))/(p - 1)$.

As in the statement of Corollary 1.2, we consider a principal \mathfrak{S}_q -bundle $P \to X$ over a compact ENR and write λ for the (q-1)-dimensional vector bundle $P \times_{\mathfrak{S}_q} L$ over X. Let μ be some odd-dimensional Euclidean vector bundle over X.

Lemma 3.1. Suppose that p is a prime and that q is not a power of p. If q is odd, then p does not divide $N(\lambda)$; if q is even, then p does not divide $N(\lambda \oplus \mu)$.

Proof. Choose a point $v \in S(L)$ that is fixed by some Sylow *p*-subgroup or, equivalently, so that its \mathfrak{S}_q -orbit D has size d which is not divisible by p.

For example, if p^s is the highest power of p dividing q, we could take $v = \overline{e}_J / \|\overline{e}_J\|$, where $J \subseteq I$ has cardinality p^s . Then $D = \{\overline{e}_K / \|\overline{e}_K\| \mid K \subseteq I, \#K = p^s\}$ and $d = \binom{q}{p^s}$.

Take $\tilde{X} = P \times_{\mathfrak{S}_q} D \subseteq P \times_{\mathfrak{S}_q} S(L) = S(\lambda) \subseteq S(\lambda \oplus \mu)$. By Proposition 2.13, $S(\lambda)$ and $S(\lambda \oplus \mu)$ admit self-maps of degree 1 - d. So, if q is odd, $N(\lambda)$ divides d, and, if q is even, $N(\lambda \oplus \mu)$ divides d.

Lemma 3.2. Suppose that q is a power $q = p^s$, $s \ge 1$, of a prime p. If p is odd, then $N(\lambda)$ is either 1 or p; if p = 2, then $N(\lambda \oplus \mu)$ is either 1 or 2.

Proof. For we can take $D = \{\overline{e}_J \mid \#J = p^{s-1}\}$. Then $\#D = \binom{p^s}{p^{s-1}}$ is divisible by p but not p^2 . (For $\alpha_p(p^{s-1}) + \alpha_p(p^s - p^{s-1}) - \alpha_p(p^s) = p - 1$.)

Proposition 3.3. Suppose that $q = p^s$ is a prime power, $s \ge 1$, and $k \ge 1$. Then there exists a principal \mathfrak{S}_q -bundle $P \to X$ admitting no \mathfrak{S}_q -map $P \to S(\mathbb{R}^k \otimes L)$.

Proof. Let G be an elementary abelian p-group of order p^s . Fixing a bijection $I \to G$, we can identify G with a subgroup of \mathfrak{S}_q such that the restriction V of $\mathbb{R}^k \otimes L$ to G satisfies $V^G = 0$. According to Proposition 2.16 there exists a principal G-bundle Q such that there is no G-map $Q \to S(V)$. Then there is no \mathfrak{S}_q -map $P = Q \times_G \mathfrak{S}_q \to S(L)$.

A similar argument to that in Proposition 2.20 shows that in Corollary 1.2 when k = 1 and q is even some additional condition is essential.

Proposition 3.4. Take $q = 2p^s$, where p > 2 is an odd prime and $s \ge 1$. Then there exists a principal \mathfrak{S}_{2p^s} -bundle $P \to X$ such that there is no \mathfrak{S}_{2p^s} -map $P \to S(L)$.

Proof. Let $G = C_2 \times H$ be the product of a cyclic group of order 2 and an elementary abelian *p*-group *H* of order p^s . Fix a bijection $I \to G$ to identify *G* with a subgroup of \mathfrak{S}_{2p^s} and the restriction of *L* to *G* with

$$V = A \oplus B \oplus (A \otimes B)$$

where $A = \mathbb{R}$ is the non-trivial C_2 -module with the generator acting as -1 and $B = \mathbb{R}[H]/\mathbb{R}$, of dimension $p^s - 1$, is the reduced regular representation of H.

Taking $H = C_p \times \cdots \times C_p$ (where again C_p is the group of *p*th roots of unity in \mathbb{C}), let *R* be the free *H*-space $S(\mathbb{C}^m) \times \cdots \times S(\mathbb{C}^m)$ with the action $(h_i) \cdot (v_i) = (h_i v_i)$ for *m* sufficiently large, say $m = 2p^s - 1$. Let *Q* be the free *G*-space $S(\mathbb{R}^2 \otimes A) \times R$ of dimension (2m-1)s+1, and let α and β be the real line bundle and (p^s-1) -dimensional real vector bundle over $X = Q/G = S(\mathbb{R}^2 \otimes A)/C_2 \times R/H$ associated with the modules *A* and *B* respectively. We take $P = Q \times_G \mathfrak{S}_{2p^s}$, so that over $X = P/\mathfrak{S}_{2p^s}$

$$\lambda = \alpha \oplus \beta \oplus (\alpha \otimes \beta).$$

Suppose that η is a $(2p^s - 2)$ -dimensional non-orientable real vector bundle over $X = Q/G = S^1 \times R/H$. Now, as in the proof of Proposition 2.20, the group

$$H^{2(p^s-1)}(X; \mathbb{Z}(\alpha)) = H^{2p^s-3}(R/H; \mathbb{F}_2)$$

is trivial. So $e(\eta) \in H^{2(p^s-1)}(X; \mathbb{Z}(\alpha))$ is zero. Recall that, for a real vector bundle ξ of even dimension 2n, the *n*th Pontryagin class is equal to the square of the Euler

class: $p_n(\xi) = (-1)^n c_{2n}(\mathbb{C} \otimes \xi) = (-1)^n e(\mathbb{C} \otimes \xi) = e(\xi \oplus \xi) = e(\xi)^2$. (The sign relates the orientations on $\mathbb{C} \otimes \xi$ and $\xi \oplus \xi$.) So $p_{p^s-1}(\eta) = e(\eta)^2 \in H^{4(p^s-1)}(X; \mathbb{Z})$ is zero. But $p_{p^s-1}(\lambda)$ is non-zero, because

$$p_{p^s-1}(\beta \oplus \beta) = e(\beta \oplus \beta)^2 = e(\beta)^4 \neq 0,$$

by the choice of m. So λ cannot be isomorphic to $\mathbb{R} \oplus \eta$.

There is, thus, no G-map
$$Q \to S(V)$$
 and no \mathfrak{S}_{2p^s} -map $P \to S(L)$.

But it is shown in [2] that these (Propositions 3.3 and 3.4) are the only exceptional cases.

Theorem 3.5. ([2, Theorem 1.1]). Suppose that q is even, not a power of 2, and not twice a power of an odd prime. Then, for any principal \mathfrak{S}_q -bundle $P \to X$, there exists an \mathfrak{S}_q -equivariant map $P \to S(L)$.

The first example is $q = 12 = 2^3 + 2^2 = 3^2 + 3^1$.

Proof. Let Γ denote the set of primes $p \leq q$ and $\Gamma_2 \subseteq \Gamma$ the set of primes $p \leq q$ with $\alpha_p(q) = 2$.

For each prime $p \in \Gamma$, choose a partition $I = \bigsqcup J_i$ as above into $\alpha_p(q)$ subsets, and let G_p be the subgroup of $G = \mathfrak{S}_q$ that preserves each subset J_i : G_p = $\prod_i \mathfrak{S}_{J_i} \leqslant \mathfrak{S}_I.$

If $\alpha_p(q) > 2$, we can choose two points x_p and $y_p \neq \pm x_p$ in S(L), of the form $\sum_i t_i \overline{e}_{J_i}$, where $t_i \neq t_j$ for $i \neq j$ and $\sum_i t_i = 0$, with stabilizer exactly G_p . Since G_p contains a Sylow *p*-subgroup of \mathfrak{S}_q , the the order d_p of the \mathfrak{S}_q -orbit of x_p is not divisible by p.

If $\alpha_p(q) = 2$, say $q = p^s + p^t$, where s > t we choose a single point x_p with stabilizer exactly G_p . Note that, in this case, $d_p = \#(G/G_p) = {p^s + p^t \choose n^t}$ is congruent to 1 (mod p). (For $(1+T)^{p^s+p^t} = (1+T^{p^t})^{p^{s-t}+1}$ in $\mathbb{F}_p[T]$.) Also, p divides $d_{p'}$ for $p' \neq p.$ (If $r_1 + \ldots + r_a = p^s + p^t$, where $r_i \ge 1$, then $\nu_p((r_1 + \ldots + r_a)!/(r_1! \cdots r_a!)) = r_a$ $(\alpha_p(r_1) + \ldots + \alpha_p(r_a) - 2)/(p-1).)$

Notice that the G-orbits of the points x_p are disjoint. We write $\tilde{X}_p = P \times_G G x_p$ and $X = \bigsqcup_p X_p \subseteq S(\lambda)$. Following the notation of Proposition 2.19, we have a vector bundle η over \tilde{X} of even dimension q-2. Let η_p be the restriction of η to \tilde{X}_p . Then $N(\eta_p) = 1$ if $p \notin \Gamma_2$, because y_p gives a section of $S(\eta_p)$, and $N(\eta_p) = 1$ or p if $p \in \Gamma_2$, by Lemma 3.2.

Now, (following the argument in [2]) we notice that $c = 1 - \sum_{p \in \Gamma} d_p$ is divisible by each $p \in \Gamma_2$. We can write $1 = \sum_{p \in \Gamma} n_p d_p$, and so $1 = \sum_{p \in \Gamma} (1 + cn_p) d_p =$ $\sum_{p \in \Gamma} m_p d_p$, where $m_p = 1 + cn_p$ is congruent to $1 \pmod{p}$ if $p \in \Gamma_2$.

The proof is completed by applying Proposition 2.19.

4. Euler classes and stable homotopy theory

Stable cohomotopy will be denoted by ω^* , and we make the abbreviation

$$\omega^0(X; -\xi) = \tilde{\omega}^0(X^{-\xi})$$

for the reduced stable cohomotopy of the Thom space $X^{-\xi}$ of the virtual bundle $-\xi$. Thus $\omega^0(X; -\xi)$ may be identified with the stable cohomotopy group $\omega^{N-n}(X^{\xi^{\perp}})$ where ξ^{\perp} is the orthogonal complement of ξ included in a trivial bundle $X \times \mathbb{R}^N$ of sufficiently high dimension N. Similar notation in cohomology H^* with integral coefficients would write $H^0(X; -\xi)$, using the Thom isomorphism, for $H^n(X; \mathbb{Z}(\xi))$.

The stable cohomotopy Euler class $\gamma(\xi) \in \omega^0(X; -\xi)$ of ξ is defined like the classical Euler class $e(\xi) \in H^n(X; \mathbb{Z}(\xi))$. Further details may be found, for example, in [9, II, Section 4].

The stable cohomotopy Euler class

$$\gamma(\xi) \in \omega^0(X; -\xi),$$

like $e(\xi)$, is zero if the sphere bundle $S(\xi)$ has a section. In a metastable range the converse is true, by Freundenthal's suspension theorem.

Proposition 4.1. Suppose that dim $X < 2(\dim \xi - 1)$. If $\gamma(\xi) = 0$, then $S(\xi)$ admits a section.

The vanishing of the cohomology Euler class $e(\xi) \in H^n(X; \mathbb{Z}(\xi))$ is sufficient for the existence of a section if dim $X \leq \dim \xi$.

Proposition 4.2. (See, for example, [9, II, Lemma 12.45]). Suppose that $\pi : \tilde{X} \to X$ is a d-fold cover. If the pullback $S(\pi^*\xi) \to \tilde{X}$ of the sphere bundle of ξ admits a section, then

$$d^k \gamma(\xi) = 0 \in \omega^0(X; -\xi)$$

for $k \ge 1$ sufficiently large.

Proof. Since $S(\pi^*\xi)$ admits a section, the stable cohomotopy Euler class $\pi^*\gamma(\xi) = \gamma(\pi^*\xi) \in \omega^0(\tilde{X}; -\pi^*\xi)$ is zero. Hence $\pi_!\pi^*\gamma(\xi) = \pi_!(1)\gamma(\xi) \in \omega^0(X; -\xi)$ is zero, where $\pi_! : \omega^0(\tilde{X}) \to \omega^0(X)$ is the transfer (or direct image) homomorphism. Now $\pi_!(1) \in \omega^0(X)$ can be written as d-x where x is nilpotent. Thus $d\gamma(\xi) = x \cdot \gamma(\xi)$, and $d^k\gamma(\xi) = x^k \cdot \gamma(\xi)$ is zero for large k. \Box

Remark 4.3. The same argument in cohomology rather than stable cohomotopy shows that $d \cdot e(\xi) = 0 \in H^n(X; \mathbb{Z}(\xi)).$

Remark 4.4. If $\tilde{X} \to X$ is embedded in $S(\xi)$, so defining a *d*-valued section, then $S(\pi^*\xi)$, as we observed in Section 2, clearly has a section.

Proposition 4.5. Suppose that $S(\xi)$ has a self-map f of degree 1 - d. Then

$$d^k \gamma(\xi) = 0 \in \omega^0(X; -\xi)$$

for $k \ge 1$ sufficiently large.

Proof. We have $\gamma(\xi) = [f] \cdot \gamma(\xi)$, where $[f] \in \omega^0(X)$ is the class represented by f. Now [f] = 1 - d + x, where x is nilpotent. So the result follows as before. \Box

Corollary 4.6. If dim ξ is odd, then $2^k \gamma(\xi) = 0$ for k large.

Proof. The antipodal map $-1: S(\xi) \to S(\xi)$ has degree -1.

Proposition 4.7. If f is the map constructed in Propositions 2.13 and 2.19 from a d-valued section of $S(\xi)$ specified by a d-fold cover $\pi : \tilde{X} \to X$ and fibrewise embedding $\iota : \tilde{X} \hookrightarrow S(\xi)$ over X and a map $h : S(\eta) \to S(\eta)$, then

$$[f] = 1 - \pi_!([h]) \in \omega^0(X),$$

where $[h] \in \omega^0(\tilde{X})$ is the class represented by h.

Proof. The section $s_{\tilde{X},h}$ in Construction 2.18 realizes the fibrewise Umkehr homomorphism

$$\iota_!: \omega^0(\tilde{X}) \to \omega^0(S(\xi); -\tau_X S(\xi)),$$

where $\tau_X S(\xi)$ is the fibrewise tangent bundle of $S(\xi) \to X$. (See, for example, [9, II, Chapter 3].) Thus, $[s_{\tilde{x},h}] = \iota_!([h])$. The projection π is the composition of ι and the projection $\sigma : S(\xi) \to X$. Hence, $\sigma_![s_{\tilde{x},h}] = \pi_![h]$.

Recall that two sections s_0, s_1 of the pullback $S(\sigma^*\xi)$ determine a difference class $\delta(s_0, s_1) \in \omega^{-1}(S(\xi); -\xi)$ constructed as an obstruction to the existence of a homotopy between the two sections.

The class

$$[s_{\tilde{x},h}] \in \omega^0(S(\xi); -\tau_X S(\xi)) = \omega^{-1}(S(\xi); -\xi)$$

can be expressed as $\delta(s_{\emptyset}, s_{\tilde{X},h})$, where s_{\emptyset} is the section at ∞ of $(\tau_X S(\xi))_{S(\xi)}^+ = S(\mathbb{R} \oplus \tau_X S(\xi))$. In the Gysin sequence

$$\cdots \to \omega^{-1}(S(\xi); -\xi) \xrightarrow{\delta} \omega^0(X) \xrightarrow{\gamma(\xi)} \omega^0(X; -\xi) = \omega^0(D(\xi), S(\xi); -\xi) \to \cdots$$

of the sphere bundle $S(\xi)$, that is, the exact sequence of the pair $(D(\xi), S(\xi))$, the coboundary δ coincides with the Umkehr map $\sigma_!$ and so maps the difference class to $\sigma_![s_{\tilde{X},h}] = \pi_![h]$.

Given any two self-maps $f_0, f_1 : S(\xi) \to S(\xi)$, the difference class $\delta(s_0, s_1)$ of the corresponding sections s_0 and s_1 of $S(\sigma^*\xi)$ maps, in the Gysin sequence, to the difference $\gamma(\xi, s_0) - \gamma(\xi, s_1)$ of the relative Euler classes $\gamma(\xi, s_0)$ and $\gamma(\xi, s_1)$ in $\omega^0(D(\xi), S(\xi); -\xi)$. (See [9, II, Lemma 4.12].) The relative Euler class $\gamma(\xi, s_i)$ is an obstruction to extending $f_i : S(\xi) \to S(\xi)$ to a map $D(\xi) \to S(\xi)$ and corresponds to $[f_i] \in \omega^0(X)$. Applying this to the self-maps $f_0 = 1$ and $f_1 = f$, which correspond, respectively, to the sections s_{\emptyset} and $s_{\bar{X},h}$, we obtain the required identity $1 - [f] = \pi_1[h]$.

Thus far, the arguments have applied to a general vector bundle ξ over a compact ENR X. We now specialize to the case that $\xi = P \times_G V$ as in the Introduction.

Corollary 4.8. Let p be a prime. Suppose that for a Sylow p-subgroup $H \leq G$ the fixed subspace V^H is non-zero. Then the p-local Euler class $\gamma(\xi) \in \omega^0(X; -\xi)_{p}$ is zero.

Proof. Take $\pi : \tilde{X} \to X$ in Proposition 4.2 to be the covering projection $P/H \to P/G$ with order d = #(G/H) prime to p.

Corollary 4.9. Suppose that for each prime p dividing #G the fixed subspace V^H of a Sylow p-subgroup H of G is non-zero. Then the stable cohomotopy Euler class $\gamma(\xi) \in \omega^0(X; -\xi)$ is zero.

Proof. Since, by Corollary 4.8, $\gamma(\xi)$ vanishes at any prime p, we conclude that the stable cohomotopy Euler class of ξ is zero.

Corollary 4.10. Suppose that for each prime p dividing #G the fixed subspace V^H of a Sylow p-subgroup H of G is non-zero. If dim X < 2(n-1), then the sphere bundle $S(\xi) \to X$ has a section.

Proof. Since $\gamma(\xi) = 0 \in \omega^0(X; -\xi)$, by Corollary 4.9, the conclusion follows from Proposition 4.1.

Proof of Theorem 1.1(b). If n is odd, $\gamma(\xi)$ is 2-torsion, by Corollary 4.6. So it enough to show that the 2-local Euler class in $\omega^0(X; -\xi)_{(2)}$ is zero.

Theorem 4.11. Let $r \ge 1$ be a positive integer such that dim X < 2(n-r) and r < n. Suppose that, for each prime p dividing #G if n is even, for p = 2 if n is odd and #G is even, the dimension of the fixed subspace V^H of a Sylow p-subgroup H of G is at least r. Then there exists a G-map

$$P \to \mathcal{O}(\mathbb{R}^r, V)$$

to the Stiefel manifold of orthogonal r-frames in V.

Proof. In this range of dimensions the Stiefel bundle

$$O(\mathbb{R}^r,\xi) = P \times_G O(\mathbb{R}^r,V) \to X$$

has a section if and only if the stable cohomotopy Euler class of the tensor product $\eta \otimes \xi$ of the Hopf line bundle η over the real projective space $P(\mathbb{R}^r)$ of \mathbb{R}^r with ξ ,

$$\gamma(\eta \otimes \xi) \in \omega^0(P(\mathbb{R}^r) \times X; -\eta \otimes \xi),$$

is zero. (See, for example, [7, Theorem 1.5].)

For a Sylow *p*-subgroup *H* of *G*, consider the finite cover $\pi : \tilde{X} = P/H \to X = P/G$ with order prime to *p*. The pullback $\pi^*\xi$ has a trivial subbundle of dimension *r* and $S(\eta \otimes \pi^*\xi)$ has a section. So $\gamma(\eta \otimes \pi^*\xi) \in \omega^0(P(\mathbb{R}^r) \times \tilde{X}; -\eta \otimes \pi^*\xi)$ is zero. From Proposition 4.2 applied to $\eta \otimes \xi$, we deduce that $\gamma(\eta \otimes \xi)$ is zero at the prime (*p*). Hence, using Corollary 4.6 if *n* is odd, we conclude that $\gamma(\eta \otimes \xi)$ is zero. \Box

References

- S. Avvakumov and R. Karasev, Envy-free division using mapping degree. Mathematika 67 (2021), 36–53.
- [2] S. Avvakumov and S. Kudrya, Vanishing of all equivariant obstructions and the mapping degree. Discrete and Computational Geometry 66 (2021), 1202–1216.
- [3] S. Avvakumov, R. Karasev and A. Skopenkov, Stronger counterexamples to the topological Twerberg conjecture. arXiv math.AT 1907.11183 (2019).
- [4] P. V. M. Blagojević and G. M. Ziegler, Convex equipartitions via equivariant obstruction theory. Israel J. Math. 200 (2014), 49–77.
- [5] P. V. M. Blagojević and G. M. Ziegler, Beyond the Borsuk-Ulam theorem: The topological Tverberg story. In: A Journey Through Discrete Mathematics. A tribute to Jiří Matoušek (eds. M. Loebl, J. Nešetřil, R. Thomas), 273–341, Springer, Cham, 2017.
- [6] G. Carlsson, Equivariant stable homotopy and Segal's Burnside ring conjecture. Ann. of Math. 120 (1984), 189–224.
- [7] M. C. Crabb, On the KO_{ℤ/2}-Euler class, I. Proc. R. Soc. Edinburgh Sect. A 117 (1991), 115–137.
- [8] M. C. Crabb, The topological Tverberg theorem and related topics. J. Fixed Point Theory Appl. 12 (2012), 1–25.
- [9] M. C. Crabb and I. M. James, *Fibrewise homotopy theory*. Springer Monographs in Mathematics. London: Springer, 1998.
- [10] M. C. Crabb and J. Jaworowski, Aspects of the Borsuk-Ulam theorem. J. Fixed Point Theory Appl. 13 (2013), 459–488.
- [11] T. tom Dieck, Transformation groups. Walter de Gruyter, Berlin, 1987.
- [12] D. McDuff, Configuration spaces of positive and negative particles. Topology 14 (1975), 91– 107.
- [13] J. W. Milnor, Topology from the differentiable viewpoint. Princeton University Press, Princeton, New Jersey, 1997.
- [14] H. J. Munkholm, On the Borsuk–Ulam theorem for \mathbb{Z}_{p^a} -actions on S^{2n-1} and maps $S^{2n-1} \to \mathbb{R}^m$. Osaka J. Math. 7 (1970), 451–456.
- [15] J. L. Noakes, Symmetric overmaps. Proc. Amer. Math. Soc. 56 (1976), 333-336.

[16] D. Sullivan, Geometric topology. Localization, periodicity and Galois symmetry. The 1970 MIT Notes: edited by A. Ranicki. Springer, Dordrecht, 2005.

Institute of Mathematics,, University of Aberdeen, Aberdeen AB24 3UE, UK $\mathit{Email}\ address: \texttt{m.crabb@abdn.ac.uk}$