### STABILITY THEOREMS FOR POSITIVELY GRADED DOMAINS

## SOURJYA BANERJEE

ABSTRACT. Given a commutative Noetherian graded domain  $R=\bigoplus_{i\geq 0}R_i$  of dimension  $d\geq 2$  with  $\dim(R_0)\geq 1$ , we prove that any unimodular row of length d+1 in R can be completed to the first row of an invertible matrix  $\alpha$  such that  $\alpha$  is homotopic to the identity matrix. Utilizing this result, it has been established that if  $I\subset R$  is an ideal satisfying  $\mu(I/I^2)=\operatorname{ht}(I)=d$ , then any set of generators of  $I/I^2$  lifts to a set of generators of I, where  $\mu(-)$  denotes the minimal number of generators. Consequently, any projective R-module of rank d with trivial determinant splits into a free factor. We also improve the injective stability of  $K_1(R)$ . Finally, we prove that for any projective R-module P of rank d, if the Quillen ideal of P is non-zero, then P is cancellative.

#### 1. Introduction

We commence by recalling an old question of Murthy [4]. Let  $A=\bigoplus_{i\geq 0}R_i$  be a normal positively graded finitely generated algebra over  $R_0$ , where  $R_0=k$  is a field. Then Murthy asked whether  $K_0(A)\cong\mathbb{Z}$ . Bloch provided a counterexample to this question by considering  $A=\frac{\mathbb{C}[X,Y,Z]}{\langle Z^2-X^3-Y^7\rangle}$ . However, if (1) k is an algebraically closed field of characteristic p>0, (2) A is a Cohen-Macaulay ring of dimension 2, and (3) the vertex (corresponding to the ideal  $\bigoplus_{i\geq 1}R_i$ ) is the only singularity of Spec(A), then Srinivas [37, Corollary 1.3] showed that Murthy's question has an affirmative answer. Therefore, using the cancellation theorem of Murthy and Swan [30], it follows that every projective A-module is free. Hence, this improves the existing stability theorems for projective modules over such graded algebras of dimension 2. This exposition is borrowed from [37].

In 1987, Lindel [24, Theorem 1.3] improved Quillen's Local-Global Principle from polynomial rings to positively graded rings. This, in particular, initiated the study of projective modules over a higher dimensional graded ring from a more algebraic point of view. Keeping track of this theme, in this article we aim to extend existing stability theorems for projective modules such as [31] from polynomial rings to an arbitrary positively graded domain of dimension  $\geq 2$ .

For the remainder of the Introduction, let us assume that (1) R is a commutative Noetherian ring with  $1 \neq 0$  of finite (Krull) dimension  $m \geq 2$ , and (2)  $A = \bigoplus_{i \geq 0} R_i$  is a commutative Noetherian (non-trivially) graded domain of dimension  $d \geq 2$  such that  $\dim(R_0) \geq 1$ .

1.1. Unimodular rows. Recall that a row vector  $v=(v_1,\ldots,v_n)\in R^n$  is called a unimodular row if  $\sum_{i=1}^n \lambda_i v_i=1$  for some  $\lambda_i\in R$ . The set of all unimodular rows in R of length n is denoted by  $\mathrm{Um}_n(R)$ . For all  $n\geq m+2$ , elementary facts (such as

Date: May 8, 2024.

<sup>2020</sup> Mathematics Subject Classification. 19A13, 13C10, 19B10, 19A15, 13A02.

Key words and phrases. Quillen patching, generalized dimension function, cancellation, efficient generation, unimodular element, monoid ring.

the prime avoidance lemma) can be employed to demonstrate that any unimodular row of length n can be completed to the first row of an invertible matrix, specifically an elementary matrix. Previous literature includes well-known examples that establish the optimality of this bound in general. However, certain classes of rings, such as affine algebras over an algebraically closed field of characteristic 0, have the potential to improve this bound [39]. Nonetheless, the complete classification of all Noetherian rings of dimension m over which all unimodular rows of length m+1 can be completed to the first row of an invertible (simultaneously elementary) matrix remains a formidable task until now. In this context, we prove the following theorem [for the proof we refer to Theorem 3.5 and Corollary 3.8].

**Theorem 1.1.** Let  $C = S^{-1}A$ , where  $S \subset A$  is a multiplicative set. Any unimodular row in C of length d+1 can be completed to the first row of an invertible matrix  $\alpha$ . Moreover, the matrix  $\alpha$  can be chosen in such a way that there exists  $\theta(T) \in GL_{d+1}(C[T])$  such that  $\theta(0) = Id$  and  $\theta(1) = \alpha$  (in this case  $\alpha$  will be called homotopic to the identity matrix). Consequently, any stably free C-module of rank d is free.

Since our assertion here is about graded rings, it is uncertain whether the wellknown Suslin's factorial row reduction technique, as employed in [39], can be applied. One possible approach is to first extend Roitman's degree reduction techniques [34] from polynomial rings to positively graded rings, then use it to apply the factorial row technique in graded rings. But such an improvement of Roitman's degree reduction techniques is a challenging task in itself. Instead, we have taken an alternative path. Let us choose  $v \in Um_{d+1}(A)$ . First, we point out a beautiful observation due to Plumstead. In [31, Example 2] he showed that one can glue finitely many generalized dimension functions on a suitable finite-cover of Spec(R) simply by taking their maximum. Utilizing this observation, we are able to apply stability theorem due to Eisenbud and Evans [16] to establish the completion of v in a suitable two-cover of Spec(A). Then leveraging the grading and applying a well-known homotopy trick due to Swan and Weibel, we generalize Quillen's splitting lemma [32, Theorem 1] over graded rings [see Proposition 3.2]. In particular, we have demonstrated that in the case of graded rings, the splitting can be obtained in such a way that both matrices fix the canonical vector  $e_1 (= (1, 0, ..., 0))$ . This allows us to patch two elementary matrices such that the resulting matrix simultaneously (1) becomes homotopic to the identity matrix, and (2) transforms the unimodular row v to  $e_1$ . One may wonder whether it is possible to improve Quillen's splitting lemma for an arbitrary ring. Unfortunately, as we mention in Remark 3.4, it is not feasible. In another sense, this setback highlights the significance of maintaining the grading in the patching diagram used in this article. By employing a similar line of reasoning and utilizing a Local-Global Principle, we establish in Theorem 3.10 that the injective stability of  $K_1(A)$  is d+1, thereby improving upon Vaseršteĭn's stability bound [40] over graded domains.

1.2. **On a question of Lindel.** We now shift our focus to another classical problem in the study of projective modules. Let P be a projective R-module with trivial determinant. Recall that P has a unimodular element if and only if it splits off a free summand of rank one. Due to a classical result of Serre [35] it is known that if  $\operatorname{rank}(P) > d$ , then P splits off a free summand of rank one. Again this bound is the best possible. However,

certain classes of rings, such as R[X], can improve this bound [31]. Hence, naturally identifying the class of rings over which top rank projective modules split into a free factor becomes the next big challenge.

For the remainder of the Introduction, let us assume that P is a projective A-module with trivial determinant such that  $\operatorname{rank}(P)=d$ . We recall that the Quillen set of P, denoted by  $J(R_0,P)$ , is defined as the collection of all elements  $s\in R_0$  such that  $P_s\cong \frac{P_s}{(PR_+)_s}\otimes R_s$ . It has been proven in [24, Theorem 1.3] that  $J(R_0,P)$  forms an ideal in  $R_0$ . Lindel gave a sufficient condition in the same article (see Theorem 2.5 there) for the existence of a unimodular element in P over certain graded rings. Informally, Lindel's theorem states that if (1)  $J(R_0,P)\neq 0$  and (2) the graded ring A satisfies some special criteria (e.g.  $\dim(A)-\dim(R_0)=1$ , when in addition A is an affine domain over a field), then P splits off a free summand of rank one. In [24, Remark 2.6], Lindel queries whether the second hypothesis in his theorem is necessary. We show in [24, Theorem 2.5] that both the hypotheses (1) and (2) are redundant [see Corollary 4.5]. In fact, we establish an even stronger result. Before stating our theorem, we digress momentarily.

Since 1980, a recurrent theme in this area has been to find the precise obstruction for a projective R-module of rank m to split a free factor. Recall that an ideal  $J \subset R$  is considered efficiently generated if  $\mu(J/J^2) = \mu(J)$ , where  $\mu(-)$  represents the minimal number of generators. Thanks to a significant result by Eisenbud and Evans [16], one can always find an A-linear map  $\gamma: P \to A$  such that the image ideal  $I(=\gamma(P))$  satisfies  $\mu(I/I^2) = \operatorname{ht}(I) = d$ . Notable contributions in this area have been made by seminal works such as [29] and [11] which eventually established the fact that the existence of a lift to a specific set of generators of  $I/I^2$  dictates the splitting problem for P. This provides an additional approach to tackle the splitting problem for projective modules of top rank.

In this context, we provide a sufficient criterion [see Proposition 4.2] for the efficient generation of a top height ideal in an arbitrary ring R. This empowers us to address the splitting problem for projective modules of top rank as well as the problem of efficient generation of top height ideals across various scenarios. In particular, we prove the following [for the proof we refer to Theorem 4.3, 4.7 and Corollary 4.5].

## **Theorem 1.2.** Let C and n be one of the following:

- (1) C = A and  $n = \dim(A) = d \ge 2$ .
- (2)  $C = S^{-1}A$ , where  $S \subset A$  is a multiplicative set contained in the set of all non-zero divisors in A such that  $\dim(C) = \dim(A)$  and  $n = \dim(A) = d \ge 3$ .
- (3) C = B[M], where B is a commutative Noetherian ring of dimension  $\geq 2$  and M is a finitely generated commutative cancellative (not necessarily torsion free) monoid of rank  $r \geq 1$ . We take  $n = \dim(B[M])$ .

Let  $I \subset C$  be an ideal such that  $\mu(I/I^2) = ht(I) = n$ . Then any set of generators of  $I = \langle f_1, \ldots, f_n \rangle + I^2$  lifts to a set of generators of I. Consequently, any projective C-module of rank n with trivial determinant splits off a free summand of rank one.

Readers may question the significance of the hypothesis  $\dim(R_0) \ge 1$  in this article. In Example 4.4 and 4.6 we demonstrate that this hypothesis is indeed necessary in Theorem 1.2.

- 1.3. Cancellation of projective modules of "top" rank. Recall that a projective R-module Q is considered cancellative if  $Q \oplus R^k \cong Q' \oplus R^k$  implies  $Q \cong Q'$ , where Q' is another R-module and  $k \geq 1$ . In Section 5, we explore the cancellation problem for projective modules over A. In Theorem 5.2 we establish that if  $J(R_0, P) \neq 0$ , then P is cancellative. As an interesting consequence, we demonstrate in Corollary 5.5 that over a Noetherian graded subring B of R[T] containing R with  $\dim(B) = m + 1$ , projective modules of rank m + 1 with trivial determinant are cancellative.
- 1.4. **On a question of Nori over a graded non-smooth algebra.** In section 6, we deduce some consequences of Theorem 1.2. We very briefly recall an algebraic analogy of a question asked by Nori [25].

**Question** 1.3. Let C be a smooth affine domain of dimension d over an infinite perfect field. Let  $I \subset C[T]$  be an ideal of height n such that  $I = \langle f_1, \dots, f_n \rangle + I^2T$ , where  $2n \geq d+3$ . Do there exist  $g_i \in I$  such that  $I = \langle g_1, \dots, g_n \rangle$  with  $g_i - f_i \in I^2T$ ?

This question is completely solved in [8] and [6]. Bhatwadekar, Mohan Kumar and Srinivas constructed an example [8, Example 6.4] of a non-smooth positively graded affine domain (with the degree zero subring  $\mathbb C$ ) such that over which Nori's question has a negative answer. However, when the ring has singularities, it is shown in [3] that imposing some suitable smoothness condition on the ideal  $I \cap R$  one can prevent such anomalies. Here, in Section 6, we aim to understand the underlying issue that prevents the existence of such a lift in [8, Example 6.4]. In particular, we prove the following [for details we refer to Theorem 6.1 and Corollary 6.3].

**Theorem 1.4.** Let  $A = \bigoplus_{i \geq 0} R_i$  be an affine domain (non-necessarily smooth) of dimension  $d \geq 3$  over an infinite field such that  $\frac{1}{d!} \in A$  and  $\dim(R_0) \geq 1$ . Let  $I \subset A[T]$  an ideal such that  $\mu(I/I^2T) = ht(I) = d$ . Then any set of generators of  $I/I^2T$  lifts to a set of generators of I. Consequently, any projective A[T]-module (with trivial determinant) of rank d splits off a free summand of rank one.

**Convention.** The symbol  $\mathbb N$  denotes the set of all non-negative integers, including 0. All rings considered in this article are assumed to be commutative Noetherian with  $1(\neq 0)$  having finite (Krull) dimension. Additionally, all graded rings discussed in this article are assumed to have a non-trivial  $\mathbb N$ -grading. For a graded ring  $R=\bigoplus_{i\geq 0}R_i$ , we use the notation  $R_+=\bigoplus_{i\geq 1}R_i$  to represent the irrelevant ideal in R containing all elements which can be written as a sum of homogeneous elements of degree >0. Every module considered in this article is assumed to be finitely generated.

#### 2. Preliminaries

This section summarizes several results and definitions from the literature that are frequently used in this article to prove the main theorems. We may restate or improve these results as necessary. Before proceeding further, we recall several definitions from the literature.

## **Definition** 2.1. Let *A* be a ring.

(1) Let M be an A-module. An element  $x \in M$  is said to be a *basic element* of M at a prime ideal  $\mathfrak{p} \in \operatorname{Spec}(A)$  if  $x \notin \mathfrak{p}M_{\mathfrak{p}}$ . For any  $S \subset \operatorname{Spec}(A)$ , we call x a basic element of M on S if it is a basic element of M at each prime ideal  $\mathfrak{p} \in S$ .

- (2) Let  $S \subset \operatorname{Spec}(A)$  and let  $\delta : S \to \mathbb{N}$  be a function. For two prime ideals  $\mathfrak{p}, \mathfrak{q} \in S$ , we define a partial order  $\mathfrak{p} << \mathfrak{q}$  if and only if  $\mathfrak{p} \subset \mathfrak{q}$  and  $\delta(\mathfrak{p}) > \delta(\mathfrak{q})$ . We say that  $\delta$  is a *generalized dimension function* if for any ideal  $I \subset A$ , the set  $V(I) \cap S$  has only finitely many minimal elements with respect to <<.
- (3) A positive integer r is said to be the *stable rank* of A, denoted as sr(A), if r is the smallest integer for which any  $(a_1, \ldots, a_{r+1}) \in Um_{r+1}(A)$ , there exists  $\lambda_i \in R$ ,  $i = 1, \ldots, r$  such that  $(a_1 + \lambda_1 a_{r+1}, \ldots, a_r + \lambda_r a_{r+1}) \in Um_r(R)$ .
- (4) We define  $H_n(A) = \{ \alpha \in GL_n(A) : \text{ there exists a } \theta(T) \in GL_n(A[T]) \text{ such that } \theta(0) = \text{Id and } \theta(1) = \alpha \}.$  Then  $H_n(A)$  is a normal subgroup of  $GL_n(A)$ .
- (5) Let  $I \subset A$  be an ideal and let "bar" denote going modulo I. We define  $\mathrm{Um}_n(A,I) := \{v \in \mathrm{Um}_n(A) : \overline{v} = e_1\}$  and  $\mathrm{GL}_n(A,I) := \{\alpha \in \mathrm{GL}_n(A) : \overline{\alpha} = \mathrm{Id}\}.$
- (6) Let P be a projective A-module such that P has a unimodular element. We choose  $\phi \in P^*$  and  $p \in P$  such that  $\phi(p) = 0$ . We define an endomorphism  $\phi_p$  as the composite  $\phi_p : P \to A \to P$ , where  $A \to P$  is the map sending  $1 \to p$ . Then by a *transvection* we mean an automorphism of P, of the form  $1 + \phi_p$ , where either  $\phi \in \operatorname{Um}(P^*)$  or  $p \in \operatorname{Um}(P)$ . By  $\operatorname{E}(P)$  we denote the subgroup of  $\operatorname{Aut}(P)$  generated by all transvections.

We begin by considering the following observation for a graded domain. This simple proposition plays a crucial role in the article, and therefore, we provide the proof.

**Proposition 2.2.** Let  $R = \bigoplus_{i \geq 0} R_i$  be a graded domain of dimension d. Let  $S \subset R_0$  be a multiplicative set such that  $S \cap \mathfrak{m} \neq \emptyset$ , for any maximal ideal  $\mathfrak{m} \in Spec(R_0)$ . Then the graded domain  $S^{-1}R$  does not have a graded maximal ideal  $S^{-1}\mathfrak{M}$  such that  $\mathfrak{M}$  is a maximal ideal in R. As a consequence, we get  $\dim(S^{-1}R) < d$ .

Proof. Suppose that  $\dim(R_0) = n$ . We give the proof by induction on n. First, we note that if n = 0, then  $R_0$  is a field. In this case, the statement is vacuously true.

Now, we consider the case where  $n \geq 1$ . If there does not exist such an S, then the statement is vacuously true. Therefore, we assume that such an S exists. Contrarily, we assume the existence of a graded maximal ideal, denoted as  $S^{-1}\mathfrak{M}$ , in  $S^{-1}R$ , where  $\mathfrak{M} \in \operatorname{Spec}(R)$  is a maximal ideal. There are two possibilities: either  $\mathfrak{M}$  is a graded maximal ideal or  $\mathfrak{M}$  is a maximal ideal but not a graded ideal. If  $\mathfrak{M}$  is a graded maximal ideal, then it can be expressed as  $\mathfrak{m} \oplus R_+$ , where  $\mathfrak{m}$  is a maximal ideal in  $R_0$ . Since  $S \cap \mathfrak{m} \neq \emptyset$ , it implies that  $S \cap \mathfrak{M} \neq \emptyset$ . However, this leads to a contradiction.

Now, we assume that  $\mathfrak{M}$  is not a graded ideal. Since  $S^{-1}\mathfrak{M}$  is a graded maximal ideal, it can be expressed as  $\mathfrak{m}' \oplus S^{-1}R_+$ , where  $\mathfrak{m}'$  is a maximal ideal in  $S^{-1}R_0$ . In particular, as R is Noetherian, there exists  $s \in S$  such that  $sR_+ \subset \mathfrak{M}$ . Because of  $S \cap \mathfrak{M} = \emptyset$ , it follows that  $R_+ \subset \mathfrak{M}$ .

We claim that  $\mathfrak{m}_0 := \mathfrak{M} \cap R_0 \neq 0$  is a non-zero prime ideal in  $R_0$ . We observe that, it is enough to show that  $\mathfrak{m}_0 \neq 0$ . Contrary, let us assume that  $\mathfrak{m}_0 = 0$ . Consider an element  $f \in \mathfrak{M}$ . We can write  $f = f_0 + f_1$ , where  $f_0 \in R_0$  and  $f_1 \in R_+$ . Since  $f_1 \in R_+ \subset \mathfrak{M}$ , we have  $f_0 \in \mathfrak{m}_0 = 0$ . This implies  $\mathfrak{M} = R_+$ . As a result, we get  $R_0 \cong R/R_+ \cong R/\mathfrak{M} \cong k$ , where k is a field. However, this is not possible as  $\dim(R_0) = n \geq 1$ . Hence, we establish that  $\mathfrak{m}_0 \neq 0$ .

Let "bar" denote going modulo  $\mathfrak{m}_0$ . Note that if  $\mathfrak{m}_0 \cap S \neq \emptyset$ , then we have  $S \cap \mathfrak{M} \neq \emptyset$ , which contradicts our assumption on the existence of such a maximal ideal. Hence,

without loss of generality, we may assume that  $\mathfrak{m}_0 \cap S = \emptyset$ . Since  $\mathfrak{m}_0 \neq 0$ , we have  $\dim(\overline{R_0}) \leq n - 1$ , and  $\overline{R}$  is a graded domain with dimension  $\leq d - 1$ .

Let  $\eta \in \operatorname{Spec}(\overline{R_0})$  be a maximal ideal in  $\overline{R_0}$ . Since  $\eta + \mathfrak{m}_0$  is also a maximal ideal in  $R_0$ , according to our induction hypothesis, we have  $S \cap \langle \eta + \mathfrak{m}_0 \rangle \neq \emptyset$ . This implies that  $\overline{S} \cap \eta \neq \emptyset$ . Furthermore, we observe that  $\overline{\mathfrak{M}}$  is a maximal ideal in  $\overline{R}$ . Moreover, as  $\mathfrak{m}_0 \subset R_0$  and  $\overline{S}^{-1}\overline{\mathfrak{M}} \cong \overline{S^{-1}}\overline{\mathfrak{M}}$ , the ideal  $\overline{S}^{-1}\overline{\mathfrak{M}}$  is a graded maximal ideal in  $\overline{S}^{-1}\overline{R}$  (recall that  $S^{-1}\mathfrak{M}$  is a graded ideal). However, by the induction hypothesis, there does not exist such a maximal ideal in  $\overline{S}^{-1}\overline{R}$ . This completes the induction step.

It remains to show that  $\dim(S^{-1}R) < d$ . To prove this, we note that for an arbitrary graded ring  $B = \bigoplus_{i \geq 0} B_i$ , there exists a graded maximal ideal  $\mathfrak N$  in B such that  $\operatorname{ht}(\mathfrak N) = \dim(B)$ . In  $S^{-1}R$ , any graded maximal ideal of height d is a localization of a maximal ideal in R. However, we have already demonstrated the nonexistence of such a graded maximal ideal in  $S^{-1}R$ . Therefore, the ring  $S^{-1}R$  does not have a graded maximal ideal of height d. Consequently, we obtain that  $\dim(S^{-1}R) < d$ .

We revisit a well-known homotopy map due to Swan and Weibel.

**Definition** 2.3. Let  $R = \bigoplus_{i \geq 0} R_i$ . We define the Swan-Weibel's homotopy map  $\Gamma_{SW}$ :  $R \to R[T]$  as follows: for any element  $f = a_0 + a_1 + \ldots + a_n \in R$ , we define  $\Gamma_{SW}(f) := a_0 + a_1 T + \ldots + a_n T^n \in R[T]$ , where  $a_i \in R_i$ .

**Lemma 2.4.** Let  $R = \bigoplus_{i \geq 0} R_i$ . Let  $\alpha \in GL_n(R)$  such that  $\overline{\alpha} = Id$ , where "bar" denotes going modulo the ideal  $R_+$ . Then there exists an  $\theta(T) \in GL_n(R[T])$  such that  $\theta(0) = Id$  and  $\theta(1) = \alpha$ . In other words  $\alpha \in H_n(R)$ . Moreover, if  $e_1\alpha = e_1$ , then we may choose such an  $\theta(T)$  with the property that  $e_1\theta(T) = e_1$ .

Proof. Consider the group homomorphism  $\widetilde{\Gamma_{SW}}: \operatorname{GL}_n(R) \to \operatorname{GL}_n(R[T])$  induced by  $\Gamma_{SW}$  [5, Definition 2.1]. Let us take  $\theta(T) = \widetilde{\Gamma_{SW}}(\alpha) \in \operatorname{GL}_n(R[T])$ . Then it follows that  $\theta(0) = \overline{\alpha} = \operatorname{Id}$  and  $\theta(1) = \alpha$ . Now we assume that  $e_1\alpha = e_1$ . As  $1 \in R_0$ , we have  $\Gamma_{SW}(1) = 1$ . Hence, we have  $e_1\theta(T) = e_1$ .

**Lemma 2.5.** Let  $R = \bigoplus_{i \geq 0} R_i$ . Then the map  $\Gamma_{SW} : R \to R[T]$  will induce a group homomorphism  $\widetilde{\Gamma_{SW}} : E_n(R) \to E_n(R[T])$ .

Proof. First, we observe that  $\Gamma_{SW}$  will induce a group homomorphism  $\widetilde{\Gamma_{SW}}: E_n(R) \to \operatorname{GL}_n(R[T])$  for details we refer to [5, Definition 2.1]. Therefore, it is enough to show that  $\widetilde{\Gamma_{SW}}(E_n(R)) \subset E_n(R[T])$ . Let  $E_{ij}(f) \in E_n(R)$  be an elementary matrix whose only non-zero non-diagonal entry is f at the position (i,j), where  $i \neq j$ . We write  $f(T) = \Gamma_{SW}(f)$ . Then we note that  $\widetilde{\Gamma_{SW}}(E_{ij}(f)) = E_{ij}(f(T)) \in E_n(R[T])$ . Moreover, since  $\widetilde{\Gamma_{SW}}$  is a group homomorphism and any element of  $E_n(R)$  can be written as a finite product of elements of the form  $E_{ij}(f)$ , it follows that  $\widetilde{\Gamma_{SW}}(E_n(R)) \subset E_n(R[T])$ .

The next lemma is known as one of the variants of Quillen-Suslin's Local-Global Principle and must be well-known. However, we could not find any suitable reference for the exact version required in this article. The closest reference we have found is [5, Theorem 3.8]. Therefore, we provide the proof, which is straightforward using the homotopy map  $\Gamma_{SW}$  and Suslin's Local-Global Principle [38, Lemma 3.5].

**Lemma 2.6.** Let  $R = \bigoplus_{i \geq 0} R_i$  and let "bar" denote going modulo the ideal  $R_+$ . Let  $s, t \in R_0$  be two co-maximal elements and let  $\alpha \in GL_n(R)$  such that (i)  $\overline{\alpha} = Id$ , (ii)  $\alpha_s \in E_n(R_s)$  and (iii)  $\alpha_t \in E_n(R_t)$ , where  $n \geq 3$ . Then  $\alpha \in E_n(R)$ .

Proof. We take  $\theta(T) = \widetilde{\Gamma_{SW}}(\alpha) \in \operatorname{GL}_n(R[T])$ , where  $\widetilde{\Gamma_{SW}} : \operatorname{GL}_n(R) \to \operatorname{GL}_n(R[T])$  is induced by  $\Gamma_{SW}$ . Then we note that  $\theta(0) = \operatorname{Id}$ . Moreover, it follows from Lemma 2.5 that  $(\theta(T))_s \in \operatorname{E}_n(R_s[T])$  and  $(\theta(T))_t \in \operatorname{E}_n(R_t[T])$ . Applying [38, Lemma 3.5] we obtain that  $\theta(T) \in \operatorname{E}_n(R[T])$ . Therefore, we get  $\alpha = \theta(1) \in \operatorname{E}_n(R)$ .

The following lemma is due to Plumstead, which is an adaptation of [31, Example 4], tailored to our requirements. Here, we point out that the following version has an additional conclusion compared to the version given in [1, Lemma 2.4], and this conclusion is crucially used in Lemma 4.1. However, the same proof works here as well. For the sake of completeness we give the proof.

**Lemma 2.7.** Let A be a ring of dimension d, and let s be a non-zero divisor in A such that  $\dim(A_s) \leq d-1$ . Then there exists a generalized dimension function  $\delta : Spec(A) \to \mathbb{N}$  such that  $\delta(\mathfrak{p}) \leq d-1$  for all  $\mathfrak{p} \in Spec(A)$ . Furthermore, we can choose  $\delta$  such that  $\delta(\mathfrak{p}) = \dim(A/\mathfrak{p})$  for all  $\mathfrak{p} \ni s$ .

Proof. Let  $\mathfrak{p} \in \operatorname{Spec}(A)$ . Then we note that either  $s \in \mathfrak{p}$  or  $s \notin \mathfrak{p}$ . That is, the prime ideal  $\mathfrak{p} \in V(s) \cup \operatorname{Spec}(A_s)$ , where  $V(s) = \{\mathfrak{p} \in \operatorname{Spec}(A) : s \in \mathfrak{p}\}$ . Hence  $\operatorname{Spec}(A) = S_1 \cup S_2$ , where  $S_1 = \operatorname{Spec}(A_s)$  and  $S_2 = V(s)$ . We define two functions:

- (1)  $d_1: S_1 \to \mathbb{N}$  such that  $d_1(\mathfrak{p}) = \dim(A_s/\mathfrak{p})$  for all  $\mathfrak{p} \in S_1$ ;
- (2)  $d_2: S_2 \to \mathbb{N}$  such that  $d_2(\mathfrak{p}) = \dim(A/\mathfrak{p})$  for all  $\mathfrak{p} \in S_2$ .

Let  $I \subset A$  be an ideal. We observe that the equality  $V(I) \cap S_1 = V(I_s)$ . Since A is a Noetherian ring, it follows from the primary decomposition of  $I_s$  that  $V(I) \cap S_1$  has finitely many minimal elements with respect to the partial ordering  $<<_1$  induced by  $d_1$  as defined in Definition 2.1. On the other hand  $V(I) \cap S_2 = V(I) \cap V(s) = V(I + \langle s \rangle)$ . Again, from the primary decomposition of  $I + \langle s \rangle$ , it follows that  $V(I) \cap S_2$  also has finitely many minimal elements with respect to  $<<_2$ , induced by  $d_2$ . Therefore, both  $d_1$  and  $d_2$  are generalized dimension function on  $S_1$  and  $S_2$ , respectively.

Moreover, we notice that  $d_1(\mathfrak{p}) \leq \dim(A_s) \leq d-1$  for all  $\mathfrak{p} \in \operatorname{Spec}(A_s) = S_1$ . On the other hand, since s is a non-zero divisor, we obtain that  $d_2(\mathfrak{p}) \leq \dim(A) - 1 \leq d-1$  for all  $\mathfrak{p} \in S_2$ .

Following [31, Example 2] we can define a generalized dimension function  $\delta: \operatorname{Spec}(A) \to \mathbb{N}$  such that  $\delta(\mathfrak{p}) = d_1(\mathfrak{p})$  if  $\mathfrak{p} \in S_1$ , and  $\delta(\mathfrak{p}) = \dim(A/\mathfrak{p})$  if  $\mathfrak{p} \in S_2$ . Then we note that  $d(\mathfrak{p}) \leq d-1$  for all  $\mathfrak{p} \in \operatorname{Spec}(A)$  and  $\delta(\mathfrak{p}) = \dim(A/\mathfrak{p})$  for all  $\mathfrak{p} \ni s$ . This completes the proof.

We end this section with the next theorem, which is derived from a pivotal result due to Eisenbud and Evans [16]. This has been used extensively throughout the article. This version is recollected from [31].

**Theorem 2.8.** Let A be a ring, and let  $\mathcal{P} \subset Spec(A)$  be a subset. Consider a generalized dimension function  $\delta: \mathcal{P} \to \mathbb{N}$ . Let M be an A-module satisfying  $\mu_{\mathfrak{p}}(M) \geq 1 + \delta(\mathfrak{p})$  for all  $\mathfrak{p} \in \mathcal{P}$ , where  $\mu_{\mathfrak{p}}(M)$  is the minimal number of generators of  $M_{\mathfrak{p}}$ . For a basic element  $(r,m) \in A \oplus M$  on  $\mathcal{P}$ , there exists an element  $m' \in M$  such that m + rm' is also a basic element on  $\mathcal{P}$ .

## 3. Unimodular rows

This section is devoted to establishing that any unimodular row of length d+1 over a graded domain of dimension  $d \ge 1$  can be completed to the first row of an invertible

matrix, which is homotopic to the identity matrix. We begin with an easy consequence of Lemma 2.7 and Theorem 2.8.

**Lemma 3.1.** Let A be a ring of dimension  $d \ge 1$ . Assume that, there exists a non-zero divisor  $s \in A$  such that  $\dim(A_s) < \dim(A)$ . Then  $sr(A) \le d$ .

Proof. Let  $v=(v_1,\ldots,v_{d+1})\in \mathrm{Um}_{d+1}(A)$ . Applying Lemma 2.7 we get a generalized dimension function  $\delta:\mathrm{Spec}(A)\to\mathbb{N}$  such that  $\delta(\mathfrak{p})\leq d-1$  for all  $\mathfrak{p}\in\mathrm{Spec}(A)$ . We note that v is a basic element of the free module  $A^{d+1}$ . Then applying Theorem 2.7 (taking  $M=A^d$ ) we obtain a basic element  $w=(v_1+\lambda_1v_{d+1},\ldots,v_d+\lambda_dv_{d+1})$  of  $A^d$ , for some  $\lambda_i\in A$ . Now since  $A^d$  is a free (in particular, a projective) module, every basic element is a unimodular row. This concludes the proof.

The next proposition is similar to the well-known Quillen's splitting lemma [32, Theorem 1]. Here we rephrase it in our setup with an added conclusion, which is crucial for this article.

**Proposition 3.2.** Let  $R = \bigoplus_{i \geq 0} R_i$  and  $s, t \in R_0$  such that  $\langle s \rangle + \langle t \rangle = R_0$ . Let  $\eta \in GL_n(R_{st}, (R_+)_{st})$  such that  $e_1 \eta = e_1$ . Then there exist  $\eta_1 \in GL_n(R_s, (R_+)_s)$  and  $\eta_2 \in GL_n(R_t, (R_+)_t)$  such that

- (1)  $\eta = (\eta_1)_t (\eta_2)_s$ ,
- (2)  $e_1\eta_i = e_1$ , for i = 1, 2.

Proof. Let "bar" denote going modulo the ideal  $R_+$ . We define  $\chi(X):=\widetilde{\Gamma_{SW}}(\eta)$ . Using Lemma 2.4 we obtain

$$\chi(X) \in \operatorname{GL}_n(R_{st}[X])$$
 such that  $\chi(0) = \overline{\eta} = \operatorname{Id}$  and  $e_1\chi(X) = e_1$ .

Applying Quillen's splitting lemma [32, Theorem 1, paragraph 2], one can find  $g = s^N \in R_0$  with  $N \in \mathbb{N}$  such that

$$\chi(X)\chi(gX)^{-1} \in \operatorname{GL}_n(R_s[X])$$
 and  $\chi(gX) \in \operatorname{GL}_n(R_t[X])$ .

Since  $e_1\chi(X)=e_1$ , we further obtain that  $e_1\chi(gX)=e_1$  and  $e_1\chi(X)\chi(gX)^{-1}=e_1$ . Let us define  $\underline{\eta_1}:=\chi(1)\chi(g)^{-1}$  and  $\underline{\eta_2}:=\chi(g)$ . We observe the matrix  $\chi(X)$  has the property that  $\overline{\chi(a)}=\operatorname{Id}$  for any  $a\in R_0$ . This implies  $\overline{\eta}_i=\operatorname{Id}$ , for i=1,2. Therefore, we get the following.

- (1)  $\eta = (\eta_1)_t (\eta_2)_s$ ;
- (2)  $\eta_1 \in GL_n(R_s, (R_+)_s);$
- (3)  $\eta_2 \in GL_n(R_t, (R_+)_t);$
- (4)  $e_1 \eta_i = e_1$ , for i = 1, 2.

This concludes the proof.

The next lemma concerns the patching of two invertible matrices in a graded ring. One may compare this with [31, Lemma 2].

**Lemma 3.3.** Let  $R = \bigoplus_{i \geq 0} R_i$  and  $s, t \in R_0$  such that  $\langle s \rangle + \langle t \rangle = R_0$ . Let  $v \in Um_n(R, R_+)$ . Assume that, there exist  $\alpha_1 \in GL_n(R_s, (R_+)_s)$  and  $\alpha_2 \in GL_n(R_t, (R_+)_t)$  such that  $v\alpha_i = e_1$ , for i = 1, 2. Then there exists an  $\alpha \in GL_n(R, R_+)$  such that  $v\alpha = e_1$ .

Proof. Let "bar" denote going modulo  $R_+$ . Let us define  $\eta := (\alpha_1)_t^{-1}(\alpha_2)_s \in GL_n(R_{st})$ . Then we note that  $\overline{\eta} = \operatorname{Id}$  and  $e_1 \eta = e_1$ . Applying Proposition 3.2 there exist  $\eta_1 \in GL_n(R_s, (R_+)_s)$  and  $\eta_2 \in GL_n(R_t, (R_+)_t)$  such that

- (a)  $\eta = (\eta_1)_t (\eta_2)_s$ ,
- (b)  $e_1 \eta_i = e_1$ , for i = 1, 2.

We now define  $\sigma_1 := \alpha_1 \eta_1 \in \operatorname{GL}_n(R_s, (R_+)_s)$  and  $\sigma_2 := \alpha_2 \eta_2^{-1} \in \operatorname{GL}_n(R_t, (R_+)_t)$ . Here we notice that  $v\sigma_i = e_1$  (i = 1, 2). Because of  $\eta = (\alpha_1)_t^{-1}(\alpha_2)_s = (\eta_1)_t(\eta_2)_s$ , we have  $(\sigma_1)_t = (\sigma_2)_s$ . Therefore, by [23, Proposition 2.2, page no 211] there exists a unique  $\alpha \in \operatorname{GL}_n(R, R_+)$  such that  $\alpha_s = \sigma_1$  and  $\alpha_t = \sigma_2$ . Furthermore, the matrix  $\alpha$  takes v to  $e_1$  as it is true locally.

**Notation.** Let A be a ring and let  $G \subset GL_n(A)$  be a subgroup. For any  $u, v \in Um_n(A)$ , we define  $u \sim_G v$  if there exists an  $\epsilon \in G$  such that  $u\epsilon = v$ . We denote the set  $\{v \in Um_n(A) : v \sim_G e_1\}$  by the notation  $e_1G$ .

Remark 3.4. One may wonder whether it is possible to improve Quillen's splitting lemma (for an arbitrary ring) in such a way that both the splitting matrices fix the canonical vector  $e_1$ . Unfortunately, achieving such an improvement is not feasible. To illustrate this, here we argue as follows: consider a ring A of dimension  $d \geq 2$ . We show that such an improvement of Quillen's splitting lemma will ultimately lead to the conclusion that  $\mathrm{Um}_{d+1}(A) = e_1\mathrm{SL}_{d+1}(A)$ , which as discussed in the Introduction is not true. We choose a  $v \in \mathrm{Um}_{d+1}(A)$ . Then we can always find a non-zero divisor  $s \in R$  such that  $v\alpha_1 = e_1$ , for some  $\alpha_1 \in \mathrm{E}_{d+1}(A_s)$ . As  $s \in \mathrm{Jac}(A_{1+\langle s \rangle})$  is a non-zero divisor, it is not difficult to establish that  $v\alpha_2 = e_1$ , for some  $\alpha_2 \in \mathrm{E}_{d+1}(A_t)$  and  $t \in 1 + \langle s \rangle$ . Now, if the elementary matrix  $\eta = (\alpha_1)_t^{-1}(\alpha_2)_s$  splits in such a way that each of its splitting matrices fixes  $e_1$ , then applying the arguments given in Lemma 3.3 one can obtain an  $\alpha \in \mathrm{SL}_{d+1}(A)$  such that  $v\alpha = e_1$ .

**Theorem 3.5.** Let  $R = \bigoplus_{i \geq 0} R_i$  be a graded domain of dimension  $d \geq 2$  such that  $\dim(R_0) \geq 1$ . Then for any  $v \in Um_{d+1}(R, R_+)$  there exists an  $\alpha \in GL_{d+1}(R, R_+)$  such that  $v\alpha = e_1$ . As a consequence

$$Um_{d+1}(R) = e_1 H_{d+1}(R).$$

Proof. Let  $v \in \operatorname{Um}_{d+1}(R)$  and  $S = R_0 \setminus \{0\}$ . Let "bar" denote going modulo  $R_+$ . As R has a non-trivial grading the ideal  $R_+ \neq 0$ . In particular, we get  $\operatorname{ht}(R_+) \geq 1$ . From standard stability results (e.g. Prime avoidance lemma) there exists  $\overline{\kappa} \in \operatorname{E}_{d+1}(\overline{R})$  such that  $\overline{v\kappa} = \overline{e}_1$ . Since the canonical map  $\operatorname{E}_{d+1}(R) \to \operatorname{E}_{d+1}(\overline{R})$  is surjective, one can alter v by an elementary matrix and may further assume that  $\overline{v} = \overline{e}_1$ . Now if there exists an  $\alpha \in \operatorname{GL}_{d+1}(R, R_+)$  such that  $v\alpha = e_1$ , then it follows from Lemma 2.4 and the fact  $\operatorname{E}_{d+1}(R) \subset \operatorname{H}_{d+1}(R)$  that  $\operatorname{Um}_{d+1}(R) = e_1\operatorname{H}_{d+1}(R)$ . Hence, to prove the theorem it is enough to find such an  $\alpha$ .

Applying Proposition 2.2 we get  $\dim(S^{-1}R) \leq d-1$ . In particular, by standard stability theorems (e.g. one can use prime avoidance lemma) to obtain that  $\mathrm{sr}(S^{-1}R) \leq d$ . Therefore, we can find an  $s \in S$  such that  $v \sim_{\mathrm{E}_{d+1}(R_s)} e_1$ . Let  $\alpha_1 \in \mathrm{E}_{d+1}(R_s)$  be such that  $v\alpha_1 = e_1$ . Furthermore, we may replace  $\alpha_1$  by  $\alpha_1\overline{\alpha}_1^{-1}$  and assume that  $\overline{\alpha}_1 = \mathrm{Id}$ .

Let  $\mathfrak{T} = \{1 + sr : r \in R_0\}$  and  $B = \mathfrak{T}^{-1}R$ . Since  $\mathfrak{T} \subset R_0$ , the ring B is also a graded ring. Moreover, we note that  $s \in Jac(\mathfrak{T}^{-1}R_0)$ . Hence, applying Proposition 2.2

it follows that  $\dim(B_s) \leq d-1$ . Therefore, by Lemma 3.1 we get  $\mathrm{sr}(B) \leq d$ . Thus, there exists an  $\alpha_2 \in \mathrm{E}_{d+1}(B)$  such that  $v\alpha_2 = e_1$ . As again we may replace  $\alpha_2$  by  $\alpha_2\overline{\alpha}_2^{-1}$  and further assume that  $\overline{\alpha}_2 = \mathrm{Id}$ . We can find  $t \in \mathfrak{T}$  such that  $\alpha_2 \in \mathrm{E}_{d+1}(R_t)$ .

Now applying Lemma 3.3 we can find an  $\alpha \in GL_{d+1}(R, R_+)$  such that  $v\alpha = e_1$ . This completes the proof.

**Remark** 3.6. Let A be a regular ring of essentially finite type over a field. Then using [41, Theorem 3.3] it is not difficult to establish the fact that  $H_{n+1}(A) = E_{n+1}(A)$ , for all  $n \geq 2$ . Hence, in Theorem 3.5, additionally if we assume that R is a regular ring of essentially finite type over a field, then we get  $\operatorname{Um}_{d+1}(R) = e_1 E_{d+1}(R)$ . However, we do not know whether the regularity of R is actually necessary.

**Remark** 3.7. One can remove the hypothesis that  $\dim(R_0) \geq 1$  in Theorem 3.5 at the expense of the hypothesis that  $\frac{1}{d!} \in R$  by utilizing the Swan-Weibel's homotopy map and applying [33, Corollary 2.5]. In fact, the same yields the following: let  $R = \bigoplus_{i \geq 0} R_i$  be a graded ring of dimension d such that  $\frac{1}{d!} \in R$ . Then  $\operatorname{Um}_{d+1}(R) = e_1 \operatorname{H}_{d+1}(R)$ . It is worth noting that the removal of the hypothesis " $\frac{1}{d!} \in R$ " from a cancellation problem is highly non-trivial (cf. [39] and [17]).

In the remaining part of the section, we extend Theorem 3.5 over various rings, which are closely related to the class of graded rings considered in Theorem 3.5.

**Corollary 3.8.** Let R and d be as in Theorem 3.5 and let  $A = \mathfrak{S}^{-1}R$ , where  $\mathfrak{S} \subset R$  is a multiplicative set. Then

$$Um_{d+1}(A) = e_1 H_{d+1}(A).$$

Proof. First we comment that, since  $\mathrm{E}_{d+1}(A)\subset \mathrm{H}_{d+1}(A)$ , the only non-trivial case is when  $\dim(A)=d$ . Hence, without loss of generality, we assume that  $\dim(A)=d$ . Let us choose  $v\in\mathrm{Um}_{d+1}(A)$ . Then there exists a non-zero  $x\in R$  such that (i)  $v\in\mathrm{Um}_{d+1}(R_x)$  and (ii)  $\dim(R_x)=d$ . Now it follows from [1, Lemma 4.4] that there exists  $u\in\mathrm{Um}_{d+1}(R)$  such that  $v\sim_{\mathrm{E}_{d+1}(R_x)}u$ . Applying Theorem 3.5 we obtain that  $u\sim_{\mathrm{H}_{d+1}(R)}e_1$ . Since  $\sim_{\mathrm{H}_{d+1}(R_x)}$  is transitive, the proof concludes.

**Corollary 3.9.** Let R and d be as in Theorem 3.5. Additionally, we assume that R is an affine domain over a field. Then

$$Um_{d+1}(R[X_1,\ldots,X_n]) = e_1 SL_{d+1}(R[X_1,\ldots,X_n]).$$

Proof. We use Quillen Induction on n to prove the theorem. For n=0 this follows from Theorem 3.5. Now let us assume that n>0. Let  $v\in \mathrm{Um}_{d+1}(R[X_1,\ldots,X_n])$ . We note that  $R[X_1]=\bigoplus_{i\geq 0}R_i[X_1]$ , where  $R_i[X_1]=\{\sum_{j=1}^n a_jX_1^j:a_j\in R_i \text{ and }n\in\mathbb{N}\}$  and the sum is defined in the obvious way. As there exists a canonical surjection  $R\to R_0$ , the ring  $R_0$  is also an affine domain over the same field, say k. Let us take  $S=k[X_1]\setminus\{0\}\subset R_0[X_1]$ . Therefore, we have  $\dim(S^{-1}R_0[X_1])=\dim(R_0)$ . As any maximal ideal m of  $R_0[X_1]$  is of height equal to  $\dim(R_0)+1$ , we have  $S\cap m\neq\emptyset$ . We take  $B=S^{-1}R[X_1]$ . Then it follows from Proposition 2.2 that  $\dim(B)\leq d$ . In particular, since  $\dim(S^{-1}R_0[X_1])=\dim(R_0)$  we have  $\dim(B)=d$ . Then  $B=\bigoplus_{i\geq 0}S^{-1}R_i[X_1](=\bigoplus_{i\geq 0}B_i$  say) is also a graded affine domain over the field  $k(X_1)$  of dimension d such that  $\dim(B_0)=\dim(R_0)\geq 1$ . Applying induction hypothesis on  $B[X_2,\ldots,X_n]$  we can find a monic polynomial  $f\in S$  such that

where  $D = R[X_2, ..., X_n]$ . Then by Affine Horrocks Theorem [32, Theorem 3] the result follows.

We end this section with a theorem on the injective stability of  $K_1(R)$ , where R is a graded domain. The proof is a straightforward consequence of Lemma 2.6 and 3.1.

**Theorem 3.10.** Let R and d be as in Theorem 3.5. Additionally, for  $n \ge 1$  we further assume that R is an affine domain over a field. Then

$$SL_{d+1}(R[X_1,\ldots,X_n]) \cap E(R[X_1,\ldots,X_n]) = E_{d+1}(R[X_1,\ldots,X_n]).$$

Proof. We again apply Quillen Induction on n to prove the theorem. We give the proof in cases.

Case - 1. Let us assume that n=0. Let  $\alpha\in \mathrm{SL}_{d+2}(R)\cap \mathrm{E}(R)$  and let "bar" denote going modulo the ideal  $R_+$ . Since  $\overline{\alpha}\in \mathrm{SL}_{d+1}(R_0)$  and  $R_0\subset R$ , we may treat  $\overline{\alpha}$  as an element of  $\mathrm{SL}_{d+1}(R)$ . Moreover, we observe that as  $\mathrm{ht}(R_+)\geq 1$  we have  $\dim(R_0)<\dim(R)$ . Therefore, using standard stability theorem due to Vaseršteĭn, as stated in [40, Theorem 3.2], we obtain that (1)  $\alpha\in \mathrm{SL}_{d+1}(R)\cap \mathrm{E}_{d+2}(R)$  and (2)  $\overline{\alpha}\in \mathrm{E}_{d+1}(R_0)\subset \mathrm{E}_{d+1}(R)$ . We take  $\beta=\alpha\overline{\alpha}^{-1}$ . Then we note that  $\beta\in \mathrm{SL}_{d+1}(R)\cap \mathrm{E}_{d+2}(R)$ . Let us consider  $\mathfrak{T}=R_0\setminus\{0\}$ . Then applying Proposition 2.2 we get  $\dim(\mathfrak{T}^{-1}R)\leq d-1$ . Hence, again using [40, Theorem 3.2] on  $\mathfrak{T}^{-1}R$  we obtain that  $(\beta)_{\mathfrak{T}}\in \mathrm{E}_{d+1}(\mathfrak{T}^{-1}R)$ . There exists an  $s\in\mathfrak{T}$  such that  $\beta_s\in \mathrm{E}_{d+1}(R_s)$ .

Let  $S = \{1 + sr : r \in R_0\}$  and let  $B = S^{-1}R$ . Then again by Proposition 2.2 we obtain that  $\dim(B_s) \leq d - 1$ . Hence, using Lemma 3.1 it follows that  $\mathrm{sr}(B) \leq d$ . We again apply [40, Theorem 3.2] to obtain that  $(\beta)_S \in \mathrm{E}_{d+1}(B)$ . We choose  $t \in S$  such that  $\beta_t \in \mathrm{E}_{d+1}(R_t)$ . Now it follows from Lemma 2.6 that  $\beta \in \mathrm{E}_{d+1}(R)$ . Because of  $\overline{\alpha} \in \mathrm{E}_{d+1}(R)$ , we have  $\alpha \in \mathrm{E}_{d+1}(R)$ .

**Case - 2.** Now let us assume that n > 0. Then applying Quillen Induction on n as described in Corollary 3.9 and using [38, Corollary 5.7] suitably one may conclude the proof.

#### 4. EFFICIENT GENERATION OF IDEALS

This section is devoted to studying the efficient generation problem for top height ideals in a ring. Before presenting the main theorem of this section, we need some preparation. We begin this section with the following lemma, which is a consequence of Lemma 2.7 and Theorem 2.8.

**Lemma 4.1.** Let A be a ring of dimension  $d \ge 2$ . Assume that, there exists a non-zero divisor  $s \in A$  such that  $\dim(A_s) < d$ . Let  $I \subset A$  be an ideal such that  $\mu(I/I^2) = ht(I) = d$ . Then any set of generators of  $I = \langle f_1, \dots, f_d \rangle + I^2$  lifts to a set of generators of I.

Proof. Applying Lemma 2.7 one may obtain a generalized dimension function  $\delta: \operatorname{Spec}(A) \to \mathbb{N}$  such that  $\delta(\mathfrak{q}) \leq d-1$  for all  $\mathfrak{q} \in \operatorname{Spec}(A)$  and  $\delta(\mathfrak{q}) = \dim(A/\mathfrak{q})$  for all  $\mathfrak{q} \ni s$ . Let  $\mathfrak{p} \in \operatorname{Spec}(A)$ . Suppose that, we have  $I \subset \mathfrak{p}$ . As  $\operatorname{ht}(I) = d$  we must have  $\operatorname{ht}(\mathfrak{p}) = d$ . Since  $\dim(A_s) < d$ , the element s is in  $\mathfrak{p}$ . This implies  $\delta(\mathfrak{p}) = \dim(A/\mathfrak{p}) = 0$ . As  $A_{\mathfrak{p}}$  is a local ring we have  $\mu(IA_{\mathfrak{p}}/I^2A_{\mathfrak{p}}) = \mu(IA_{\mathfrak{p}}) = d$ . Therefore, we obtain that  $\mu(IA_{\mathfrak{p}}) + \delta(\mathfrak{p}) \leq d$ . Now if  $I \not\subset \mathfrak{p}$ , then  $\mu(IA_{\mathfrak{p}}) = 1$ . Thus also in this case we have  $\mu(IA_{\mathfrak{p}}) + \delta(\mathfrak{p}) \leq d$ . In particular, we get  $\sup\{\mu(IA_{\mathfrak{p}}) + \delta(\mathfrak{p}) : \mathfrak{p} \in \operatorname{Spec}(A)\} \leq d$ . Hence,

one may apply [31, Theorem 0] to find  $e_i \in I^2$  (i = 1, ..., d) such that  $I = \langle l_1, ..., l_d \rangle$ , where  $l_i = f_i + e_i$ . This completes the proof.

In the following proposition, we present a general criterion for the efficient generation of a top height ideal in an arbitrary ring. This criterion enables us to identify the essential requirements to apply Mohan Kumar's fundamental technique presented in [27] to solve the efficient generation problem. By doing so, we are able to provide a unified approach in Theorem 4.3.

**Proposition 4.2.** Let A be a ring of dimension  $d \geq 2$ . Let  $I \subset A$  be an ideal such that  $\mu(I/I^2) = ht(I) = d$ . Suppose that  $I = \langle f_1, \dots, f_d \rangle + I^2$ . Moreover, we assume that there exists a non-zero divisor  $s \in A$  and a multiplicative set  $S \subset \{1 + sr : r \in A\}$  such that the following hold.

- (1)  $IA_s = \langle f_1, \dots, f_d \rangle A_s + I^2 A_s$ , has a lift to a set of generators of  $IA_s$ ,
- (2)  $\dim(S^{-1}A_s) < d$ , and
- (3)  $Um_d(S^{-1}A_s) = e_1H_d(S^{-1}A_s).$

Then there exist  $F_i \in A$  such that  $I = \langle F_1, \dots, F_d \rangle$ , with  $f_i - F_i \in I^2$ .

Proof. Let  $g_i \in IA_s$  be such that  $IA_s = \langle g_1, \dots, g_d \rangle A_s$  with  $f_i - g_i \in I^2A_s$ . We observe that, if  $\dim(A_s) < d$ , then applying Lemma 4.1 the proof follows. Hence, we assume that  $\dim(A_s) = d$ . Therefore, using ([11, Lemma 5.6] and [20, Remark 5.7]) we may further assume that  $s \in \sqrt{I}$ .

Let us take  $B = S^{-1}A$ . Since  $\dim(B_s) < d$ , using Lemma 4.1 we can lift  $f_i$ 's to a set of generators of IB. In particular, we get  $l_i \in IB$  such that  $IB = \langle l_1, \ldots, l_d \rangle$  and  $l_i - f_i \in I^2$ , for  $i = 1, \ldots, d$ .

Since  $s \in \sqrt{I}$ , the row vectors  $(g_1,\ldots,g_d)$  and  $(l_1,\ldots,l_d)$  are in  $\mathrm{Um}_d(B_s)$ . Hence, by hypothesis (3) there exists an  $\epsilon \in \mathrm{H}_d(B_s)$  such that  $(g_1,\ldots,g_d)\epsilon = (l_1,\ldots,l_d)$ . As  $\epsilon \in \mathrm{H}_d(B_s)$  there exists a  $\theta(T) \in \mathrm{GL}_n(B_s[T])$  such that  $\theta(0) = \mathrm{Id}$  and  $\theta(1) = \epsilon$ . Since A is a Noetherian ring and there are only finitely many  $g_i$  and  $l_i$ , we can find  $t \in S$  such that

- (1)  $IA_t = \langle l_1, \dots, l_d \rangle$  with  $f_i l_i \in I^2 A_t$ ;
- (2)  $\theta(T) \in GL_d(A_{st}[T])$ .

Applying Quillen's splitting lemma [32, Theorem 1] we obtain  $\epsilon_1 \in \operatorname{GL}_d(A_s)$  and  $\epsilon_2 \in \operatorname{GL}_d(A_t)$  such that  $\epsilon = (\epsilon_1)_t(\epsilon_2)_s$ . Because of  $\langle s \rangle + \langle t \rangle = A$ , one may apply a standard patching to obtain  $F_i \in I$  such that  $I = \langle F_1, \dots, F_d \rangle$  with  $F_i - f_i \in I^2$  for  $i = 1, \dots, d$ .  $\square$  Now we are ready to prove the main theorem of this section.

**Theorem 4.3.** Let R and d be as in Theorem 3.5. Let A and n be one of the following:

- (1) A = R and  $n = \dim(R) = d \ge 3$ .
- (2)  $A = \mathfrak{S}^{-1}R$ , where  $\mathfrak{S} \subset R$  is a multiplicative set contained in the set of all non-zero divisors in R such that  $\dim(A) = \dim(R)$  and  $n = \dim(R) = d \geq 3$ .
- (3) A = B[M], where B is a ring of dimension  $\geq 2$  and M is a finitely generated commutative cancellative (not necessarily torsion free) monoid of rank  $r \geq 1$ . We take  $n = \dim(B[M])$ .

Let  $I \subset A$  be an ideal such that  $\mu(I/I^2) = ht(I) = n$ . Then any set of generators of  $I = \langle f_1, \ldots, f_n \rangle + I^2$  lifts to a set of generators of I.

Proof. We will show that for each of the above rings all the hypotheses of Proposition 4.2 are satisfied. We handle these three rings separately in the following cases.

Case - 1. In this case, we assume that A=R. Let us take  $\mathfrak{T}=R_0\setminus\{0\}$ . Then by Proposition 2.2 the dimension of  $\mathfrak{T}^{-1}A$  is strictly smaller than n. Hence, applying [27] we can lift  $f_i$ 's to a set of generators of  $\mathfrak{T}^{-1}I$ . Therefore, there exist  $s\in\mathfrak{T}$  and  $g_i\in A_s$  with  $I_s=\langle g_1,\ldots,g_n\rangle$  such that  $f_i-g_i\in I_s^2$  for  $i=1,\ldots,n$ . Let  $S:=\{1+sr:r\in R_0\}\subset\{1+sx:x\in A\}$ . Then again applying Proposition 2.2 we have  $\dim(S^{-1}A_s)< n$ . Moreover, we observe that since  $S\subset R_0$  and  $s\in R_0$ , the ring  $S^{-1}R_s$  retains the grading induce from R. Because of  $n=d\geq 3$ , using Theorem 3.5 we have  $\mathrm{Um}_n(S^{-1}R_s)=e_1\mathrm{H}_n(S^{-1}R_s)$ . Therefore, applying Proposition 4.2 we obtain the required lift.

Case - 2. In this case we assume that  $A=\mathfrak{S}^{-1}R$ . Let  $\mathfrak{T}$  be as considered in Case - 1. Then as it was shown in the previous case that  $\dim(\mathfrak{T}^{-1}R)< n$ , which further implies that  $\dim(\mathfrak{T}^{-1}A)< n$ . Therefore, following the arguments in the previous case, we can find a non-zero divisor  $s\in R_0$  and  $g_i\in A_s$  such that  $I_s=\langle g_1,\ldots,g_n\rangle$ , with  $f_i-g_i\in I_s^2$ . Let us take  $S=\{1+sr:r\in R_0\}\subset\{1+sx:x\in A\}$ . Then  $S^{-1}A_s=\mathfrak{S}^{-1}(S^{-1}R_s)$ , where  $S^{-1}R_s$  is a positively graded ring of dimension  $\leq n-1$ . Hence, applying Corollary 3.8 we get  $\mathrm{Um}_n(S^{-1}A_s)=e_1\mathrm{H}_n(S^{-1}A_s)$ . Now one may apply Proposition 4.2 to complete the proof.

Case - 3. In this case we take A=B[M]. First we note that for a monoid ring B[M] we have  $\dim(B[M])=\dim(B)+\operatorname{rank}(M)$  [13, Theorem 4.23]. Let  $\mathfrak T$  be the set of all non-zero divisors in B. Then  $\dim(\mathfrak T^{-1}B[M])=r$ . Since  $n>\dim(\mathfrak T^{-1}B[M])$  by [27] we can lift  $f_i$ 's to a set of generators of  $\mathfrak T^{-1}I$ . Therefore, there exist  $s\in\mathfrak T$  and  $g_i\in B_s[M]$  with  $I_s=\langle g_1,\ldots,g_n\rangle$  such that  $f_i-g_i\in I_s^2$  for  $i=1,\ldots,n$ . Let  $S=\{1+sr:r\in B\}$  and let  $C=S^{-1}B$ . Then as  $\dim(C_s)<\dim(B)$  we have  $\dim(S^{-1}B_s[M])=\dim(C_s[M])< n$ . Therefore, using [18, Theorem 1.1] we get  $\operatorname{Um}_n(C_s[M])=e_1\operatorname{E}_n(C_s[M])$ . Now one may apply Proposition 4.2 to complete the proof.

We now provide an example that proves the necessity of the hypothesis  $\dim(R_0) > 0$  in Theorem 4.3 (1). We essentially use the example constructed by Bhatwadekar, Mohan Kumar and Srinivas [8, Example 6.4] in which they provided a non-smooth graded domain (with the degree zero subring a field) over which Nori's question has a negative answer.

**Example** 4.4. Consider the graded domain  $B = \frac{\mathbb{C}[X,Y,Z,W]}{\langle X^5+Y^5+Z^5+W^5\rangle}$ . By [8, Example 6.4] there exist (1) an ideal  $I \subset B[T]$  such that  $\mu(I/I^2T) = \operatorname{ht}(I) = 3$  and (2) a set of generators  $I = \langle f_1, f_2, f_3 \rangle + I^2T$ , which does not lift to a set of generators of I. Let  $S = \mathbb{C}[T] \setminus \{0\}$  and  $C = S^{-1}B[T]$ . Moreover, one may observe that I does not contain a monic polynomial in T. As if it did, then by [25, Theorem 2.1], one could lift  $I = \langle f_1, f_2, f_3 \rangle + I^2T$  to a set of generators for I. Then C is a graded domain of dimension 3 such that the degree zero subring of C is the field  $\mathbb{C}(T)$ , and IC is an ideal of C of height 3 such that  $IC = \langle f_1, f_2, f_3 \rangle C + I^2C$  does not lift to a set of generators of IC. As if such a lift exists then by [14, Theorem 3.10] one can lift  $I = \langle f_1, f_2, f_3 \rangle + I^2T$  to a set of generators of I, which is not true by (2).

The following is an interesting consequence of the previous theorem. For monoid rings, this is an improvement of [21, Theorem 3.4].

**Corollary 4.5.** Let A and n be as in Theorem 4.3. Let P be a projective A-module with trivial determinant of rank n. Then P has a unimodular element.

Proof. By a theorem due to Eisenbud and Evans [16] we can find a surjection  $P \rightarrow I \subset A$  such that ht(I) = d. Now the result follows from Theorem 4.3 and applying subtraction principle [11, Corollary 3.5].

**Example** 4.6. Here we show that the hypothesis  $\dim(R_0) \geq 1$  is also necessary in Corollary 4.5, where  $A = \bigoplus_{i>0} R_i$ . Let C, S, B, I and  $f_i$  be as in Example 4.4. Recall that the *d*-th Euler class group  $E^d(D[T])$  and the weak Euler class group  $E^d_0(D[T])$  as defined in [14], where D is a ring of dimension  $d \geq 3$  such that  $\mathbb{Q} \subset D$ . We consider  $(I,\omega_I) \in E^d(B[T])$ , where  $\omega_I$  is the local orientation induced by  $I = \langle f_1, f_2, f_3 \rangle +$  $I^2$ . Applying [9, Theorem 2.7] we can find a projective B[T]-module P (with trivial determinant) of rank 3 and a surjection  $\theta: P \to I$ . We claim that  $S^{-1}P$  does not have a unimodular element. First, we note that if  $S^{-1}P$  has a unimodular element, then there exists  $f \in \mathbb{C}[T] \setminus \{0\}$  such that  $P_f$  has a unimodular element. But then it follows from [10, Theorem 3.4] that P has a unimodular element. Hence, to prove our claim it is enough to show that P does not have a unimodular element. We fix a trivialization  $\chi: \wedge^3 P \xrightarrow{\sim} B[T]$ . Then it follows from [14, Proposition 5.8] and [2, Theorem 3.4] that  $E^3(B[T]) \cong E_0^3(B[T])$ . In particular, this give us  $e(P,\chi) = (I,\omega_I)$ in  $E^3(B[T])$ . Moreover, using [14, Corollary 4.11] we obtain that P has a unimodular element if and only if  $(I, \omega_I) = 0$  in  $E^3(B[T])$ . Now if  $(I, \omega_I) = 0$  in  $E^3(B[T])$ , then one may also lift  $IC = \langle f_1, f_2, f_3 \rangle C + I^2 C$  to a set of generators of IC. However, as it is shown in Example 4.4 that this is not feasible. Hence, the module *P* does not have a unimodular element.

In the next theorem we extend Theorem 4.3 and Corollary 4.5 to the case where the dimension of the graded ring is 2.

**Theorem 4.7.** Let R be as in Theorem 3.5 and  $\dim(R) = 2$ . Let P be a projective R-module of rank 2 with trivial determinant. Suppose  $I \subset R$  is an ideal such that  $I = \langle f_1, f_2 \rangle + I^2$ . Then

- (1) P is a free module and
- (2) there exist  $F_i \in I$  such that  $I = \langle F_1, F_2 \rangle$ , with  $F_i f_i \in I^2$ .

Proof. We consider  $S=R_0\setminus\{0\}$ . Then by Lemma 2.2 we get  $\dim(S^{-1}R)\leq 1$ . Since determinant of P is trivial, it follows from [35] that the module  $S^{-1}P$  is free. As P is finitely generated module over a Noetherian ring there exists an  $s\in S$  such that  $P_s$  is a free module. Let us take  $T=\{1+sr:r\in R_0\}$  and  $B=T^{-1}R$ . Then applying Lemma 2.2 we obtain that  $\dim(B_s)\leq 1$ . By Lemma 2.7 and Theorem 2.8 we get that, the module  $T^{-1}P$  has a unimodular element. Moreover, as determinant of P is trivial, the module  $T^{-1}P$  is free. Thus, there exists an element  $t\in T$  such that  $P_t$  is a free module. Therefore, the Quillen ideal  $J(R_0,P)$  of P is  $R_0$ . In other words, we have  $P\cong \frac{P}{PR_+}\otimes R$ . As  $\dim(R_0)=1$ , again by [35] the  $R_0$ -module  $\frac{P}{PR_+}$  has a unimodular element and hence free. Implying that P is free.

Now we consider  $I=\langle f_1,f_2\rangle+I^2$ . By [26] there exists  $e\in I^2$  such that  $I=\langle f_1,f_2,e\rangle$  where  $e(1-e)\in\langle f_1,f_2\rangle$ . Then  $I_e=R_e=\langle 1,0\rangle$  and  $I_{1-e}=\langle f_1,f_2\rangle_{1-e}$ . Since any unimodular row of length two can be completed to an invertible matrix, using a standard patching argument we obtain a projective R-module Q of rank 2 with trivial

determinant and a surjection  $\gamma: Q \to J$  such that  $\gamma$  locally lifts  $\{f_1, f_2\}$ . Now as Q is free by the previous case, the theorem concludes.

**Remark** 4.8. One may observe that in our circumstances we have been able to establish that Theorem 3.5 implies the main results of this section (e.g., 4.3, 4.5, 4.7). However, one cannot expect this to happen for an arbitrary ring. For an example we refer to [28, Remark 3.8].

4.1. **Precise obstruction.** Let  $R = \bigoplus_{i \geq 0} R_i$  be a graded domain of dimension  $d \geq 3$ , satisfying  $\dim(R_0) = 0$  and  $\frac{1}{d!} \in R$ . In this subsection, we aim to provide a necessary and sufficient condition for the splitting of projective R-modules of rank d with a trivial determinant. As the idea used in the proof is well-established due to R. Sridharan [36], we will present only a sketch of the proof.

**Theorem 4.9.** Let  $R=\bigoplus_{i\geq 0}R_i$  be a graded domain of dimension  $d\geq 3$  such that  $\dim(R_0)=0$  and  $\frac{1}{d!}\in R$ . Let P be a projective R-modules of rank d with trivial determinant. Suppose that there exists an R-linear surjection  $\alpha:P\to I$ , where  $I\subset R$  is an ideal of height d. Then P splits into a free summand of rank one if and only if  $\mu(I)=d$ .

Proof. First, we note that if P splits into a free summand of rank one, then  $\mu(I)=d$  follows using [26, Lemma 1]. For a detailed proof, we refer to [11, Corollary 4.4]. Hence, we assume that  $\mu(I)=d$ . Let  $I=\langle a_1,\ldots,a_d\rangle$  be a set of generators for I. We fix an isomorphism  $\chi:R\cong \wedge^d P$ . Let the pair  $\alpha,\chi$  induce  $I=\langle b_1,\ldots,b_d\rangle+I^2$ . We observe that in view of [11, Corollary 3.4], to prove the theorem, it suffices to show that there exist  $c_i\in I$  such that  $I=\langle c_1,\ldots,c_d\rangle$  and  $b_i-c_i\in I^2$ ,  $i=1,\ldots,d$ . The remaining part of the proof is devoted to showing only this.

Considering that two sets of generators of  $I/I^2$  may differ only by an invertible matrix in R/I, we find  $\widetilde{\theta} \in \mathrm{GL}_d(R/I)$  connecting the two surjections  $(R/I)^d \twoheadrightarrow I/I^2$  induced by  $a_i$ 's and  $b_i$ 's. We choose a lift  $\theta \in \mathrm{M}_{d \times d}(R)$  of  $\widetilde{\theta}$  and an element  $u \in R$  such that  $\det(\theta)u-1 \in I$ . Now, we consider the unimodular row  $w=(u,a_2,-a_1,a_3,\ldots,a_d) \in \mathrm{Um}_{d+1}(R)$ . Since  $\frac{1}{d!} \in R$  by Remark 3.7 the unimodular row w can be completed to the first row of a matrix in  $\mathrm{SL}_{d+1}(R)$ . Hence, using [2, Proposition 7.4], we can find a matrix  $\tau \in \mathrm{M}_{d \times d}(R)$  with  $\det(\tau) = \det(\theta)$  modulo I, such that  $(a_1,\ldots,a_d)\tau = (f_1,\ldots,f_d)$ , where  $I = \langle f_1,\ldots,f_d \rangle$ .

Let "bar" denote going modulo I. It follows from the construction of  $\tau$  that  $\overline{\theta}^{-1}\overline{\tau} \in \mathrm{SL}_d(R/I)$ . As  $\dim(R/I) = 0$  and  $d \geq 3$ , we have  $\mathrm{SL}_d(R/I) = \mathrm{E}_d(R/I)$ . Hence we can find a lift  $\gamma \in \mathrm{E}_d(R)$  of  $\overline{\theta}^{-1}\overline{\tau}$ . Then the required  $c_i's$  are defined as  $(c_1,\ldots,c_d):=(f_1,\ldots,f_d)\gamma$ .

# 5. CANCELLATION OF PROJECTIVE MODULES

This section is devoted to investigating the cancellation property of projective modules over a graded ring. We begin with a lemma, which is an analogy of [31, Lemma 2] in our setup.

**Lemma 5.1.** Let  $R = \bigoplus_{i \geq 0} R_i$  and M, M' be R-modules. Suppose that there exist  $s, t \in R_0$  be co-maximal and isomorphisms  $\sigma_1 : M_s \xrightarrow{\sim} M'_s$  and  $\sigma_2 : M_t \xrightarrow{\sim} M'_t$  such that

- (1)  $(\sigma_1)_t \equiv (\sigma_2)_s \mod (R_+)_{st};$
- (2)  $M_{st}$  is a free module.

Then there exists an isomorphism  $\sigma: M \xrightarrow{\sim} M'$  such that (i)  $\sigma_s \equiv \sigma_1 \mod (R_+)_s$  and (ii)  $\sigma_t \equiv \sigma_2 \mod (R_+)_t$ .

Proof. Let  $\operatorname{rank}(M_{st})=n$  and let "bar" denote going modulo  $R_+$ . Since  $M_{st}$  is free there exists an isomorphism  $\tau:M_{st}\stackrel{\sim}{\to} R^n_{st}$ . For an arbitrary isomorphism  $\gamma:M_{st}\stackrel{\sim}{\to} M_{st}$  we now consider the following commutative diagram

$$M_{st} \xrightarrow{\gamma} M_{st}$$

$$\downarrow^{\tau} \qquad \qquad \downarrow^{\tau}$$

$$R_{st}^{n} \xrightarrow{\widetilde{\gamma}} R_{st}^{n}$$

where  $\tilde{\gamma} = \tau \gamma \tau^{-1}$ . We will call  $\tilde{\gamma}$  is induced from  $\gamma$  and  $\tau$ .

We take  $\gamma = (\sigma_1)_t^{-1} \circ (\sigma_2)_s : M_{st} \xrightarrow{\sim} M_{st}$ . Then from (2) it follows that  $\overline{\gamma} = \operatorname{Id}$ . We consider the isomorphism  $\widetilde{\gamma} \in \operatorname{GL}_n(R_{st})$  induced from  $\gamma$  and  $\tau$ . Since  $\overline{\gamma} = \operatorname{Id}$ , we have  $\overline{\widetilde{\gamma}} = \operatorname{Id}$ . Applying Lemma 2.4 we can get a matrix, say  $\widetilde{\theta(T)} \in \operatorname{GL}_n(R_{st}[T])$  such that  $\widetilde{\theta(0)} = \operatorname{Id}$  and  $\widetilde{\theta(1)} = \widetilde{\gamma}$ . Let us take  $\theta(T) = (\tau \otimes R_{st}[T])^{-1}\widetilde{\theta(T)}(\tau \otimes R_{st}[T])$ . Then we observe that  $\theta(T) \in \operatorname{Aut}(M_{st}[T])$  such that  $\theta(0) = \operatorname{Id}$  and  $\theta(1) = \gamma$ . Now we define the isomorphism  $\phi(T) = (\sigma_1 \otimes R_{st}[T]) \circ \theta(T) : M_{st}[T] \xrightarrow{\sim} M'_{st}[T]$ . Then the proof follows from applying [31, Lemma 1].

Now we present the main theorem of the section.

**Theorem 5.2.** Let R and d be as in Theorem 3.5. Let P be a projective R-module of rank d such that  $J(R_0, P) \neq 0$ . Then P is cancellative.

Proof. First we comment that since  $J(R_0,P)\neq 0$  there exists an  $s\in R_0\setminus\{0\}$  such that  $P_s$  is a free module. To see this let us choose a non-zero element  $k\in R_0$  such that  $P_k$  is an extended projective module from  $R_0$ . Consider the multiplicative set  $T=R_0\setminus\{0\}$ . Since  $P_k$  is extended, the module  $T^{-1}P_k$  is free. Now as P is finitely generated we may choose a suitable multiple s=kl, for some  $l\in T$  and ensure the existence of such an s.

Let  $(f,p)\in \mathrm{Um}(R\oplus P)$ . Since R has a non-trivial grading, the ideal  $R_+$  is non-zero. In particular, the height of  $R_+$  is  $\geq 1$ . Hence, going modulo a non-zero element  $g\in R_+$  and altering (f,p) suitably via an element of  $\mathrm{Aut}(R\oplus P)$  we may assume that  $f-1\in \langle g\rangle$  and  $p\in \langle g\rangle P$  [cf. Theorem 3.5, first paragraph]. We take  $P'=\frac{R\oplus P}{(f,p)R}$ . Then to prove the theorem it is enough to show that there exists an isomorphism  $\sigma:P\stackrel{\sim}{\to}P'$ . Moreover, we comment on an observation that finding a  $\sigma:P\stackrel{\sim}{\to}P'$  such that  $\overline{\sigma}=\mathrm{Id}$  is equivalent to find an  $\alpha\in\mathrm{Aut}(R\oplus P)$  such that  $\alpha(f,p)=(1,0)$  and  $\overline{\alpha}=\mathrm{Id}$ .

Let "bar" denote going modulo  $R_+$  as well as  $PR_+$ . As  $P_s$  is free, by Theorem 3.5 we can find  $\alpha_1 \in \operatorname{Aut}(R_s \oplus P_s)$  such that (1)  $\alpha_1(f,p) = (1,0)$  and (2)  $\overline{\alpha}_1 = \operatorname{Id}$ . Then  $\alpha_1$  will induce an isomorphism  $\sigma_1 : P_s \xrightarrow{\sim} P'_t$  such that  $\overline{\sigma}_1 = \operatorname{Id}$ .

Let  $S=\{1+sr:r\in R_0\}$ . We denote  $B=S^{-1}R$ ,  $L=S^{-1}P$  and  $L'=S^{-1}P'$ . We note that  $s\in \operatorname{Jac}(S^{-1}R_0)$ . Therefore, by Proposition 2.2 we get  $\dim(B_s)\leq d-1$ . Hence, using Lemma 2.7 we can obtain a generalized dimension function  $\delta:\operatorname{Spec}(B)\to\mathbb{N}$  such that  $\delta(\mathfrak{p})\leq d-1$  for all  $\mathfrak{p}\in\operatorname{Spec}(B)$ . Since  $p\in\langle g\rangle P$ , we note that  $(f,p)\in\operatorname{Um}(R\oplus\langle g\rangle P)$ . Moreover, the module  $\langle g\rangle L$  is a projective B-module of rank d. Hence, applying Theorem 2.8 we can find  $p'\in P$  such that  $q:=p+gfp'\in\operatorname{Um}(\langle g\rangle L)$ . Moreover, as  $q\in\langle g\rangle L$  and  $f-1\in\langle g\rangle$  one may obtain an  $\alpha_2\in\operatorname{Aut}(B\oplus L)$  such that  $\alpha_2(f,p)=(1,0)$  and  $\overline{\alpha}_2=\operatorname{Id}$ . Then  $\alpha_2$  will induce an isomorphism  $\sigma_2:L\overset{\sim}{\to} L'$  such that  $\overline{\sigma}_2=\operatorname{Id}$ . Since

all modules are finitely generated (over a Noetherian ring) there exists an isomorphism  $\sigma_2: P_t \xrightarrow{\sim} P_t'$  such that  $\overline{\sigma}_2 = \mathrm{Id}$ , for some  $t \in S$ .

Now applying Lemma 5.1 we get the required isomorphism  $\sigma: P \xrightarrow{\sim} P'$  such that  $\overline{\sigma} = \mathrm{Id}$ . This completes the proof.

**Remark** 5.3. Let R and P be as in Theorem 5.2, and we consider  $S = R_0 \setminus \{0\}$ . Then  $J(R_0, P) \neq 0$  if and only if  $S^{-1}P$  is free. If  $S^{-1}R_0 = \overline{\mathbb{Q}}$  and  $\wedge^d P \cong R$ , then it follows from [22, Theorem 6.4.2] and [35] that  $J(R_0, P) \neq 0$ .

**Remark** 5.4. It would be interesting to know whether the hypothesis  $J(R_0, P) \neq 0$  in Theorem 5.2 is necessary or not.

We now discuss an interesting consequence of Theorem 5.2.

**Corollary 5.5.** Let R be an integral domain of dimension  $d \ge 1$ , and A be a graded subring of R[T] containing R such that  $\dim(A) = d + 1$ . Let P be a projective A-module of rank d + 1, so that the determinant of P is extended from the base ring R. Then P is cancellative.

Proof. Let us take  $\mathfrak{T}=R\setminus\{0\}$ . Since  $\dim(A)=d+1$ , there exists an  $a\in\mathfrak{T}$  such that  $A_a$  contains a monic polynomial in T. Then  $\mathfrak{T}^{-1}A\hookrightarrow(\mathfrak{T}^{-1}R)[T]$  is an integral extension. This further implies that for any multiplicative set  $S\subset R$  containing a, we must have  $\dim(S^{-1}A)=\dim(S^{-1}R)+1$ . Therefore, in view of Theorem 5.2 it is enough to show that  $\operatorname{ht}(J(R,P))\geq 1$ . To prove this we observe that, since  $\dim(\mathfrak{T}^{-1}R)=0$ , we have  $\dim(\mathfrak{T}^{-1}A)=1$ . As the determinant of P is extended from R, applying [35] the module  $\mathfrak{T}^{-1}P$  is a free  $\mathfrak{T}^{-1}A$ -module. Hence, there exists an element  $s\in\mathfrak{T}$  such that  $P_s$  is a free  $A_s$ -module. That is, the non-zero element  $s\in J(R,P)$ .

## 6. APPLICATIONS

This section is devoted to establishing some consequences of Theorem 4.3.

6.1. On a question of Nori: non-smooth graded case. Let  $R = \bigoplus_{i \geq 0} R_i$  be an affine graded domain of dimension  $d \geq 3$  over a field k such that  $\mathbb{Q} \subset k$  and  $\dim(R_0) \geq 1$ . In the next theorem, we show that Nori's question [25] on homotopy of sections of projective modules has an affirmative answer over R even without the smoothness assumption.

**Theorem 6.1.** Let R be as in Theorem 3.5 and  $d \ge 3$ . Moreover, we assume that R is an affine algebra over an infinite field such that  $\frac{1}{d!} \in R$ . Let  $I \subset R[T]$  an ideal such that  $\mu(I/I^2T) = ht(I) = d$ . Then any set of generators of  $I/I^2T$  lifts to a set of generators of I.

Proof. If I contains a monic polynomial in T, then the result follows from [25]. Hence, without loss of generality, we may assume that I does not contain a monic polynomial in T. Let  $I = \langle f_1, \ldots, f_d \rangle + I^2T$ . First, we comment that in [14, Theorem 3.10] the hypothesis that the ring containing  $\mathbb Q$  can be weakened by assuming the ring contains an infinite field such that d! is invertible. We denote  $R(T) = \mathfrak{T}^{-1}R[T]$ , where  $\mathfrak{T}$  be the ring consisting of all monic polynomials in R[T]. In view of [14, Theorem 3.10] it is enough to prove that there exist  $F_i \in IR(T)$ , such that  $IR(T) = \langle F_1, \ldots, F_d \rangle$  and  $f_i - F_i \in I^2R(T)$ . The proof is devoted to establishing only this.

Consider the multiplicative set  $S=\{f\in R_0[T]: f \text{ is a monic polynomial}\}$  and let  $B=S^{-1}R[T]$ . We consider the grading  $R[T]=\bigoplus_{i\geq 0}R_i[T]$ . Let  $\mathfrak m$  be a maximal ideal in

 $R_0[T]$ . Then  $\operatorname{ht}(\mathfrak{m})=\dim(R_0)+1$ . Therefore, from Suslin's monic polynomial theorem (see [23, Chapter III, §3, 3.3, page no 108]) we have  $S\cap\mathfrak{m}\neq\emptyset$ . Then by Proposition 2.2 we have  $\dim(B)\leq d$ . Since  $S\subset R_0[T]$ , the ring  $B=\bigoplus_{i\geq 0}S^{-1}R_i[T](=\bigoplus_{i\geq 0}B_i$  say) is also a graded domain of dimension d such that  $\dim(B_0)=\dim(S^{-1}R_0[T])=\dim(R_0)\geq 1$ . As I is not containing a monic polynomial we have  $\operatorname{ht}(IB)\geq d$ . Moreover, since T is a unit in B, we have  $IB=\langle f_1,\ldots,f_d\rangle B+I^2B$ . Now, applying Theorem 4.3 we obtain  $F_i\in IB$  such that  $IB=\langle F_1,\ldots,F_d\rangle B$  and  $f_i-F_i\in I^2B$ . Since B is a subring of R(T) we get  $IR(T)=\langle F_1,\ldots,F_d\rangle R(T)$  such that  $f_i-F_i\in I^2R(T)$ . This concludes the proof.

**Remark** 6.2. It follows from [8, Example 6.4] that in Theorem 6.1 the hypothesis  $\dim(R_0) > 0$  is necessary.

**Corollary 6.3.** Let R and d be as in Theorem 6.1. Let  $I \subset R[T]$  an ideal such that  $\mu(I/I^2) = ht(I) = d$ . Then any set of generators of  $I/I^2$  lifts to a set of generators of I.

Proof. Let  $I=\langle f_1,\ldots,f_d\rangle+I^2$  and let  $I(0)=\{f(0):f\in R[T]\}$ . Then as R contains an infinite field without loss of generality we may assume that  $I(0)\subset R$  is an ideal of height  $\geq d$  (for details see the proof of [12, Theorem 3.4]). If  $\operatorname{ht}(I(0))>d$ , then we can always lift any set of generators of  $I(0)/I(0)^2$ . Now if  $\operatorname{ht}(I(0))=d$ , then applying Theorem 4.3 there exist  $a_i\in I$  such that  $I(0)=\langle a_1,\ldots,a_d\rangle$ , with  $f_i(0)-a_i\in I(0)^2$ , for  $i=1,\ldots,d$ . Hence, by [8, Remark 3.9] there exist  $g_i\in I$  such that  $I=\langle g_1,\ldots,g_d\rangle+I^2T$  with  $f_i-g_i\in I^2$  and  $g_i(0)=a_i$ , for  $i=1,\ldots,d$ . Now the result follows from Theorem 6.1.

**Corollary 6.4.** Let R and d be as in Theorem 6.1. Let P be a projective R[T]-module with trivial determinant of rank d. Then P has a unimodular element.

Proof. Applying a theorem due to Eisenbud and Evans [16] we can find a surjection  $P \to I \subset A[T]$  such that  $\operatorname{ht}(I) = d$ . Then the result follows from Corollary 6.3 and subtraction principle as stated in [14, Corollary 4.13] (taking  $Q = (R[T])^{d-1}$ ,  $I_1 = R[T]$  and  $I_2 = I$ ).

Remark 6.5. One can remove the restriction on the base field in Corollary 6.4 in the following way: let P be a projective R[T]-module of rank d with trivial determinant. Recall that the ring R(T) is obtained by localizing R[T] with respect to the multiplicative set consisting of all monic polynomials in R[T]. Then, in view of [7, Theorem 5.2 and Remark 5.3], it is enough to show that the modules P/TP and  $P \otimes R(T)$  have unimodular elements. Let S be the multiplicative set consisting of all monic polynomials in  $R_0[T]$ . Then, it follows from Corollary 4.5 that P/TP and  $S^{-1}P$  (and hence  $P \otimes R(T)$ ) have unimodular elements.

6.2. Generating ideals up to projective equivalence. Recall that, two ideals I and J in a ring A are said to be *projectively equivalent* if some power of I and some power (usually different) of J have the same integral closure. The following theorem is an improvement of [19] in our setup.

**Theorem 6.6.** Let A and n be as in Theorem 4.3. Let  $I \subset A$  be an ideal of height  $\geq 2$ . Then there exists an ideal  $J \subset A$  projectively equivalent to I satisfying  $\mu(J) \leq n$ .

Proof. First we observe that combining the results [15, Proposition 2.2] and Theorem 4.3 one can prove the following: let  $K \subset A$  be an ideal such that (i)  $\mu(K/K^2) \leq n$  and (ii)  $\operatorname{ht}(K) \geq 2$ . Then  $\mu(K) \leq n$ . Applying [19] we obtain an ideal  $J \subset A$  such that (1) I and J are projectively equivalent, (2)  $\operatorname{ht}(J) \geq 2$  and (3)  $\mu(J/J^2) \leq n$ . Now it follows from the previously mentioned observation that  $\mu(J) \leq n$ .

#### ACKNOWLEDGMENT

I thank Mrinal Kanti Das for suggesting Theorem 6.6.

#### REFERENCES

- [1] S. Banerjee. Subrings of polynomial rings and the conjectures of Eisenbud and Evans. *Journal of Algebra*, 641:85–104, Mar. 2024. doi:10.1016/j.jalgebra.2023.10.039.
- [2] S. Banerjee and M. K. Das. Splitting criteria for projective modules over polynomial algebras. June 2022. arXiv: 2206.06819.
- [3] S. Banerjee and M. K. Das. On a question of Nori: Obstructions, improvements, and applications. *Journal of Algebra*, 635:271–299, Dec. 2023. doi:10.1016/j.jalgebra.2023.08.003.
- [4] H. Bass. Some problems in "classical" algebraic K-theory. In *Lecture Notes in Mathematics*, pages 1–73. Springer Berlin Heidelberg, 1973. doi:10.1007/bfb0073718.
- [5] R. Basu and M. K. Singh. Quillen-Suslin theory for classical groups: revisited over graded rings. In *Categorical, homological and combinatorial methods in algebra*, volume 751 of *Contemp. Math.*, pages 5–18. Amer. Math. Soc., [Providence], RI, [2020] ©2020. doi:10.1090/conm/751/15113.
- [6] S. M. Bhatwadekar and M. K. Keshari. A question of Nori: Projective generation of ideals. *K-Theory*, 28(4):329–351, apr 2003. doi:10.1023/a:1026217116072.
- [7] S. M. Bhatwadekar, H. Lindel, and R. A. Rao. The Bass-Murthy question: Serre dimension of Laurent polynomial extensions. *Inventiones Mathematicae*, 81(1):189–203, feb 1985. doi:10.1007/bf01388777.
- [8] S. M. Bhatwadekar and R. Sridharan. Projective generation of curves in polynomial extensions of an affine domain and a question of Nori. *Inventiones Mathematicae*, 133(1):161–192, jun 1998. doi:10.1007/s002220050243.
- [9] S. M. Bhatwadekar and R. Sridharan. Projective generation of curves in polynomial extensions of an affine domain (II). *K-Theory*, 15(3):293–300, nov 1998. doi:10.1023/a:1007739731247.
- [10] S. M. Bhatwadekar and R. Sridharan. Zero cycles and the Euler class groups of smooth real affine varieties. *Inventiones Mathematicae*, 136(2):287–322, a 1999. doi:10.1007/s002220050311.
- [11] S. M. Bhatwadekar and R. Sridharan. The Euler class group of a Noetherian ring. *Compositio Mathematica*, 122(2):183–222, 2000. doi:10.1023/a:1001872132498.
- [12] S. M. Bhatwadekar and R. Sridharan. On a question of Roitman. *J. Ramanujan Math. Soc*, 16(1):45–61, 2001.
- [13] W. Bruns and J. Gubeladze. *Polytopes, rings, and K-theory*. Springer monographs in mathematics. Springer, 2009. doi:10.1007/b105283.
- [14] M. K. Das. The Euler class group of a polynomial algebra. *Journal of Algebra*, 264(2):582–612, jun 2003. doi:10.1016/s0021-8693(03)00240-0.
- [15] M. K. Das and R. Sridharan. Good invariants for bad ideals. *Journal of Algebra*, 323(12):3216–3229, 2010. doi:10.1016/j.jalgebra.2010.04.006.
- [16] D. Eisenbud and E. G. Evans, Jr. Generating modules efficiently: theorems from algebraic *K*-theory. *Journal of Algebra*, 27:278–305, 1973. doi:10.1016/0021-8693 (73) 90106-3.
- [17] J. Fasel, R. A. Rao, and R. G. Swan. On stably free modules over affine algebras. *Publications mathématiques de l'IHÉS*, 116(1):223–243, jun 2012. doi:10.1007/s10240-012-0041-y.
- [18] J. Gubeladze. Unimodular rows over monoid rings. *Advances in Mathematics*, 337:193–215, oct 2018. doi:10.1016/j.aim.2018.08.011.
- [19] D. Katz. Generating ideals up to projective equivalence. *Proceedings of the American Mathematical Society*, 120(1):79–83, 1994. doi:10.2307/2160169.

- [20] M. K. Keshari. Euler class group of a Noetherian ring. M-Phil Thesis-2001 (52 pages)., Aug. 2001. arXiv:1408.2645.
- [21] M. K. Keshari and M. A. Mathew. On Serre dimension of monoid algebras and Segre extensions. *Journal of Pure and Applied Algebra*, 226(9):107058, sep 2022. doi:10.1016/j.jpaa.2022.107058.
- [22] A. Krishna and V. Srinivas. Zero cycles on singular varieties. In *Algebraic cycles and motives*. *Vol.* 1, volume 343 of *London Math. Soc. Lecture Note Ser.*, pages 264–277. Cambridge Univ. Press, Cambridge, 2007. doi:10.1017/CB09780511721496.007.
- [23] T. Y. Lam. Serre's Problem on Projective Modules. Springer Berlin Heidelberg, 2006. doi:10.1007/978-3-540-34575-6.
- [24] H. Lindel. On projective modules over positively graded rings. *Tata Inst. Fund. Res., Bombay,* 11:251–273, 1987.
- [25] S. Mandal (with an appendix by M. V. Nori). Homotopy of sections of projective modules. *J. Algebraic Geom.* 1 (1992), (4):639–646, 1992.
- [26] N. Mohan Kumar. Complete intersections. *Journal of Mathematics of Kyoto University*, 17(3):533–538, 1977. doi:10.1215/kjm/1250522714.
- [27] N. Mohan Kumar. On two conjectures about polynomial rings. *Inventiones Mathematicae*, 46(3):225–236, oct 1978. doi:10.1007/bf01390276.
- [28] N. Mohan Kumar, M. P. Murthy and A. Roy. A cancellation theorem for projective modules over finitely generated rings. Algebraic geometry and commutative algebra, Vol. I, 281–287, Kinokuniya, Tokyo, 1988.
- [29] M. P. Murthy. Zero cycles and projective modules. *Annals of Mathematics*, 140(2):405–434, sep 1994. doi:10.2307/2118605.
- [30] M. P. Murthy and R. G. Swan. Vector bundles over affine surfaces. *Inventiones Mathematicae*, 36(1):125–165, dec 1976. doi:10.1007/bf01390007.
- [31] B. Plumstead. The conjectures of Eisenbud and Evans. *American Journal of Mathematics*, 105(6):1417–1433, 1983. doi:10.2307/2374448.
- [32] D. Quillen. Projective modules over polynomial rings. *Inventiones Mathematicae*, 36:167–171, 1976. doi:10.1007/BF01390008.
- [33] R. A. Rao. The Bass-Quillen conjecture in dimension three but characteristic  $\neq 2,3$  via a question of A. Suslin. *Inventiones Mathematicae*, 93(3):609–618, 1988. doi:10.1007/BF01410201.
- [34] M. Roitman. On stably extended projective modules over polynomial rings. *Proceedings of the American Mathematical Society*, 97(4):585–589, 1986. doi:10.1090/s0002-9939-1986-0845969-9.
- [35] J.-P. Serre. Modules projectifs et espaces fibrés à fibre vectorielle. *Séminaire Dubreil. Algèbre et théorie des nombres*, 11(2):1–18, 1957-1958. URL: http://eudml.org/doc/111153.
- [36] R. Sridharan. Non-vanishing sections of algebraic vector bundles. *Journal of Algebra*, 176(3):947–958, sep 1995. doi:10.1006/jabr.1995.1279.
- [37] V. Srinivas. Vector bundles on the cone over a curve. *Compositio Mathematica*, 47(3):249–269, 1982. URL: http://www.numdam.org/item/CM\_1982\_\_47\_3\_249\_0/.
- [38] A. A. Suslin. On the structure of the special linear group over polynomial rings. *Mathematics of the USSR-Izvestiya*, 11(2):221–238, apr 1977. doi:10.1070/im1977v011n02abeh001709.
- [39] A. A. Suslin. Cancellation over affine varieties. *Journal of Soviet Mathematics*, 27(4):2974–2980, nov 1984. doi:10.1007/bf01410752.
- [40] L. N. Vaseršteĭn. On the stabilization of the general linear group over a ring. *Mathematics of the USSR-Sbornik*, 8(3):383–400, apr 1969. doi:10.1070/sm1969v008n03abeh001279.
- [41] T. Vorst. The general linear group of polynomial rings over regular rings. *Communications in Algebra*, 9(5):499–509, jan 1981. doi:10.1080/00927878108822596.

THE INSTITUTE OF MATHEMATICAL SCIENCES, HBNI, C.I.T. CAMPUS, THARAMANI, CHENNAI 600113, INDIA

Email address: sourjyab@imsc.res.in, sourjya91@gmail.com