Explicit Formula for Partial Information Decomposition

Aobo Lyu*, Andrew Clark*, and Netanel Raviv[†]

*Department of Electrical and Systems Engineering, Washington University in St. Louis, St. Louis, MO, USA [†]Department of Computer Science and Engineering, Washington University in St. Louis, St. Louis, MO, USA aobo.lyu@wustl.edu, andrewclark@wustl.edu, netanel.raviv@wustl.edu

Abstract—Mutual information between two random variables is a well-studied notion, whose understanding is fairly complete. Mutual information between one random variable and a pair of other random variables, however, is a far more involved notion. Specifically, Shannon's mutual information does not capture finegrained interactions between those three variables, resulting in limited insights in complex systems. To capture these fine-grained interactions, in 2010 Williams and Beer proposed to decompose this mutual information to information atoms, called unique, redundant, and synergistic, and proposed several operational axioms that these atoms must satisfy. In spite of numerous efforts, a general formula which satisfies these axioms has yet to be found. Inspired by Judea Pearl's do-calculus, we resolve this open problem by introducing the *do-operation*, an operation over the variable system which sets a certain marginal to a desired value, which is distinct from any existing approaches. Using this operation, we provide the first explicit formula for calculating the information atoms so that Williams and Beer's axioms are satisfied, as well as additional properties from subsequent studies in the field.

I. INTRODUCTION

Since its inception by Claude Shannon [1], mutual information has remained a pivotal measure in information theory, which finds extensive applications across multiple other domains. Extending mutual information to multivariate systems has attracted significant academic interest, but no widely agreed upon generalization exists to date. For instance, the so-called *interaction information* [2] emerged in 1960 as an equivalent notion for mutual information in multivariate systems, and yet, it provides negative values in many common systems, contradicting Shannon's viewpoint of information measures as nonnegative quantities.

Arguably the simplest multivariate setting in which Shannon's mutual information fails to capture the full complexity of the system is that of a three variable system, with *two source variables*, and *one target variable*. Mutual information between the source variables and the target variable does not provide insights about *how* the source variables influence the target variable. Specifically, in various points of the probability space the value of the target variable might be computable either:

- (a) exclusively from one source variable (but not the other);
- (b) either one of the source variables; or
- (c) both variables jointly (but not separately).

In 2010 William and Beer [3] proposed to formalize the above fine-grained interactions in a three variable system¹ using an axiomatic approach they called *Partial Information Decomposition* (PID). They proposed decomposing said mutual information to four constituent ingredients called information atoms, which capture the above possible interactions between the variables:

- (a) two *unique information atoms*, one for each source variable, which capture the information each source variable implies about the target variable, that cannot be inferred from the other;
- (b) one *redundant information atom*, which captures the information that can inferred about the target variable from either one of the source variables; and
- (c) one synergistic information atom, which captures the information that can be inferred about the target variable from both source variables jointly, but not individually.

Ref. [3] proposed a set of axioms that the above information atoms should satisfy in order to provide said insights, and follow-up works in the field identified several additional properties [4]–[7]. Yet, in spite of extensive efforts [8]–[12], a comprehensive definition of information atoms which satisfies all these axioms and properties is yet to be found.

In spite of limited understanding of the information atoms, PID has already found multiple applications in various fields. As a simple example [13, Fig. 1], one can imagine the two source variables being education level and gender, and the target variable being annual income. An exact formula for computing the information atoms would shed insightful information about the extent to which annual income is a result of education level, gender, either one, or both.

Beyond this simple example, PID has broad applications in a wide range of fields. In brain network analysis, PID (or similar ideas) has been instrumental in measuring correlations between neurons [14] and understanding complex neuronal interactions in cognitive processes [15]. For privacy and fairness studies, the synergistic concept provides insights about data disclosure mechanisms [16], [17]. In the field of causality, information decomposition can be used to distinguish and quantify the occurrence of causal emergence [18], and more.

¹William and Beer formulated their notions for any number of source variables, and yet herein we focus on three variables for simplicity. Extending our methods to multiple source variables will be addressed in future version of this paper.

In this paper, we propose an explicit PID formula that satisfies all of Williams and Beer's axioms, as well as several additional desired properties. We do so by introducing the *dooperation*, which is inspired by similar concepts in the field of causal analysis [19]–[21]. Intuitively, based on the understanding that unique information is "ideal conditional mutual information," our method first adjusts the entire probability distribution by using the do-operation in order to make the target variable identical to its conditional distribution given one source variable, and then calculates the expectation of mutual information. It is worth noting that our method is not based on any of the point-wise, localized, or optimization approaches that existing methods use.

We begin in Section II by introducing the PID framework, and its axioms and properties. We continue in Section III by introducing our do-operation and the definition of unique information, from which all other definitions follow, and prove that all axioms and properties are satisfied. We discuss the intrinsic meaning of our definition in Section IV, and provide all proofs in the appendix.

II. FRAMEWORK, AXIOMS, AND PROPERTIES

The following notational conventions are observed throughout this article: X, \mathcal{X}, x (similarly Y, \mathcal{Y}, y etc.) denote a random variable, its corresponding (finite) alphabet, and an element of that alphabet, respectively. The distribution of Xis denoted by \mathcal{D}_X , the joint distribution of X and Y is denoted by $\mathcal{D}_{X,Y}$, and the distribution of X given Y = y is denoted by $\mathcal{D}_{X|Y=y}$.

For random variables X, Y, Z, the quantity I((X, Y); Z) captures the amount of information that one *target variable* Z shares with the *source variables* (X, Y), but provides no further information regarding finer interactions between the three variables. To gain more subtle insights into the interactions between Z and (X, Y), [3] proposed to further decompose I((X, Y); Z) into *information atoms*. Specifically, the shared information between Z and (X, Y) should contain a *redundant* information atom, two *unique* information atoms, and one *synergistic* information atom (see Figure 1).

The redundant information atom $\operatorname{Red}(X, Y \to Z)$ (also called "shared") represents the information which either X or Y imply about Z. The unique information atom $\operatorname{Un}(X \to Z|Y)$ represents the information individually contributed to Z by X, but not by Y (similarly $\operatorname{Un}(Y \to Z|X)$). The synergistic information atom $\operatorname{Syn}(X, Y \to Z)$ (also called "complementary"), represents the information that can only be known about Z through the *joint* observation of X and Y, but cannot be provided by either one of them separately. Together, we must have that

$$I((X,Y);Z) = \operatorname{Red}(X,Y \to Z) + \operatorname{Syn}(X,Y \to Z) + \operatorname{Un}(X \to Z|Y) + \operatorname{Un}(Y \to Z|X).$$
(1)

We refer to (1) as *Partial Information Decomposition* (PID).

Moreover, since the redundant atom together with one of the unique atoms constitute all information that one source



Fig. 1. A pictorial representation of Partial Information Decomposition (1), where I((X, Y); Z) is decomposed to its finer information atoms, the synergistic $Syn(X, Y \to Z)$ (also called "complementary"), the redundant $Red(X, Y \to Z)$ (also called "shared"), and the two directional unique components $Un(X \to Z|Y)$ and $Un(Y \to Z|X)$. The summation of the redundant atom and one of the unique atoms must be equal to the corresponding mutual information, as described in Eq. (2).

variable implies about the target variable, it must be the case that their summation equals the mutual information between the two, i.e., that

$$I(X;Z) = \operatorname{Red}(X, Y \to Z) + \operatorname{Un}(X \to Z|Y), \text{ and}$$

$$I(Y;Z) = \operatorname{Red}(X, Y \to Z) + \operatorname{Un}(Y \to Z|X).$$
(2)

In a similar spirit, the synergistic information atom and one of the unique information atoms measure shared information between the target variable and one of the source variables, while excluding the other source variable. Therefore, the summation of these quantities should coincide with the wellknown definition of conditional mutual information, i.e.,

$$I(Z; X|Y) = \operatorname{Syn}(X, Y \to Z) + \operatorname{Un}(X \to Z|Y), \text{ and}$$

$$I(Z; Y|X) = \operatorname{Syn}(X, Y \to Z) + \operatorname{Un}(Y \to Z|X).$$
(3)

Eqs. (2), and (3) are the foundation of an axiomatic approach towards an operational definition of the information atoms. These equations form the first in a series of axioms, presented next, which were raised in previous works on the topic [3], [22], [23]. Such axiomatic approach was also taken in the past in order to shed light on Shannon's mutual information [24].

Axiom 1 (Information atoms relationship). *Partial Information Decomposition* (1) *satisfies* (2) *and* (3).

Notice that it suffices to specify the definition of any one of the information atoms, and the definitions for the remaining atoms follow from Axiom 1. Consequently, [3], [25] chose to specify Red, and provided three additional axioms which Red should satisfy.

The first additional axiom is *commutativity* of the source variables, which implies that the order of the source variables must not affect the value of the redundant information.

Axiom 2 (Commutativity). Partial Information Decomposition satisfies $\operatorname{Red}(X, Y \to Z) = \operatorname{Red}(Y, X \to Z)$.

The second is *monotonicity*, which implies that the redundant information is non-increasing when adding a source variable, since the newly added variable cannot increase the redundancy between the original variables. We sidestep the discussion about monotonicity with more than two variables, which is not our focus in this paper, even though it can be easily obtained by extending our definition to more than two source variables.

The third is *self-redundancy*, which defines the redundant information from *one* source variable to the target variable (i.e., $\text{Red}(X \rightarrow Z)$) as the mutual information between them. In the case of two source variables considered herein, monotonicity and self-redundancy merge into the following single axiom.

Axiom 3 (Monotonicity and self-redundancy). Partial Information Decomposition satisfies $\operatorname{Red}(X, Y \to Z) \leq \min\{I(X; Z), I(Y; Z)\}$.

Notice that Axiom 3, alongside Axiom 1 (specifically (2)), imply that Un is a nonnegative quantity. The nonnegativity of Red is stated in [3], [25] as a separate axiom, shown next.

Axiom 4 (Nonnegativity). *Partial Information Decomposition* satisfies $\operatorname{Red}(X, Y \to Z) \ge 0$.

The nonnegativity of Syn is normally not listed as an axiom, since it is debatable if it should or should not be nonnegative; we will show that our method yields nonnegative Syn under the *closed-system assumption* (i.e., H(Z|X, Y) = 0) in Section III, and further discussion is given in Section IV.

Besides, subsequent to [3], [25], studies suggested two additional properties, *additivity* and *continuity* [12], [26]. Additivity implies that whenever independent variable systems are considered, the joint information measures should be the sum of the information measures of each individual system. This is the case, for instance, in joint entropy of two independent variables.

Property 1 (Additivity). *Partial Information Decomposition* of two independent systems $\mathcal{D}_{X,Y,Z}$ and $\mathcal{D}_{\bar{X},\bar{Y},\bar{Z}}$ satisfy

$$\begin{aligned} &\operatorname{Un}((X,X) \to (Z,Z)|(Y,Y)) \\ &= \operatorname{Un}(X \to Z|Y) + \operatorname{Un}(\bar{X} \to \bar{Z}|\bar{Y}), \text{ and} \\ &\operatorname{F}((X,\bar{X}),(Y,\bar{Y}) \to (Z,\bar{Z})) = \\ &\operatorname{F}(X,Y \to Z) + \operatorname{F}(\bar{X},\bar{Y} \to \bar{Z}), \end{aligned}$$

for every $F \in \{Red, Syn\}$.

Continuity implies that small changes in the probability distribution lead to small changes in the value of the information measure. It ensures that the measure behaves predictably and is a key property in information theory, particularly for measures like entropy and mutual information.

Property 2 (Continuity). Red, Un, and Syn are continuous functions from the underlying joint distributions $\mathcal{D}_{X,Y,Z}$ to \mathbb{R} .

In addition, another well-known property is *independent identity* [9], which asserts that in a system of two independent source variables and a target variable which equals to their joint distribution, the redundant information should be zero.

Property 3 (Independent Identity). If I(X, Y) = 0 and Z = (X, Y), then $\operatorname{Red}(X, Y \to Z) = 0$.

We mention that several important properties can be inferred from the above. For example, the non-negativity of Un can be obtained from Axiom 1 and Axiom 3 as mentioned earlier; the commutativity of Syn follows from Axiom 1 and Axiom 2; the difference between (2) and (3) is often called *consistency* [12], etc.

Finally, we emphasize once again that none of the existing operational definitions of the information atoms satisfy all of the above. A comprehensive list of violations is beyond the page limit of this paper, and yet we briefly mention that Axiom 4 (nonnegativity) is violated by [8], [9], [27] (although some sources do not refer to non-negativity as a requirement); Property 1 (additivity) is violated by all works except [12], [28], [8], and [29] according to [26]; Property 3 (independent identity) is violated by [3]; Property 2 (continuity) is violated by [8], [11], [28], [29], etc.

III. PROPOSED INFORMATION DECOMPOSITION DEFINITION

In this section, we present our operational definition of Un, from which the definitions of the remaining information atoms follow. Then, we explain the logic behind this definition, and prove that it satisfies all the axioms and properties proposed in Section II.

A. Definition of Information Atoms

Given a system X, Y, Z define a new random variable X' over the alphabet \mathcal{X} via its conditional joint distribution as follows:

$$\Pr(X' = x, Z = z | Y = y)$$

$$\triangleq \Pr(X = x | Z = z) \Pr(Z = z | Y = y), \quad (4)$$

meaning, for every $x \in \mathcal{X}$,

$$\Pr(X'=x) = \sum_{y,z\in\mathcal{Y}\times\mathcal{Z}} \Pr(X'=x, Z=z|Y=y) \Pr(Y=y).$$

The variable X' is well-defined since all probabilities are nonnegative, and since

$$\sum_{x \in \mathcal{X}} \Pr(X' = x)$$

=
$$\sum_{x, y, z \in \mathcal{X} \times \mathcal{Y} \times \mathcal{Z}} \Pr(X' = x, Z = z | Y = y) \Pr(Y = y)$$

=
$$\sum_{y \in \mathcal{Y}} \Pr(Y = y) = 1.$$

Definition 1 (Unique Information). The unique information from X to Z given Y is $Un(X \rightarrow Z|Y) = I(X'; Z|Y)$.

The definitions for the remaining information atoms are then implied by Axiom 1 as follows.

Definition 2 (Redundant Information). *The Redundant Information from X and Y to Z is defined as:*

$$\operatorname{Red}(X, Y \to Z) = I(X; Z) - \operatorname{Un}(X \to Z|Y).$$

Definition 3 (Synergistic Information). The synergistic information from X and Y to Z is defined as:

$$\operatorname{Syn}(X, Y \to Z) = I(X; Z|Y) - \operatorname{Un}(X \to Z|Y).$$

It should be noted that Definition 2 and Definition 3 strictly depend on the order of the source variables; the commutativity of Red (Axiom 2) will be addressed in the sequel, and the commutativity of Syn follows from Axiom 1 and Axiom 2 as mentioned earlier.

B. Intuitive Explanation of Definition 1.

Our definition of unique information Un is derived from a newly defined *do-operation*.

Definition 4 (Do-operation). Given $\mathcal{D}_{X,Z}$ and \mathcal{D}_C such that the support of \mathcal{D}_C is contained in the support of \mathcal{D}_Z , let $do(\mathcal{D}_{X,Z}|\mathcal{D}_C) = \mathcal{D}_{A,C}$, where

$$\Pr(A = x, C = z) = \Pr(X = x | Z = z) \Pr(C = z)$$
(5)

for all $x, z \in \mathcal{X} \times \mathcal{Z}$.

In Lemma 8, which is given and proved in Appendix A, it is shown that $\mathcal{D}_{A,C}$ in Definition 4 is well-defined in the sense that the right marginal of $\mathcal{D}_{A,C}$ is identical to \mathcal{D}_C . Therefore, there is no ambiguity in referring to both the input distribution and the right marginal of the output distribution by the same letter C. This operation receives $\mathcal{D}_{X,Z}$ and \mathcal{D}_C , and outputs a joint distribution $\mathcal{D}_{A,C}$ whose right marginal is \mathcal{D}_C , and $\mathcal{D}_{A|C=z} = \mathcal{D}_{X|Z=z}$ for all $z \in \mathcal{Z}$. Using the do-operation, Un can be defined equivalently as follows:

Definition 5 (Unique Information, equivalent definition). For $y \in \mathcal{Y}$ let C_y be a random variable with distribution $\mathcal{D}_{C_y} = \mathcal{D}_{Z|Y=y}$, and let $\mathcal{D}_{A_y,C_y} = do(\mathcal{D}_{X,Z}|\mathcal{D}_{C_y})$. The unique information from X to Z given Y is defined as:

$$\operatorname{Un}(X \to Z|Y) = \sum_{y \in \mathcal{Y}} \Pr(Y = y) I(A_y; C_y)$$

The proof of equivalence is simple, and is given in Appendix B. The idea behind Definition 5 is that by setting C = (Z|Y = y) for some $y \in \mathcal{Y}$ in Definition 4, and then by averaging the resulting mutual information values over all $y \in \mathcal{Y}$, we eliminate the effect of Y from the directional dependence between X and Z, without changing the directional dependence itself. A similar idea exists in Judea Pearl's do-calculus [20] (also [21]), where a node in a Bayesian network is set to a certain value while removing all incoming dependencies to that node, thereby distilling the causal relationship between that value and the remainder of the network.

C. Satisfaction of axioms and properties

To show that our definition satisfies the axioms and properties mentioned in Section II, we require the following technical lemma. The proof is given in Appendix C.

Lemma 1. Following the notations of Definition 1, we have that H(X|Z) = H(X'|Z) = H(X'|Z,Y), and that H(X') = H(X).

Corollary 1. Unique information (Def. 1) can be written as:

$$Un(X \to Z|Y) = I(X'; Z|Y) = X(X'|Y) - H(X'|Z, Y) = H(X'|Y) - H(X|Z).$$

Corollary 2. By Lemma 1, and since conditioning reduces entropy, we have that $H(X'|Y) \leq H(X') = H(X)$.

Based on the above lemmas and corollaries, we are in a position to prove that our definition of Un satisfies the required axioms.

1) Proof of Axiom 1, Information atoms relationship: Follows immediately from Definition 2 and Definition 3.

2) *Proof of Axiom 2, Commutativity:* First, Definition 1 and Definition 2 provide the following equivalent way for computing Red, which is proved in Appendix D.

Lemma 2. Redundant information (Def. 2) can alternatively be written as $\operatorname{Red}(X, Y \to Z) = I(X'; Y)$.

Similarly, by switching between X and Y in Definition 2 we have that $\operatorname{Red}(Y, X \to Z) = I(Y; Z) - \operatorname{Un}(Y \to Z|X)$; based on Lemma 2, this equals to I(X; Y'), where Y' is defined analogously to X' in (4), i.e.,

$$Pr(Y' = y, Z = z | X = x) = Pr(Y = y | Z = z) Pr(Z = z | X = x).$$
(6)

Then, we can conclude the commutativity of redundant information through the following lemma, which is proved in Appendix E.

Lemma 3 (Commutativity of Redundant Information). For X' and Y' as above, we have that I(X';Y) = I(X;Y').

Combining Lemma 2 and Lemma 3 readily implies the commutativity of Red, i.e., $\operatorname{Red}(X, Y \to Z) = \operatorname{Red}(Y, X \to Z)$.

3) Proof of Axiom 3, Monotonicity and self-redundancy: According to Definition 1, Un is nonnegative as conditional mutual information. Therefore, $\operatorname{Red}(X, Y \to Z) \leq I(X; Z)$ by Definition 2. Similarly, by Definition 2 we have $\operatorname{Red}(Y, X \to Z) = I(Y; Z) - \operatorname{Un}(Y \to Z|X)$, and hence $\operatorname{Red}(Y, X \to Z) \leq I(Y; Z)$. Since Red is symmetric by Axiom 2, it follows that

$$\operatorname{Red}(X, Y \to Z) \le \min\{I(X; Z), I(Y; Z)\}.$$

4) *Proof of Axiom 4, Nonnegativity:* We begin by showing that Red is nonnegative, for which we require the following lemma, proved in Appendix F.

Lemma 4. Unique information (Def. 1) is bounded from above by mutual information, i.e.,

 $\mathrm{Un}(X \to Z|Y) \le I(X;Z).$

Then, nonnegativity of Red follows from Lemma 4.

Corollary 3 (Nonnegativity of Redundant Information). *Redundant information (Def. 2) is nonnegative, i.e.,*

$$\operatorname{Red}(X, Y \to Z) \ge 0.$$

Finally, we remark that even though it is not a required axiom, non-negativity of Syn can be proved under an additional assumption as follows.

Remark 1 (Nonnegativity of Synergistic Information). Suppose H(Z|X,Y) = 0 (closed system assumption). Since $Un(X \rightarrow Z|Y) = I(X';Z|Y) \leq H(Z|Y)$, it follows from Definition 3 that $Syn(X, Y \rightarrow Z) \geq 0$.

5) *Proof of Property 1, Additivity:* The following lemma is proved in Appendix G.

Lemma 5 (Additivity of Unique Information). For two independent sets of variables X, Y, Z and $\overline{X}, \overline{Y}, \overline{Z}$, unique information (Def. 1) is additive:

$$Un((X,\bar{X}) \to (Z,\bar{Z})|(Y,\bar{Y})) = Un(X \to Z|Y) + Un(\bar{X} \to \bar{Z}|\bar{Y})$$
(7)

Since mutual information and conditional entropy are additive in the above sense, by Definition 2 and Definition 3, alongside Lemma 5, Red and Syn are additive as well.

6) *Proof of Property 2, Continuity:* We begin by showing that Red is continuous, for which we require the following lemma proved in Appendix H.

Lemma 6 (Continuity of Redundant Information). *The redundant information (Def. 2) is a continuous function of the input distribution* $\mathcal{D}_{X,Y,Z}$ *to* \mathbb{R} .

By Definition 2 and Definition 3, the continuity of Un and Syn can also be derived.

7) *Proof of Property 3, Independent Identity:* The following lemma is proved in Appendix I.

Lemma 7. The operator Red satisfies Property 3.

IV. DISCUSSION

In this paper, we proposed an explicit operational formula for PID, which is distinct from any existing approach, and proved that it satisfies all axioms and properties. In this section we provide further intuitive explanation for our approach.

First, we wish to elucidate the role that our do-operation plays in the definition of Un (Definition 5). In a sense, the do-operation can be understood as adjusting the marginal distribution of the Z variable of $\mathcal{D}_{X,Z}$, while impacting its connections with other variables as little as possible. This understanding can be confirmed by Lemma 1, which shows that the expected value of the conditional entropy after the do-operation retains its original value. This resembles the invariance implied in Shannon's communication model [1], where the conditional entropy of the output given the input is not affected by the input distribution. From this perspective, Z and X can be regarded as the input and output of the channel, respectively, that indicates their "relationship." The do-operation changes the distribution of the input Z, but does not change the channel's characteristic (i.e., H(X|Z)).

Based on this, Definition 5 realizes the intuition that unique information should represent the relationship between source variable and target variable given the other source variable. So, we use the do-operation to control the marginal distribution of the target variable Z to its conditional distribution given the value y of some source variable Y, and then use the expectation of mutual information $\sum_{y} \Pr(Y = y)I(A_y; C_y)$ to capture the "connection" between the specific source variable X and target variable Z given Y after the do-operation.

The reason this method can partition I(X; Z|Y) to Syn and Un (Def. 3), is that the do-operation eliminates highorder relations between Y and X, Z, i.e. Syn. Specifically, conditional mutual information relies on the joint conditional probability $\mathcal{D}_{(X,Z)|y}$, in expectation over all $y \in \mathcal{Y}$. This distribution includes both the conditional influence of Y on X, Z, but also has a simultaneous influence on the relationship between X and Z.

However, Definition 5 of unique information retains the relationship between X and Z without influence from Y by using the conditional probability $\mathcal{D}_{Z|y}$, in expectation over all $y \in \mathcal{Y}$, to perform the do-operation, which only reflects the conditional influence of Y on Z. Therefore, the expectation of mutual information $\sum_{y} \Pr(Y = y)I(A_y; C_y)$ can accurately quantify the unique information, which represents the pure conditional mutual relationship.

In addition to the above analysis of do-operations in unique information, Lemma 2 also brings another perspective worth discussing. Redundant information can be understood as the mutual information I(X';Y) (or I(X;Y')) obtained by changing the joint probability distribution $\mathcal{D}_{X,Y}$ according to $\mathcal{D}_{X,Y,Z}$ without changing the marginal distribution \mathcal{D}_X (or \mathcal{D}_Y) according to Lemma 1 (H(X') = H(X)).

As mentioned earlier, our definition of Syn might be negative, unless the system is closed (i.e., H(Z|X,Y) = 0, Remark 1). While Un and Red represent the information shared by one or two source variables with the target variable, Syn represents the information provided to the target variable by the "cooperation" of source variables. It is an accepted aphorism that cooperation does not necessarily increase outcome, and hence it might be the case that negative values of Syn conform with intuition. However, the reason why this explanation is no longer necessary in a closed system, as well as alternative interpretations of Syn that are nonnegative, remain to be studied.

Acknowledgments. The authors would like to thank an anonymous reviewer whose suggestions greatly simplified the paper. This research was supported by AFOSR grants FA9550-22-1-0054 and FA9550-23-1-0208.

REFERENCES

- Claude Elwood Shannon. A mathematical theory of communication. ACM SIGMOBILE mobile computing and communications review, 5(1):3–55, 2001.
- [2] Satosi Watanabe. Information theoretical analysis of multivariate correlation. *IBM Journal of research and development*, 4(1):66–82, 1960.
- [3] Paul L Williams and Randall D Beer. Nonnegative decomposition of multivariate information. arXiv preprint arXiv:1004.2515, 2010.
- [4] Robin AA Ince. The partial entropy decomposition: Decomposing multivariate entropy and mutual information via pointwise common surprisal. arXiv preprint arXiv:1702.01591, 2017.
- [5] Pedro AM Mediano, Fernando Rosas, Robin L Carhart-Harris, Anil K Seth, and Adam B Barrett. Beyond integrated information: A taxonomy of information dynamics phenomena. *arXiv preprint arXiv:1909.02297*, 2019.
- [6] Aobo Lyu, Bing Yuan, Ou Deng, Mingzhe Yang, Andrew Clark, and Jiang Zhang. System information decomposition. arXiv preprint arXiv:2306.08288, 2023.
- [7] Thomas F Varley. Generalized decomposition of multivariate information. arXiv preprint arXiv:2309.08003, 2023.
- [8] Virgil Griffith, Edwin KP Chong, Ryan G James, Christopher J Ellison, and James P Crutchfield. Intersection information based on common randomness. *Entropy*, 16(4):1985–2000, 2014.
- [9] Robin AA Ince. Measuring multivariate redundant information with pointwise common change in surprisal. *Entropy*, 19(7):318, 2017.
- [10] Nils Bertschinger, Johannes Rauh, Eckehard Olbrich, and Jürgen Jost. Shared information—new insights and problems in decomposing information in complex systems. In *Proceedings of the European conference* on complex systems 2012, pages 251–269. Springer, 2013.
- [11] Malte Harder, Christoph Salge, and Daniel Polani. Bivariate measure of redundant information. *Physical Review E*, 87(1):012130, 2013.
- [12] Nils Bertschinger, Johannes Rauh, Eckehard Olbrich, Jürgen Jost, and Nihat Ay. Quantifying unique information. *Entropy*, 16(4):2161–2183, 2014.
- [13] Pradeep Kr Banerjee, Johannes Rauh, and Guido Montúfar. Computing the unique information. In 2018 IEEE International Symposium on Information Theory (ISIT), pages 141–145. IEEE, 2018.
- [14] Elad Schneidman, William Bialek, and Michael J Berry. Synergy, redundancy, and independence in population codes. *Journal of Neuroscience*, 23(37):11539–11553, 2003.

- [15] Thomas F Varley, Maria Pope, Maria Grazia, Joshua, and Olaf Sporns. Partial entropy decomposition reveals higher-order information structures in human brain activity. *Proceedings of the National Academy of Sciences*, 120(30):e2300888120, 2023.
- [16] Borzoo Rassouli, Fernando E Rosas, and Deniz Gündüz. Data disclosure under perfect sample privacy. *IEEE Transactions on Information Forensics and Security*, 15:2012–2025, 2019.
- [17] Faisal Hamman and Sanghamitra Dutta. Demystifying local and global fairness trade-offs in federated learning using partial information decomposition. arXiv preprint arXiv:2307.11333, 2023.
- [18] Fernando E Rosas, Pedro AM Mediano, Henrik J Jensen, Anil K Seth, Adam B Barrett, Robin L Carhart-Harris, and Daniel Bor. Reconciling emergences: An information-theoretic approach to identify causal emergence in multivariate data. *PLoS computational biology*, 16(12):e1008289, 2020.
- [19] Judea Pearl. Causal diagrams for empirical research. *Biometrika*, 82(4):669–688, 1995.
- [20] Judea Pearl. Causality. Cambridge University Press, 2009.
- [21] Erik P Hoel, Larissa Albantakis, and Giulio Tononi. Quantifying causal emergence shows that macro can beat micro. *Proceedings of the National Academy of Sciences*, 110(49):19790–19795, 2013.
- [22] Paul L Williams. Information dynamics: Its theory and application to embodied cognitive systems. PhD thesis, Indiana University, 2011.
- [23] Virgil Griffith and Christof Koch. Quantifying synergistic mutual information. In *Guided self-organization: inception*, pages 159–190. Springer, 2014.
- [24] Imre Csiszár. Axiomatic characterizations of information measures. *Entropy*, 10(3):261–273, 2008.
- [25] Joseph T Lizier, Benjamin Flecker, and Paul L Williams. Towards a synergy-based approach to measuring information modification. In 2013 IEEE Symposium on Artificial Life (ALIFE), pages 43–51. IEEE, 2013.
- [26] Johannes Rauh, Pradeep Kr Banerjee, Eckehard Olbrich, Guido Montúfar, and Jürgen Jost. Continuity and additivity properties of information decompositions. *International Journal of Approximate Reasoning*, 161:108979, 2023.
- [27] Conor Finn and Joseph T Lizier. Pointwise partial information decompositionusing the specificity and ambiguity lattices. *Entropy*, 20(4):297, 2018.
- [28] Virgil Griffith and Tracey Ho. Quantifying redundant information in predicting a target random variable. *Entropy*, 17(7):4644–4653, 2015.
- [29] Artemy Kolchinsky. A novel approach to the partial information decomposition. *Entropy*, 24(3):403, 2022.

APPENDIX

A. Proof of the completeness of Definition 4.

In this part, we will show that do-operation's output is a probability distribution with the same marginal distribution as its input.

Lemma 8. For $\mathcal{D}_{X,Z}$ and \mathcal{D}_C as in Definition 4, the output $\Pr(A, C = x, z)$ of (5) describes a probability distribution, *i.e.*,

$$0 \le \Pr(A = x, C = z) \le 1, \text{ and}$$
$$\sum_{z \in \mathcal{X} \times \mathcal{Z}} \Pr(A = x, C = z) = 1.$$

Furthermore, the marginal distribution \mathcal{D}_C of the output $\mathcal{D}_{A,C}$ is equal to the input (call it $\mathcal{D}_{C'}$ in this section), i.e.,.

$$\sum_{x \in \mathcal{X}} \Pr(A = x, C = z) = \Pr(C' = z).$$

Proof. We begin by showing $0 \leq \Pr(A, C = (x, z)) \leq 1$. By Definition 4,

$$Pr(A = x, C = z)$$

= Pr(X = x|Z = z) Pr(C' = z). (8)

Since both terms in (8) are between 0 and 1, so is Pr(A = x, C = z).

We continue by showing that

$$\sum_{x,z} \Pr(A = x, C = z)$$

$$\stackrel{\text{Def. 4}}{=} \sum_{x,z} \Pr(X = x | Z = z) \Pr(C' = z)$$

$$= \sum_{z \in \mathcal{Z}} \Pr(C' = z) \sum_{x \in \mathcal{X}} \Pr(X = x | Z = z)$$

$$= \sum_{z \in \mathcal{Z}} \Pr(C' = z) = 1.$$
(9)

In (9), we can also conclude that $\sum_{x \in \mathcal{X}} \Pr(A = x, C = z) = \Pr(C' = z)$, which implies that the input $\mathcal{D}_{C'}$ is equal to the marginal distribution \mathcal{D}_C of the output $\mathcal{D}_{A,C}$.

B. Proof of the equivalence of Definitions 1 and 5.

Lemma 9. Definition 1 and Definition 5 are equivalent.

Proof. For the purpose of the proof, let

$$\operatorname{Un}(X \to Z|Y) = \sum_{y \in \mathcal{Y}} \Pr(Y = y) I(A_y; C_y), \text{ and } (10)$$

$$\operatorname{Un}'(X \to Z|Y) = I(X';Z|Y). \tag{11}$$

where $\mathcal{D}_{C_y} = \mathcal{D}_{Z|Y=y}$. By Definition 4 we have $\mathcal{D}_{A_y,C_y} = do(\mathcal{D}_{X,Z}|\mathcal{D}_{C_y})$, i.e.,

$$\Pr(A_y = x, C_y = z) = \Pr(X = x | Z = z) \Pr(Z = z | Y = y)$$

$$\stackrel{(4)}{=} \Pr(X' = x, Z = z | Y = y).$$

Therefore, we have

$$(10) = \sum_{y \in \mathcal{Y}} \Pr(Y = y) I(X', Z | Y = y)$$
$$= I(X', Z | Y) = \operatorname{Un}'(X \to Z | Y).$$

C. Proof of Lemma 1.

Proof. By Definition 1, for every $x, y, z \in \mathcal{X} \times \mathcal{Y} \times \mathcal{Z}$, we have

$$Pr(X' = x, Z = z | Y = y) =$$

= Pr(X = x | Z = z) Pr(Z = z | Y = y), (12)

and therefore

$$\frac{\Pr(X' = x, Z = z | Y = y)}{\Pr(Z = z | Y = y)} = \Pr(X = x | Z = z),$$

which can be simplified to

$$\Pr(X' = x | Z = z, Y = y) = \Pr(X = x | Z = z).$$

Therefore, we have that H(X'|Z = z, Y = y) = H(X|Z = z) for every $y, z \in \mathcal{Y} \times \mathcal{Z}$, which implies that H(X'|Z, Y) = H(X|Z).

Also, from (12) we have

$$Pr(X' = x, Z = z, Y = y)$$

= Pr(X = x|Z = z) Pr(Z = z, Y = y),

and summation over all $y \in \mathcal{Y}$ yields

$$Pr(X' = x, Z = z) = Pr(X = x | Z = z) Pr(Z = z)$$

= Pr(X = x, Z = z), (13)

which readily implies that H(X'|Z = z) = H(X|Z = z) for every $z \in \mathbb{Z}$, and hence H(X'|Z) = H(X|Z). Also, by (13), we have

$$Pr(X' = x) = \sum_{z \in \mathcal{Z}} Pr(X' = x, Z = z)$$
$$= \sum_{z \in \mathcal{Z}} Pr(X = x, Z = z)$$
$$= Pr(X = x),$$

which implies that H(X') = H(X).

D. Proof of Lemma 2.

Proof. By Definition 2, we have:

$$\operatorname{Red}(X, Y \to Z) = I(X, Z) - \operatorname{Un}(X \to Z|Y)$$
$$= (I(X, Z) + H(X|Z)) - (\operatorname{Un}(X \to Z|Y) + H(X|Z))$$
$$= H(X) - (\operatorname{Un}(X \to Z|Y) + H(X|Z))$$
(14)

Since Corollary 1 states that

$$\operatorname{Un}(X \to Z|Y) = H(X'|Y) - H(X|Z),$$

it follows that

$$(14) = H(X) - H(X'|Y)$$

= $^{\text{Lem. 1}} H(X') - H(X'|Y) = I(X';Y). \square$

E. Proof of Lemma 3.

Proof. To prove that I(X';Y) = I(X;Y'), it suffices to show that Pr(X' = x, Y = y, Z = z) = Pr(X = x, Y' = y, Z = z) for all $x, y, z \in \mathcal{X} \times \mathcal{Y} \times \mathcal{Z}$. We have

$$Pr(X' = x, Y = y, Z = z)$$

= Pr(X' = x, Z = z|Y = y) Pr(Y = y)
$$\stackrel{(4)}{=} Pr(X = x|Z = z) Pr(Z = z|Y = y) Pr(Y = y)$$

= Pr(X = x|Z = z) Pr(Z = z, Y = y)
= Pr(X = x|Z = z) Pr(Y = y|Z = z) Pr(Z = z), (15)

in which X and Y are symmetric. The proof is concluded by following similar steps in reversed order, i.e.,

$$\begin{aligned} (15) &= \Pr(X = x, Z = z) \Pr(Y = y | Z = z) \\ &= \Pr(Z = z | X = x) \Pr(Y = y | Z = z) \Pr(X = x) \\ &\stackrel{(6)}{=} \Pr(Y' = y, Z = z | X = x) \Pr(X = x) \\ &= \Pr(X = x, Y' = y, Z = z). \end{aligned}$$

F. Proof of Lemma 4.

Proof. By Corollary 1, Lemma 4 is equivalent to

$$H(X'|Y) - H(X|Z) \le I(X,Z)$$
$$H(X'|Y) \le H(X),$$

which coincides with Corollary 2, and the proof follows. \Box

G. Proof of Lemma 5.

Proof. By Definition 1, we have:

$$Un((X, X) \to (Z, Z)|(Y, Y)) = I((X, \bar{X})'; (Z, \bar{Z})|(Y, \bar{Y}))$$
(16)

where

$$Pr((X, \bar{X})', (Z, \bar{Z}) = (x, \bar{x}), (z, \bar{z})|(Y, \bar{Y}) = (y, \bar{y}))$$

=
$$Pr((X, \bar{X}) = (x, \bar{x})|(Z, \bar{Z}) = (z, \bar{z}))$$

$$\cdot Pr((Z, \bar{Z}) = (z, \bar{z})|(Y, \bar{Y}) = (y, \bar{y})).$$
(17)

Since X, Y, Z are independent from $\overline{X}, \overline{Y}, \overline{Z}$, we have

$$\begin{aligned} (17) &= \Pr(X = x | Z = z) \Pr(\bar{X} = \bar{x} | \bar{Z} = \bar{z}) \\ &\cdot \Pr(Z = z | Y = y) \Pr(\bar{Z} = \bar{z} | \bar{Y} = \bar{y}) \\ &\stackrel{(4)}{=} \Pr(X' = x, Z = z | Y = y) \Pr(\bar{X}', \bar{Z} = \bar{x}, \bar{z} | \bar{Y} = \bar{y}), \end{aligned}$$

which implies that X', Z|Y = y and $\bar{X}', \bar{Z}|\bar{Y} = \bar{y}$ are also independent for every $y, \bar{y} \in \mathcal{Y} \times \bar{\mathcal{Y}}$. Therefore, we have

$$= I(X', Z|Y) + I(\bar{X}', \bar{Z}|\bar{Y})$$

= Un(X' \rightarrow Z|Y) + Un(\bar{X}' \rightarrow \bar{Z}|\bar{Y})

H. Proof of Lemma 6.

Proof. Recall that Lemma 2 states that $\operatorname{Red}(X, Y \to Z) = I(X'; Y)$. Therefore, since I(X'; Y) is a continuous function of $\mathcal{D}_{X',Y}$, it suffices to prove that the mapping $\mathcal{F}(\mathcal{D}_{X,Y,Z}) = \mathcal{D}_{X',Y}$ is continuous, which holds by (4).

I. Proof of Lemma 7.

Proof. By Def. 1, we have $Un(X \to Z|Y) = I(X';Z|Y)$, where

$$Pr(X' = x, Z = z | Y = y) = Pr(X = x | Z = z) Pr(Z = z | Y = y).$$
(18)

Since Z = (X, Y), Eq. (18) is zero whenever $z \neq (x, y)$, and otherwise

$$(18) = \Pr(X = x | (X, Y) = (x, y)) \Pr((X, Y) = (x, y) | Y = y)$$

= $\Pr(X = x | Y = y).$ (19)

Also, since X and Y are independent, we have

$$(19) = \Pr(X = x).$$

Therefore, we have

$$\mathrm{Un}(X\to Z|Y)=I(X';Z|Y)=H(X)=I(X;Z),$$

which by Definition 2 implies that

$$\operatorname{Red}(X, Y \to Z) = I(X; Z) - \operatorname{Un}(X \to Z|Y) = 0. \quad \Box$$