Clique-free t-matchings in degree-bounded graphs*

Katarzyna Paluch^a, Mateusz Wasylkiewicz^a

^aInstitute of Computer Science, University of Wrocław, Wrocław, Poland

Abstract

We consider problems of finding a maximum size/weight t-matching without forbidden subgraphs in an undirected graph G with the maximum degree bounded by t+1, where t is an integer greater than 2. Depending on the variant forbidden subgraphs denote certain subsets of t-regular complete partite subgraphs of G. A graph is complete partite if there exists a partition of its vertex set such that every pair of vertices from different sets is connected by an edge and vertices from the same set form an independent set. A clique K_{t+1} and a bipartite clique $K_{t,t}$ are examples of complete partite graphs. These problems are natural generalizations of the triangle-free and square-free 2-matching problems in subcubic graphs. In the weighted setting we assume that the weights of edges of G are V-ertex-induced on every forbidden subgraph. We present simple and fast combinatorial algorithms for these problems. The presented algorithms are the first ones for the weighted versions, and for the unweighted ones, are faster than those known previously. Our approach relies on the use of gadgets with so-called V-edges. A V-edge of edge V-edge is, informally speaking, a half of V-econtaining exactly one of its endpoints.

Keywords: restricted t-matching, partite complete graph, combinatorial algorithm, half-edge, gadget

1. Introduction

We consider several variants of the problem of finding a maximum size or weight t-matching without forbidden subgraphs.

Given a positive integer t, a subset M of edges of an undirected simple graph is a t-matching if every vertex is incident to at most t edges of M. t-matchings belong to a wider class of b-matchings, where for every vertex v in the set of vertices V of the graph, we are given a natural number b(v) and a subset of edges is a b-matching if every vertex is incident to at most b(v) of its edges. A b-matching of maximum size/weight can be found in polynomial time by a reduction to a classical matching.

In the problems that we study in this paper we are given an undirected graph G=(V,E) in which each vertex has degree at most t+1 and the goal is to find a maximum size/weight t-matching that does not contain certain complete t-regular partite graphs. A graph $H=(V_H,E_H)$ is **complete partite** if there exists a partition V_1,V_2,\ldots,V_p of V_H of size $p\geq 2$ such that $E_H=\{(v,u):\exists i\neq j\ v\in V_i\wedge u\in V_j\}$. Each such V_i is called a **color class** of H. We denote the i-th color class of H by $V_i(H)$. If all color classes of H have the same size, we denote H by K_q^p where $q=|V_1|$. Notice that K_q^p is t-regular for t=(p-1)q. Observe that a clique K_a consisting of a vertices and a bipartite clique $K_{a,b}$ with color classes consisting of a and b vertices are special cases of complete partite graphs. We say that a t-matching of G is **restricted** if it does not contain an edge set of any K_{t+1} and $K_{t,t}$ of G and that it is K_q^p -free for t=(p-1)q if it does not contain an edge set of any subgraph of G isomorphic to t-regular K_q^p . The restricted t-matching problem and the K_q^p -free t-matching problem consist in finding a maximum size restricted/ K_q^p -free t-matching, respectively. Depending on the variant of the problem, we refer to respective subgraphs K_q^p , K_{t+1} and $K_{t,t}$ as **forbidden** subgraphs. We say that the restricted/ K_q^p -free t-matching problem is **bounded** if every vertex of the input graph is assumed to have degree at most t+1. Polynomial time algorithms for the bounded restricted t-matching

^{*}Partially supported by Polish National Science Center grant 2018/29/B/ST6/02633.

Email addresses: abraka@cs.uni.wroc.pl (Katarzyna Paluch), mateusz.wasylkiewicz@cs.uni.wroc.pl (Mateusz Wasylkiewicz)

problem and the bounded K_q^p -free t-matching problem were presented by Bérczi and Végh [2] and Kobayashi and Yin [18], respectively.

In the weighted versions of these problems, each edge e is associated with a nonnegative weight w(e) and we are interested in computing a restricted/ K_q^p -free t-matching of maximum weight. For a subgraph H of G, we say that the weight function is **vertex-induced on** H if real (possibly negative) values called **potentials** can be assigned to the vertices of H in such a way that the weight of every edge of H is equal to the sum of the potentials of its endpoints. Vornberger [27] showed that the weighted restricted 2-matching problem is \mathcal{NP} -hard. However, the weighted **bounded** restricted 2-matching problem is solvable in polynomial time when the weights of the edges are vertex-induced on every subgraph isomorphic to a square, which was shown by Paluch and Wasylkiewicz [23].

1.1. Our results

We present simple combinatorial algorithms for the weighted and unweighted versions of both the bounded restricted t-matching problem and the bounded K_q^p -free t-matching problem, both for $t \geq 3$. In the weighted setting we assume that the weights of the edges are nonnegative and vertex-induced on every forbidden subgraph. In these algorithms, instead of calculating restricted or K_q^p -free t-matching directly, we calculate its complement first. Such complement has fewer edges than sought-after restricted or K_q^p -free t-matching which results in faster running time of the algorithm. To accomplish this, we augment some forbidden subgraphs of the input graph with gadgets containing so-called half-edges and define a function b on the set of vertices in such a way that, any b-matching in the thus constructed graph G' yields a complement of restricted or K_q^p -free t-matching. Half-edges have already been introduced in [21] and used in several subsequent papers. The presented algorithms are the first ones for the weighted versions of these problems. The running time of our algorithms is $\mathcal{O}(\min\{nm\log n, n^3\})$. Moreover, for the unweighted version, our algorithms are faster than those known previously. The algorithm for the unweighted bounded restricted t-matching problem given by Bérczi and Végh [2] runs in $\mathcal{O}(t^3n + nm\log m)$ time whereas the algorithm for the unweighted bounded K_q^p -free t-matching problem given by Kobayashi and Yin [18] runs in $\mathcal{O}(pq^3n + nm\log m)$ time. For the unweighted versions of both problems, the running times of our algorithms are $\mathcal{O}(\sqrt{nm})$. Previous approaches to these problems relied on the shrinking of parts of or subsets of forbidden subgraphs and because of this do not lend themselves to the weighted setting.

The algorithms presented in this paper extend and generalize those from [23]. The generalizations require some new ideas as well as making observations regarding the structure. It turns out that the assumption $t \geq 3$ allows us to simplify some arguments. The most involved is the case of q=2 in the bounded K_q^p -free t-matching problem. Notice that this case was considered separately also in the previous algorithm for this problem given by Kobayashi and Yin [18].

1.2. Related work

Restricted 2-matchings and restricted t-matchings are classical problems of combinatorial optimization. Notice that a 2-matching is a set of vertex-disjoint cycles and paths whereas a restricted 2-matching is a 2-matching whose every cycle has length at least five since K_3 is a triangle, i.e. cycle of length three, and $K_{2,2}$ is a square, i.e. cycle of length four. Therefore, the restricted 2-matching problem can be used to approximate some variants of the travelling salesman problem (see [7] for details). Hartvigsen gave a complicated algorithm for the problem of finding a maximum size triangle-free 2-matching. Papadimitriou [4] showed that it is \mathcal{NP} -hard to find a maximum size 2matching without any cycle of length at most five. Recently, Kobayashi [16] gave a polynomial algorithm for finding a maximum weight 2-matching that does not contain any triangle from a given set of forbidden edge-disjoint triangles. Polynomial algorithms for the square-free and/or triangle-free 2-matchings in subcubic graphs were presented in [1, 2, 12–14, 18, 23]. The weighted version of the restricted 2-matching problem is \mathcal{NP} -hard for general weights, even in graphs which are both subcubic and bipartite (see [1]), therefore an assumption is made in all of these algorithms that the weight function is vertex-induced on every forbidden square. Such weight functions can be seen as a generalization of weight functions which assign values to the vertices instead of the edges. Regarding the squarefree 2-matching problem in general graphs, Nam [20] constructed a complex algorithm for it for graphs, in which all squares are vertex-disjoint. Finding a polynomial algorithm for the square-free 2-matching problem in general graphs is an open problem.

A natural generalization of the restricted 2-matching problem is the restricted t-matching problem. Since the weighted version of the restricted 2-matching problem is \mathcal{NP} -hard already, in the weighted version of the restricted

t-matching problem an assumption is typically made that the weight function is vertex-induced on every forbidden subgraph. Polynomial algorithms for restricted t-matchings in bipartite graphs were presented in [11, 19, 22, 24, 25]. In all previous algorithms for the weighted version of the restricted t-matching problem in bipartite graphs, an assumption is made that the weight function is vertex-induced on every forbidden $K_{t,t}$. It is an interesting question to generalize the known results for the square-free and/or triangle-free 2-matching problem in subcubic graphs to restricted t-matchings. Since all of these algorithms used the fact that every node of a forbidden subgraph is incident to at most one edge which does not belong to this forbidden subgraph, it is natural to consider restricted t-matching problem in graphs of the maximum degree at most t+1, i.e. the bounded restricted t-matching problem. As stated before, the unweighted version of this problem was considered by Bérczi and Végh [2].

It is worth mentioning that the bounded restricted t-matching problem has some connections to the (vertex)connectivity augmentation problem in which we want to make a given undirected graph G k-connected by adding a minimum number of different new edges. Let n be the number of the vertices of G. As pointed out by Bérczi and Végh [2], this problem is equivalent to finding a maximum size t-matching which contains no complete bipartite subgraphs $K_{a,b}$ consisting of a+b=t+2 vertices in the complement of G for t=n-k-1. A special case of the connectivity augmentation problem is increasing connectivity by one problem when given graph G is already (k-1)-connected. There exists a polynomial algorithm for this case given by Végh [26] which works in $\mathcal{O}(kn^7)$ time. Notice that the maximum degree of the complement of a (k-1)-connected graph is at most t+1 for t=n-k-1, therefore the increasing connectivity by one problem can be reduced to a problem of finding a maximum size t-matching which contains no complete bipartite subgraphs of size t+2 in the graphs of the maximum degree at most t+1. Weighted versions of these problems are also of interest since, in the same paper, Végh considered a generalization of the increasing connectivity by one problem when the edges of the complement of G are given weights and we want to minimize the total sum of added edges. This version is \mathcal{NP} -hard for general weights, however Végh presented a polynomial algorithm for the weighted version if the weights are vertex-induced on the whole complement of G.

It turns out that the polynomial solvability of the restricted t-matching problem has connections to jump systems. It was conjectured by Cunningham [5] that for every natural number k, the degree sequences of 2-matchings with no cycles of length at most k of any graph form a jump system if and only if there exists a polynomial algorithm for finding such 2-matchings of maximum size. The only open case of this conjecture is k=4. It was shown by Kobayashi et al. [17] that the degree sequences of restricted 2-matchings of any graph form a jump system which suggests that there should exist a polynomial algorithm for the restricted 2-matching problem in general graphs. A generalization of Cunningham's conjecture to the t-matchings seems to be also true. In the same paper, Kobayashi et al. proved that the degree sequences of restricted t-matchings in any graph form a jump system. In fact, they proved an even stronger result. They showed that the degree sequences of t-matchings with no subgraphs from a given family \mathcal{H} of t-regular connected graphs form a jump system in any graph if and only if every member of \mathcal{H} is a complete partite graph. Moreover, it was shown by Kobayashi and Yin [18] that this problem is \mathcal{NP} -hard, even in the graphs of maximum degree at most t+1, if \mathcal{H} is a singleton of any t-regular connected graph which is not complete partite. Notice that a t-regular complete partite graph is exactly K_q^p for t=(p-1)q, hence it is natural to consider K_q^p -free t-matchings for such values of t.

The weighted versions of restricted t-matchings were studied in the context of the M-concave functions on constant-parity jump systems, which can be seen as discrete counterparts of the submodular functions. Kobayashi et al. [17] proved that the degree sequences of maximum weight restricted t-matchings in a bipartite graph form an M-concave function on the constant-parity jump system if and only if the weight function is vertex-induced on every forbidden subgraph. The algorithm of Bérczi and Kobayashi [1] for maximum weight square-free 2-matchings in subcubic graphs is based on the fact that the degree sequences of these 2-matchings form an M-concave function in any subcubic graph if the weight function is vertex-induced on every square. On the other hand, the degree sequences of maximum weight triangle-free 2-matchings form an M-concave function in any graph, which was shown by Kobayashi [15]. However, it is unknown if there exists a polynomial algorithm for finding maximum weight triangle-free 2-matchings in general graphs.

2. Preliminaries

Let G = (V, E) be an undirected graph with vertex set V and edge set E. We denote the number of vertices of G by n and the number of edges of G by m. We assume that all graphs are simple, i.e., they contain neither loops

nor parallel edges. We denote an edge connecting vertices v and u by (v,u). For a subgraph H of G, we denote the vertex set of H by V(H) and the edge set by E(H). Given a weight function $w:E\to\mathbb{R}$, the **weight of** H, denoted by w(H), is defined as the sum of weights of edges of H. For an edge set $F\subseteq E$ and $v\in V$, we denote by $\deg_F(v)$ the number of edges of F incident to v. With a slight abuse of notation, for a graph H, we sometimes write $\deg_H(v)$ instead of $\deg_{E(H)}(v)$. For any natural number k, we say that G is k-regular if every vertex of G has degree exactly k. For a vertex set $A\subseteq V$, we denote by G[A] a subgraph of G induced by G[A] and G[A] and G[A] and G[A] and G[A] and G[A] are isomorphic by G[A] and G[A] are isomorphic by G[A] and G[A] are isomorphic by G[A] are isomorphic by G[A] and G[A] are isomorphic by G[A] and G[A] are isomorphic by G[A] are isomorphic by G[A] and G[A] are isomorphic by G[A] are isomorphic by G[A] and G[A] are isomorphic by G[A] are isomorphic by G[A] and G[A] are isomorphic by G[A] and G[A] and G[A] are isomorphic by G[A] are isomorphic by G[A] and G[A] are i

An instance of each of the two problems that we consider in the paper consists of an undirected graph G=(V,E) whose every vertex has degree at least one and at most t+1, and a weight function $w:E\to\mathbb{R}_{\geq 0}$. In the bounded K_q^p -free t-matching problem we are also given natural numbers $p\geq 2$, $q\geq 1$ such that t=(p-1)q. In the bounded restricted (resp. K_q^p -free) t-matching problem we assume that w is vertex-induced on every K_{t+1} and $K_{t,t}$ (resp. on every K_q^p) of G and the goal is to find a maximum weight restricted (resp. K_q^p -free) t-matching of G.

For a weight function $w: E \to \mathbb{R}$ which is vertex-induced on some subgraph H of G, we say that a function $r_H: V(H) \to \mathbb{R}$ which assigns a potential to every vertex of H is a **potential function** of H.

We say that an edge set $M \subseteq E$ is a **co-t-matching** of G if $E \setminus M$ is a t-matching of G. Additionally, we say that a co-t-matching $M \subseteq E$ is **covering** (resp. K_q^p -covering) if $E \setminus M$ is a restricted (resp. K_q^p -free) t-matching of G. We say that an edge set $M \subseteq E$ covers a forbidden subgraph H of G, or that H is covered by M, if M contains at least one edge of H. Notice that if the degree of every vertex of G is at most t+1, then an edge set $M \subseteq E$ is a co-t-matching of G if every vertex of G of degree t+1 is incident to at least one edge of M. Moreover, M is a K_q^p -covering (resp. covering) co-t-matching if, in addiction, M covers every K_q^p (resp. every K_{t+1} and $K_{t,t}$) of G. A K_q^p -)covering co-t-matching M of G is said to be **minimum weight** if there is no K_q^p -)covering co-t-matching K_q^p - of weight smaller than K_q^p -)covering co-t-matching K_q^p -covering co-t-matching K_q^p -matching K_q^p -covering co-t-matching K_q^p -covering co-t-matching

We will need to compute a b-matching of a graph G where we are given vectors $l,b \in \mathbb{N}^V$ and a weight function $w: E \to \mathbb{R}$. (We allow negative weights here.) For a vertex $v \in V$, [l(v),b(v)] is said to be a **capacity interval** of v. An edge set $M \subseteq E$ is said to be an (l,b)-matching if $l(v) \le \deg_M(v) \le b(v)$ for every $v \in V$. Given an (l,b)-matching M and an edge $e = (u,v) \in M$, we say that u is **matched to** v in M. Moreover, we say that a vertex v of G is **unmatched** in M if v is not incident to any edge of M. A **matching** and a **perfect matching** of G is any (l,b)-matching of G where the capacity interval of every vertex v of G is equal to, respectively, [0,1] and [1,1]. An (l,b)-matching M is said to be a **maximum weight** (l,b)-matching if there is no (l,b)-matching M' of G of weight greater than w(M). A maximum weight (l,b)-matching can be computed efficiently.

Theorem 1 ([9]). There is an algorithm that, given a multigraph G = (V, E) (i.e. G may contain parallel edges), a weight function $w : E \to \mathbb{R}$ and vectors $l, b \in \mathbb{N}^V$, finds a maximum weight (l, b)-matching of G in time $O\left(\left(\sum_{v \in V} b(v)\right) \min\{|E| \log |V|, |V|^2\}\right)$, assuming that G admits any (l, b)-matching.

We define a *minimum weight* (l,b)-matching of G analogously. Notice that calculating a minimum weight (l,b)-matching of G can be reduced to calculating a maximum weight (l,b)-matching of G by negating the weight of every edge of G.

3. Outline of the Algorithm

We give an outline of our algorithms for the weighted versions of the bounded restricted and the bounded K_q^p -free t-matching problem assuming that the weight function is vertex-induced on every forbidden subgraph. Recall that we assume that $t \geq 3$. In the K_q^p -free t-matching problem we additionally assume that t = (p-1)q.

Algorithm 1 Computing a maximum weight restricted or K_q^p -free t-matching of graph G of maximum degree at most t+1 given a weight function $w: E \to \mathbb{R}_{>0}$.

- 1: Construct an auxiliary multigraph G' = (V', E') consisting of O(n) vertices and O(m) edges by augmenting some forbidden subgraphs of G with gadgets containing half-edges. (Both gadgets and half-edges are defined later.)
- 2: Define a weight function $w': E' \to \mathbb{R}$ and vectors $l, b \in \mathbb{N}^{V'}$.
- 3: Compute a minimum weight (l, b)-matching M' of G'.
- 4: Construct a co-t-matching \bar{M} of G by replacing all half-edges of M' with some edges of G in such a way that $w(\bar{M}) \leq w'(M')$.
- 5: Add some edges of the remaining forbidden subgraphs to \bar{M} by replacing some of its edges with other ones without increasing the weight of \bar{M} .
- 6: Return $E \setminus \bar{M}$.

The precise construction of G', w', l and b for the bounded restricted (resp. K_q^p -free) t-matching problem is given in Section 4 (resp. Section 5).

For the bounded restricted t-matching problem, an implementation of Step 4 is given in Theorem 3 whereas an implementation of Step 5 consists of subsequent applications of Lemma 5. An implementation of these steps for the bounded K_a^p -free t-matching problem is given in Theorem 4.

Claim 1. Algorithm 1 runs in time $\mathcal{O}(\min\{nm\log n, n^3\})$ in the weighted variant and $\mathcal{O}(\sqrt{n}m)$ in the unweighted. We prove Claim 1 in Section 6.

4. The bounded restricted t-matching problem

In this section we address the weighted bounded restricted t-matching problem. Recall that we assume that $t \ge 3$ and the nonnegative weight function w is vertex-induced on every K_{t+1} and $K_{t,t}$ of G.

We build a graph G' = (V', E') together with a weight function $w' : E' \to \mathbb{R}$, in which subgraphs, called *gadgets*, are added to some forbidden subgraphs of G. The precise construction of gadgets is the following.

Let H be any K_{t+1} of G. We define a **gadget for** H. Let $v_1, v_2, \ldots, v_{t+1}$ be vertices of H. We introduce one new vertex u_H called a **subdivision vertex**. For each vertex v_i of H we add edge (v_i, u_H) called a **half-edge** of H. We set the capacity interval of u_H to [2, 2].




Figure 1: A gadget for K_{t+1} for t = 5.

Let H be any $K_{t,t}$ of G. We define a **gadget for** H. Let $a_1, a_2, \ldots, a_t \in V_1(H)$ and $b_1, b_2, \ldots, b_t \in V_2(H)$. We introduce two new subdivision vertices u_H^1 and u_H^2 . We connect u_H^1 with all vertices of $V_1(H)$ by half-edges. Symmetrically, we connect u_H^2 with all vertices of $V_2(H)$ by half-edges. We set the capacity intervals of u_H^1 and u_H^2 to [1,1].

The main ideas behind these gadgets are the following. An (l,b)-matching M' of G' is to represent roughly a covering co-t-matching \bar{M} of G. We want to ensure that at least one edge of H belongs to \bar{M} . Notice that M'

contains exactly two half-edges of H incident to some vertices z_1 and z_2 of H. We add an edge (z_1, z_2) to \bar{M} . Moreover, we add to \bar{M} all edges of H which belong to M'. In this way, H is guaranteed to be covered by \bar{M} .

Regarding weights w' of the edges in the gadgets, we assign them as follows. Let r_H be any potential function of H. Every half-edge of H incident to vertex v_i gets weight $r_H(v_i)$.




Figure 2: A gadget for $K_{t,t}$ for t=3.

4.1. Disjoint forbidden subgraphs

In this subsection we consider a case when all forbidden subgraphs of G are pairwise vertex-disjoint. We show how to drop this assumption later, in Subsection 4.2.

The precise construction of G', w', l and b is the following. We start off with G' = G. We add a gadget to every forbidden subgraph of G. This operation is well-defined because these forbidden subgraphs are pairwise vertex-disjoint. We set the capacity interval of every vertex v of degree t+1 of G to [1,t+1]. We set the capacity interval of every other vertex v of G to $[0,\deg_G(v)]$. We set the weight of every edge e of G' which belongs to G to w(e). In theorems below we show the correspondence between covering co-t-matchings of G and (l,b)-matchings of G'.

Theorem 2. Let \bar{M} be any covering co-t-matching of G. Then we can find an (l,b)-matching M' of G' such that $w'(M') = w(\bar{M})$.

Proof. We initialize M' as the empty set. We add every edge of \bar{M} that does not belong to any forbidden subgraph of G to M'.

Consider any $H=K_{t+1}$ of G. Since \bar{M} is covering, there exists an edge of H which belongs to \bar{M} . If more than one edge of H belongs to \bar{M} , we choose one of them. Suppose that we chose $(z_1,z_2)\in \bar{M}$. Then we add edges (z_1,u_H) and (z_2,u_H) to M'. We add every other edge of H which belongs to \bar{M} to M'.

Consider any $H = K_{t,t}$ of G. Since \overline{M} is covering, there exists an edge (a,b) of H which belongs to \overline{M} . Suppose that $a \in V_1(H)$ and $b \in V_2(H)$. We add edges (a,u_H^1) and (b,u_H^2) to M'. We add every other edge of H which belongs to \overline{M} to M'.

Since the weight of any replaced edge in G is equal to the sum of the weights of its corresponding half-edges in G', we get that $w'(M') = w(\bar{M})$.

Theorem 3. Let M' be any (l,b)-matching of G'. Then we can find a covering co-t-matching \bar{M} of G such that $w(\bar{M}) \leq w'(M')$.

Proof. We initialize \bar{M} as the empty set. We add every edge of M' which belongs to G to \bar{M} . For every forbidden subgraph of G we add some of its edges to \bar{M} .

Let H be any K_{t+1} of G. Notice that exactly two vertices of H, say v_1 and v_2 , are matched to u_H in M'. We add (v_1, v_2) to \bar{M} if it does not belong to \bar{M} already.

For any $H = K_{t,t}$ of G we proceed analogously, because in the gadget for H the subdivision vertices u_H^1 and u_H^2 are matched to two vertices $a \in V_1(H)$ and $b \in V_2(H)$.

4.2. Non-disjoint forbidden subgraphs

In the following lemmas we examine the ways in which forbidden subgraphs of G may overlap.

Lemma 1. Let H_1 and H_2 be any two different K_{t+1} 's of G with a common vertex. Then $H_1 \cap H_2 = K_t$.

Proof. Let v be a common vertex of H_1 and H_2 . It has at most t+1 neighbours and exactly t of them belong to H_1 and t to H_2 . Therefore, at least t-1 neighbours of v belong to both H_1 and H_2 , so H_1 and H_2 have at least t common vertices. Since H_1 and H_2 are different, they cannot have t+1 common vertices.

Lemma 2. Let H_1 and H_2 be any two different $K_{t,t}$'s of G with a common vertex. Then $H_1 \cap H_2$ contains $K_{t-1,t-1}$.

Proof. Let $v \in V_1(H_1)$ be a common vertex of H_1 and H_2 . W.l.o.g. we can assume that $v \in V_1(H_2)$ (otherwise we can rename the color classes of H_2). The vertex v is incident to exactly t edges of H_1 and t edges of H_2 , hence at least t-1 of these edges belong to $H_1 \cap H_2$. Let U denote their endpoints. Then $U \setminus \{v\} \subseteq V_2(H_1) \cap V_2(H_2)$ and $|U \setminus \{v\}| \ge t-1 \ge 1$. This implies that there exists a vertex $v' \in V_2(H_1) \cap V_2(H_2)$ and thus, by repeating the same argument, there are at least t-1 vertices in $V_1(H_1) \cap V_1(H_2)$.

Lemma 3. Let H_1 and H_2 be any, respectively, K_{t+1} and $K_{t,t}$ of G with a common vertex. Then t=3 and $H_1 \cap H_2 = K_{2,2}$.

Proof. Let v_1 be a common vertex of H_1 and H_2 . W.l.o.g we can assume that $v_1 \in V_1(H_2)$. Repeating the same argument as in the proofs of Lemma 1 and Lemma 2, we get that at least $t-1 \geq 1$ vertices of $V_2(H_2)$ belong to H_1 . Similarly, at least t-1 vertices of $V_1(H_2)$ belong to H_1 , so H_1 has at least H_2 vertices. Therefore, H_2 belong to H_3 so H_4 has exactly two common vertices with each color class of H_2 .

We define the following family of **problematic** subgraphs of G.

Definition 1. Let H be any forbidden K_{t+1} of G. We say that H is **unproblematic** if H has a common vertex with another forbidden subgraph H' of G such that:

1.
$$H' = K_{t+1}$$
 and $w(H) \le w(H')$, or

2.
$$H' = K_{t,t}$$
.

Otherwise, we say that H is **problematic**.

Definition 2. Let H be any forbidden $K_{t,t}$ of G. We say that H is unproblematic if H has a common vertex with another forbidden subgraph $H' = K_{t,t}$ of G such that $w(H) \leq w(H')$.

Otherwise, we say that H is **problematic**.

Lemma 4. Any two different problematic subgraphs of G are vertex-disjoint.

Proof. Consider any two different forbidden cliques H_1 and H_2 with a common vertex. We show that at least one of H_1, H_2 is unproblematic.

If both H_1 and H_2 are K_{t+1} 's and $w(H_1) \leq w(H_2)$, then H_1 is unproblematic. Similarly, if both H_1 and H_2 are $K_{t,t}$'s and $w(H_1) \leq w(H_2)$, then H_1 is unproblematic.

If
$$H_1 = K_{t+1}$$
 and $H_2 = K_{t,t}$, then by Point 2 of Definition 1, H_1 is unproblematic.

To find a minimum weight covering co-t-matching of G, we use the graph G' from Subsection 4.1. However, we only replace every problematic subgraph by its gadget. In the such constructed G' we compute an (l,b)-matching of G', which corresponds to a co-t-matching \bar{M} of G. The co-t-matching \bar{M} may not cover some unproblematic forbidden subgraphs. Below we show that it is possible to modify \bar{M} so that it covers every forbidden subgraph.

Lemma 5. Let \bar{M} be any co-t-matching of G. If \bar{M} does not cover some unproblematic subgraph H, then we can find a co-t-matching \bar{N} of G such that $w(\bar{N}) \leq w(\bar{M})$ which covers H and every forbidden subgraph covered by \bar{M} .

Proof. If H is a K_{t+1} of G sharing a vertex with some $H' = K_{t,t}$ of G, then by Lemma 3, t = 3, $H = K_4$, $H' = K_{3,3}$ of G and $H \cap H' = K_{2,2}$. Let $v_1 \in V(H) \cap V_1(H')$, $v_2 \in V(H) \cap V_2(H')$, $u_1 \in V_1(H') \setminus V(H)$ and $u_2 \in V_2(H') \setminus V(H)$. We set $\bar{N} = \bar{M} \setminus \{(v_1, u_2), (v_2, u_1)\} \cup \{(v_1, v_2), (u_1, u_2)\}$. Notice that the only edge of H' incident to u_1 or u_2 which can possibly not belong to \bar{M} is (u_1, u_2) since every vertex of H is incident to t = 3 edges of H which do not belong to \bar{M} . Thus, both u_1 and u_2 are incident to some edge of \bar{N} . Moreover, $w(\bar{N}) \leq w(\bar{M})$ because w is vertex-induced on H'.

If H is a K_{t+1} of G which does not share a vertex with any $K_{t,t}$ of G, then H shares a vertex with another $H'=K_{t+1}$ of G such that $w(H)\leq w(H')$. Let u (resp. u') be the only vertex of H (resp. H') which does not belong to H' (resp. H). Notice that every edge of H' incident to u' belongs to \bar{M} , so $\deg_{\bar{M}}(u')\geq t$. Since $w(H)\leq w(H')$, the sum of the weights of edges of H incident to u is not greater than the sum of weights of edges of H' incident to u'. Therefore, there exists a common vertex of H and H', say z, such that $w(u,z)\leq w(u',z)$. We define \bar{N} as \bar{M} with the edge (u',z) replaced by (u,z). As a result, \bar{N} covers H. It still covers H' since every vertex of $V(H)\cap V(H')$ different than z is matched to u' in \bar{N} . Notice that \bar{N} is still a co-t-matching since $\deg_{\bar{N}}(u')=\deg_{\bar{M}}(u')-1\geq t-1\geq 1$ since $t\geq 3$.

Let H be a $K_{t,t}$ of G sharing a vertex with another $H' = K_{t,t}$ of G such that $w(H) \leq w(H')$. If $H \cap H' = K_{t-1,t-1}$, we proceed analogously as in the case when $H = K_{t+1}$ shares a vertex with $H' = K_{t,t}$. Otherwise, $H \cap H' = K_{t,t-1}$ and we proceed analogously as in the case when $H = K_{t+1}$ shares a vertex with another $H' = K_{t+1}$.

5. The bounded K_q^p -free t-matching problem

In this section we consider the weighted bounded K_q^p -free t-matching problem. Recall that t=(p-1)q for this problem. Let us observe that for q=1, $K_q^p=K_{t+1}$ and for p=2, $K_q^p=K_{t,t}$. Since the bounded K_q^p -free t-matching problem for these cases can be solved by a slight modification of the algorithm for the bounded restricted t-matching problem, we assume that $q\geq 2$ and $p\geq 3$. Recall that we assume that the weight function w is vertexinduced on every subgraph of G isomorphic to K_q^p .

We denote the empty graph on n vertices by \hat{I}_n . For graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ such that V_1 and V_2 are disjoint, we say that a graph $G_1 \times G_2 = (V_1 \cup V_2, E_1 \cup E_2 \cup (V_1 \times V_2))$ is a **product** of G_1 and G_2 .

Similarly as in Section 4, we build a graph G' = (V', E') together with a weight function $w' : E' \to \mathbb{R}$, in which some K_a^p 's of G are augmented with gadgets.

5.1. Case $q \ge 3$

The case of $q \ge 3$ is a little simpler than that of q = 2. We consider it first.

Let H be any K_q^p of G. We define a **gadget for** H. We introduce p subdivision vertices $u_H^1, u_H^2, \ldots, u_H^p$ and a **global vertex** z_H . For every i, we connect u_H^i with every vertex of $V_i(H)$ by half-edges and with z_H (see Fig. 3). Let r_H be any potential function of H. A half-edge of H incident to vertex v_i gets weight $r_H(v_i)$. The remaining edges of the gadget get weight 0. We set the capacity interval of vertex z_H to [p-2, p-2] and of every subdivision vertex of the gadget for H to [1,1].




Figure 3: A gadget for K_q^p for p=3 and q=3.

Lemma 6. Let H_1 and H_2 be any two different K_q^p 's of G with a common vertex. Then $H_1 \cap H_2$ contains $I_{q-1} \times K_q^{p-1}$.

Proof. Let V_1, V_2, \ldots, V_p be the color classes of H_1 and $v \in V_1$ a common vertex of H_1 and H_2 . We first show that at most one vertex of H_1 does not belong to H_2 . Since v is adjacent to every vertex of V_2, V_3, \ldots, V_p via (p-1)q=t edges and $\deg_G(v) \leq t+1$, there exists at most one other edge incident to v. Exactly t of all edges incident to v belong to H_2 , therefore all vertices of V_2, V_3, \ldots, V_p except at most one belong to H_2 as well. Suppose that if there exists a vertex of H_1 which does not belong to H_2 , then it belongs to V_p . Let u be any vertex of V_2 . Since v is a vertex of v and hence v is v in the example of v in the exist v in v in the exist v in

Claim 2. If all vertices of one color class V_i belong to H_2 , then they belong to the same color class of H_2 .

Proof. Suppose to the contrary that two vertices $v_1, v_2 \in V_i$ belong to different color classes of H_2 . This means that H_2 contains an edge (v_1, v_2) . Let v_3 be any other vertex of V_i (it exists since $q \geq 3$). It cannot belong to the same color class of H_2 as both v_1 and v_2 . Thus it is adjacent to at least one of v_1, v_2 . Suppose that v_3 is a neighbour of v_1 . This means that v_1 is incident to t+2 edges of G: t edges of H_1 , t and t and t and t and t are contradiction.

By the above claim H_1 and H_2 have different vertex sets because otherwise they would be equal. We have already assumed that if H_1 contains a vertex not contained in H_2 , then it belongs to V_p . Thus by the above claim, we conclude that $V_1, V_2, \ldots, V_{p-1}$ are color classes of H_2 and q-1 vertices of V_p belong to H_2 , which proves that $H_1 \cap H_2 = I_{q-1} \times K_q^{p-1}$.

Definition 3. Let H be any K_q^p of G. We say that H is **unproblematic** if there exists another $H' = K_q^p$ of G that shares a vertex with H and such that $w(H) \leq w(H')$.

Lemma 7. Any two different problematic K_q^p 's of G are vertex-disjoint.

Proof. Consider any two different K_q^p 's H_1 and H_2 of G with a common vertex. By Lemma 6, $H_1 \cap H_2$ contains $I_{q-1} \times K_q^{p-1}$. Assume that $w(H_1) \leq w(H_2)$. Then H_1 fulfills Definition 1.

Theorem 4. For any K_q^p -covering co-t-matching \bar{M} of G there exists an (l,b)-matching M' of G' such that $w'(M') = w(\bar{M})$.

For any (l,b)-matching M' of G' there exists a K_q^p -covering co-t-matching \bar{M} of G such that $w(\bar{M}) \leq w'(M')$.

Proof. We prove the first sentence of Theorem 4 in the beginning. Let H be any K_q^p of G. Since \overline{M} covers H, some edge of H belongs to \overline{M} . If more than one edge of H belongs to \overline{M} , we choose one of them. Suppose that we chose $(a,b)\in \overline{M}$. Assume that $a\in V_1(H)$ and $b\in V_2(H)$. Then we add edges (a,u_H^1) and (b,u_H^2) to M'. Additionally, we add p-2 edges, (z_H,u_H^i) for every i>2, to M'. We add every other edge of H to M'.

We prove the second sentence of Theorem 4 now. Let H be any K_q^p of G. Notice that exactly two subdivision vertices of the gadget for H are matched to vertices of G in M', because z_H is matched to exactly p-2 other subdivision vertices. Assume that u_H^1 and u_H^2 are not matched to z_H in M'. Therefore, they are matched to some vertices a and b, respectively, of G. We add an edge (a,b) and every other edge of H which belongs to M' to M. Hence, M covers every problematic K_q^p of G.

However, \bar{M} may not cover some unproblematic K_q^p 's of G. Let H by any such K_q^p . From Definition 3, there exists a $H' = K_q^p$ of G such that $H \cap H' = I_{q-1} \times K_q^{p-1}$ and $w(H) \leq w(H')$. Let u (resp. u') be only vertex of H (resp. H') which does not belong to H' (resp. H). Rest of the proof is identical to covering unproblematic K_{t+1} which shares some vertex with another K_{t+1} of G in the proof of Lemma 5.

5.2. Case q = 2

Next we consider the case of q=2. Notice that t=2p-2 in this case. It turns out that two different K_2^p 's of G can have the same set of vertices.

Lemma 8. Let H_1 and H_2 be two K_2^p 's of G with a common vertex. Then either:

1.
$$H_1 \cap H_2 = I_1 \times K_2^{p-1}$$
, or

2. H_1 and H_2 have the same set of vertices and at least two color classes of H_1 induce an edge in G.

Proof. Observe that the first paragraph of the proof of Lemma 6 holds in this case because we did not assume that $q \ge 3$ there. Therefore, $H_1 \cap H_2$ contains $I_{q-1} \times K_q^{p-1}$. If $H_1 \cap H_2 = I_{q-1} \times K_q^{p-1}$, we are done.

If H_1 and H_2 have the same set of vertices, we need to show that for some two different color classes of H_1 , its vertices are connected via an edge of G. Since H_1 and H_2 are different, there exists a color class of H_1 consisting of vertices v_1 and v_2 which belong to different color classes of H_2 . Since an edge (v_1, v_2) belongs to H_2 , the color class of H_1 consisting of v_1 and v_2 is one of the color classes we are looking for. Let u be the other vertex which belongs to the same color class of H_2 as v_1 . The color class of H_1 to which u belongs is the second class from the claim. \square

Definition 4. Let A be any subset of 2p vertices of G. We say that G[A] is **dense** if it contains at least two different forbidden K_2^p 's. The set $C_{G[A]} = \{a \in A : N(a) \subset A \land |N(A)| = t+1\}$ is called the **core** of G[A].

Lemma 9. The size of the core of any dense subgraph of G is even and at least four.

Proof. Consider two different K_2^p 's H_1 and H_2 with the same vertex set A. Notice that the core of A is equal to the sum of color classes of H_1 whose induce an edge in G, so its size is even. Moreover, its size is at least four, because by Lemma 8, there are at least two such color classes.

To guarantee that every K_2^p contained in a dense subgraph of G is covered by the computed co-t-matching, we introduce a separate gadget for a dense subgraph. These gadgets are used alongside gadgets for problematic K_2^p 's. (We generalize Definition 3 and the gadget for K_q^p to the case q=2 in a straightforward manner.) In the following lemma we prove that any dense subgraph of G is vertex-disjoint both with any other dense subgraph and with any other problematic K_2^p 's of G.

Lemma 10. Let H be any dense graph of G. Then any K_2^p of G which shares a vertex with H has the same set of vertices as H.

Proof. Let H' be any K_2^p of G which has a common vertex with H. Suppose that the set of vertices of H' is different than V(H). Since there exists some K_2^p of H, we can use Lemma 8 and conclude that all but one vertex of H' belongs to H. Let $v \in V(H') \setminus V(H)$. Notice that vertex v is adjacent to at least 2p-2 vertices of H. By Lemma 9, the core of H consists of at least four vertices, which means that at least two of these vertices are adjacent to v - a contradiction with the definition of the core.

The following observation describes all K_2^p 's contained in a dense subgraph of G.

Observation 1. Consider any dense subgraph H of G with the core C_H . Let H' be any K_2^p of H. Then the color classes of any K_2^p of H consist of the color classes of H' which are not contained in C_H and some partition of C_H into two element subsets. Moreover, any such partition of C_H forms the color classes of exactly one K_2^p of H.

Lemma 11. The weight function w is vertex-induced on any dense subgraph H of G. Moreover, the potential function of any K_2^p of H is also a potential function r_H of H.

Proof. We show that the potential functions of any two different K_2^p 's of H are the same. It ends proof since, by Observation 1, every edge of H belongs to some K_2^p of H.

Let H_1 and H_2 be any two different K_2^p 's of G. Let r_1 and r_2 be any potential funcions of H_1 and H_2 , respectively. Consider any vertex v of H. Assume that $v \in V(H_1)$. We show that there exists a triangle (v, v_1, v_2) which belongs to both H_1 and H_2 . In the light of Observation 9 it implies that $r_1(v) = r_2(v)$.

Set v_1 to any vertex of $V_2(H_1)$ which belongs to other color class of H_2 than v. If there exists some vertex of $V_3(H_1)$ (this color class exists since $p \geq 3$) which belongs to the same color class of H_2 as neither v nor v_1 , we choose it as v_2 .

Otherwise, both vertices of $V_3(H_1)$ belong to the same color class of H_2 as v and v_1 , respectively. In this case we choose v_2 in such a way that it belongs to the same color class of H_2 as v_1 and we change v_1 to another vertex of $V_2(H_1)$. Since q=2, v_1 has to be in different color class of H_2 than v and v_2 .

To construct a gadget for a dense subgraph of G we make use of the following lemma. In our applications, M will typically be a set of edges of a co-t-matching of G which belong to H.

Lemma 12. Let H be any dense subgraph of G with the core C_H of size 2k. Assume that every vertex of C_H is incident to at least one edge of a subset $M \subseteq E(H)$. Then the following statements are equivalent.

- 1. M covers every K_2^p of H.
- 2. M is not a perfect matching of $G[C_H]$.
- 3. some vertex of C_H is incident to at least two edges of M, or some edge of M is incident to some vertex of $V(H) \setminus C_H$.

Proof. We prove the three following implications.

- $(1) \Rightarrow (2)$. We prove it by contraposition. Assume that M is a perfect matching of $G[C_H]$. Then M forms a partition of some $H' = K_2^p$ of H together with color classes of any K_2^p of H whose are not contained in C_H . Notice that H' is not covered by M.
- $(2) \Rightarrow (3)$. We prove it by contraposition. Assume that every vertex of C_H is incident to at most one edge of M and every edge of M has both endpoints in C_H . From our assumption, every vertex of C_H is incident to exactly one edge of M. Therefore, M is a perfect matching of $G[C_H]$.
- $(3)\Rightarrow (1)$. Let H' be any K_2^p of H. We will show that some edge of M belongs to E(H'). We consider two cases. If some vertex v of C_H is incident to two different edges of M, say (v,u_1) and (v,u_2) , then at least one of these edges belongs to E(H') since u_1 or u_2 belongs to a different color class of H' than v. If some edge e of M is incident to some vertex of $V(H)\setminus C_H$, then e belongs to E(H') since, by Observation 1, it belongs to every K_2^p of H.

Let H be any dense subgraph of G with the core C_H of size 2k. Let $V_1, V_2, \ldots, V_{p-k}$ be the color classes of any K_2^p of H whose are not contained in C_H . We define $G_H - a$ gadget for H. By Lemma 11, w is vertex-induced on H. Let r_H be the potential function of H. We define $c_H - a$ center of H as a vertex of C_H of the minimum potential. If there many such vertices, we choose any of them. If the potential of c_H is negative, then the gadget for H is empty. We show later in the proof of Theorem 6 that every K_2^p of such dense subgraph H is covered by the constructed co-t-matching M.

We assume now that the potential of c_H is nonnegative. In such a case, we introduce p-k+1 subdivision vertices $u_H^c, u_H^1, u_H^2, \ldots, u_H^{p-k}$ and a global vertex z_H . For every i, we connect u_H^i with every vertex of V_i by half-edges and with z_H . We connect u_H^c with c_H by two half-edges and with z_H by two edges (see Fig. 4). A half-edge of H incident to vertex v gets weight $r_H(v)$. Notice that both half-edges incident to c_H get weight $r_H(c_H) \geq 0$. The remaining edges of the gadget get weight $v_H(v)$. We set the capacity interval of vertex v_H^c to v_H^c 0, of vertex v_H^c 1 to v_H^c 2, and of every other subdivision vertex of the gadget for v_H^c 3.

The main idea behind this gadget is the following. In the vein of Lemma 12, we want to ensure that some vertex of C_H is incident to some two edges of \bar{M} or there exists an edge of $\bar{M} \cap E(H)$ incident to some vertex of $V(H) \setminus C_H$. If some vertex of C_H is incident to at least two edges of \bar{M} , then we can assume that it is a center of H since \bar{M} is not minimum otherwise. If some edge of \bar{M} has exactly one endpoint in C_H , then we can assume that it is c_H by replacing some edges of \bar{M} if necessary. Therefore, it is sufficient to guarantee that exactly two half-edges of H are present in M'. If both of these half-edges are adjacent to c_H , it corresponds to including some two edges incident to c_H in \bar{M} by "splitting" some edge of M' whose both endpoints belong to C_H . Otherwise, these half-edges correspond to some edge of H with some endpoint in $V(H) \setminus C_H$.

Theorem 5. Let \bar{M} be a minimum weight K_q^p -covering co-t-matching of G. Then we can find an (l,b)-matching M' of G' such that $w'(M') = w(\bar{M})$.

Proof. Let H be any dense subgraph of G with the core C_H of size 2k. Define $\bar{M}_H = \bar{M} \cap E(H)$. We add every edge of \bar{M}_H to M'. If the gadget for H is empty, we are done. Therefore, we can assume that the potential of the center of H is nonnegative. The main idea is to find an edge $e = (v_1, v_2)$ of \bar{M}_H such that $v_1, v_2 \in V(H) \setminus C_H \cup \{c_H\}$ and to replace it in M' by two half-edges – one incident to v_1 and one incident to v_2 . Then we add p - k edges incident




Figure 4: A gadget for a dense subgraph consisting of eight vertices with the core of size four whose center c_H has nonnegative potential.

to z_H to M' in order to satisfy the degree contraints of the subdivision vertices of the gadget for H. If such edge e exists, we are done. Therefore, we assume that such edge does not exist.

If some vertex $v \in C_H \setminus \{c_H\}$ is matched to some vertex $u \in V(H) \setminus C_H$, we consider two cases. If v is incident to another edge of \bar{M}_H besides (v,u), we change \bar{M} by replacing (v,u) by (c_H,u) . Notice that this replacement does not increase the weight of \bar{M} since c_H has the minimum potential among the vertices of C_H . Moreover, \bar{M} is still a co-t-matching of G after this operation because v is still incident to some edge of \bar{M} . If (v,u) is the only edge of \bar{M}_H incident to v, let v' be any vertex of C_H matched to c_H in \bar{M}_H . Notice that v' exists since \bar{M} is a co-t-matching of G. From our assumption, $v \neq v'$ and $(v,v') \notin \bar{M}$. We replace (v,u) and (c_H,v') in \bar{M} by (c_H,u) and (v,v'). Notice that this replacement does not affect the weight of \bar{M} because, by Lemma 11, w is vertex-induced on H. In both cases, we choose (c_H,u) as the edge e.

If no vertex of $C_H \setminus \{c_H\}$ is matched to a vertex of $V(H) \setminus C_H$, then every edge of \overline{M}_H has both endpoints in C_H . Recall that \bar{M} is K_q^p -covering, so by Lemma 12, some vertex of C_H is incident to two edges of \bar{M}_H . By Lemma 11, w is vertex-induced on H, so $w(\bar{M}_H) = \sum_{v \in C_H} r_H(v) \deg_{\bar{M}_H}(v)$. The main idea is to replace \bar{M}_H in \bar{M} by a subset N of k+1 edges of $G[C_H]$ in which c_H has degree at least two in such a way that only the vertices of smallest potential in C_H have degree higher than one in N. It guarantees that $w(N) \leq w(\bar{M}_H)$ since, from our assumption, the potentials of the vertices of C_H are nonnegative. We consider two cases. If some vertex of C_H is incident to at least three edges of \bar{M}_H , then N consists of three different edges of $G[C_H]$ incident to c_H and a perfect matching of a subgraph induced by the remaining k-4 vertices of C_H . Since the degree of c_H in N is three and the degree of every other vertex of C_H in N is one, we can conclude that $w(N) \leq w(\overline{M}_H)$. Otherwise, some two vertices of C_H have degree two in \bar{M}_H . Let c_H' be the vertex of $C_H \setminus \{c_H\}$ of the minimum potential. Then N consists of an edge (c_H, c'_H) and any perfect matching of $G[C_H]$ which does not contain edge (c_H, c'_H) . Since the degrees of both c_H and c'_H in N are two and the degree of every other vertex of C_H in N is one, we can conclude that $w(N) \leq w(\overline{M}_H)$. In both cases, we remove M_H from M' and we choose any two different edges (c_H, v_1) and (c_H, v_2) of N. We add every other edge of N to M'. Moreover, we add an edge (v_1, v_2) and both half-edges incident to c_H to M'. At the end, we add every edge of the gadget for G_H which connects z_H and any subdivision vertex different than u_L^c to M'. Notice that there are exactly p - k such edges.

Theorem 6. Let M' be a minimum weight (l,b)-matching of G'. Then we can find a K_q^p -covering co-t-matching \bar{M} of G such that $w(\bar{M}) \leq w'(M')$.

Proof. We assume that M' has a minimum number of edges among all minimum weight (l,b)-matchings of G'. We initialize \bar{M} as the empty set. We add every edge of M' which belongs to G to \bar{M} . We consider problematic K_2^p 's of G analogously to K_a^p 's in the proof of Theorem 4.

At the beginning we prove that \bar{M} covers every K_2^p contained in any dense subgraph of G whose center has negative potential. Suppose that \bar{M} does not cover some K_2^p of a dense subgraph H with the core C_H such that $r_H(c_H) < 0$. Notice that every vertex of C_H is incident to some edge of $\bar{M}_H = \bar{M} \cap E(H)$ because it has degree t+1 in G and in G'. Therefore, we can use Lemma 12 and conclude that \bar{M}_H is a perfect matching of $G[C_H]$. From the construction of \bar{M} , $M'_H = M' \cap E(H)$ is a perfect matching of $G[C_H]$ as well. Define $M'' = M' \setminus M'_H \cup \{(c_H, v) : d_H \in M' \cap E(H) : d_H \in M'' \cap E(H) : d_H \in M''$

 $v \in C_H \setminus \{c_H\}\}$. Notice that M'' is an (l,b)-matching of G' such that w'(M'') < w'(M') since $r_H(c_H) < 0$. A contradiction with the minimality of M'.

Consider any dense subgraph H of G with the core C_H whose center has nonnegative potential. From the construction of the gadget for H, exactly two its half-edges belong to M'. If these half-edges are incident to two different vertices u_1 and u_2 of G, then we add an edge (u_1, u_2) to \bar{M} unless it belongs already. Notice that u_1 or u_2 belongs to $V(H) \setminus C_H$, so by Lemma 12, \bar{M} covers all K_2^p 's of H. Otherwise, both half-edges of the gadget for H which belong to M' are incident to c_H .

If there exists an edge (v_1, v_2) of M' such that both v_1 and v_2 belong to $C_H \setminus \{c_H\}$, we remove (v_1, v_2) from \bar{M} . Then we add edges (c_H, v_1) and (c_H, v_2) to \bar{M} unless they belong to \bar{M} already. From now we assume that there is no such edge.

We claim that at most one vertex of $C_H \setminus \{c_H\}$ is matched to c_H in M'. Suppose that some two different edges (c_H, v_1) and (c_H, v_2) belong to M' where v_1 and v_2 belong to C_H . From our assumption, (v_1, v_2) does not belong to M', so we can construct an (l, b)-matching of G' of weight smaller than w'(M') by replacing (c_H, v_1) and (c_H, v_2) in M' by (v_1, v_2) . A contradiction with the minimality of M'. Notice that we use the assumption that $r_H(c_H) \geq 0$ here.

Since every vertex of $C_H \setminus \{c_H\}$ is incident to at least one edge of M' and the core of H consists of at least four vertices, there exist two edges (v_1,u_1) and (v_2,u_2) which belong to M' such that v_1 and v_2 belong to C_H whereas u_1 and u_2 belong to $V(H) \setminus C_H$. We replace edges (v_1,u_1) and (v_2,u_2) in \overline{M} by edges (v_1,v_2) , (c_H,u_1) and (c_H,u_2) unless they belong to \overline{M} already.

6. Running time analysis

In this section we consider the running time of Algorithm 1 for both the bounded restricted t-matching problem and the bounded K_q^p -free t-matching problem. Since all problematic and dense subgraphs of G are pairwise vertex-disjoint, graph G' constructed in Step 1 of Algorithm 1 has O(n) vertices and O(m) edges indeed. We prove that Step 3 of Algorithm 1 can be implemented to run in time $\mathcal{O}(\min\{nm\log n, n^3\})$ in the weighted variant (see Subsection 6.1) and $\mathcal{O}(\sqrt{n}m)$ in the unweighted (see Subsection 6.2).

Observe that we have to find all forbidden subgraphs of G in order to build G' and remove unproblematic subgraphs from the acquired co-t-matching \bar{M} of G in Step 5 of Algorithm 1. Moreover, we need to know which forbidden subgraphs of G have common vertices in order to determine which of them are problematic. We prove that this can be done in $\mathcal{O}(m)$ time in Subsection 6.3.

In the weighted setting, we also have to calculate and compare the weights of forbidden subgraphs of G which are not part of any dense subgraph in order to determine which of them are problematic. Notice that, for any such two different forbidden subgraphs H and H' of G which have a common vertex, we can obtain the weight of H' from the weight of H in $\mathcal{O}(t)$ time because the vertex sets of H and H' differ by at most two vertices. We initially calculate the weights of some vertex-disjoint forbidden subgraphs of G naively and we use these results to obtain the weights of the remaining forbidden subgraphs.

On the other hand, we have to know the potential function of every problematic and dense subgraph of G in order to define the weights of half-edges in the gadgets in Step 2 of Algorithm 1. Since we do not assume that the forbidden subgraphs are given as a part of the input, we cannot assume that the potential functions of forbidden subgraphs are given. However, we show that the potential function of any forbidden subgraph of G can be extracted from the weight function w in $\mathcal{O}(t^2)$ time in Subsection 6.4. By Lemma 11, the potential function of any dense subgraph H of G can be found by extracting the potential function of any forbidden subgraph contained in H. Therefore, the potential functions of all problematic and dense subgraphs of G can be found in $\mathcal{O}(m)$ time because they are vertex-disjoint.

We claim that Step 5 of Algorithm 1 can be implemented to run in $\mathcal{O}(m)$ time since all forbidden subgraphs of G which are not covered by co-t-matching \bar{M} are vertex-disjoint. It follows from the simple observation that any two different forbidden subgraph H_1 and H_2 with a common vertex have some common vertex v of degree exactly t+1 whose every incident edge belongs to H_1 or H_2 . Since \bar{M} is a co-t-matching of G, such vertex v has to be incident to some edge of \bar{M} , so \bar{M} covers H_1 or H_2 .

We discuss briefly how G is stored in the memory by Algorithm 1. For every vertex of G, it stores a list of its neighbours, in no particular order. In the weighted variant, every entry in these lists also contains the weight of the corresponding edge.

6.1. Computing a minimum weight (l, b)-matching of G' in the weighted case

In this subsection, we present how to find a minimum weight (l,b)-matching of G' in $\mathcal{O}(\min\{nm\log n, n^3\})$ time. We use the algorithm from Theorem 1 for G', l and b with the weight function -w, i.e. we set the weight of every edge e of G' to -w'(e). However, this approach results in $\mathcal{O}(\min\{m^2\log n, n^2m\})$ running time since sum of the upper bounds of the capacity intervals of the vertices of G' is $\mathcal{O}(m)$. We show how to modify the algorithm from Theorem 1 to work in $\mathcal{O}(\min\{nm\log n, n^3\})$ time in our case. To explain this, we have to briefly present how this algorithm works.

At the beginning, the algorithm constructs an auxiliary graph G^* by creating two copies of graph G' and connecting the corresponding copies v_1 and v_2 of every vertex v of G' by b(v) - l(v) paths of length three. We call these paths *added paths* (of vertex v). Both edges incident to endpoints of an added path P are said to be *outermost*. The remaining edge of P is said to be *middle*. Both endpoints of the middle edge of P are said to be *central*. The algorithm calculates a maximum weight (b^*, b^*) -matching of G^* where $b^*(v_i) = b(v)$ for every copy v_i of vertex v of G' and $b^*(u) = 1$ for every other vertex u of G^* . The weight function w^* of G^* is defined as follows. The weight of any copy of an edge e of G' is equal to -w'(e). Let $W = \max_{e \in E(G')}\{|w'(e)|\}$. For every added path, its middle edge gets weight 2W and both outermost edges get weight W. It is easy to check that a maximum weight (l,b)-matching of G' with the weight function -w' corresponds to a maximum weight (b^*,b^*) -matching of G^* with the weight function w^* .

The algorithm finds a maximum weight (b^*,b^*) -matching of G^* as follows. It simulates the classical algorithm for finding a maximum weight matching, given by Edmonds [6] and improved by Gabow [8] and Galil, Micali and Gabow [10], in an auxiliary graph \hat{G} . Graph \hat{G} is constructed from G^* by splitting every its vertex v into $\deg_{G^*}(v)$ vertices called *external vertices* (of v). The external vertices of v are incident to $\deg_{G^*}(v)$ edges – each vertex to one edge – corresponding to all the edges incident to v in G^* . With a slight abuse of notation, we identify every edge of G^* with its corresponding edge in \hat{G} . Every external vertex of v which is incident to some outermost edge of some added path is said to be *improper*. All other external vertices of v are said to be *proper*. Futhermore, $\deg_{G^*}(v) - b^*(v)$ additional vertices, called *internal vertices* (of v) are added to \hat{G} . Every internal vertex of v is connected to every external vertex of v by an edge. We refer to a graph induced by all external and internal vertices of v as v0 as v1 as v2 as v3 as v4 as v5. The weight function v6 of v6 is defined as follows. Every edge of v6 which corresponds to some edge v6 as v7 gets weight v8. Every edge incident to some internal vertex gets weight v8.

It is well known (see [3] for example) that every b^* -matching of G^* corresponds to some matching of G in which every internal vertex is matched. Moreover, every (b^*, b^*) -matching of G^* corresponds to some perfect matching of G. Hence, the algorithm computes a maximum weight perfect matching of G. A matching G of G is said to be **extreme** if it has the maximum weight among all matchings of G of size $|\hat{M}|$. The algorithm computes extreme matchings of G consisting of exactly G exactly G edges, for G exactly G exactly G edges, for G exactly G exac

$$\frac{1}{2} \sum_{v \in V(G^*)} b^*(v) = \sum_{v \in V(G')} b(v) + \frac{1}{2} \sum_{v \in V(G')} (b(v) - l(v)) = O(m).$$

The main idea behind the speed-up of the presented algorithm is to initialize it with extreme matching M_1 of \hat{G} which has only $\mathcal{O}(n)$ unmatched vertices instead of M_0 . Such an improvement results in $\mathcal{O}(n)$ phases instead of

 $\mathcal{O}(m)$, which gives the desired running time. In fact, we choose M_1 which is a maximum weight (not necessarily perfect) matching of \hat{G} . It guarantees that M_1 is an extreme matching. M_1 roughly correponds to a matching N of \hat{G} which contains both outermost edges of every added path. Of course, for every internal vertex v of \hat{G} , N contains exactly one edge incident to v which is not incident to any other edge of N. Notice that the number of vertices of \hat{G} unmatched in N is equal to $2\sum_{v\in V(G')}l(v)=\mathcal{O}(n)$. However, N does not have to be an extreme matching of \hat{G} . Observe that the weight of the middle edge of an added path is equal to the total weight of its outermost edges. Therefore, it may happen that replacing outermost edges of N of some added path P by the middle edge of P and some copy of some edge of P which has the positive weight results in a matching of P of size P and weight greater than P the weight with regard to P if and only if P with the weight function P. Hence, any edge P of P has positive weight with regard to P if and only if P is negative. Notice that only half-edges of the gadgets in P may have negative weights since we assume that the weight P of any edge P of P is nonnegative. A half-edge P has negative weight P if its endpoint which belongs to P has negative potential. Therefore, we include some half-edges in P in such a way that the number of vertices of P unmatched in P is still P to P in such a way that the number of vertices of P unmatched in P is still P in such a way that the number of vertices of P unmatched in P is still P in such a way that the number of vertices of P unmatched in P is still P in such a way that the number of vertices of P unmatched in P is still P in such a way that the number of vertices of P unmatched in P is still P the number of vertices of P unmatched in P is still P the number of vertices of P unmatched in P is stil

Now we give the exact construction of M_1 . We initialize M_1 as the empty set. We partition the edge set of \hat{G} into disjoint subsets, called **parts** of \hat{G} . For every part E_0 of \hat{G} , we choose a subset $M \subseteq E_0$, called a **representation** of E_0 , such that, for every matching N of \hat{G} , $\hat{w}(M) \ge \hat{w}(N \cap E_0)$. We add the representation of every part of \hat{G} to M_1 . We choose the representations of parts of \hat{G} in such a way that M_1 is a matching of \hat{G} . It is easy to check that M_1 is indeed a maximum weight matching of \hat{G} .

Now we present a partition of the edge set of \hat{G} into parts of \hat{G} and its representations.

- 1. Consider any copy u_i in G^* of a subdivision vertex u of G'. Observe that no added path of u was added to G^* , so all external vertices of u_i are proper. Notice that the external vertices of u_i are incident to some half-edges and possibly to some edges of weight zero. Furthermore, the substitute of u_i consists of $\deg_{G'}(u)$ external and $\deg_{G'}(u) b(u)$ internal vertices. We construct part E_{u_i} of \hat{G} from the edge set of the substitute of u_i and all edges of positive weight (with regard to \hat{w}) which are incident to some external vertex of u_i . The representation of E_{u_i} consists of up to b(u) copies of half-edges incident to u of the greatest positive weights and one incident edge of the substitute of u_i per every internal vertex of u_i . If there are less than b(u) such half-edges, the copies of all of them are added to the representation of E_{u_i} . Of course, we choose this representation in such a way that its two different edges are not incident.
- 2. Consider any copy z_i in G^* of a global vertex z of G'. Recall that all edges incident to z in G' have weight zero. We construct part E_{z_i} from the edge set of the substitute of z_i . Its representation consists of one incident egde per every internal vertex of z_i .
- 3. Consider any vertex v of G. For every copy v_i of v in G^* , we construct part E_{v_i} from the edge set of the substitute of v_i . Moreover, for every added path P of v, we construct part E_P from all edges of P and all edges of the substitutes of both central vertices of P. Recall that v is incident to at most one half-edge in G'. Hence, at most one proper external vertex u_0 of v_i is incident to some edge which belongs to the representation of some part of \hat{G} . Observe that, if u_0 exists, we cannot include any edge incident to u_0 in the representation of E_{v_i} . The construction of the representations of these parts depends on the degree of v in G.
 - Consider a case where $\deg_G(v) = t + 1$. Notice that the substitute of any copy v_i of v consists of t internal vertices, t + 1 proper external vertices and t improper external vertices. The representation of E_{v_i} consists of exactly one edge incident to u and some proper external vertex of u_i different than u_0 , per every internal vertex u of v_i .
 - Consider a case where $d=\deg_G(v)\leq t$. Notice that the substitute of any copy v_i of v consists of d internal vertices, d proper external vertices and d improper external vertices. If u_0 does not exist, the representation of E_{v_i} consists of one edge incident to u and some proper external vertex of v_i , per every internal vertex u of v_i . If u_0 exists, the representation of E_{v_i} consists of d edges. Every edge of this representation is incident to some internal vertex of v_i . On the other hand, d-1 of these edges are incident to d-1 proper external vertices of v_i different than u_0 . One remaining edge is incident to an improper external vertex v_i' of v_i . We choose v_1' and v_2' in such a way that both of them are incident to the outermost edges of the same added path of v.

For every added path P of v, we choose the representation of E_P in the following way. If both outermost edges of P are incident to v_1' or v_2' , the representation of E_P consists of the middle edge of P and one edge from the edge set of every substitute of a center of P. Otherwise, the representation of E_P consists of the outermost edges of P and, again, one edge from the edge set of every substitute of a center of P. Notice that the total weight of the representation of E_P is equal to 6W.

4. Consider any edge e of \hat{G} which does not belong to any part of \hat{G} yet. Observe that $\hat{w}(e) \leq 0$. We construct part E_e which consists of single edge e. Its representation is empty.

It remains to show that the number of vertices of \hat{G} unmatched in M_1 is $\mathcal{O}(n)$. Notice that all internal and improper external vertices in \hat{G} are matched in M_1 . Hence, it is sufficient to consider only proper external vertices. Observe that, for every copy u_i in G^* of any subdivision vertex u of G', at most b(u) proper external vertices of u_i are unmatched in M_1 . Moreover, for every copy z_i in G^* of any global vertex z of G', there are exactly b(z) proper external vertices of z_i unmatched in M_1 . Therefore, there are $\mathcal{O}(t)$ proper external vertices unmatched in M_1 per every gadget in G'. For any copy v_i in G^* of any vertex v of G, at most one proper external vertex of v_i is unmatched in M_1 . It exists only if $\deg_G(v) = t+1$ and no proper external vertex of v_i is incident to a half-edge which belongs to the representation of some part of \hat{G} . Therefore, there are $\mathcal{O}(n)$ proper external vertices in \hat{G} which are unmatched in M_1 .

6.2. Computing a minimum weight (l, b)-matching of G' in the unweighted case

In this subsection, we present how to find a minimum weight (l,b)-matching of G' in $\mathcal{O}(\sqrt{n}m)$ time in the unweighted case, i.e. when w(e)=1 for every edge e of G. Notice that w is vertex-induced in this case since we can set the potential of every vertex of a forbidden subgraph to $\frac{1}{2}$. Hence, the weight w'(e) of every half-edge e of G' is defined as $\frac{1}{2}$. We claim that the minimum weight (l,b)-matching of G' is a **minimum cardinality** (l,b)-matching of G', i.e. an (l,b)-matching of G' with the minimum number of edges among all (l,b)-matchings of G'. If follows from the following lemma.

Lemma 13. Let M' be any (l, b)-matching of G'. Then

$$|M'| - w'(M') = k_{t+1} + k_{t,t} + (p-1)k_q^p + \sum_{H-a \text{ dense subgraph of } G} (p - \frac{|C_H|}{2} + 1),$$

where k_{t+1} , $k_{t,t}$ and k_q^p denote the number of the gadgets for, correspondingly, K_{t+1} 's, $K_{t,t}$'s and K_q^p 's in G'.

Proof. Notice that every edge of M' which does not belong to any gadget contributes one to both |M'| and w'(M'). Hence, |M'| - w'(M') is equal to the sum of values $|M'_H| - w'(M'_H)$ for all forbidden subgraphs H of G for which the gadget was added in G', where M'_H denotes a set of all edges of the gadget for H which belong to M'. Notice that for every problematic K_{t+1} and $K_{t,t}$ H of G, M'_H consists of exactly two half-edges, hence $|M'_H| - w'(M'_H) = 1$ then. Similarly, for every $H = K_q^p$ of G, $|M'_H| = p$ and $w'(M'_H) = 1$. In the end, for any dense subgraph H of G with the core of size 2k, we have $|M'_H| = p - k + 2$ and $w'(M'_H) = 1$.

We define a *maximum cardinality* (l,b)-matching of G' analogously. To calculate a minimum cardinality (l,b)-matching of G' we use the following algorithm given by Gabow.

Theorem 7 ([9]). There is an algorithm that, given a multigraph G = (V, E) (i.e. G may contain parallel edges) and vectors $l, b \in \mathbb{N}^V$, finds a maximum cardinality (l, b)-matching of G in time $O\left(\sqrt{\sum_{v \in V} b(v)}|E|\right)$, assuming that G admits any (l, b)-matching.

It turns out that calculating a minimum cardinality (l, b)-matching can easily be reduced to calculating a maximum cardinality (l, b)-matching.

Lemma 14. There is an algorithm that, given a multigraph G = (V, E) (i.e. G may contain parallel edges) and vectors $l, b \in \mathbb{N}^V$, finds a **minimum** cardinality (l, b)-matching of G in time $O\left(\sqrt{\sum_{v \in V} b(v)}|E|\right)$, assuming that G admits any (l, b)-matching.

Proof. We construct an auxiliary graph G^* together with vectors $l^*, b^* \in \mathbb{N}^{V(G^*)}$. We start off with $G^* = G$. For every vertex v of G, we add an auxiliary vertex v' to G^* . We connect v with v' by b(v) - l(v) parallel edges in G^* . We set $l^*(v) = b^*(v) = b(v)$, $l^*(v') = 0$ and $b^*(v') = b(v) - l(v)$. We claim that every (l, b)-matching of G corresponds to an (l^*, b^*) -matching of G^* . Indeed, for any (l, b)-matching M of G, we can construct an (l^*, b^*) -matching M^* of G^* by adding, for every vertex v of G, $b(v) - \deg_M(v)$ copies of an edge (v, v') to M. Notice that

$$|M^*| = |M| + \sum_{v \in V} (b(v) - \deg_M(v)) = \left(\sum_{v \in V} b(v)\right) - |M|,$$

so a minimum cardinality (l,b)-matching of G corresponds to a maximum cardinality (l^*,b^*) -matching of G^* . Hence, we can calculate the solution using the algorithm from Theorem 7 on G^* , l^* and b^* . Notice that $\sum_{v \in V(G^*)} b^*(v) \leq 2\sum_{v \in V(G)} b(v)$. Moreover, we can assume that $b(v) \leq \deg_G(v)$ for every vertex v of G. Both of these facts guarantee the desired running time. \square

We can calculate a minimum cardinality (l,b)-matching of G' by applying Lemma 14 directly. However, it works in $\mathcal{O}(m^{3/2})$ time since the sum of the upper bounds of the capacity intervals in G' is O(m). We can calculate it faster using the following lemma.

Lemma 15. Let M' be any (l,b)-matching of G'. Then there exists a minimum cardinality (l,b)-matching N' of G' such that $\deg_{N'}(v) \leq \deg_{M'}(v)$ for every vertex v of G'.

We postpone the proof of Lemma 15 since it is quite technical. The main idea behind calculating a minimum cardinality (l,b)-matching of G' is to find some (l,b)-matching M' of G' which consists of $\mathcal{O}(n)$ edges. Such M' can be constructed in the following way. We initialize M' as the empty set. For every vertex v of degree t+1 in G, we add any edge of G incident to v to M'. Additionally, we have to add some edges of the gadgets in order to satisfy the degree constrains of the global and subdivision vertices. It is easy to check that O(t) edges of every gadget are sufficient, hence M' has $\mathcal{O}(n)$ edges in total. Using Lemma 15, we can find a minimum cardinality (l,b)-matching of G' by running the algorithm from Lemma 14 for G and vectors $l,b'\in\mathbb{N}^{V(G)}$ such that $b'(v)=\deg_M(v)$ for every vertex v of G'. Since $\sum_{v\in V}b'(v)=2|M'|=\mathcal{O}(n)$, the total runtime of this procedure is $\mathcal{O}(\sqrt{nm})$.

To prove Lemma 15, we use the classical notion of alternating paths and cycles. Let M be an (l,b)-matching of a graph G. An edge belonging to M will be referred to as an M-edge and an edge not belonging to M as a non-M-edge. An M-alternating path P is any sequence of vertices (v_1, v_2, \ldots, v_k) such that edges on P are alternately M-edges and non-M-edges and no edge occurs on P more than once. An M-alternating cycle C has the same definition as an M-alternating path except that $v_1 = v_k$ and additionally $(v_{k-1}, v_k) \in M$ if and only if $(v_1, v_2) \notin M$. Note that an M-alternating path or cycle may go through some vertices more than once but via different edges. An M-alternating path is called M-augmenting if it begins and ends with a non-M-edge. With a slight abuse of notation, we identify an M-alternating path or cycle with its edge set. For subsets $A, B \subseteq E$, we define a symmetric difference $A \oplus B = (A \setminus B) \cup (B \setminus A)$.

Proof of Lemma 15. Let N be any minimum cardinality (l,b)-matching of G'. It is well-known that $M' \oplus N$ can be partitioned into a collection $\mathcal P$ of pairwise edge-disjoint paths and cycles which are both M'-alternating and N-alternating. We assume that partition $\mathcal P$ is minimal, i.e. no two paths or cycles of $\mathcal P$ can be merged into one M'-alternating and N-alternating path or cycle. From the definition of N, $|N| \leq |M'|$, so $\mathcal P$ contains k = |M'| - |N| different paths P_1, P_2, \ldots, P_k which are both M'-alternating and N-augmenting. We say that these paths are special. Define $N' = M' \oplus P_1 \oplus P_2 \oplus \ldots \oplus P_k$, i.e. N' consists of all M'-edges which do not belong to any special path and all N-edges which belong to some special path. We claim that N' is the sought-after minimum cardinality (l,b)-matching of G'. At first, notice that |N'| = |M'| - k = |N|. Moreover, observe that adding P_i to the symmetric difference with M' decrease only the degree of both endpoints of P_i while leaving the degrees of other vertices of G' unchanged. Therefore, $\deg_{N'}(v) \leq \deg_{M'}(v) \leq b(v)$ for every vertex v of G'.

It remains to show that $\deg_{N'}(v) \geq l(v)$ for every vertex v of G'. From our previous observation, it is sufficient to prove this only for endpoints of special paths. Consider any vertex v of G' which is an endpoint of $p \geq 1$ special paths. If some special path has both endpoints in v, we calculate it twice. Notice that v is incident to exactly p different M'-edges in G' which belong to some special paths. We claim that the number of remaining N-edges incident to v

in G' is not greater than the number of remaining M'-edges incident to v in G'. Indeed, if there are more N-edges than M'-edges among the remaining edges incident to v in G', then v is an endpoint of some M'-alternating and N-alternating path P' of $\mathcal P$ such that an edge of P' which is incident to v in G' belongs to N. Notice that we can merge P' and any special path which has an endpoint in v into one path - a contradition with the minimality of $\mathcal P$. From our claim, $\deg_N(v) \leq \deg_{M'}(v) - p$. Hence, $\deg_{M'}(v) = \deg_{M'}(v) - p \geq \deg_N(v) \geq l(v)$ because N is an (l,b)-matching of G'.

6.3. Finding all forbidden subgraphs of G

In this subsection, we present Algorithm 2 which finds all K_q^p 's of G in $\mathcal{O}(m)$ time, given some natural numbers $p \geq 2$ and $q \geq 1$ such that t = (p-1)q. It also returns which pairs of K_q^p 's of G have some common vertices. Applying this algorithm to Algorithm 1 for the bounded K_q^p -free t-matching problem is fairly straightforward. For the bounded restricted t-matching problem, we run Algorithm 2 twice – once to find all K_{t+1} 's of G, and again to find all $K_{t,t}$'s of G. However, we have to find also all pairs consisting of K_{t+1} and $K_{t,t}$ of G which have a common vertex. This can be done in $\mathcal{O}(m)$ time since Lemma 3 indicates that for every such pair, t = 3 and the vertex set of K_{t+1} is a subset of the vertex set of $K_{t,t}$.

We say that a graph is **bounded** if its every vertex has degree at most t+1. The following lemma gives an important subprocedure of Algorithm 2. We postpone its proof since it is quite technical.

Lemma 16. Let v be any vertex of a bounded graph \tilde{G} . We can find in $\mathcal{O}(t^2)$ time all K_q^p 's of \tilde{G} which contain v.

The main idea behind Algorithm 2 is to find two different vertices v_1 and v_2 of G which have $\Theta(t)$ common neighbours using Lemma 17 given below. We check if v_1 or v_2 belongs to some K_q^p of G using Lemma 16. Recall that if some two K_q^p 's of G have a common vertex, then they have at least pq-2 common vertices (see Lemmas 1, 2, 6 and 8). Therefore, if we find some $H=K_q^p$ of G, we can find all K_q^p 's of G which have some common vertex with H in $\mathcal{O}(t^2)$ time. In such a case, we return H and all K_q^p 's of G which have a common vertex with H. Then we remove the vertex set of H from G and we continue to search for K_q^p 's in the rest of the graph. Observe that if v_1 and v_2 do not belong to any K_q^p of G, then none of the common neighbours of v_1 and v_2 belongs to any K_q^p of G as well. In this case, we remove v_1 , v_2 , and all their common neighbours from G, and we search for K_q^p 's in the rest of the graph. In both cases, we remove $\Theta(t)$ vertices in $\mathcal{O}(t^2)$ time, which means that we spend $\mathcal{O}(t)$ time per vertex on average. Since no vertex of degree less than t belongs to some K_q^p of G, we can remove such vertices from G at the beginning. Therefore, we can conclude that the running time of Algorithm 2 is $\mathcal{O}(m)$.

Lemma 17. Let v_1 be any vertex of a bounded graph \tilde{G} . There exists a procedure which works in $\mathcal{O}(t)$ time and it either concludes that v_1 does not belong to any K_q^p of \tilde{G} , or returns a vertex $v_2 \neq v_1$ of \tilde{G} which has at least $\max\{p-2,1\}q$ common neighbours with v_1 .

Proof. The procedure works in the following way. It processes any two different neighbours of v_1 , say u_1 and u_2 (or all neighbours of v_1 , if there are less than two of them). The procedure processes every vertex $u \in \{u_1, u_2\}$ as follows. It checks any two different neighbours of u different than v_1 , say z_1 and z_2 , as candidates for v_2 (or all of them, if there are less than two such vertices). For every $z \in \{z_1, z_2\}$, the procedure checks if v_1 and v_2 have at least $\max\{p-2,1\}q$ common neighbours. If so, the procedure returns $v_2=z$. If none of the vertices $v_2=z$ satisfies this condition, the procedure returns that $v_1=z$ does not belong to any $v_2=z$. It is easy to notice that this procedure can be implemented to run in $v_2=z$ time. Its correctness follows from the following claims.

Claim 3. Assume that v_1 belongs to some $H = K_q^p$ of G. Then the procedure given above considers some u and z such that both edges (v_1, u) and (u, z) belong to H.

Proof. We consider the execution of the presented procedure. Since v_1 belongs to t-regular H, v_1 has at least $t \geq 3$ neighbours in \tilde{G} , so u_1 and u_2 exist. Recall that \tilde{G} is bounded, so $\deg_{\tilde{G}}(v_1) \leq t+1$. Therefore, at most one edge of \tilde{G} incident to v_1 does not belong to H. It implies that at least one of edges (v_1, u_1) and (v_1, u_2) belongs to H. Consider $u \in \{u_1, u_2\}$ such that (v_1, u) belongs to H. We can use similar arguments for u as for v_1 to prove that both v_1, v_2 exist and that at least one of edges v_1, v_2 belongs to v_1, v_2 belongs to v_2, v_3 such that v_1, v_2 belongs to v_3 .

Claim 4. Assume that some two different edges (w_1, t) and (w_2, t) belong to the same K_q^p of \tilde{G} . Then w_1 and w_2 have at least $\max\{p-2, 1\}q$ common neighbours.

Proof. Let H be any K_q^p of G which contains both (w_1,t) and (w_2,t) . Assume that $t \in V_1(H)$. Then both w_1 and w_2 belong to $V(H) \setminus V_1(H)$. We consider the following two cases.

- 1. If w_1 and w_2 belong to the same color class of H, say $V_2(H)$, then they have at least (p-1)q common neighbours all vertices of $V(H) \setminus V_2(H)$.
- 2. If w_1 and w_2 belong to different color classes of H, say $V_2(H)$ and $V_3(H)$ respectively, then they have at least (p-2)q common neighbours all vertices of $V(H) \setminus (V_2(H) \cup V_3(H))$.

In both cases, w_1 and w_2 have at least (p-2)q common neighbours. However, case 2. cannot happen if p=2, so we can conclude that w_1 and w_2 have at least $\max\{p-2,1\}q$ common neighbours.

Now we present a pseudocode of Algorithm 2 and a proof of its correctness.

Algorithm 2 Finding all K_a^p 's of G

- 1: Initialize \tilde{G} as G. Remove all vertices of degree less than t in G from \tilde{G} .
- 2: If \tilde{G} is empty, stop the algorithm.
- 3: Run the procedure from Lemma 17 on \tilde{G} and any its vertex v_1 .
- 4: If v_1 does not belong to any K_q^p of G, remove v_1 from G and go to Step 2.
- 5: Otherwise, let v_2 be a vertex found in Step 3. Run the procedure from Lemma 16 on \tilde{G} and v_i , for every $i \in \{1,2\}$.
- 6: If none of the vertices v_1 , v_2 belong to any K_q^p of \hat{G} , remove v_1 , v_2 and all its common neighbours from G and go to Step 2.
- 7: Otherwise, let H be any K_q^p found in Step 5. Return H and all K_q^p 's of \tilde{G} which have some common vertex with H. Return that every two different returned K_q^p 's have a common vertex. Remove H from \tilde{G} and go to Step 2.

Claim 5. Algorithm 2 finds all K_q^p 's of G in $\mathcal{O}(m)$ time. Moreover, it finds all pairs of K_q^p 's of G which have a common vertex.

Proof. At first, we prove that Algorithm 2 finds all K_q^p 's of G. We claim that during its execution an invariant is maintained that all K_q^p 's of G which were not returned yet are present in \tilde{G} . To show this, it is sufficient to prove that every time some vertex v is removed from \tilde{G} , all K_q^p 's of \tilde{G} which contain v were returned by Algorithm 2 already. We consider all steps of Algorithm 2 where some vertices are removed from \tilde{G} :

- Step 1: notice that no vertex removed from \tilde{G} belongs to any K_q^p of G since K_q^p is t-regular,
- Step 4: we remove v_1 which does not belong to any K_a^p of \tilde{G} ,
- Step 6: if some common neighbour of v_1 and v_2 belongs to some K_q^p of \tilde{G} , then also v_1 or v_2 belongs to some K_q^p of \tilde{G} , but we assume that both v_1 and v_2 do not belong to any K_q^p of \tilde{G} ,
- Step 7: we remove V(H) from \hat{G} , but we return all K_q^p 's of \hat{G} which contain some vertex of V(H).

Now we prove that Algorithm 2 finds all pairs of K_q^p 's of G which have a common vertex. We return some K_q^p 's only in Step 7. Recall that all these K_q^p 's do not belong to \tilde{G} after Step 7 anymore. Notice that it is sufficient to show that all these K_q^p 's do not have any common vertices with all other K_q^p 's of \tilde{G} . It follows from the following claim.

Claim 6. Let H_1 , H_2 and H_3 be any K_q^p 's of a bounded graph \tilde{G} . Assume that H_1 has a common vertex with H_2 and a common vertex with H_3 . Then H_2 and H_3 have a common vertex.

Proof. Recall that, by Lemmas 1, 2, 6 and 8, any two different K_q^p 's of \tilde{G} which have a common vertex have at least pq-2 common vertices. Moreover, if p>2, then they have at least pq-1 common vertices.

Consider a case where p > 2. Then at most one vertex of H_1 does not belong to H_2 and at most one vertex of H_1 does not belong to H_3 . Hence, there are at least $pq - 2 = t + q - 2 \ge 2$ vertices of H_1 which belong to both H_2 and H_3 . Any of these vertices is a common vertex of H_2 and H_3 .

Consider a case where p=2. Then at most two vertices of H_1 do not belong to H_2 and at most two do not belong to H_3 . Hence, there are at least $2q-4=2t-4\geq 2$ vertices of H_1 which belong to both H_2 and H_3 .

It remains to show that Algorithm 2 works in $\mathcal{O}(m)$ time. It is easy to implement Step 1 in $\mathcal{O}(m)$ time. We divide the rest of the execution of Algorithm 2 into phases. Every phase works in $\mathcal{O}(tk)$ time and ends with the removal of k vertices from \tilde{G} , for some natural positive number k. Notice that it implies that Algorithm 2 works in $\mathcal{O}(tn_t)$ time where n_t denotes the number of the vertices of degree at least t in G, but from the handshaking lemma, $tn_t \leq 2m$.

The partition of the rest of the execution of Algorithm 2 into phases is simple. For every removal of some vertices from \tilde{G} , its phase consists of this removal and all steps preceding it until the previous removal. We prove that every such phase works in $\mathcal{O}(tk)$ time indeed, where k denotes the number of removed vertices. For the removal in Step 4, by Lemma 17, its phase works in $\mathcal{O}(t)$ time and we remove one vertex. For the removal in Steps 6 and 7, by Lemmas 16, 17 and our previous observations, its phase works in $\mathcal{O}(t^2)$ time and we remove $\Theta(t)$ vertices.

Now we present the proof of Lemma 16. We prove it for cases where $K_q^p = K_{t+1}$, $K_q^p = K_2^p$ and for the remaining cases separately. At first, we give some definitions. Let G = (V, E) be any graph. A **complement** of G is a graph H = (V, E'), where E' consists of all pairs of different vertices of V which are not connected by any edge from E. Graph G is said to a **star** if $E = \{(v_0, v) : v \in V \setminus \{v_0\}\}$ for some vertex v_0 of V. In such a case, vertex v_0 is called a **center** of G. Graph G is said to be **empty** if its edge set is empty. For vertex-disjoint sets $A, B \subseteq V$, G[A, B] denotes a bipartite graph $(A \cup B, E')$ where E' consists of all edges of G whose one endpoint belongs to G and another one to G. A **bipartite complement** of G[A, B] is a bipartite graph $(A \cup B, E'')$ where $E'' = \{(a, b) : a \in A \land b \in B \land (a, b) \notin E\}$. Any vertex $v \in V$ is said to be a **cut vertex** of G if a removal of G increases the number of the connected components of G. A cut vertex G is said to be **nontrivial** if every connected component which appears after the removal of G from G consists of at least two vertices. We extend the definition of G to cases where G is a graph whose vertex set is empty whereas G is an empty graph which consists of G vertices.

Proof of Lemma 16 for K_{t+1} 's. Let e_1 and e_2 be any two different edges incident to v in \tilde{G} . (If v has the degree less than two in \tilde{G} , then v clearly cannot belong to any t-regular K_{t+1} of \tilde{G} since $t \geq 3$.) Because the degree of v in \tilde{G} is not greater than t+1, any K_{t+1} of \tilde{G} which contains v has to contain e_1 or e_2 . For each edge $e \in \{e_1, e_2\}$, we find all K_{t+1} 's of \tilde{G} which contain e. Observe that if e belongs to some K_{t+1} of \tilde{G} , then both its endpoints must have at least t-1 common neighbours.

If both endpoints of e have exactly t-1 common neighbours, we check if these common neighbours and both endpoints of e form some K_{t+1} of \tilde{G} . Observe that checking if a given subset A of t+1 vertices form a K_{t+1} in a bounded graph \tilde{G} can be implemented to run in $\mathcal{O}(t^2)$ time – we can construct the complement of $\tilde{G}[A]$ and check if it contains no edges. Indeed, for every vertex u of A, we can mark all its neighbours in \tilde{G} in $\mathcal{O}(t)$ time. All unmarked vertices of $A \setminus \{u\}$ are all the neighbours of u in the complement of $\tilde{G}[A]$.

If both endpoints of e have exactly t common vertices, we check if they form a K_{t+2} of \tilde{G} together with both endpoints of e. If not, we make use of the following observation to find all K_{t+1} 's in a subgraph of \tilde{G} induced by these vertices.

Observation 2. Let A be any set of t + 2 different vertices of \tilde{G} . Let H be any K_{t+1} of \tilde{G} such that $V(H) \subseteq A$. Then the complement of $\tilde{G}[A]$ is a star with a center in the only vertex of A which does not belong to H.

Therefore, it is sufficient to check if the complement of \tilde{G} induced on these t+2 vertices is a star. If so, we get at most two different K_{t+1} 's of \tilde{G} which contain e, depending on how many centers this star has. (Notice that a star consisting of a single edge has two different centers.) Again, we can construct such complement in $\mathcal{O}(t^2)$ time.

Proof of Lemma 16 for K_q^p 's *for* $p \ge 2$ *and* $q \ge 3$. We consider any two different edges e_1 and e_2 incident to v in \tilde{G} , similarly as in the proof for K_{t+1} 's. We find all K_q^p 's of \tilde{G} which contain e_1 or e_2 .

Definition 5. Let $e = (v_0, v_1)$ be any edge of \tilde{G} . We denote by $\Gamma(e)$ a set of common neighbours of both endpoints of e. For $i \in \{0, 1\}$, we denote by $\Gamma_i(e)$ a set of all neighbours of v_i different than v_{1-i} which do not belong to $\Gamma(e)$.

Observe that the endpoints of any edge e of \tilde{G} belong to neither $\Gamma(e)$ nor $\Gamma_i(e)$. Since \tilde{G} is bounded, $|\Gamma(e)| + |\Gamma_i(e)| \le t$ for every $i \in \{0, 1\}$.

Observation 3. For any edge e of any K_q^p of \tilde{G} , $(p-2)q \leq |\Gamma(e)| \leq (p-2)q + 2$.

Because of Observation 3, it is sufficient to consider the following three cases while considering an edge $e=(v_0,v_1)\in\{e_1,e_2\}$ as a candidate for an edge of some K_q^p of \tilde{G} . We assume that both v_0 and v_1 have degree exactly t+1 in \tilde{G} since we can add artificial vertices if necessary.

1. $|\Gamma(e)| = (p-2)q + 2$ and $|\Gamma_i(e)| = q - 2$.

Since the number of all neighbours of each endpoint of e (including v_0 and v_1) is not greater than pq, we can find sought-after K_q^p using the following observation.

Observation 4. Let A be any set of pq different vertices of \tilde{G} . Let H be any K_q^p of \tilde{G} such that $V(H) \subseteq A$. Then the complement of G[A] consists of exactly p connected components, each of which consists of exactly q vertices. Moreover, the vertex set of each of these components is some color class of H.

2. $|\Gamma(e)| = (p-2)q$ and $|\Gamma_i(e)| = q$.

We check if there exists a K_q^{p-2} H' in $\tilde{G}[\Gamma(e)]$ similarly as in Case 1. If so, we check if it is possible to extend H' to a valid K_q^p , using q vertices of each set $\Gamma_i(e) \cup \{v_{1-i}\}$ as another color class. To do this, we search for $K_{t,t}$'s of \tilde{G} whose each color class is contained in $\Gamma_i(e) \cup \{v_{1-i}\}$, for some $i \in \{1,2\}$, using the following observation.

Observation 5. Let A_1 and A_2 be any two vertex-disjoint subsets of vertices of \tilde{G} , each of size exactly t+1. Let H be any $K_{t,t}$ of \tilde{G} such that $V_i(H) \subseteq A_i$, for every $i \in \{1,2\}$. Then the bipartite complement of $\tilde{G}[A_1,A_2]$ consists of two (possibly empty) stars with centers in vertices of, respectively, $A_1 \setminus V_1(H)$ and $A_2 \setminus V_2(H)$.

We check if it is possible to merge H' and found $K_{t,t}$'s into a single K_q^p of \tilde{G} . Since each set $\Gamma_i(e) \cup \{v_{1-i}\}$ contains q+1 vertices, we can find several valid K_q^p 's of \tilde{G} in this case.

3. $|\Gamma(e)| = (p-2)q + 1$ and $|\Gamma_i(e)| = q - 1$.

We check if there exists a K_a^{p-2} H' in $\tilde{G}[\Gamma(e)]$ using the following observation.

Observation 6. Let A be any set of pq + 1 different vertices of \tilde{G} . Let H be any K_q^p of \tilde{G} such that $V(H) \subseteq A$. Then the only vertex of A which does not belong to H is only either isolated vertex or nontrivial cut vertex of the complement of $\tilde{G}[A]$.

We check if both sets $\Gamma_i(e) \cup \{v_{1-i}\}$ form other color classes of sought-after K_q^p . Moreover, we check if it possible to replace some vertex of one of these sets by the only vertex of $\Gamma(e)$ which does not belong to H' to get a valid K_q^p of \tilde{G} .

Proof of Lemma 16 for K_2^p 's *for* $p \ge 3$. We find all K_2^p 's of \tilde{G} which contain v similarly to K_q^p 's. We need the following slight modifications to Observation 4 and Observation 6.

Observation 7. Let A be any subset of 2p different vertices of \tilde{G} . Let H be any K_2^p of \tilde{G} such that $V(H) \subseteq A$. Then the complement of $\tilde{G}[A]$ is a (possibly empty) matching.

Observation 8. Let A be any set of 2p+1 different vertices of \tilde{G} . Let H be any K_2^p of \tilde{G} such that $V(H) \subseteq A$. Then there exist a constant number of vertices v such that $\tilde{G}[A \setminus \{v\}]$ is a matching. Moreover, all of these vertices can be found in $\mathcal{O}(t^2)$ time.

6.4. Finding the potential functions

In this subsection, we present how to find the potential function of any forbidden subgraph of G. We use the following observations to do so.

Observation 9. Consider any triangle t of G. Then there exists exactly one potential function of t.

The potential function of any K_{t+1} of G can be calculated in $\mathcal{O}(t^2)$ time by covering the set of its vertices by triangles and by using Observation 9.

Observation 10. Consider any $H = K_{t,t}$ of G and let r_H be any potential function of H. For any real number δ , define the following potential function of H:

$$r_H^{\delta}(v) = \left\{ \begin{array}{ll} r_H(v) + \delta & \text{if } v \in V_1(H), \\ r_H(v) - \delta & \text{if } v \in V_2(H). \end{array} \right.$$

Then the set of all potential functions of H is a set of all r_H^δ over all real numbers δ .

We can find any potential function of any $H = K_{t,t}$ of G by defining the potential of some vertex of H arbitrarily which determines potentials of all other vertices of H. It is easy to see that this procedure can be implemented to run in $\Theta(t)$ time.

For $p \ge 3$, we can find the potential function of any K_q^p of G in $\mathcal{O}(t^2)$ time using Observation 9 in the same way as for K_{t+1} 's.

7. Conclusion

We presented the first polynomial algorithms for the weighted versions of the bounded restricted and K_q^p -free t-matching problems in which the weight function is vertex-induced on every forbidden subgraph. Moreover, these algorithms for the unweighted versions are faster than those known previously.

It is imaginable that our algorithms can be adapted to find maximum size or weight t-matchings which contain no complete bipartite subgraphs consisting of t+2 vertices in the graphs of maximum degree at most t+1, possibly leading to a faster algorithm for the problem of increasing connectivity by one.

Based on our result, one can suspect that the degree sequences of maximum weight restricted t-matchings form an M-concave function on the constant-parity jump system in any graph of maximum degree at most t+1 if the weight function is vertex-induced on every forbidden subgraph.

References

- [1] K. Bérczi and Y. Kobayashi. An algorithm for (n-3)-connectivity augmentation problem: Jump system approach. *Journal of Combinatorial Theory, Series B*, 102(3):565–587, 2012.
- [2] K. Bérczi and L. Végh. Restricted b-matchings in degree-bounded graphs. In *Integer Programming and Combinatorial Optimization*, pages 43–56, 2010.
- [3] C. Berge. Graphs and Hypergraphs. North-Holland Publishing Company, 1973.
- [4] G. Cornuéjols and W. Pulleyblank. A matching problem with side conditions. Discrete Mathematics, 29(2):135–159, 1980.
- [5] W. Cunningham. Matching, matroids, and extensions. Mathematical Programming, 91:515-542, 2002.
- [6] J. Edmonds. Maximum matching and a polyhedron with 0, 1-vertices. *Journal of Research of the National Bureau of Standards Section B Mathematics and Mathematical Physics*, pages 125–130, 1965.
- [7] M. Fisher, G. Nemhauser, and L. Wolsey. An analysis of approximations for finding a maximum weight hamiltonian circuit. *Operations Research*, 27(4):799–809, 1979.
- [8] H. Gabow. An efficient implementation of Edmonds' algorithm for maximum weight matching on graphs. Technical report, University of Colorado, 1975.

- [9] H. Gabow. An efficient reduction technique for degree-constrained subgraph and bidirected network flow problems. In *Proceedings of the Fifteenth Annual ACM Symposium on Theory of Computing*, pages 448–456, 1983.
- [10] Z. Galil, S. Micali, and H. Gabow. An $O(EV \log V)$ algorithm for finding a maximal weighted matching in general graphs. SIAM Journal on Computing, 15:120–130, 1986.
- [11] D. Hartvigsen. Finding maximum square-free 2-matchings in bipartite graphs. *Journal of Combinatorial Theory, Series B*, 96(5):693–705, 2006
- [12] D. Hartvigsen and Y. Li. Maximum cardinality simple 2-matchings in subcubic graphs. SIAM Journal on Optimization, 21(3):1027–1045, 2011
- [13] D. Hartvigsen and Y. Li. Polyhedron of triangle-free simple 2-matchings in subcubic graphs. Mathematical Programming, 138:43-82, 2013.
- [14] Y. Kobayashi. A simple algorithm for finding a maximum triangle-free 2-matching in subcubic graphs. *Discrete Optimization*, 7:197–202, 2010
- [15] Y. Kobayashi. Triangle-free 2-matchings and M-concave functions on jump systems. Discrete Applied Mathematics, 175:35–42, 2014.
- [16] Y. Kobayashi. Weighted triangle-free 2-matching problem with edge-disjoint forbidden triangles. In *Integer Programming and Combinatorial Optimization*, pages 280–293, 2020.
- [17] Y. Kobayashi, J. Szabo, and K. Takazawa. A proof of Cunningham's conjecture on restricted subgraphs and jump systems. *Journal of Combinatorial Theory, Series B*, 102:948–966, 2012.
- [18] Y. Kobayashi and X. Yin. An algorithm for finding a maximum t-matching excluding complete partite subgraphs. *Discrete Optimization*, 9(2):98–108, 2012.
- [19] M. Makai. On maximum cost K_{t,t}-free t-matchings of bipartite graphs. SIAM Journal on Discrete Mathematics, 21:349–360, 2007.
- [20] Y. Nam. Matching Theory: Subgraphs with Degree Constraints and other Properties. PhD thesis, University of British Columbia, 1994.
- [21] K. Paluch, K. Elbassioni, and A. van Zuylen. Simpler approximation of the maximum asymmetric traveling salesman problem. In 29th International Symposium on Theoretical Aspects of Computer Science, pages 501–506, 2012.
- [22] K. Paluch and M. Wasylkiewicz. Restricted t-matchings via half-edges. In 29th Annual European Symposium on Algorithms, pages 73:1–73:17, 2021.
- [23] K. Paluch and M. Wasylkiewicz. A simple combinatorial algorithm for restricted 2-matchings in subcubic graphs via half-edges. *Information Processing Letters*, 171, 2021.
- [24] G. Pap. Combinatorial algorithms for matchings, even factors and square-free 2-factors. Mathematical Programming, 110:57-69, 2007.
- [25] K. Takazawa. A weighted $K_{t,t}$ -free t-factor algorithm for bipartite graphs. Mathematics of Operations Research, 34(2):351–362, 2009.
- [26] L. Végh. Augmenting undirected node-connectivity by one. SIAM Journal on Discrete Mathematics, 25(2):695-718, 2011.
- [27] O. Vornberger. Easy and hard cycle covers. Technical report, Universität Paderborn, 1980.