On Smale's 17th problem over the reals

Andrea Montanari^{*} Eliran Subag[†]

May 6, 2024

Abstract

We consider the problem of efficiently solving a system of n non-linear equations in \mathbb{R}^d . Addressing Smale's 17th problem stated in 1998, we consider a setting whereby the n equations are random homogeneous polynomials of arbitrary degrees. In the complex case and for n = d-1, Beltrán and Pardo proved the existence of an efficient randomized algorithm and Lairez recently showed it can be de-randomized to produce a deterministic efficient algorithm. Here we consider the real setting, to which previously developed methods do not apply. We describe an algorithm that efficiently finds solutions (with high probability) for $n = d - O(\sqrt{d \log d})$. If the maximal degree is very large, we also give an algorithm that works up to n = d - 1.

Contents

| 1 | Introduction and main result | 2 |
|---|--|-----------|
| | 1.1 Approximate solutions | 4 |
| | 1.2 Optimization | 4 |
| | 1.3 Hessian descent: the case of moderate $p_{\text{max}} \leq d^2$ | 5 |
| | 1.4 Brute-force search: the case of large $p_{\text{max}} > d^2 \dots \dots$ | 6 |
| | 1.5 Smale's 17th problem in the complex case | 7 |
| 2 | Preliminaries | 8 |
| 3 | Bounds on the Hessian | 14 |
| 4 | Minimal singular value at solutions | 21 |
| | 4.1 Upper bound on volume in expectation | 22 |
| | 4.2 Lower bound on the volume of a neighborhood | 23 |
| | 4.3 Proof of Proposition 4.1 | 25 |
| | 4.4 Proof of Proposition 4.2 | 25 |
| 5 | Good events | 26 |
| 6 | Sufficient conditions for approximate solutions | 27 |
| | 6.1 The case of moderate $p_{\max} \leq d^2$ | 31 |
| | 6.2 The case of large $p_{\text{max}} > d^2$ | 31 |
| 7 | Sub-routines and their complexity | 32 |

^{*}Department of Statistics and Department of Mathematics, Stanford University

[†]Department of Mathematics, Weizmann Institute of Science

| 8 | Analysis of Hessian descent: Proof of Theorem 1 | 34 |
|---|--|-----------|
| | 8.1 Proof of Theorem 1 | 30 |
| 9 | Analysis of brute-force search: Proof of Theorem 2 | 37 |
| A | Tools: Linear algebra | 39 |
| В | Tools: Random matrix theory | 39 |

1 Introduction and main result

In 1998 Steve Smale published a list of 'Mathematical problems for the next century' [Sma98]. Quoting from the original, his 17th problem asked:

"Can a zero of n complex polynomial equations in n unknowns be found approximately, on the average, in polynomial time with a uniform algorithm?"

The precise setting of the problem, originally introduced in the so-called Bézout series [SS93a, SS93b, SS93c, SS96, SS94] of Shub and Smale, is as follows. First, the assumption that the polynomials are homogeneous is made. In this case, for the set of solutions to be discrete, if the number of equations is n, the number of variables needs to be d = n + 1, and one either has to restrict to solutions in the projective space as in the Bézout series, or equivalently to solutions on the unit sphere as we shall in the current work. Second, the question is a probabilistic one. Namely, it concerns a random polynomial system $\mathbf{F}(\mathbf{x}) := (F_1(\mathbf{x}), \ldots, F_n(\mathbf{x}))$ consisting of independent polynomials

$$F_i(\boldsymbol{x}) := \sum_{k_1 + \dots + k_d = p_i} \left(\frac{p_i!}{k_1! \cdots k_d!} \right)^{\frac{1}{2}} a_{k_1, \dots, k_d}^{(i)} x_1^{k_1} \cdots x_d^{k_d},$$
(1)

where $\boldsymbol{x} = (x_1, \ldots, x_d) \in \mathbb{C}^d$, $a_{k_1, \ldots, k_d}^{(i)}$ are i.i.d. complex standard Gaussian variables and p_1, \ldots, p_n is some sequence of degrees. By classical results of Abel and Galois, polynomials generically do not have closed-form solutions. The 17th problem therefore asks for an algorithm that finds an 'approximate solution'. The latter is defined as a point on the unit sphere such that (projected) Newton's method started from it converges immediately and quadratically fast to a solution, see Section 1.1. The model of computation allows storing real numbers and complexity is measured by the number of read/writes and elementary operations with real numbers (formally, a Blum-Shub-Smale machine [BSS89]). To avoid technicalities we will also assume that the square root of a positive real number can be computed in a single time unit. The algorithm receives as input the coefficients of the system \boldsymbol{F} and is required to have a polynomial time complexity on average in $N := \sum_{i=1}^{n} {d+p_i-1 \choose p_i}$, the total number of coefficients. Roughly speaking, a uniform algorithm is an algorithm that can be implemented as a single program, whereas for non-uniform algorithms the implementation may depend on the input (in our setting, on d, n and p_1, \ldots, p_n). For the precise definition, see [BCSS98] and [BP09, Section 1.3].

In its original form, Smale's 17th problem was recently solved after several major breakthroughs [BP09, BP08, BS09, BC11, Lai17] (see more below in Section 1.5). We, however, will be interested in the real version of it. Indeed, after stating the problem in [Sma98], Smale also mentioned that the same problem is very interesting, and even more difficult, in the real case. Namely, when the coefficients $a_{k_1,\ldots,k_d}^{(i)}$ are assumed to be real Gaussian variables instead of complex and one searches for solutions on the unit sphere in \mathbb{R}^d instead of \mathbb{C}^d . We note that in the real case

the specific choice of the combinatorial factor above makes $F_i(\boldsymbol{x})$ rotationally invariant in law. Namely, $F_i(\boldsymbol{x}) \stackrel{\mathcal{L}}{=} F_i(O\boldsymbol{x})$ as processes for any orthogonal matrix O, where $\stackrel{\mathcal{L}}{=}$ denotes equality in law. Henceforth, we shall assume this setting. Moreover, we will consider an under-determined variant of Smale's 17th problem in the real case, by allowing the number of equations n to be strictly smaller than d-1. We still keep the notation n for the number of equations and d for the number of variables. Our main results are polynomial time algorithms to find approximate solutions, which we now state in two separate regimes. We denote by $p_{\max} = \max_{i \leq n} p_i$ the maximal degree, by $\boldsymbol{a} := \left(\left(\frac{p_i!}{k_1! \cdots k_d!} \right)^{\frac{1}{2}} a_{k_1, \dots, k_d}^{(i)} \right)_{i \leq n, k_1 + \cdots + k_d = p_i}$ the set of real Gaussian coefficients, and by $\mathbb{S}^{d-1} \subset \mathbb{R}^d$ the unit sphere. From now on, we shall always assume that $p_i \geq 2$ for all i, since for $p_i = 1$ the zero set of $F_i(\boldsymbol{x})$ is the orthogonal space to $(a_1^{(i)}, \dots, a_d^{(i)})$ and, thanks to rotational invariance, via a pre-processing stage by a change of basis we may remove $F_i(\boldsymbol{x})$ from the system and reduce the dimension by one.

Theorem 1 (Moderate p_{\max}). Assume that $p_{\max} \leq d^2$. For some absolute constants A, C, C_0 , if $n = \lfloor d - A(d \log d)^{1/2} \rfloor$ then there exists an algorithm which takes as inputs the random coefficients **a** and returns as output $\mathbf{x}^{\text{alg}} \in \mathbb{S}^{d-1}$ such that the following holds:

(i) The real complexity χ of the algorithm is bounded by

$$\chi \le C_0 N d^{9/2} p_{\max}^4 (1 + p_{\max}/d) \log(p_{\max})^2.$$
⁽²⁾

(ii) The output \mathbf{x}^{alg} is an approximate solution with probability at least $1 - Ce^{-d/C}$.

Theorem 2 (Large p_{\max}). Assume that $p_{\max} > d^2$. For some absolute constants C, C_0 , if n = d - 1 then there exists an algorithm which takes as inputs the random coefficients **a** and returns as output $\mathbf{x}^{\text{alg}} \in \mathbb{S}^{d-1}$ such that the following holds:

(i) The real complexity χ of the algorithm is bounded by

$$\chi \le N dp_{\max} \cdot \left(C_0 d^5 p_{\max}^2 \log(p_{\max}) \right)^d \le N^3 p_{\max}^3 (d!)^2 \left(C_0 d^5 \log(p_{\max}) \right)^d.$$
(3)

(ii) The output \mathbf{x}^{alg} is an approximate solution with probability at least $1 - Cd^{-\frac{1}{2}}$.

The two algorithms will be described in Sections 1.3 and 1.4 below. They output an approximate solution on an event of high probability which we describe in Section 5. While in the complex case Smale's problem seem to ask for an algorithm whose complexity is bounded on average but always finds an approximate solution, in the real case this is not possible. Indeed, if p_i is even, with positive probability $\mathbf{F}_i(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$. In fact, only very recently it was shown [Sub23] that a solution exists with high probability as $d \to \infty$ (previously, this was known only in the i.i.d. case $p_i \equiv p$ [Wsc05]). We emphasize that our algorithms are not randomized, in the sense that they do not use a source of randomness (other than the coefficients as input).

The main discrepancy between our results and Smale's problem over the reals is, of course, that Theorem 1 about the moderate case does not work up to n = d-1, but only for $n = d-O(\sqrt{d \log d})$. To the best of our knowledge, however, the real case of Smale's problem is completely open and this is a significant progress. It also seems to be more difficult than the complex case which was solved only recently after intensive research going back at least to the works of Shub and Smale from the early 90s (see Section 1.5).

1.1 Approximate solutions

Since we work on the unit sphere $\mathbb{S}^{d-1} := \{ \boldsymbol{x} \in \mathbb{R}^d : \|\boldsymbol{x}\| = 1 \}$, we define Newton's method by taking a step in an orthogonal direction and projecting back to the sphere. This definition is basically a variant of Newton's Method in the projective space, defined by Shub [Shu93] for polynomial systems of n = d - 1 equations in d complex variables. Here we also allow for n < d - 1.

Given a matrix $\mathbf{A} \in \mathbb{R}^{n_2 \times n_1}$ or a linear operator $\mathbf{A} : V_1 \to V_2$, with $\dim(V_i) = n_i$ (we will often identify the two objects), for $n_1 \leq n_2$ we denote by $\sigma_{\min}(\mathbf{A}) := \min\{\|\mathbf{A}\mathbf{x}\| : \|\mathbf{x}\| = 1\}$ the minimum singular value of \mathbf{A} . For $n_1 > n_2$, $\sigma_{\min}(\mathbf{A}) := \sigma_{\min}(\mathbf{A}^{\mathsf{T}})$.

minimum singular value of \mathbf{A} . For $n_1 > n_2$, $\sigma_{\min}(\mathbf{A}) := \sigma_{\min}(\mathbf{A}^{\mathsf{T}})$. We denote by $\mathbf{DF}(\mathbf{x}) := (\partial_j F_i(\mathbf{x}))_{ij} \in \mathbb{R}^{n \times d}$ the differential or Jacobian matrix and by $\mathsf{T}_{\mathbf{x}} \subset \mathbb{R}^d$ the orthogonal space to \mathbf{x} . Given a point $\mathbf{x} \in \mathbb{S}^{d-1}$ we define Newton's operator as follows. The matrix $\mathbf{DF}(\mathbf{x})$ defines a linear operator and its restriction to a subspace is defined as an operator in the obvious way. If $\sigma_{\min}(\mathbf{DF}(\mathbf{x})|_{\mathsf{T}_{\mathbf{x}}}) > 0$, let $\mathbf{v} \in \mathbb{R}^d$ be the unique solution to

$$F(x) + DF(x)v = 0 \quad \text{subject to} \quad v \perp x, \ v \perp \ker(DF(x)), \tag{4}$$

where ker $(\boldsymbol{A}) := \{ \boldsymbol{y} : \boldsymbol{A}\boldsymbol{y} = 0 \}$, and define

$$\Phi_{\rm NM}(\boldsymbol{x}) := \frac{\boldsymbol{x} + \boldsymbol{v}}{\|\boldsymbol{x} + \boldsymbol{v}\|_2} \,. \tag{5}$$

Otherwise, if $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x})|_{\mathsf{T}_{\boldsymbol{x}}}) = 0$, arbitrarily define

$$\Phi_{\mathrm{NM}}(oldsymbol{x}):=oldsymbol{x}$$
 .

Definition 1.1 (Approximate solution). Let $\mathbf{x} \in \mathbb{S}^{d-1}$ and define the sequence $\mathbf{x}^0 = \mathbf{x}$, $\mathbf{x}^{i+1} = \Phi_{\text{NM}}(\mathbf{x}^i)$ by Newton's method. We say that \mathbf{x} is an approximate solution of the system \mathbf{F} if there exists $\mathbf{z} \in \mathbb{S}^{d-1}$ such that $\mathbf{F}(\mathbf{z}) = 0$ and $\|\mathbf{x}^i - \mathbf{z}\|_2 \leq 2^{1-2^i} \|\mathbf{x}^0 - \mathbf{z}\|_2$ for all $i \geq 0$.

1.2 Optimization

We define the "energy function"

$$H(\boldsymbol{x}) := \frac{1}{2} \|\boldsymbol{F}(\boldsymbol{x})\|^2.$$
(6)

Of course, F(x) = 0 if and only if H(x) = 0. In other words, the solutions of the polynomial system are exactly the minimizers of the energy function. This simple observation allows us to change perspective and try to algorithmically minimize a real-valued function over the sphere \mathbb{S}^{d-1} , instead of working with a vector-valued function.

Problems not very different from this have been studied recently in the context of spin glass models. In the terminology of spin glass theory, $F_i(\mathbf{x})$ as in (1) is the energy function of the spherical *p*-spin model (with $p_i = p$), up to normalization. For the spin glass model, it is customary to consider $F_i(\mathbf{x})$ as a function on the sphere of radius \sqrt{d} and multiply it by $d^{-(p-1)/2}$. More generally, for mixed *p*-spin the energy is defined as a linear combinations of the pure *p*-spin energies with deterministic coefficients. Given a model, those coefficients are fixed and one is interested in the $d \to \infty$ asymptotics.

In [Sub18] an algorithm for minimizing the energy of a spherical mixed *p*-spin model by Hessian descent was proposed and analyzed. The algorithm constructs a discrete path $\boldsymbol{x}_0, \ldots, \boldsymbol{x}_k$ from the origin to the sphere with $\|\boldsymbol{x}_i\| = \sqrt{i \cdot d/k}$, whose increments at each step $\boldsymbol{x}_{i+1} - \boldsymbol{x}_i$ are orthogonal to the position \boldsymbol{x}_i , and are approximately in the span of the eigenvectors corresponding to the minimal eigenvalues of the Hessian of the energy. Related Approximate Message Passing algorithms for Ising

models, which use the same energy functions but on the hyper-cube $\{+1, -1\}^d$, were proposed and studied in [Mon19, EAMS21]. Both types of algorithms [Sub18, Mon19, EAMS21] were proved to be optimal within a certain large class of appropriately defined Lipschitz algorithms in [HS21].

In light of the close connections to the spherical spin glass models, it is natural to wonder whether the Hessian descent algorithm of [Sub18] can be adapted to minimize $H(\mathbf{x})$. In an earlier work [MS23] we studied this question for $F_i(\mathbf{x})$ that correspond to general mixed models. In other words, the $F_i(\mathbf{x})$ studied in [MS23] are given by linear combinations of the functions as in (1). The functions $F_i(\mathbf{x})$ were also allowed to have a zero order term. Namely, one of the terms in the linear combination was given by $\gamma_0 W_i$ where $\gamma_0 \geq 0$ and $W_i \sim \mathsf{N}(0,1)$. We assumed there that the functions $F_i(\mathbf{x})$ are i.i.d. and our goal was only to find points \mathbf{x}_* such that $H(\mathbf{x}_*) = o(n)$ asymptotically, while the typical value is $H(\mathbf{x}) = \Theta(n)$ for most points on the sphere. The objective $H(\mathbf{x}_*) = o(n)$ is better suited for the methods developed for algorithmic optimization of spin glasses [Sub18, Mon19, EAMS21].

In the present work, we adapt the Hessian descent algorithm to the problem of finding approximate solutions.

1.3 Hessian descent: the case of moderate $p_{\text{max}} \leq d^2$

In the present paper, the functions $F_i(\boldsymbol{x})$ are homogeneous, and therefore changing the norm $\|\boldsymbol{x}\|$ does not change the problem (apart from a scaling factor). As mentioned above, the earlier papers [Sub18, MS23] construct a path in the interior of the unit ball, hence effectively changing the energy landscape as $\|\boldsymbol{x}\|$ increases. This is irrelevant in the homogeneous case. Instead, we may take an orthogonal step in \mathbb{R}^d and project back to the unit sphere \mathbb{S}^{d-1} .

Moreover, here we wish to achieve energy values much smaller than in the spin glass optimization problems, and we therefore have to take steps which depend on the value of the energy at each step and run the algorithm with a much larger number of steps. Below $C_1, c_0 > 0$ are absolute constants which will be determined in Lemma 2.1 and the proof of Theorem 1. In pseudo-code, the algorithm we use in the moderate $p_{\text{max}} \leq d^2$ setting of the Theorem 1 is given below.

Algorithm 1: Hessian descent

Input: The coefficients $\boldsymbol{a} := \left(\left(\frac{p_i!}{k_1! \cdots k_d!} \right)^{\frac{1}{2}} a_{k_1, \dots, k_d}^{(i)} \right)_{i \le n, k_1 + \dots + k_d = p_i}$, number of iterations kOutput: $\boldsymbol{x}^{\text{HD}} \in \mathbb{S}^{d-1}$ Initialize $\boldsymbol{x}_0 = (1, 0, \dots, 0) \in \mathbb{S}^{d-1}$; for $i \in \{0, \dots, k-1\}$ do Find $\boldsymbol{v}_i \perp \boldsymbol{x}_i$, $\|\boldsymbol{v}_i\| = 1$, s.t.: $\langle \nabla^2 H(\boldsymbol{x}_i), \boldsymbol{v}_i^{\otimes 2} \rangle \leq \frac{1}{2} \min_{\boldsymbol{u} \perp \boldsymbol{x}_i, \|\boldsymbol{u}\| = 1} \langle \nabla^2 H(\boldsymbol{x}_i), \boldsymbol{u}^{\otimes 2} \rangle$; $\delta_i = \left(\frac{1}{30C_1} \frac{1}{p_{\max}^4 \log(p_{\max})} \frac{\sqrt{(d-n)H(\boldsymbol{x}_i)}}{d} \wedge \frac{1}{p_{\max}} \right)^{1/2}$; $s_i = \text{sign} \left(H(\boldsymbol{x}_i + \delta_i \boldsymbol{v}_i) - H(\boldsymbol{x}_i - \delta_i \boldsymbol{v}_i)\right)$; $\boldsymbol{y}_{i+1} = \boldsymbol{x}_i - s_i \delta_i \boldsymbol{v}_i$; $\boldsymbol{x}_{i+1} = \frac{\boldsymbol{y}_{i+1}}{\|\boldsymbol{y}_{i+1}\|}$; if $H(\boldsymbol{x}_{i+1}) \leq p_{\max}^{-cod}$ then \mid return $\boldsymbol{x}^{\text{HD}} = \boldsymbol{x}_{k}$ Above we use the convention that sign(0) = 1.

Remark 1.1. In the pseudo-code above we allowed ourselves to be imprecise, to improve readability. Specifically, we did not explain how we 'find' the vector v_i or when it is even possible. More accurately, our algorithm will use a sub-routine which we define in Section 7. On an event of overwhelmingly high probability, uniformly in x_i , the sub-routine outputs a vector v_i as required.

To explain the basic idea, consider a Taylor expansion of $H(\mathbf{x})$ of degree 3 around \mathbf{x}_i . The choice of the sign s_i in the pseudocode above guarantees that the first and third order terms in the expansion may only decrease the energy as we move from \mathbf{x}_i to \mathbf{y}_{i+1} . Hence, assuming an appropriate bound on the fourth order derivative along the path from from \mathbf{x}_i to \mathbf{y}_{i+1} , we may conclude a decrease in the energy by an amount which we can explicitly bound using the condition

$$\langle \nabla^2 H(\boldsymbol{x}_i), \boldsymbol{v}_i^{\otimes 2} \rangle \leq rac{1}{2} \min_{\boldsymbol{u} \perp \boldsymbol{x}_i, \|\boldsymbol{u}\|=1} \langle \nabla^2 H(\boldsymbol{x}_i), \boldsymbol{u}^{\otimes 2} \rangle.$$

Since $F_i(\boldsymbol{x})$ are homogeneous, projecting \boldsymbol{y}_{i+1} to \boldsymbol{x}_{i+1} can only reduce the energy.

The algorithm of Theorem 1 runs the Hessian descent algorithm for $k = Cd^{3/2}p_{\text{max}}^4 \log(p_{\text{max}})^2$ iterations, for some absolute constant C. We will prove that the algorithm outputs an approximate solution when a certain 'good' event \mathcal{E} holds. The event \mathcal{E} is defined in Section 5 by: (1) upper bounds on the derivatives of $H(\mathbf{x})$ of different orders and certain Lipschitz constants uniformly over \mathbb{S}^{d-1} ; (2) a uniform lower bound on the smallest eigenvalue of $\nabla^2 H(\mathbf{x})$ restricted to the orthogonal space; and (3) a lower bound on the singular value of $DF(\mathbf{x})$ uniformly over the solutions of the system $F(\mathbf{x}) = 0$.

Using (3), we will show quantitatively in Section 6 that in order for \boldsymbol{x} to be an approximate solution, it is sufficient for it to be close enough to some exact solution $\boldsymbol{\alpha}$. In our analysis of the Hessian descent algorithm in Theorem 4, we will prove bounds for $H(\boldsymbol{x}_i)$ and $\|\boldsymbol{x}_i - \boldsymbol{x}_j\|$, which holds for any $i \geq 1$ (even if we produce an infinite sequence \boldsymbol{x}_i by running the Hessian descent indefinitely). These bounds imply that $\boldsymbol{x}_{\infty} = \lim_{i \to \infty} \boldsymbol{x}_i$ is well-defined and is a solution $\boldsymbol{F}(\boldsymbol{x}_{\infty}) = 0$, and that for k as above, for some $i \leq k$, $\|\boldsymbol{x}_i - \boldsymbol{x}_{\infty}\|$ is small enough to conclude that \boldsymbol{x}_i is an approximate solution.

1.4 Brute-force search: the case of large $p_{\text{max}} > d^2$

The algorithm in this case is much simpler. Since the input size

$$N \ge \binom{p_{\max} + d - 1}{d - 1} \ge \left(\frac{p_{\max}}{d}\right)^{d - 1},\tag{7}$$

is extremely large, we can perform a brute-force search by sampling the space on a dense enough net. We look for a point where the norm, or energy, is small and the minimal singular value of the projected differential is not too small. As the minimal singular value $\sigma_{\min}(\mathbf{A})$ of a matrix \mathbf{A} is equal to the minimal solution of the characteristic polynomial, we can only approximate it. What we mean below by an approximation of $\sigma_{\min}(\mathbf{DF}(\mathbf{x})|_{\mathsf{T}_{\mathbf{x}}})$ will be made precise in Section 7 where we will also define a sub-routine that computes the approximation. In the proof of Theorem 2 we will show that, for appropriate choice of the parameters δ and η , with high probability the algorithm in the following pseudo-code returns an approximate solution.

Algorithm 2: Brute-force search

Data: The coefficients $\boldsymbol{a} := \left(\left(\frac{p_i!}{k_1!\cdots k_d!}\right)^{\frac{1}{2}} a_{k_1,\dots,k_d}^{(i)} \right)_{i \le n, k_1 + \dots + k_d = p_i}$, parameters $\delta, \bar{\eta}$ **Result:** $\boldsymbol{x}^{\text{BF}} \in \mathbb{S}^{d-1}_{\delta}$ Set $\delta_0 = \frac{\delta}{2C_0 d^5 p_{\max}^3 \log(p_{\max})}, t = \frac{\delta}{d^4 p_{\max}^2 \sqrt{\log(p_{\max})}};$ for $\boldsymbol{y} \in \mathcal{N} := \{\boldsymbol{z} = (z_1, \dots, z_d) \in [-1, 1]^d : z_i / \delta_0 \in \mathbb{Z}\}$ do $\begin{vmatrix} \boldsymbol{x} = \boldsymbol{y} / \| \boldsymbol{y} \|; \\ \text{Compute } s_{\min}(\boldsymbol{x}) \text{ an approximation of } \sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x}) \|_{\mathsf{T}_{\boldsymbol{x}}}); \\ \text{if } \| \boldsymbol{F}(\boldsymbol{x}) \| \le t, s_{\min}(\boldsymbol{x}) \ge \frac{\bar{\eta}}{d^{7/4}} \text{ then} \\ \mid \text{ return } \boldsymbol{x}^{\text{BF}} = \boldsymbol{x} \\ \text{ end} \\ \text{return } \boldsymbol{x}^{\text{BF}} = (1, 0, \dots, 0)$

1.5 Smale's 17th problem in the complex case

The study of Smale's 17th problem actually started in Shub and Smale's 'Bézout series' [SS93a, SS93b, SS93c, SS96, SS94], several years before it was posed in [Sma98]. This sequence of papers studies a homotopy continuation method, which in turn builds on numerical analysis approaches that were studied since the seventies [Dre78, GZ79, Kel78]. The basic idea of this approach is to define an interpolation $t\mathbf{F} + (1-t)\mathbf{F}$, $t \in [0,1]$, between the system one wishes to solve \mathbf{F} and another system $ar{F}$ of homogeneous polynomials of the same degrees with a root x_0 which we know of. If the system $ar{F}$ is not singular, then a.s. the number of roots in the projective space of each of the systems F = 0 and $\bar{F} = 0$ is exactly the Bézout number $\prod_{i=1}^{N-1} p_i$. On the unit sphere the number of solutions is double. These roots come in pairs $(\boldsymbol{x}(0), \boldsymbol{x}(1))$, where $\boldsymbol{x}(0)$ solves $\boldsymbol{F}(\boldsymbol{x}(0)) = 0$ and x(1) solves F(x(1)) = 0. Each pair is connected by a smooth path $[0,1] \ni t \mapsto x(t)$ of solutions to $tF + (1-t)\bar{F} = 0$. Homotopy based algorithms start at time t = 0 with \bar{F} and x(0) and iteratively increase the time parameter t by a small step while trying to keep track of the root by an iteration of Newton's method, in order to end at time t = 1 with F and an approximate root x_1 . The size of the step was analyzed in terms of a *condition number* in the Bézout series and in [BP11, BC11, Shu09, ABB⁺16]. However, major difficulties remained in how to choose the system F and root x_0 to start from.

In the last paper in the Bézout series [SS94], Shub and Smale proved that there exists a good system and a root to start from, but their argument was not constructive and no practical way (an algorithm) to pick such a system and a root was provided. In major breakthroughs [BP08, BP09], Beltrán and Pardo had an ingenious idea for how to choose a good system and a starting solution at random, which led to a randomized polynomial time algorithm (on average). This solved Smale's problem, up to the point of using randomization as opposed to a deterministic algorithm. Another breakthrough was made by Bürgisser and Cucker [BC11] who performed a smoothed analysis of the algorithm of Beltrán and Pardo, leading to a deterministic algorithm of (almost polynomial) complexity $N^{O(\log \log N)}$. Finally, Lairez [Lai17] found a remarkable way to de-randomize the algorithm of Beltrán and Pardo by using the random input itself to generate the initial pair. The combination of those works gave a solution to Smale's problem in the complex case. Important improvements to the latter algorithms were recently established in [BCL23, Lai20].

2 Preliminaries

For any ℓ , let $\mathbf{G}^{(\ell)} = (G_{i_1,\dots,i_{p_\ell}}^{(\ell)})_{i_1,\dots,i_{p_\ell} \leq d}$ be a tensor with i.i.d. entries $G_{i_1,\dots,i_{p_\ell}}^{(\ell)} \sim \mathsf{N}(0,1)$. We will also assume that $\mathbf{G}^{(\ell)}$ are independent for different values of ℓ . We define $\overline{\mathbf{G}}^{(\ell)} = (\overline{G}_{i_1,\dots,i_{p_\ell}}^{(\ell)})_{i_1,\dots,i_{p_\ell} \leq d}$ the symmetrization of $\mathbf{G}^{(\ell)}$ by

$$\overline{G}_{i_1,\dots,i_{p_\ell}}^{(\ell)} := (p_\ell!)^{-1} \sum_{\pi \in S_{p_\ell}} G_{i_{\pi(1)},\dots,i_{\pi(p_\ell)}}^{(\ell)}.$$
(8)

Observe that

$$\left(\frac{p_{\ell}!}{k_{1}!\cdots k_{d}!}\right)^{\frac{1}{2}}a_{k_{1},\dots,k_{d}}^{(\ell)} \quad \text{and} \quad \sum_{\mathcal{J}(k_{1},\dots,k_{d})}\overline{G}_{i_{1},\dots,i_{p_{\ell}}}^{(\ell)} = \sum_{\mathcal{J}(k_{1},\dots,k_{d})}G_{i_{1},\dots,i_{p_{\ell}}}^{(\ell)} \tag{9}$$

have the same law, where $\mathcal{J}(k_1, \ldots, k_d)$ is the set of indices i_1, \ldots, i_{p_ℓ} such that $x_{i_1} \cdots x_{i_{p_\ell}} = x_1^{k_1} \cdots x_d^{k_d}$. Hence we can assume that they are defined on the same probability space such that both sides of (9) are equal and thus

$$F_{\ell}(\boldsymbol{x}) = \langle \boldsymbol{G}^{(\ell)}, \boldsymbol{x}^{\otimes p_{\ell}} \rangle = \langle \overline{\boldsymbol{G}}^{(\ell)}, \boldsymbol{x}^{\otimes p_{\ell}} \rangle.$$
(10)

We will often work with this representation for the random polynomials. As the representation in (1) requires less variables to describe $F_{\ell}(\boldsymbol{x})$, it is used to define the input in the algorithmic problem. Yet another equivalent way to describe the polynomials is through their covariance function

$$\mathbb{E}\left[F_i(\boldsymbol{x}^1)F_j(\boldsymbol{x}^2)\right] = \delta_{ij}\xi_i\left(\langle \boldsymbol{x}^1, \boldsymbol{x}^2\rangle\right), \quad \text{where } \xi_i(t) := t^{p_i}.$$
(11)

Recall that we denote by $DF(\mathbf{x}) := (\partial_j F_i(\mathbf{x}))_{ij} \in \mathbb{R}^{n \times d}$ the differential or Jacobian matrix and by $\mathsf{T}_{\mathbf{x}} \subset \mathbb{R}^d$ the orthogonal space to \mathbf{x} , which we identify with the tangent space to the sphere of radius \sqrt{q} , $\mathbb{S}^{d-1}(\sqrt{q})$ at \mathbf{x} , when $\|\mathbf{x}\|_2^2 = q > 0$. We often view $\nabla^k F_1(\mathbf{x}) = (\partial_{i_1,\ldots,i_k} F_1(\mathbf{x}))_{i_1\ldots,i_k \leq d}$ as a k-th order tensor $\nabla^k F_1(\mathbf{x}) \in (\mathbb{R}^d)^{\otimes k}$ and $\nabla^k F(\mathbf{x}) = (\partial_{i_1,\ldots,i_k} F_\ell(\mathbf{x}))_{\ell \leq n,i_1\ldots,i_k \leq d}$ as a (k + 1)-th order tensor $\nabla^k F(\mathbf{x}) \in \mathbb{R}^n \otimes (\mathbb{R}^d)^{\otimes k}$. The operator norm of a tensor $\mathbf{T} \in \mathbb{R}^{n_1} \otimes \cdots \otimes \mathbb{R}^{n_k}$ is

$$\|\boldsymbol{T}\|_{\mathrm{op}} := \max_{\boldsymbol{v}_1 \in \mathbb{S}^{n_1 - 1}} \cdots \max_{\boldsymbol{v}_k \in \mathbb{S}^{n_k - 1}} \langle \boldsymbol{T}, \boldsymbol{v}_1 \otimes \cdots \otimes \boldsymbol{v}_k \rangle.$$
(12)

Finally, we denote by $\mathsf{B}^d(r) := \{ \boldsymbol{x} \in \mathbb{R}^d : \|\boldsymbol{x}\| \leq r \}$ the ball of radius r, and let $\mathsf{B}^d(r_1, r_2) := \mathsf{B}^d(r_2) \setminus \mathsf{B}^d(r_1)$.

Throughout, we will write $\boldsymbol{W} \sim \mathsf{GOE}(N)$ if $\boldsymbol{W} = \boldsymbol{W}^{\mathsf{T}}$ and $(W_{ij})_{i \leq j \leq N}$ are independent with $W_{ii} \sim \mathsf{N}(0,2), W_{ij} \sim \mathsf{N}(0,1)$ for i < j. We write $\boldsymbol{Z} \sim \mathsf{GOE}(M,N)$ if $(Z_{ij})_{i \leq M,j \leq N}$ are independent with $Z_{ij} \sim \mathsf{N}(0,1)$. It is useful to recall that (under the present setting) $F_{\ell}(\boldsymbol{x})$ and its derivatives are jointly Gaussian and

$$\mathbb{E}\left\{\frac{\partial}{\partial x_{i_1}}\cdots\frac{\partial}{\partial x_{i_k}}F_{\ell}(\boldsymbol{x})\frac{\partial}{\partial y_{j_1}}\cdots\frac{\partial}{\partial y_{j_m}}F_{\ell}(\boldsymbol{y})\right\} = \frac{\partial}{\partial x_{i_1}}\cdots\frac{\partial}{\partial x_{i_k}}\frac{\partial}{\partial y_{j_1}}\cdots\frac{\partial}{\partial y_{j_m}}\mathbb{E}\left\{F_{\ell}(\boldsymbol{x})F_{\ell}(\boldsymbol{y})\right\}.$$
 (13)

Our first preliminary result provides bounds on the norm of F(x) and its derivatives.

Lemma 2.1. Assume that $n \leq d$. Then, for any $k_{\max} \geq 2$ there exist constants C_*, C_1 depending on k_{\max} , such that, with probability at least $1 - C_* \exp(-d/C_*)$

$$\max_{\boldsymbol{x}\in\mathsf{B}^{d}(1)}\|\boldsymbol{F}(\boldsymbol{x})\|_{2} \leq C_{1}\sqrt{d\log(p_{\max})},$$
(14)

$$\max_{\boldsymbol{x}\in\mathsf{B}^{d}(1)}\|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\|_{\mathrm{op}} \leq C_{1}p_{\mathrm{max}}\sqrt{d\log(p_{\mathrm{max}})}\,,\tag{15}$$

$$\max_{\boldsymbol{x}\in\mathsf{B}^{d}(1)}\left\|\nabla^{k}\boldsymbol{F}(\boldsymbol{x})\right\|_{\mathrm{op}} \leq C_{1}p_{\mathrm{max}}^{k}\sqrt{d\log p_{\mathrm{max}}}, \quad \forall 2 \leq k \leq k_{\mathrm{max}},$$
(16)

$$\max_{\boldsymbol{x}\in\mathsf{B}^{d}(1)}\left\|\nabla^{k}H(\boldsymbol{x})\right\|_{\mathrm{op}} \leq C_{1}p_{\mathrm{max}}^{k}d\log p_{\mathrm{max}}. \quad \forall 1\leq k\leq k_{\mathrm{max}}.$$
(17)

In fact, it will be clear from the proof that the last two inequalities follow for $2 \le k \le K$ with general K, if one allows the constants C_*, C_1 to depend on K.

Proof. It will be enough to prove a bound on the probability for each of the inequalities (14)-(17) separately, and conclude the lemma by a union bound.

Proof of Eq. (14). Consider the following Gaussian process indexed by $u \in \mathbb{S}^{n-1}$, $x \in \mathbb{S}^{d-1}$:

$$Z_0(\boldsymbol{u}, \boldsymbol{x}) := \langle \boldsymbol{u}, \boldsymbol{F}(\boldsymbol{x}) \rangle, \qquad (18)$$

and notice that, using homogeneity,

$$\max_{\boldsymbol{x}\in\mathsf{B}^{d}(1)}\|\boldsymbol{F}(\boldsymbol{x})\|_{2} = \max_{\boldsymbol{u}\in\mathbb{S}^{n-1},\boldsymbol{x}\in\mathbb{S}^{d-1}}Z_{0}(\boldsymbol{u},\boldsymbol{x}).$$
(19)

With $e_1 = (1, 0, ..., 0)$ denoting the standard basis element, define the spherical cap $\operatorname{Cap}_d(\pi/4) := \{ x \in \mathbb{S}^{d-1} : \langle x, e_1 \rangle \ge 1/\sqrt{2} \}$ and note that for any $x_1, x_2 \in \operatorname{Cap}_d(\pi/4), \langle x_1, x_2 \rangle \ge 0$. We claim that to prove (14), it is enough to prove that for some absolute $C_1, C_2, C_3 > 0$,

$$\mathbb{P}\left(M_0 := \max_{\boldsymbol{u} \in \mathbb{S}^{n-1}, \boldsymbol{x} \in \mathsf{Cap}_d(\pi/4)} Z_0(\boldsymbol{u}, \boldsymbol{x}) > C_1 p_{\max} \sqrt{d \log(p_{\max})}\right) < C_2 \exp(-C_3 d/C_*).$$
(20)

Indeed, by the Borell-TIS inequality, by increasing C_1 we can make sure that the same bound holds with C_3 is as large as we wish. Thus, noting that \mathbb{S}^{d-1} can be covered by $e^{C'd}$ rotations of $\mathsf{Cap}_d(\pi/4)$ for some absolute constant C' > 0, by rotational invariance and a union bound we obtain the required bound with \boldsymbol{x} maximized over \mathbb{S}^{d-1} instead of $\mathsf{Cap}_d(\pi/4)$.

In order to prove Eq. (20), we compute the canonical distance of this process to get (using $\xi_i(1) = 1$)

$$\begin{aligned} d_0(\boldsymbol{u}_1, \boldsymbol{x}_1; \boldsymbol{u}_2, \boldsymbol{x}_2)^2 &:= \mathbb{E}\left\{ \left(\langle \boldsymbol{u}_1, \boldsymbol{F}(\boldsymbol{x}_1) \rangle - \langle \boldsymbol{u}_2, \boldsymbol{F}(\boldsymbol{x}_2) \rangle \right)^2 \right\} \\ &= \sum_{i=1}^n \left\{ u_{1,i}^2 - 2u_{1,i}u_{2,i}\,\xi_i(\langle \boldsymbol{x}_1, \boldsymbol{x}_2 \rangle) + u_{2,i}^2 \right\} \\ &= \frac{1}{2} \sum_{i=1}^n (u_{1,i}^2 + u_{2,i}^2) \left(2 - 2\xi_i(\langle \boldsymbol{x}_1, \boldsymbol{x}_2 \rangle) \right) + \sum_{i=1}^n (u_{1,i} - u_{2,i})^2 \xi_i(\langle \boldsymbol{x}_1, \boldsymbol{x}_2 \rangle) \\ &\leq \max_{i \leq n} \left(2 - 2\xi_i(\langle \boldsymbol{x}_1, \boldsymbol{x}_2 \rangle) \right) + \|\boldsymbol{u}_1 - \boldsymbol{u}_2\|_2^2 \\ &\leq \xi_{\min}(\|\boldsymbol{x}_1\|^2) - 2\xi_{\min}(\langle \boldsymbol{x}_1, \boldsymbol{x}_2 \rangle) + \xi_{\min}(\|\boldsymbol{x}_2\|^2) + \|\boldsymbol{u}_1 - \boldsymbol{u}_2\|_2^2, \end{aligned}$$

where $\xi_{\min}(q) := \min_{i \leq n} \xi_i(q) = q^{p_{\max}}$ and we assume that $\boldsymbol{x}_1, \boldsymbol{x}_2 \in \mathsf{Cap}_d(\pi/4)$ so that $\langle \boldsymbol{x}_1, \boldsymbol{x}_2 \rangle \geq 0$. We recognize that the right-hand side is the canonical distance of the process $\overline{Z}(\boldsymbol{u}, \boldsymbol{x}) := \langle \boldsymbol{g}, \boldsymbol{u} \rangle + F_{\max}(\boldsymbol{x})$, where $\boldsymbol{g} \sim \mathsf{N}(0, \boldsymbol{I}_n)$ and F_{\max} is the centered Gaussian process with covariance $\langle \boldsymbol{x}_1, \boldsymbol{x}_2 \rangle^{p_{\max}}$. Using [MS23, Proposition A1], we get

$$\mathbb{E} \max_{\boldsymbol{u} \in \mathbb{S}^{n-1} \boldsymbol{x} \in \mathbb{S}^{d-1}} \overline{Z}(\boldsymbol{u}, \boldsymbol{x}) \le \mathbb{E} \|\boldsymbol{g}\| + \mathbb{E} \max_{\boldsymbol{x} \in \mathbb{S}^{d-1}} F_{\max}(\boldsymbol{x}) \le C\sqrt{n} + C\sqrt{d\log p_{\max}}.$$
 (21)

Hence, for M_0 as in (20), using the Sudakov-Fernique inequality we get, for $n \leq d$,

$$\mathbb{E}M_0 \le C\sqrt{d\log p_{\max}} \,. \tag{22}$$

On the other hand, $Z_0(\boldsymbol{u}, \boldsymbol{x})$ is a Lipschitz function of \boldsymbol{G} , hence so is M_0 , whence, by Gaussian concentration

$$\mathbb{P}(M_0 \ge \mathbb{E}M_0 + t) \le e^{-t^2/2}.$$
(23)

Equation (20), and thus (14), follow from the last two displays.

Proof of Eq. (15). We have

$$\max_{\boldsymbol{x}\in\mathsf{B}^{d}(1)} \|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\|_{\mathrm{op}} \stackrel{(a)}{\leq} \max_{\boldsymbol{x}\in\mathsf{B}^{d}(1)} \max_{\boldsymbol{v}\in\mathbb{S}^{d-1}:\langle\boldsymbol{v},\boldsymbol{x}\rangle=0} \|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{v}\| + \max_{\boldsymbol{x}\in\mathsf{B}^{d}(1)\setminus\mathbf{0}} \frac{\|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{x}\|}{\|\boldsymbol{x}\|}$$

$$\stackrel{(b)}{\leq} \max_{\boldsymbol{x},\boldsymbol{v}\in\mathbb{S}^{d-1}:\langle\boldsymbol{v},\boldsymbol{x}\rangle=0} \|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{v}\| + \max_{\boldsymbol{x}\in\mathbb{S}^{d-1}} \|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{x}\|$$

$$\stackrel{(c)}{\leq} \max_{(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v})\in\mathcal{A}_{n,d}} Z_{1}(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v}) + \max_{\boldsymbol{x}\in\mathbb{S}^{d-1}} \|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{x}\|, \qquad (24)$$

where (a) follows by triangle inequality, (b) by homogeneity, and in (c) we defined:

$$\mathcal{A}_{n,d} := \left\{ (\boldsymbol{u}, \boldsymbol{x}, \boldsymbol{v}) \in \mathbb{S}^{n-1} \times (\mathbb{S}^{d-1})^2 : \langle \boldsymbol{v}, \boldsymbol{x} \rangle = 0 \right\},$$
(25)

$$Z_1(\boldsymbol{u}, \boldsymbol{x}, \boldsymbol{v}) := \langle \boldsymbol{u}, \boldsymbol{D} \boldsymbol{F}(\boldsymbol{x}) \boldsymbol{v} \rangle.$$
(26)

The second term in the upper bound (24) is easily treated since

$$(\boldsymbol{DF}(\boldsymbol{x})\boldsymbol{x})_i = p_i \langle \boldsymbol{G}^{(i)}, \boldsymbol{x}^{\otimes p_i} \rangle = p_i F_i(\boldsymbol{x}),$$

and therefore

$$\mathbb{E} \| \boldsymbol{D} \boldsymbol{F}(\boldsymbol{x}) \boldsymbol{x} \| \le p_{\max} \mathbb{E} \| \boldsymbol{F}(\boldsymbol{x}) \| \le C p_{\max} \sqrt{d \log(p_{\max})} \,.$$
(27)

It is therefore sufficient to bound the first term in Eq. (24).

We next compute the canonical distance associated to the process Z_1 . For $(u_s, x_s, v_s) \in \mathcal{A}_{n,d}$, $s \in \{1, 2\}$, we get

$$d_1(\boldsymbol{u}_1, \boldsymbol{v}_1, \boldsymbol{x}_1; \boldsymbol{u}_2, \boldsymbol{v}_2, \boldsymbol{x}_2)^2 = \Delta_1(\boldsymbol{u}_1, \boldsymbol{v}_1, \boldsymbol{x}_1; \boldsymbol{u}_2, \boldsymbol{v}_2, \boldsymbol{x}_2) + \Delta_2(\boldsymbol{u}_1, \boldsymbol{v}_1, \boldsymbol{x}_1; \boldsymbol{u}_2, \boldsymbol{v}_2, \boldsymbol{x}_2),$$
(28)

$$\Delta_{1}(\boldsymbol{u}_{1},\boldsymbol{v}_{1},\boldsymbol{x}_{1};\boldsymbol{u}_{2},\boldsymbol{v}_{2},\boldsymbol{x}_{2}) := -2\langle \boldsymbol{x}_{1},\boldsymbol{v}_{2}\rangle\langle \boldsymbol{x}_{2},\boldsymbol{v}_{1}\rangle\sum_{i=1}^{n}u_{1,i}u_{2,i}\xi_{i}''(\langle \boldsymbol{x}_{1},\boldsymbol{x}_{2}\rangle),$$

$$\Delta_{2}(\boldsymbol{u}_{1},\boldsymbol{v}_{1},\boldsymbol{x}_{1};\boldsymbol{u}_{2},\boldsymbol{v}_{2},\boldsymbol{x}_{2}) := \sum_{i=1}^{n}u_{1,i}^{2}\xi_{i}'(1) + \sum_{i=1}^{n}u_{2,i}^{2}\xi_{i}'(1) - 2\langle \boldsymbol{v}_{1},\boldsymbol{v}_{2}\rangle\sum_{i=1}^{n}u_{1,i}u_{2,i}\xi_{i}'(\langle \boldsymbol{x}_{1},\boldsymbol{x}_{1}\rangle). \quad (29)$$

Denoting by \boldsymbol{P}_s^{\perp} the projector orthogonal to $\boldsymbol{x}_s,$ we have

$$\Delta_{1}(\boldsymbol{u}_{1}, \boldsymbol{v}_{1}, \boldsymbol{x}_{1}; \boldsymbol{u}_{2}, \boldsymbol{v}_{2}, \boldsymbol{x}_{2}) \leq 2 \max_{i \leq n} \xi_{i}^{\prime\prime}(1) \|\boldsymbol{P}_{1}^{\perp}\boldsymbol{x}_{2}\| \|\boldsymbol{P}_{2}^{\perp}\boldsymbol{x}_{1}\| \\ = 2p_{\max}(p_{\max} - 1)(1 - \langle \boldsymbol{x}_{1}, \boldsymbol{x}_{2} \rangle^{2}) \\ \leq 2p_{\max}^{2} \|\boldsymbol{x}_{1} - \boldsymbol{x}_{2}\|^{2}.$$
(30)

On the other hand,

$$\Delta_2(\boldsymbol{u}_1, \boldsymbol{v}_1, \boldsymbol{x}_1; \boldsymbol{u}_2, \boldsymbol{v}_2, \boldsymbol{x}_2) \leq \sum_{i=1}^n u_{1,i}^2 \xi_i'(1) + \sum_{i=1}^n u_{2,i}^2 \xi_i'(1) - 2\sum_{i=1}^n u_{1,i} u_{2,i} \xi_i'(\langle \boldsymbol{x}_1, \boldsymbol{x}_1 \rangle) + p_{\max} \|\boldsymbol{v}_1 - \boldsymbol{v}_2\|^2$$

Putting the above bounds together and using Sudakov-Fernique inequality, we see that

$$\mathbb{E} \max_{(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v})\in\mathcal{A}_{n,d}} Z_1(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v}) \leq \mathbb{E} \max_{(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v})\in\mathcal{A}_{n,d}} \left\{ \sqrt{2}p_{\max} \langle \boldsymbol{g}, \boldsymbol{x} \rangle + \sqrt{p_{\max}} \langle \boldsymbol{h}, \boldsymbol{v} \rangle + \langle \boldsymbol{u}, \widehat{\boldsymbol{F}}(\boldsymbol{x}) \rangle \right\},$$
(31)

where $\boldsymbol{g}, \boldsymbol{h}$ are mutually independent standard normal vectors, independent of $\hat{\boldsymbol{F}}$, a centered Gaussian process (taking values in \mathbb{R}^n) with covariance $\mathbb{E}\{\hat{F}_i(\boldsymbol{x}_1)\hat{F}_j(\boldsymbol{x}_2)\} = \delta_{ij}\xi'_i(\langle \boldsymbol{x}_1, \boldsymbol{x}_2 \rangle)$. By the same argument used for \boldsymbol{F} , we have $\mathbb{E}\max_{\boldsymbol{x}\in\mathbb{S}^{d-1}} \|\hat{\boldsymbol{F}}(\boldsymbol{x})\| \leq Cp_{\max}\sqrt{d\log p_{\max}}$. We then have

$$\mathbb{E} \max_{(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v})\in\mathcal{A}_{n,d}} Z_1(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v}) \leq C p_{\max}\sqrt{d} + \mathbb{E} \max_{\boldsymbol{x}\in\mathbb{S}^{d-1}} \|\widehat{\boldsymbol{F}}(\boldsymbol{x})\| \leq C p_{\max}\sqrt{d\log p_{\max}}.$$

Using this bound together with (27), we thus obtain

$$\mathbb{E} \| \boldsymbol{D} \boldsymbol{F}(\boldsymbol{x}) \|_{\text{op}} \le C p_{\max} \sqrt{d \log p_{\max}} \,. \tag{32}$$

The tail probability is controlled by Gaussian concentration as in the case of $\max_{\boldsymbol{x} \in \mathbb{S}^{d-1}} \|\boldsymbol{F}(\boldsymbol{x})\|$ treated above.

Proof of Eq. (16). To generalize the argument at the previous point, we define the process

$$Z_k(\boldsymbol{u}, \boldsymbol{x}, \boldsymbol{v}) := \sum_{i=1}^n u_i \langle \nabla^k F_i(\boldsymbol{x}), \boldsymbol{v}^{\otimes k} \rangle.$$
(33)

The quantity we want to upper bound is

$$\max_{\boldsymbol{x}\in\mathsf{B}^{d}(1)}\left\|\nabla^{k}\boldsymbol{F}(\boldsymbol{x})\right\|_{\mathrm{op}} = \max_{\boldsymbol{x}\in\mathsf{B}^{d}(1)}\max_{\boldsymbol{u}\in\mathbb{S}^{n-1}}\max_{\boldsymbol{v}\in\mathbb{S}^{d-1}}Z_{k}(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v}).$$
(34)

Note that we can always decompose $\boldsymbol{v} = a\boldsymbol{x} + \boldsymbol{v}_{\perp}$ where $\langle \boldsymbol{v}_{\perp}, \boldsymbol{x} \rangle = 0$, and therefore¹

$$Z_k(\boldsymbol{u}, \boldsymbol{x}, \boldsymbol{v}) := \sum_{j=0}^k \binom{k}{j} a^{k-j} \sum_{i=1}^n u_i \langle \nabla^k F_i(\boldsymbol{x}) \{ \boldsymbol{x}^{k-j} \}, \boldsymbol{v}_{\perp}^{\otimes j} \rangle.$$
(35)

By homogeneity, $\nabla^k F_i(\boldsymbol{x}) \{ \boldsymbol{x}^{k-j} \} = (p_i - j) \cdots (p_i - k + 1) \nabla^j F_i(\boldsymbol{x})$, and therefore all terms except the one with j = k can be controlled by bounds on lower order derivatives k. We therefore need to bound $\mathbb{E} \max_{(\boldsymbol{u}, \boldsymbol{x}, \boldsymbol{v}) \in \mathcal{A}_{n,d}} Z_k(\boldsymbol{u}, \boldsymbol{x}, \boldsymbol{v})$, i.e. to consider the case $\langle \boldsymbol{v}, \boldsymbol{x} \rangle = 0$.

Define the covaraince function

$$C_{k}(\boldsymbol{u}_{1},\boldsymbol{v}_{1},\boldsymbol{x}_{1};\boldsymbol{u}_{2},\boldsymbol{v}_{2},\boldsymbol{x}_{2}) := \mathbb{E}\left\{Z_{k}(\boldsymbol{u}_{1},\boldsymbol{x}_{1},\boldsymbol{v}_{1})Z_{k}(\boldsymbol{u}_{2},\boldsymbol{x}_{2},\boldsymbol{v}_{2})\right\} = \sum_{i=1}^{n} u_{1,i}u_{2,i}\partial_{\boldsymbol{v}_{1}}^{k}\partial_{\boldsymbol{v}_{2}}^{k}\xi_{i}(\langle\boldsymbol{x}_{1},\boldsymbol{x}_{2}\rangle)$$
$$= \sum_{j_{1},j_{2}}\hat{a}_{j_{1},j_{2},k}\sum_{i=1}^{n} u_{1,i}u_{2,i}\xi_{i}^{(j_{1}+j_{2})}(\langle\boldsymbol{x}_{1},\boldsymbol{x}_{2}\rangle)\langle\boldsymbol{v}_{1},\boldsymbol{x}_{2}\rangle^{j_{1}+j_{2}-k}\langle\boldsymbol{v}_{2},\boldsymbol{x}_{1}\rangle^{j_{1}+j_{2}-k}\langle\boldsymbol{v}_{1},\boldsymbol{v}_{2}\rangle^{2k-j_{1}-j_{2}}$$

¹Given symmetric tensors $\boldsymbol{A} \in (\mathbb{R}^n)^{\otimes k}$, $\boldsymbol{B} \in (\mathbb{R}^n)^{\otimes (k-j)}$, we denote by $\boldsymbol{T} = \boldsymbol{A}\{\boldsymbol{B}\}$ the tensor with components $T_{i_1...i_j} = \sum_{i_{j+1}...i_k} A_{i_1...i_k} B_{i_{j+1}...i_k}$.

$$=\sum_{j}a_{j,k}\sum_{i=1}^{n}u_{1,i}u_{2,i}\xi_{i}^{(j)}(\langle \boldsymbol{x}_{1},\boldsymbol{x}_{2}\rangle)\langle \boldsymbol{v}_{1},\boldsymbol{x}_{2}\rangle^{j-k}\langle \boldsymbol{v}_{2},\boldsymbol{x}_{1}\rangle^{j-k}\langle \boldsymbol{v}_{1},\boldsymbol{v}_{2}\rangle^{2k-j},$$

where $\partial_{\boldsymbol{v}_i}$ denotes the directional derivative corresponding to \boldsymbol{v}_i w.r.t. the coordinates of \boldsymbol{x}_i , $\hat{a}_{j_1,j_2,k}$ and $a_{j,k}$ are combinatorial factors and the summation is over j or j_1, j_2 such that all the exponents above are non-negative. Define $\hat{\boldsymbol{C}}_k(\boldsymbol{u}_1, \boldsymbol{v}_1, \boldsymbol{x}_1; \boldsymbol{u}_2, \boldsymbol{v}_2, \boldsymbol{x}_2)$ by the same expression as above with summation only over $j \neq k$ and note that, for $(\boldsymbol{u}, \boldsymbol{x}, \boldsymbol{v}) \in \mathcal{A}_{n,d}$, we have that

$$egin{aligned} \hat{m{C}}_k(m{u},m{v},m{x};m{u},m{v},m{x}) &= 0, \ m{C}_k(m{u},m{v},m{x};m{u},m{v},m{x}) &= a_{k,k}\sum_{i=1}^n u_i^2 \xi_i^{(k)}(1). \end{aligned}$$

Thus, for $(u_i, x_i, v_i) \in A_{n,d}$, the squared canonical distance of $Z_k(u, x, v)$ is given by

$$d_{k}(\boldsymbol{u}_{1}, \boldsymbol{v}_{1}, \boldsymbol{x}_{1}; \boldsymbol{u}_{2}, \boldsymbol{v}_{2}, \boldsymbol{x}_{2})^{2} = \boldsymbol{C}_{k}(\boldsymbol{u}_{1}, \boldsymbol{v}_{1}, \boldsymbol{x}_{1}; \boldsymbol{u}_{1}, \boldsymbol{v}_{1}, \boldsymbol{x}_{1}) - 2\boldsymbol{C}_{k}(\boldsymbol{u}_{1}, \boldsymbol{v}_{1}, \boldsymbol{x}_{1}; \boldsymbol{u}_{2}, \boldsymbol{v}_{2}, \boldsymbol{x}_{2}) + \boldsymbol{C}_{k}(\boldsymbol{u}_{2}, \boldsymbol{v}_{2}, \boldsymbol{x}_{2}; \boldsymbol{u}_{2}, \boldsymbol{v}_{2}, \boldsymbol{x}_{2})$$

$$\hat{\boldsymbol{C}}_{k}(\boldsymbol{u}_{1}, \boldsymbol{v}_{1}, \boldsymbol{x}_{1}; \boldsymbol{u}_{2}, \boldsymbol{v}_{2}, \boldsymbol{x}_{2}) + \hat{\boldsymbol{C}}_{k}(\boldsymbol{u}_{2}, \boldsymbol{v}_{2}, \boldsymbol{x}_{2}; \boldsymbol{u}_{2}, \boldsymbol{v}_{2}, \boldsymbol{x}_{2})$$

$$(26)$$

$$= C_k(u_1, v_1, x_1; u_1, v_1, x_1) - 2C_k(u_1, v_1, x_1; u_2, v_2, x_2) + C_k(u_2, v_2, x_2; u_2, v_2, x_2)$$
(36)

$$+ a_{k,k} \Big(\sum_{i=1}^{n} u_{1,i}^2 \xi_i^{(k)}(1) + \sum_{i=1}^{n} u_{2,i}^2 \xi_i^{(k)}(1) - 2 \sum_{i=1}^{n} u_{1,i} u_{2,i} \xi_i^{(k)}(\langle \boldsymbol{x}_1, \boldsymbol{x}_2 \rangle) \langle \boldsymbol{v}_1, \boldsymbol{v}_2 \rangle^k \Big) \,.$$
(37)

The expression in (36) is bounded by

$$A_k \max_{i \leq n, j \leq k} \xi_i^{(2j)}(1) |\langle \boldsymbol{v}_1, \boldsymbol{x}_2 \rangle \langle \boldsymbol{v}_2, \boldsymbol{x}_1 \rangle| \leq A_k p_{\max}^{2k} \|\boldsymbol{x}_1 - \boldsymbol{x}_2\|^2,$$

where A_k is a combinatorial factor and we used the same bound as in (30). On the other hand, assuming that $\langle \boldsymbol{v}_1, \boldsymbol{v}_2 \rangle \geq 0$, the expression in (37) is bounded by

$$a_{k,k} \Big(\sum_{i=1}^{n} u_{1,i}^{2} \xi_{i}^{(k)}(1) + \sum_{i=1}^{n} u_{2,i}^{2} \xi_{i}^{(k)}(1) - 2 \sum_{i=1}^{n} u_{1,i} u_{2,i} \xi_{i}^{(k)}(\langle \boldsymbol{x}_{1}, \boldsymbol{x}_{2} \rangle) \Big) + k p_{\max}^{k} \| \boldsymbol{v}_{1} - \boldsymbol{v}_{2} \|^{2}.$$

Hence, by the Sudakov-Fernique inequality

$$\mathbb{E} \max_{(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v})\in\mathcal{A}_{n,d}:\boldsymbol{v}\in\mathsf{Cap}_d(\pi/4)} Z_k(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v}) \leq \mathbb{E} \max_{(\boldsymbol{u},\boldsymbol{x},\boldsymbol{v})\in\mathcal{A}_{n,d}} \left\{ \sqrt{A_k} p_{\max}^k \langle \boldsymbol{g}, \boldsymbol{x} \rangle + \sqrt{k p_{\max}^k} \langle \boldsymbol{h}, \boldsymbol{v} \rangle + \langle \boldsymbol{u}, \widehat{\boldsymbol{F}}(\boldsymbol{x}) \rangle \right\},$$

where $\boldsymbol{g}, \boldsymbol{h}$ are mutually independent standard normal vectors, independent of $\hat{\boldsymbol{F}}$, a centered Gaussian process (taking values in \mathbb{R}^n) with covariance $\mathbb{E}\{\hat{F}_i(\boldsymbol{x}_1)\hat{F}_j(\boldsymbol{x}_2)\} = \delta_{ij}\xi_i^{(k)}(\langle \boldsymbol{x}_1, \boldsymbol{x}_2 \rangle)$. The proof of Eq. (16) thus follows by a similar argument to that following (31) and the argument used in the proof of Eq. (14) to move from $\mathsf{Cap}_d(\pi/4)$ to the whole sphere.

Proof of Eq. (17). We use

$$\left\|
abla^k H(oldsymbol{x})
ight\|_{ ext{op}} \leq 2^k \max_{0 \leq i \leq k} \left\|
abla^i oldsymbol{F}(oldsymbol{x})
ight\|_{ ext{op}} \left\|
abla^{k-i} oldsymbol{F}(oldsymbol{x})
ight\|_{ ext{op}},$$

and therefore this claim follows from the previous ones.

For any $\boldsymbol{x} \in \mathbb{R}^d \setminus \{0\}$, we let $\boldsymbol{U}_{\boldsymbol{x}} \in \mathbb{R}^{d \times (d-1)}$ be an arbitrary matrix whose columns form a basis of $\mathsf{T}_{\boldsymbol{x}}$. For $\boldsymbol{x}_1, \boldsymbol{x}_2 \in \mathbb{R}^d \setminus \{0\}$, define $\boldsymbol{U}_{\boldsymbol{x}_1, \boldsymbol{x}_2} := \boldsymbol{R}_{\boldsymbol{x}_1, \boldsymbol{x}_2} \boldsymbol{U}_{\boldsymbol{x}_1}$, where $\boldsymbol{R}_{\boldsymbol{x}_1, \boldsymbol{x}_2}$ is the rotation that keeps unchanged the space orthogonal to $\boldsymbol{x}_1, \boldsymbol{x}_2$, and maps $\boldsymbol{x}_1/\|\boldsymbol{x}_1\|_2$ to $\boldsymbol{x}_2/\|\boldsymbol{x}_2\|_2$. We will need the following geometric fact.

Lemma 2.2. For any non-zero $\boldsymbol{x}_1, \boldsymbol{x}_2 \in \mathsf{B}^d(1)$, we have

$$\| \boldsymbol{U}_{\boldsymbol{x}_1, \boldsymbol{x}_2} - \boldsymbol{U}_{\boldsymbol{x}_1} \|_{\text{op}} \le \frac{\| \boldsymbol{x}_1 - \boldsymbol{x}_2 \|_2}{\| \boldsymbol{x}_1 \|_2 \wedge \| \boldsymbol{x}_2 \|_2}.$$
 (38)

Proof. Note that $\|(\boldsymbol{R}_{\boldsymbol{x}_1,\boldsymbol{x}_2} - \boldsymbol{I})\boldsymbol{v}\|_2$ is constant for any unit vector $\boldsymbol{v} \in \text{span}(\boldsymbol{x}_1,\boldsymbol{x}_2)$, and vanishes if $\boldsymbol{v} \perp \text{span}(\boldsymbol{x}_1,\boldsymbol{x}_2)$. By using $\boldsymbol{v} = \boldsymbol{v}_1 := \boldsymbol{x}_1/\|\boldsymbol{x}_1\|_2$, we then have

$$egin{aligned} & ig\| m{U}_{m{x}_1,m{x}_2} - m{U}_{m{x}_1} ig\|_{ ext{op}} & \leq \|m{R}_{m{x}_1,m{x}_2} - m{I} ig\|_{ ext{op}} \ & = \|(m{R}_{m{x}_1,m{x}_2} - m{I})m{v}_1 \|_2 \ & = \Big\| rac{m{x}_1}{\|m{x}_1\|} - rac{m{x}_2}{\|m{x}_2\|} \Big\|_2 \ & \leq rac{\|m{x}_1 - m{x}_2\|_2}{\|m{x}_1\|_2 \wedge \|m{x}_2\|_2} \,. \end{aligned}$$

Definition 2.3. For $\Omega \subseteq \mathbb{R}^d$, we define the following Lipschitz constants:

$$\operatorname{Lip}(\boldsymbol{F};\Omega) := \sup_{\boldsymbol{x}_1 \neq \boldsymbol{x}_2 \in \Omega} \frac{\|\boldsymbol{F}(\boldsymbol{x}_1) - \boldsymbol{F}(\boldsymbol{x}_2)\|}{\|\boldsymbol{x}_1 - \boldsymbol{x}_2\|_2},$$
(39)

$$\operatorname{Lip}(\boldsymbol{DF};\Omega) := \sup_{\boldsymbol{x}_1 \neq \boldsymbol{x}_2 \in \Omega} \frac{\|\boldsymbol{DF}(\boldsymbol{x}_1) - \boldsymbol{DF}(\boldsymbol{x}_2)\|_{\operatorname{op}}}{\|\boldsymbol{x}_1 - \boldsymbol{x}_2\|_2},$$
(40)

$$\operatorname{Lip}(\nabla^{2} \boldsymbol{F}; \Omega) := \sup_{\boldsymbol{x}_{1} \neq \boldsymbol{x}_{2} \in \Omega} \max_{\ell \leq n} \frac{\|\nabla^{2} F_{\ell}(\boldsymbol{x}_{1}) - \nabla^{2} F_{\ell}(\boldsymbol{x}_{2})\|_{\operatorname{op}}}{\|\boldsymbol{x}_{1} - \boldsymbol{x}_{2}\|_{2}}.$$
(41)

We also define the following Lipschitz constants for projections onto the tangent space:

$$\operatorname{Lip}_{\perp}(\boldsymbol{DF};\Omega) := \sup_{\boldsymbol{x}_{1} \neq \boldsymbol{x}_{2} \in \Omega} \frac{\|\boldsymbol{DF}(\boldsymbol{x}_{1})\boldsymbol{U}_{\boldsymbol{x}_{1}} - \boldsymbol{DF}(\boldsymbol{x}_{2})\boldsymbol{U}_{\boldsymbol{x}_{1},\boldsymbol{x}_{2}}\|_{\operatorname{op}}}{\|\boldsymbol{x}_{1} - \boldsymbol{x}_{2}\|_{2}},$$
(42)

$$\operatorname{Lip}_{\perp}(\nabla^{2} \boldsymbol{F}; \Omega) := \sup_{\boldsymbol{x}_{1} \neq \boldsymbol{x}_{2} \in \Omega} \max_{\ell \leq n} \frac{\|\boldsymbol{U}_{\boldsymbol{x}_{1}}^{\mathsf{T}} \nabla^{2} F_{\ell}(\boldsymbol{x}_{1}) \boldsymbol{U}_{\boldsymbol{x}_{1}} - \boldsymbol{U}_{\boldsymbol{x}_{1}, \boldsymbol{x}_{2}}^{\mathsf{T}} \nabla^{2} F_{\ell}(\boldsymbol{x}_{2}) \boldsymbol{U}_{\boldsymbol{x}_{1}, \boldsymbol{x}_{2}} \|_{\operatorname{op}}}{\|\boldsymbol{x}_{1} - \boldsymbol{x}_{2}\|_{2}}.$$
 (43)

The next lemma is an immediate consequence of Lemma 2.1 and Lemma 2.2.

Lemma 2.4. Assume that $n \leq d$. Then there exist absolute constants C_0 , C_* such that the following hold with probability at least $1 - C_* \exp(-d/C_*)$:

$$\operatorname{Lip}(\boldsymbol{F}; \mathsf{B}^{d}(1)) \le C_0 p_{\max} \sqrt{d \log(p_{\max})}, \qquad (44)$$

$$\operatorname{Lip}(\boldsymbol{DF}; \mathsf{B}^{d}(1)) \leq C_{0} p_{\max}^{2} \sqrt{d \log(p_{\max})}, \quad \operatorname{Lip}_{\perp}(\boldsymbol{DF}; \mathsf{B}^{d}(\rho, 1)) \leq \frac{C_{0}}{\rho} p_{\max}^{2} \sqrt{d \log(p_{\max})}, \quad (45)$$

$$\operatorname{Lip}(\nabla^{2}\boldsymbol{F};\mathsf{B}^{d}(1)) \leq C_{0}p_{\max}^{3}\sqrt{d\log(p_{\max})}, \quad \operatorname{Lip}_{\perp}(\nabla^{2}\boldsymbol{F};\mathsf{B}^{d}(\rho,1)) \leq \frac{C_{0}}{\rho}p_{\max}^{3}\sqrt{d\log(p_{\max})}.$$
(46)

Proof. The bounds on $\operatorname{Lip}(\boldsymbol{F}; \mathsf{B}^d(1))$, $\operatorname{Lip}(\boldsymbol{DF}; \mathsf{B}^d(1))$, $\operatorname{Lip}(\nabla^2 \boldsymbol{F}; \mathsf{B}^d(1))$ follow immediately from Lemma 2.1. The bounds on $\operatorname{Lip}_{\perp}(\boldsymbol{DF}; \mathsf{B}^d(1))$, $\operatorname{Lip}_{\perp}(\nabla^2 \boldsymbol{F}; \mathsf{B}^d(1))$ are proved similarly and we limit ourselves to the last one. Writing $\boldsymbol{U}_1 := \boldsymbol{U}_{\boldsymbol{x}_1}, \boldsymbol{U}_2 := \boldsymbol{U}_{\boldsymbol{x}_1, \boldsymbol{x}_2}$, and assuming without loss of generality $\|\boldsymbol{x}_1\|_2 \leq \|\boldsymbol{x}_2\|_2$,

$$\max_{\ell \leq n} \| \boldsymbol{U}_1^\mathsf{T} \nabla^2 F_\ell(\boldsymbol{x}_1) \boldsymbol{U}_1 - \boldsymbol{U}_2^\mathsf{T} \nabla^2 F_\ell(\boldsymbol{x}_2) \boldsymbol{U}_2 \|_{\mathrm{op}}$$

$$\leq \max_{\ell \leq n} \left\| \nabla^2 F_{\ell}(\boldsymbol{x}_1) - \nabla^2 F_{\ell}(\boldsymbol{x}_2) \right\|_{\text{op}} + 3 \max_{\ell \leq n} \left\| \nabla^2 F_{\ell}(\boldsymbol{x}_1) \right\|_{\text{op}} \left\| \boldsymbol{U}_1 - \boldsymbol{U}_2 \right\|_{\text{op}}$$

$$\leq \max_{\ell \leq n} \left\| \nabla^2 F_{\ell}(\boldsymbol{x}_1) - \nabla^2 F_{\ell}(\boldsymbol{x}_2) \right\|_{\text{op}} + 3C \max_{\ell \leq n} \frac{\left\| \nabla^2 F_{\ell}(\boldsymbol{x}_1) \right\|_{\text{op}}}{\|\boldsymbol{x}_1 - \boldsymbol{x}_2\|_2}$$

$$\leq C_0 p_{\max}^3 \sqrt{d \log(p_{\max})} \|\boldsymbol{x}_1 - \boldsymbol{x}_2\|_2 + \frac{C_0}{\rho} p_{\max}^2 \sqrt{d \log(p_{\max})} \|\boldsymbol{x}_1 - \boldsymbol{x}_2\|_2 .$$

3 Bounds on the Hessian

In this section we derive the main lower bound on the eigenvalues of the Hessian, which will be used in the analysis of the Hessian descent algorithm.

Denote by $\mathcal{H}(\boldsymbol{x}) := \nabla^2 H(\boldsymbol{x})|_{\mathsf{T}_{\boldsymbol{x}}}$ the restriction of the Hessian on the tangent space. In a matrix representation, this is given by

$$\mathcal{H}(\boldsymbol{x}) = \sum_{\ell=1}^{n} F_{\ell}(\boldsymbol{x}) \boldsymbol{U}_{\boldsymbol{x}}^{\mathsf{T}} \nabla^{2} F_{\ell}(\boldsymbol{x}) \boldsymbol{U}_{\boldsymbol{x}} + \boldsymbol{U}_{\boldsymbol{x}}^{\mathsf{T}} \boldsymbol{D} \boldsymbol{F}(\boldsymbol{x})^{\mathsf{T}} \boldsymbol{D} \boldsymbol{F}(\boldsymbol{x}) \boldsymbol{U}_{\boldsymbol{x}}$$
(47)

$$=: \mathcal{H}_1(\boldsymbol{x}) + \mathcal{H}_2(\boldsymbol{x}) \,. \tag{48}$$

We omit the calculation for the next lemma, which is similar e.g. to [ABAČ13, MS23, Sub17, Sub23]. Recall that $\xi_i(t) := t^{p_i}$.

Lemma 3.1. Define for $k \in \mathbb{N}$, $q \in [0,1]$ define $S_k(q) := \operatorname{diag}\left(\sqrt{\xi_i^{(k)}(q)} : i \leq n\right)$, where $\xi_i^{(k)}$ is the k-th derivative of ξ_i . For a fixed $\mathbf{x} \in \mathbb{R}^d$ with $\|\mathbf{x}\|_2^2 = q$, we have $F(\mathbf{x}) = S_0(q) \mathbf{g}$, $DF(\mathbf{x})U_{\mathbf{x}} = S_1(q) \mathbf{Z}$, $U_{\mathbf{x}}^{\mathsf{T}} \nabla^2 F_\ell(\mathbf{x}) U_{\mathbf{x}} = \sqrt{\xi_\ell''(q)} \mathbf{W}_\ell$. Where $\mathbf{g}, (\mathbf{W}_\ell)_{\ell \leq n}, \mathbf{Z}$ are mutually independent with

$$\boldsymbol{g} \sim \mathsf{N}(0, \boldsymbol{I}_n), \quad \boldsymbol{W}_{\ell} \sim \mathsf{GOE}(d-1), \quad \boldsymbol{Z} \sim \mathsf{GOE}(n, d-1).$$
 (49)

As a consequence

$$\mathcal{H}(\boldsymbol{x}) = \|\boldsymbol{S}_0(q)\boldsymbol{S}_2(q)\boldsymbol{g}\|_2 \boldsymbol{W} + \boldsymbol{Z}^{\mathsf{T}}\boldsymbol{S}_1(q)^2 \boldsymbol{Z}, \qquad (50)$$

$$H(\boldsymbol{x}) = \frac{1}{2} \|\boldsymbol{S}_0(q)\boldsymbol{g}\|_2^2,$$
(51)

where

$$(\boldsymbol{g}, \boldsymbol{W}, \boldsymbol{Z}) \sim \mathsf{N}(0, \boldsymbol{I}_n) \otimes \mathsf{GOE}(d-1) \otimes \mathsf{GOE}(n, d-1).$$
 (52)

We denote by $\lambda_i(\mathbf{M})$ the *i*-th smallest eigenvalue of a symmetric matrix \mathbf{M} .

Theorem 3. For $t \in (0,1)$, define $\xi'_{\min}(t) := \min_{i \leq n} \xi'_i(t)$ and $\xi''_{\min}(t) := \min_{i \leq n} \xi''_i(t)$. Then for any $s \in \mathbb{N}$, there exist constants C_* , $C_{\#}$, and $d_0 = d_0(s)$ such that the following holds. Define $A_{\#}, B_{\#}$ via

$$A_{\#} = C_{\#} \left(1 \vee \sqrt{\frac{\log p_{\max}}{\log d}} \vee \sqrt{\frac{-\log[\xi'_{\min}(\rho^2) \wedge \rho]}{\log d}} \right) , \tag{53}$$

$$B_{\#} = C_{\#} \left(\sqrt{\log p_{\max}} \vee \sqrt{-\log(\xi_{\min}''(\rho^2)\rho^2 \wedge 1)} \vee \sqrt{\log d} \right).$$
(54)

If $n \leq d - 4A_{\#} (d \log d)^{1/2}$, $n \leq d - B_{\#} d^{1/2}$, then the following holds for all $d \geq d_0(s)$:

$$\mathbb{P}\Big(\forall \boldsymbol{x} \in \mathsf{B}^{d}(\rho, 1) \ \lambda_{s}(\boldsymbol{\mathcal{H}}(\boldsymbol{x})) \leq -\frac{1}{10}\sqrt{(d-n)\xi_{\min}^{\prime\prime}(\rho^{2})H(\boldsymbol{x})}\Big) \geq 1 - C_{*}e^{-d/C_{*}}.$$
(55)

Proof. For $\boldsymbol{\alpha} \in \mathbb{R}^n$, define the symmetric form $\boldsymbol{M}(\boldsymbol{x}; \boldsymbol{\alpha})$ on $\mathsf{T}_{\boldsymbol{x}}$ by

$$\boldsymbol{M}(\boldsymbol{x};\boldsymbol{\alpha}) = \sum_{\ell=1}^{n} \alpha_{\ell} \nabla^{2} F_{\ell}(\boldsymbol{x}) \Big|_{\mathsf{T}_{\boldsymbol{x}}}.$$
(56)

Let $v_1(x), \ldots, v_{d-1}(x)$ be right singular vectors of $DF(x)|_{\mathsf{T}_x}$ corresponding to singular values $\sigma_1(DF(x)|_{\mathsf{T}_x}) \geq \sigma_2(DF(x)|_{\mathsf{T}_x}) \geq \cdots \geq \sigma_{d-1}(DF(x)|_{\mathsf{T}_x})$ (whereby we include zero singular values). For $m \in \{1, \ldots, n+1\}$, define

$$V_m(\boldsymbol{x}) := \operatorname{span} \left(\boldsymbol{v}_{n-m+2}(\boldsymbol{x}), \dots, \boldsymbol{v}_{d-1}(\boldsymbol{x}) \right).$$
(57)

Let $\operatorname{null}(DF(x)|_{\mathsf{T}_x}) \subseteq \mathsf{T}_x$ denote the null space of $DF(x)|_{\mathsf{T}_x}$ (in the basis U_x , this is equivalent to the null space of $DF(x)U_x$). Since $\dim(\operatorname{null}(DF(x)|_{\mathsf{T}_x})) \ge d-1-n$, we have

$$V_1(\boldsymbol{x}) \subseteq \operatorname{null}(\boldsymbol{DF}(\boldsymbol{x})|_{\mathsf{T}_{\boldsymbol{x}}}), \tag{58}$$

$$V_1(\boldsymbol{x}) \subseteq V_2(\boldsymbol{x}) \subseteq \cdots \subseteq V_{n+1}(\boldsymbol{x}).$$
(59)

(with $V_1(\boldsymbol{x}) = \operatorname{null}(\boldsymbol{DF}(\boldsymbol{x})|_{\mathsf{T}_{\boldsymbol{x}}})$ if $\boldsymbol{DF}(\boldsymbol{x})|_{\mathsf{T}_{\boldsymbol{x}}}$ has full row rank.) We will define $d_m := \dim(V_m(\boldsymbol{x})) = d - n + m - 2$ and, for each m,

$$L_{k,m}(\boldsymbol{x};\boldsymbol{\alpha}) := \lambda_k \left(\boldsymbol{M}(\boldsymbol{x};\boldsymbol{\alpha})|_{V_m(\boldsymbol{x})} \right).$$
(60)

where $\lambda_{\ell}(\mathbf{A})$ denotes the ℓ -th smallest eigenvalue of matrix \mathbf{A} .

Defining $\boldsymbol{lpha}_*(\boldsymbol{x}) := \boldsymbol{F}(\boldsymbol{x}) / \| \boldsymbol{F}(\boldsymbol{x}) \|_2$, we then have

$$\lambda_s(\mathcal{H}(\boldsymbol{x})) \le \lambda_s(\mathcal{H}_1(\boldsymbol{x})|_{\text{null}(\boldsymbol{DF}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}})})$$
(61)

$$= \|\boldsymbol{F}(\boldsymbol{x})\|_{2} \cdot \lambda_{s}(\boldsymbol{M}(\boldsymbol{x};\boldsymbol{\alpha}_{*}(\boldsymbol{x}))|_{\text{null}(\boldsymbol{DF}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}})})$$
(62)

$$\leq \|\boldsymbol{F}(\boldsymbol{x})\|_{2} \cdot \lambda_{s}(\boldsymbol{M}(\boldsymbol{x};\boldsymbol{\alpha}_{*}(\boldsymbol{x})|_{V_{1}(\boldsymbol{x})})$$
(63)

$$= \|\boldsymbol{F}(\boldsymbol{x})\|_2 \cdot L_{s,1}(\boldsymbol{x}; \boldsymbol{\alpha}_*(\boldsymbol{x})).$$
(64)

Further notice that, by the variational representation of eigenvalues, we have, for each k, m,

$$L_{k,m}(\boldsymbol{x};\boldsymbol{\alpha}) \le L_{k+1,m+1}(\boldsymbol{x};\boldsymbol{\alpha}).$$
(65)

To see this, writing for simplicity $M = M(s; \alpha)$ and $V_m = V_m(x)$, note that

$$\lambda_{k+1}(\boldsymbol{M}|_{V_{m+1}}) = \max_{\substack{U \subseteq V_{m+1} \\ \dim(U) = d_{m+1} - k}} \lambda_1(\boldsymbol{M}|_U)$$
$$\geq \max_{\substack{U \subseteq V_m \\ \dim(U) = d_m - (k-1)}} \lambda_1(\boldsymbol{M}|_U)$$
$$= \lambda_k(\boldsymbol{M}|_{V_m}),$$

where the inequality holds, since $V_m \subseteq V_{m+1}$ and $d_{m+1} - k = d_m - (k-1)$. We then have

$$\mathrm{P}_{\mathrm{good}} := \mathbb{P}\Big(\forall \boldsymbol{x} \in \mathsf{B}^d(\rho, 1) \; \lambda_s(\boldsymbol{\mathcal{H}}(\boldsymbol{x})) \leq -\frac{1}{10} \sqrt{(d-n)\xi_{\min}''(\rho^2)H(\boldsymbol{x})} \Big)$$

$$\geq \mathbb{P}\Big(\max_{\boldsymbol{x}\in\mathsf{B}^{d}(\rho,1)}\max_{\boldsymbol{\alpha}\in\mathbb{S}^{n-1}}L_{s,1}(\boldsymbol{x};\boldsymbol{\alpha})\leq -\frac{1}{10}\sqrt{\xi_{\min}''(\rho^{2})(d-n)}\Big).$$
(66)

Now note that, for fixed $\boldsymbol{\alpha}, \boldsymbol{x}$, with $\|\boldsymbol{\alpha}\|_2 = 1$, we have

$$\left(\boldsymbol{M}(\boldsymbol{x};\boldsymbol{\alpha}),\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})|_{\mathsf{T}_{\boldsymbol{x}}}\right) \stackrel{\mathrm{d}}{=} \left(\nu(\boldsymbol{\alpha};\|\boldsymbol{x}\|_{2}^{2})\cdot\boldsymbol{W},\,\boldsymbol{S}_{1}(\|\boldsymbol{x}\|_{2}^{2})\cdot\boldsymbol{G}\right),\tag{67}$$

$$\nu(\boldsymbol{\alpha};q) = \left(\sum_{i=1}^{n} \alpha_i^2 \xi_i''(q)\right)^{1/2},\tag{68}$$

where $\boldsymbol{W} \sim \mathsf{GOE}(d-1)$, $\boldsymbol{G} \sim \mathsf{GOE}(n, d-1)$ are independent random matrices (cf. Lemma 3.1). Hence $\boldsymbol{M}(\boldsymbol{x}; \boldsymbol{\alpha})|_{V_m(\boldsymbol{x})} \sim \nu(\boldsymbol{\alpha}; q)\mathsf{GOE}(d_m)$, $q = \|\boldsymbol{x}\|_2^2$. Since $\|\boldsymbol{\alpha}\|_2 = 1$, we have $\nu(\boldsymbol{\alpha}; q) \geq \sqrt{\xi''_{\min}(q)}$. Therefore, by Lemma B.1, there exists an absolute constant $C_2 > 0$ such that, for any $k \leq 3(d-n)/8$,

$$\mathbb{P}\Big(L_{k,m}(\boldsymbol{x};\boldsymbol{\alpha}) \ge -\frac{1}{5}\sqrt{\xi_{\min}''(\|\boldsymbol{x}\|_2^2)(d-n)}\Big) \le e^{-C_2(d-n)^2}.$$
(69)

Define the following events, depending on absolute constants $C_0, C_1 > 0$, which will be chosen below

$$\mathcal{E}_{1}(C_{0}) := \left\{ \left| L_{s,1}(\boldsymbol{x};\boldsymbol{\alpha}_{1}) - L_{s,1}(\boldsymbol{x};\boldsymbol{\alpha}_{2}) \right| \le C_{0} dp_{\max}^{3} \|\boldsymbol{\alpha}_{1} - \boldsymbol{\alpha}_{2}\|_{2} \ \forall \boldsymbol{x} \in \mathsf{B}^{d}(1) \,\forall \boldsymbol{\alpha}_{1}, \boldsymbol{\alpha}_{2} \in \mathbb{S}^{n-1} \right\},$$

$$\tag{70}$$

$$\mathcal{E}_{2,m}(C_1,\rho) := \left\{ L_{s,1}(\boldsymbol{x}_1; \boldsymbol{\alpha}) \le L_{m+s-1,m}(\boldsymbol{x}_0; \boldsymbol{\alpha}) + \frac{C_1 d^2 p_{\max}^5}{\xi_{\min}'(\rho^2)^{1/2} \rho} \| \boldsymbol{x}_0 - \boldsymbol{x}_1 \|_2 \ \forall \boldsymbol{x}_0, \boldsymbol{x}_1 \in \mathsf{B}^d(\rho, 1) \, \forall \boldsymbol{\alpha} \in \mathbb{S}^{n-1} \right\},\tag{71}$$

where we recall that $\xi'_{\min}(t) := \min_{i \le n} \xi'_i(t)$. Further, define

$$\mathcal{E}_2(C_1,\rho) := \mathcal{E}_{2,m=\lfloor (d-n)/4 \rfloor}(C_1,\rho).$$
(72)

Let $N^d(\eta)$ be an η -net in $\mathsf{B}^d(\rho, 1) \times \mathbb{S}^{n-1}$. Denoting by $(\boldsymbol{x}^{\eta}, \boldsymbol{\alpha}^{\eta})$ the projection of $(\boldsymbol{x}, \boldsymbol{\alpha}) \in \mathsf{B}^d(\rho, 1) \times \mathbb{S}^{n-1}$ onto $N^d(\eta)$, we have that, on $\mathcal{E}_1(C_0) \cap \mathcal{E}_2(C_1, \rho)$,

$$L_{s,1}(\boldsymbol{x};\boldsymbol{\alpha}) \le L_{m+s-1,m}(\boldsymbol{x}^{\eta};\boldsymbol{\alpha}^{\eta}) + \frac{(C_0 + C_1)d^2 p_{\max}^5 \eta}{1 \wedge [\rho \xi'_{\min}(\rho^2)^{1/2}]}.$$
(73)

Taking

$$\eta = \frac{1}{10(C_0 + C_1)d^2 p_{\max}^5} \xi_{\min}''(\rho^2)^{1/2} \{ [\rho \xi_{\min}'(\rho^2)^{1/2}] \wedge 1 \},$$
(74)

this implies $L_{s,1}(\boldsymbol{x};\boldsymbol{\alpha}) \leq L_{m+s-1,m}(\boldsymbol{x}^{\eta};\boldsymbol{\alpha}^{\eta}) + \xi_{\min}''(\rho^2)^{1/2}/10$. Then continuing from Eq. (66), with $m = \lfloor (d-n)/4 \rfloor$ we obtain, for C, C_2 absolute constants and $s \leq (d-n)/8$,

$$\begin{aligned} \mathbf{P}_{\text{good}} &\geq 1 - |N^{d}(\eta)| \max_{(\boldsymbol{x}, \boldsymbol{\alpha}) \in N^{d}(\eta)} \mathbb{P} \left(L_{m+s-1, m}(\boldsymbol{x}; \boldsymbol{\alpha}) \geq -\frac{1}{5} \sqrt{\xi_{\min}^{\prime\prime}(\rho^{2})(d-n)} \right) - \mathbb{P}(\mathcal{E}_{1}^{c}(C_{0})) - \mathbb{P}(\mathcal{E}_{2}^{c}(C_{1}, \rho)) \\ &\geq 1 - \left(\frac{C}{\eta}\right)^{2d} e^{-C_{2}(d-n)^{2}} - \mathbb{P}(\mathcal{E}_{1}^{c}(C_{0})) - \mathbb{P}(\mathcal{E}_{2}^{c}(C_{1}, \rho)) \\ &\geq 1 - e^{-\Lambda d} - C_{*} e^{-d/C_{*}} , \end{aligned}$$

where the last step follows from Lemma 3.2 and Lemma 3.3 (since $n \leq d - 4A_{\#}(d \log d)^{1/2}$ holds by assumption), and $\Lambda = C_2(d-n)^2/d - 2\log(C/\eta)$ is bounded by

$$\Lambda \ge C_2 \frac{(d-n)^2}{d} - C' \left(\log d \lor \log p_{\max} \lor \log \frac{1}{\xi_{\min}''(\rho^2)\rho^2} \right), \tag{75}$$

for some absolute C', where we used the fact that $p_{\max}\xi'_i(\rho^2) \ge \xi''_i(\rho^2)\rho^2$. The requirement that $n \le d - B_{\#} d^{1/2}$ with sufficiently large $C_{\#}$ guarantees that $\Lambda > c$ for some absolute c > 0.

Lemma 3.2. Under the assumptions of Theorem 3, there exists an absolute constant C_* such that, defining \mathcal{E}_1 as per Eq. (70), we have

$$\mathbb{P}(\mathcal{E}_1^c(C_*)) \le C_* e^{-d/C_*} \,. \tag{76}$$

Proof. By Weyl's inequality,

$$\begin{split} \left| L_{s,1}(\boldsymbol{x};\boldsymbol{\alpha}_1) - L_{s,1}(\boldsymbol{x};\boldsymbol{\alpha}_2) \right| &\leq \left\| \boldsymbol{M}(\boldsymbol{x};\boldsymbol{\alpha}_1) - \boldsymbol{M}(\boldsymbol{x};\boldsymbol{\alpha}_2) \right\|_{\text{op}} \\ &\leq \sqrt{n} \|\boldsymbol{\alpha}_1 - \boldsymbol{\alpha}_2\|_2 \cdot \max_{\ell \leq n} \left\| \nabla^2 F_{\ell}(\boldsymbol{x}) |_{\mathsf{T}_{\boldsymbol{x}}} \right\|_{\text{op}} \\ &\leq C \sqrt{n} \|\boldsymbol{\alpha}_1 - \boldsymbol{\alpha}_2\|_2 p_{\max}^2 \sqrt{d \log p_{\max}} \,, \end{split}$$

where the last step holds with probability at least $1 - C \exp(-d/C)$ by Lemma 2.1.

Lemma 3.3. Under the assumptions of Theorem 3, there exist absolute constants $C_{\#}, C_* > 0$ such that the following holds. Define \mathcal{E}_2 as per Eq. (71), and let $A_{\#}$ be defined by

$$A_{\#} = C_{\#} \left(1 \vee \sqrt{\frac{\log p_{\max}}{\log d}} \vee \sqrt{\frac{-\log(\xi_{\min}'(\rho^2) \wedge \rho)}{\log d}} \right) . \tag{77}$$

If $n \leq d - A_{\#} (d \log d)^{1/2}$, $m = \lfloor (d - n)/4 \rfloor$ then we have

$$\mathbb{P}(\mathcal{E}_{2,m}^{c}(C_{*},\rho)) \le C_{*} e^{-d/C_{*}}.$$
(78)

Before proving Lemma 3.3, it is useful to state and prove an auxiliary result.

Lemma 3.4. For $t \in (0,1)$, let $\xi'_{\min}(t) := \min_{i \leq n} \xi'_i(t)$. For $\mathbf{x} \in \mathsf{B}^d(1)$, denote by $\sigma_1(\mathbf{x}) \geq \sigma_2(\mathbf{x}) \geq \cdots \geq \sigma_{d-1}(\mathbf{x})$ the singular values of $\mathbf{DF}(\mathbf{x})|_{\mathsf{T}_{\mathbf{x}}}$ (including the vanishing ones).

Then there exist absolute constants C_* , $C_{\#}$, $\varepsilon_0 > 0$ such that the following holds, with $A_{\#}$ defined as per Eq. (77). For any $\varepsilon \in (0, \varepsilon_0]$, $\rho \in (0, 1)$, $m \ge (d - n)\varepsilon$ and $n \le d - A_{\#}\sqrt{\varepsilon^{-1}d(\log d)}$, we have

$$\mathbb{P}\Big(\forall \boldsymbol{x} \in \mathsf{B}^{d}(\rho, 1): \ \sigma_{n-m+1}(\boldsymbol{x}) \geq \frac{1}{C_{*}} \sqrt{\xi_{\min}'(\|\boldsymbol{x}\|_{2}^{2})} (\sqrt{d} - \sqrt{n}) \Big) \geq 1 - C_{*} e^{-d/C_{*}} .$$
(79)

Proof. If $\mathbf{a}_i \in \mathbb{R}^{(d-1)}$, $i \leq n+1$, define $Q_{ij} := \langle \mathbf{a}_i, \mathbf{a}_j \rangle$, and $\mathbf{Q}_{\ell} := (Q_{ij})_{i,j \leq \ell} \in \mathbb{R}^{\ell \times \ell}$, then, by the variational principle, for any $i \leq n$

$$\lambda_i(\boldsymbol{Q}_{n+1}) \le \lambda_i(\boldsymbol{Q}_n) \,. \tag{80}$$

Note that $\sigma_{n-m+1}(\boldsymbol{x})^2 = \lambda_{m-1}(\boldsymbol{Q}_n)$ for $\boldsymbol{Q}_n := \boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}})^{\mathsf{T}}$. In particular, removing one row of $\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})$ cannot increase these eigenvalues. Hence, without loss of generality we can

assume throughout the proof that $n \ge d/2$. Also note that since m is an integer, we may assume that $\varepsilon(d-n) \ge 1$.

First consider a fixed point $\boldsymbol{x} \in \mathsf{B}^{d}(1)$. Recall that, by Lemma 3.1, $\boldsymbol{DF}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}} = \boldsymbol{S}^{(1)}(\|\boldsymbol{x}\|_{2}^{2})\boldsymbol{Z}$, with $\boldsymbol{Z} \sim \mathsf{GOE}(n, d-1)$. Denoting the eigenvalues of $\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}$ by $\lambda_{1}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq \lambda_{2}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq \cdots \leq \lambda_{n}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}})$, we have that, for any $1 \leq m \leq n$,

$$\xi_{\min}'(\|\boldsymbol{x}\|^2)\lambda_{k(\varepsilon)}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq \xi_{\min}'(\|\boldsymbol{x}\|^2)\lambda_m(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq \sigma_{n-m+1}(\boldsymbol{x})^2,$$
(81)

where we define $k(\varepsilon) = \lfloor \varepsilon(d-n) \rfloor$.

By Lemma B.2, with N = d - 1, M = n, $\ell = k(\varepsilon)$, there exists constants $\varepsilon_0 > 0$, $\Delta_* > 0$, such that for all $\varepsilon \in (0, \varepsilon_0)$,

$$\mathbb{P}\left(\lambda_{k(\varepsilon)}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq 4\Delta_{*}^{2}(\sqrt{d} - \sqrt{n})^{2}\right) \leq e^{-\varepsilon(d-n)^{2}},\tag{82}$$

where we used that $\sqrt{d} - \sqrt{n} \leq \sqrt{d-1} - \sqrt{n-1}$. Therefore, for $n \leq d - Ad^{1/2}(\log d)^{1/2}$, we have

$$\mathbb{P}\Big(\lambda_{k(\varepsilon)}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq 4\Delta_*^2(\sqrt{d} - \sqrt{n})^2\Big) \leq \exp\left(-A^2\varepsilon(d\log d)\right).$$
(83)

Let $N^d(\eta)$ be an η -net in $\mathsf{B}^d(\rho, 1)$. For $\boldsymbol{x} \in \mathsf{B}^d(\rho, 1)$, let $\boldsymbol{x}^{\eta} = \operatorname{argmin}_{\boldsymbol{y} \in N^d(\eta)} \|\boldsymbol{x} - \boldsymbol{y}\|_2$ be its projection onto the net. Then, using $\sqrt{d} - \sqrt{n} \ge 1/(2\sqrt{d})$,

$$\begin{aligned} \mathbb{P}\Big(\exists \boldsymbol{x} \in \mathsf{B}^{d}(\rho, 1): \ \left|\sigma_{n-m+1}(\boldsymbol{x}) - \sigma_{n-m+1}(\boldsymbol{x}^{\eta})\right| &\geq \Delta_{*}\sqrt{\xi_{\min}'(\|\boldsymbol{x}\|_{2}^{2})}(\sqrt{d} - \sqrt{n})\Big) \\ &\leq \mathbb{P}\Big(\operatorname{Lip}_{\perp}(\boldsymbol{D}\boldsymbol{F}; \mathsf{B}^{d}(1)) \geq \frac{\Delta_{*}}{2\eta\sqrt{d}}\sqrt{\xi_{\min}'(\rho^{2})}\Big) \\ &\leq C_{*}e^{-d/C_{*}} ,\end{aligned}$$

where the last inequality holds by Lemma 2.4 for $\eta < C d^{-1} p_{\max}^{-3} \rho \sqrt{\xi'_{\min}(\rho^2)}$ with C a sufficiently small constant, and $C_* > 0$ an absolute constant. For any $m \ge k(\varepsilon)$ we therefore get, using (81) and (83),

$$\begin{split} \mathbb{P}\Big(\exists \boldsymbol{x} \in \mathsf{B}^{d}(\rho, 1) : \sigma_{n-m+1}(\boldsymbol{x}) \leq \Delta_{*}\sqrt{\xi_{\min}'(\|\boldsymbol{x}\|_{2}^{2})}(\sqrt{d} - \sqrt{n})\Big) \\ &\leq \mathbb{P}\Big(\exists \boldsymbol{x} \in N^{d}(\eta) : \ \sigma_{n-m+1}(\boldsymbol{x}) \leq 2\Delta_{*}\sqrt{\xi_{\min}^{(1)}(\|\boldsymbol{x}\|_{2}^{2})}(\sqrt{d} - \sqrt{n})\Big) + C_{*}e^{-d/C_{*}} \\ &\leq |N^{d}(\eta)|\exp\Big(-A^{2}\varepsilon(d\log d)\Big) + C_{*}e^{-d/C_{*}} \\ &\leq \exp\Big(d\log\frac{10}{\eta} - A^{2}\varepsilon d\log d\Big) + C_{*}e^{-d/C_{*}} \,. \end{split}$$

We then choose $\eta = C d^{-1} p_{\max}^{-3} \rho \sqrt{\xi'_{\min}(\rho^2)}$ and $A = A_{\#}/\sqrt{\varepsilon}$, with $C_{\#}$ a sufficiently large absolute constant to conclude the proof.

We are now in position to prove Lemma 3.3.

Proof of Lemma 3.3. Denote by \mathcal{G}_1 and \mathcal{G}_2 the high probability events of Lemmas 2.1 and 2.4. Further, let \mathcal{G}_3 be the high probability event of Lemma 3.4 with $m = \lfloor (d-n)/4 \rfloor$ and some fixed $\varepsilon < 1/4$. We suppose that $n \leq d - A_{\#}\sqrt{d(\log d)}$. Since ε is fixed, by increasing $C_{\#}$ if needed, we may assume that the conclusion of Lemma 3.4 holds. Namely, that \mathcal{G}_3 holds with probability at least $1 - C_* e^{-d/C_*}$. To complete the proof of the lemma, we will prove that the constant $c_2 > 0$ can be chosen so that $\mathcal{E}_2(c_2, \rho) \supseteq \mathcal{G}_1 \cap \mathcal{G}_2 \cap \mathcal{G}_3$. Hence, we hereafter assume that events $\mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3$ hold.

In the following we will identify $\mathsf{T}_{\boldsymbol{x}}$ with $V^{\perp}(\boldsymbol{x})$ (the orthogonal complement of \boldsymbol{x}) in the obvious way, and therefore identify $\mathsf{T}_{\boldsymbol{x}_1}$, $\mathsf{T}_{\boldsymbol{x}_2}$ when $\boldsymbol{x}_1 = \alpha \boldsymbol{x}_2$ for $\alpha \in \mathbb{R} \setminus \{0\}$. We also recall the rotation $\boldsymbol{R}_{\boldsymbol{x}_1,\boldsymbol{x}_2} \in \mathbb{R}^{d \times d}$ defined Section 2, which maps $\mathsf{T}_{\boldsymbol{x}_1}$ to $\mathsf{T}_{\boldsymbol{x}_2}$. We will use the same notation for the restriction $\boldsymbol{R}_{\boldsymbol{x}_1,\boldsymbol{x}_2}$: $\mathsf{T}_{\boldsymbol{x}_1} \to \mathsf{T}_{\boldsymbol{x}_2}$, which is the parallel transport on the sphere, along the geodesic connecting the two points $\boldsymbol{x}_1/\|\boldsymbol{x}_1\|$ and $\boldsymbol{x}_2/\|\boldsymbol{x}_2\|$. Note that $\boldsymbol{R}_{\boldsymbol{x}_1,\boldsymbol{x}_2} = \boldsymbol{R}_{\boldsymbol{x}_2,\boldsymbol{x}_1}^{-1} = \boldsymbol{R}_{\boldsymbol{x}_2,\boldsymbol{x}_1}^{\mathsf{T}}$

Fix two points $\boldsymbol{x}_0, \boldsymbol{x}_1 \in \mathsf{B}^d(1)$, and define the following linear operators:

$$\boldsymbol{D}_0 := \boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}_0)\big|_{\mathsf{T}_{\boldsymbol{x}_0}}, \quad \boldsymbol{D}_1 := \boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}_1)\big|_{\mathsf{T}_{\boldsymbol{x}_1}}\boldsymbol{R}_{\boldsymbol{x}_0,\boldsymbol{x}_1},$$
(84)

$$\boldsymbol{M}_0 := \boldsymbol{M}(\boldsymbol{x}_0; \boldsymbol{\alpha}), \quad \boldsymbol{M}_1 := \boldsymbol{R}_{\boldsymbol{x}_0, \boldsymbol{x}_1}^{\mathsf{T}} \boldsymbol{M}(\boldsymbol{x}_1; \boldsymbol{\alpha}) \boldsymbol{R}_{\boldsymbol{x}_0, \boldsymbol{x}_1}.$$
(85)

We view D_0, D_1 as linear operators $D_0, D_1 : \mathsf{T}_{\boldsymbol{x}_0} \to \mathbb{R}^n$, and M_0, M_1 as symmetric forms $M_0, M_1 : \mathsf{T}_{\boldsymbol{x}_0} \times \mathsf{T}_{\boldsymbol{x}_0} \to \mathbb{R}$. We note that below restrictions $M_i|_V$ to a linear subspace $V \subset \mathsf{T}_{\boldsymbol{x}_0}$ are defined as $V \times V \to \mathbb{R}$ symmetric forms.

Recall definition (57), which we repeat here for the reader's convenience

$$V_m(\boldsymbol{x}) := \operatorname{span} \left(\boldsymbol{v}_{n-m+2}(\boldsymbol{x}), \dots, \boldsymbol{v}_{d-1}(\boldsymbol{x}) \right) \subseteq \mathsf{T}_{\boldsymbol{x}},$$
(86)

where $v_1(x), \ldots, v_{d-1}(x)$ denote the right singular vectors of $DF(x)|_{\mathsf{T}_x}$ (corresponding to the singular values in decreasing order). In particular, letting $v_{a,1}, \ldots, v_{a,d-1}$ denote the right singular vectors of D_a , we introduce the shorthands

$$V_{0,m} := V_m(\boldsymbol{x}_0) = \operatorname{span}(\boldsymbol{v}_{0,n-m+2}, \dots, \boldsymbol{v}_{0,d-1}), \qquad (87)$$

$$V_{1,s} := \operatorname{span}\left(\boldsymbol{v}_{1,n-s+2}, \dots, \boldsymbol{v}_{1,d-1}\right).$$
(88)

Note that we have $V_{1,s} = \mathbf{R}_{\mathbf{x}_1,\mathbf{x}_0} V_s(\mathbf{x}_1)$ and therefore we obtain the following identities:

$$L_{k,m}(\boldsymbol{x}_0;\boldsymbol{\alpha}) = \lambda_k(\boldsymbol{M}_0|_{V_{0,m}}), \quad L_{s,1}(\boldsymbol{x}_1;\boldsymbol{\alpha}) = \lambda_s(\boldsymbol{M}_1|_{V_{1,1}}).$$
(89)

Recall that $d_m := \dim(V_{0,m}) = d - n + m - 2$. Let $\mathbf{E}_0 \in \mathbb{R}^{d \times d_m}$ be a matrix whose columns form an orthonormal basis of $V_{0,m}$, and $\mathbf{E}_1 \in \mathbb{R}^{d \times d_1}$ a matrix whose columns form an orthonormal basis of $V_{1,1}$. Then we can rewrite the above formulas as

$$L_{k,m}(\boldsymbol{x}_0;\boldsymbol{\alpha}) = \lambda_k(\boldsymbol{E}_0^{\mathsf{T}}\boldsymbol{M}_0\boldsymbol{E}_0), \quad L_{s,1}(\boldsymbol{x}_1;\boldsymbol{\alpha}) = \lambda_s(\boldsymbol{E}_1^{\mathsf{T}}\boldsymbol{M}_1\boldsymbol{E}_1).$$
(90)

By the variational representation of eigenvalues, for $m > s, k \ge s$, we have

$$L_{k,m}(\boldsymbol{x}_0; \boldsymbol{\alpha}) = \max_{\boldsymbol{Q} \in \mathcal{O}(d_m, d_m - k + s)} \lambda_s \left(\boldsymbol{Q}^\mathsf{T} \boldsymbol{E}_0^\mathsf{T} \boldsymbol{M}_0 \boldsymbol{E}_0 \boldsymbol{Q} \right),$$
(91)

where the maximization is over the Stiefel manifold $\mathcal{O}(d_m, d_m - k + s) \subseteq \mathbb{R}^{d_m \times (d_m - k + s)}$ of matrices whose columns form an orthonormal frame.

Let $\overline{E}_0 \in \mathbb{R}^{d \times (d-1-d_m)}$ be such that $[E_0|\overline{E}_0] \in \mathbb{R}^{d \times (d-1)}$ is an orthonormal basis of T_{x_0} . Note that $E_0 E_0^{\mathsf{T}}$ and $\overline{E}_0 \overline{E}_0^{\mathsf{T}}$ are the projections to the column space of E_0 and \overline{E}_0 , respectively. We then have

$$\boldsymbol{E}_{1}^{\mathsf{T}}\boldsymbol{M}_{1}\boldsymbol{E}_{1} = \boldsymbol{E}_{1}^{\mathsf{T}}\boldsymbol{E}_{0}\boldsymbol{E}_{0}^{\mathsf{T}}\boldsymbol{M}_{1}\boldsymbol{E}_{0}\boldsymbol{E}_{0}^{\mathsf{T}}\boldsymbol{E}_{1} + \boldsymbol{E}_{1}^{\mathsf{T}}\overline{\boldsymbol{E}}_{0}\overline{\boldsymbol{E}}_{0}^{\mathsf{T}}\boldsymbol{M}_{1}\boldsymbol{E}_{0}\boldsymbol{E}_{0}^{\mathsf{T}}\boldsymbol{E}_{1} \\ + \boldsymbol{E}_{1}^{\mathsf{T}}\boldsymbol{E}_{0}\boldsymbol{E}_{0}^{\mathsf{T}}\boldsymbol{M}_{1}\overline{\boldsymbol{E}}_{0}\overline{\boldsymbol{E}}_{0}^{\mathsf{T}}\boldsymbol{E}_{1} + \boldsymbol{E}_{1}^{\mathsf{T}}\overline{\boldsymbol{E}}_{0}\overline{\boldsymbol{E}}_{0}^{\mathsf{T}}\boldsymbol{M}_{1}\overline{\boldsymbol{E}}_{0}\overline{\boldsymbol{E}}_{0}^{\mathsf{T}}\boldsymbol{E}_{1}$$

$$\leq \boldsymbol{A}^{\mathsf{T}} \boldsymbol{E}_{0}^{\mathsf{T}} \boldsymbol{M}_{1} \boldsymbol{E}_{0} \boldsymbol{A} + 2 \| \boldsymbol{E}_{1}^{\mathsf{T}} \overline{\boldsymbol{E}}_{0} \|_{\text{op}} \| \boldsymbol{M}_{1} \|_{\text{op}} \cdot \boldsymbol{I} + \| \boldsymbol{E}_{1}^{\mathsf{T}} \overline{\boldsymbol{E}}_{0} \|_{\text{op}}^{2} \| \boldsymbol{M}_{1} \|_{\text{op}} \cdot \boldsymbol{I} \\ \leq \boldsymbol{A}^{\mathsf{T}} \boldsymbol{E}_{0}^{\mathsf{T}} \boldsymbol{M}_{1} \boldsymbol{E}_{0} \boldsymbol{A} + 3 \| \boldsymbol{E}_{1}^{\mathsf{T}} \overline{\boldsymbol{E}}_{0} \|_{\text{op}} \| \boldsymbol{M}_{1} \|_{\text{op}} \cdot \boldsymbol{I} ,$$

where we defined $\mathbf{A} := \mathbf{E}_0^{\mathsf{T}} \mathbf{E}_1 \in \mathbb{R}^{d_m \times d_1}$, $\|\mathbf{A}\|_{\text{op}} \leq 1$ and we write $\mathbf{B}_1 \preceq \mathbf{B}_2$ if $\mathbf{B}_2 - \mathbf{B}_1$ is non-negative definite. Continuing from the previous sequence of inequalities,

$$\boldsymbol{E}_{1}^{\mathsf{T}}\boldsymbol{M}_{1}\boldsymbol{E}_{1} \preceq \boldsymbol{A}^{\mathsf{T}}\boldsymbol{E}_{0}^{\mathsf{T}}\boldsymbol{M}_{0}\boldsymbol{E}_{0}\boldsymbol{A} + \|\boldsymbol{M}_{0} - \boldsymbol{M}_{1}\|_{\mathrm{op}} \cdot \boldsymbol{I} + 3\|\boldsymbol{E}_{1}^{\mathsf{T}}\overline{\boldsymbol{E}}_{0}\|_{\mathrm{op}}\|\boldsymbol{M}_{1}\|_{\mathrm{op}} \cdot \boldsymbol{I}.$$
(92)

Let $\boldsymbol{A} = \boldsymbol{Q}_* \boldsymbol{S} \boldsymbol{O}^\mathsf{T}$, $\boldsymbol{Q}_* \in \mathcal{O}(d_m, d_1)$, $\boldsymbol{O} \in \mathcal{O}(d_1, d_1)$, $\boldsymbol{S} \succeq \boldsymbol{0}$ diagonal, be the reduced singular value decomposition of \boldsymbol{A} (if rank $(\boldsymbol{A}) < d_1$, orthogonal columns are added to $\boldsymbol{Q}_*, \boldsymbol{O}$), and define $\boldsymbol{A}_0 := \boldsymbol{Q}_* \boldsymbol{O}^\mathsf{T}$. Noting that $\|\boldsymbol{S}\|_{\text{op}} \leq 1$ we have

$$egin{aligned} \|oldsymbol{A}-oldsymbol{A}_0\|_{ ext{op}} &= \|oldsymbol{I}-oldsymbol{S}^2\|_{ ext{op}} \ &= \|oldsymbol{I}-oldsymbol{A}^{\mathsf{T}}oldsymbol{E}_0\|_{ ext{op}} = \|oldsymbol{I}-oldsymbol{E}_1^{\mathsf{T}}oldsymbol{E}_0\|_{ ext{op}} \ &= \|oldsymbol{\overline{E}}_0^{\mathsf{T}}oldsymbol{E}_1\|_{ ext{op}}^2, \end{aligned}$$

where for the last equality we used that $I = E_0 E_0^{\mathsf{T}} + \overline{E}_0 \overline{E}_0^{\mathsf{T}} + x_0 x_0^{\mathsf{T}} / ||x_0||^2$ and $x_0^{\mathsf{T}} E_1 = 0$. Hence

$$\boldsymbol{A}^{\mathsf{T}}\boldsymbol{E}_{0}^{\mathsf{T}}\boldsymbol{M}_{0}\boldsymbol{E}_{0}\boldsymbol{A} \leq \boldsymbol{A}_{0}^{\mathsf{T}}\boldsymbol{E}_{0}^{\mathsf{T}}\boldsymbol{M}_{0}\boldsymbol{E}_{0}\boldsymbol{A}_{0} + 3\|\overline{\boldsymbol{E}}_{0}^{\mathsf{T}}\boldsymbol{E}_{1}\|_{\scriptscriptstyle \mathrm{op}}^{2}\|\boldsymbol{M}_{0}\|_{\scriptscriptstyle \mathrm{op}} \cdot \boldsymbol{I}.$$

Substituting this in Eq. (92) and using $\|\boldsymbol{E}_0^{\mathsf{T}}\boldsymbol{E}_1\|_{\text{op}}^2 \leq 1$, we get

$$\boldsymbol{E}_{1}^{\mathsf{T}}\boldsymbol{M}_{1}\boldsymbol{E}_{1} \preceq \boldsymbol{O}\boldsymbol{Q}_{*}^{\mathsf{T}}\boldsymbol{E}_{0}^{\mathsf{T}}\boldsymbol{M}_{0}\boldsymbol{E}_{0}\boldsymbol{Q}_{*}\boldsymbol{O}^{\mathsf{T}} + \|\boldsymbol{M}_{0} - \boldsymbol{M}_{1}\|_{\mathrm{op}} \cdot \boldsymbol{I} + 3\|\boldsymbol{E}_{1}^{\mathsf{T}}\overline{\boldsymbol{E}}_{0}\|_{\mathrm{op}}(\|\boldsymbol{M}_{0}\|_{\mathrm{op}} + \|\boldsymbol{M}_{1}\|_{\mathrm{op}}) \cdot \boldsymbol{I},$$

and therefore

$$\lambda_{s}(\boldsymbol{E}_{1}^{\mathsf{T}}\boldsymbol{M}_{1}\boldsymbol{E}_{1}) \leq \lambda_{s}(\boldsymbol{Q}_{*}^{\mathsf{T}}\boldsymbol{E}_{0}^{\mathsf{T}}\boldsymbol{M}_{0}\boldsymbol{E}_{0}\boldsymbol{Q}_{*}) + \|\boldsymbol{M}_{0} - \boldsymbol{M}_{1}\|_{\text{op}} + 3\|\boldsymbol{E}_{1}^{\mathsf{T}}\overline{\boldsymbol{E}}_{0}\|_{\text{op}}(\|\boldsymbol{M}_{0}\|_{\text{op}} + \|\boldsymbol{M}_{1}\|_{\text{op}}).$$

By Eq. (90) we have $\lambda_s(\boldsymbol{E}_1^{\mathsf{T}}\boldsymbol{M}_1\boldsymbol{E}_1) = L_{s,1}(\boldsymbol{x}_1;\boldsymbol{\alpha})$. Further, by Eq. (91), $\lambda_s(\boldsymbol{Q}_*^{\mathsf{T}}\boldsymbol{E}_0^{\mathsf{T}}\boldsymbol{M}_0\boldsymbol{E}_0\boldsymbol{Q}_*) \leq L_{m+s-1,m}(\boldsymbol{x}_0;\boldsymbol{\alpha})$. We thus obtained

$$L_{s,1}(\boldsymbol{x}_{1};\boldsymbol{\alpha}) \leq L_{m+s-1,m}(\boldsymbol{x}_{0};\boldsymbol{\alpha}) + \|\boldsymbol{M}_{0} - \boldsymbol{M}_{1}\|_{\text{op}} + 3\|\boldsymbol{E}_{1}^{\mathsf{T}}\overline{\boldsymbol{E}}_{0}\|_{\text{op}}(\|\boldsymbol{M}_{0}\|_{\text{op}} + \|\boldsymbol{M}_{1}\|_{\text{op}}).$$
(93)

We finally bound the error terms on the right-hand side of Eq. (93). Recall the definitions of Eqs. (84), (85). We now choose an orthonormal basis on T_{x_0} , which we write as an orthogonal matrix $U_0 \in \mathcal{O}(d, d-1)$, and an orthonormal basis for T_{x_1} , given by $U_1 = \mathbf{R}_{x_0,x_1}U_0$. We then obtain (identifying operators with their matrix representation)

$$D_0 := DF(x_0)U_0, \quad D_1 := DF(x_1)R_{x_0,x_1}U_0 = DF(x_1)U_1,$$
 (94)

$$\boldsymbol{M}_{0} := \sum_{\ell=1}^{n} \alpha_{\ell} \boldsymbol{U}_{0}^{\mathsf{T}} \nabla^{2} F_{\ell}(\boldsymbol{x}_{0}) \boldsymbol{U}_{0}, \quad \boldsymbol{M}_{1} := \sum_{\ell=1}^{n} \alpha_{\ell} \boldsymbol{U}_{1}^{\mathsf{T}} \nabla^{2} F_{\ell}(\boldsymbol{x}_{1}) \boldsymbol{U}_{1}.$$
(95)

We then have

$$\|\boldsymbol{M}_{0} - \boldsymbol{M}_{1}\|_{\text{op}} \leq \|\boldsymbol{\alpha}\|_{1} \max_{\ell \leq n} \|\boldsymbol{U}_{0}^{\mathsf{T}} \nabla^{2} F_{\ell}(\boldsymbol{x}_{0}) \boldsymbol{U}_{0} - \boldsymbol{U}_{1}^{\mathsf{T}} \nabla^{2} F_{\ell}(\boldsymbol{x}_{1}) \boldsymbol{U}_{1}\|_{\text{op}}$$
(96)

$$\leq \frac{C_0}{\rho} dp_{\max}^3 \sqrt{\log(p_{\max})} \|\boldsymbol{x}_0 - \boldsymbol{x}_1\|, \qquad (97)$$

where we used $\|\boldsymbol{\alpha}\|_1 \leq \sqrt{n} \leq \sqrt{d}$. The last inequality holds on the event \mathcal{G}_2 of Lemma 2.4.

By a similar argument,

$$\|\boldsymbol{M}_1\|_{\rm op} \le \|\boldsymbol{\alpha}\|_1 \max_{\ell \le n} \|\nabla^2 F_\ell(\boldsymbol{x}_0)\|_{\rm op} \le C_1 dp_{\max}^2 \sqrt{\log(p_{\max})} \,.$$
(98)

on the event \mathcal{G}_1 of Lemma 2.1, and similarly for $\|\boldsymbol{M}_0\|_{\text{op}}$.

Further,

$$\|\boldsymbol{D}_{0} - \boldsymbol{D}_{1}\|_{\text{op}} = \|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}_{0})\boldsymbol{U}_{0} - \boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}_{1})\boldsymbol{U}_{1}\|_{\text{op}}$$

$$\leq \frac{C_{0}}{\rho}p_{\text{max}}^{2}\sqrt{d\log(p_{\text{max}})}\|\boldsymbol{x}_{0} - \boldsymbol{x}_{1}\|,$$
(99)

where the last step we used once more the fact that event \mathcal{G}_2 of Lemma 2.4 holds.

Finally, by Corollary A.1 (applied with $A_i = D_i$, and d replaced by d - 1), we get

$$\begin{aligned} \left\| \boldsymbol{E}_{1}^{\mathsf{T}} \overline{\boldsymbol{E}}_{0} \right\|_{\mathrm{op}} &\leq \frac{1}{\sigma_{n-m+1}(\boldsymbol{D}_{0})} \left\| \boldsymbol{D}_{0} - \boldsymbol{D}_{1} \right\|_{\mathrm{op}} \\ &\leq \frac{(a)}{\leq} \frac{C_{0}/C_{*}}{(\sqrt{d} - \sqrt{n})\xi'_{\min}(\|\boldsymbol{x}_{0}\|^{2})^{1/2}\rho} p_{\max}^{2} \sqrt{d\log(p_{\max})} \|\boldsymbol{x}_{0} - \boldsymbol{x}_{1}\| \end{aligned} \tag{100}$$

$$\stackrel{(b)}{\leq} \frac{2C_0/C_* dp_{\max}^2 \sqrt{\log(p_{\max})}}{\xi_{\min}'(\|\boldsymbol{x}_0\|^2)^{1/2} \rho} \|\boldsymbol{x}_0 - \boldsymbol{x}_1\|.$$
(101)

Here (a) holds by the bound (99), on the event \mathcal{G}_3 of Lemma 3.4 (with $m = \lfloor (d-n)/4 \rfloor$) and (b) since, we have $\sqrt{d} - \sqrt{n} \ge (d-n)/(2\sqrt{d}) \ge 1/(2\sqrt{d})$.

Using the bounds (97), (98), and (101) in Eq. (93), we finally obtain

$$\begin{split} L_{s,1}(\boldsymbol{x}_{1};\boldsymbol{\alpha}) &\leq L_{m+s-1,m}(\boldsymbol{x}_{0};\boldsymbol{\alpha}) \frac{C_{0}}{\rho} dp_{\max}^{3} \sqrt{\log(p_{\max})} \|\boldsymbol{x}_{0} - \boldsymbol{x}_{1}\| + \frac{Cd^{2}p_{\max}^{4}\log(p_{\max})}{\xi_{\min}'(\|\boldsymbol{x}_{0}\|^{2})^{1/2}\rho} \|\boldsymbol{x}_{0} - \boldsymbol{x}_{1}\| \\ &\leq L_{m+s-1,m}(\boldsymbol{x}_{0};\boldsymbol{\alpha}) + \frac{C'd^{2}p_{\max}^{5}}{\xi_{\min}'(\rho^{2})^{1/2}\rho} \|\boldsymbol{x}_{0} - \boldsymbol{x}_{1}\|, \end{split}$$

for some absolute constants C, C' > 0, which concludes the proof.

4 Minimal singular value at solutions

In Section 6 we will prove that w.h.p. any point sufficiently close to a solution is an approximate solution. The distance from the solution will be controlled by Lipschitz constants which we have already analyzed in Section 2, and the minimal singular value of the Jacobian DF(x) at the solution which we study in this section.

In the analysis of the brute-force search algorithm (Algorithm 2) that applies to $p_{\text{max}} > d^2$, we will use the following proposition, controlling the *maximal* value of $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}})$ over all solutions.

Proposition 4.1. Suppose n = d - 1. For some universal constants $C, \eta > 0$,

$$\mathbb{P}\left(\exists \boldsymbol{x} \in \mathbb{S}^{d-1} : \boldsymbol{F}(\boldsymbol{x}) = 0, \, \sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}}) \ge \frac{\eta}{d^{7/4}}\right) \ge 1 - \frac{C}{\sqrt{d}}.$$
(102)

In contrast, in the analysis of the Hessian descent algorithm (Algorithm 1) for $p_{\text{max}} \leq d^2$, we will use the following proposition, concerning the *minimal* value of $\sigma_{\min}(DF(x)U_x)$ over the solutions. **Proposition 4.2.** There exist universal constants C, C' > 0 such that, defining, for $d/2 \le n \le d-1$,

$$\tau := \left(e^{Cd} p_{\max}^{d - \frac{3}{4}n - 1} \log(p_{\max})^{\frac{1}{4}(d - n - 1)} d^{\frac{1}{4}(d - n)} \right)^{-1},$$
(103)

we have that

$$\mathbb{P}\left(\min\left\{\sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}}):\,\boldsymbol{x}\in\mathbb{S}^{d-1},\,\boldsymbol{F}(\boldsymbol{x})=0\right\}\geq\tau\right)\geq1-C'\exp(-d/C').$$
(104)

4.1 Upper bound on volume in expectation

Consider the set

$$\Psi(\tau) := \left\{ \boldsymbol{x} \in \mathbb{S}^{d-1} : \boldsymbol{F}(\boldsymbol{x}) = 0, \, \sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}}) \leq \tau \right\}.$$

Denote by Vol_k the k-dimensional Hausdorff measure. We start by proving an upper bound for the volume of $\Psi(\tau)$ in expectation. This bound will be used in the proof of both propositions above.

Lemma 4.3. Let $Z \sim \text{GOE}(n, d-1)$. Then,

$$\mathbb{E}\operatorname{Vol}_{d-1-n}(\Psi(\tau)) \leq \operatorname{Vol}_{d-1}(\mathbb{S}^{d-1}) \prod_{i \leq n} \sqrt{\frac{p_i}{2\pi}} \mathbb{E}\left(|\det(\boldsymbol{Z}\boldsymbol{Z}^{\intercal})|^{1/2} \mathbb{1}\left\{ \sigma_{\min}(\boldsymbol{Z}) \leq \tau \right\} \right).$$
(105)

Moreover, if we assume that $\tau \leq 1/n$, then for some universal constant $C_1 > 0$,

$$\mathbb{E}\operatorname{Vol}_{d-1-n}(\Psi(\tau)) \leq \operatorname{Vol}_{d-1}(\mathbb{S}^{d-1}) \prod_{i \leq n} \sqrt{\frac{p_i}{2\pi}} \mathbb{E}\left(|\det(\boldsymbol{Z}\boldsymbol{Z}^{\intercal})|^{1/2} \right) \cdot C_1 d^{5/2} \tau \mathbb{P}\left(\sigma_{\min}(\boldsymbol{Z}) \leq \tau\right).$$
(106)

Proof. The expectation $\mathbb{E}\text{Vol}_{d-1-n}(\Psi(\tau))$ can be expressed by a variant of the Kac-Rice formula, e.g. as in [AW09, Theorem 6.8] (see [Sub23, Remark 3] for a discussion on the regularity conditions required for its application). Precisely, abbreviating $\boldsymbol{\Delta}(\boldsymbol{x}) = \boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}}$, the formula gives

$$\begin{split} & \mathbb{E}\operatorname{Vol}_{d-1-n}(\Psi(\tau)) \\ &= \int_{\mathbb{S}^{d-1}} \varphi_{F(\boldsymbol{x})}(0) \mathbb{E}\Big[|\det(\boldsymbol{\Delta}(\boldsymbol{x})\boldsymbol{\Delta}(\boldsymbol{x})^{\mathsf{T}})|^{1/2} \mathbb{1}\{\sigma_{\min}(\boldsymbol{\Delta}(\boldsymbol{x})) \leq \tau\} \ \Big| \ \boldsymbol{F}(\boldsymbol{x}) = 0 \Big] \mathrm{d}\operatorname{Vol}_{d-1}(\boldsymbol{x}) \\ &= \operatorname{Vol}_{d-1}(\mathbb{S}^{d-1})(2\pi)^{-n/2} \mathbb{E}\Big[|\det(\boldsymbol{\Delta}(\boldsymbol{x})\boldsymbol{\Delta}(\boldsymbol{x})^{\mathsf{T}})|^{1/2} \mathbb{1}\{\sigma_{\min}(\boldsymbol{\Delta}(\boldsymbol{x})) \leq \tau\} \ \Big| \ \boldsymbol{F}(\boldsymbol{x}) = 0 \Big] \,, \end{split}$$

where $\varphi_{F(x)}(0) = (2\pi)^{-n/2}$ is the density of the random vector F(x) evaluated at the origin and, using symmetry, in the second line x is an arbitrary point in \mathbb{S}^{d-1} .

Recall that by Lemma 3.1, conditional on $F(\mathbf{x}) = 0$, $\Delta(\mathbf{x}) \stackrel{d}{=} S_1(1)\mathbf{Z}$ where $S_1(1) := \text{diag}(\sqrt{p_i}: i \leq n)$ and $\mathbf{Z} \sim \text{GOE}(n, d-1)$. Note that $\sigma_{\min}(S_1(1)\mathbf{Z}) \geq \sigma_{\min}(\mathbf{Z})$, since $\sigma_{\min}(\mathbf{A}) = \min_{\mathbf{v} \in \mathbb{S}^{n-1}} \max_{\mathbf{u} \in \mathbb{S}^{d-2}} \mathbf{v}^{\mathsf{T}} \mathbf{A} \mathbf{u}$ for any $\mathbf{A} \in \mathbb{R}^{n \times (d-1)}$. This proves (105).

The random matrix $\mathbf{Z}\mathbf{Z}^{\intercal}$ is a Wishart matrix. Recall that its eigenvalues $\lambda_1 \leq \cdots \leq \lambda_n$ have density $c_d\phi_{n,d-1}$ where c_d is a normalizing constant and

$$\phi_{n,d-1}(x_1,\ldots,x_n) = \prod_i x_i^{(d-n-2)/2} e^{-x_i/2} \prod_{i< j} |x_i - x_j|$$
(107)

on $B_n := \{(x_1, \dots, x_n) : 0 < x_1 < \dots < x_n\}$. Since

$$\phi_{n,d-1}(x_1,\ldots,x_n) = x_1^{(d-n-2)/2} e^{-x_1/2} \prod_{i=2}^n |x_i - x_1| \phi_{n-1,d-2}(x_2,\ldots,x_n),$$

by conditioning on $\sigma_{\min}(\mathbf{Z})^2 = \lambda_1$,

$$\mathbb{E}\Big[\left|\det(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}})\right|^{1/2} \left|\sigma_{\min}(\boldsymbol{Z}) \leq \tau\right] = \int_{0}^{\tau^{2}} \frac{\mathbb{E}\Big[\prod_{i=1}^{n-1} \bar{\lambda}_{i}^{1/2} |\bar{\lambda}_{i} - t| \mathbb{1}\{\bar{\lambda}_{1} > t\}\Big]}{\mathbb{E}\Big[\prod_{i=1}^{n-1} |\bar{\lambda}_{i} - t| \mathbb{1}\{\bar{\lambda}_{1} > t\}\Big]} f_{n,d-1}(t)\sqrt{t} \,\mathrm{d}t\,, \quad (108)$$

where $\bar{\lambda}_1 \leq \cdots \leq \bar{\lambda}_{n-1}$ are the eigenvalues of $\bar{Z}\bar{Z}^{\mathsf{T}}$ with $\bar{Z} \sim \mathsf{GOE}(n-1, d-2)$ and $f_{n,d-1}(t)$ is the density of λ_1 normalized so that $\int_0^{\tau^2} f_{n,d-1}(t) dt = 1$. The numerator on the right-hand side of (108) is bounded from above by

$$\mathbb{E}\left[|\det(\bar{\boldsymbol{Z}}\bar{\boldsymbol{Z}}^{\mathsf{T}})|^{3/2}
ight].$$

Using the density (107), we can write the denominator as

$$c_{d-1} \int_{B_{n-1}+t} \prod_{i=1}^{n-1} |x_i - t| x_i^{(d-n-2)/2} e^{-x_i/2} \prod_{i < j} |x_i - x_j| dx_1 \cdots dx_{n-1}$$

$$\geq c_{d-1} \int_{B_{n-1}} \prod_{i=1}^{n-1} |x_i| x_i^{(d-n-2)/2} e^{-(x_i+t)/2} \prod_{i < j} |x_i - x_j| dx_1 \cdots dx_{n-1} \geq \frac{1}{2} \mathbb{E} \Big[|\det(\bar{\boldsymbol{Z}}\bar{\boldsymbol{Z}}^{\mathsf{T}})| \Big],$$

where by an abuse of notation we denote by $B_{n-1} + t$ the Minkowski sum of B_{n-1} and $\{(t, \ldots, t)\}$ and the inequality holds if t < 1/n.

Recall that for any matrix $\mathbf{A} \in \mathbb{R}^{M \times N}$, $|\det(\mathbf{A}\mathbf{A}^{\intercal})|^{1/2} = \prod_{i \leq M} ||\Theta_i(\mathbf{A})||$, where $\Theta_i(\mathbf{A})$ is the projection of the *i*-th row of \boldsymbol{A} onto the orthogonal space to its first i-1 rows. Hence, $|\det(\bar{\boldsymbol{Z}}\bar{\boldsymbol{Z}}^{\mathsf{T}})| = \prod_{i=1}^{n-1} \chi_{d-1-i}^2$ and $|\det(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}})| = \prod_{i=1}^n \chi_{d-i}^2$ in distribution, where χ_k^2 are i.i.d. chi-squared variables of k degrees of freedom. Using this, one can verify that, for absolute C > 0,

$$\frac{\mathbb{E}\left[|\det(\bar{\boldsymbol{Z}}\bar{\boldsymbol{Z}}^{\mathsf{T}})|^{3/2}\right]}{\mathbb{E}\left[|\det(\bar{\boldsymbol{Z}}\bar{\boldsymbol{Z}}^{\mathsf{T}})|\right]\mathbb{E}\left[|\det(\bar{\boldsymbol{Z}}\bar{\boldsymbol{Z}}^{\mathsf{T}})|^{1/2}\right]} \le Cd^2 \quad \text{and} \quad \frac{\mathbb{E}\left[|\det(\bar{\boldsymbol{Z}}\bar{\boldsymbol{Z}}^{\mathsf{T}})|^{1/2}\right]}{\mathbb{E}\left[|\det(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}})|^{1/2}\right]} \le C/\sqrt{d}.$$

Combining the above with (108) and plugging back into (105) completes the proof.

4.2Lower bound on the volume of a neighborhood

Recall the definitions of $\operatorname{Lip}(DF; \Omega)$ and $\operatorname{Lip}(DF; \Omega)$ in Eqs. (40), (42).

Lemma 4.4. Let $c, \tau, r > 0$ such that r < 1/10 and $(1-r)^{p_{\max}\tau} - rc\sqrt{d} \ge \tau/2$. On the event that

$$\operatorname{Lip}_{\perp}(\boldsymbol{DF}; \mathbb{S}^{d-1}(1)) \vee \operatorname{Lip}(\boldsymbol{DF}; \mathsf{B}^{d}(1)) \leq c\sqrt{d},$$
(109)

if there exists a point $\mathbf{x} \in \mathbb{S}^{d-1}(1)$ such that $\mathbf{F}(\mathbf{x}) = 0$ and $\sigma_{\min}(\mathbf{DF}(\mathbf{x})\mathbf{U}_{\mathbf{x}}) \in [\tau, 2\tau]$ then

$$\operatorname{Vol}_{d-n-1}(\Psi(3\tau)) \ge \left(\frac{r}{2\sqrt{2}}\right)^{d-n-1} \operatorname{Vol}_{d-n-1}(\mathsf{B}^{d-n-1}(1)).$$
 (110)

Proof. Assume throughout the proof that the event in (109) occurs and let x be a point as in the lemma. Since the F_i are homogeneous, DF(x)x = 0. Setting t = 1 - r, for z = tx we have that $DF(z) = \operatorname{diag}(t^{p_i-1}: 1 \le i \le n)DF(z) \text{ and } \sigma_{\min}(DF(z)) = \sigma_{\min}(DF(z)U_z) \in t^{p_{\max}}[\tau, 2\tau].$

Let $V := \{ \boldsymbol{D} \boldsymbol{F}(\boldsymbol{z})^{\mathsf{T}} \cdot \boldsymbol{v} : \boldsymbol{v} \in \mathbb{R}^n \} \subset \mathsf{T}_{\boldsymbol{z}}$, let $\boldsymbol{T} \in \mathbb{R}^{d \times n}$ be a matrix whose columns are an orthonormal basis of V and let $\boldsymbol{K} \in \mathbb{R}^{d \times (d-n-1)}$ be a matrix whose columns are an orthonormal basis of $V^{\perp} \cap \mathsf{T}_{\boldsymbol{z}}$. For $(\boldsymbol{a}, \boldsymbol{b}) \in \mathbb{R}^{d-n-1} \times \mathbb{R}^n$, define

$$\ddot{F}(a, b) = F(z + K \cdot a + T \cdot b),$$

and note that

$$D_1 \hat{F}(\boldsymbol{a}, \boldsymbol{b}) := \left(\frac{\mathrm{d}}{\mathrm{d}a_j} \hat{F}_i(\boldsymbol{a}, \boldsymbol{b})\right)_{i \le n, j \le d-n-1} = DF(\boldsymbol{z} + \boldsymbol{K} \cdot \boldsymbol{a} + \boldsymbol{T} \cdot \boldsymbol{b}) \cdot \boldsymbol{K},$$
$$D_2 \hat{F}(\boldsymbol{a}, \boldsymbol{b}) := \left(\frac{\mathrm{d}}{\mathrm{d}b_j} \hat{F}_i(\boldsymbol{a}, \boldsymbol{b})\right)_{i \le n, j \le n} = DF(\boldsymbol{z} + \boldsymbol{K} \cdot \boldsymbol{a} + \boldsymbol{T} \cdot \boldsymbol{b}) \cdot \boldsymbol{T}.$$

Since $\lambda_{\min}(\boldsymbol{D}_2 \hat{\boldsymbol{F}}(0,0)) = \sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{z}))$ and, for i = 1, 2,

$$\begin{split} \|\boldsymbol{D}_{i}\hat{\boldsymbol{F}}(\boldsymbol{a},\boldsymbol{b}) - \boldsymbol{D}_{i}\hat{\boldsymbol{F}}(0,0)\|_{\text{op}} &\leq \|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{z} + \boldsymbol{K}\cdot\boldsymbol{a} + \boldsymbol{T}\cdot\boldsymbol{b}) - \boldsymbol{D}\boldsymbol{F}(\boldsymbol{z})\|_{\text{op}} \\ &\leq \|\boldsymbol{K}\cdot\boldsymbol{a} + \boldsymbol{T}\cdot\boldsymbol{b}\|\operatorname{Lip}(\boldsymbol{D}\boldsymbol{F};\mathsf{B}^{d}(1)) \leq c\sqrt{d}\|(\boldsymbol{a},\boldsymbol{b})\|\,, \end{split}$$

if $\|(\boldsymbol{a}, \boldsymbol{b})\| = \|\boldsymbol{K} \cdot \boldsymbol{a} + \boldsymbol{T} \cdot \boldsymbol{b}\| \le r$ then, using the assumption $(1-r)^{p_{\max}\tau} - rc\sqrt{d} \ge \tau/2$,

$$\sigma_{\min}(\boldsymbol{D}_{2}\hat{\boldsymbol{F}}(\boldsymbol{a},\boldsymbol{b})), \ \sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{z}+\boldsymbol{K}\cdot\boldsymbol{a}+\boldsymbol{T}\cdot\boldsymbol{b})) \geq \tau/2.$$
(111)

Suppose that $(\boldsymbol{a}, \boldsymbol{b})$ is a point such that $\|(\boldsymbol{a}, \boldsymbol{b})\| < r$ and $\hat{\boldsymbol{F}}(\boldsymbol{a}, \boldsymbol{b}) = 0$. Then by the implicit function theorem, there is a unique continuously differentiable function g from an open neighborhood of \boldsymbol{a} to \mathbb{R}^n such that $g(\boldsymbol{a}) = \boldsymbol{b}$ and for any \boldsymbol{a}' in this neighborhood $\hat{\boldsymbol{F}}(\boldsymbol{a}', g(\boldsymbol{a}')) = 0$. From compactness, there exists a closed set $A \subset \mathbb{R}^{d-n-1}$ containing the origin and a unique $g: A \to \mathbb{R}^n$ continuous on A and continuously differentiable on its interior A^o such that $g(\boldsymbol{0}) = \boldsymbol{0}$ and $\hat{\boldsymbol{F}}(\boldsymbol{a}, g(\boldsymbol{a})) = 0$ for any $\boldsymbol{a} \in A$, and such that $\|(\boldsymbol{a}, g(\boldsymbol{a}))\| < r$ on A^o and $\|(\boldsymbol{a}, g(\boldsymbol{a}))\| = r$ on $\partial A = A \setminus A^o$. Moreover, by the implicit function theorem, on A^o ,

$$\boldsymbol{D}g(\boldsymbol{a}) := \left(\frac{\mathrm{d}}{\mathrm{d}a_j}g_i(\boldsymbol{a})\right)_{i \le n, j \le d-n-1} = -\boldsymbol{D}_2\hat{\boldsymbol{F}}(\boldsymbol{a}, g(\boldsymbol{a}))^{-1} \cdot \boldsymbol{D}_1\hat{\boldsymbol{F}}(\boldsymbol{a}, g(\boldsymbol{a}))^{-1}$$

Since $\boldsymbol{D}_1 \hat{\boldsymbol{F}}(\boldsymbol{0}, g(\boldsymbol{0})) = \boldsymbol{0} \in \mathbb{R}^{n \times (d-n-1)}$, on $\|(\boldsymbol{a}, g(\boldsymbol{a}))\| < r$,

$$\|\boldsymbol{D}_1 \hat{\boldsymbol{F}}(\boldsymbol{a}, g(\boldsymbol{a}))\|_{ ext{op}} \leq c\sqrt{d} \cdot \|(\boldsymbol{a}, g(\boldsymbol{a}))\| < \tau/2$$

and therefore, by Eq. (111),

$$\|\boldsymbol{D}g(\boldsymbol{a})\|_{\text{op}} < 1.$$

Define $B = \{ \boldsymbol{a} \in \mathbb{R}^{d-n-1} : \|\boldsymbol{a}\| < r/\sqrt{2} \}$. Obviously, $A \supset B$ for otherwise we would have some $\boldsymbol{a} \in \partial A \cap B$ for which $\|(\boldsymbol{a}, g(\boldsymbol{a}))\| = r$ while $\|\boldsymbol{a}\| \vee \|g(\boldsymbol{a})\| < r/\sqrt{2}$. Define $P(\boldsymbol{y}) := \boldsymbol{y}/\|\boldsymbol{y}\|$ and $\boldsymbol{y}(\boldsymbol{a}) = \boldsymbol{z} + \boldsymbol{K} \cdot \boldsymbol{a} + \boldsymbol{T} \cdot g(\boldsymbol{a})$. We have that

$$\left\{ \boldsymbol{y} \in \mathbb{S}^{d-1} : \boldsymbol{F}(\boldsymbol{y}) = 0 \right\} \supset P\left(\left\{ \boldsymbol{y}(\boldsymbol{a}) : \boldsymbol{a} \in B \right\} \right).$$
(112)

We note that (using that r < 1/10) for any $a_1, a_2 \in B$,

$$\|P(\boldsymbol{y}(\boldsymbol{a}_{1})) - P(\boldsymbol{y}(\boldsymbol{a}_{2}))\| \ge \frac{1}{2} \left\| \frac{\boldsymbol{y}(\boldsymbol{a}_{1})}{\|\boldsymbol{y}(\boldsymbol{a}_{1})\| \wedge \|\boldsymbol{y}(\boldsymbol{a}_{2})\|} - \frac{\boldsymbol{y}(\boldsymbol{a}_{2})}{\|\boldsymbol{y}(\boldsymbol{a}_{1})\| \wedge \|\boldsymbol{y}(\boldsymbol{a}_{2})\|} \right\| \ge \frac{1}{2} \|\boldsymbol{a}_{1} - \boldsymbol{a}_{2}\|.$$
(113)

It thus follows that

$$\operatorname{Vol}_{d-n-1}\left(P\left(\left\{\boldsymbol{y}(\boldsymbol{a}):\,\boldsymbol{a}\in B\right\}\right)\right) \ge 2^{-(d-n-1)}\operatorname{Vol}_{d-n-1}(B)$$
$$= \left(\frac{r}{2\sqrt{2}}\right)^{d-n-1}\operatorname{Vol}_{d-n-1}\left(\mathsf{B}^{d-n-1}(1)\right).$$

Moreover, for any \boldsymbol{a} in the set on the right-hand side of (112), if $\boldsymbol{w} = P(\boldsymbol{y}(\boldsymbol{a}))$ then $\|\boldsymbol{w} - \boldsymbol{x}\| \leq 2r$ and thus $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{w})\boldsymbol{U}_{\boldsymbol{w}}) \leq 2\tau + 2rc\sqrt{d} \leq 3\tau$.

4.3 **Proof of Proposition 4.1**

We assume throughout this proof that n = d-1, and therefore $\operatorname{Vol}_{d-1-n}(\Psi(\tau)) =: |\Psi(\tau)|$ is just the cardinality of the set $\Psi(\tau)$. Note that $\Psi(\infty) = \{ \boldsymbol{x} \in \mathbb{S}^{d-1} : \boldsymbol{F}(\boldsymbol{x}) = 0 \}$. By Eq. (2.2) of [Sub23], we have that Eq. (105) holds with equality for $\tau = \infty$, namely

$$\mathbb{E} |\Psi(\infty)| = \operatorname{Vol}_{d-1}(\mathbb{S}^{d-1}) \prod_{i \le n} \sqrt{\frac{p_i}{2\pi}} \mathbb{E} \left(|\det(\mathbf{Z}\mathbf{Z}^{\intercal})|^{1/2} \right).$$
(114)

Using Eq. (106) in Lemma 4.3 and Corollary B.4, we obtain that, for some absolute constant C_0 ,

$$\frac{\mathbb{E}|\Psi(\tau)|}{\mathbb{E}|\Psi(\infty)|} \le C_0 d^3 \tau^2$$

Hence, by Markov's inequality,

$$\mathbb{P}\Big(|\Psi(\tau)| \ge \frac{1}{2}\mathbb{E}|\Psi(\infty)|\Big) \le 2C_0 d^3 \tau^2 \,.$$

By Theorem 2 of [Sub23], for some absolute constant C_1 ,

$$\mathbb{P}\Big(|\Psi(\infty)| < \frac{1}{2}\mathbb{E}|\Psi(\infty)|\Big) \le C_1 d^{-1/2}.$$

The proof is completed since

$$\mathbb{P}(\Psi(\infty) \setminus \Psi(\tau) \neq \varnothing) \ge 1 - C_1 d^{-1/2} - 2C_0 d^3 \tau^2.$$

4.4 **Proof of Proposition 4.2**

Let C_0 and C_1 be the absolute constants as in Lemmas 2.4 and 4.3 and denote by \mathcal{E} the event that (45) occurs. For any $\tau < 1/n$, from Lemmas 4.3 and 4.4 and Markov's inequality,

$$\mathbb{P}\left(\mathcal{E} \cap \left\{ \exists \boldsymbol{x} \in \mathbb{S}^{d-1} : \boldsymbol{F}(\boldsymbol{x}) = 0, \, \sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}}) \in [\tau, 2\tau] \right\} \right) \leq M_{d,n}(\tau) \,,$$

where

$$M_{d,n}(\tau) := \frac{\operatorname{Vol}_{d-1}(\mathbb{S}^{d-1}) \prod_{i \le n} \sqrt{\frac{p_i}{2\pi}} \mathbb{E}\left(|\det(\mathbf{Z}\mathbf{Z}^{\intercal})|^{1/2} \right) \cdot C_1 d^{5/2} 3\tau \mathbb{P}\left(\sigma_{\min}(\mathbf{Z}) \le 3\tau\right)}{\left(\frac{r}{2\sqrt{2}}\right)^{d-n-1} \operatorname{Vol}_{d-n-1}\left(\mathsf{B}^{d-n-1}(1)\right)}, \quad (115)$$

provided that that

$$r < \frac{1}{10}, \qquad (1-r)^{p_{\max}} \tau - rC_0 p_{\max}^2 \sqrt{\log(p_{\max})d} \ge \frac{\tau}{2}.$$
 (116)

To satisfy these two inequalities, we shall assume without loss of generality that $C_0 > 1$, and set

$$r = \frac{\tau}{4C_0 p_{\max}^2 \sqrt{\log(p_{\max})d}}.$$
(117)

Recall that $\operatorname{Vol}_{k-1}(\mathbb{S}^{k-1}) = 2\pi^{k/2}/\Gamma(\frac{k}{2})$ and $\operatorname{Vol}_k(\mathsf{B}^k(1)) = \pi^{(k/2)}/\Gamma(k/2+1)$ and $\mathbb{E}|\det(\mathbf{Z}\mathbf{Z}^{\intercal})|^{1/2} = 2^{n/2}\Gamma(d/2)/\Gamma((d-n)/2)$ (e.g., by the relation to chi-variables as in the proof of Lemma 4.3). Hence,

$$\frac{\operatorname{Vol}_{d-1}(\mathbb{S}^{d-1})\prod_{i\leq n}\sqrt{\frac{1}{2\pi}}\mathbb{E}\left(|\det(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}})|^{1/2}\right)}{\operatorname{Vol}_{d-n-1}\left(\mathsf{B}^{d-n-1}(1)\right)}\leq c\sqrt{d-n}\,,$$

for some universal c > 0. By Corollary B.4, for universal c' > 0,

$$\mathbb{P}\Big(\sigma_{\min}(\mathbf{Z}) \le 3\tau\Big) \le \Big(\frac{c'\tau}{\sqrt{d-1} - \sqrt{n-1}}\Big)^{d-n}.$$

Combining the above and substituting r as in (117), we have that $M_{d,n}(\tau)$, defined in Eq. (115), is bounded as

$$M_{d,n}(\tau) \le c_0^{d-n} p_{\max}^{\frac{n}{2}} d^{5/2} \sqrt{d-n\tau} \left(\frac{p_{\max}^2 \sqrt{\log(p_{\max})d}}{\tau}\right)^{d-n-1} \left(\frac{\tau}{\sqrt{d-1}-\sqrt{n-1}}\right)^{d-n} \le \tau^2 \cdot c_0^{d-n} p_{\max}^{\frac{n}{2}} d^3 \left(p_{\max}^2 \sqrt{\log(p_{\max})d}\right)^{d-n-1} =: \tau^2 A,$$

for some universal constant $c_0 > 0$.

By a union bound,

$$\mathbb{P}\left(\mathcal{E} \cap \left\{\exists \boldsymbol{x} \in \mathbb{S}^{d-1}: \boldsymbol{F}(\boldsymbol{x}) = 0, \, \sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}}) \leq \tau\right\}\right) \leq \sum_{i=1}^{\infty} (2^{-i}\tau)^2 A < \tau^2 A.$$

Since $\mathbb{P}(\mathcal{E}) \geq 1 - C_* \exp(-d/C_*)$ by Lemma 2.4, taking τ as in (103) with large enough C, this completes the proof.

5 Good events

Our algorithms output an approximate solution on an event with high probability, which we now define precisely. We treat the two cases $p_{\max} \leq d^2$ and $p_{\max} > d^2$ separately. Denote the left-hand side of (44), (45) and (46) by $\operatorname{Lip}^{(0)}(\mathbf{F})$, $\operatorname{Lip}^{(1)}(\mathbf{F})$ and $\operatorname{Lip}^{(2)}(\mathbf{F})$, respectively, with $\Omega = \mathsf{B}^d(1)$. With $\rho = 1$ and $A_{\#}$ and $B_{\#}$ as in Theorem 3, define $A = 4A_{\#} \vee B_{\#}$ so that $A(d \log d)^{1/2} \geq 4A_{\#}(d \log d)^{1/2}$ and $A(d \log d)^{1/2} \geq B_{\#}d^{1/2}$. Let C, C_0 and C_1 be the absolute constants from Proposition 4.2 and Lemmas 2.4 and 2.1, and let $\eta > 0$ be the absolute constant as in Proposition 4.1. As in (103), we define

$$\tau := \left(e^{Cd} p_{\max}^{d-\frac{3}{4}n-1} \log(p_{\max})^{\frac{1}{4}(d-n-1)} d^{\frac{1}{4}(d-n+5)} \right)^{-1}$$

Definition 5.1 (Good event for $p_{\max} \leq d^2$). Suppose that $p_{\max} \leq d^2$ and $\frac{d}{2} \leq n \leq d - A(d \log d)^{1/2}$. We denote by $\mathcal{E} = \mathcal{E}(d, n)$ the event that:

1. $\max_{\boldsymbol{x}\in\mathsf{B}^d(1)}\|\boldsymbol{F}(\boldsymbol{x})\|_2 \leq C_1\sqrt{d\log(p_{\max})}.$

2.
$$\operatorname{Lip}^{(i)}(\mathbf{F}) \leq C_0 p_{\max}^{1+i} \sqrt{d \log(p_{\max})}$$
, for $i = 0, 1, 2$.
3. $\min \left\{ \sigma_{\min}(\mathbf{DF}(\mathbf{x})\mathbf{U}_{\mathbf{x}}) : \mathbf{x} \in \mathbb{S}^{d-1}, \mathbf{F}(\mathbf{x}) = 0 \right\} \geq \tau$.
4. $\forall \mathbf{x} \in \mathbb{S}^{d-1} : \lambda_1(\mathcal{H}(\mathbf{x})) \leq -\frac{1}{10}\sqrt{(d-n)H(\mathbf{x})}$.
5. $\forall \mathbf{x} \in \mathsf{B}^d(1, 1+1/p_{\max}), 2 \leq k \leq 4 : \|\nabla^k H(\mathbf{x})\|_{\operatorname{op}} \leq 9C_1 dp_{\max}^k \log(p_{\max})$.
6. For $\mathbf{x}^0 = (1, 0, \dots, 0), H(\mathbf{x}^0) \leq d$.

Lemma 5.2. Assume the setting of Definition 5.1. Then, for some universal constant C > 0,

$$\mathbb{P}\left(\mathcal{E}(d,n)\right) \ge 1 - C\exp(-d/C).$$

Proof. It is enough to prove the bound on the probability for each of the events in Items 1-6 of Definition 5.1 separately. For Items 1-5 the bound follows from Lemma 2.1, Lemma 2.4, Proposition 4.2 and Theorem 3, where for Item 5 we used the fact that for t > 1, $\|\nabla^4 H(t\boldsymbol{x})\|_{\text{op}} \leq t^{2p_{\text{max}}} \|\nabla^4 H(\boldsymbol{x})\|_{\text{op}}$, since $\nabla^k F_i(t\boldsymbol{x}) = t^{p_i-k} \nabla^k F_i(\boldsymbol{x})$. The bound for Item 6 is trivial.

Definition 5.3 (Good event for $p_{\max} > d^2$). Suppose that $p_{\max} > d^2$ and n = d - 1. Denote by $\mathcal{E} = \mathcal{E}(d)$ the event that:

- 1. $\operatorname{Lip}^{(i)}(F) \leq C_0 p_{\max}^{1+i} \sqrt{d \log(p_{\max})}$, for i = 0, 1. 2. $\max_{\boldsymbol{x} \in \mathsf{B}^d(1)} \| \boldsymbol{D} F(\boldsymbol{x}) \|_{\operatorname{op}} \leq C_1 p_{\max} \sqrt{d \log(p_{\max})}$.
- 3. There exists $\boldsymbol{\alpha} \in \mathbb{S}^{d-1}$ such that $\boldsymbol{F}(\boldsymbol{\alpha}) = 0$ and $\sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{\alpha})\boldsymbol{U}_{\boldsymbol{\alpha}}) \geq \eta/d^{7/4}$.

Lemma 5.4. Assume the setting of Definition 5.3. Then, for some universal constant C > 0,

$$\mathbb{P}\left(\mathcal{E}(d)\right) \ge 1 - Cd^{-1/2}$$

Proof. Here too it will be enough to prove the bound for each item separately. Items 1 and 2 follow from Lemma 2.4 and Lemma 2.1. Item 3 follows from Proposition 4.1

6 Sufficient conditions for approximate solutions

Throughout the section, we consider the Newton iteration introduced in Section 1.1, and denoted formally by

$$\boldsymbol{x}^{0} = \boldsymbol{x}, \quad \boldsymbol{x}^{k+1} = \Phi_{\text{NM}}(\boldsymbol{x}^{k}),$$
(118)

for a general initialization $\boldsymbol{x} \in \mathbb{S}^{d-1}$. More explicitly, we have $\Phi_{\text{NM}}(\boldsymbol{x}^k) = (\boldsymbol{x}^k + \boldsymbol{v}^k)/\|\boldsymbol{x}^k + \boldsymbol{v}^k\|$, where \boldsymbol{v}^k solves

$$\boldsymbol{F}(\boldsymbol{x}^{k}) + \boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k})\boldsymbol{v}^{k} = 0 \quad \text{subject to} \quad \boldsymbol{v}^{k} \perp \boldsymbol{x}^{k}, \, \boldsymbol{v}^{k} \perp \ker(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k})) \,. \tag{119}$$

In this section we will always work on the event that $\sigma_{\min}(DF(x^k)|_{\mathsf{T}_x}) > 0$, so that a solution to the above indeed exits. Assuming that there exists at least one non-zero solution to F(x) = 0, we will denote by

$$oldsymbol{lpha}^k := rgmin\left\{ \|oldsymbol{x}^k - oldsymbol{lpha}\|:\,oldsymbol{lpha} \in \mathbb{S}^{d-1}, oldsymbol{F}(oldsymbol{lpha}) = 0
ight\}$$

the closest solution to x^k (if more than one solution with minimal distance exists, choose one arbitrarily). We will prove sufficient conditions for a point $x \in \mathbb{S}^{d-1}$ to be an approximate solution. We shall treat the two cases $p_{\max} \leq d^2$ and $p_{\max} > d^2$ separately. In both we will rely on the following.

Proposition 6.1 (Quadratic convergence). Assume that $F(\mathbf{x}) = 0$ has at least one non-zero solution. Let $L := \operatorname{Lip}(\mathbf{DF}; \mathbb{S}^{d-1}) \vee \operatorname{Lip}_{\perp}(\mathbf{DF}; \mathbb{S}^{d-1})$ as defined in (40) and (42), $M := \sup_{\mathbf{x} \in \mathbb{S}^{d-1}} \|\mathbf{DF}(\mathbf{x})\mathbf{x}\|$ and $T := \inf_{k \ge 0} \sigma_{\min}(\mathbf{DF}(\mathbf{\alpha}^k)|_{\mathsf{T}_{\mathbf{\alpha}^k}})$. Assume that T > 0 and set B := 3 + (8L + 4M)/T. If the initial point $\mathbf{x}^0 \in \mathbb{S}^{d-1}$ satisfies $\|\mathbf{x}^0 - \mathbf{\alpha}^0\| \le 1/B$, then

$$\forall k \ge 0: \quad \|\boldsymbol{x}^{k+1} - \boldsymbol{\alpha}^{k+1}\| \le \|\boldsymbol{x}^{k+1} - \boldsymbol{\alpha}^{k}\| \le B \|\boldsymbol{x}^{k} - \boldsymbol{\alpha}^{k}\|^{2}.$$
(120)

Proof. The first inequality in (120) follows by definition. We claim that to prove the proposition, it will be enough to show that

$$\forall k \ge 0, \quad \|\boldsymbol{x}^k + \boldsymbol{v}^k - \boldsymbol{\alpha}^k\| \le B \|\boldsymbol{x}^k - \boldsymbol{\alpha}^k\|^2.$$
(121)

Indeed, this will imply that $\|\boldsymbol{x}^k + \boldsymbol{v}^k - \boldsymbol{\alpha}^k\| \le \|\boldsymbol{\alpha}^k\| = 1$ and $\langle \boldsymbol{\alpha}^k, \boldsymbol{x}^k + \boldsymbol{v}^k \rangle \ge 0$, from which one can easily verify that

$$\|oldsymbol{x}^{k+1}-oldsymbol{lpha}^k\|\leq \|oldsymbol{x}^k+oldsymbol{v}^k-oldsymbol{lpha}^k\|$$
 .

To prove (121) by induction, we let $k \ge 0$ be an arbitrary integer number. Assuming that (121) holds for any $0 \le i \le k - 1$, we will show that it holds for k. Note that by assumption, $\|\boldsymbol{x}^0 - \boldsymbol{\alpha}^0\| \le 1/B$. Since we assume (121) for $0 \le i \le k - 1$, we have from the above that for $0 \le i \le k$, $\|\boldsymbol{x}^i - \boldsymbol{\alpha}^i\|$ is a decreasing sequence and therefore also $\|\boldsymbol{x}^k - \boldsymbol{\alpha}^k\| \le 1/B$.

Denote $V(\boldsymbol{x}) := \ker(\boldsymbol{DF}(\boldsymbol{x})|_{\mathsf{T}_{\boldsymbol{x}}}) \subset \mathsf{T}_{\boldsymbol{x}}$ and let $V(\boldsymbol{x})^{\perp} := \{\boldsymbol{u} \in \mathsf{T}_{\boldsymbol{x}}^{\top} : \boldsymbol{u} \perp V(\boldsymbol{x})\}$ be its orthogonal complement in $\mathsf{T}_{\boldsymbol{x}}$. Denote by $P_{V(\boldsymbol{x}^{k})}, P_{V(\boldsymbol{x}^{k})}^{\perp}$ and $P_{\boldsymbol{x}^{k}}$ the orthogonal projection matrices onto $V(\boldsymbol{x}^{k}), V(\boldsymbol{x}^{k})^{\perp}$ and \boldsymbol{x}^{k} . We claim that

$$\begin{split} \sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x}^{k})\big|_{\mathsf{T}_{\boldsymbol{x}^{k}}}) &= \sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x}^{k})\boldsymbol{U}_{\boldsymbol{x}^{k}}) = \sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x}^{k})P_{V(\boldsymbol{x}^{k})}^{\perp}),\\ \sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha}^{k})\big|_{\mathsf{T}_{\boldsymbol{\alpha}^{k}}}) &= \sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha}^{k})\boldsymbol{U}_{\boldsymbol{\alpha}^{k}}) = \sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha}^{k})). \end{split}$$

The first equality in both lines follows by definition. From homogeneity of $F_i(\boldsymbol{x})$ and since $\boldsymbol{F}(\boldsymbol{\alpha}^k) = 0$, the rows of $\boldsymbol{DF}(\boldsymbol{\alpha}^k)$ are orthogonal to $\boldsymbol{\alpha}^k$. The second equality in the second line follows from this. The second equality in the first line holds whenever $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x}^k)\boldsymbol{U}_{\boldsymbol{x}^k}) > 0$. Indeed, using $\|\boldsymbol{x}^k - \boldsymbol{\alpha}^k\| \leq \frac{1}{B}$, $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x}^k)\boldsymbol{U}_{\boldsymbol{x}^k}) \geq T - L\|\boldsymbol{x}^k - \boldsymbol{\alpha}^k\| \geq T/2$. Note that by assumption, $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha}^k)) \geq T$.

Hence, to prove (121), using that $P_{V(\boldsymbol{x}^k)} + P_{V(\boldsymbol{x}^k)}^{\perp} + P_{\boldsymbol{x}^k} = \boldsymbol{I}$ it will be sufficient to show that

$$\|P_{V(\boldsymbol{x}^{k})}^{\perp}(\boldsymbol{x}^{k}+\boldsymbol{v}^{k}-\boldsymbol{\alpha}^{k})\| \leq \frac{2(L+M)}{\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x}^{k})P_{V(\boldsymbol{x}^{k})}^{\perp})} \|\boldsymbol{x}^{k}-\boldsymbol{\alpha}^{k}\|^{2}, \qquad (122)$$

$$\|P_{V(\boldsymbol{x}^{k})}(\boldsymbol{x}^{k}+\boldsymbol{v}^{k}-\boldsymbol{\alpha}^{k})\| \leq \left(2+\frac{4L}{\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha}^{k}))}\right)\|\boldsymbol{x}^{k}-\boldsymbol{\alpha}^{k}\|^{2}, \qquad (123)$$

$$\|P_{\boldsymbol{x}^{k}}(\boldsymbol{x}^{k}+\boldsymbol{v}^{k}-\boldsymbol{\alpha}^{k})\| \leq \frac{1}{2}\|\boldsymbol{x}^{k}-\boldsymbol{\alpha}^{k}\|^{2}.$$
(124)

Denote by $d(\boldsymbol{x}, \boldsymbol{y}) = \arccos(\langle \boldsymbol{x}, \boldsymbol{y} \rangle)$ be the geodesic distance on \mathbb{S}^{d-1} and let $t_k := d(\boldsymbol{\alpha}^k, \boldsymbol{x}^k)$. For $t \in [0, t_k]$, let $\boldsymbol{x}^k(t) \in \mathbb{S}^{d-1}$ be a geodesic from \boldsymbol{x}^k to $\boldsymbol{\alpha}^k$ with constant speed 1 (so that, in particular, $\boldsymbol{x}^k(0) = \boldsymbol{x}^k$ and $\boldsymbol{x}^k(t_k) = \boldsymbol{\alpha}^k$). Using (119) we write

$$0 = \boldsymbol{F}(\boldsymbol{\alpha}^{k}) = \boldsymbol{F}(\boldsymbol{x}^{k}) + \int_{0}^{t_{k}} \frac{\mathrm{d}\boldsymbol{F}(\boldsymbol{x}^{k}(t))}{\mathrm{d}t} \mathrm{d}t$$

$$= -\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k})\boldsymbol{v}^{k} + \int_{0}^{t_{k}} \boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k}(t))\dot{\boldsymbol{x}}^{k}(t)\mathrm{d}t$$

$$= -\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k})P_{V(\boldsymbol{x}^{k})}^{\perp}\boldsymbol{v}^{k} + \boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k})P_{V(\boldsymbol{x}^{k})}^{\perp}(\boldsymbol{\alpha}^{k} - \boldsymbol{x}^{k})$$

$$+ \underbrace{\int_{0}^{t_{k}} \left[\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k}(t)) - \boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k})\right]\dot{\boldsymbol{x}}^{k}(t)\mathrm{d}t}_{=:\boldsymbol{R}_{1}}$$

$$+ \underbrace{\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k})P_{V(\boldsymbol{x}^{k})}(\boldsymbol{\alpha}^{k} - \boldsymbol{x}^{k})}_{=:\boldsymbol{R}_{2}} + \underbrace{\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k})P_{\boldsymbol{x}^{k}}(\boldsymbol{\alpha}^{k} - \boldsymbol{x}^{k})}_{=:\boldsymbol{R}_{3}}, \qquad (125)$$

where integrals of vector valued functions are carried out elementwise.

By the definition of $V(\boldsymbol{x})$, $\boldsymbol{DF}(\boldsymbol{x}^k)P_{V(\boldsymbol{x}^k)}$ is the zero matrix and $\boldsymbol{R}_2 = 0$. Moreover,

$$\|\boldsymbol{R}_1\| \leq t_k \sup_{t \leq t_k} \|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^k(t)) - \boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^k)\|_{\text{op}} \leq d(\boldsymbol{\alpha}^k, \boldsymbol{x}^k) \cdot L \|\boldsymbol{x}^k - \boldsymbol{\alpha}^k\| \leq 2L \|\boldsymbol{x}^k - \boldsymbol{\alpha}^k\|^2$$

and

$$\|\boldsymbol{R}_3\| \leq \|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^k)\boldsymbol{x}^k\| \cdot |\langle \boldsymbol{x}^k, \boldsymbol{x}^k - \boldsymbol{\alpha}^k \rangle| = \|\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^k)\boldsymbol{x}^k\| \cdot \|\boldsymbol{x}^k - \boldsymbol{\alpha}^k\|^2/2 \leq M \|\boldsymbol{x}^k - \boldsymbol{\alpha}^k\|^2.$$

Continuing from (125) we have that

$$0 = oldsymbol{D}oldsymbol{F}(oldsymbol{x}^k) P_{V(oldsymbol{x}^k)}^{\perp}(oldsymbol{lpha}^k - oldsymbol{x}^k - oldsymbol{v}^k) + oldsymbol{R}\,,$$

where $\mathbf{R} := \mathbf{R}_1 + \mathbf{R}_2 + \mathbf{R}_3$ and $\|\mathbf{R}\| \le 2(L+M) \|\mathbf{x}^k - \mathbf{\alpha}^k\|^2$, from which (122) follows.

Next, we prove (123). Since we assume that T > 0, note that by the implicit function theorem, for any $\boldsymbol{u} \in V(\boldsymbol{\alpha}^k)$ there exist small $\varepsilon > 0$ and a function $\boldsymbol{\varphi} : (-\varepsilon, \varepsilon) \to \mathbb{R}^d$ such that $\|\boldsymbol{\varphi}(t)\| = O(t^2)$ and, for any $t \in (-\varepsilon, \varepsilon)$,

$$\boldsymbol{\gamma}(t) := \frac{\boldsymbol{\alpha}^k + t\boldsymbol{u} + \boldsymbol{\varphi}(t)}{\|\boldsymbol{\alpha}^k + t\boldsymbol{u} + \boldsymbol{\varphi}(t)\|} \in \mathbb{S}^{d-1}$$

is a solution (i.e., $F(\gamma(t)) = 0$). If there existed vector $\boldsymbol{u} \in V(\boldsymbol{\alpha}^k)$ such that $\langle \dot{\boldsymbol{x}}^k(t_k), \boldsymbol{u} \rangle < 0$, we would have $\|\boldsymbol{\gamma}(t) - \boldsymbol{x}^k\| < \|\boldsymbol{\alpha}^k - \boldsymbol{x}^k\|$ for sufficiently small t > 0. We conclude that no such vector exists, i.e., that $\dot{\boldsymbol{x}}^k(t_k) \perp V(\boldsymbol{\alpha}^k)$ or, equivalently, $P_{V(\boldsymbol{\alpha}^k)}\dot{\boldsymbol{x}}^k(t_k) = 0$. From homogeneity of $F_i(\boldsymbol{x})$, at the solution $\boldsymbol{\alpha}^k$ we have that $\ker(\boldsymbol{DF}(\boldsymbol{\alpha}^k)) = V(\boldsymbol{\alpha}^k) \oplus \operatorname{span}\{\boldsymbol{\alpha}^k\}$. Since $\dot{\boldsymbol{x}}^k(t_k) \perp \boldsymbol{\alpha}^k$, we also have that $P_{\ker(\boldsymbol{DF}(\boldsymbol{\alpha}^k))}\dot{\boldsymbol{x}}^k(t_k) = 0$.

Using that $\boldsymbol{v}^k \perp V(\boldsymbol{x}^k)$, we write

$$P_{V(\boldsymbol{x}^{k})}(\boldsymbol{x}^{k} + \boldsymbol{v}^{k} - \boldsymbol{\alpha}^{k}) = P_{V(\boldsymbol{x}^{k})}(\boldsymbol{x}^{k} - \boldsymbol{\alpha}^{k}) = -\int_{0}^{t_{k}} P_{V(\boldsymbol{x}^{k})} \dot{\boldsymbol{x}}^{k}(t) dt$$
$$= -\int_{0}^{t_{k}} P_{V(\boldsymbol{x}^{k})} (\dot{\boldsymbol{x}}^{k}(t) - \dot{\boldsymbol{x}}^{k}(t_{k})) dt - t_{k} P_{V(\boldsymbol{x}^{k})} \dot{\boldsymbol{x}}^{k}(t_{k})$$

Noting that $\|\dot{\boldsymbol{x}}^k(t) - \dot{\boldsymbol{x}}^k(t_k)\|$ is maximal over $[0, t_k]$ for t = 0 and recalling that $\|\dot{\boldsymbol{x}}^k(t_k)\| = 1$,

$$\|P_{V(\boldsymbol{x}^{k})}(\boldsymbol{x}^{k}+\boldsymbol{v}^{k}-\boldsymbol{\alpha}^{k})\| \leq t_{k}\|\dot{\boldsymbol{x}}^{k}(0)-\dot{\boldsymbol{x}}^{k}(t_{k})\|+t_{k}\|P_{V(\boldsymbol{x}^{k})}(\boldsymbol{I}-P_{\ker(\boldsymbol{DF}(\boldsymbol{\alpha}^{k}))})\|_{\text{op}}.$$

Since $t_k \leq 2 \| \boldsymbol{x}^k - \boldsymbol{\alpha}^k \|$, the first term is bounded by

$$t_k \| \dot{\boldsymbol{x}}^k(0) - \dot{\boldsymbol{x}}^k(t_k) \| = t_k \| \boldsymbol{x}^k(0) - \boldsymbol{x}^k(t_k) \| \le 2 \| \boldsymbol{x}^k - \boldsymbol{\alpha}^k \|^2.$$

For any $\boldsymbol{x} \in \mathbb{S}^{d-1}$, $V(\boldsymbol{x}) \subset \ker(\boldsymbol{DF}(\boldsymbol{x}))$. Thus,

$$\|P_{V(\boldsymbol{x}^{k})}(\boldsymbol{I} - P_{\ker(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{\alpha}^{k}))})\|_{\text{op}} \leq \|P_{\ker(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^{k}))}(\boldsymbol{I} - P_{\ker(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{\alpha}^{k}))})\|_{\text{op}}.$$
(126)

Applying Theorem 5 (specifically, Eq. (145)) with $A_0 = DF(x^k)$, $A_1 = DF(\alpha^k)$ and $k(a) = \operatorname{rank}(A_a)$ so that $\Delta = \sigma_{\min}(DF(\alpha^k)) > 0$ (by Eq. (104)), we obtain that the right-hand side of (126) is bounded as follows:

$$\begin{split} \|P_{\ker(\boldsymbol{DF}(\boldsymbol{x}^{k}))}(\boldsymbol{I} - P_{\ker(\boldsymbol{DF}(\boldsymbol{\alpha}^{k}))})\|_{\text{op}} &\leq 2 \frac{\|\boldsymbol{DF}(\boldsymbol{x}^{k}) - \boldsymbol{DF}(\boldsymbol{\alpha}^{k})\|}{\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha}^{k}))} \\ &\leq \frac{2L\|\boldsymbol{x}^{k} - \boldsymbol{\alpha}^{k}\|}{\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha}^{k}))} \,. \end{split}$$

Combining the above and using again that $t_k \leq 2 \| \boldsymbol{x}^k - \boldsymbol{\alpha}^k \|$, we obtain (123).

Finally, (124) follows since

$$\|P_{\boldsymbol{x}^k}(\boldsymbol{x}^k + \boldsymbol{v}^k - \boldsymbol{\alpha}^k)\| = \|P_{\boldsymbol{x}^k}(\boldsymbol{x}^k - \boldsymbol{\alpha}^k)\| = 1 - \langle \boldsymbol{x}^k, \boldsymbol{\alpha}^k \rangle = \frac{1}{2} \|\boldsymbol{x}^k - \boldsymbol{\alpha}^k\|^2.$$

Corollary 6.2. Assume that F(x) = 0 has at least one non-zero solution. Define L and M as in Proposition 6.1 and

$$T := \sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x}^0)\big|_{\mathsf{T}_{\boldsymbol{x}^0}}) - 5L\|\boldsymbol{x}^0 - \boldsymbol{\alpha}^0\|.$$

Assume that T > 0 and set B := 3 + (8L + 4M)/T. If the initial point $\mathbf{x}^0 \in \mathbb{S}^{d-1}$ satisfies $\|\mathbf{x}^0 - \mathbf{\alpha}^0\| \le 1/4B$, then

$$\forall k \ge 0: \quad \|\boldsymbol{x}^{k+1} - \boldsymbol{\alpha}^{k+1}\| \le \|\boldsymbol{x}^{k+1} - \boldsymbol{\alpha}^{k}\| \le B \|\boldsymbol{x}^{k} - \boldsymbol{\alpha}^{k}\|^{2}.$$
(127)

Proof. As in Proposition 6.1, we will prove (127) by induction on k. Here, however, we do not assume the uniform bound on the minimal singular value at α^k and $k \ge 0$.

We start with k = 0. By definition, $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha}^0)|_{\mathsf{T}_{\boldsymbol{\alpha}^0}}) > T$. Observe that the proof of (127) in Proposition 6.1 for k = 0 only relied the latter bound on the singular value at $\boldsymbol{\alpha}^0$ but did not use the same bound for $\boldsymbol{\alpha}^i$ with i > 0. Hence, by exactly the same argument as in Proposition 6.1, (127) follows for k = 0.

Now, assume that (127) holds for any $0 \le i \le k - 1$. Then, by induction, for any $1 \le i \le k$,

$$\|\boldsymbol{x}^{i} - \boldsymbol{\alpha}^{i}\| \le \|\boldsymbol{x}^{i} - \boldsymbol{\alpha}^{i-1}\| \le \|\boldsymbol{x}^{0} - \boldsymbol{\alpha}^{0}\| \left(B\|\boldsymbol{x}^{0} - \boldsymbol{\alpha}^{0}\|\right)^{2^{i}-1} \le \|\boldsymbol{x}^{0} - \boldsymbol{\alpha}^{0}\| \left(\frac{1}{4}\right)^{2^{i}-1}$$
(128)

and therefore

$$\|\boldsymbol{x}^{i} - \boldsymbol{x}^{i-1}\| \le \|\boldsymbol{x}^{i} - \boldsymbol{\alpha}^{i-1}\| + \|\boldsymbol{x}^{i-1} - \boldsymbol{\alpha}^{i-1}\| \le \frac{5}{4}\|\boldsymbol{x}^{0} - \boldsymbol{\alpha}^{0}\| \left(\frac{1}{4}\right)^{2^{i-1}-1}.$$
(129)

Thus,

$$\| oldsymbol{lpha}^i - oldsymbol{x}^0 \| \le \| oldsymbol{lpha}^i - oldsymbol{x}^i \| + \| oldsymbol{x}^i - oldsymbol{x}^0 \| + \| oldsymbol{x}^0 - oldsymbol{lpha}^0 \|$$

$$\leq \|\boldsymbol{x}^{0} - \boldsymbol{\alpha}^{0}\| \left[\left(\frac{1}{4}\right)^{2^{i-1}} + \frac{5}{4} \sum_{j=1}^{i} \left(\frac{1}{4}\right)^{2^{j-1}-1} + 1 \right] \leq 5 \|\boldsymbol{x}^{0} - \boldsymbol{\alpha}^{0}\|.$$

And we conclude that

$$\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha}^{i})\big|_{\mathsf{T}_{\boldsymbol{\alpha}^{i}}}) \ge \sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x}^{0})\big|_{\mathsf{T}_{\boldsymbol{x}^{0}}}) - 5L\|\boldsymbol{x}^{0} - \boldsymbol{\alpha}^{0}\| = T.$$
(130)

The proof of (127) for k follows by the same induction argument we used in the proof of Proposition 6.1, which relied on the bound $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha}^i)|_{\mathsf{T}_{\boldsymbol{\alpha}^i}}) \geq T$ with $i \leq k$ only and not on the same bound for i > k.

6.1 The case of moderate $p_{\text{max}} \leq d^2$

We prove the following sufficient condition for an approximate solution.

Proposition 6.3. Assume the same setting and notation as in Definition 5.1 and that points 1, 2 and 3 in the definition hold. Let τ be defined as in (103) and let

$$B := \frac{c p_{\max}^2 \sqrt{d \log p_{\max}}}{\tau}, \qquad c := 15(C_0 \vee C_1 \vee 1).$$
(131)

If $\|\boldsymbol{x} - \boldsymbol{\alpha}\| \leq \frac{1}{4B}$ and $\boldsymbol{F}(\boldsymbol{\alpha}) = \boldsymbol{0}$ for some $\boldsymbol{x}, \boldsymbol{\alpha} \in \mathbb{S}^{d-1}$, then \boldsymbol{x} is an approximate solution.

Proof. Let $\mathbf{x}^0 = \mathbf{x}$ and define $\mathbf{x}^{i+1} = \Phi_{\text{NM}}(\mathbf{x}^i)$ by Newton's method as in Eqs. (118), (119). Let L, M and T be as defined in Proposition 6.1. By Points 2 and 3 of Definition 5.1, $L \leq C_0 p_{\max}^2 \sqrt{d \log(p_{\max})}$ and $T \geq \tau$. Note that for any $\mathbf{x} \in \mathbb{S}^{d-1}$, the *i*-th element of $DF(\mathbf{x})\mathbf{x}$ is the radial derivative of F_i at \mathbf{x} which is equal to $p_i F_i(\mathbf{x})$, since F_i is homogeneous. Thus, by Point 1 of Definition 5.1, $M \leq C_1 p_{\max} \sqrt{d \log(p_{\max})}$. Hence, for B as defined in (131), $\frac{8L+4M}{T} + 3 \leq B$. By Proposition 6.1, it follows that (120) holds.

Therefore, as in (128) and (129),

$$\|m{x}^k - m{lpha}^k\| \le \|m{x}^0 - m{lpha}^0\| \Big(rac{1}{4}\Big)^{2^k-1} ext{ and } \|m{x}^k - m{x}^{k-1}\| \le rac{5}{4} \|m{x}^0 - m{lpha}^0\| \Big(rac{1}{4}\Big)^{2^{k-1}-1}.$$

Thus, the limit $\bar{\boldsymbol{\alpha}} = \lim_{k \to \infty} \boldsymbol{x}^k \lim_{k \to \infty} \boldsymbol{\alpha}^k$ is well-defined, $\boldsymbol{F}(\bar{\boldsymbol{\alpha}}) = 0$ and, for $k \ge 1$,

$$\|\boldsymbol{x}^{k} - \bar{\boldsymbol{\alpha}}\| \leq \sum_{j=k}^{\infty} \|\boldsymbol{x}^{j+1} - \boldsymbol{x}^{j}\| \leq \frac{5}{4} \|\boldsymbol{x}^{0} - \boldsymbol{\alpha}^{0}\| \sum_{j=k}^{\infty} 4^{1-2^{j}} \leq \left(\frac{5}{4}\right)^{2} \|\boldsymbol{x}^{0} - \boldsymbol{\alpha}^{0}\| 4^{1-2^{k}} \leq \|\boldsymbol{x}^{0} - \bar{\boldsymbol{\alpha}}\| 2^{1-2^{k}}.$$

6.2 The case of large $p_{\text{max}} > d^2$

Corollary 6.4. Let $n \leq d-1$ be arbitrary and assume that the bounds of Points 1 and 2 in Definition 5.3 occur for some constants C_0 and C_1 . Then, there exists $\delta = \delta(C_0, C_1, \bar{\eta})$ such that any $\mathbf{x} \in \mathbb{S}^{d-1}$ that satisfies

$$\|\boldsymbol{F}(\boldsymbol{x})\| \leq \frac{\delta}{d^4 p_{\max}^2 \sqrt{\log(p_{\max})}} \quad and \quad \sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}}) \geq \frac{\bar{\eta}}{2d^{7/4}}$$
(132)

is an approximate solution.

Proof. Assume that $\boldsymbol{x} \in \mathbb{S}^{d-1}$ satisfies (132), for some $0 < \delta$ which will be assumed to be small enough whenever needed and may depend on C_0 , C_1 and $\bar{\eta}$. First, we will show that there exists a solution close to \boldsymbol{x} by appealing to [MS23, Lemma E.1] about gradient flow.

Since, assuming δ is small,

$$\begin{split} \operatorname{Lip}_{\perp}(\boldsymbol{D}\boldsymbol{F};\mathbb{S}^{d-1}) \|\boldsymbol{F}(\boldsymbol{x})\| &\leq C_0 p_{\max}^2 \sqrt{d\log(p_{\max})} \cdot \frac{\delta}{d^4 p_{\max}^2 \sqrt{\log(p_{\max})}} \\ &\leq \frac{\bar{\eta}^2}{16d^{7/2}} \leq \frac{1}{4} \big(\sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}})\big)^2 \,, \end{split}$$

by [MS23, Lemma E.1], gradient flow $\boldsymbol{x}(t)$ on \mathbb{S}^{d-1} started from $\boldsymbol{x}(0) = \boldsymbol{x}$ converges to a solution $\boldsymbol{\alpha} := \lim_{t \to \infty} \boldsymbol{x}(t)$. Namely, $\boldsymbol{F}(\boldsymbol{\alpha}) = \boldsymbol{0}$. By Eq. (162) of [MS23],

$$\|\boldsymbol{x} - \boldsymbol{\alpha}\| \le 2 \frac{\|\boldsymbol{F}(\boldsymbol{x})\|}{\sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}})} \le \frac{4\delta}{\bar{\eta}d^{9/4}p_{\max}^2\sqrt{\log(p_{\max})}}.$$

For Newton's method started from $\mathbf{x}^0 = \mathbf{x}$, in the notation of Corollary 6.2, $L = \text{Lip}^{(1)}(\mathbf{F}) \leq C_0 p_{\text{max}}^2 \sqrt{d \log(p_{\text{max}})}$, $M \leq C_1 p_{\text{max}} \sqrt{d \log(p_{\text{max}})}$ (see the proof of Proposition 6.3 for the last inequality),

$$T \ge \frac{\bar{\eta}}{2d^{7/4}} - 5C_0 p_{\max}^2 \sqrt{d\log(p_{\max})} \cdot \frac{4\delta}{\bar{\eta}d^{9/4} p_{\max}^2 \sqrt{\log(p_{\max})}} \ge \frac{\bar{\eta} - 40C_0 \delta/\bar{\eta}}{2d^{7/4}}$$

Therefore, for small δ , we have T > 0 and

$$B \le 8(C_0 + C_1)p_{\max}^2 \sqrt{d\log(p_{\max})} \cdot \frac{d^{7/4}}{\bar{\eta}/4} + 3$$

and

$$\|\boldsymbol{x}^0 - \boldsymbol{\alpha}\| \le \frac{1}{4B}$$

Hence, (127) follows, by Corollary 6.2. The fact that x is an approximate solution follows by the same argument in the last paragraph of the proof of Proposition 6.3.

7 Sub-routines and their complexity

In this section we define and analyze sub-routines that are used by the main algorithms. The first is for finding vectors v_i as in Algorithm 1. Its pseudo-code is given below. Below e_i denotes the standard basis of \mathbb{R}^d and C_1 and c_0 are constants as in Lemma 2.1 and Algorithm 1. (The value of c_0 will be determined in the proof of Theorem 1.) The absolute constant $c = c(C_1, c_0) > 0$ in the pseudo-code will be determined in the proof of Lemma 7.1.

Algorithm 3: Find a good direction

Input: The coefficients \boldsymbol{a} , a point $\boldsymbol{x} \in \mathbb{S}^{d-1}$ Output: A unit vector $\boldsymbol{v} \in \mathbb{S}^{d-1}$ orthogonal to \boldsymbol{x} $\mu = 9C_1 dp_{\max}^2 \log(p_{\max});$ $\boldsymbol{A} = (\boldsymbol{I} - \boldsymbol{x} \boldsymbol{x}^{\mathsf{T}})(\mu \boldsymbol{I} - \nabla^2 H(\boldsymbol{x}))(\boldsymbol{I} - \boldsymbol{x} \boldsymbol{x}^{\mathsf{T}});$ k = 1;while $k < e^{c(d+\log p_{\max})}$ do $| \boldsymbol{A} = \boldsymbol{A}^2, \ k = 2k;$ end $i_0 = \operatorname{argmin}_{i \leq d} \langle \nabla^2 H(\boldsymbol{x}), (\boldsymbol{A} \boldsymbol{e}_i / \| \boldsymbol{A} \boldsymbol{e}_i \|)^{\otimes 2} \rangle;$ return $\boldsymbol{v} = \boldsymbol{A} \boldsymbol{e}_{i_0} / \| \boldsymbol{A} \boldsymbol{e}_{i_0} \|$

In our running time analysis below, we compare the number of operation with the input size N (the number of coefficients to specify the function F). Recall that

$$N = \sum_{i=1}^{n} \binom{d+p_{i}-1}{p_{i}}.$$
(133)

Lemma 7.1. The running time of Algorithm 3 is $O(Nd^3(1 + p_{\max}/d))$. On the event that Points 4 and 5 of Definition 5.1 hold, for any $\boldsymbol{x} \in \mathbb{S}^{d-1}$ such that $H(\boldsymbol{x}) \geq p_{\max}^{-c_0d}$, the algorithm outputs a vector \boldsymbol{v} such that $\boldsymbol{v} \perp \boldsymbol{x}$, $\|\boldsymbol{v}\| = 1$ and

$$\langle \nabla^2 H(\boldsymbol{x}_i), \boldsymbol{v}^{\otimes 2} \rangle \leq \frac{1}{2} \min_{\boldsymbol{u} \perp \boldsymbol{x}, \|\boldsymbol{u}\|=1} \langle \nabla^2 H(\boldsymbol{x}), \boldsymbol{u}^{\otimes 2} \rangle.$$
 (134)

Proof. Let v_1, \ldots, v_{d-1} and $\lambda_1 \leq \cdots \leq \lambda_{d-1}$ be the eigenvectors and eigenvalues of $\nabla^2 H(\boldsymbol{x})|_{\mathsf{T}_{\boldsymbol{x}}}$. On the event as in the statement of the lemma, $|\lambda_i| \leq \mu$ and $\lambda_1 \leq -\frac{1}{10}\sqrt{(d-n)H(\boldsymbol{x})}$. We have that

$$oldsymbol{A}^koldsymbol{e}_i = \sum_{j\leq d-1} (\mu-\lambda_j)^k \langle oldsymbol{e}_i,oldsymbol{v}_j
angle oldsymbol{v}_j$$

and

$$\begin{split} \langle \nabla^2 H(\boldsymbol{x}), (\boldsymbol{A}^k \boldsymbol{e}_i / \| \boldsymbol{A}^k \boldsymbol{e}_i \|)^{\otimes 2} \rangle &= \frac{\sum_{j \le d-1} (\mu - \lambda_j)^{2k} \langle \boldsymbol{e}_i, \boldsymbol{v}_j \rangle^2 \lambda_j}{\sum_{j \le d-1} (\mu - \lambda_j)^{2k} \langle \boldsymbol{e}_i, \boldsymbol{v}_j \rangle^2} \\ &\le \frac{\sum_{j \in I} \left(\frac{\mu - \lambda_j}{\mu - \lambda_1}\right)^{2k} \langle \boldsymbol{e}_i, \boldsymbol{v}_j \rangle^2 \lambda_j + \mu e^{-\frac{k|\lambda_1|}{4\mu}}}{\sum_{j \in I} \left(\frac{\mu - \lambda_j}{\mu - \lambda_1}\right)^{2k} \langle \boldsymbol{e}_i, \boldsymbol{v}_j \rangle^2 + e^{-\frac{k|\lambda_1|}{4\mu}}}, \end{split}$$

provided that the numerator in the second ratio is negative, where I is the set of indices such that $\lambda_j \leq \frac{3}{4}\lambda_1$ and we used that, for $j \notin I$,

$$\frac{\mu - \lambda_j}{\mu - \lambda_1} \le 1 - \frac{|\lambda_1|}{8\mu} \le e^{-\frac{|\lambda_1|}{8\mu}}$$

For some i, $\langle \boldsymbol{e}_i, \boldsymbol{v}_1 \rangle^2 \geq 1/d$. Hence, (134) follows since $|\lambda_1|/(10d) \geq \mu e^{-\frac{k|\lambda_1|}{4\mu}}$ and $1/(10d) \geq e^{-\frac{k|\lambda_1|}{4\mu}}$ if $k < e^{c(d+\log p_{\max})}$ and $c = c(C_1, c_0)$ is sufficiently large.

By induction on p_i , one can check that computing all products of the form $x_1^{k_1} \cdots x_d^{k_d}$ that appear in (1) requires $c_1 \binom{d+p_1}{p_i}$ operations for some constant c_1 . Hence, calculating $F_i(\boldsymbol{x})$ from the coefficients \boldsymbol{a} requires $c_1N(1 + p_{\max}/d)$ operations at most. Any first or second order derivative of $F_i(\boldsymbol{x})$ can be obtained with the same running time. Therefore, \boldsymbol{A} can be computed in $c'_1Nd^3(1 + p_{\max}/d)$ operations. The number of iterations in the while-loop is bounded by $1+c(d+\log p_{\max})/\log 2$ and each iteration requires $O(d^3)$ operations. The complexity of running the while-loop can therefore be absorbed in the previous term.

The next sub-routines approximate the maximal and minimal singular value of a matrix. Denote by ω_L the time complexity for $L \times L$ matrix multiplication.

Lemma 7.2. There are an absolute constant c > 0 and an algorithm that, given a matrix $\mathbf{A} \in \mathbb{R}^{M \times L}$, computes in time complexity $c(\log \log M \cdot \omega_M + \omega_L))$ a value

$$s_{\max} \in \left[\frac{1}{2}\sigma_{\max}(\mathbf{A}), \sigma_{\max}(\mathbf{A})\right].$$
 (135)

Proof. Define $\mathbf{B} = \mathbf{A}\mathbf{A}^{\mathsf{T}}$ and denote its eigenvalues by $0 \leq \lambda_1 \leq \cdots \leq \lambda_M$. Let $k = 2^s$ with $s = \lceil \log_2 \log M \rceil$. We claim that $s_{\max} := \max_{i \leq N} \|\mathbf{B}^k \mathbf{e}_i\|^{1/k}$ satisfies $\mu \in [\frac{1}{2}\lambda_M, \lambda_M]$. Indeed, denoting by \mathbf{v}_j the eigenvector corresponding to $\lambda_j, \lambda_M^{2k} \geq \|\mathbf{B}^k \mathbf{e}_i\|^2 = \sum_j \lambda_j^{2k} \langle \mathbf{e}_i, \mathbf{v}_j \rangle^2 \geq \lambda_M^{2k} \langle \mathbf{e}_i, \mathbf{v}_M \rangle^2$. The claim thus follows since $\max_i \langle \mathbf{e}_i, \mathbf{v}_M \rangle^2 \geq 1/M$. Exploiting the fact that k is a power of 2 as in Algorithm 3, the time complexity for computing μ is as stated in the lemma. We note that computing the k-th root for k a power of two can be done by iterative computing the square root, which we assume as part of our computational model.

Lemma 7.3. There are an absolute constant c > 0 and an algorithm that, given a symmetric matrix $\mathbf{A} \in \mathbb{R}^{M \times L}$ and $\kappa > 1$, computes in time complexity $c((\log \kappa + \log \log M)\omega_M + \omega_L))$ a value

$$s_{\min} \in \left[\sigma_{\min}(\mathbf{A}), \sigma_{\min}(\mathbf{A}) + \sqrt{2\sigma_{\max}(\mathbf{A})(1 - M^{-1/2\kappa})}\right].$$
 (136)

Proof. Define $\mathbf{B} = \mathbf{A}\mathbf{A}^{\mathsf{T}}$, λ_i and s_{\max} as in the previous proof. Define $\mathbf{D} = 2s_{\max}\mathbf{I} - \mathbf{B}$ and set $\kappa' = 2^{\lceil \log_2 \kappa \rceil}$. Similarly to the previous proof, $S := \max_i \|\mathbf{D}^{\kappa'} \mathbf{e}_i\|^{1/\kappa'} \in [M^{-1/2\kappa'}(2s_{\max} - \lambda_1), 2s_{\max} - \lambda_1]$. Therefore, $s_{\min} := \sqrt{2s_{\max} - S}$ satisfies (136).

The time complexity for computing s_{\max} is $O(\log \log M \cdot \omega_M + \omega_L))$. Exploiting the fact that κ' is a power of 2 as in Algorithm 3, the running time for computing s_{\min} given s_{\max} is $O(\log(\kappa)\omega_M)$.

8 Analysis of Hessian descent: Proof of Theorem 1

We start with an analysis of the Hessian Descent algorithm (Algorithm 1). In the theorem below we will assume that $(\boldsymbol{x}_i, \boldsymbol{y}_i, \boldsymbol{v}_i, \delta_i)_{i\geq 0}$ is an infinite sequence as follows. For i = 0, $\boldsymbol{x}_0 = \boldsymbol{y}_0 = (1, 0, \dots, 0)$. For any $i \geq 0$, \boldsymbol{v}_i is a unit vector such that $\boldsymbol{v}_i \perp \boldsymbol{x}_i$,

$$\langle \nabla^2 H(\boldsymbol{x}_i), \boldsymbol{v}_i^{\otimes 2} \rangle \le -\frac{1}{20} \sqrt{(d-n)H(\boldsymbol{x}_i)},$$
(137)

 $\boldsymbol{y}_{i+1} = \boldsymbol{x}_i \pm \delta_i \boldsymbol{v}_i$, with the sign chosen such that $H(\boldsymbol{y}_{i+1}) \leq \min\{H(\boldsymbol{x}_i + \delta_i \boldsymbol{v}_i), H(\boldsymbol{x}_i - \delta_i \boldsymbol{v}_i)\}$, and $\boldsymbol{x}_{i+1} = \boldsymbol{y}_{i+1} / \|\boldsymbol{y}_{i+1}\|$. δ_i is as defined in Algorithm 1.

Theorem 4. Suppose the sequence $(\boldsymbol{x}_i, \boldsymbol{y}_i, \boldsymbol{v}_i, \delta_i)_{i \geq 0}$ is as above and that the event in Point 5 of Definition 5.1 occurs. For some universal constants c_1, c_2 we have the following. If

$$\sqrt{(d-n)H(\boldsymbol{x}_i)} \ge 30C_1 dp_{\max}^3 \log(p_{\max})$$
(138)

then

$$H(\boldsymbol{x}_{i+1}) - H(\boldsymbol{x}_i) \le -\frac{3}{8}C_1 dp_{\max}^2 \log(p_{\max}).$$
(139)

Otherwise, if (138) does not hold, for any $j \ge i$,

$$H(\boldsymbol{x}_j) \le H(\boldsymbol{x}_i) \cdot \left(1 - \frac{c_1}{p_{\max}^4 \log(p_{\max})} \frac{d-n}{d}\right)^{j-i}$$
(140)

and

$$\left\|\boldsymbol{x}_{i} - \boldsymbol{x}_{j}\right\|_{2} \leq c_{2} H(\boldsymbol{x}_{i})^{\frac{1}{4}} p_{\max}^{2} \sqrt{d \log(p_{\max})} \,.$$

$$(141)$$

Proof. By Taylor's theorem,

$$H(\boldsymbol{y}_{i+1}) - H(\boldsymbol{x}_{i}) \stackrel{(a)}{\leq} \sum_{k=1}^{3} \frac{\delta_{i}^{k}}{k!} \langle \nabla^{k} H(\boldsymbol{x}_{i}), \boldsymbol{v}_{i}^{\otimes k} \rangle + \frac{\delta_{i}^{4}}{4!} 3C_{1} dp_{\max}^{4} \log(p_{\max})$$

$$\stackrel{(b)}{\leq} \frac{\delta_{i}^{2}}{2} \langle \nabla^{2} H(\boldsymbol{x}_{i}), \boldsymbol{v}_{i}^{\otimes 2} \rangle + \frac{\delta_{i}^{4}}{4!} 3C_{1} dp_{\max}^{4} \log(p_{\max})$$

$$\stackrel{(c)}{\leq} -\frac{\delta_{i}^{2}}{40} \sqrt{(d-n)H(\boldsymbol{x}_{i})} + \frac{\delta_{i}^{4}}{4!} 3C_{1} dp_{\max}^{4} \log(p_{\max})$$

$$\stackrel{(d)}{\leq} -\frac{\delta_{i}^{2}}{80} \sqrt{(d-n)H(\boldsymbol{x}_{i})}, \qquad (142)$$

where for (a) we used that $\delta_i \leq \sqrt{1/p_{\text{max}}}$ and Point 5 of Definition 5.1 and C_1 is as there, for (b) we used the fact that the same bound holds whether we take $\mathbf{y}_{i+1} = \mathbf{x}_i + \mathbf{v}_i$ or $\mathbf{y}_{i+1} = \mathbf{x}_i - \mathbf{v}_i$ and we choose the sign that minimizes $H(\mathbf{y}_{i+1})$, for (c) we used (137), and (d) follows from the choice of δ_i as in Algorithm 1.

We have that $\delta_i = \sqrt{1/p_{\text{max}}}$ if and only if (138) holds, in which case, using homogeneity for the first inequality,

$$H(\boldsymbol{x}_{i+1}) - H(\boldsymbol{x}_i) \le H(\boldsymbol{y}_{i+1}) - H(\boldsymbol{x}_i) \le -\frac{3}{8}C_1 dp_{\max}^2 \log(p_{\max}),$$

and (139) follows.

Now, suppose (138) does not hold for some given *i*. Then, using (142) and the definition of δ_i ,

$$H(\boldsymbol{x}_{i+1}) - H(\boldsymbol{x}_i) \le H(\boldsymbol{y}_{i+1}) - H(\boldsymbol{x}_i) \le -\frac{1}{2400C_1} \frac{1}{p_{\max}^4 \log(p_{\max})} \frac{d-n}{d} H(\boldsymbol{x}_i) \le -\frac{1}{2400C_1} \frac{d-n}{d} H(\boldsymbol{x}_i) \le -\frac{1}{240C_1} \frac{d-n}{d} H(\boldsymbol$$

and for $c_1 = 1/2400C_1$,

$$H(\boldsymbol{x}_{i+1}) \leq H(\boldsymbol{x}_i) \left(1 - \frac{c_1}{p_{\max}^4 \log(p_{\max})} \frac{d-n}{d}\right),$$

which completes the proof of (140).

For any j > i,

$$\|m{x}_j - m{x}_i\| \le \sum_{k=i}^{j-1} \|m{x}_{k+1} - m{x}_k\| \le \sum_{k=i}^{j-1} \|m{y}_{k+1} - m{x}_k\| = \sum_{k=i}^{j-1} \delta_k$$

$$= \left(\frac{1}{30C_{1}}\frac{1}{p_{\max}^{4}\log(p_{\max})}\frac{\sqrt{(d-n)}}{d}\right)^{\frac{1}{2}}\sum_{k=i}^{j-1}H(\boldsymbol{x}_{k})^{\frac{1}{4}}$$

$$\leq H(\boldsymbol{x}_{i})^{\frac{1}{4}}\left(\frac{1}{30C_{1}}\frac{1}{p_{\max}^{4}\log(p_{\max})}\frac{\sqrt{(d-n)}}{d}\right)^{\frac{1}{2}}\left(1-\left(1-\frac{c_{1}}{p_{\max}^{4}\log(p_{\max})}\frac{d-n}{d}\right)^{\frac{1}{4}}\right)^{-1}$$

$$\leq H(\boldsymbol{x}_{i})^{\frac{1}{4}}\left(\frac{1}{30C_{1}}\frac{1}{p_{\max}^{4}\log(p_{\max})}\frac{\sqrt{(d-n)}}{d}\right)^{\frac{1}{2}}\left(\frac{1}{4}\frac{c_{1}}{p_{\max}^{4}\log(p_{\max})}\frac{d-n}{d}\right)^{-1},$$

$$\leq (141) \text{ with } c_{2} = \sqrt{8/(15C_{1}c_{1}^{2})}.$$

which proves (141) with $c_2 = \sqrt{8/(15C_1c_1^2)}$.

8.1 Proof of Theorem 1

Consider the Hessian Descent algorithm (Algorithm 1), with $k = C'_0 d^{3/2} p_{\text{max}}^4 \log(p_{\text{max}})^2$ iterations for some absolute $C'_0 > 0$ to be determined below. Recall that to find a vector \boldsymbol{v}_i we use the sub-routine from Section 7. By Lemma 7.1, the time complexity for a single iteration is $O(Nd^3(1 + p_{\text{max}}/d))$ and the total time complexity is

$$\chi \le C_0 N d^{9/2} p_{\max}^4 (1 + p_{\max}/d) \log(p_{\max})^2$$

for appropriate C_0 , which proves (2).

Next, we will assume that the good event $\mathcal{E}(d, n)$ of Definition 5.1 occurs and prove that the output $\boldsymbol{x}^{\text{HD}}$ is an approximate solution (recall Definition 1.1). Indeed, in light of the lower bound on $\mathbb{P}(\mathcal{E}(d, n))$ in Lemma 5.2, this will complete the proof of the theorem.

Suppose that i_0 is the largest index for which x_{i_0} is computed before the algorithm terminates. By Lemma 7.1, on $\mathcal{E}(d, n)$, for all indices $i < i_0$,

$$\langle
abla^2 H(\boldsymbol{x}_i), \boldsymbol{v}_i^{\otimes 2}
angle \leq rac{1}{2} \min_{\boldsymbol{u} \perp \boldsymbol{x}_i, \|\boldsymbol{u}\| = 1} \langle
abla^2 H(\boldsymbol{x}_i), \boldsymbol{u}^{\otimes 2}
angle.$$

From that bound of Point 4 Definition 5.1, v_i therefore also satisfies (137). By the same point in Definition 5.1, for $i \ge i_0$ there exists v_i such that (137) holds. Hence, we may extend the sequence $(x_i, y_i, v_i, \delta_i)_{i \le i_0}$ produced by the Algorithm 1 to an infinite sequence $(x_i, y_i, v_i, \delta_i)_{i \ge 0}$ as in the setting of Theorem 4.

From (139) and the bound $H(x_0) \leq d$ as in Point 6 of Definition 5.1, for some absolute constant c > 0, for any $i \geq cd$ the condition in (138) does not hold.

By (140) and (141), \boldsymbol{x}_i is a Cauchy sequence, and $\boldsymbol{\alpha} := \lim_{i \to \infty} \boldsymbol{x}_i \in \mathbb{S}^{d-1}$ is an exact solution, $\boldsymbol{F}(\boldsymbol{\alpha}) = \boldsymbol{0}$. Moreover, for any $i \geq cd$,

$$H(\boldsymbol{x}_i) \leq d \cdot \left(1 - \frac{c_1}{p_{\max}^4 \log(p_{\max})} \frac{d-n}{d}\right)^{i-cd}.$$

Therefore, if the number of steps k is as above with sufficiently large C'_0 , then of some $i \leq k$ we have that $H(\boldsymbol{x}_i) \leq p_{\max}^{-c_0 d}$, and the algorithm outputs $\boldsymbol{x}^{\text{HD}} = \boldsymbol{x}_i$. By (141),

$$\|\boldsymbol{x}^{\text{HD}} - \boldsymbol{\alpha}\| \le c_2 p_{\max}^{-c_0 d/4} p_{\max}^2 \sqrt{d \log(p_{\max})}$$

If c_0 is sufficiently large, then in the notation of Proposition 6.3 we have that $\|\boldsymbol{x}^{\text{HD}} - \boldsymbol{\alpha}\| \leq 1/4B$ and $\boldsymbol{x}^{\text{HD}}$ is therefore an approximate solution.

9 Analysis of brute-force search: Proof of Theorem 2

Recall the good event $\mathcal{E} = \mathcal{E}(d)$ of Definition 5.3. Below we use the same notation as in the latter definition and Algorithm 2. We assume that the algorithm is run with $\bar{\eta} = \eta/2$ for η as in Definition 5.3 and $\delta = \delta(C_0, C_1, \bar{\eta}) \wedge \eta/2$, where $\delta(C_0, C_1, \bar{\eta})$ is as in Corollary 6.4.

Equipped with the sub-routines of Lemma 7.3, we now make precise the approximation of $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x})|_{\mathsf{T}_{\boldsymbol{x}}})$ in Algorithm 2. Namely, we define $s_{\min}(\boldsymbol{x})$ as the output of the algorithm of Lemma 7.3 with input $\boldsymbol{A} = \boldsymbol{DF}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}}$ and $\kappa = 2\bar{\eta}^{-1}C_1p_{\max}\sqrt{\log p_{\max}}d^{9/4}\log d$, with an arbitrary matrix $\boldsymbol{U}_{\boldsymbol{x}}$ whose columns form an orthonormal basis of $\mathsf{T}_{\boldsymbol{x}}$.

The number of iterations of the for-loop is $\lceil 2/\delta_0 \rceil^d$ at most. By the explanation in the proof of Lemma 7.1, $\|\mathbf{F}(\mathbf{x})\|$ and $\mathbf{DF}(\mathbf{x})$ can be computed from \mathbf{a} in $O(N(1 + p_{\max}/d)nd)$ operations. By Lemma 7.3, $s_{\min}(\mathbf{x})$ can be computed in $O(\log \kappa + \log \log d\omega_d)$ operations. Hence, the total complexity of the algorithm is bounded, for some absolute constant c, by

$$(cd^5p_{\max}^2\log(p_{\max}))^d Nd^2(1+p_{\max}/d).$$

By Lemma 5.4, the probability of event $\mathcal{E}(d)$ is at least $1 - C/\sqrt{d}$. To complete the proof, we will show that on \mathcal{E} , Algorithm 2 outputs an approximate solution. The value of κ above was chosen so that, by Lemma 7.3,

$$s_{\min}(\boldsymbol{x}) \in \left[\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x})\big|_{\mathsf{T}_{\boldsymbol{x}}}), \sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x})\big|_{\mathsf{T}_{\boldsymbol{x}}}) + \frac{\bar{\eta}}{2d^{7/4}}\right].$$

Suppose that on one of the iterations of the for-loop, a point $\boldsymbol{x} \in \mathbb{S}^{d-1}$ such that $\|\boldsymbol{F}(\boldsymbol{x})\| \leq t$ and $s_{\min}(\boldsymbol{x}) \geq \frac{\bar{\eta}}{d^{7/4}}$ is found. Then, $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{x})|_{\mathsf{T}_{\boldsymbol{x}}}) \geq \frac{\bar{\eta}}{2d^{7/4}}$. Hence, by Corollary 6.4 the output will be an approximate solution.

It remains to show that on \mathcal{E} , a point as above is indeed found in the for-loop (instead of returning $\boldsymbol{x}^{\text{BF}} = (1, 0, \dots, 0)$). Namely, denoting by $\bar{\mathcal{N}}$ the projection of \mathcal{N} to \mathbb{S}^{d-1} , we need to show that there exists some $\boldsymbol{x} \in \bar{\mathcal{N}}$ such that $\|\boldsymbol{F}(\boldsymbol{x})\| \leq t$ and $s_{\min}(\boldsymbol{x}) \geq \frac{\bar{\eta}}{d^{7/4}}$. The set \mathcal{N} over which the for-loop runs is a $\sqrt{d}\delta_0$ -net of the cube $[-1, 1]^d$. Suppose that \boldsymbol{w} is an arbitrary point in \mathbb{S}^{d-1} and let \boldsymbol{y} be the closest point to it in \mathcal{N} . Since $\|\boldsymbol{y} - \boldsymbol{y}/\|\boldsymbol{y}\|\| \leq \|\boldsymbol{w} - \boldsymbol{y}\| \leq \sqrt{d}\delta_0$, the projection $\bar{\mathcal{N}}$ is a $2\sqrt{d}\delta_0$ -net of \mathbb{S}^{d-1} .

By Point 3 of Definition 5.3, there exists some solution $\boldsymbol{\alpha} \in \mathbb{S}^{d-1}$ such that $\sigma_{\min}(\boldsymbol{DF}(\boldsymbol{\alpha})\boldsymbol{U}_{\boldsymbol{\alpha}}) \geq \eta/\sqrt{d}$. Let \boldsymbol{x} be the closest point to $\boldsymbol{\alpha}$ in $\bar{\mathcal{N}}$. Then, from Points 1 and 2 of Definition 5.3,

$$\|\boldsymbol{F}(\boldsymbol{x})\| \leq \|\boldsymbol{F}(\boldsymbol{\alpha})\| + 2\sqrt{d}\delta_0 \operatorname{Lip}^{(0)}(\boldsymbol{F}) \leq \frac{2\delta C_0}{d^4 p_{\max}^2 \sqrt{\log(p_{\max})}} = t,$$

and

$$s_{\min}(\boldsymbol{x}) \ge \sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{x})\boldsymbol{U}_{\boldsymbol{x}}) \ge \sigma_{\min}(\boldsymbol{D}\boldsymbol{F}(\boldsymbol{\alpha})\boldsymbol{U}_{\boldsymbol{\alpha}}) - 2\sqrt{d}\delta_{0}\operatorname{Lip}^{(1)}(\boldsymbol{F})$$
$$\ge \frac{\eta}{d^{7/4}} - \frac{\delta}{d^{4}p_{\max}\sqrt{\log(p_{\max})}} \ge \frac{\eta}{2d^{7/4}} = \frac{\bar{\eta}}{d^{7/4}}.$$

This completes the proof.

Acknowledgements

AM was supported by the NSF through award DMS-2031883, the Simons Foundation through Award 814639 for the Collaboration on the Theoretical Foundations of Deep Learning, the NSF

grant CCF2006489 and the ONR grant N00014-18-1-2729. ES was supported by the Israel Science Foundation theough grant No. 2055/21 and a research grant from the Center for Scientific Excellence at the Weizmann Institute of Science. ES is the incumbent of the Skirball Chair in New Scientists.

A Tools: Linear algebra

Theorem 5 (Wedin [Wed72]). Let $A_0, A_1 \in \mathbb{R}^{m \times n}$ have singular value decomposition (for $a \in \{0,1\}$)

$$\boldsymbol{A}_a = \boldsymbol{U}_a \boldsymbol{\Sigma}_a \boldsymbol{V}_a^{\mathsf{T}} \,, \tag{143}$$

with Σ_a containing the singular values in decreasing order. Further let $U_{a,+} \in \mathbb{R}^{m \times k(a)}$, $V_{a,+} \in \mathbb{R}^{n \times k(a)}$, be formed by the first k(a) columns of U_a , V_a , so that

$$\boldsymbol{U}_{a} = \begin{bmatrix} \boldsymbol{U}_{a,+} | \boldsymbol{U}_{a,-} \end{bmatrix}, \qquad \boldsymbol{V}_{a} = \begin{bmatrix} \boldsymbol{V}_{a,+} | \boldsymbol{V}_{a,-} \end{bmatrix}.$$
(144)

Finally assume $\Delta \equiv \sigma_{k(1)}(\mathbf{A}_1) - \sigma_{k(0)+1}(\mathbf{A}_0) > 0$. Let $\mathbf{P}_a = \mathbf{V}_{a,+}\mathbf{V}_{a,+}^{\mathsf{T}}$ (respectively $\mathbf{Q}_a = \mathbf{U}_{a,+}\mathbf{U}_{a,+}^{\mathsf{T}}$) denote the projector onto the right singular space (left singular space) corresponding to large singular values of \mathbf{A}_a . Then we have

$$\|(\boldsymbol{I}_{n}-\boldsymbol{P}_{0})\boldsymbol{P}_{1}\|_{\text{op}} \leq \frac{1}{\Delta} \left\{ \|(\boldsymbol{I}-\boldsymbol{Q}_{0})(\boldsymbol{A}_{0}-\boldsymbol{A}_{1})\boldsymbol{P}_{1}\|_{\text{op}} \vee \|\boldsymbol{Q}_{1}(\boldsymbol{A}_{0}-\boldsymbol{A}_{1})(\boldsymbol{I}-\boldsymbol{P}_{0})\|_{\text{op}} \right\}, \quad (145)$$

If instead we have $\Delta \equiv \sigma_{k(0)}(\mathbf{A}_0) - \sigma_{k(1)+1}(\mathbf{A}_1) > 0$, then

$$\left\| \boldsymbol{P}_{0}(\boldsymbol{I}_{n} - \boldsymbol{P}_{1}) \right\|_{\text{op}} \leq \frac{1}{\Delta} \left\{ \left\| (\boldsymbol{I} - \boldsymbol{Q}_{1})(\boldsymbol{A}_{0} - \boldsymbol{A}_{1})\boldsymbol{P}_{0} \right\|_{\text{op}} \lor \left\| \boldsymbol{Q}_{0}(\boldsymbol{A}_{0} - \boldsymbol{A}_{1})(\boldsymbol{I} - \boldsymbol{P}_{1}) \right\|_{\text{op}} \right\}.$$
 (146)

Corollary A.1. Let $A_0, A_1 \in \mathbb{R}^{n \times d}$ and denote by $v_1(a), \ldots, v_n(a)$, $a \in \{0, 1\}$ two bases of right singular vectors of these matrices, with associated singular values $\sigma_1(A_a) \geq \cdots \geq \sigma_d(A_a)$ (including vanishing singular values). Assume n < d and define, for $m \geq 1$,

$$V_{0,m} = \operatorname{span}(\boldsymbol{v}_{n-m+2}(0), \dots, \boldsymbol{v}_d(0)), \qquad (147)$$

$$V_{1,1} = \operatorname{span}(\boldsymbol{v}_{n+1}(1), \dots, \boldsymbol{v}_d(1)).$$
(148)

(In particular, $V_{1,1}$ is a subspace of the null space of A_1 .) Finally, denote by $E_1 \in \mathbb{R}^{d \times (d-n)}$ a matrix whose columns form an orthonormal basis of $V_{1,1}$, and by $\overline{E}_0 \in \mathbb{R}^{d \times (d-n+m-1)}$ a matrix whose columns form an orthonormal basis of $V_{0,m}^{\perp}$.

Then we have

$$\left\|\boldsymbol{E}_{1}^{\mathsf{T}}\overline{\boldsymbol{E}}_{0}\right\|_{\mathrm{op}} \leq \frac{1}{\sigma_{n-m+1}(\boldsymbol{A}_{0})} \|\boldsymbol{A}_{0}-\boldsymbol{A}_{1}\|_{\mathrm{op}}.$$
(149)

B Tools: Random matrix theory

Given the symmetric matrix $\mathbf{M} \in \mathbb{R}^{n \times n}$, we denote by $\lambda_1(\mathbf{M}) \leq \lambda_2(\mathbf{M}) \leq \cdots \leq \lambda_n(\mathbf{M})$ its the eigenvalues in increasing order. We denote by $F_{\rm sc}(t) = \frac{1}{2\pi} \int_{-2}^{(t \wedge 2) \vee (-2)} \sqrt{4 - x^2} \, \mathrm{d}x$ the semicircle distribution.

Lemma B.1. For any $t \in (-2,2)$ and $\varepsilon > 0$ there exists a constant $C_0 = C_0(t,\varepsilon)$ such that for $W \sim \text{GOE}(N)$ and any k such that $k/N \leq F_{sc}(t) - \varepsilon$,

$$\mathbb{P}\Big(\lambda_k(\boldsymbol{W}) > t\sqrt{N}\Big) \le C_0 \, e^{-N^2/C_0} \,. \tag{150}$$

Proof. Since $\lambda_k(\mathbf{W}) > t\sqrt{N}$ if and only if $\#\{\lambda_i(\mathbf{W}) \le t\sqrt{N}\} < k$, the lemma follows from the large deviation principle for the empirical measure of eigenvalues of a GOE matrix proved in [BAG97, Theorem 2.1.1].

Lemma B.2. For all $c_0 > 0$, there exists $C_*(c_0)$ such that the following holds. If $\mathbb{Z} \sim \mathsf{GOE}(M, N)$, $c_0 N < M \leq N$ and $1 \leq \ell \leq M/2$ then, for any $\Delta > 0$,

$$\mathbb{P}\left(\lambda_{\ell}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq \Delta^{2}\left(\sqrt{N} - \sqrt{M-1}\right)^{2}\right) \leq \left(C_{*}(c_{0})\Delta\right)^{\ell(N-M+\ell)}.$$
(151)

Proof. Let $f(\boldsymbol{u}) = f(u_1, \ldots, u_M)$ denote the joint density of the ordered eigenvalues $\lambda_1 \leq \cdots \leq \lambda_M$, $\lambda_i = \lambda_i (\boldsymbol{Z} \boldsymbol{Z}^{\mathsf{T}})$. We have [AGZ09, Proposition 4.1.3]

$$f(\boldsymbol{u}) = \frac{1}{Z_{M,N}} \prod_{i=1}^{M} u_i^{(N-M-1)/2} e^{-u_i/2} \prod_{i < j} (u_j - u_i) \, \mathbf{1}_{\mathbb{J}_M}(\boldsymbol{u}) \,, \tag{152}$$

where $\mathbb{J}_M = \mathbb{J}_M(0)$ and $\mathbb{J}_M(a) \subseteq \mathbb{R}^M$ is the set $\mathbb{J}_M(a) := \{ \boldsymbol{u} : a \leq u_1 \leq \cdots \leq u_M \}$, and

$$Z_{M,N} = \left(\frac{2^N}{\pi}\right)^{M/2} \prod_{i=1}^M \Gamma\left(\frac{N-i+1}{2}\right) \Gamma\left(\frac{M-i+1}{2}\right).$$
(153)

We denote joint density of the lowest ℓ eigenvalues $\boldsymbol{u}_{\leq \ell} = (u_1, \ldots, u_\ell)$ by f_ℓ . It is given by

$$f_{\ell}(\boldsymbol{u}_{\leq \ell}) = \frac{F_{\ell}(\boldsymbol{u}_{\leq \ell})}{Z_{M,N}} \int_{\mathbb{J}_{M-\ell}(u_{\ell})} \prod_{i=\ell+1}^{M} \prod_{m=1}^{\ell} (u_{i} - u_{m}) \prod_{i=\ell+1}^{M} u_{i}^{(N-M-1)/2} e^{-u_{i}/2} \prod_{\ell < i < j} (u_{j} - u_{i}) \mathrm{d}\boldsymbol{u}_{>\ell} \,,$$

$$F_{\ell}(\boldsymbol{u}_{\leq \ell}) := \prod_{i=1}^{\ell} u_{i}^{\frac{N-M-1}{2}} e^{-\frac{u_{i}}{2}} \prod_{i < j \leq \ell} (u_{j} - u_{i}) \, \mathbf{1}_{\mathbb{J}_{\ell}}(\boldsymbol{u}_{\leq \ell}) \,.$$

We can bound the integral as

$$f_{\ell}(\boldsymbol{u}_{\leq \ell}) \leq \frac{F_{\ell}(\boldsymbol{u}_{\leq \ell})}{Z_{M,N}} \int_{\mathbb{J}_{M-\ell}(u_{\ell})} \prod_{i=\ell+1}^{M} u_{i}^{(N-M+2\ell-1)/2} e^{-u_{i}/2} \prod_{\ell < i < j} (u_{j} - u_{i}) \mathrm{d}\boldsymbol{u}_{>\ell}$$
$$\leq \frac{F_{\ell}(\boldsymbol{u}_{\leq \ell})}{Z_{M,N}} \int_{\mathbb{J}_{M-\ell}(0)} \prod_{i=\ell+1}^{M} u_{i}^{(N-M+2\ell-1)/2} e^{-u_{i}/2} \prod_{\ell < i < j} (u_{j} - u_{i}) \mathrm{d}\boldsymbol{u}_{>\ell}$$
$$= \frac{Z_{M-\ell,N+\ell}}{Z_{M,N}} F_{\ell}(\boldsymbol{u}_{\leq \ell}) \,.$$

Therefore

$$\mathbb{P}\left(\lambda_{\ell}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq \delta\right) \leq \frac{Z_{M-\ell,N+\ell}}{Z_{M,N}} \int F_{\ell}(\boldsymbol{u}_{\leq \ell}) \mathbf{1}_{u_{\ell} \leq \delta} \mathrm{d}\boldsymbol{u}_{\leq \ell}$$

$$\leq \frac{Z_{M-\ell,N+\ell}}{Z_{M,N}} \frac{1}{\ell!} \int \prod_{i=1}^{\ell} u_i^{\frac{N-M-1}{2}} \prod_{i < j \leq \ell} |u_j - u_i| \prod_{i=1}^{\ell} \mathbf{1}_{0 \leq u_i \leq \delta} \mathrm{d}\boldsymbol{u}_{\leq \ell}$$

$$\leq \frac{Z_{M-\ell,N+\ell}}{Z_{M,N}} \frac{1}{\ell!} \mathscr{S}_{\ell} \left(\frac{N-M+1}{2}, 1, \frac{1}{2}\right) \cdot \delta^{\ell(N-M+\ell)/2} ,$$

where $\mathscr{S}_{\ell}(\alpha, \beta, \gamma)$ is Selberg's integral [AGZ09, Theorem 2,5.8]. In particular, we have

$$\mathscr{S}_{\ell}(\alpha, 1, 1/2) = \prod_{i=0}^{\ell-1} \frac{\Gamma(\alpha + (i/2))\Gamma(1 + (i/2))\Gamma((3+i)/2)}{\Gamma(\alpha + 1 + (\ell + i - 1)/2)\Gamma(3/2)}.$$

After reordering we can write the above bound as

$$\mathbb{P}\Big(\lambda_{\ell}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq \delta\Big) \leq \frac{1}{\Gamma(3/2)^{\ell}\ell!} \frac{Q_{M,N,\ell}}{R_{M,N,\ell}} S_{M,N,\ell} T_{M,N,\ell} \cdot \delta^{\ell(N-M+\ell)/2},$$
(154)

where

$$Q_{M,N,\ell} := 2^{-\frac{\ell}{2}(N-M) - \frac{\ell^2}{2}} \pi^{\ell/2},$$

$$R_{M,N,\ell} := \prod_{i=1}^{2\ell} \Gamma((N-M+i)/2),$$

$$S_{M,N,\ell} := \prod_{i=1}^{\ell} \frac{\Gamma((N+i)/2)}{\Gamma((M-\ell+i)/2)},$$

$$T_{M,N,\ell} := \prod_{i=1}^{\ell} \frac{\Gamma((N-M+i)/2)}{\Gamma((N-M+\ell+1+i)/2)} \cdot \prod_{i=1}^{\ell} \Gamma((i+1)/2) \Gamma((i+2)/2).$$

We estimate the above quantities in the following lemma.

Lemma B.3. For all $c_0 > 0$, there exists a constant $C_1 = C_1(c_0)$ such that the following holds. If $c_0 N < M \le N$ and $1 \le \ell \le M/2$, then

$$\log \frac{T_{M,N,\ell}}{R_{M,N,\ell}} \le -\ell(N - M + \ell) \log(N - M + \ell) + C_1 \ell(N - M + \ell),$$
(155)

$$\log S_{M,N,\ell} \le \ell \frac{N - M + \ell}{2} \log M + C_1 \ell (N - M + 1).$$
(156)

The proof of this lemma is postponed. We will now use it to prove the claim of the present one. Continuing from Eq. (154) (and denoting by C constants that depend on c_0 and can change from line to line):

$$\mathbb{P}\left(\lambda_{\ell}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq \Delta^{2}\left(\sqrt{N} - \sqrt{M-1}\right)^{2}\right) \leq \mathbb{P}\left(\lambda_{\ell}(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}}) \leq C\Delta^{2}\frac{(N-M)^{2}}{N}\right) \\
\leq \frac{T_{M,N,\ell}}{R_{M,N,\ell}} S_{M,N,\ell} \cdot (C\Delta)^{\ell(N-M+\ell)} \cdot \left(\frac{(N-M)^{2}}{N}\right)^{\ell(N-M+\ell)/2} \\
\leq (C\Delta)^{\ell(N-M+\ell)} \exp(E_{M,N,\ell}),$$

where, for C_1, C_2 depending on c_0 ,

$$E_{M,N,\ell} := -\ell(N - M + \ell) \log(N - M + \ell) + \ell \frac{N - M + \ell}{2} \log M + C_1 \ell (N - M + \ell) + \ell (N - M + \ell) \log(N - M) - \frac{1}{2} \ell (N - M + \ell) \log N \leq C_2 \ell (N - M + \ell) .$$

Substituting above, and adjusting the constant C yields the claim.

We now complete the last proof by proving Lemma B.3.

Proof of Lemma B.3. Recalling the Legendre duplication formula, $\Gamma(z)\Gamma(z+\frac{1}{2}) = 2^{1-2z}\sqrt{\pi} \Gamma(2z)$, we have that

$$\frac{T_{M,N,\ell}}{R_{M,N,\ell}} = \prod_{i=1}^{\ell} \frac{\Gamma((i+1)/2)\Gamma((i+2)/2)}{\Gamma((N-M+\ell+1+i)/2)\Gamma((N-M+\ell+i)/2)}$$
$$= 2^{\ell(N-M+\ell-1)} \prod_{i=1}^{\ell} \frac{\Gamma(i+1)}{\Gamma(N-M+\ell+i)}.$$

Using $|\log(n!) - n\log n| < Cn$ for some absolute constant C > 0, (155) follows since

$$\log\left(\frac{T_{M,N,\ell}}{R_{M,N,\ell}}\right) \le 2(C + \log 2)\ell(N - M + 2\ell) \\ + \sum_{i=1}^{\ell} \left(i\log(i) - (N - M + \ell + i - 1)\log(N - M + \ell + i - 1)\right) \\ \le 2(C + \log 2)\ell(N - M + 2\ell) - \ell(N - M + \ell - 1)\log(N - M + \ell).$$

In order to prove Eq. (156), we use Stirling's formula to get, for some absolute constant C > 0,

$$\begin{split} \log S_{M,N,\ell} &\leq \sum_{i=1}^{\ell} \left\{ \frac{N+i}{2} \log \frac{N+i}{2} - \frac{N+i}{2} - \frac{1}{2} \log \frac{N+i}{2} \\ &- \frac{M-\ell+i}{2} \log \frac{M-\ell+i}{2} + \frac{M-\ell+i}{2} + \frac{1}{2} \log \frac{M-\ell+i}{2} \right\} + C\ell \\ &= \sum_{i=1}^{\ell} \left\{ \frac{N+i}{2} \log \frac{N+i}{M-\ell+i} + \frac{N-M+\ell}{2} \log \frac{M-\ell+i}{2} \right\} \\ &- \sum_{i=1}^{\ell} \left\{ \frac{N-M+\ell}{2} + \frac{1}{2} \log \frac{N+i}{M-\ell+i} \right\} + C\ell \\ &\stackrel{(a)}{\leq} - \sum_{i=1}^{\ell} \frac{N+i}{2} \log \left(1 - \frac{N-M+\ell}{N+i} \right) + \ell \frac{N-M+\ell}{2} \log M + C\ell \\ &\stackrel{(b)}{\leq} \ell \frac{N-M+\ell}{2} \log M + C\ell + C_1(c_0)\ell(N-M) \,, \end{split}$$

where in (a) we omitted the negative sum and for (b) we applied the inequality $-\log(1-x) \le Cx$ for $0 \le x \le 1 - C^{-1}$ with $C = (1-x)^{-1}$ to $x = (N - M + \ell)/(N + i)$.

Since $(\sigma_{\min}(\boldsymbol{Z}))^2 = \lambda_1(\boldsymbol{Z}\boldsymbol{Z}^{\mathsf{T}})$, the following is an immediate consequence of Lemma B.2.

Corollary B.4. In the setting of Lemma B.2,

$$\mathbb{P}\Big(\sigma_{\min}(\mathbf{Z}) \le \varepsilon \big(\sqrt{N} - \sqrt{M-1}\big)\Big) \le (C_*(c_0)\varepsilon)^{N-M+1}.$$
(157)

References

[ABAČ13] Antonio Auffinger, Gérard Ben Arous, and Jiri Černý, Random matrices and complexity of spin glasses, Comm. Pure Appl. Math. 66 (2013), no. 2, 165–201. MR 2999295

- [ABB⁺16] Diego Armentano, Carlos Beltrán, Peter Bürgisser, Felipe Cucker, and Michael Shub, Condition length and complexity for the solution of polynomial systems, Found. Comput. Math. 16 (2016), no. 6, 1401–1422. MR 3579713
- [AGZ09] Greg W. Anderson, Alice Guionnet, and Ofer Zeitouni, An introduction to random matrices, Cambridge University Press, 2009.
- [AW09] J.-M. Azaïs and M. Wschebor, Level sets and extrema of random processes and fields, John Wiley & Sons, Inc., Hoboken, NJ, 2009. MR 2478201
- [BAG97] G. Ben Arous and A. Guionnet, Large deviations for wigner's law and voiculescu's noncommutative entropy, Probability Theory and Related Fields 108 (1997), no. 4, 517–542.
- [BC11] Peter Bürgisser and Felipe Cucker, On a problem posed by Steve Smale, Ann. of Math.
 (2) 174 (2011), no. 3, 1785–1836. MR 2846491
- [BCL23] Peter Bürgisser, Felipe Cucker, and Pierre Lairez, *Rigid continuation paths II. structured polynomial systems*, Forum Math. Pi **11** (2023), Paper No. e12, 44. MR 4575356
- [BCSS98] Lenore Blum, Felipe Cucker, Michael Shub, and Steve Smale, *Complexity and real computation*, Springer-Verlag, New York, 1998, With a foreword by Richard M. Karp. MR 1479636
- [BP08] Carlos Beltrán and Luis Miguel Pardo, On Smale's 17th problem: a probabilistic positive solution, Found. Comput. Math. 8 (2008), no. 1, 1–43. MR 2403529
- [BP09] _____, Smale's 17th problem: average polynomial time to compute affine and projective solutions, J. Amer. Math. Soc. **22** (2009), no. 2, 363–385. MR 2476778
- [BP11] _____, Fast linear homotopy to find approximate zeros of polynomial systems, Found. Comput. Math. **11** (2011), no. 1, 95–129. MR 2754191
- [BS09] Carlos Beltrán and Michael Shub, Complexity of Bezout's theorem. VII. Distance estimates in the condition metric, Found. Comput. Math. 9 (2009), no. 2, 179–195. MR 2496559
- [BSS89] Lenore Blum, Michael Shub, and Steve Smale, On a theory of computation and complexity over the real numbers: NP-completeness, recursive functions and universal machines, Bull. Amer. Math. Soc. (N.S.) 21 (1989), no. 1, 1–46. MR 974426
- [Dre78] Franz-Josef Drexler, Eine Methode zur Berechnung sämtlicher Lösungen von Polynomgleichungssystemen, Numer. Math. 29 (1977/78), no. 1, 45–58. MR 483386
- [EAMS21] Ahmed El Alaoui, Andrea Montanari, and Mark Sellke, Optimization of mean-field spin glasses, The Annals of Probability 49 (2021), no. 6, 2922–2960.
- [GZ79] C. B. García and W. I. Zangwill, Finding all solutions to polynomial systems and other systems of equations, Math. Programming 16 (1979), no. 2, 159–176. MR 527572
- [HS21] Brice Huang and Mark Sellke, *Tight Lipschitz Hardness for Optimizing Mean Field Spin Glasses*, arXiv:2110.07847 (2021).
- [Kel78] Herbert B Keller, *Global homotopies and newton methods*, Recent advances in numerical analysis, Elsevier, 1978, pp. 73–94.

- [Lai17] Pierre Lairez, A deterministic algorithm to compute approximate roots of polynomial systems in polynomial average time, Found. Comput. Math. 17 (2017), no. 5, 1265– 1292. MR 3709332
- [Lai20] _____, Rigid continuation paths I. Quasilinear average complexity for solving polynomial systems, J. Amer. Math. Soc. **33** (2020), no. 2, 487–526. MR 4073867
- [Mon19] Andrea Montanari, Optimization of the sherrington-kirkpatrick hamiltonian, 2019 IEEE 60th Annual Symposium on Foundations of Computer Science (FOCS), IEEE, 2019, pp. 1417–1433.
- [MS23] Andrea Montanari and Eliran Subag, Solving overparametrized systems of random equations: I. model and algorithms for approximate solutions, arXiv:2306.13326 (2023).
- [Shu93] Michael Shub, Some remarks on Bezout's theorem and complexity theory, From Topology to Computation: Proceedings of the Smalefest, Springer, New York, 1993, pp. 443–455. MR 1246139
- [Shu09] _____, Complexity of Bezout's theorem. VI. Geodesics in the condition (number) metric, Found. Comput. Math. 9 (2009), no. 2, 171–178. MR 2496558
- [Sma98] Steve Smale, Mathematical problems for the next century, Mathematical Intelligencer 20 (1998), no. 2, 7–15.
- [SS93a] Michael Shub and Steve Smale, Complexity of Bézout's theorem. I. Geometric aspects, J. Amer. Math. Soc. 6 (1993), no. 2, 459–501. MR 1175980
- [SS93b] _____, Complexity of Bezout's theorem. II. Volumes and probabilities, Computational algebraic geometry (Nice, 1992), Progr. Math., vol. 109, Birkhäuser Boston, Boston, MA, 1993, pp. 267–285. MR 1230872
- [SS93c] _____, Complexity of Bezout's theorem. III. Condition number and packing, vol. 9, 1993, Festschrift for Joseph F. Traub, Part I, pp. 4–14. MR 1213484
- [SS94] _____, Complexity of Bezout's theorem. V. Polynomial time, vol. 133, 1994, Selected papers of the Workshop on Continuous Algorithms and Complexity (Barcelona, 1993), pp. 141–164. MR 1294430
- [SS96] _____, Complexity of Bezout's theorem. IV. Probability of success; extensions, SIAM J. Numer. Anal. 33 (1996), no. 1, 128–148. MR 1377247
- [Sub17] Eliran Subag, The complexity of spherical p-spin models—a second moment approach, Ann. Probab. **45** (2017), no. 5, 3385–3450. MR 3706746
- [Sub18] _____, Following the ground-states of full-RSB spherical spin glasses, arXiv:1812.04588 (2018).
- [Sub23] _____, Concentration for the zero set of random polynomial systems, arXiv:2303.11924 (2023).
- [Wed72] Per-Åke Wedin, Perturbation bounds in connection with singular value decomposition, BIT Numerical Mathematics **12** (1972), no. 1, 99–111.
- [Wsc05] Mario Wschebor, On the Kostlan-Shub-Smale model for random polynomial systems. Variance of the number of roots, J. Complexity **21** (2005), no. 6, 773–789. MR 2182444