

## Choked precessing jets in tidal disruption events and high-energy neutrinos

QI-RUI YANG,<sup>1,2</sup> JIAN-HE ZHENG,<sup>1,2</sup> RUO-YU LIU,<sup>1,2</sup> AND XIANG-YU WANG<sup>1,2</sup>

<sup>1</sup>*School of Astronomy and Space Science, Nanjing University, Nanjing 210023, People's Republic of China*

<sup>2</sup>*Key laboratory of Modern Astronomy and Astrophysics (Nanjing University),  
Ministry of Education, Nanjing 210023, People's Republic of China*

### ABSTRACT

It has been suggested that relativistic jets might have been commonly formed in tidal disruption events (TDEs), but those with relatively weak power could be choked by the surrounding envelope. The discovery of high-energy neutrinos possibly associated with some normal TDEs may support this picture in the hypothesis that the neutrinos are produced by choked jets. Recently, it was noted that disrupted stars generally have misaligned orbits with respect to the supermassive black hole spin axis and highly misaligned precessing jets are more likely to be choked. Here we revisit the jet break-out condition for misaligned precessing jets by considering the jet could be collimated by the cocoon pressure while propagating in the disk wind envelope. The jet head opening angle decreases as the jet propagates in the envelope, but the minimum power of a successful jet remains unchanged in terms of the physical jet power. We further calculate the neutrino flux from choked precessing jets, assuming that the cocoon energy does not exceed the kinetic energy of the disk wind. We find that neutrino flux from highly misaligned choked jets is sufficient to explain the neutrinos from AT2019aal, while it is marginal to explain the neutrinos from AT2019dsg and AT2019fdr. The latter could be produced by weakly misaligned choked jets, since the duty cycle that the jet sweeps across increases as the misaligned angle decreases. We also show that the population of choked TDE jets could contribute to  $\sim 10\%$  of the observed diffuse neutrino flux measured by IceCube.

### 1. INTRODUCTION

A couple of TDEs with relativistic jets (hereafter “jetted TDEs”) have been observed, including Swift J1644+57 (Bloom et al. 2011), Swift J2058+05 (Cenko et al. 2012), Swift J1112+82 (Brown et al. 2015), and AT2022cmc (Andreoni et al. 2022). The volumetric rate of on-axis jetted TDEs has been inferred to be  $\sim 0.03 \text{Gpc}^{-3} \text{yr}^{-1}$  (Burrows et al. 2011; Andreoni et al. 2022). If we correct for a beaming factor  $f_b \sim 0.01$  (adopting a typical half-opening angle of  $\theta_j \sim 0.1$ ), the inferred intrinsic rate of jetted TDEs is  $\sim 3 \text{Gpc}^{-3} \text{yr}^{-1}$ . This is a very small fraction ( $\sim 1\%$ ) of the rate of all TDEs observed in various surveys,  $R \sim 300 \text{Gpc}^{-3} \text{yr}^{-1}$  (van Velzen 2018; Sazonov et al. 2021; Yao et al. 2023). The isotropic equivalent jet luminosity of these jetted TDEs is in the range of  $\sim 10^{47} - 10^{48} \text{ergs}^{-1}$ . Since the black hole accretion is expected to produce a continuous distribution of jet luminosity, as has been seen in active

galactic nuclei (AGN) jets, the fact that no less powerful jets have been found in TDEs so far is a puzzle.

Wang & Liu (2016) suggested that, while powerful jets, such as that in Swift J1644+57, can successfully break out from the surrounding envelope, weak jets could be choked during the propagation inside the envelope. This could explain why no less powerful jets have been found in TDEs. Like successful jets, the choked jet can also accelerate cosmic rays through shocks and further produce neutrinos via interaction with the surrounding dense photons (Wang et al. 2011; Wang & Liu 2016). Recent claims that high-energy neutrino events detected by IceCube are coincident with three TDE candidates (AT2019dsg, AT2019fdr and AT2019aal) stimulated the study on neutrino production in TDEs. Various production sites of high-energy neutrinos have been proposed, including successful jets (Liu et al. 2020; Winter & Lunardini 2021), choked jets (Zheng et al. 2023; Mukhopadhyay et al. 2023), disk outflow (Murase et al. 2020; Wu et al. 2022; Winter & Lunardini 2023), and the AGN core (Hayasaki & Yamazaki 2019; Murase et al. 2020). In particular, Zheng et al. (2023) explore the parameter space of the jets that can produce de-

tectable neutrino flux while being choked in the expanding envelope, finding that the cumulative neutrino numbers of AT2019fdp and AT2019aalc are consistent with the expected range imposed by observations, although the allowed parameter space for AT2019dsg is small. Mukhopadhyay et al. (2023) discussed high-energy neutrino production in delayed choked jets, and find that the a delay of the jet launch can make the jets more easily choked and can increasing the allowed neutrino flux.

Very recently, Teboul & Metzger (2023) and Lu et al. (2023) noted that disrupted stars generally have misaligned orbits with respect to the supermassive black hole (SMBH) spin axis and the accretion disk fed by the fallback stellar debris undergoes Lense-Thirring precession around the BH spin axis. They found that that precessing jets can only break out of the wind envelope confinement if the misalignment angles  $\theta_{LS}$  (between the jet axis and the BH spin axis) is less than a few times the jet opening angle  $\theta_j$ , explaining the tiny fraction of jetted TDEs. In most TDEs with  $\theta_{LS} \gg \theta_j$ , the jets are initially choked by the disk wind and may only break out later when the disk finally aligns itself with the spin axis due to the damping of the precession (Teboul & Metzger 2023; Lu et al. 2023).

In this paper, we study the neutrino production in choked TDE jets when the jet is precessing. Teboul & Metzger (2023) and Lu et al. (2023) both analytically study the propagation of a misaligned precessing jet inside the disk wind. Lu et al. (2023) also carry out three-dimensional relativistic hydrodynamic simulations of a precessing jet. In §2, we first revisit the jet propagation and the break-out condition for misaligned precessing jets by considering that the jet could be collimated by the cocoon pressure while it is propagating in the envelope. In §3, we calculate the neutrino flux produced by choked precessing jets and compare with the observations of three TDEs. We also calculate the diffuse neutrino emission from the population of choked TDEs. Finally we give discussions and conclusions in §4.

## 2. JET PROPAGATION AND THE BREAK-OUT CONDITION FOR PRECESSING JETS

After the stellar disruption, an accretion disk is formed in the orbital plane of the star whose normal is generally misaligned by an angle  $\theta_{LS}$  with respect to the SMBH spin axis. The accretion disk undergoes Lense-Thirring precession around the black hole spin axis and the jet is aligned with the instantaneous disk angular momentum (Teboul & Metzger 2023; Lu et al. 2023). Matter falls back to the disk at a highly super-

Eddington accretion rate  $\dot{M}$ , driving a disk wind of velocity  $\beta_w = v_w/c$  and mass-loss rate  $\dot{M}_w = f_w \dot{M}$ , where  $f_w < 1$  is the fraction of the mass fallback rate lost in the wind. Metzger & Stone (2016) argued that only a small fraction the returning debris ends up reaching the inner circular disk and accreting onto the SMBH, with the vast majority becoming unbound in a wind outflow. Due to the rapid precession, these disk winds will inevitably interact and collide with each other on large scales, leading to a quasi-spherical outflow.

We consider a relativistic jet with a kinetic luminosity  $L_j$ , an initial half-opening angle  $\theta_0$  and a bulk Lorentz factor  $\Gamma = 1/\theta_0$ . As the jet advances in the quasi-spherical wind envelope, the jet drives a bow shock ahead of it. The jet is capped by a termination shock, and a reverse shock propagates back into the jet, where the jet is decelerated and heated. The hot head material spills sideways, forming a cocoon that engulfs the jet and collimate it. The velocity of the jet head is determined by the balance of pressure between ambient gas and the jet head (Matzner 2003). In a moving wind envelope, we can express the velocity in the frame of the wind envelope by a coordinate transformation (e.g., Ioka & Nakamura 2018; Hamidani et al. 2020; Gottlieb & Nakar 2022; Lu et al. 2023):

$$\beta_h - \beta_w \simeq \frac{\beta_j - \beta_w}{1 + \tilde{L}^{-1/2}}, \quad (1)$$

where


$$\tilde{L} = \frac{L_j}{\Sigma_j \rho_w(r_h) c^3} = \frac{L_{j,\text{iso}} \beta_w}{M_w c^2}. \quad (2)$$

Here  $\beta_h$  is the jet head velocity in unit of the speed of light (measured in the lab frame),  $r_h$  is the radius of the jet head,  $\rho_w$  is wind density and  $\Sigma_j = \pi r_h^2 \theta_j^2$  is the jet cross-section at the location of the head with  $\theta_j$  being the jet head opening angle.  $L_{j,\text{iso}} = 2L_j/\theta_j^2$  is the isotropic equivalent luminosity and  $\dot{M}_w = 4\pi r_h^2 \rho_w \beta_w c$  is disk wind mass-loss rate.

Due to precession, the jet becomes episodic in a given direction. For such an episodic jet, if the timescale for the jet head to break out from the wind envelope is longer than the active timescale of the jet episode, the jet is choked (Lu et al. 2023). Following Lu et al. (2023), we consider that, along a fixed direction, the jet is “on” for a duration of  $t_{\text{on}}$  and “off” for a duration of  $T_{\text{prec}} - t_{\text{on}}$ , where  $T_{\text{prec}}$  is the precessional period. The jet head will catch up with the outer edge of the wind after a time

$$t_{\text{bo}} = \frac{\beta_w (T_{\text{prec}} - t_{\text{on}})}{\beta_h - \beta_w}. \quad (3)$$

The active timescale of the jet episode is the time for the reverse shock crossing the jet episode, which is (Lu



**Figure 1.** Evolution of  $\theta_j/\theta_0$  as a function of the jet efficiency  $\eta$  for various initial jet opening angles  $\theta_0$ .

et al. 2023)

$$t_{\text{act}} = \frac{\beta_j t_{\text{on}}}{\beta_j - \beta_h}. \quad (4)$$

A successful jet break-out requires  $t_{\text{bo}}/t_{\text{act}} < 1$ , which can be converted to the following form

$$\frac{L_{j,\text{iso}}}{\beta_w \dot{M}_w c^2} > \left( \frac{1 - \xi_{\text{duty}}}{\xi_{\text{duty}}} \right)^2, \quad (5)$$

where  $\xi_{\text{duty}} = t_{\text{on}}/T_{\text{prec}}$  is the duty cycle representing the fraction that the jet is "on" along a fixed direction of the whole precession period, which can be expressed as  $\xi_{\text{duty}} \approx \theta_j/(\pi \sin \theta_{\text{LS}})$  when  $\theta_{\text{LS}} \gg \theta_j$  (Lu et al. 2023). Thus the break-out condition is (Lu et al. 2023)

$$L_{j,\text{iso}} \gtrsim \beta_w \dot{M}_w c^2 \frac{\pi^2 \sin^2 \theta_{\text{LS}}}{\theta_j^2}. \quad (6)$$

Jets may be collimated by the cocoon pressure while it is propagating in the wind envelope. For the precession jet in our case, we assume that during the "on" period, the jet propagation is almost straight and the wind envelope is quasi-steady. The critical parameter  $\tilde{L}$  determines evolution of the jet (Bromberg et al. 2011). The jet is collimated if the collimation shock converges below the jet head, which occurs when  $\tilde{L} < \theta_0^{-4/3}$ . The opening angle of the jet head is thus given by (Bromberg et al. 2011)

$$\theta_j = \begin{cases} \tilde{L}^{1/4} \theta_0^2 = \left( \frac{2\eta \beta_w}{f_w} \right)^{1/6} \theta_0^{4/3}, & \tilde{L} < 1 \\ \tilde{L}^{3/4} \theta_0^2 = \left( \frac{2\eta \beta_w}{f_w} \right)^{3/10} \theta_0^{4/5}, & 1 < \tilde{L} < \theta_0^{-4/3} \\ \theta_0 & \tilde{L} > \theta_0^{-4/3} \end{cases} \quad (7)$$

where  $\eta = L_j/\dot{M}c^2$  is the jet efficiency. Fig.1 shows the evolution of  $\theta_j$  as a function of the jet efficiency  $\eta$ . It

indicates that for low jet efficiency, the jet head opening angle is significantly lower than the initial opening angle of the jet as the result of the collimation. The parameter  $\tilde{L}$  also indicates the velocity of jet head. For  $\tilde{L} < 1$ , the jet head is non-relativistic and vice versa.

We would like to stress that due to the collimation, the jet head opening angle  $\theta_j$  is smaller than the initial jet opening angle  $\theta_0$ , and as a result,  $L_{j,\text{iso}}$  may change due to collimation. So we express the break-out condition using the beam-corrected jet luminosity


$$L_j \geq \frac{1}{2} \dot{M} f_w c^2 \beta_w \pi^2 \sin^2 \theta_{\text{LS}}. \quad (8)$$

This condition is independent of the jet opening angle and holds regardless of whether the collimation effect is taken into account. This is because a decreasing opening angle  $\theta_j$  leads to a higher  $L_{j,\text{iso}}$  and also a shorter jet activity period  $t_{\text{act}}$  due to jet precession, which cancel with each other.


The wind speed  $v_w$  is roughly the local Keplerian speed of the accretion disk at the launching radius of the disk wind. For typical TDEs of a solar-like star by a  $10^6 - 10^7 M_\odot$  SMBH,  $v_w \simeq 0.1c$  (Metzger & Stone 2016; Lu et al. 2023). The discovery of highly blueshifted ( $v = 0.05c$ ) broad Balmer and metastable helium absorption lines in the optical spectra of the TDE AT2018zr is consistent with this wind speed (Hung et al. 2019). We show the break-out condition in Fig.2 for an example jet with  $\theta_0 = 1/3$ ,  $f_w = 0.5$  and  $\beta_w = 0.1$ . From top to the bottom (corresponding to a decreasing  $L_j/Mc^2$ ), the various color regions in Fig.2 represent the uncollimated break-out jet (grey region), collimated break-out jet with a relativistic jet head (red region), the collimated choked jet with a relativistic jet head (pink region), the collimated break jet with a sub-relativistic jet head (dark yellow region), the collimated choked jet with a sub-relativistic jet head (yellow region). Eq.(8) implies that break-out condition is independent of  $\theta_0$  or  $\theta_j$ . With fixed  $f_w$ ,  $\beta_w$  and misaligned angle  $\theta_{\text{LS}}$ , the jet efficiency  $\eta = L_j/\dot{M}c^2$  entirely determines whether the precessing jet breaks out or not.

We show the position of a jetted TDE like Swift J1644+57 in Fig.2. The luminosity of Swift J1644+57 is about  $10^{48}$  erg  $s^{-1}$  and the opening angle of the jet is assumed to be  $\theta_j = 0.1$ . For a wind mass-loss rate  $2M_\odot \text{yr}^{-1}$  (Lu et al. 2023), we derive  $\eta = 0.023$ . Swift J1644 can only have a successful jet break-out provided that the misalignment angle is sufficiently small ( $\theta_{\text{LS}} \lesssim 15^\circ$ ), consistent with the result in Lu et al. (2023).

For an easy comparison with the results in Lu et al. (2023), we express the break-out condition using the



**Figure 2.** Break-out condition in terms of the jet efficiency  $L_j/\dot{M}_w c^2$  and misaligned angle  $\theta_{LS}$  for the parameter values of  $\theta_0 = 0.33$ ,  $f_w = 0.5$ ,  $\beta_w = 0.1$ . The gray solid line shows the jet break-out criterion. The darker regions indicate break-out jet, whereas the lighter regions indicate choked jet. The yellow regions, the red regions and the gray region represent collimated jets with non-relativistic jet head, collimated jets with relativistic jet head and uncollimated jets, respectively. The horizontal black solid line shows the case of Swift J1644+57 with a jet efficiency of  $\eta = 0.023$ .



**Figure 3.** Break-out condition in terms of the dimensionless term  $L_{j,iso}/\beta_w \dot{M}_w c^2$  and misaligned angle  $\theta_{LS}$  using the same parameters as those in Fig. 2. The solid gray line represents the break-out criterion when the collimation caused by the cocoon pressure is taken into account, while the blue dot-dashed line shows the break-out condition without taking into account the collimation effect (i.e., assuming that jets are conical). The meanings of the regions with different colors are the same as that in Fig. 2.

isotropic luminosity  $L_{j,iso}$  by using Eq.(7) and Eq.(6):

$$\frac{L_{j,iso}}{\beta_w \dot{M}_w c^2} > \begin{cases} (\beta_w^{1/2} \pi^2 \sin^2 \theta_{LS} \theta_0^{-4})^{2/3} / \beta_w, & \tilde{L} < 1 \\ (\beta_w^{-1/2} \pi^2 \sin^2 \theta_{LS} \theta_0^{-4})^{2/5} / \beta_w, & 1 < \tilde{L} < \theta_0^{-4/3} \\ \pi^2 \sin^2 \theta_{LS} \theta_0^{-2}, & \tilde{L} > \theta_0^{-4/3} \end{cases} \quad (9)$$

The corresponding break-out condition and various regimes of the jet are shown in Fig. 3. The gray solid line represents the break-out criterion when the jet collimation effect is taken into account, while the blue dot-dashed line represents the case assuming a conical jet structure. A much larger  $L_{j,iso}$  is allowed for choked precessing jets when the jet collimation effect is properly taken into account.

### 3. HIGH-ENERGY NEUTRINOS FROM CHOKED PRECESSING JETS

The shocks in choked jets will accelerate cosmic rays to ultra-high energies (Farrar & Gruzinov 2009; Wang et al. 2011; Wang & Liu 2016) and further produce high-energy neutrinos (Wang & Liu 2016; Mukhopadhyay et al. 2023; Lu et al. 2023). Internal shocks may occur due to the internal collisions within the jets, resulted from the inhomogeneity in the jet velocity. Temporal variability with  $\delta t \approx 100s$  has been seen in the X-ray emission of the jetted TDE Swift J1644+57, which is thought to be generated by internal shocks (Burrows et al. 2011). We take a fiducial physical value  $\Gamma \sim 1/\theta_0$  for the jet Lorentz factor, and hence the collision occurs at  $R_{int} \approx \Gamma^2 c \delta t = 3 \times 10^{14} \text{cm} \Gamma_1^2$ . Internal shocks that propagate into the low-density jets are collisionless, although they locate inside the optically thick envelope (Wang & Liu 2016).

One important process for neutrino production in choked jets is the interaction between the cosmic rays and the thermal photons generated in the wind envelope (i.e., the  $p\gamma$  process). To calculate the neutrino flux, we need to know the density of thermal photons in the jet. Since the jet size is much smaller than that of the wind envelope, the density of thermal photons around the jet should be roughly equal to the density of thermal photons in the envelope at the same radius. We use the same treatment of thermal photons as that in Zheng et al. (2023). The photosphere temperature can be obtained from observations of TDEs. Then from the energy transferring equation, we can derive the temperature inside the envelope, which follows  $T \propto r^{-(n+1)/4}$  for  $r \ll R_{ph}$  (Roth et al. 2016), where  $R_{ph}$  is the photosphere radius. For the wind density, we have  $n = 2$  and  $T \propto r^{-3/4}$ . The typical energy of protons interacting with thermal photons with temperature  $T = 10^5 - 10^6 \text{K}$  is  $\epsilon_p = 0.1 \text{GeV}^2 / 3kT = 0.6 - 6 \text{PeV}$  and the neutrino

energy is correspondingly  $\epsilon_\nu = 0.05\epsilon_p = 30 - 300\text{TeV}$ , consistent with the measured energy of three neutrinos possibly associated with TDEs.

Considering that the duty cycle of the neutrino radiation cone for a precessing jet is  $\chi_{\text{duty}} = (1/\Gamma)/\pi\sin\theta_{\text{LS}}$ , the fluence of neutrinos from this precessing jet is

$$\epsilon_\nu \mathcal{F}_\nu \approx \int_0^{t_d/(1+z)} \frac{L_p f_{p\gamma} e^{-\frac{\epsilon_p}{\epsilon_{p,\text{max}}}}}{32\pi D_L^2 \ln(\epsilon_{p,\text{max}}/\epsilon_{p,\text{min}})} \chi_{\text{duty}} dt, \quad (10)$$

where  $t_d$  is the neutrino trigger time,  $L_p = \epsilon_p L_j \Gamma^2$  is the isotropic proton luminosity (assuming a neutrino radiation cone of  $1/\Gamma$ ),  $\epsilon_p$  is the fraction of the jet energy converted into relativistic protons and  $\Gamma$  is the bulk Lorentz factor of the jet.  $f_{p\gamma}$  is the fraction of energy loss by the  $p\gamma$  process, which can be obtained following Zheng et al. (2023). Assuming that the jet power scales with the accretion rate, the jet luminosity evolves with time as  $L_j \propto (t/\tau)^{-\frac{5}{3}}$  after the peak (Krolik & Piran 2012; Piran et al. 2015), where  $\tau$  is the characteristic fallback timescale, which is roughly the orbital period of the most bound debris (Gezari 2021), i.e.,


$$\tau = 41 M_{\text{BH},6}^{1/2} \left( \frac{M_\star}{M_\odot} \right)^{-0.1} \text{ day}, \quad (11)$$

with  $M_{\text{BH}}$  being the mass of SMBH and  $M_\star$  being the mass of the disrupted star. We take the minimum energy of protons as  $\epsilon_{p,\text{min}} = \Gamma m_p c^2$  and assume equipartition factors  $\epsilon_B = 0.1$  and  $\epsilon_p = 0.2$  in the calculation. Then we can obtain the cumulative muon neutrino number between 10 TeV and 1PeV by using

$$N_\nu = \int_{10\text{TeV}}^{1\text{PeV}} A_{\text{eff}}(\epsilon_\nu) \mathcal{F}_\nu d\epsilon_\nu. \quad (12)$$

where  $A_{\text{eff}}$  is the effective area of IceCube (Blaufuss et al. 2019).

We calculate the expected neutrino number from precessing jets as the function of the misaligned angle and jet power, as shown in Fig.4. We take fiducial values of  $\Gamma = 3$ ,  $\theta_0 = 0.33$  and  $\tau = 41\text{day}$  (corresponding to a SMBH with  $10^6 M_\odot$ ) for the jets. We take a reference value ( $D = 220\text{Mpc}$ ) for the TDE distance, equal to the distance of AT2019dsg. As expected, the neutrino flux increases as the jet power increases. The neutrino flux also increases as the misaligned angle decreases, because the duty cycle that the jet sweeps across increases as the misaligned angle decreases. The black solid line represents the break-out condition given by Eq.(7). The maximum neutrino number that precessing choked TDEs can produce is  $N_\nu \sim 0.1$ . The black dot-dashed line shows the case of TDEs like Swift J1644+57, which has




**Figure 4.** Color map of neutrinos numbers as the function of misaligned angle versus jet power in a general case. The solid black line represents the breakout condition, and the dot-dashed black line corresponds to a jet similar to Swift J1644 with a jet luminosity of  $10^{46}\text{erg s}^{-1}$ . The left side of the black solid line corresponds to the region that jets will not break through the envelope and vice versa.


a high luminosity of  $10^{46}\text{erg s}^{-1}$ . For this case, the maximum neutrino number produced by the choked jet is about  $N_\nu \sim 0.1$ .

### 3.1. Could the neutrinos from AT2019dsg, AT2019fdr and AT2019aalc be produced by choked precessing jets?

We explore the parameter space of the precessing jets that can produce detectable neutrino flux in three TDE candidates (AT2019dsg, AT2019fdr and AT2019aalc) while being choked in the wind envelope. To calculate the maximum neutrino flux in the three TDEs, we need to know the maximum luminosity allowed for choked jets, which can be obtained from Eq.(8). Meanwhile, the maximum luminosity could also be limited by the maximally allowed deposited energy by choked jets, since this deposited energy will eventually transfer to the wind envelope via the cocoon. A detailed numerical study is needed to understand how the deposited energy by choked jets affects the wind envelope and the radiation. For simplicity, we assume that the cocoon energy does not exceed the kinetic energy of the disk wind, otherwise the kinetic energy of the disk wind will increase significantly. It will be shown below that the critical jet luminosity limited by the maximally allowed deposited energy by choked jets is much smaller than the jet luminosity limited by the jet break-out condition for the three TDEs, so the real constraints on the neutrino flux come from the maximally allowed deposited energy.



**Figure 5.** Color map of the expected neutrino numbers for three TDEs (AT2019dsg, AT2019fdr, AT2019aalc) as a function of the jet luminosity  $L_j$  and initial opening angle  $\theta_0$  with a fixed misaligned angle  $\theta_{LS} = \pi/4$ . The white dashed line represent the minimal required neutrino numbers inferred from observations. The black vertical line represents the maximum jet luminosity limited by the break-out criterion represented as  $L_{BO}$ , while the white solid line represents the maximum jet luminosity limited by the wind power ( $\frac{1}{2}\dot{M}_w v_w^2$ ).



**Figure 6.** Color map of the expected neutrino numbers for three TDEs (AT2019dsg, AT2019fdr, AT2019aalc) as a function of the jet luminosity  $L_j$  and misaligned angle  $\theta_{LS}$  with a fixed initial jet half-opening angle of  $\theta_0 = 0.1$ . The meanings of each lines are the same as that in Fig.5.

The mass of the disrupted stars in three TDEs is not well-known. Similar to Zheng et al. (2023), we assume the mass of the disrupted stars scales with the bolometric energy, i.e.,  $M_\star \approx E_{bol} \approx L_{acc,max}\tau$  (Metzger & Stone 2016). Then the masses of the disrupted stars are estimated to be  $0.8M_\odot$ ,  $20M_\odot$  and  $4M_\odot$  for AT2019dsg, AT2019fdr and AT2019aalc, respectively. Another important parameter is the mass-loss rate  $\dot{M}_w = f_w \dot{M}$  in the disk wind, where  $\dot{M} = M_\star/(3\tau)$  is the mass fallback rate. The characteristic fallback timescale  $\tau$  for three TDE AT2019dsg, AT2019fdr, AT2019aalc are 40 days, 180 days and 140 days respectively (van Velzen et al. 2021).

We assume that the deposited cocoon energy should not exceed the kinetic energy of the disk wind too much, otherwise the wind kinetic energy will be significantly increased. The power of wind is represented by  $L_w = \frac{1}{2}\dot{M}_w v_w^2$ . In the following calculation, we set  $\beta_w = 0.1$  and  $f_w = 0.5$ . For AT2019dsg, the maxi-

um jet luminosity limited by a choked jet can reach  $L_j = 1.7 \times 10^{46} \text{ erg s}^{-1}$ . However, the wind kinetic power of AT2019dsg is estimated to be only  $3.5 \times 10^{44} \text{ erg s}^{-1}$ , which actually limits the allowed maximum jet luminosity. Similarly, for AT2019fdr, the maximum luminosity of a choked jet reaches  $L_j = 9.5 \times 10^{46} \text{ erg s}^{-1}$ , while the wind kinetic power is  $1.9 \times 10^{45} \text{ erg s}^{-1}$ . For AT2019aalc, the maximum jet luminosity for a choked jet is  $L_j = 2.5 \times 10^{46} \text{ erg s}^{-1}$ , while the wind kinetic luminosity is  $4.9 \times 10^{45} \text{ erg s}^{-1}$ .

The deposited cocoon energy can be expressed as

$$E_c = \int L_j(1 - \beta_h) dt. \quad (13)$$

If we consider the wind power for three TDEs as the jet luminosity, the cocoon energy, using Eq.(1) and Eq.(13), is estimated to be  $1.4 \times 10^{51} \text{ erg}$ ,  $3.5 \times 10^{52} \text{ erg}$  and  $7 \times 10^{51} \text{ erg}$  for AT2019dsg, AT2019fdr and AT2019aalc, respectively.

The expected neutrino numbers of three TDEs are shown by the color intensity in Fig.5. The dashed white lines show the parameter values corresponding to the minimum neutrino number of each TDE required by observations at 90% confidence(Stein et al. 2021). The solid black lines show the critical jet luminosity limited by the break-out condition and the solid white lines show the jet luminosity limited by the power of the disk wind. Fig.5 shows that precessing choked jet model can explain the neutrinos from AT2019aalc, while it can only marginally explain the neutrinos from AT2019dsg and AT2019fdr for a fixed large misaligned angle  $\theta_{LS} = \pi/4$ . Fig.6 illustrates the impact of the misaligned angle  $\theta_{LS}$  on the expected neutrino number. A weakly misaligned angle can result in a high duty cycle, consequently increase the neutrino flux. For AT2019dsg and AT2019fdr, misaligned angles less than 0.4 radians can meet the requirement of the observed neutrino flux. The most optimistic estimate of the neutrino number in the weakly misaligned jet scenario can reach  $N_\nu = 0.02$ ,  $N_\nu = 0.015$  and  $N_\nu = 0.12$  for AT2019dsg, AT2019fdr and AT2019aalc respectively.

### 3.2. Diffuse neutrino emission from Choked TDE jets

If choked precessing jets are common in normal TDEs, neutrinos from these choked precessing jets may contribute to the all-sky diffuse neutrino background. A rough estimate of the diffuse muon neutrino emission from these jets, using the above constraints and the updated event rate of normal TDEs (Yao et al. 2023), can be obtained by

$$\begin{aligned} \varepsilon_\nu^2 \Phi_\nu &\approx \frac{c}{4\pi H_0} f_z E_\nu \rho_0 \\ &\approx 3 \times 10^{-9} f_z f_{p\gamma} E_{p,51} \left( \frac{\rho_0}{310 \text{Gpc}^{-3} \text{yr}^{-1}} \right) \quad (14) \\ &\text{GeV cm}^{-2} \text{s}^{-1} \text{sr}^{-1}, \end{aligned}$$

where  $E_\nu = f_{p\gamma} E_p / 8 \ln(\epsilon_{p,\max}/\epsilon_{p,\min})$  is the muon neutrino energy released by a single TDE event,  $E_p$  is total CR proton energy injected by one TDE event,  $\rho_0$  is the local event rate of TDEs,  $H_0$  is the Hubble constant, and  $f_z$  is the correction factor for the contribution from high redshift TDEs, which is uncertain from current observations. For the TDE event rates, a latest demography study suggested that the local TDE event rate is  $\rho_0 \simeq 310 \text{Gpc}^{-3} \text{yr}^{-1}$  (Yao et al. 2023). A typical solar mass TDE can eject half of the mass into the disk wind at a velocity  $v_w \simeq 0.1c$ (Metzger & Stone 2016), leading to a kinetic energy of  $E_k \approx 2.3 \times 10^{51} (f_w/0.5)(v_w/0.1c)^2$  erg in the disk wind. Noting that this energy does not violate the constraints from radio observations of some TDEs, as the ejecta

is still in the free-expansion phase(Matsumoto & Piran 2021). Radio observations show that the kinetic energy of the jets in Swift J1644+57 and AT2022cmc is about  $E_k \sim 10^{51} - 10^{52}$  erg (Cendes et al. 2023). This motivated us to assume that the kinetic energy of the choked jet is comparable to the kinetic energy of the disk wind, so the CR proton energy in one TDE event is  $E_p \sim 4.5 \times 10^{50} (v_w/0.1c)^2$  erg if taking  $\epsilon_p = 0.2$ . Assuming  $f_{p\gamma} \simeq 1$  and  $f_z \simeq 1$ , the single flavor neutrino flux contributed by choked TDE jets is approximately  $\sim 10^{-9} \text{GeV cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$ , which is about  $\sim 10\%$  of the diffuse neutrino flux at 30-300 TeV (Abbasi et al. 2022).

## 4. SUMMARY AND DISCUSSIONS

We revisited the jet break-out condition for misaligned precessing jets in TDEs by considering that the jet could be collimated by the cocoon pressure while the jet is propagating in the quasi-isotropic disk wind. We find that the collimation effect is important for choked precessing jets. As a result, the jet head opening angle decreases as the jet propagates in the envelope, and the isotropic jet luminosity  $L_{j,\text{iso}}$  increases correspondingly. Nevertheless, the break-out condition does not change if we consider the beam-corrected jet luminosity, which is simply  $L_j \geq \frac{1}{2} \dot{M} f_w c^2 \beta_w \pi^2 \sin^2 \theta_{LS}$ . Thus, for known values of  $f_w$ ,  $\beta_w$  and  $\theta_{LS}$ , the break-out condition depends only on the jet efficiency  $\eta = L_j / \dot{M} c^2$ . For typical parameter values of  $f_w = 0.5$ ,  $\beta_w = 0.1$  and  $\theta_{LS} = \pi/4$ , successful jets imply  $\eta > 10^{-1}$ . Such high-efficiency jets may not be common in TDEs, so the majority of TDE jets are choked, explaining the small fraction of TDEs that have powerful jets. At later times, the jets could possibly break out if the disk finally aligns itself with the spin axis due to the damping of the precession (Teboul & Metzger 2023; Lu et al. 2023).

We further study the neutrino emission from choked precessing jets. Protons accelerated by shocks inside the choked precessing jet interact with the dense thermal photons in the wind envelope and produce high-energy neutrinos. In Fig.5 and Fig.6 we show the cumulative neutrino numbers for three TDEs in the parameter space of  $\theta_0 - L_j$  and  $\theta_{LS} - L_j$ . Although precession allows a much larger jet luminosity before break-out (compared with the aligned jet case discussed in Zheng et al. (2023)), the real physical jet power can not be so large, otherwise the deposited energy by choked jets would exceed the wind power. For the same jet luminosity, the observed neutrino flux from a choked precessing jet is reduced by a factor represented by the duty cycle due to precession, which is  $\theta_0 / (\pi \sin \theta_{LS})$ . We propose that the neutrinos from AT2019dsg, AT2019fdr are more likely to be produced by weakly misaligned choked jets,

and neutrinos from AT2019aal are probably emitted by misaligned choked precessing jets. We also calculate the all-sky diffuse neutrino flux from all choked TDE jets and find that it constitutes a fraction of  $\sim 10\%$  of the total neutrino background flux.

In this work, we assumed that the energy of the choked jet is limited by the kinetic energy of the disk wind. However, it is unknown how the deposited energy by

choked jets affects the wind envelope and the radiation. A detailed numerical study is needed to verify this assumption. If the maximally allowed energy for the choked jets is larger, the neutrinos flux will become larger correspondingly.

- 1 We thank Wenbing Lu and Weihua Lei for valuable discussions. The work is supported by the NSFC under
- 2 grants Nos. 12333006, 12121003 and 12393852.
- 3

## REFERENCES

- Abbasi, R., Ackermann, M., Adams, J., et al. 2022, *ApJ*, 928, 50, doi: [10.3847/1538-4357/ac4d29](https://doi.org/10.3847/1538-4357/ac4d29)
- Andreoni, I., Coughlin, M. W., Perley, D. A., et al. 2022, *Nature*, 612, 430, doi: [10.1038/s41586-022-05465-8](https://doi.org/10.1038/s41586-022-05465-8)
- Blaufuss, E., Kintscher, T., Lu, L., & Tung, C. F. 2019, in *International Cosmic Ray Conference*, Vol. 36, 36th International Cosmic Ray Conference (ICRC2019), 1021. <https://arxiv.org/abs/1908.04884>
- Bloom, J. S., Giannios, D., Metzger, B. D., et al. 2011, *Science*, 333, 203, doi: [10.1126/science.1207150](https://doi.org/10.1126/science.1207150)
- Bromberg, O., Nakar, E., Piran, T., & Sari, R. 2011, *ApJ*, 740, 100, doi: [10.1088/0004-637X/740/2/100](https://doi.org/10.1088/0004-637X/740/2/100)
- Brown, G. C., Levan, A. J., Stanway, E. R., et al. 2015, *MNRAS*, 452, 4297, doi: [10.1093/mnras/stv1520](https://doi.org/10.1093/mnras/stv1520)
- Burrows, D. N., Kennea, J. A., Ghisellini, G., et al. 2011, *Nature*, 476, 421, doi: [10.1038/nature10374](https://doi.org/10.1038/nature10374)
- Cendes, Y., Berger, E., Alexander, K. D., et al. 2023, arXiv e-prints, arXiv:2308.13595, doi: [10.48550/arXiv.2308.13595](https://doi.org/10.48550/arXiv.2308.13595)
- Cenko, S. B., Krimm, H. A., Horesh, A., et al. 2012, *ApJ*, 753, 77, doi: [10.1088/0004-637X/753/1/77](https://doi.org/10.1088/0004-637X/753/1/77)
- Farrar, G. R., & Gruzinov, A. 2009, *ApJ*, 693, 329, doi: [10.1088/0004-637X/693/1/329](https://doi.org/10.1088/0004-637X/693/1/329)
- Gezari, S. 2021, *ARA&A*, 59, 21, doi: [10.1146/annurev-astro-111720-030029](https://doi.org/10.1146/annurev-astro-111720-030029)
- Gottlieb, O., & Nakar, E. 2022, *MNRAS*, 517, 1640, doi: [10.1093/mnras/stac2699](https://doi.org/10.1093/mnras/stac2699)
- Hamidani, H., Kiuchi, K., & Ioka, K. 2020, *MNRAS*, 491, 3192, doi: [10.1093/mnras/stz3231](https://doi.org/10.1093/mnras/stz3231)
- Hayasaki, K., & Yamazaki, R. 2019, *ApJ*, 886, 114, doi: [10.3847/1538-4357/ab44ca](https://doi.org/10.3847/1538-4357/ab44ca)
- Hung, T., Cenko, S. B., Roth, N., et al. 2019, *ApJ*, 879, 119, doi: [10.3847/1538-4357/ab24de](https://doi.org/10.3847/1538-4357/ab24de)
- Ioka, K., & Nakamura, T. 2018, *Progress of Theoretical and Experimental Physics*, 2018, 043E02, doi: [10.1093/ptep/pty036](https://doi.org/10.1093/ptep/pty036)
- Krolik, J. H., & Piran, T. 2012, *ApJ*, 749, 92, doi: [10.1088/0004-637X/749/1/92](https://doi.org/10.1088/0004-637X/749/1/92)
- Liu, R.-Y., Xi, S.-Q., & Wang, X.-Y. 2020, *PhRvD*, 102, 083028, doi: [10.1103/PhysRevD.102.083028](https://doi.org/10.1103/PhysRevD.102.083028)
- Lu, W., Matsumoto, T., & Matzner, C. D. 2023, arXiv e-prints, arXiv:2310.15336, doi: [10.48550/arXiv.2310.15336](https://doi.org/10.48550/arXiv.2310.15336)
- Matsumoto, T., & Piran, T. 2021, *MNRAS*, 507, 4196, doi: [10.1093/mnras/stab2418](https://doi.org/10.1093/mnras/stab2418)
- Matzner, C. D. 2003, *MNRAS*, 345, 575, doi: [10.1046/j.1365-8711.2003.06969.x](https://doi.org/10.1046/j.1365-8711.2003.06969.x)
- Metzger, B. D., & Stone, N. C. 2016, *MNRAS*, 461, 948, doi: [10.1093/mnras/stw1394](https://doi.org/10.1093/mnras/stw1394)
- Mukhopadhyay, M., Bhattacharya, M., & Murase, K. 2023, arXiv e-prints, arXiv:2309.02275, doi: [10.48550/arXiv.2309.02275](https://doi.org/10.48550/arXiv.2309.02275)
- Murase, K., Kimura, S. S., Zhang, B. T., Oikonomou, F., & Petropoulou, M. 2020, *ApJ*, 902, 108, doi: [10.3847/1538-4357/abb3c0](https://doi.org/10.3847/1538-4357/abb3c0)
- Piran, T., Sądowski, A., & Tchekhovskoy, A. 2015, *MNRAS*, 453, 157, doi: [10.1093/mnras/stv1547](https://doi.org/10.1093/mnras/stv1547)
- Roth, N., Kasen, D., Guillochon, J., & Ramirez-Ruiz, E. 2016, *ApJ*, 827, 3, doi: [10.3847/0004-637X/827/1/3](https://doi.org/10.3847/0004-637X/827/1/3)
- Sazonov, S., Gilfanov, M., Medvedev, P., et al. 2021, *MNRAS*, 508, 3820, doi: [10.1093/mnras/stab2843](https://doi.org/10.1093/mnras/stab2843)
- Stein, R., van Velzen, S., Kowalski, M., et al. 2021, *Nature Astronomy*, 5, 510, doi: [10.1038/s41550-020-01295-8](https://doi.org/10.1038/s41550-020-01295-8)
- Teboul, O., & Metzger, B. D. 2023, *ApJL*, 957, L9, doi: [10.3847/2041-8213/ad0037](https://doi.org/10.3847/2041-8213/ad0037)
- van Velzen, S. 2018, *ApJ*, 852, 72, doi: [10.3847/1538-4357/aa998e](https://doi.org/10.3847/1538-4357/aa998e)
- van Velzen, S., Gezari, S., Hammerstein, E., et al. 2021, *ApJ*, 908, 4, doi: [10.3847/1538-4357/abc258](https://doi.org/10.3847/1538-4357/abc258)
- Wang, X.-Y., & Liu, R.-Y. 2016, *PhRvD*, 93, 083005, doi: [10.1103/PhysRevD.93.083005](https://doi.org/10.1103/PhysRevD.93.083005)
- Wang, X.-Y., Liu, R.-Y., Dai, Z.-G., & Cheng, K. S. 2011, *PhRvD*, 84, 081301, doi: [10.1103/PhysRevD.84.081301](https://doi.org/10.1103/PhysRevD.84.081301)
- Winter, W., & Lunardini, C. 2021, *Nature Astronomy*, 5, 472, doi: [10.1038/s41550-021-01305-3](https://doi.org/10.1038/s41550-021-01305-3)
- . 2023, *ApJ*, 948, 42, doi: [10.3847/1538-4357/acbe9e](https://doi.org/10.3847/1538-4357/acbe9e)

Wu, H.-J., Mou, G., Wang, K., Wang, W., & Li, Z. 2022, MNRAS, 514, 4406, doi: [10.1093/mnras/stac1621](https://doi.org/10.1093/mnras/stac1621)  
Yao, Y., Ravi, V., Gezari, S., et al. 2023, ApJL, 955, L6, doi: [10.3847/2041-8213/acf216](https://doi.org/10.3847/2041-8213/acf216)

Zheng, J.-H., Liu, R.-Y., & Wang, X.-Y. 2023, ApJ, 954, 17, doi: [10.3847/1538-4357/ace71c](https://doi.org/10.3847/1538-4357/ace71c)