

**COMPACT MINIMAL SUBMANIFOLDS OF
THE RIEMANNIAN SYMMETRIC SPACES
 $\mathbf{SU}(n)/\mathbf{SO}(n)$, $\mathbf{Sp}(n)/\mathbf{U}(n)$, $\mathbf{SO}(2n)/\mathbf{U}(n)$, $\mathbf{SU}(2n)/\mathbf{Sp}(n)$
VIA COMPLEX-VALUED EIGENFUNCTIONS**

JOHANNA MARIE GEGENFURTNER AND SIGMUNDUR GUDMUNDSSON

ABSTRACT. In this work we construct new multi-dimensional families of compact minimal submanifolds, of the classical Riemannian symmetric spaces $\mathbf{SU}(n)/\mathbf{SO}(n)$, $\mathbf{Sp}(n)/\mathbf{U}(n)$, $\mathbf{SO}(2n)/\mathbf{U}(n)$ and $\mathbf{SU}(2n)/\mathbf{Sp}(n)$, of codimension two.

1. INTRODUCTION

The study of minimal submanifolds of a given ambient space plays a central role in differential geometry. This has a long, interesting history and has attracted the interests of profound mathematicians for many generations. The famous Weierstrass-Enneper representation formula, for minimal surfaces in Euclidean 3-space, brings *complex analysis* into play as a useful tool for the study of these beautiful objects.

This was later generalised to the study of minimal surfaces in much more general ambient manifolds via *conformal harmonic immersions*. The next result follows from the seminal paper [6] of Eells and Sampson from 1964. For this see also Proposition 3.5.1 of [2].

Theorem 1.1. *Let $\phi : (M^m, g) \rightarrow (N, h)$ be a smooth conformal map between Riemannian manifolds. If $m = 2$ then ϕ is harmonic if and only if the image is minimal in (N, h) .*

This result has turned out to be very useful in the construction of minimal surfaces in Riemannian symmetric spaces of various types. For this we refer to [5], [7], [19], [3] and [4], just to name a few.

In their work [1] from 1981, Baird and Eells have shown that complex-valued harmonic morphisms from Riemannian manifolds are useful tools for the study of minimal submanifolds of codimension two. This can be seen as dual to the above mentioned generalisation of the Weierstrass-Enneper representation. Harmonic morphisms are the much studied *horizontally conformal* harmonic maps. For an introduction to the theory of harmonic morphisms we recommend the book [2], by Baird and Wood, and the regularly updated online bibliography [11].

2020 *Mathematics Subject Classification.* 53C35, 53C43, 58E20.

Key words and phrases. minimal submanifolds, eigenfunctions, symmetric spaces.

Theorem 1.2. [1] *Let $\phi : (M, g) \rightarrow \mathbb{C}$ be a complex-valued harmonic morphism from a Riemannian manifold. Then every regular fibre of ϕ is a minimal submanifold of (M, g) of codimension two.*

The recent work [13], introduces a method for constructing minimal submanifolds of Riemannian manifolds, via *submersions*, see Theorem 3.1. Then this scheme is employed to provide compact examples in several important cases.

The main ingredients for this new procedure are the so called *complex-valued eigenfunctions* on the Riemannian ambient space. These are functions which are eigen both with respect to the classical *Laplace-Beltrami* and the so called *conformality* operator, see Section 2.

In the current study we continue the investigation and apply the new above mentioned method to the classical Riemannian symmetric spaces

$$\mathbf{SU}(n)/\mathbf{SO}(n), \mathbf{Sp}(n)/\mathbf{U}(n), \mathbf{SO}(2n)/\mathbf{U}(n), \mathbf{SU}(2n)/\mathbf{Sp}(n).$$

In these cases we find multi-dimensional families of compact minimal submanifolds of codimension two, see Theorems 8.2, 9.2, 10.2 and 11.2. Readers interested in further details are referred to [8].

2. EIGENFUNCTIONS AND EIGENFAMILIES

Let (M, g) be an m -dimensional Riemannian manifold and $T^{\mathbb{C}}M$ be the complexification of the tangent bundle TM of M . We extend the metric g to a complex bilinear form on $T^{\mathbb{C}}M$. Then the gradient $\nabla\phi$ of a complex-valued function $\phi : (M, g) \rightarrow \mathbb{C}$ is a section of $T^{\mathbb{C}}M$. In this situation, the well-known complex linear *Laplace-Beltrami operator* (alt. *tension field*) τ on (M, g) acts locally on ϕ as follows

$$\tau(\phi) = \operatorname{div}(\nabla\phi) = \sum_{i,j=1}^m \frac{1}{\sqrt{|g|}} \frac{\partial}{\partial x_j} \left(g^{ij} \sqrt{|g|} \frac{\partial \phi}{\partial x_i} \right).$$

For two complex-valued functions $\phi, \psi : (M, g) \rightarrow \mathbb{C}$ we have the following well-known fundamental relation

$$\tau(\phi\psi) = \tau(\phi)\psi + 2\kappa(\phi, \psi) + \phi\tau(\psi),$$

where the complex bilinear *conformality operator* κ is given by

$$\kappa(\phi, \psi) = g(\nabla\phi, \nabla\psi).$$

Locally this satisfies

$$\kappa(\phi, \psi) = \sum_{i,j=1}^m g^{ij} \cdot \frac{\partial \phi}{\partial x_i} \frac{\partial \psi}{\partial x_j}.$$

Definition 2.1. [14] Let (M, g) be a Riemannian manifold. Then a complex-valued function $\phi : M \rightarrow \mathbb{C}$ is said to be a (λ, μ) -*eigenfunction* if it is eigen

both with respect to the Laplace-Beltrami operator τ and the conformality operator κ i.e. there exist complex numbers $\lambda, \mu \in \mathbb{C}$ such that

$$\tau(\phi) = \lambda \cdot \phi \quad \text{and} \quad \kappa(\phi, \phi) = \mu \cdot \phi^2.$$

A set $\mathcal{E} = \{\phi_i : M \rightarrow \mathbb{C} \mid i \in I\}$ of complex-valued functions is said to be a (λ, μ) -*eigenfamily* on M if there exist complex numbers $\lambda, \mu \in \mathbb{C}$ such that for all $\phi, \psi \in \mathcal{E}$ we have

$$\tau(\phi) = \lambda \cdot \phi \quad \text{and} \quad \kappa(\phi, \psi) = \mu \cdot \phi \psi.$$

For the standard odd-dimensional round spheres we have the following eigenfamilies based on the classical real-valued spherical harmonics.

Example 2.2. [13] Let S^{2n-1} be the odd-dimensional unit sphere in the standard Euclidean space $\mathbb{C}^n \cong \mathbb{R}^{2n}$ and define $\phi_1, \dots, \phi_n : S^{2n-1} \rightarrow \mathbb{C}$ by

$$\phi_j : (z_1, \dots, z_n) \mapsto \frac{z_j}{\sqrt{|z_1|^2 + \dots + |z_n|^2}}.$$

Then the tension field τ and the conformality operator κ on S^{2n-1} satisfy

$$\tau(\phi_j) = -(2n-1) \cdot \phi_j \quad \text{and} \quad \kappa(\phi_j, \phi_k) = -1 \cdot \phi_j \cdot \phi_k.$$

For the standard complex projective space $\mathbb{C}P^n$ we have a complex multi-dimensional eigenfamily, in a similar way.

Example 2.3. [13] Let $\mathbb{C}P^n$ be the standard n -dimensional complex projective space. For a fixed integer $1 \leq \alpha < n+1$ and some $1 \leq j \leq \alpha < k \leq n+1$ define the function $\phi_{jk} : \mathbb{C}P^n \rightarrow \mathbb{C}$ by

$$\phi_{jk} : [z_1, \dots, z_{n+1}] \mapsto \frac{z_j \cdot \bar{z}_k}{z_1 \cdot \bar{z}_1 + \dots + z_{n+1} \cdot \bar{z}_{n+1}}.$$

Then the tension field τ and the conformality operator κ on $\mathbb{C}P^n$ satisfy

$$\tau(\phi_{jk}) = -4(n+1) \cdot \phi_{jk} \quad \text{and} \quad \kappa(\phi_{jk}, \phi_{lm}) = -4 \cdot \phi_{jk} \cdot \phi_{lm}.$$

In recent years, explicit eigenfamilies of complex-valued functions have been found on all the classical compact Riemannian symmetric spaces. For this see Table 1.

The next result shows that a given eigenfamily \mathcal{E} on (M, g) induces a large collection $\mathcal{P}_d(\mathcal{E})$ of such objects. This is particularly interesting in the light of Table 1.

Theorem 2.4. [10] *Let (M, g) be a Riemannian manifold and the set of complex-valued functions*

$$\mathcal{E} = \{\phi_i : M \rightarrow \mathbb{C} \mid i = 1, 2, \dots, n\}$$

be a finite eigenfamily i.e. there exist complex numbers $\lambda, \mu \in \mathbb{C}$ such that for all $\phi, \psi \in \mathcal{E}$

$$\tau(\phi) = \lambda \cdot \phi \quad \text{and} \quad \kappa(\phi, \psi) = \mu \cdot \phi \psi.$$

U/K	λ	μ	Eigenfunctions
$\mathbf{SO}(n)$	$-\frac{(n-1)}{2}$	$-\frac{1}{2}$	see [14]
$\mathbf{SU}(n)$	$-\frac{n^2-1}{n}$	$-\frac{n-1}{n}$	see [16]
$\mathbf{Sp}(n)$	$-\frac{2n+1}{2}$	$-\frac{1}{2}$	see [12]
$\mathbf{SU}(n)/\mathbf{SO}(n)$	$-\frac{2(n^2+n-2)}{n}$	$-\frac{4(n-1)}{n}$	see [15]
$\mathbf{Sp}(n)/\mathbf{U}(n)$	$-2(n+1)$	-2	see [15]
$\mathbf{SO}(2n)/\mathbf{U}(n)$	$-2(n-1)$	-1	see [15]
$\mathbf{SU}(2n)/\mathbf{Sp}(n)$	$-\frac{2(2n^2-n-1)}{n}$	$-\frac{2(n-1)}{n}$	see [15]
$\mathbf{SO}(m+n)/\mathbf{SO}(m) \times \mathbf{SO}(n)$	$-(m+n)$	-2	see [9]
$\mathbf{U}(m+n)/\mathbf{U}(m) \times \mathbf{U}(n)$	$-2(m+n)$	-2	see [10]
$\mathbf{Sp}(m+n)/\mathbf{Sp}(m) \times \mathbf{Sp}(n)$	$-2(m+n)$	-1	see [10]

TABLE 1. Eigenfamilies on the classical compact irreducible Riemannian symmetric spaces.

Then the set of complex homogeneous polynomials of degree d

$$\mathcal{P}_d(\mathcal{E}) = \{P : M \rightarrow \mathbb{C} \mid P \in \mathbb{C}[\phi_1, \phi_2, \dots, \phi_n], P(\alpha \cdot \phi) = \alpha^d \cdot P(\phi), \alpha \in \mathbb{C}\}$$

is an eigenfamily on M such that for all $P, Q \in \mathcal{P}_d(\mathcal{E})$ we have

$$\tau(P) = (d\lambda + d(d-1)\mu) \cdot P \quad \text{and} \quad \kappa(P, Q) = d^2\mu \cdot P \cdot Q.$$

We conclude this section with the following two results, particularly useful in the above mentioned situations of compact Riemannian symmetric spaces.

Proposition 2.5. *Let $\pi : (\hat{M}, \hat{g}) \rightarrow (M, g)$ be a Riemannian submersion between Riemannian manifolds. Further let $\phi : (M, g) \rightarrow \mathbb{C}$ be a smooth function and $\hat{\phi} : (\hat{M}, \hat{g}) \rightarrow \mathbb{C}$ be the composition $\hat{\phi} = \phi \circ \pi$. Then the*

corresponding tension fields τ and conformality operators κ satisfy

$$\tau(\hat{\phi}) = \tau(\phi) \circ \pi \quad \text{and} \quad \kappa(\hat{\phi}, \hat{\psi}) = \kappa(\phi, \psi) \circ \pi.$$

Proof. The arguments needed here can be found in [15] □

In the sequel, we shall apply the following immediate consequence of Proposition 2.5.

Corollary 2.6. *Let $\pi : (\hat{M}, \hat{g}) \rightarrow (M, g)$ be a Riemannian submersion. For a complex-valued function smooth function $\phi : (M, g) \rightarrow \mathbb{C}$ let $\hat{\phi} : (\hat{M}, \hat{g}) \rightarrow \mathbb{C}$ be the composition $\hat{\phi} = \phi \circ \pi$. Then the following statements are equivalent*

- (i) $\phi : M \rightarrow \mathbb{C}$ is a (λ, μ) -eigenfunction on M ,
- (ii) $\hat{\phi} : \hat{M} \rightarrow \mathbb{C}$ is a (λ, μ) -eigenfunction on \hat{M} .

3. MINIMAL SUBMANIFOLDS VIA EIGENFUNCTIONS

The recent paper [13] provides a new application of complex-valued eigenfunctions. This is a method for constructing minimal submanifolds of codimension two.

Theorem 3.1. [13] *Let $\phi : (M, g) \rightarrow \mathbb{C}$ be a complex-valued eigenfunction on a Riemannian manifold, such that $0 \in \phi(M)$ is a regular value for ϕ . Then the fibre $\phi^{-1}(\{0\})$ is a minimal submanifold of M of codimension two.*

The main aim of our work is to apply Theorem 3.1 in several of the interesting cases when the Riemannian manifold (M, g) is one of the classical compact symmetric spaces.

The next result, from Riedler and Siffert's paper [18] supplies us with a straightforward way of checking whether an eigenfunction, on a compact and connected Riemannian manifold, attains the required value $0 \in \mathbb{C}$.

Theorem 3.2. [18] *Let (M, g) be a compact and connected Riemannian manifold and let $\phi : M \rightarrow \mathbb{C}$ be a (λ, μ) -eigenfunction not identically zero. Then the following are equivalent.*

- (1) $\lambda = \mu$.
- (2) $|\phi|^2$ is constant.
- (3) $\phi(x) \neq 0$ for all $x \in M$.

As an obvious consequence we have the following.

Corollary 3.3. *If $\phi : M \rightarrow \mathbb{C}$ is a complex-valued (λ, μ) -eigenfunction on a compact and connected Riemannian manifold (M, g) such that $\lambda \neq \mu$, then there exists $x \in M$ such that $\phi(x) = 0$.*

4. THE GENERAL LINEAR GROUP $\mathbf{GL}_n(\mathbb{C})$

In this section we now turn our attention to the concrete Riemannian matrix Lie groups embedded as subgroups of the complex general linear group.

The group of linear automorphism of \mathbb{C}^n is the complex general linear group $\mathbf{GL}_n(\mathbb{C}) = \{z \in \mathbb{C}^{n \times n} \mid \det z \neq 0\}$ of invertible $n \times n$ matrices with its standard representation

$$z \mapsto \begin{bmatrix} z_{11} & \cdots & z_{1n} \\ \vdots & \ddots & \vdots \\ z_{n1} & \cdots & z_{nn} \end{bmatrix}.$$

Its Lie algebra $\mathfrak{gl}_n(\mathbb{C})$ of left-invariant vector fields on $\mathbf{GL}_n(\mathbb{C})$ can be identified with $\mathbb{C}^{n \times n}$ i.e. the complex linear space of $n \times n$ matrices. We equip $\mathbf{GL}_n(\mathbb{C})$ with its natural left-invariant Riemannian metric g induced by the standard Euclidean inner product $\mathfrak{gl}_n(\mathbb{C}) \times \mathfrak{gl}_n(\mathbb{C}) \rightarrow \mathbb{R}$ on its Lie algebra $\mathfrak{gl}_n(\mathbb{C})$ satisfying

$$g(Z, W) \mapsto \Re \text{ trace}(Z \cdot \bar{W}^t).$$

For $1 \leq i, j \leq n$, we shall by E_{ij} denote the element of $\mathbb{R}^{n \times n}$ satisfying

$$(E_{ij})_{kl} = \delta_{ik} \delta_{jl}$$

and by D_t the diagonal matrices $D_t = E_{tt}$. For $1 \leq r < s \leq n$, let X_{rs} and Y_{rs} be the matrices satisfying

$$X_{rs} = \frac{1}{\sqrt{2}}(E_{rs} + E_{sr}), \quad Y_{rs} = \frac{1}{\sqrt{2}}(E_{rs} - E_{sr}).$$

For the real vector space $\mathfrak{gl}_n(\mathbb{C})$ we then have the canonical orthonormal basis $\mathcal{B}^{\mathbb{C}} = \mathcal{B} \cup i\mathcal{B}$, where

$$\mathcal{B} = \{Y_{rs}, X_{rs} \mid 1 \leq r < s \leq n\} \cup \{D_t \mid t = 1, 2, \dots, n\}.$$

Let G be a classical Lie subgroup of $\mathbf{GL}_n(\mathbb{C})$ with Lie algebra \mathfrak{g} inheriting the induced left-invariant Riemannian metric, which we shall also denote by g . In the cases considered in this paper, $\mathcal{B}_{\mathfrak{g}} = \mathcal{B}^{\mathbb{C}} \cap \mathfrak{g}$ will be an orthonormal basis for the subalgebra \mathfrak{g} of $\mathfrak{gl}_n(\mathbb{C})$. By employing the Koszul formula for the Levi-Civita connection ∇ on (G, g) , we see that for all $Z, W \in \mathcal{B}_{\mathfrak{g}}$ we have

$$\begin{aligned} g(\nabla_Z Z, W) &= g([W, Z], Z) \\ &= \Re \text{ trace}((WZ - ZW)\bar{Z}^t) \\ &= \Re \text{ trace}(W(Z\bar{Z}^t - \bar{Z}^t Z)) \\ &= 0. \end{aligned}$$

If $Z \in \mathfrak{g}$ is a left-invariant vector field on G and $\phi : U \rightarrow \mathbb{C}$ is a local complex-valued function on G then the k -th order derivatives $Z^k(\phi)$ satisfy

$$Z^k(\phi)(p) = \frac{d^k}{ds^k}(\phi(p \cdot \exp(sZ))) \Big|_{s=0},$$

This implies that the tension field τ and the conformality operator κ on G fulfill

$$\tau(\phi) = \sum_{Z \in \mathcal{B}_{\mathfrak{g}}} (Z^2(\phi) - \nabla_Z Z(\phi)) = \sum_{Z \in \mathcal{B}_{\mathfrak{g}}} Z^2(\phi),$$

$$\kappa(\phi, \psi) = \sum_{Z \in \mathcal{B}_{\mathfrak{g}}} Z(\phi) \cdot Z(\psi),$$

where $\mathcal{B}_{\mathfrak{g}}$ is the orthonormal basis $\mathcal{B}^{\mathbb{C}} \cap \mathfrak{g}$ for the Lie algebra \mathfrak{g} .

5. THE SPECIAL ORTHOGONAL GROUP $\mathbf{SO}(n)$

In this section we introduce the basic tools, related to the special orthogonal group $\mathbf{SO}(n)$, needed for our later constructions. The classical Lie group $\mathbf{SO}(n)$ is given by

$$\mathbf{SO}(n) = \{x \in \mathbf{GL}_n(\mathbb{R}) \mid x \cdot x^t = I_n, \det x = 1\}.$$

For this we use its standard n -dimensional representation on \mathbb{C}^n with

$$x \mapsto \begin{bmatrix} x_{11} & \cdots & x_{1n} \\ \vdots & \ddots & \vdots \\ x_{n1} & \cdots & x_{nn} \end{bmatrix}.$$

The Lie algebra $\mathfrak{so}(n)$ of $\mathbf{SO}(n)$ is the set of real skew-symmetric matrices

$$\mathfrak{so}(n) = \{X \in \mathfrak{gl}_n(\mathbb{R}) \mid X + X^t = 0\}$$

and for this we have the canonical orthonormal basis

$$\mathcal{B}_{\mathfrak{so}(n)} = \{Y_{rs} \mid 1 \leq r < s \leq n\}.$$

The gradient $\nabla \phi$ of a complex-valued function $\phi : \mathbf{SO}(n) \rightarrow \mathbb{C}$ is an element of the complexified tangent bundle $T^{\mathbb{C}}\mathbf{SO}(n)$. This satisfies

$$\nabla \phi = \sum_{Y \in \mathcal{B}_{\mathfrak{so}(n)}} Y(\phi) \cdot Y.$$

We now consider the functions $x_{j\alpha} : \mathbf{SO}(n) \rightarrow \mathbb{C}$ with $x_{j\alpha} : x \mapsto e_j \cdot x \cdot e_{\alpha}^t$. For any tangent vector $Y \in B_{\mathbf{u}(n)}$ we have

$$Y(x_{j\alpha}) : x \mapsto e_j \cdot x \cdot Y \cdot e_{\alpha}^t = \sum_{k=1}^n x_{jk} \cdot Y_{k\alpha}.$$

This implies that

$$Y_{rs}(x_{j\alpha}) = \frac{1}{\sqrt{2}} (x_{jr} \cdot \delta_{\alpha s} - x_{js} \cdot \delta_{\alpha r}). \quad (5.1)$$

6. THE UNITARY GROUPS $\mathbf{U}(n)$ AND $\mathbf{SU}(n)$

The unitary group $\mathbf{U}(n)$ is the compact subgroup of $\mathbf{GL}_n(\mathbb{C})$ given by

$$\mathbf{U}(n) = \{z \in \mathbb{C}^{n \times n} \mid z \cdot \bar{z}^t = I\}.$$

For its standard complex representation $\mathbf{U}(n) \rightarrow \mathbf{GL}_n(\mathbb{C})$ on \mathbb{C}^n we use the notation

$$z \mapsto \begin{bmatrix} z_{11} & \cdots & z_{1n} \\ \vdots & \ddots & \vdots \\ z_{n1} & \cdots & z_{nn} \end{bmatrix}.$$

The Lie algebra $\mathfrak{u}(n)$ of $\mathbf{U}(n)$ consists of the skew-Hermitian matrices i.e.

$$\mathfrak{u}(n) = \{Z \in \mathbb{C}^{n \times n} \mid Z + \bar{Z}^t = 0\}.$$

The left-invariant metric on $\mathbf{GL}_n(\mathbb{C})$ induces the standard biinvariant Riemannian metric g on $\mathbf{U}(n)$, with

$$g(Z, W) = \Re \operatorname{trace}(Z \cdot \bar{W}^t).$$

The canonical orthonormal basis for the Lie algebra $\mathfrak{u}(n)$ is then given by

$$\mathcal{B}_{\mathfrak{u}(n)} = \{Y_{rs}, iX_{rs} \mid 1 \leq r < s \leq n\} \cup \{iD_t \mid t = 1, \dots, n\}.$$

The special unitary group $\mathbf{SU}(n)$ is the subgroup of $\mathbf{U}(n)$ satisfying

$$\mathbf{SU}(n) = \{z \in \mathbf{U}(n) \mid \det(z) = 1\}$$

with Lie algebra

$$\mathfrak{su}(n) = \{Z \in \mathfrak{u}(n) \mid \operatorname{trace} Z = 0\}.$$

For the Lie algebra $\mathfrak{u}(n)$ of $\mathbf{U}(n)$ we have the orthogonal decomposition

$$\mathfrak{u}(n) = \mathfrak{su}(n) \oplus \mathfrak{m},$$

where the one dimensional subspace \mathfrak{m} is generated by the unit vector

$$E = i I_n / \sqrt{n}.$$

7. THE QUATERNIONIC UNITARY GROUP $\mathbf{Sp}(n)$

The group $\mathbf{Sp}(n)$ is the intersection of the unitary group $\mathbf{U}(2n)$ and the standard representation of the quaternionic general linear group $\mathbf{GL}_n(\mathbb{H})$ in $\mathbb{C}^{2n \times 2n}$ given by

$$(z + jw) \mapsto q = \begin{bmatrix} z & w \\ -\bar{w} & \bar{z} \end{bmatrix}. \quad (7.1)$$

The Lie algebra $\mathfrak{sp}(n)$ of $\mathbf{Sp}(n)$ satisfies

$$\mathfrak{sp}(n) = \left\{ \begin{bmatrix} Z & W \\ -\bar{W} & \bar{Z} \end{bmatrix} \in \mathbb{C}^{2n \times 2n} \mid Z^* + Z = 0, W^t - W = 0 \right\}.$$

We now introduce the following notation for the elements of the orthonormal basis $\mathcal{B}_{\mathfrak{sp}(n)}$ of the Lie algebra $\mathfrak{sp}(n)$:

$$\begin{aligned} Y_{rs}^a &= \frac{1}{\sqrt{2}} \begin{bmatrix} Y_{rs} & 0 \\ 0 & Y_{rs} \end{bmatrix}, X_{rs}^a = \frac{1}{\sqrt{2}} \begin{bmatrix} iX_{rs} & 0 \\ 0 & -iX_{rs} \end{bmatrix}, \\ X_{rs}^b &= \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & iX_{rs} \\ iX_{rs} & 0 \end{bmatrix}, X_{rs}^c = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & X_{rs} \\ -X_{rs} & 0 \end{bmatrix}, \\ D_t^a &= \frac{1}{\sqrt{2}} \begin{bmatrix} iD_t & 0 \\ 0 & -iD_t \end{bmatrix}, D_t^b = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & iD_t \\ iD_t & 0 \end{bmatrix}, D_t^c = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & D_t \\ -D_t & 0 \end{bmatrix}. \end{aligned}$$

Here $1 \leq r < s \leq n$ and $1 \leq t \leq n$.

8. THE SYMMETRIC SPACE $\mathbf{SU}(n)/\mathbf{SO}(n)$

The purpose of this section is to prove Theorem 8.2 and thereby construct a new multi-dimensional family of compact minimal submanifolds of the homogeneous quotient manifold $\mathbf{SU}(n)/\mathbf{SO}(n)$, which carries the structure of a compact Riemannian symmetric space. It is well-known that the natural projection $\pi : \mathbf{SU}(n) \rightarrow \mathbf{SU}(n)/\mathbf{SO}(n)$ from the special unitary group $\mathbf{SU}(n)$ is a Riemannian submersion. This means that we can apply Corollary 2.6 in this situation.

From the work [15], of Gudmundsson, Siffert and Sobak, we have the following construction of eigenfunctions.

Proposition 8.1. [15] *For $n \geq 2$, let a be a non-zero element of \mathbb{C}^n and $A \in \mathbb{C}^{n \times n}$ with $A = a^t a$. Further let $\phi_a : \mathbf{SU}(n)/\mathbf{SO}(n) \rightarrow \mathbb{C}$ be the complex-valued function induced by its $\mathbf{SO}(n)$ -invariant lift $\hat{\phi}_a : \mathbf{SU}(n) \rightarrow \mathbb{C}$ given by*

$$\hat{\phi}_a(z) = \text{trace}(Azz^t).$$

Then ϕ_a is an eigenfunction on $\mathbf{SU}(n)/\mathbf{SO}(n)$ with eigenvalues λ and μ satisfying

$$\lambda = -\frac{2(n^2 + n - 2)}{n} \quad \text{and} \quad \mu = -\frac{4(n - 1)}{n}.$$

With our next result we construct a complex $(n - 1)$ -dimensional family

$$\mathcal{F} = \{\phi_a^{-1}(\{0\}) \mid [a] \in \mathbb{C}P^{n-1}\}$$

of minimal submanifolds of $\mathbf{SU}(n)/\mathbf{SO}(n)$ of codimension two.

Theorem 8.2. *For $n \geq 2$, let a be a non-zero element of \mathbb{C}^n and $A \in \mathbb{C}^{n \times n}$ with $A = a^t a$. Further let $\phi_a : \mathbf{SU}(n)/\mathbf{SO}(n) \rightarrow \mathbb{C}$ be the complex-valued eigenfunction induced by its $\mathbf{SO}(n)$ -invariant lift $\hat{\phi}_a : \mathbf{SU}(n) \rightarrow \mathbb{C}$ given by*

$$\hat{\phi}_a(z) = \text{trace}(Azz^t).$$

Then ϕ_a is a submersion and the compact fibre $\phi_a^{-1}(\{0\})$ over $0 \in \mathbb{C}$ is a minimal submanifold of $\mathbf{SU}(n)/\mathbf{SO}(n)$ of codimension two.

Proof. For the Lie algebra $\mathfrak{su}(n)$ of $\mathbf{SU}(n)$ we have the orthogonal decomposition $\mathfrak{su}(n) = \mathfrak{so}(n) \oplus \mathfrak{ip}$ where the subspace

$$\mathfrak{p} = \{X \in \mathbb{R}^{n \times n} \mid X = X^t \text{ and } \text{trace } X = 0\}$$

is generated by the elements of

$$\{X_{rs}, D_{rs} \mid 1 \leq r < s \leq n\}.$$

This means that if a point $z \in \mathbf{SU}(n)$ is a critical point for $\hat{\phi} : \mathbf{SU}(n) \rightarrow \mathbb{C}$ then, for all $1 \leq r < s \leq n$, we have

$$\begin{aligned} X_{rs}(\hat{\phi}_a) &= \text{trace}(X_{rs}^t \cdot z^t \cdot A \cdot z + z^t \cdot A \cdot z \cdot X_{rs}) \\ &= \text{trace}((X_{rs}^t + X_{rs}) \cdot z^t \cdot a^t \cdot a \cdot z) \\ &= 2 \text{trace}(a \cdot z \cdot X_{rs} \cdot z^t \cdot a^t) \\ &= 2\sqrt{2} \left(\sum_{k=1}^n a_k z_{kr} \right) \left(\sum_{k=1}^n a_k z_{ks} \right) \\ &= 0. \end{aligned}$$

Hence, for any choice $1 \leq r < s \leq n$, either

$$\sum_{k=1}^n a_k z_{kr} = 0 \quad \text{or} \quad \sum_{k=1}^n a_k z_{ks} = 0. \quad (8.1)$$

Now there exists an integer $1 \leq t \leq n$ such that

$$\sum_{k=1}^n a_k z_{kt} \neq 0, \quad (8.2)$$

otherwise the rows of the matrix $z \in \mathbf{SU}(n)$ would be linearly dependent, which is impossible. It then follows from statement (8.1) that if $m \neq t$, then

$$\sum_{k=1}^n a_k z_{km} = 0.$$

Similarly, we see that at a critical point $z \in \mathbf{SU}(n)$ we have $D_{rs}(\phi) = 0$ for all $1 \leq r < s \leq n$. This is equivalent to

$$\left(\sum_{k=1}^n a_k z_{kr} + \sum_{k=1}^n a_k z_{ks} \right) \cdot \left(\sum_{k=1}^n a_k z_{kr} - \sum_{k=1}^n a_k z_{ks} \right) = 0.$$

Applying our previous findings, we see that

$$\sum_{k=1}^n a_k z_{kt} = 0,$$

which contradicts equation (8.2). We conclude that the eigenfunction $\phi_a : \mathbf{SU}(n)/\mathbf{SO}(n) \rightarrow \mathbb{C}$ has no critical points. According to Corollary 3.2, the origin $0 \in \mathbb{C}$ lies in the image of ϕ_a . It then immediately follows from Theorem 3.1 that the fibre $\phi_a^{-1}(\{0\})$ is minimal. \square

9. THE SYMMETRIC SPACE $\mathbf{Sp}(n)/\mathbf{U}(n)$

In this section we prove Theorem 9.2 and thereby construct a new multi-dimensional family of compact minimal submanifolds of the Riemannian symmetric space $\mathbf{Sp}(n)/\mathbf{U}(n)$. It is well-known that the natural projection $\pi : \mathbf{Sp}(n) \rightarrow \mathbf{Sp}(n)/\mathbf{U}(n)$ from the quaternionic unitary group $\mathbf{Sp}(n)$ is a Riemannian submersion. This means that we can employ Corollary 2.6 in this situation.

Here we use the standard representation of the quaternionic unitary group $\mathbf{Sp}(n)$ on \mathbb{C}^{2n} given by equation (7.1). Then we identify the unitary group with following subgroup of $\mathbf{Sp}(n)$

$$\mathbf{U}(n) \cong \left\{ \begin{bmatrix} x & y \\ -y & x \end{bmatrix} \in \mathbb{R}^{2n \times 2n} \mid xx^t + yy^t = I_n, \quad xy^t = yx^t \right\}.$$

For the Lie algebra $\mathfrak{sp}(n)$ of $\mathbf{Sp}(n)$ we have the orthogonal decomposition $\mathfrak{sp}(n) = \mathfrak{u}(n) \oplus \mathfrak{p}$ where the subspace \mathfrak{p} is generated by the elements of its following orthonormal basis

$$\begin{aligned} \mathcal{B}_{\mathfrak{p}} = & \left\{ X_{rs}^a = \frac{i}{2} \begin{bmatrix} X_{rs} & 0 \\ 0 & -X_{rs} \end{bmatrix}, D_t^a = \frac{i}{\sqrt{2}} \begin{bmatrix} D_t & 0 \\ 0 & -D_t \end{bmatrix}, \right. \\ & X_{rs}^b = \frac{i}{2} \begin{bmatrix} 0 & X_{rs} \\ X_{rs} & 0 \end{bmatrix}, D_t^b = \frac{i}{\sqrt{2}} \begin{bmatrix} 0 & D_t \\ D_t & 0 \end{bmatrix} \\ & \left. \mid 1 \leq r < s \leq n, 1 \leq t \leq n \right\}. \end{aligned}$$

Let $\sigma : \mathbf{Sp}(n) \rightarrow \mathbf{Sp}(n)$ be the map given by $\sigma(q) = q q^t$. Then it is easily seen that σ is $\mathbf{U}(n)$ -invariant i.e. for all $q \in \mathbf{Sp}(n)$ and $z \in \mathbf{U}(n)$ we have

$$\sigma(qz) = qz z^t q^t = q q^t = \sigma(q).$$

The following result on eigenfunctions on the quotient space $\mathbf{Sp}(n)/\mathbf{U}(n)$ can be found in [15].

Proposition 9.1. [15] *Let a be a non-zero element of \mathbb{C}^{2n} and $A \in \mathbb{C}^{2n \times 2n}$ with $A = a^t a$. Further let $\phi_a : \mathbf{Sp}(n)/\mathbf{U}(n) \rightarrow \mathbb{C}$ be the complex-valued function induced by its $\mathbf{U}(n)$ -invariant lift $\hat{\phi}_a : \mathbf{Sp}(n) \rightarrow \mathbb{C}$ given by*

$$\hat{\phi}_a(q) = \text{trace}(A q q^t).$$

Then ϕ_a is an eigenfunction on $\mathbf{Sp}(n)/\mathbf{U}(n)$ with eigenvalues

$$\lambda = -2(n+1) \quad \text{and} \quad \mu = -2.$$

With the following result we construct a complex $(2n-1)$ -dimensional family

$$\mathcal{F} = \{\phi_a^{-1}(\{0\}) \mid [a] \in \mathbb{C}P^{2n-1}\}$$

of minimal submanifolds of $\mathbf{Sp}(n)/\mathbf{U}(n)$ of codimension two. This is our main result of this section.

Theorem 9.2. *Let a be a non-zero element of \mathbb{C}^{2n} and $A \in \mathbb{C}^{2n \times 2n}$ with $A = a^t a$. Further let $\phi_a : \mathbf{Sp}(n)/\mathbf{U}(n) \rightarrow \mathbb{C}$ be the complex-valued eigenfunction induced by its $\mathbf{U}(n)$ -invariant lift $\hat{\phi}_a : \mathbf{Sp}(n) \rightarrow \mathbb{C}$ given by*

$$\hat{\phi}_a(q) = \text{trace}(Aqq^t).$$

Then ϕ_a is a submersion and the compact fibre $\phi_a^{-1}(\{0\})$ over $0 \in \mathbb{C}$ is a minimal submanifold of $\mathbf{Sp}(n)/\mathbf{U}(n)$ of codimension two.

Proof. Let us assume that $q \in \mathbf{Sp}(n)$ is a critical point for the function $\hat{\phi}_a$ i.e. that the gradient

$$\nabla \hat{\phi}_a(q) = \sum_{X \in \mathcal{B}_p} X(\hat{\phi}_a(q)) \cdot X = 0.$$

By setting $D_t^a(\phi_a) = 0$ and $D_t^b(\phi_a) = 0$, for all $1 \leq t \leq n$, we obtain the following relations

$$\left(\sum_{k=1}^n a_k q_{kt} \right) \cdot \left(\sum_{k=1}^n a_k q_{k,n+t} \right) = 0 \quad (9.1)$$

and

$$\left(\sum_{k=1}^n a_k q_{kt} \right)^2 - \left(\sum_{k=1}^n a_k q_{k,n+t} \right)^2 = 0, \quad (9.2)$$

Equation 9.1 implies that, for all $1 \leq t \leq n$, either

$$\sum_{k=1}^n a_k q_{kt} = 0 \quad \text{or} \quad \sum_{k=1}^n a_k q_{k,n+t} = 0.$$

By plugging this into the equation 9.1 we then yield

$$\sum_{k=1}^n a_k q_{kt} = 0 \quad \text{and} \quad \sum_{k=1}^n a_k q_{k,n+t} = 0,$$

for all $1 \leq t \leq n$. But this is clearly impossible.

This shows that the eigenfunction $\hat{\phi}_a$ has no critical points and hence is a submersion. It follows from Corollary 3.2 that the origin $0 \in \mathbb{C}$ lies in the image of ϕ_a . Theorem 3.1 implies that the fibre $\phi_a^{-1}(\{0\})$ is a minimal submanifold of $\mathbf{Sp}(n)/\mathbf{U}(n)$ of codimension two. \square

10. THE SYMMETRIC SPACE $\mathbf{SO}(2n)/\mathbf{U}(n)$

In this section we prove Theorem 10.2 and thereby construct a new multi-dimensional family of compact minimal submanifolds of the Riemannian symmetric space $\mathbf{SO}(2n)/\mathbf{U}(n)$. It is well-known that the natural projection $\pi : \mathbf{SO}(2n) \rightarrow \mathbf{SO}(2n)/\mathbf{U}(n)$ from the special orthogonal group $\mathbf{SO}(2n)$ is a Riemannian submersion. This means that we can apply Corollary 2.6 in this situation.

For the unitary group $\mathbf{U}(n)$ we have a well-known natural group monomorphism

$$\psi : \mathbf{U}(n) \rightarrow \mathbf{SO}(2n), \quad \text{with } \psi : z = x + iy \mapsto \begin{pmatrix} x & -y \\ y & x \end{pmatrix},$$

embedding it into the special orthogonal group $\mathbf{SO}(2n)$. Here $x, y \in \mathbb{R}^{n \times n}$ are the real and imaginary parts of $z \in \mathbf{U}(n)$. In this section we identify $\mathbf{U}(n)$ with the image $\psi(\mathbf{U}(n))$ which clearly is a subgroup of $\mathbf{SO}(2n)$.

For the Lie algebra $\mathfrak{so}(2n)$ we have the orthogonal decomposition

$$\mathfrak{so}(2n) = \mathfrak{u}(n) \oplus \mathfrak{p},$$

where $\mathfrak{u}(n)$ is the Lie algebra of $\mathbf{U}(n)$ as a subgroup of $\mathbf{SO}(2n)$. A natural orthonormal basis for the vector space \mathfrak{p} is given by

$$\mathcal{B}_{\mathfrak{p}} = \left\{ Y_{rs}^a = \frac{1}{\sqrt{2}} \begin{pmatrix} Y_{rs} & 0 \\ 0 & -Y_{rs} \end{pmatrix}, Y_{rs}^b = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & Y_{rs} \\ Y_{rs} & 0 \end{pmatrix} \mid 1 \leq r < s \leq n \right\}.$$

Let I_n denote the $n \times n$ identity matrix and J_n be the standard complex structure on \mathbb{R}^{2n} given by

$$J_n = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}.$$

Then consider the map $\sigma : \mathbf{SO}(2n) \rightarrow \mathbf{SO}(2n)$ satisfying

$$\sigma(x) = x \cdot J_n \cdot x^t.$$

A simple calculation shows that this map is $\mathbf{U}(n)$ -invariant i.e. for any $y \in \mathbf{U}(n)$ and $x \in \mathbf{SO}(2n)$ we have

$$\sigma(xy) = xyJ_ny^tx^t = xJ_nx^t = \sigma(x).$$

This shows that for any skew-symmetric matrix $A \in \mathbb{C}^{2n \times 2n}$ the function $\hat{\phi}_A : \mathbf{SO}(2n) \rightarrow \mathbb{C}$, given by

$$\hat{\phi}_A(x) = \text{trace}(A\sigma(x)) = \text{trace}(x^tAxJ_n),$$

is $\mathbf{U}(n)$ -invariant and hence induces a map $\phi_A : \mathbf{SO}(2n)/\mathbf{U}(n) \rightarrow \mathbb{C}$ on the the quotient space. Note that in this case the skew-symmetry condition can be assumed without loss of generality, since $\sigma(x)$ is skew-symmetric.

We now have the following slightly improved version of Proposition 5.1. of [15].

Proposition 10.1. *For $n \geq 2$, let a, b be two linearly independent elements of \mathbb{C}^{2n} satisfying the condition*

$$(a, a)(b, b) - (a, b)^2 = 0.$$

Let $A \in \mathbb{C}^{2n \times 2n}$ be the skew-symmetric matrix

$$A = \sum_{j, \alpha=1}^{2n} a_j b_{\alpha} Y_{j\alpha}.$$

Further let $\phi_{a,b} : \mathbf{SO}(2n)/\mathbf{U}(n) \rightarrow \mathbb{C}$ be the complex-valued function induced by its $\mathbf{U}(n)$ -invariant lift $\hat{\phi}_{a,b} : \mathbf{SO}(2n) \rightarrow \mathbb{C}$ given by

$$\hat{\phi}_{a,b}(x) = \text{trace}(AxJ_n x^t).$$

Then $\phi_{a,b}$ is an eigenfunction on $\mathbf{SO}(2n)/\mathbf{U}(n)$ with the eigenvalues

$$\lambda = -2(n-1) \quad \text{and} \quad \mu = -1.$$

Proof. The arguments needed here are exactly the same as those employed for Proposition 5.1. in [15]. \square

We now state Theorem 10.2, which is our main result of this section.

Theorem 10.2. *For $n \geq 2$, let $a = u + iv \in \mathbb{C}^{2n}$ and $b \in \mathbb{R}^{2n}$ be two linearly independent elements, such that $(a, a) = (a, b) = 0$. Let $A \in \mathbb{C}^{2n \times 2n}$ be the skew-symmetric matrix*

$$A = \sum_{j,\alpha=1}^{2n} a_j b_\alpha Y_{j\alpha}.$$

Further let $\phi_{a,b} : \mathbf{SO}(2n)/\mathbf{U}(n) \rightarrow \mathbb{C}$ be the complex-valued eigenfunction induced by its $\mathbf{U}(n)$ -invariant lift $\hat{\phi}_{a,b} : \mathbf{SO}(2n) \rightarrow \mathbb{C}$ given by

$$\hat{\phi}_{a,b}(x) = \text{trace}(AxJ_n x^t).$$

Then the origin $0 \in \mathbb{C}$ is a regular value of $\phi_{a,b}$, so the compact fibre $\phi_{a,b}^{-1}(\{0\})$ is a minimal submanifold of $\mathbf{SO}(2n)/\mathbf{U}(n)$ of codimension two.

Proof. From the definition $\hat{\phi}_{a,b}(x) = \text{trace}(AxJ_n x^t)$ of $\hat{\phi}_{a,b}$, it is easily seen that

$$\hat{\phi}_{a,b}(x) = \frac{2}{\sqrt{2}} \cdot \sum_{t=1}^n ((a, x_{n+t})(b, x_t) - (a, x_t)(b, x_{n+t})).$$

Since $x \in \mathbf{SO}(2n)$, we know that its columns x_1, x_2, \dots, x_{2n} form an orthonormal basis for the vector space \mathbb{R}^{2n} . This gives the equation

$$(a, b) = \sum_{k=1}^{2n} (a, x_k)(b, x_k) = 0.$$

Corollary 3.2 implies that there exists an element $x \in \mathbf{SO}(2n)$ satisfying $\hat{\phi}_{a,b}(x) = 0$. Let us now assume that the element $x \in \mathbf{SO}(2n)$ is in the fibre $\hat{\phi}_{a,b}^{-1}(\{0\})$ i.e.

$$\sum_{t=1}^n ((a, x_{n+t})(b, x_t) - (a, x_t)(b, x_{n+t})) = 0.$$

The orthogonal complement \mathfrak{p} of the subspace $\mathfrak{u}(n)$ in the Lie algebra $\mathfrak{so}(2n)$ is generated by the elements

$$Y_{rs}^a = \begin{pmatrix} Y_{rs} & 0 \\ 0 & -Y_{rs} \end{pmatrix}, Y_{rs}^b = \begin{pmatrix} 0 & Y_{rs} \\ Y_{rs} & 0 \end{pmatrix} \in \mathcal{B}_{\mathfrak{p}},$$

where $1 \leq r < s \leq n$. A simple computation shows that for $Y_{rs}^a, Y_{rs}^b \in \mathcal{B}_p$ the conditions

$$\Re Y_{rs}^a(\hat{\phi}_{a,b}) = 0 \quad \text{and} \quad \Re Y_{rs}^b(\hat{\phi}_{a,b}) = 0$$

are equivalent to

$$(u, x_r)(b, x_{n+s}) + (u, x_{n+r})(b, x_s) = (u, x_s)(b, x_{n+r}) + (u, x_{n+s})(b, x_r),$$

$$(u, x_r)(b, x_s) + (u, x_{n+s})(b, x_{n+r}) = (u, x_s)(b, x_r) + (u, x_{n+r})(b, x_{n+s}),$$

respectively. The non vanishing vector $a = u + i v \in \mathbb{C}^{2n}$ is isotropic i.e. $(a, a) = 0$, hence $|u| = |v| \neq 0$. This implies that there exists an $r \in \{1, 2, \dots, 2n\}$ such that $(u, x_r) \neq 0$. Without loss of generality we can assume that $(u, x_1) \neq 0$. Then with $s = 2, 3, \dots, n$ we yield the following system of $2n = 2(n-1) + 2$ equations

$$0 = (u, x_r)(b, x_{n+s}) + (u, x_{n+r})(b, x_s) - (u, x_s)(b, x_{n+r}) - (u, x_{n+s})(b, x_r),$$

$$0 = (u, x_r)(b, x_s) + (u, x_{n+s})(b, x_{n+r}) - (u, x_s)(b, x_r) - (u, x_{n+r})(b, x_{n+s}),$$

$$0 = \Re(a, b) = \sum_{k=1}^{2n} (u, x_k)(b, x_k),$$

$$0 = \Re \hat{\phi}_{a,b}(x) = - \sum_{k=1}^n ((u, x_{n+k})(b, x_k) - (u, x_k)(b, x_{n+k})).$$

This can be written in the form

$$M_n \cdot \begin{bmatrix} (b, x_1) \\ (b, x_2) \\ \vdots \\ (b, x_{2n}) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

By reordering the columns of the matrix $M_n \in \mathbb{R}^{2n \times 2n}$ it can be brought on the form

$$M_n \mapsto \tilde{M}_n = (u, x_1) \cdot I_{2n} + S_n,$$

where I_{2n} is the identity matrix and $S_n \in \mathbb{R}^{2n \times 2n}$ is skew-symmetric, see Appendix A. This means that the matrix M_n is invertible. A lengthy and rather tedious calculation shows that $M_n \in \mathbb{R}^{2n \times 2n}$ satisfies

$$|\det M_n| = |\det \tilde{M}_n| = ((u, x_1)^2 + (u, x_{n+1})^2)^{n-2} \cdot |u|^4 \neq 0.$$

This implies that the non-zero vector $b \in \mathbb{R}^{2n}$ vanishes. This contradicts the fact that an element $x \in \hat{\phi}_{a,b}^{-1}(\{0\})$ is a critical point of the $\mathbf{U}(n)$ -invariant function $\hat{\phi}_{a,b} : \mathbf{SO}(2n) \rightarrow \mathbb{C}$. It then follows that the compact fibre $\phi_{a,b}^{-1}(\{0\})$ is regular. According to Theorem 3.1 it is minimal as well. \square

11. THE SYMMETRIC SPACE $\mathbf{SU}(2n)/\mathbf{Sp}(n)$

In this section we prove Theorem 11.2 and thereby construct a new multi-dimensional family of compact minimal submanifolds of the Riemannian symmetric space $\mathbf{SU}(2n)/\mathbf{Sp}(n)$. It is well-known that the natural projection $\pi : \mathbf{SU}(2n) \rightarrow \mathbf{SU}(2n)/\mathbf{Sp}(n)$ from the special unitary group $\mathbf{SU}(2n)$ is a Riemannian submersion. This means that we can employ Corollary 2.6 in this situation.

For the quaternionic unitary group $\mathbf{Sp}(n)$ we have the well-known group monomorphism

$$\psi : \mathbf{Sp}(n) \rightarrow \mathbf{SU}(2n), \quad \text{with } \psi : q = z + jw \mapsto \begin{pmatrix} z & -\bar{w} \\ w & \bar{z} \end{pmatrix},$$

embedding $\mathbf{Sp}(n)$ into the special orthogonal group $\mathbf{SU}(2n)$. Here $z, w \in \mathbb{C}^{n \times n}$. In this section we identify $\mathbf{Sp}(n)$ with the image $\psi(\mathbf{Sp}(n))$ which clearly is a subgroup of $\mathbf{SU}(2n)$.

For the Lie algebra $\mathfrak{su}(2n)$ we have the orthogonal decomposition

$$\mathfrak{su}(2n) = \mathfrak{sp}(n) \oplus \mathfrak{p},$$

where $\mathfrak{sp}(n)$ is the Lie algebra of $\mathbf{Sp}(n)$ as a subgroup of $\mathbf{SU}(2n)$. Then the vector space \mathfrak{p} is generated by the elements of the set

$$\begin{aligned} \mathcal{S} = & \left\{ Y_{rs}^a = \frac{1}{2} \begin{pmatrix} Y_{rs} & 0 \\ 0 & -Y_{rs} \end{pmatrix}, X_{rs}^a = \frac{1}{2} \begin{pmatrix} iX_{rs} & 0 \\ 0 & iX_{rs} \end{pmatrix}, Y_{rs}^b = \frac{1}{2} \begin{pmatrix} 0 & Y_{rs} \\ Y_{rs} & 0 \end{pmatrix}, \right. \\ & Y_{rs}^c = \frac{1}{2} \begin{pmatrix} 0 & iY_{rs} \\ -iY_{rs} & 0 \end{pmatrix} \mid 1 \leq r < s \leq n \Big\} \\ & \cup \left\{ D_{rs}^a = \frac{1}{2} \begin{pmatrix} iD_{rs} & 0 \\ 0 & iD_{rs} \end{pmatrix} \mid 1 \leq r < s \leq n \right\}, \end{aligned}$$

where $D_{rs} = \frac{1}{\sqrt{2}} \cdot (E_{rr} - E_{ss})$.

Let us now consider the map $\sigma : \mathbf{SU}(2n) \rightarrow \mathbf{SU}(2n)$ satisfying

$$\sigma(x) = z J_n z^t.$$

A simple calculation shows that this map is $\mathbf{Sp}(n)$ -invariant i.e. for any $q \in \mathbf{Sp}(n)$ and $z \in \mathbf{SU}(2n)$ we have

$$\sigma(zq) = zqJ_nq^tz^t = zJ_nz^t = \sigma(z).$$

This shows that for any skew-symmetric matrix $A \in \mathbb{C}^{2n \times 2n}$ the function $\hat{\phi}_A : \mathbf{SU}(2n) \rightarrow \mathbb{C}$, given by

$$\hat{\phi}_A(z) = \text{trace}(AzJ_nz^t),$$

is $\mathbf{Sp}(n)$ -invariant and hence induces a map $\phi_A : \mathbf{SU}(2n)/\mathbf{Sp}(n) \rightarrow \mathbb{C}$ on the quotient space. Note that in this case the skew-symmetry condition can be assumed without loss of generality, since $\sigma(z)$ is skew-symmetric.

The following useful result can be found in [15].

Proposition 11.1. [15] *For $n \geq 2$, let a, b be two linearly independent elements of \mathbb{C}^{2n} and $A \in \mathbb{C}^{2n \times 2n}$ be the skew-symmetric matrix*

$$A = \sum_{j, \alpha=1}^{2n} a_j b_\alpha Y_{j\alpha}.$$

Further let $\phi_{a,b} : \mathbf{SU}(2n)/\mathbf{Sp}(n) \rightarrow \mathbb{C}$ be the complex-valued function induced by its $\mathbf{Sp}(n)$ -invariant lift $\hat{\phi}_{a,b} : \mathbf{SU}(2n) \rightarrow \mathbb{C}$ given by

$$\hat{\phi}_{a,b}(z) = \text{trace}(AzJ_n z^t).$$

Then $\phi_{a,b}$ is an eigenfunction on $\mathbf{SU}(2n)/\mathbf{Sp}(n)$ with eigenvalues

$$\lambda = -\frac{2(2n^2 - n - 1)}{n} \quad \text{and} \quad \mu = -\frac{2(n - 1)}{n}.$$

We now state Theorem 11.2, which is our main result of this section.

Theorem 11.2. *For $n \geq 2$, let a, b be two linearly independent elements of \mathbb{C}^{2n} and $A \in \mathbb{C}^{2n \times 2n}$ be the skew-symmetric matrix*

$$A = \sum_{j, \alpha=1}^{2n} a_j b_\alpha Y_{j\alpha}.$$

Further let $\phi_{a,b} : \mathbf{SU}(2n)/\mathbf{Sp}(n) \rightarrow \mathbb{C}$ be the complex-valued eigenfunction induced by its $\mathbf{Sp}(n)$ -invariant lift $\hat{\phi}_{a,b} : \mathbf{SU}(2n) \rightarrow \mathbb{C}$ given by

$$\hat{\phi}_{a,b}(z) = \text{trace}(AzJ_n z^t).$$

Then the origin $0 \in \mathbb{C}$ is a regular value of $\phi_{a,b}$ and the compact fibre $\phi_{a,b}^{-1}(\{0\})$ is a minimal submanifold of $\mathbf{SU}(2n)/\mathbf{Sp}(n)$.

Proof. Let us assume that $z \in \mathbf{SU}(2n)$ is an element of the fibre $\hat{\phi}_{a,b}^{-1}(\{0\})$ i.e. $\hat{\phi}_{a,b}(z) = 0$. The statement of Corollary 3.2 guarantees the existence of such a point z . Further assume that z is a critical point of $\hat{\phi}_{a,b} : \mathbf{SU}(2n) \rightarrow \mathbb{C}$ which means that $Y(\hat{\phi}_{a,b}(z)) = 0$ for every element in the orthogonal complement \mathfrak{p} of $\mathfrak{sp}(n)$ in $\mathfrak{su}(2n)$. It can now be shown that for all $1 \leq r < s \leq n$,

$$Y_{rs}^a(\phi_{a,b}) = 0, \quad Y_{rs}^b(\phi_{a,b}) = 0, \quad X_{rs}^a(\phi_{a,b}) = 0, \quad Y_{rs}^c(\phi_{a,b}) = 0$$

if and only if

$$\begin{aligned} 0 &= (a, z_{n+s})(b, z_r) - (a, z_{n+r})(b, z_s) + (a, z_s)(b, z_{n+r}) - (a, z_r)(b, z_{n+s}), \\ 0 &= (a, z_s)(b, z_r) - (a, z_r)(b, z_s) - (a, z_{n+s})(b, z_{n+r}) + (a, z_{n+r})(b, z_{n+s}), \\ 0 &= (a, z_{n+s})(b, z_r) + (a, z_{n+r})(b, z_s) - (a, z_s)(b, z_{n+r}) - (a, z_r)(b, z_{n+s}), \\ 0 &= (a, z_s)(b, z_r) - (a, z_r)(b, z_s) + (a, z_{n+s})(b, z_{n+r}) - (a, z_{n+r})(b, z_{n+s}). \end{aligned}$$

From the first and the third equations we immediately see that for all $1 \leq r < s \leq n$, we have

$$(a, z_r)(b, z_{n+s}) = (a, z_{n+s})(b, z_r), \quad (11.1)$$

$$(a, z_s)(b, z_{n+r}) = (a, z_{n+r})(b, z_s). \quad (11.2)$$

Similarly, the second and the fourth equations tell us that for all $1 \leq r < s \leq n$,

$$(a, z_r)(b, z_s) = (a, z_s)(b, z_r), \quad (11.3)$$

$$(a, z_{n+r})(b, z_{n+s}) = (a, z_{n+s})(b, z_{n+r}). \quad (11.4)$$

Since the columns of \bar{z} form a basis of \mathbb{C}^{2n} , there exists an index $1 \leq t \leq 2n$ such that

$$\langle a, \bar{z}_t \rangle = (a, z_t) \neq 0.$$

Without loss of generality, we can assume that $1 \leq t \leq n$. Equations 11.1 and 11.3 now show that for all $1 \leq s \leq n$, $s \neq t$, it holds that

$$(b, z_{n+s}) = \frac{(a, z_{n+s})}{(a, z_t)} \cdot (b, z_t) \quad \text{and} \quad (b, z_s) = \frac{(a, z_s)}{(a, z_t)} \cdot (b, z_t).$$

Now note that $\phi_{a,b}(z) = 0$ if and only if

$$0 = (a, z_{n+1})(b, z_1) + \cdots + (a, z_{2n})(b, z_n) - (a, z_1)(b, z_{n+1}) - \cdots - (a, z_n)(b, z_{2n}).$$

Entering our latest findings into this equation, we note that

$$\begin{aligned} 0 &= (a, z_{n+t})(b, z_t) + \sum_{s \neq t} (a, z_{n+s}) \cdot \frac{(a, z_s)}{(a, z_t)} \cdot (b, z_t) \\ &\quad - (a, z_t)(b, z_{n+t}) - \sum_{s \neq t} (a, z_s) \cdot \frac{(a, z_{n+s})}{(a, z_t)} \cdot (b, z_t) \\ &= (a, z_{n+t})(b, z_t) - (a, z_t)(b, z_{n+t}). \end{aligned}$$

Thus, we also have that

$$(b, z_{n+t}) = \frac{(a, z_{n+t})}{(a, z_t)} \cdot (b, z_t).$$

To summarise, we have now established that for every $1 \leq s \leq 2n$,

$$\langle b, \bar{z}_s \rangle = (b, z_s) = \frac{(a, z_s)}{(a, z_t)} \cdot (b, z_t) = \frac{(b, z_t)}{(a, z_t)} \cdot \langle a, \bar{z}_s \rangle.$$

Thus,

$$b = \sum_{s=1}^{2n} \langle b, \bar{z}_s \rangle \bar{z}_s = \frac{(b, z_t)}{(a, z_t)} \cdot \sum_{s=1}^{2n} \langle a, \bar{z}_s \rangle \bar{z}_s = \frac{(b, z_t)}{(a, z_t)} \cdot a.$$

This is a contradiction, since we have assumed that the elements a and b are linearly independent. We conclude that $\phi_{a,b}$ is regular over 0. By Theorem 3.1, $\phi_{a,b}^{-1}(\{0\})$ is a minimal submanifold of $\mathbf{SU}(2n)/\mathbf{Sp}(n)$. \square

12. ACKNOWLEDGEMENTS

The authors are grateful to Thomas Jack Munn for useful discussions on this work.

APPENDIX A. THE LINEAR SYSTEM FOR $\mathbf{SO}(2n)/\mathbf{U}(n)$ **The Case $n = 2$.**

Here we assume that $(u, x_1) \neq 0$ i.e. $r = 1$ and $s = 2$. Then obtain the following system of $4 = 2(2 - 1) + 2$ equations

$$\begin{bmatrix} -(u, x_4) & (u, x_3) & -(u, x_2) & (u, x_1) \\ -(u, x_2) & (u, x_1) & (u, x_4) & -(u, x_3) \\ (u, x_1) & (u, x_2) & (u, x_3) & (u, x_4) \\ (u, x_3) & (u, x_4) & -(u, x_1) & -(u, x_2) \end{bmatrix} \cdot \begin{bmatrix} (b, x_1) \\ (b, x_2) \\ (b, x_3) \\ (b, x_4) \end{bmatrix} = 0.$$

The matrix M_2 can then be transformed into $\tilde{M}_2 = (u, x_1) \cdot I_4 + S_2$.

$$\tilde{M}_2 = \begin{bmatrix} (u, x_1) & (u, x_3) & -(u, x_4) & (u, x_2) \\ -(u, x_3) & (u, x_1) & -(u, x_2) & -(u, x_4) \\ (u, x_4) & (u, x_2) & (u, x_1) & -(u, x_3) \\ -(u, x_2) & (u, x_4) & (u, x_3) & (u, x_1) \end{bmatrix}$$

It can be shown that $|\det M_2| = |\det \tilde{M}_2| = |u|^4 \neq 0$.

The Case $n = 3$.

Here we assume that $(u, x_1) \neq 0$ i.e. $r = 1$ and $s = 2, 3$ and yield the following matrix $M_3 \in \mathbb{R}^{6 \times 6}$

$$M_3 = \begin{bmatrix} -(u, x_5) & (u, x_4) & 0 & -(u, x_2) & (u, x_1) & 0 \\ -(u, x_2) & (u, x_1) & 0 & (u, x_5) & -(u, x_4) & 0 \\ -(u, x_6) & 0 & (u, x_4) & -(u, x_3) & 0 & (u, x_1) \\ -(u, x_3) & 0 & (u, x_1) & (u, x_6) & 0 & -(u, x_4) \\ (u, x_1) & (u, x_2) & (u, x_3) & (u, x_4) & (u, x_5) & (u, x_6) \\ (u, x_4) & (u, x_5) & (u, x_6) & -(u, x_1) & -(u, x_2) & -(u, x_3) \end{bmatrix}.$$

The matrix M_3 can then be transformed into $\tilde{M}_3 = (u, x_1) \cdot I_6 + S_3$.

$$\tilde{M}_3 = \begin{bmatrix} (u, x_1) & (u, x_4) & 0 & 0 & -(u, x_5) & (u, x_2) \\ -(u, x_4) & (u, x_1) & 0 & 0 & -(u, x_2) & -(u, x_5) \\ 0 & 0 & (u, x_1) & (u, x_4) & -(u, x_6) & (u, x_3) \\ 0 & 0 & -(u, x_4) & (u, x_1) & -(u, x_3) & -(u, x_6) \\ (u, x_5) & (u, x_2) & (u, x_6) & (u, x_3) & (u, x_1) & -(u, x_4) \\ -(u, x_2) & (u, x_5) & -(u, x_3) & (u, x_6) & (u, x_4) & (u, x_1) \end{bmatrix}.$$

$$|\det M_3| = |\det \tilde{M}_3| = ((u, x_1)^2 + (u, x_4)^2) \cdot |u|^4 \neq 0.$$

The General Case $n \geq 2$.

Here we assume that $(u, x_1) \neq 0$ i.e. $r = 1$ and $s = 2, 3, \dots, n$ and then obtain the following system of $2n = 2(n - 1) + 2$ equations

$$\begin{aligned} 0 &= (u, x_r)(b, x_{n+s}) + (u, x_{n+r})(b, x_s) - (u, x_s)(b, x_{n+r}) - (u, x_{n+s})(b, x_r), \\ 0 &= (u, x_r)(b, x_s) + (u, x_{n+s})(b, x_{n+r}) - (u, x_s)(b, x_r) - (u, x_{n+r})(b, x_{n+s}), \end{aligned}$$

$$\begin{aligned}
0 &= \Re(a, b) = \sum_{k=1}^{2n} (u, x_k) (b, x_k), \\
0 &= \Re \hat{\phi}_{a,b}(x) = - \sum_{k=1}^n ((u, x_{n+k}) (b, x_k) - (u, x_k) (b, x_{n+k})).
\end{aligned}$$

The corresponding matrix M_n can then be transformed into $\tilde{M}_n = (u, x_1) \cdot I_{2n} + S_n$. Here the matrix $M_n \in \mathbb{R}^{2n \times 2n}$ satisfies

$$|\det M_n| = ((u, x_1)^2 + (u, x_{n+1})^2)^{n-2} \cdot |u|^4 \neq 0.$$

REFERENCES

- [1] P. Baird, J. Eells, *A conservation law for harmonic maps*, Geometry Symposium Utrecht 1980, Lecture Notes in Mathematics **894**, Springer (1981), 1-25.
- [2] P. Baird and J. C. Wood, *Harmonic morphisms between Riemannian manifolds*, London Math. Soc. Monogr. **29**, Oxford Univ. Press (2003).
- [3] F. E. Burstall, J. H. Rawnsley, *Twistor theory for Riemannian symmetric spaces. With applications to harmonic maps of Riemann surfaces* Lecture Notes in Math. **1424**, Springer (1990)
- [4] F.E. Burstall, M.A. Guest, *Harmonic two-spheres in compact symmetric spaces, revisited*, Math. Ann. **309** (1997), 541-752.
- [5] E. Calabi, *Minimal immersions of surfaces in euclidean spheres*, J. Differential Geom. **1** (1967), 111-125.
- [6] J. Eells, J. H. Sampson, *Harmonic mappings of Riemannian manifolds*, Amer. J. Math. **86** (1964), 109-160.
- [7] J. Eells, J.C. Wood, *Harmonic maps from surfaces to complex projective spaces*, Adv. in Math. **49**, (1983), 217-263.
- [8] J. M. Gegenfurtner, *Minimal Submanifolds of the Classical Compact Riemannian Symmetric Spaces*, Master's dissertation, University of Lund (2024).
www.matematik.lu.se/matematiklu/personal/sigma/students/Johanna-Marie-Gegenfurtner-MSc.pdf
- [9] E. Ghandour, S. Gudmundsson, *Explicit p-harmonic functions on the real Grassmannians*, Adv. Geom. **23** (2023), 315–321.
- [10] E. Ghandour, S. Gudmundsson, *Explicit harmonic morphisms and p-harmonic functions from the complex and quaternionic Grassmannians*, Ann. Global Anal. Geom. **64** (2023), 15.
- [11] S. Gudmundsson, *The Bibliography of Harmonic Morphisms*, www.matematik.lu.se/matematiklu/personal/sigma/harmonic/bibliography.html
- [12] S. Gudmundsson, S. Montaldo, A. Ratto, *Biharmonic functions on the classical compact simple Lie groups*, J. Geom. Anal. **28** (2018), 1525-1547.
- [13] S. Gudmundsson, T. J. Munn, *Minimal submanifolds via complex-valued eigenfunctions*, J. Geom. Anal. **34** (2024), 190.
- [14] S. Gudmundsson, A. Sakovich, *Harmonic morphisms from the classical compact semisimple Lie groups*, Ann. Global Anal. Geom. **33** (2008), 343-356.
- [15] S. Gudmundsson, A. Siffert, M. Sobak, *Explicit proper p-harmonic functions on the Riemannian symmetric spaces $\mathbf{SU}(n)/\mathbf{SO}(n)$, $\mathbf{Sp}(n)/\mathbf{U}(n)$, $\mathbf{SO}(2n)/\mathbf{U}(n)$, $\mathbf{SU}(2n)/\mathbf{Sp}(n)$* , J. Geom. Anal. **32** (2022), 147
- [16] S. Gudmundsson, M. Sobak, *Proper r-harmonic functions from Riemannian manifolds*, Ann. Global Anal. Geom. **57** (2020), 217-223.
- [17] S. Helgason, *Differential Geometry, Lie groups and Symmetric Spaces*, Pure and Applied Mathematics **80**, Academic Press (1978).

- [18] O. Riedler, A. Siffert, *Global eigenfamilies on compact manifolds*, preprint, University of Münster (2024).
- [19] K. Uhlenbeck, *Harmonic maps into Lie groups: classical solutions of the chiral model*, J. Differential Geom. **30** (1989), 1–50.

MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF LUND, BOX 118, LUND 221 00, SWEDEN

Email address: `Johanna.Gegenfurtner@gmail.com`

MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF LUND, BOX 118, LUND 221 00, SWEDEN

Email address: `Sigmundur.Gudmundsson@math.lu.se`