A CENTRAL LIMIT THEOREM ASSOCIATED WITH A SEQUENCE OF POSITIVE LINE BUNDLES

AFRIM BOJNIK AND OZAN GÜNYÜZ

ABSTRACT. We prove a central limit theorem for random currents of integration along the zero divisors of standard Gaussian holomorphic sections in a sequence of holomorphic line bundles with Hermitian metrics of class \mathscr{C}^3 over a compact Kähler manifold. In the course of our analysis, we derive first-order asymptotics and upper decay estimates for near and off-diagonal Bergman kernels, respectively. These results are essential for determining the statistical properties of the zeros of random holomorphic sections.

1. INTRODUCTION

Over recent years, there has been a growing interest in probabilistic challenges within the scope of complex algebraic or analytic geometry. Such problems initially arose from studying the zeros of random polynomials and other analytic functions, analyzed by mathematicians under different assumptions about random coefficients. The field gained momentum with the foundational works of Bloch-Pólya[BP], Littlewood-Offord [LO43], Hammersley [HAM56], Kac [Kac43], and Erdös-Turán [ET50], and has now become a classical area of study. A significant breakthrough was achieved by Shiffman and Zelditch [SZ99], who extended these ideas into a complex geometric setting. They demonstrated that the zeros of random sections of high tensor powers of a positive line bundle L on a projective manifold X tend to become uniformly distributed in accordance with the natural measure from line bundle L. In a series of papers, these authors further delved into the correlations, variance, expected distribution and other statistical properties related to zeros of holomorphic sections, as elaborated in references [BSZ, SZ08, SZ10]. In this framework, Dinh and Sibony [DS06] innovated a method for analyzing zero distribution by applying formalism of meromorphic transforms from complex dynamics, and set convergence speed bounds in the compact case, enhancing Shiffman and Zelditch's initial results. For a thorough understanding of zero distribution, especially in the light of broader probability distributions and various geometric scenarios, we invite the reader to look into the survey [BCHM18]. Furthermore, it is important to emphasize that random polynomials and holomorphic sections have significant role in the field of statistical physics. Holomorphic random sections, in particular, are employed as an important model for understanding quantum chaos, and physicists have conducted extensive research to analyze the distribution of their zero points see for example [BBL, Hann, NV98].

Alongside these developments, investigation of the central limit theorem in the context of smooth linear statistics, such as integrals of smooth test forms over zero divisors of random holomorphic sections, is another intriguing challenge. In this domain, the work by Sodin and Tsirelson [ST] is particularly noteworthy. They established the following asymptotic normality result for Gaussian random polynomials and analytic functions in the complex plane. This result stands as a crucial instrument that will play an essential role in our study presented in this paper.

Date: May 7, 2024.

²⁰²⁰ Mathematics Subject Classification. Primary: 32A60, 60F05, 32A25; Secondary: 53C55.

Key words and phrases. random holomorphic sections, central limit theorem, compact Kähler manifolds.

A. Bojnik was partially supported by TÜBİTAK grant ARDEB-3501/118F049.

2

1.1. Theorem of Sodin and Tsirelson. Given a sequence $\{\gamma_j\}_{j=1}^{\infty}$ of complex-valued measurable functions on a measure space (G, σ) such that

(1.1)
$$\sum_{j=1}^{\infty} |\gamma_j(x)|^2 = 1 \text{ for any } x \in G,$$

following [ST] (and also [SZ10]), a *normalized complex Gaussian process* is defined to be a complexvalued random function $\alpha(x)$ on a measure space (G, σ) in the following form

(1.2)
$$\alpha(x) = \sum_{j=1}^{\infty} b_j \gamma_j(x),$$

where the coefficients b_j are i.i.d. centered complex Gaussian random variables with variance one. The covariance function of $\alpha(x)$ is defined by

(1.3)
$$\mathcal{C}(x,y) = \mathbb{E}[\alpha(x)\overline{\alpha(y)}] = \sum_{j=1}^{\infty} \gamma_j(x)\overline{\gamma_j(y)}.$$

A simple observation gives that $|\beta(x,y)| \leq 1$ and $\beta(x,x) = 1.$

Consider a sequence $\{\alpha_j\}_{j=1}^{\infty}$ of normalized complex Gaussian processes on a finite measure space (G, σ) , and let $\lambda(\rho) \in L^2(\mathbb{R}^+, e^{\frac{-\rho^2}{2}}\rho d\rho)$. Suppose $\phi : G \to \mathbb{R}$ is a bounded and measurable function, we will focus on the following non-linear functionals that also serve as random variables in this context.

(1.4)
$$\mathcal{F}_{p}^{\phi}(\alpha_{p}) = \int_{G} \lambda(|\alpha_{p}(x)|)\phi(x)d\sigma(x).$$

The next theorem (Theorem 2.2 of [ST]) was proved by Sodin and Tsirelson.

Theorem 1.1. For each n = 1, 2, ..., let $C_p(r, s)$ be the covariance functions for the complex Gaussian processes. Assume that the two conditions below hold for all $\nu \in \mathbb{N}$:

(i)

$$\liminf_{n \to \infty} \frac{\int_G \int_G |\mathcal{C}_p(r,s)|^{2\nu} \phi(r) \phi(s) d\sigma(r) d\sigma(s)}{\sup_{r \in G} \int_G |\mathcal{C}_p(r,s)| d\sigma(s)} > 0.$$

(ii)

$$\lim_{n \to \infty} \sup_{r \in G} \int_{G} |\mathcal{C}_{p}(r,s)| d\sigma(s) = 0$$

Then the distributions of the random variables

$$\frac{\mathcal{F}_p^{\phi}(\alpha_p) - \mathbb{E}[\mathcal{F}_p^{\phi}(\alpha_p)]}{\sqrt{\operatorname{Var}[\mathcal{F}_p^{\phi}(\alpha_p)]}}$$

converges weakly to the normal distribution $\mathcal{N}(0,1)$ as $p \to \infty$. If λ is increasing, then it is sufficient for (i) to hold only for $\nu = 1$.

The proof is based on applying the method of moments, a fundamental tool in probability theory, coupled with the use of the diagram technique. This approach facilitates the computation of moments for non-linear functionals, which are then compared to the moments found in a standard Gaussian distribution. Such an approach is a classical one in establishing the central limit theorem for non-linear functionals within Gaussian fields. We also remark that the condition (ii) ensures that $\operatorname{Var}[\mathcal{F}_p^{\phi}(\alpha_p)] \to 0$ as $p \to \infty$. There are two extensions of Theorem 1.1 in different contexts. The first, due to the work of Shiffman and Zelditch ([SZ10]), applies to the prequantum line bundle setting. This involves random holomorphic sections of a Hermitian line bundle (with C^3 Hermitian metrics) over a compact Kähler manifold, where the first Chern form and Kähler form satisfy the prequantum line bundle condition. The second extension is studied in [Bay16], where Bayraktar expanded Theorem 1.1 to encompass general random polynomials in \mathbb{C}^n with techniques of weighted pluripotential theory.

In both these adaptations, the asymptotic behavior of the normalized Bergman kernel, particularly off-diagonal and near-diagonal, proves to be critical. This kernel, functioning as a covariance function of normalized complex Gaussian processes, is pivotal to the analyses. Separately, Nazarov and Sodin ([NS]) explored the asymptotic normality of linear statistics of zeros in Gaussian entire functions on \mathbb{C} . Their approach is broader, considering measurable bounded test functions and the clustering of *k*-point correlation functions.

The primary objective of the current study is to apply Theorem 1.1 to a broader framework than previously discussed scenarios, specifically to standard Gaussian holomorphic sections in a sequence of positive holomorphic line bundles with class C^3 Hermitian metrics on a compact Kähler manifold. To achieve this, we start by setting up our geometric and probabilistic framework, building upon the foundations presented in earlier studies [CLMM], [CMM]. The key to our analysis is the upper decaying estimate of the off-diagonal Bergman kernel and the first order asymptotics of the Bergman kernel function. In this context, for the asymptotic distribution and variance estimates of smooth linear statistics of random zero sets in any codimension under more general probability measures, we refer the interested reader to [BG].

2. BACKGROUND

Let (X, ω) be a compact Kähler manifold, with $\dim_{\mathbb{C}} X = n$, and let $\{(L_p, h_p)\}_{p\geq 1}^{\infty}$ be a sequence of positive line bundles with Hermitian \mathscr{C}^3 -metrics h_p . In this paper, we employ the concepts of Nakano and Griffiths positivity for vector bundles. It is important to note that Nakano positivity implies Griffiths positivity in vector bundles, though the reverse is not always true (refer to [Dem12] for more details). However, in the case of line bundles, these two notions of positivity coincide. Therefore, in our context, we define positivity for a holomorphic line bundle (Hermitian, but not necessarily with \mathscr{C}^{∞} -metrics), denoted as (L, h), in the following way: a line bundle is considered positive or semi-positive if its local weight functions φ , which are of class \mathscr{C}^3 and corresponding to the metric h, are strictly plurisubharmonic and plurisubharmonic, respectively.

Since our focus is on positive line bundles, the associated local weight functions φ are strictly plurisubharmonic. This means that they belong to the class \mathscr{C}^3 and satisfy $dd^c\varphi > 0$ at every point. According to a well-known characterization of ampleness (refer to Corollary 13.3 of [Dem12]), the line bundles we are currently investigating are ample. This also indicates, following Kodaira's projectivity criterion, that any compact Kähler manifold with a positive line bundle is projective.

We denote the global holomorphic sections of L_p by $H^0(X, L_p)$. In this context, we consider an inner product on the space of smooth sections $\mathscr{C}^{\infty}(X, L_p)$, using the metric h_p and the volume form $\frac{\omega^n}{n!}$ on X. This inner product is defined as

(2.1)
$$\langle s_1, s_2 \rangle_p := \int_X \langle s_1(x), s_2(x) \rangle_{h_p} \frac{\omega^n}{n!}$$

and the norm of a section s is given by $||s||_p^2 := \langle s, s \rangle_p$. Based on the Cartan-Serre finiteness theorem, we know that $H^0(X, L_p)$ is finite-dimensional. We denote the dimension of this space as $d_p := \dim H^0(X, L_p)$. Subsequently, we consider $\mathcal{L}^2(X, L_p)$, which is the completion of $\mathscr{C}^\infty(X, L_p)$ under this norm, forming a Hilbert space of square-integrable sections of L_p . Next, we define the orthogonal projection operator $K_p : \mathcal{L}^2(X, L_p) \to H^0(X, L_p)$. The Bergman kernel, $K_p(x, y)$, is the integral kernel of this projection. If $\{S_j^p\}_{j=1}^{d_p}$ is an orthonormal basis for $H^0(X, L_p)$, by using reproducing property of $K_p(x, y)$, we express $K_p(x, y)$ in terms of this basis as follows:

(2.2)
$$K_p(x,y) = \sum_{j=1}^{d_p} S_j^p(x) \otimes S_j^p(y)^* \in L_{p,x} \otimes L_{p,y}^*.$$

where $S_j^p(y)^* = \langle ., S_j^p(y) \rangle_{h_p} \in L_{p,y}^*$. The restriction of the Bergman kernel to the diagonal of X is called the Bergman kernel function of $H^0(X, L_p)$, which we denote by $K_p(x) := K_p(x, x)$, and (2.2) becomes

(2.3)
$$K_p(x) = \sum_{j=1}^{d_p} |S_j^p(x)|_{h_p}^2.$$

The Bergman kernel function has the dimensional density property, namely

$$\int_X K_p(x) \frac{\omega^n}{n!} = d_p.$$

In addition, it satisfies the following variational principle

(2.4)
$$K_p(x) = \max\{|S(x)|_{h_p}^2 : S \in H^0(X, L_p), \ ||S||_p = 1\}.$$

This holds for every $x \in X$ for which $\varphi_p(x) > -\infty$, with φ_p denoting a local weight, defined above, for the metric h_p in the vicinity of x.

We also define the normalized Bergman kernel as follows:

$$\widehat{\mathcal{K}}_p(x,y) := \frac{|K_p(x,y)|_{h_{p,x} \otimes h_{p,y}}}{\sqrt{K_p(x)}\sqrt{K_p(y)}}.$$

which will be important throughout.

2.1. Reference Covers. (U, z), $z = (z_1, ..., z_n)$, will indicate local coordinates centered at a point $x \in X$. The closed polydisk around $y \in U$ of equilateral radius (r, ..., r), r > 0, is given by

 $P^{n}(y,r) := \{ z \in U : |z_{j} - y_{j}| \le r, \ j = 1, 2, \dots, m \}.$

The coordinates (U, z) are said to be Kähler at $y \in U$ in case

(2.5)
$$\omega_z = \frac{i}{2} \sum_{j=1}^m dz_j \wedge \overline{dz_j} + O(|z-y|^2) \sum_{j,k} dz_j \wedge d\overline{z_k} \text{ on } U_{\overline{z_k}}$$

Definition 2.1. A reference cover of X is defined as follows: for j = 1, 2, ..., N, a set of points $x_j \in X$ and

- (a) Stein open simply connected coordinate neighborhoods $(U_j, w^{(j)})$ centered at $x_j \equiv 0$.
- (b) $R_j > 0$ such that $P^n(x_j, 2R_j) \Subset U_j$ and for every $y \in P^n(x_j, 2R_j)$ there exist coordinates on U_j which are Kähler at y.

(c)
$$X = \bigcup_{j=1}^{N} P^n(x_j, R_j)$$

We will write $R = \min R_i$ once a reference cover is provided.

It is not difficult to see how one can construct a reference cover. Indeed, first, for $x \in X$, take a Stein open simply connected neighborhood (for instance, a round ball in \mathbb{C}^n) U of $0 \in \mathbb{C}^n$, where $x \equiv 0$ under a determined chart. Choose some R > 0 so that $P^n(x, R) \Subset U$ and for every $y \in P^n(x, R)$ there exist Kähler coordinates (U, z) at y. The compactness of X implies that there exist finitely many points $\{x_j\}_{j=1}^N$ such that the three conditions above are satisfied.

We take into consideration the differential operators D_w^{α} , $\alpha \in \mathbb{N}^{2n}$ on U_j , corresponding to the real coordinates associated to $w = w^j$. For $\varphi \in \mathscr{C}^k(U_j)$, we define

(2.6)
$$\|\varphi\|_{k} = \|\varphi\|_{k,w} = \sup\{|D_{w}^{\alpha}\varphi(w)| : w \in P^{n}(x_{j}, 2R_{j}), |\alpha| \le k\}.$$

Let (L, h) be a Hermitian holomorphic line bundle on X, i.e., the metric h is smooth. For $k \leq l$, write

$$||h||_{k,U_i} = \inf \{ ||\varphi_j||_k : \varphi_j \in \mathscr{C}^l(U_j) \text{ is a weight of } h \text{ on } U_j \},\$$

and

$$||h||_{k} = \max\{1, ||h||_{k, U_{j}} : 1 \le j \le N\}.$$

 φ_j is said to be a weight of h on U_j if there exists a holomorphic frame e_j of L on U_j such that $|e_j|_h = e^{-\varphi_j}$.

Lemma 2.2. Let a reference cover of X be given. Then there exists a constant D > 0 relying on the reference cover with the following property: When provided with any Hermitian line bundle (L, h) on X, any $j \in \{1, ..., N\}$ and any $x \in P^n(x_j, R_j)$, there exist coordinates $z = (z_1, ..., z_n)$ on $P^n(x, R)$ which are centered at $x \equiv 0$ and Kähler coordinates for x such that

- (i) $dV \leq (1+Dr^2)\frac{\omega^n}{n!}$ and $\frac{\omega^n}{n!} \leq (1+Dr^2)dV$ hold on $P^n(x,r)$ for any r < R where dV = dV(z) is the Euclidean volume relative to the coordinates z,
- (ii) (L,h) has a weight φ on $P^n(x,R)$ with $\varphi(z) = \Re t(z) + \sum_{j=1}^n \lambda_j |z_j|^2 + \widetilde{\varphi}(z)$, where t is a holomorphic polynomial of degree at most 1, $\lambda_j \in \mathbb{R}$ and $|\widetilde{\varphi}(z)| \leq D' ||h||_3 |z|^3$ for $z \in P^n(x,R)$.

Proof. By the definition of a reference cover, there exist coordinates z on U_j which are Kähler for $x \in P^n(x_j, R_j)$. Then $\omega = \sum_{l=1}^n dz_l \wedge d\overline{z}_l + O(|z-x|^2) \sum_{j,k} dz_j \wedge d\overline{z}_k$ and (i) holds with a constant D_j uniform for $x \in P^n(x_j, R_j)$. Let e_j be a frame of L on U_j , φ a weight of h on U_j with $|e_j|_h = e^{-\varphi}$ and $||\varphi||_{3,z} \leq 2||h||_3$. By translation, we may assume x = 0 and write $\varphi(z) = \Re t(z) + \varphi_2(z) + \varphi_3(z)$, where t(z) is a holomorphic polynomial of degree ≤ 1 in z, $\varphi_2(z) = \sum_{k,l=1}^n \mu_{kl} z_k \overline{z}_l$ and $\Re f(z) + \varphi_2(z)$ is the Taylor polynomial of order 2 of φ at 0. In order to estimate $\varphi_3(z)$, let $||\varphi||_{3,z}$ be the supremum norm of the derivatives of φ of order 3 on $P^n(x_j, R_j)$ in the z-coordinates. Then, by (2.6), there exists a constant D'_j being uniform on $P^n(x_j, R_j)$ such that $||\varphi||_{3,z} \leq D'_j ||\varphi||_{3,w} \leq 2D'_j ||h||_3$, which also gives that $||\varphi_3(z)| \leq 2D'_j ||h||_3 |z|^3$ for all $z \in P^n(x, R)$.

Applying a unitary change of coordinates, we may suppose that $\varphi(\zeta) = \Re t(\zeta) + \sum_{j=1}^{n} \lambda_j^p |\zeta_j|^2 + \widetilde{\varphi}(\zeta)$. Under these coordinates, $\frac{w^n}{n!}$ and $\widetilde{\varphi}(\zeta)$ verify the required estimates with a uniform constant D_j for $x \in P^n(x_j, R_j)$, as unitary transformations preserve distances. Finally putting $D' = \max_{1 \le j \le N} D'_j$ finishes the proof.

Let (X, ω) be a compact Kähler manifold of $\dim_{\mathbb{C}} X = n$. Following [CLMM], the diophantine approximation of the Kähler form ω is defined as follows:

(2.7)
$$\frac{1}{A_p}c_1(L_p,h_p) = \omega + O(A_p^{-a}) \text{ in the } \mathscr{C}^0\text{-topology as } p \to \infty,$$

where a > 0, $A_p > 0$ and $\lim_{p\to\infty} A_p = \infty$. This means that

(2.8)
$$\left\|\frac{1}{A_p}c_1(L_p,h_p) - \omega\right\|_{\mathscr{C}^0} = O(A_p^{-a})$$

where $A_p > 0$, a > 0 and $\lim_{p\to\infty} A_p = +\infty$. This was first considered in [CLMM] in the \mathscr{C}^{∞} -norm topology induced by the Levi-Civita connection ∇^{TX} because the authors deal with the complete asymptotic expansion of the Bergman kernel restricted to the diagonal. We do not need such a strong topology, in fact, only the \mathscr{C}^0 -norm (or continuous norm) topology will be sufficient for us.

Let $\{U_j\}_{j=1}^N$ be a finite subcover of X. Locally, on each U_j , we have the following representations

(2.9)
$$\frac{1}{A_p}c_1(L_p,h_p)(z) = i\sum_{k,j}\frac{1}{\pi}\frac{1}{A_p}\alpha_{kj}(z)dz_k \wedge d\overline{z_j}$$

and

(2.10)
$$\omega(z) = i \sum_{k,j} \chi_{kj}(z) dz_k \wedge d\overline{z_j}.$$

Here $[\chi_{kj}(z)]_{kj}$ is a positive definite Hermitian matrix, because in a Kähler manifold, we always have a strictly plurisubharmonic local potential function ψ so that

$$\chi_{kj}(z) = \frac{\partial^2 \psi}{\partial z_j \partial \overline{z_k}}(z) \in \mathbb{R}, \text{ for each } 1 \le k, j \le n.$$

Similarly, since line bundles L_p are positive, by the definition of positivity, $[\alpha_{kj}(z)]_{kj} = \left[\frac{\partial^2 \varphi_p}{\partial z_j \partial \overline{z_k}}(z)\right]_{kj}$ is a positive definite Hermitian matrix , where φ_p is the corresponding local weight function for h_p . Note that in particular $\alpha_{kj}(z) \in \mathbb{R}$ for every $1 \leq k, j \leq n$.

Let us fix some U_l taken from the subcover. By the diophantine approximation condition (2.7) on $\overline{U_l}$, for any $\epsilon > 0$, there exists some $p_0 = p_0(\epsilon) \in \mathbb{N}$ such that, for all $p \ge p_0$,

(2.11)
$$-\epsilon \le \frac{1}{\pi A_p} \alpha_{kj}^{(p)}(z) - \chi_{kj}(z) \le \epsilon$$

for all $z \in \overline{U_l}$. Take, for example,

$$\epsilon = \frac{1}{4} (\min_{j,k} \min_{z \in \overline{U_j}} \chi_{kj}(z)).$$

Then (2.11) gives

$$\frac{3}{4}\chi_{kj}(z) \le \frac{1}{\pi A_p} \alpha_{kj}^{(p)}(z) \le \frac{5}{4}\chi_{kj}(z).$$

Summing this last inequality over $idz_k \wedge d\overline{z_j}$, we have, for all $z \in \overline{U_l} \subset X$ and for all $p \ge p_0$

$$i\sum_{k,j}\frac{3\chi_{kj}(z)}{4}dz_k \wedge d\overline{z_j} \le i\sum_{k,j}\frac{1}{\pi A_p}\alpha_{kj}^{(p)}(z)dz_k \wedge d\overline{z_j} \le i\sum_{k,j}\frac{5\chi_{kj}(z)}{4}dz_k \wedge d\overline{z_j},$$

which concludes that

(2.12)
$$\frac{3\omega}{4} \le \frac{1}{A_p} c_1(L_p, h_p) \le \frac{5\omega}{4}.$$

for $p \ge p_0$. This will be useful in the proof of Theorem 4.1 and Theorem 4.2.

We also observe that, at the point $x \equiv 0$ where we have the Kähler coordinates by (2.5), we have

$$\omega_x = i \sum_{j=1}^n \frac{1}{2} dz_j \wedge d\overline{z_j}.$$

Also, by using the local representation of $c_1(L_p, h_p)$ and Lemma 2.2,

(2.13)
$$c_1(L_p,h_p)_x = dd^c \varphi_p(0) = i \sum_{j=1}^n \frac{\lambda_j^p}{\pi} dz_j \wedge d\overline{z_j}.$$

Diophantine approximation (2.7) implies

(2.14)
$$\lim_{p \to \infty} \frac{\lambda_j^p}{\pi A_p} = \frac{1}{2} \text{ for } j = 1, 2, \dots, n,$$

which in turn gives

(2.15)
$$\lim_{p \to \infty} \frac{\lambda_1^p \dots \lambda_n^p}{A_p^n} = (\frac{\pi}{2})^n.$$

In order to measure the distance between any two points x, y on the compact Kähler manifold (X, ω) , we use the Riemannian distance, which is defined as follows: As is well-known, the Kähler form ω and the complex structure J on X compatible with ω determine a Riemannian metric g on X by $g(u, v) := \omega(u, Jv)$ for all $u, v \in TX$. Given a piecewise smooth curve $\gamma : [a, b] \to X$ with $\gamma(a) = x$ and $\gamma(b) = y$, the length $L(\gamma)$ of the curve γ is given by

$$L(\gamma) = \int_{a}^{b} \sqrt{g_{\gamma(t)}(\dot{\gamma}(t), \dot{\gamma}(t))} dt$$

and the Riemannian distance d is defined by

$$d(x,y) = \inf \{L(\gamma) : \gamma(a) = x, \ \gamma(b) = y\}.$$

3. Demailly's L^2 -estimations for $\overline{\partial}$ operator

Essential for proving both the upper decay estimate of the Bergman Kernel and the first order asymptotics of the Bergman kernel function in our current diophantine setting, we follow the approaches in [CMM] and [BCM] to provide first certain L^2 estimations for solutions of the $\overline{\partial}$ equation, and then derive a weighted estimate for these solutions.

Theorem 3.1. ([Dem82], Théorème 5.1) Let (X, ω) be a Kähler manifold with $\dim_{\mathbb{C}} X = n$ having a complete Kähler metric. Let (L, h) be a singular Hermitian holomorphic line bundles such that $c_1(L, h) \ge 0$. Then for any form $g \in L^2_{n,1}(X, L, loc)$ verifying

(3.1)
$$\overline{\partial}g = 0, \quad \int_X |g|_h^2 \frac{\omega^n}{n!} < \infty$$

there is $u \in L^2_{n,0}(X, L, loc)$ with $\overline{\partial} u = g$ such that

(3.2)
$$\int_X |u|_h^2 \frac{\omega^n}{n!} \le \int_X |g|_h^2 \frac{\omega^n}{n!}.$$

Theorem 3.2. Let X be a complete Kähler manifold with $\dim_{\mathbb{C}} X = n$ and let ω be a Kähler form (not necessarily complete) on X such that its Ricci form $\operatorname{Ric}_{\omega} \geq -2\pi T_0 \omega$ on X for some constant $T_0 > 0$. Let (L_p, h_p) be a sequence of holomorphic line bundles on X with Hermitian metrics h_p of class \mathscr{C}^3 such that (2.7) holds and there is a $p_0 \in \mathbb{N}$ such that $A_p \geq 4T_0$ for all $p > p_0$. If $p > p_0$ and $f \in L^2_{0,1}(X, L_p, loc)$ satisfies $\overline{\partial} f = 0$ and $\int_X |f|^2_{h_p} \frac{\omega^n}{n!} < \infty$, then there exists $u \in L^2_{0,0}(X, L_p, loc)$ such that $\overline{\partial} u = f$ and $\int_X |u|^2_{h_p} \frac{\omega^n}{n!} \leq \frac{2}{A_p} \int_X |f|^2_{h_p} \frac{\omega^n}{n!}$. *Proof.* By the diophantine approximation relation (2.7), fix some $p_0 \in \mathbb{N}$ so that the assertions in the theorem are satisfied and also for all $p > p_0$, $\frac{3\omega}{2} \ge \frac{1}{A_p}c_1(L_p, h_p) \ge \frac{3\omega}{4}$. Let $L_p = F_p \otimes K_X$, where $F_p = L_p \otimes K_X^{-1}$. The canonical line bundle K_X is endowed with the metric h^{K_X} induced by ω . If $g_p = h_p \otimes h^{K_X^{-1}}$ is the induced metric on F_p , then, since $c_1(K_X, h_{K_X}) = -\frac{1}{2\pi} \operatorname{Ric}_{\omega}$ and $A_p \ge 4T_0$ for all $p > p_0$, (3.3)

$$\frac{1}{A_p}c_1(F_p, g_p) = \frac{1}{A_p}c_1(L_p, h_p) - \frac{1}{A_p}c_1(K_X, h^{K_X}) = \frac{1}{A_p}c_1(L_p, h_p) + \frac{1}{2\pi A_p}Ric_\omega \ge \frac{3\omega}{4} - \frac{\omega}{4} = \frac{\omega}{2} \ge 0$$

for all $p > p_0$. On the other hand, there exists a natural isometry,

$$\Psi = \sim : \Lambda^{0,q}(T^*(X)) \otimes L_p \to \Lambda^{n,q}(T^*(X)) \otimes F_p$$

by

(3.4)
$$\Psi(s) = \tilde{s} = (w^1 \wedge \ldots \wedge w^n \wedge s) \otimes (w_1 \wedge \ldots \wedge w_n),$$

where w_1, \ldots, w_n is a local orthonormal frame of $T^{(1,0)}(X)$ and $\{w^1, \ldots, w^n\}$ is the dual frame. This operator Ψ commutes with the action of $\overline{\partial}$. Now for a form $f \in L^2_{0,1}(X, L_p, loc)$ satisfying $\overline{\partial}f = 0$ and $\int_X |f|^2_{h_p} \frac{\omega^n}{n!} < \infty$, obviously we have $\int_X \frac{2}{A_p} |f|^2_{h_p} \frac{\omega^n}{n!} < \infty$. By using the isometry Ψ , we can find $\Psi(f) = F \in L^2_{n,1}(X, F_p, loc)$ with $\overline{\partial}F = \overline{\partial}\Psi(f) = \Psi\overline{\partial}f = 0$ and $\int_X \frac{2}{A_p} |F|^2_{h_p} \frac{\omega^n}{n!} < \infty$ since isometries preserve the L^2 -norm. By Theorem 3.1, there exists $\tilde{f} \in L^2_{n,0}(X, F_p, loc)$ such that $\overline{\partial}\tilde{f} = F$ and $\int_X |\tilde{f}|^2_{h_p} \frac{\omega^n}{n!} \leq \int_X \frac{2}{A_p} |F|^2_{h_p} \frac{\omega^n}{n!}$. Taking $u := \Psi^{-1}\tilde{f}$ and $f = \Psi^{-1}(F)$ finishes the proof since Ψ^{-1} is an isometry as well.

Theorem 3.3. Let (X, ω) be a compact Kähler manifold, $\dim_{\mathbb{C}} X = n$ and let $\{(L_p, h_p)\}_{p\geq 1}$ be a sequence of holomorphic line bundles on X with \mathscr{C}^3 Hermitian metrics as before such that the diophantine approximation condition (2.7) holds. Then there exists $p_0 \in \mathbb{N}$ such that if u_p are real-valued functions of class \mathscr{C}^2 on X such that

(3.5)
$$\|\overline{\partial}u_p\|_{L^{\infty}(X)} \leq \frac{\sqrt{A_p}}{8}, \ dd^c u_p \geq -\frac{A_p}{4}\omega,$$

then

(3.6)
$$\int_{X} |v|_{h_{p}}^{2} e^{2u_{p}} \frac{\omega^{n}}{n!} \leq \frac{16}{3A_{p}} \int_{X} |\overline{\partial}v|_{h_{p}}^{2} e^{2u_{p}} \frac{\omega^{n}}{n!}$$

holds for $p > p_0$ and for every \mathscr{C}^1 -smooth section v of L_p which is orthogonal to $H^0(X, L_p)$ with respect to the inner product induced by h_p and ω^n .

Proof. As in the proof of Theorem 3.2, via the diophantine convergence assumption (2.7), we first fix some $p_0 \in \mathbb{N}$ so that for any (fixed) $p > p_0$, one can get $\frac{3\omega}{4} \leq \frac{1}{A_p}c_1(L_p,h_p) \leq \frac{3\omega}{2}$ and $A_p \geq 4T_0$. The main idea is to use Theorem 3.2. To this end, let us fix a constant $T_0 > 0$ so that $\operatorname{Ric}_{\omega} \geq -2\pi T_0 \omega$ on X. Using the real-valued functions u_p given in the assumptions of theorem, we consider the metrics $g_p := e^{-2u_p}h_p$ on L_p . From (2.3.5) in [MM1](p. 98) and the second relation in (3.5) yield the following

$$c_1(L_p, g_p) = c_1(L_p, h_p) + dd^c u_p \ge \frac{3A_p\omega}{4} - \frac{A_p\omega}{4} = \frac{A_p\omega}{2}$$

If we define an inner product by using g_p in $L^2(X, L_p)$ as $(s_1, s_2)_{g_p} := \int_X \langle s_1, s_2 \rangle_{g_p} \frac{\omega^n}{n!}$, we see, by the relation $g_p = e^{-2u_p}h_p$, for every $s \in H^0(X, L_p)$,

$$(e^{u_p}v,s)_{g_p} = \int_X \langle e^{2u_p}v,s\rangle_{g_p} \frac{\omega^n}{n!} = \int_X \langle v,s\rangle_{h_p} \frac{\omega^n}{n!} = 0$$

for every \mathscr{C}^1 -smooth section v of L_p .

Write

(3.7)
$$\beta = \overline{\partial}(e^{2u_p}v) = e^{2u_p}(2\overline{\partial}u_p \wedge v + \overline{\partial}v),$$

Since $\overline{\partial}\beta = 0$ and by assumptions on u_p and v, it follows immediately that $\beta \in L^2_{0,1}(X, L_p, loc)$, so by Theorem 3.2, there exists $\tilde{v} \in L^2_{0,0}(X, L_p, loc)$ such that $\overline{\partial}\tilde{v} = \beta$ and

(3.8)
$$\int_X |\tilde{v}|_{g_p}^2 \frac{\omega^n}{n!} \le \frac{2}{A_p} \int_X |\beta|_{g_p}^2 \frac{\omega^n}{n!}.$$

Since $e^{2u_p}v$ is orthogonal to $H^0(X, L_p)$ for every $v \in \mathscr{C}^1(X, L_p)$, by writing $\tilde{v} = e^{2u_p}v + s$ for some $s \in H^0(X, L_p)$, one can observe

(3.9)
$$(e^{2u_p}v + s, e^{2u_p}v + s)_{g_p} = \int_X |\tilde{v}|_{g_p}^2 \frac{\omega^n}{n!} = \int_X (|e^{2u_p}v|_{g_p}^2 + |s|_{g_p}^2) \frac{\omega^n}{n!} \ge \int_X |e^{2u_p}v|_{g_p}^2 \frac{\omega^n}{n!}$$

From (3.8) and (3.9), we have

(3.10)
$$\int_{X} |e^{2u_{p}}v|_{g_{p}}^{2} \frac{\omega^{n}}{n!} \leq \int_{X} |\tilde{v}|_{g_{p}}^{2} \frac{\omega^{n}}{n!} \leq \frac{2}{A_{p}} \int_{X} |\beta|_{g_{p}}^{2} \frac{\omega^{n}}{n!}$$

Let us now estimate $|\beta|_{g_p}^2$ from above. By (3.7) and the first upper bound in (3.5), we obtain the following

$$(3.11) \qquad |\beta|_{g_p}^2 = e^{2u_p} |2\overline{\partial}u_p \wedge v + \overline{\partial}v|_{h_p}^2 \le 2e^{2u_p} (4|\overline{\partial}u_p \wedge v|_{h_p}^2 + |\overline{\partial}v|_{h_p}^2 \le 2e^{2u_p} (\frac{A_p}{16}|v|_{h_p}^2 + |\overline{\partial}v|_{h_p}^2),$$

where, in the first estimation, we use an elementary inequality for norms: $|x + y|^2 \le 2(|x|^2 + |y^2|)$. Finally, putting (3.11) into (3.10) finishes the proof.

4. BERGMAN KERNEL ESTIMATIONS

Theorem 4.1. Let (X, ω) be a compact Kähler manifold with $\dim_{\mathbb{C}} X = n$. Let $\{(L_p, h_p)\}_{p\geq 1}^{\infty}$ be a sequence of holomorphic line bundles with Hermitian metrics h_p of class \mathscr{C}^3 such that (2.7) holds. Assume that $\eta_p = \frac{\|h_p\|_3}{\sqrt{A_p}} \to 0$ as $p \to \infty$. Then we have

(4.1)
$$\lim_{p \to \infty} \frac{K_p(x)}{A_p^n} = 1$$

Proof. We begin by taking a reference cover of the Kähler manifold X, as in Definition 2.1. Selecting $x \in X$ and a corresponding *z*-coordinate system based on Lemma 2.2 at $x \in X$. Then

(4.2)
$$\varphi_p(z) = \Re t_p(z) + \varphi'_p(z) + \widetilde{\varphi_p}(z), \quad \varphi'_p(z) = \sum_{j=1}^n \lambda_j^p |z_j|^2,$$

 φ_p is the weight for the Hermitian metric h_p on $P^n(x, R)$ satisfying the condition (ii) in Lemma 2.2 and t_p is the polynomial of degree at most 2. Let e_p be a local frame of L_p on U_j with the norm $|e_p|_{h_p} = e^{-\varphi_p}$. Next, we choose $R_p \in (0, R/2)$, which we will determine later.

To estimate the norm of a section $S \in H^0(X, L_p)$ at the point $x \equiv 0$, we consider $S = f e_p$, where f is a holomorphic function on $P^n(x, R)$. Utilizing the sub-averaging property for plurisubharmonic functions, we obtain:

(4.3)
$$|S(x)|_{h_p}^2 = |f(0)e^{-t_p(0)}|^2 = |f(0)|^2 e^{-2\Re t_p(0)} \le \frac{\int_{P^n(0,R_p)} |f|^2 e^{-2\Re t_p} e^{-2\varphi'_p} e^{-2\widetilde{\varphi_p}} \frac{\omega^n}{n!}}{\int_{P^n(0,R_p)} e^{-2\varphi'_p} e^{-2\widetilde{\varphi_p}} \frac{\omega^n}{n!}}$$

For the right-hand side of (4.3), by Lemma 2.2, there exists a constant D > 0 such that $-D||h_p||_3|z|^3 \leq \widetilde{\varphi_p}(z) \leq D||h_p||_3|z|^3$, and by considering (2.1) and (4.2) we have,

$$\frac{\int_{P^n(0,R_p)} |f|^2 e^{-2\Re t_p} e^{-2\varphi'_p} e^{-2\widetilde{\varphi_p}} \frac{\omega^n}{n!}}{\int_{P^n(0,R_p)} e^{-2\varphi'_p} e^{-2\widetilde{\varphi_p}} \frac{\omega^n}{n!}} \leq \frac{\int_{P^n(0,R_p)} |f|^2 e^{-2\varphi_p} \frac{\omega^n}{n!}}{\int_{P^n(0,R_p)} e^{-2\varphi'_p} e^{-2\widetilde{\varphi_p}} \frac{dV}{1+DR_p^2}} \leq \frac{(1+DR_p^2) e^{2D||h_p||_3R_p^3} \|S\|_p^2}{\int_{P^n(0,R_p)} e^{-2\varphi'_p} dV}$$

Combining the above inequality with (4.3) yields

(4.4)
$$|S(x)|_{h_p}^2 = |f(0)e^{-t_p(0)}|^2 = |f(0)|^2 e^{-2\Re t_p(0)} \le \frac{(1+DR_p^2)e^{2D}||h_p||_3R_p^3}{\int_{P^n(0,R_p)} e^{-2\varphi'_p}dV} |S||_p^2$$

Let us now estimate the integral in the denominator of (4.4). To do this, we consider the Gaussian-type integrals of finite radius,

(4.5)
$$F(\rho) := \int_{|\xi| \le \rho} e^{-2|\xi|^2} dm(\xi) = \frac{\pi}{2} (1 - e^{-2\rho^2}),$$

where dm is the Lebesgue measure on \mathbb{C} . It is easy to see that F is an increasing function of ρ . We also write

(4.6)
$$F(\infty) := \lim_{\rho \to \infty} \int_{|\xi| \le \rho} e^{-2|\xi|^2} dm(\xi) := \int_{\mathbb{C}} e^{-2|\xi|^2} dm(\xi) = \frac{\pi}{2}.$$

Since

(4.7)
$$\int_{P^n(0,R_p)} e^{-2\varphi'_p} dV = \int_{P^n(0,R_p)} e^{-2(\lambda_1^p |z_1|^2 + \dots + \lambda_n^p |z_n|^2)} dV(z)$$

it is enough to treat the integral

(4.8)
$$\int_{\Delta(0,R_p)} e^{-2\lambda_j^p |z_j|^2} dm(z_j)$$

in order to get a lower bound for the integral (4.7), where $\Delta(0, R_p)$ is the unit closed disk in \mathbb{C} . By the relation (2.12), there exists $p_1 \in \mathbb{N}$ such that, for all $p > p_1$,

(4.9)
$$\frac{3A_p}{4}\omega_x \le c_1(L_p, h_p)_x \le \frac{5A_p}{4}\omega_x,$$

which, on account of (2.1) and (2.13), leads to

(4.10)
$$\frac{3\pi A_p}{8} \le \lambda_j^p \le \frac{5\pi A_p}{8}$$

Let us go back to the integral (4.7),

$$\int_{\Delta(0,R_p)=\{|z_j| \le R_p\}} e^{-2\lambda_j^p |z_j|^2} dm(z_j),$$

which, by a change of variable $\left(\sqrt{\lambda_j^p} z_j = w_j\right)$, equals the following

$$\frac{1}{\lambda_j^p} \int_{\{|w_j| \le R_p \sqrt{\lambda_j^p}\}} e^{-2|w_j|^2} dm(w_j).$$

Now, (4.10) gives $\sqrt{\frac{\lambda_j^p}{A_p}} > \sqrt{\frac{3\pi}{8}} > 1$, which gives $R_p \sqrt{A_p} \le R_p \sqrt{\lambda_j^p}$. Combining this with the fact that F is increasing, we get

(4.11)
$$\int_{\Delta(0,R_p)=\{|z_j|\leq R_p\}} e^{-2\lambda_j^p |z_j|^2} dm(z_j) = \frac{1}{\lambda_j^p} \int_{\{|w_j|\leq R_p\sqrt{\lambda_j^p}\}} e^{-2|w_j|^2} dm(w_j) \geq \frac{F(R_p\sqrt{A_p})}{\lambda_j^p}.$$

Consequently, from (4.7), we have

$$\int_{P^n(0,R_p)} e^{-2\varphi'_p} dV \ge \frac{F(R_p\sqrt{A_p})^n}{\lambda_1^p \dots \lambda_n^p}.$$

Inserting this last inequality in (4.4) give

(4.12)
$$|S(x)|_{h_p}^2 \le \frac{(1+DR_p^2)e^{2D\,\|h_p\|_3\,R_p^3}}{F(R_p\sqrt{A_p})^n}\lambda_1^p\dots\lambda_n^p\,\|S\|_p^2$$

If we take the supremum in (4.12) for all $S \in H^0(X, L_p)$ with $||S||_p = 1$ and use the variational principle (2.4) for K_p , we get

(4.13)
$$K_p(x) \le \frac{(1 + DR_p^2)e^{2D \|h_p\|_3 R_p^3}}{F(R_p\sqrt{A_p})^n} \lambda_1^p \dots \lambda_n^p$$

for any $R_p \in (0, \frac{R}{2})$.

We will now determine a lower bound for K_p by employing L^2 estimations obtained earlier as Theorem 3.2. Let $\kappa : \mathbb{C}^n \to [0, 1]$ be a cut-off function with a compact support in $P^n(0, 2)$, $\kappa = 1$ on $P^n(0, 1)$. By defining $\kappa_p(z) := \kappa(\frac{z}{R_p})$, we consider $H = \kappa_p e^{t_p} e_p$, which is a (smooth) section of L_p , and $|H(x)|_{h_p}^2 = |\kappa_p(x)|^2 e^{2\Re t_p(x)} e^{-\varphi_p(x)}$. We estimate $||H||_p$ from above as follows:

(4.14)
$$\|H\|_{p}^{2} \leq \int_{P^{n}(0,2R_{p})} e^{2\Re t_{p}(x)} e^{-2\varphi_{p}(x)} \frac{\omega_{x}^{n}}{n!} = \int_{P^{n}(0,2R_{p})} e^{-2\varphi_{p}'(x)} e^{-2\widetilde{\varphi}_{p}(x)} \frac{\omega_{x}^{n}}{n!}$$

By using Lemma 2.2 (i) along with the relations (4.6) and (4.11) on the integral at the very right end of the inequality (4.14), we get the following

$$(4.15) \qquad \|H\|_p^2 \le (1+4DR_p^2)e^{16D\|h_p\|_3R_p^3} \int_{P^n(0,2R_p)} e^{-2\varphi_p'} dV \le (1+4DR_p^2)e^{16D\|h_p\|_3R_p^3} \left(\frac{\pi}{2}\right)^n \frac{1}{\lambda_1^p \dots \lambda_n^p}.$$

Let us define $\Phi = \overline{\partial}H$. Noting that $\|\overline{\partial}\kappa_p\|^2 = \|\overline{\partial}\kappa\|^2/R_p^2$, where $\|\overline{\partial}\kappa\|$ is the supremum of $|\overline{\partial}\kappa|$, we deduce the following inequality:

$$\|\Phi\|_{p}^{2} \leq \int_{P^{n}(0,2R_{p})} |\overline{\partial}\kappa_{p}|^{2} e^{-2\varphi_{p}'} e^{-2\widetilde{\varphi}_{p}} \frac{\omega^{n}}{n!} \leq \frac{\|\overline{\partial}\kappa\|^{2}}{R_{p}^{2}} (\frac{\pi}{2})^{n} \frac{(1+4D\,R_{p}^{2})e^{16\,D\,\|h_{p}\|_{3}\,R_{p}^{3}}}{\lambda_{1}^{p}\dots\lambda_{n}^{p}}$$

As $A_p \to \infty$, by using Theorem 3.2, there exists $p > p_0$ such that, for all $p > p_0$, we can find a smooth section Γ of L_p as a solution to the $\overline{\partial}$ -equation for Φ such that $\overline{\partial}\Gamma = \Phi = \overline{\partial}H$ and

(4.16)
$$\|\Gamma\|_{p}^{2} \leq \frac{2}{A_{p}} \|\Phi\|_{p}^{2} \leq \frac{2\|\overline{\partial}\kappa\|}{A_{p}R_{p}^{2}} (\frac{\pi}{2})^{n} \frac{(1+4DR_{p}^{2})e^{(16D\|h_{p}\|_{3}R_{p}^{3})}}{\lambda_{1}^{p}\dots\lambda_{n}^{p}}$$

Given that $H = e_p$ is holomorphic on $P^n(0, R_p)$, Γ is holomorphic on $P^n(0, R_p)$ as well since $\overline{\partial}\Gamma = \overline{\partial}H = 0$ on $P^n(0, R_p)$. Applying estimate (4.12) to Γ on $P^n(0, R_p)$ leads us to the following inequality

(4.17)
$$|\Gamma(x)|_{h_p}^2 \le \frac{(1+DR_p^2)e^{2D\|h_p\|_3R_p^3}}{F(R_p\sqrt{A_p})^n}\lambda_1^p\dots\lambda_n^p\|G\|_p^2$$

(4.18)
$$\leq \frac{2\|\overline{\partial}\kappa\|^2}{A_p R_p^2 F (R_p \sqrt{A_p})^n} (\frac{\pi}{2})^n (1 + 4 D R_p^2)^2 e^{18 D \|h_p\|_3 R_p^3}.$$

Now we will construct a new section $\Lambda := H - \Gamma \in H^0(X, L_p)$. Then, by a basic inequality $|S_1 - S_2|_{h_p}^2 \ge (|S_1|_{h_p} - |S_2|_{h_p})^2$ for norms applied to Λ , combined with (4.17) and the observation $|F(x)|_{h_p} = 1$, we get

(4.19)
$$\begin{aligned} |\Lambda(x)|_{h_p}^2 &\geq (|H(x)|_{h_p} - |\Gamma(x)|_{h_p})^2 \\ &\geq \left(1 - (\frac{\pi}{2})^{n/2} \frac{\sqrt{2} \|\overline{\partial}\kappa\| (1 + 4DR_p^2)}{R_p \sqrt{A_p} F(R_p \sqrt{A_p})^{n/2}} e^{9D\|h_p\|_3 R_p^3} \right)^2. \end{aligned}$$

On the other hand, by (4.15) and (4.16) together with the triangle inequality, we obtain

$$(4.20) \|\Lambda\|_p^2 \le (\|H\|_p + \|\Gamma\|_p)^2 \le (\frac{\pi}{2})^n \frac{1}{\lambda_1^p \dots \lambda_n^p} (1 + 4DR_p^2) e^{16D\|h_p\|_3R_p^3} (1 + \frac{\sqrt{2}\|\overline{\partial}\kappa\|}{R_p\sqrt{A_p}})^2$$

To simplify what we have done so far, we write

(4.21)
$$B_1(R_p) := \left(1 - \left(\frac{\pi}{2}\right)^{n/2} \frac{\sqrt{2} \|\overline{\partial}\kappa\| (1 + 4DR_p^2)}{R_p \sqrt{A_p} F(R_p \sqrt{A_p})^{n/2}} e^{9D\|h_p\|_3 R_p^3}\right)^2$$

and

(4.22)
$$B_2(R_p) := (1 + 4 D R_p^2) e^{16 D \|h_p\|_3 R_p^3} (1 + \frac{\sqrt{2} \|\partial \kappa\|}{R_p \sqrt{A_p}})^2.$$

The variational property (2.4) combined with (4.19) and (4.20) implies

(4.23)
$$K_p(x) \ge \frac{|\Lambda(x)|_{h_p}^2}{\|\Lambda\|_p^2} \ge \frac{\lambda_1^p \dots \lambda_n^p}{(\frac{\pi}{2})^n} \cdot \frac{B_1(R_p)}{B_2(R_p)}$$

For the upper bound (4.13), as above, we put

(4.24)
$$B_3(R_p) := \left(\frac{\pi/2}{F(R_p\sqrt{A_p})}\right)^n (1+D\,R_p^2) \, e^{2\,D\,\|h_p\|_3\,R_p^3}.$$

Observe that

(4.25)
$$(\frac{\pi}{2})^n K_p(x) \le B_3(R_p) \lambda_1^p \dots \lambda_n^p.$$

By our hypothesis $\eta_p = \frac{\|h_p\|_3}{\sqrt{A_p}} \to 0$, and now we determine R_p in the following way

$$R_p := \eta_p^{1/3} ||h_p||_3^{-1/3} = \frac{\eta_p^{-2/3}}{\sqrt{A_p}},$$

which means

$$\eta_p = ||h_p||_3 R_p^3, \ \eta_p^{-2/3} = R_p \sqrt{A_p}.$$

Since $||h_p||_3 \ge 1$ from Subsection 2.1, we have $R_p \le \eta_p^{1/3}$, and so $R_p \to 0$ when $p \to \infty$. All in all, based on R_p , it follows from the quantities $B_1(R_p)$, $B_2(R_p)$ and $B_3(R_p)$ that we find uniform upper and lower bounds for K_p depending only on η_p

(4.26)
$$\frac{B_1(R_p)}{B_2(R_p)} \ge 1 - D'\eta_p^{2/3} \text{ and } B_3(R_p) \le 1 + D'\eta_p^{2/3}.$$

Here D' > 0 denotes a constant that merely depends on the reference cover. We finally consider the following inequality that holds for all $p > p_0$

(4.27)
$$\frac{1}{A_p^n} (\frac{\pi}{2})^n \frac{\lambda_1^p \dots \lambda_n^p}{(\frac{\pi}{2})^n} \cdot \frac{B_1(R_p)}{B_2(R_p)} \le (\frac{\pi}{2})^n \frac{K_p(x)}{A_p^n} \le \frac{1}{A_p^n} B_3(R_p) \lambda_1^p \dots \lambda_n^p$$

which, in light of the findings (2.15), (4.23), (4.25) and (4.26), finishes the proof.

Relying on the proof presented in [BCM], which incorporates methods from [Brn], [CMM] and [Li], we provide a proof for the off-diagonal decay estimate of the Bergman kernels $K_p(x, y)$ associated with the corresponding line bundles (L_p, h_p) in the current setting of diophantine approximation.

Theorem 4.2. Let (X, ω) be a compact Kähler manifold with $\dim_{\mathbb{C}} X = n$. Let $\{(L_p, h_p)\}_{p \ge 1}$ be a sequence of holomorphic line bundles with Hermitian metrics h_p of class \mathscr{C}^3 such that (2.7) is satisfied. Write $\eta_p = \frac{\|h_p\|_3}{\sqrt{A_p}} \to 0$ when $p \to \infty$. Then there exist constants $G, B > 0, p_0 \ge 1$ such that for every $x, y \in X$ and $p > p_0$, the following estimation holds true

(4.28)
$$|K_p(x,y)|_{h_p}^2 \le G e^{-B\sqrt{A_p d(x,y)}} A_p^{2n}$$

Proof. Initially, we select a reference cover for X in accordance with the earlier definition above and choose a large enough $p_0 \in \mathbb{N}$ such that

$$R_p := \frac{1}{\sqrt{A_p}} < \frac{R}{2}$$

and Theorem 3.2 and Theorem 3.3 are valid for all $p > p_0$.

Let $y \in X$ and r > 0. Write

$$B(y,r) := \{ x \in X : d(y,x) < r \},\$$

which is the ball of radius r > 0 centered at y. Choose a constant $\theta \ge 1$ so that for any $y \in X$,

$$P^n(y, R_p) \subseteq B(y, \theta R_p),$$

where $P^n(y, R_p)$ is the (closed) polydisk centered at y given by the coordinates centered at y in view of Lemma 2.2.

<u>*Claim:*</u> There exists a constant D' > 1 such that if $y \in X$, so $y \in P^n(x_j, R_j)$ for some j and z-coordinates centered at y are due to Lemma 2.2, then

(4.29)
$$|S(y)|_{h_p}^2 \le D' A_p^n \int_{P^n(y,R_p)} |S|_{h_p}^2 \frac{\omega^n}{n!}$$

where, as above, $P^n(y, R_p)$ is the (closed) polydisk centered at $y \equiv 0$ in the z-coordinates and S is an arbitrary continuous section of L_p on X which is holomorphic on $P^n(y, R_p)$.

Proof of the Claim: By Lemma 2.2(ii), (L_p, h_p) has a weight φ_p on $P^n(y, R)$ such that

(4.30)
$$\varphi_p(z) = \Re t_p(z) + \varphi'_p(z) + \widetilde{\varphi_p}(z),$$

where $t_p(z), \varphi_p'(z) = \sum_{l=1}^n \lambda_l^p |z_l|^2$ and $\widetilde{\varphi_p}(z)$ satisfies the inequality

(4.31)
$$-D\|h_p\|_3|z|^3 \le \widetilde{\varphi_p}(z) \le D\|h_p\|_3|z|^3$$

for $z \in P^n(y, R)$ (Recall that $R = \min R_j$). Let e_p be a frame of L_p on U_j so that $S = f e_p$, where f is a holomorphic function on $P^n(y, R_p)$ and $|e_p|_{h_p} = e^{-\varphi_p}$. As in the beginning of the proof of Theorem 4.1, we have first the relation, which is nothing but (4.4)

$$(4.32) |S(y)|_{h_p}^2 = |f(0)e^{-t_p(0)}|^2 = |f(0)|^2 e^{-2\Re t_p(0)} \le \frac{(1+DR_p^2)e^{2D||h_p||_3R_p^3} ||S||_p^2}{\int_{P^n(0,R_p)} e^{-2\varphi'_p} dV}.$$

Since

(4.33)
$$\int_{P^n(0,R_p)} e^{-2\varphi'_p} dV = \int_{P^n(0,R_p)} e^{-2(\lambda_1^p |z_1|^2 + \ldots + \lambda_n^p |z_n|^2)} dV(z),$$

as we have done in the proof of Theorem 4.1, it will be sufficient for us to find a lower bound for the integral

(4.34)
$$\int_{\Delta(0,R_p)} e^{-2\lambda_j^p |z_j|^2} dm(z_j)$$

in order to get a lower bound for the whole integral (4.33), where $\Delta(0, R_p)$ is the unit closed disk in \mathbb{C} . By the relation (4.10), there exists $p_1 \in \mathbb{N}$ such that, for all $p > p_1$, $\frac{3\pi A_p}{8} \le \lambda_j^p \le \frac{5\pi A_p}{8}$, and as before, $\sqrt{\frac{\lambda_j^p}{A_p}} > \sqrt{\frac{3\pi}{8}} > 1$. We also observe that

(4.35)
$$F(1) = \frac{\pi}{2}(1 - \frac{1}{e^2}) > \frac{\pi}{2}\frac{21}{25} > 1$$

since $\frac{5}{2} < e < 3$. By the same argument used in the proof of Theorem 4.1, we get

$$\int_{\Delta(0,R_p)=\{|z_j| \le \frac{1}{\sqrt{A_p}}\}} e^{-2\lambda_j^p |z_j|^2} dm(z_j) = \frac{1}{\lambda_j^p} \int_{\left\{|w_j| \le \sqrt{\frac{\lambda_j^p}{A_p}}\right\}} e^{-2|w_j|^2} dm(w_j) \ge \frac{1}{\lambda_j^p},$$

since F(1) > 1 by (4.35) and F is increasing. Consequently, from (4.33), we have

(4.36)
$$\int_{P^n(0,R_p)} e^{-2\varphi'_p} dV \ge \frac{1}{\lambda_1^p \dots \lambda_n^p}$$

Inserting (4.36) into the sub-mean estimation (4.32) and using (4.10), one has

(4.37)
$$|S(y)|_{h_p}^2 \le (1 + DR_p^2) e^{2D||h_p||_3 R_p^3} \lambda_1^p \dots \lambda_n^p \int_{P^n(y,R_p)} |S|_{h_p}^2 \frac{\omega^n}{n!}$$

(4.38)
$$\leq (1 + DR_p^2) e^{2D \|h_p\|_3 R_p^3} (\frac{5\pi}{8})^n A_p^n \int_{P^n(y,R_p)} |S|_{h_p}^2 \frac{\omega^n}{n!}$$

As $R_p = \frac{1}{\sqrt{A_p}} \to 0$ and by our assumption that $\eta_p = \frac{\|h_p\|_3}{\sqrt{A_p}} \to 0$ when $p \to \infty$, one can find a constant D' > 1 such that, for a large enough $p_2 \in \mathbb{N}$,

$$(1 + DR_p^2)e^{2D\|h_p\|_3 R_p^3} (\frac{5\pi}{8})^n \le D'$$

for all $p > \max\{p_0, p_1, p_2\}$. Hence

(4.39)
$$|S(y)|_{h_p}^2 \le D' A_p^n \int_{P^n(y,R_p)} |S|_{h_p}^2 \frac{\omega^n}{n!}$$

which completes the proof of the claim. Let us fix $x \in X$. Then there exists $S_p = S_{p,x} \in H^0(X, L_p)$ such that

$$|S_p(y)|_{h_p}^2 = |K_p(x,y)|_{h_p}^2$$

for all $y \in X$. By Theorem 4.1, there exists a constant D'' > 1 and $p_3 \in \mathbb{N}$ such that, for all $p > p_3$,

where D'' is some constant that depends only on the reference cover. On the other hand,

(4.41)
$$||S_p||_p^2 = \int_X |S_p(y)|_{h_p}^2 \frac{\omega^n(y)}{n!} = \int_X |K_p(x,y)|_{h_p}^2 \frac{\omega^n(y)}{n!} = K_p(x)$$

In the rest of the proof, we proceed with the near-diagonal and off-diagonal estimations of $K_p(x, y)$.

For the near-diagonal estimation, let $y \in X$ and $d(x, y) \leq \frac{4\theta}{\sqrt{A_p}}$. By the variational property (2.4) of $K_p(x)$, the inequality (4.40) and (5.10), we have (4.42)

$$|K_p(x,y)|_{h_p}^2 = |S_p(y)|_{h_p}^2 \le K_p(y) ||S_p||_p^2 \le K_p(x) K_p(y) \le (D'')^2 A_p^{2n} \le e^{4\theta} (D'')^2 A_p^{2n} e^{-\sqrt{A_p d(x,y)}}.$$

We go on with the far off-diagonal estimation. Let $y \in X$, and this time, consider

$$\delta := d(x, y) > 4\theta \frac{1}{\sqrt{A_p}} = 4\theta R_p$$

By the choice of S_p and the claim in the beginning of the proof, we get

(4.43)
$$|S_p(y)|_{h_p}^2 = |K_p(x,y)|_{h_p}^2 \le A_p^n \int_{P^n(y,R_p)} |K_p(x,\zeta)|_{h_p}^2 \frac{\omega_{\zeta}^n}{n!}.$$

We observe that the inclusions

(4.44)
$$P^n(x,R_p) \subset B(x,\frac{\delta}{4}) \text{ and } P^n(y,R_p) \subset \{\zeta \in X : d(x,\zeta) > \frac{3\delta}{4}\}$$

hold.

Let β be a non-negative smooth function on X with the following properties:

$$\beta(\zeta) = 1 \text{ if } d(x,\zeta) \ge \frac{3\delta}{4}$$
$$\beta(\zeta) = 0 \text{ if } d(x,\zeta) \le \frac{\delta}{2}$$
$$\overline{\partial}\beta(\zeta)|^2 \le \frac{c}{\delta^2}\beta(\zeta) \text{ for some } c > 0.$$

According to these data, we first have

(4.45)
$$\int_{P^n(y,R_p)} |K_p(x,\zeta)|_{h_p}^2 \frac{\omega_{\zeta}^n}{n!} \le \int_X |K_p(x,\zeta)|_{h_p}^2 \beta(\zeta) \frac{\omega_{\zeta}^n}{n!}.$$

Using the variational property for $K_p(x)$, the right-hand side of the inequality (4.45) takes the following form:

$$\max\{|K_p(\beta S)|_{h_p}^2 : S \in H^0(X, L_p), \ \int_X |S|_{h_p}^2 \beta \, \frac{\omega^n}{n!} = 1\},\$$

where

$$K_p(\beta S)(x) = \int_X K_p(x,\zeta)(\beta(\zeta) S(\zeta)) \frac{\omega_{\zeta}^n}{n!}$$

is the Bergman projection of the smooth section βS to $H^0(X, L_p)$. Note that when $\beta \equiv 1$, the usual variational formula (2.4) is obtained. Therefore, if we manage to estimate $|K_p(\beta S)|_{h_p}^2$, then we will be done. To this end, to find an upper bound for $|K_p(\beta S)|_{h_p}^2$, we use the decomposition of the space $L^2(X, L_p)$ as below:

(4.46)
$$L^2(X, L_p) = H^0(X, L_p) \oplus Y.$$

(Since $L^2(X, L_p)$ is an Hilbert space and $H^0(X, L_p)$ is a closed subspace of it, such an orthogonal complementary subspace Y always exists). Since $\beta S \in \mathscr{C}^{\infty}(X, L_p) \hookrightarrow L^2(X, L_p)$, it follows from the decomposition (4.46) that there exists an element $u \in Y$ such that

$$u = \beta S - K_p(\beta S).$$

Owing to the inclusion $P^n(x, R_p) \subset B(x, \frac{\delta}{4})$ from (4.44) and $\beta(\zeta) = 0$ for $\zeta \in B(x, \frac{\delta}{2})$ given previously in the proof, we readily have $\beta = 0$ on $P^n(x, R_p)$, so $u = \beta S - K_p(\beta S) = -K_p(\beta S)$, which is holomorphic on $P^n(x, R_p)$ (defined by the coordinates centered at x provided by Lemma 2.2). Therefore, by using the claim in the beginning, one has

(4.47)
$$|K_p(\beta S)|_{h_p}^2 = |u(x)|_{h_p}^2 \le D' A_p^n \int_{P^n(x,R_p)} |u|_{h_p}^2 \frac{\omega^n}{n!}$$

We provide an upper bound for the integral on the right-hand side of (4.47) by Theorem 3.3. For this purpose, let $\tau : [0, \infty) \to (-\infty, 0]$ be a smooth function defined as follows:

$$\tau(x) := \begin{cases} 0, & \text{if } x \le \frac{1}{4} \\ -x, & \text{if } x \ge \frac{1}{2}. \end{cases}$$

Write $\phi_{\delta}(x) = \delta \tau(\frac{x}{\delta})$. Observe that τ' and τ'' have compact supports within the set $[\frac{1}{4}, \frac{1}{2}]$, and so are ϕ'_{δ} and ϕ''_{δ} . This means that there exists a constant $M_0 > 0$ such that $|\phi'_{\delta}(x)| \leq M_0$ and $|\phi''_{\delta}(x)| \leq \frac{M_0}{\delta}$ for all $x \geq 0$. Define the function

$$v_p(\zeta) := \epsilon \sqrt{A_p} \phi_{\delta}(d(x,\zeta)).$$

Since ϕ'_{δ} and ϕ''_{δ} are smooth and have compact supports, we can find a constant $M_1 > 0$ such that

$$(4.48) \|\overline{\partial}v_p\|_{L^{\infty}(X)} \le M_1 \epsilon \sqrt{A_p}$$

(4.49)
$$dd^{c}v_{p} \geq -\frac{M_{1}}{\delta}\epsilon\sqrt{A_{p}}\omega \geq -\frac{M_{1}}{4\theta}\epsilon A_{p}\omega$$

because of the inequality $\delta > 4\theta \frac{1}{\sqrt{A_p}}$. Now we can choose $\epsilon = \frac{1}{8M_1}$ for the conditions of Theorem 3.3 to hold. Since $\tau(x) = 0$ for $x \leq \frac{1}{4}$, we have that

$$v_p(\zeta) = \epsilon \sqrt{A_p} \phi_{\delta}(d(x,\zeta)) = \epsilon \sqrt{A_p} \,\delta \,\tau(\frac{d(x,\zeta)}{\delta}) = 0$$

for $d(x,\zeta) \leq \frac{\delta}{4}$. Also, by the inclusion $P^n(x,R_p) \subset B(x,\frac{\delta}{4})$ given in (4.44), we get $v_p(\zeta) = 0$ on $P^n(x,R_p)$. By the definition of β , it is seen that $\overline{\partial}u = \overline{\partial}(\beta S) = \overline{\partial}\beta \wedge S$ (because S is holomorphic) has the following (compact) support

$$U_{\delta} = \{\zeta \in X : \frac{\delta}{2} \le d(x,\zeta) \le \frac{3\delta}{4}\},\$$

so, for $\zeta \in U_{\delta}$, by the definitions of U_{δ} , τ and ϕ_{δ} we obtain

$$v_p(\zeta) = \epsilon \sqrt{A_p} \,\phi_\delta(d(x,\zeta)) = \epsilon \sqrt{A_p} \,\delta \,\tau(\frac{d(x,\zeta)}{\delta}) = -\epsilon \sqrt{A_p} \,d(x,\zeta) \le -\epsilon \sqrt{A_p} \,\frac{\delta}{2}$$

By Theorem 3.3 and the definition of β , we have

$$\begin{split} \int_{P^n(x,R_p)} |u|_{h_p}^2 \frac{\omega^n}{n!} &\leq \int_X |u|_{h_p}^2 e^{v_p} \frac{\omega^n}{n!} \\ &\leq \frac{16}{3A_p} \int_{U_\delta} |\overline{\partial}(\beta S)|_{h_p}^2 e^{v_p} \frac{\omega^n}{n!} \\ &\leq \frac{16c}{3A_p \delta^2} e^{-\epsilon \sqrt{A_p}\delta} \int_{U_\delta} |S|_{h_p}^2 \beta \frac{\omega^n}{n!} \\ &\leq \frac{c}{3} e^{-\sqrt{A_p}\delta} \leq c e^{-\sqrt{A_p}\delta}. \end{split}$$

Plugging this last inequality into (4.47) gives

$$|K_p(\beta S)|_{h_p}^2 \le D' A_p^n c e^{-\epsilon \sqrt{A_p}\delta},$$

which gives, by using the inequality (4.45),

$$\int_{P^n(y,R_p)} |K_p(x,\zeta)|^2_{h_p} \frac{\omega_{\zeta}^n}{n!} \le D' A_p^n c \, e^{-\epsilon \sqrt{A_p} d(x,y)}.$$

From the inequality (4.43), we infer

(4.50)
$$|K_p(x,y)|_{h_p}^2 \le c \, (D')^2 A_p^{2n} \, e^{-\epsilon \sqrt{A_p} d(x,y)}$$

which finalizes the proof.

г				1
ь	_	_	_	4

4.1. Linearization. Let $V \subset X$, $U \subset \mathbb{C}^n$ be open subsets, and $x_0 \in V$, $0 \in U$. Let us take a (Kähler) coordinate chart as follows $\gamma : (V, x_0) \to (U, 0)$, $\gamma(x_0) = 0$. We will use the following notation, so-called linearization of the coordinates on the Kähler manifold X: For any $u, v \in \mathbb{C}^n$, we write $\gamma^{-1}(u) = x_0 + u$ and $\gamma^{-1}(v) = x_0 + v$, and

(4.51)
$$K_p(\gamma^{-1}(u), \gamma^{-1}(v)) := K_p(x_0 + u, x_0 + v).$$

Since $0 \in \mathbb{C}^n$ and \mathbb{C}^n is a complex vector space, we can write v = 0 + v and u = 0 + u. Linearization means that when we translate $0 \in \mathbb{C}^n$ by u (or by v for that matter), by thinking of $0 \in \mathbb{C}^n$ as $x_0 \in X$, we can also write $\gamma^{-1}(u) = x_0 + u$, so in local coordinates we express the difference between $\gamma^{-1}(u)$ and x_0 (not meaningful in X) by the difference u - 0 (meaningful in \mathbb{C}^n because \mathbb{C}^n is a complex vector space). This is also called the abuse of notation in, for instance, [SZ08] and [SZ10].

Modifying the argument in ([Bay16], Theorem 2.3), we consider the following holomorphic functions

$$\Gamma_{p}(u,v) = \frac{K_{p}(\frac{u}{\sqrt{A_{p}}}, \frac{\overline{v}}{\sqrt{A_{p}}})e^{-t_{p}(\frac{u}{\sqrt{A_{p}}}) - \overline{t_{p}(\frac{\overline{v}}{\sqrt{A_{p}}})}}{A_{p}^{n} e^{\frac{2}{A_{p}}\langle\Lambda_{p}u,\overline{v}\rangle}} \\ = \frac{K_{p}(\frac{u}{\sqrt{A_{p}}}, \frac{\overline{v}}{\sqrt{A_{p}}})e^{-t_{p}(\frac{u}{\sqrt{A_{p}}}) - \overline{t_{p}(\frac{\overline{v}}{\sqrt{A_{p}}})}}e^{-\widetilde{\varphi_{p}}(\frac{u}{\sqrt{A_{p}}})}e^{-\widetilde{\varphi_{p}}(\frac{\overline{v}}{\sqrt{A_{p}}})}}{A_{p}^{n} e^{\frac{2}{A_{p}}\langle\Lambda_{p}u,\overline{v}\rangle}}e^{\widetilde{\varphi_{p}}(\frac{\overline{v}}{\sqrt{A_{p}}})}e^{\widetilde{\varphi_{p}}(\frac{$$

on $\Omega = \{(u, v) : u, v \in P^n(0, R)\}$, where $\Lambda_p := Diag[\lambda_1^p, \dots, \lambda_n^p]$, which is a diagonal matrix whose diagonal entries λ_j^p are positive from the discussions in Section 2. Let $\Omega_0 := \{(u, \overline{u}) : u \in P^n(0, R)\}$. It follows from Theorem 4.1 and Lemma 2.2 (ii) that $\Gamma_p \to 1$ on Ω_0 . Observe that since Γ_p is uniformly bounded on Ω , there is a subsequence $\{\Gamma_{p_d}\}$ such that $\Gamma_{p_d} \to \Gamma_0$ uniformly on Ω , where we must have that $\Gamma_0 \equiv 1$ on Ω_0 . Since Ω_0 is a maximally totally real submanifold, we get $\Gamma_0 = 1$ on the whole Ω . Since this argument can be applied to any subsequence of Γ_p , we see that $\Gamma_p \to 1$. We make now an observation for $|\Gamma_p(u, v)|^2$ that will be used in our main theorem. Since Λ_p has positive diagonal entries, its square root $\Lambda_p^{1/2}$ is defined, so we have

(4.52)
$$\langle \Lambda_p u, \overline{v} \rangle = \langle \Lambda_p^{1/2} u, \Lambda_p^{1/2} \overline{v} \rangle.$$

$$(4.53) \qquad |\Gamma_{p}(u,v)|^{2} = \frac{|K_{p}(\frac{u}{\sqrt{A_{p}}}, \frac{\overline{v}}{\sqrt{A_{p}}})|^{2}e^{-2\Re t_{p}(\frac{u}{\sqrt{A_{p}}})}e^{-2\Re t_{p}(\frac{\overline{v}}{\sqrt{A_{p}}})}{A_{p}^{2n} e^{\frac{4}{A_{p}}\Re(\langle\Lambda_{p}u,\overline{v}\rangle)}} \\ = \frac{|K_{p}(\frac{u}{\sqrt{A_{p}}}, \frac{\overline{v}}{\sqrt{A_{p}}})|^{2}e^{-2\Re t_{p}(\frac{u}{\sqrt{A_{p}}})}e^{-2\Re t_{p}(\frac{\overline{v}}{\sqrt{A_{p}}})}{A_{p}^{2n} e^{2\Sigma_{j=1}^{n}\frac{\lambda_{j}^{p}}{A_{p}}|u_{j}|^{2}}e^{2\Sigma_{j=1}^{n}\frac{\lambda_{j}^{p}}{A_{p}}|v_{j}|^{2}}e^{-2\Sigma_{j=1}^{n}\frac{\lambda_{j}^{p}}{A_{p}}|u_{j}-\overline{v_{j}}|^{2}} \\ = \frac{|K_{p}(\frac{u}{\sqrt{A_{p}}}, \frac{\overline{v}}{\sqrt{A_{p}}})|^{2}e^{-2\varphi_{p}(\frac{u}{\sqrt{A_{p}}})}e^{-2\varphi_{p}(\frac{\overline{v}}{\sqrt{A_{p}}})}e^{\widetilde{\varphi_{p}}(\frac{\overline{v}}{\sqrt{A_{p}}})}e^{\widetilde{\varphi_{p}}(\frac{\overline{v}}{\sqrt{A_{p}}})}, \\ A_{p}^{2n} e^{-2\Sigma_{j=1}^{n}\frac{\lambda_{j}^{p}}{A_{p}}|u_{j}-\overline{v_{j}}|^{2}} \end{aligned}$$

where, in the second equality, we have used (4.52) and the polarization identity

$$\Re(\langle \mathbf{x}, \mathbf{y} \rangle) = \frac{1}{2} (\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2)$$

for the vectors $\mathbf{x} = \Lambda_p^{1/2} u$ and $\mathbf{y} = \Lambda_p^{1/2} \overline{v}$, and in the third equality, we take into account the representation $\varphi_p(z) = \Re t_p(z) + \sum_{j=1}^m \lambda_j^p |z_j|^2 + \widetilde{\varphi}_p(z)$ from Lemma 2.2 (ii). By the limit argument made above regarding the holomorphic functions Γ_p , the expression (4.53) for $|\Gamma_p(u, v)|^2$ and Lemma 2.2(ii), we have

(4.54)
$$\frac{|K_p(\frac{u}{\sqrt{A_p}}, \frac{\overline{v}}{\sqrt{A_p}})|^2 e^{-2\varphi_p(\frac{u}{\sqrt{A_p}})} e^{-2\varphi_p(\frac{\overline{v}}{\sqrt{A_p}})}}{A_p^{2n} e^{-2\Sigma_{j=1}^n \frac{\lambda_j^p}{A_p}|u_j - \overline{v_j}|^2}} \to 1$$

as $p \to \infty$. By linearization (4.51) on the coordinate polydisk $P^n(x, R) \subset U_j$, where we have the Kähler coordinates at the point $x \equiv 0$ provided by Lemma 2.2 for (4.54), we obtain

(4.55)
$$\frac{|K_p(x+\frac{u}{\sqrt{A_p}},x+\frac{\overline{v}}{\sqrt{A_p}})|^2 e^{-2\varphi_p(x+\frac{u}{\sqrt{A_p}})} e^{-2\varphi_p(x+\frac{\overline{v}}{\sqrt{A_p}})}}{A_p^{2n} e^{-2\sum_{j=1}^n \frac{\lambda_j^p}{A_p}|u_j-\overline{v_j}|^2}} \to 1 \quad \text{when } p \to \infty.$$

5. Asymptotic Normality

5.1. Random holomorphic sections and random zero currents of integration. Let $s \in H^0(X, L) \setminus \{0\}$. We denote by Z_s the set of zeros of s. Z_s is a purely 1-codimensional analytic variety of X. By the symbol $[Z_s]$, we mean the current of integration on Z_s , defined by

$$\langle [Z_s], \varphi \rangle = \int_{Z_s} \varphi_s$$

here $\varphi \in \mathcal{D}^{n-1,n-1}(X)$, which represents the space of test forms of type (n-1,n-1).

A complex random variable W is said to be standard Gaussian in case $W = X + \sqrt{-1}Y$, where X and Y are i.i.d. centered Gaussian distributions of variance 1/2. The Poincaré-Lelong formula is one indispensable instrument that will be of use in the proof of our main theorem. Let us state it for a zero current associated with the zero set of a holomorphic section $s_p \in H^0(X, L_p)$.

(5.1)
$$[Z_{s_p}] = dd^c \log |s_p|_{h_p}^2 + c_1(L_p, h_p).$$

Here and throughout $dd^c = \frac{\sqrt{-1}}{\pi} \partial \overline{\partial}$.

5.2. Asymptotic normality of random zero currents. Now we are ready to state and prove our main theorem. For the proof, we use the arguments from [SZ10].

Theorem 5.1. Let $\{(L_p, h_p)\}_{p\geq 1}^{\infty}$ be a sequence of positive holomorphic line bundles over a compact Kähler manifold (X, ω) of dimension n with diophantine condition (2.7) and Hermitian metrics of class \mathscr{C}^3 such that $\frac{\|h_p\|_3}{\sqrt{A_p}} \to 0$ as $p \to \infty$. Suppose that $H^0(X, L_p)$ is endowed with the standard Gaussian probability measure for all $p \geq 1$. Let $s_p \in H^0(X, L_p)$ and ϕ be a real valued (n-1, n-1)-form on X with \mathscr{C}^3 -coefficients and $dd^c \phi \neq 0$. Then the distributions of the random variables

(5.2)
$$\frac{\langle [Z_{s_p}], \phi \rangle - \mathbb{E}\langle [Z_{s_p}], \phi \rangle}{\sqrt{\operatorname{Var}\langle [Z_{s_p}], \phi \rangle}}$$

weakly converge towards the standard (real) Gaussian distribution $\mathcal{N}(0,1)$ as $p \to \infty$.

Proof. To begin with, we modify the information about analytic functions from the introduction to suit our present setting. The normalized Gaussian processes α_p on X will be constructed as follows: We take a measurable section e_{L_p} of L_p such that $e_{L_p} : X \to L_p$ with $|e_{L_p}(x)|_{h_p} = 1$ for any $x \in X$. We pick now an orthonormal basis $\{s_p^j\}_{j=1}^{d_p}$ of $H^0(X, L_p)$, where $s_p^j = \varphi_p^j e_{L_p}$. Let us write

(5.3)
$$f_p^j(x) = \frac{\varphi_p^j(x)}{\sqrt{K_p(x,x)}}, \ j = 1, 2, \dots, d_p.$$

Notice that $|\varphi_p^j| = |s_p^j|_{h_p}$ and $\sum_{j=1}^{d_p} |f_p^j(x)|^2 = 1$ by the relation (2.3). Therefore, we can express a normalized complex Gaussian process on X for each $p \in \mathbb{N}$ as follows:

(5.4)
$$\alpha_p = \sum_{j=1}^{d_p} b_j f_p^j$$

where the coefficients b_j are i.i.d. complex Gaussian centered random variables with variance one. We observe that a random holomorphic section $s_p = \sum_{j=1}^{d_p} b_j s_p^j$ can be represented as

(5.5)
$$s_p = \sum_{j=1}^{d_p} b_j s_p^j = \sqrt{K_p(x, x)} \alpha_p e_{L_p},$$

which indicates the presence of the normalized complex Gaussian process. The relation (5.5) gives us that

(5.6)
$$|\alpha_p(x)| = \frac{|s_p^j(x)|_{h_p}}{\sqrt{K_p(x,x)}}.$$

We proceed to compute the covariance functions C_p of the complex Gaussian processes α_p . We observe from the fact that the complex Gaussian random coefficients b_j in (5.4) are centered, i.i.d., and have variance one

(5.7)
$$\operatorname{Var}[b_j] = \mathbb{E}[|b_j|^2] = 1, \ \mathbb{E}[b_k \bar{b}_l] = 0 \ \text{if } k \neq l$$

By the relation (1.3), (5.4) and (5.7), we have

(5.8)
$$C_p(x,y) = \mathbb{E}[\sum_{j=1}^{d_p} b_j f_p^j(x) \sum_{j=1}^{d_p} b_j f_p^j(y)] = \sum_{j=1}^{d_p} f_p^j(x) \overline{f_p^j(y)}.$$

Since $\{s_p^j\}_{j=1}^{d_p}$ is an orthornormal basis, due to the representation $s_p^j = \varphi_p^j e_{L_p}$ and the relation (5.3), we have $\varphi_p^k(x)\overline{\varphi_p^l(y)} = 0$ whenever $k \neq l$. Combining this with (5.8) yields

(5.9)
$$|\mathcal{C}_p(x,y)| = \sqrt{\sum_{j=1}^{d_p} |f_p^j(x)|^2 |f_p^j(y)|^2}.$$

Next, after a series of computations, we arrive at

(5.10)
$$|K_p(x,y)|_{h_{p,x}\otimes h_{p,y}} = \sqrt{\sum_{j=1}^{d_p} |s_p^j(x)|_{h_{p,x}}^2 |s_p^j(y)|_{h_{p,y}}^2}$$

By putting together (5.9), (5.10) and (5.3), we find

(5.11)
$$\widehat{\mathcal{K}}_p(x,y) = |\mathcal{C}_p(x,y)|$$

because $|s_p^j|_{h_p} = |\varphi_p^j|$.

Take $\lambda(\rho) = \log \rho$ and $(G, \sigma) = (X, \frac{\omega^n}{n!})$. In the rest of the proof, to make the notation lighter, we will write $d\vartheta_X := \frac{w^n}{n!}$ for the (Riemannian) volume form on X. Let us fix a (n-1, n-1) real-valued form ϕ with \mathscr{C}^3 coefficients. Then

(5.12)
$$dd^c\phi = \frac{\sqrt{-1}}{\pi}\partial\bar{\partial}\phi = \psi \,d\vartheta_X,$$

where the function ψ is a real-valued \mathscr{C}^1 function on *X*. By invoking the Poincaré-Lelong formula (5.1), the non-linear random functional given by (1.4) assumes the subsequent form in our case,

(5.13)
$$\mathcal{F}_{p}^{\psi}(\alpha_{p}) = \int_{X} \left(\log |s_{p}|_{h_{p}} - \log \sqrt{K_{p}(x,x)} \right) \frac{\sqrt{-1}}{\pi} \partial \overline{\partial} \phi(x) = \langle [Z_{s_{p}}], \phi \rangle + \zeta_{p,L_{p}},$$

where $\zeta_{p,L_p} := \langle -c_1(L_p, h_p) - \log \sqrt{K_p(x, x)}, \phi \rangle$, namely, ζ_{p,L_p} is some constant depending only on the line bundle L_p and the dimension of the Kähler manifold X. Thanks to a standard property of variance, the expression (5.13) shows that $\mathcal{F}_p^{\psi}(\alpha_p)$ and $\langle [Z_{s_p}], \phi \rangle$ have the equal variances.

For the remaining part of the proof, our goal is to validate the fulfillment of the requirements (i) and (ii) of Theorem 1.1 for the current setting. First, with $\lambda(\rho) = \log \rho$ being increasing, we only consider the case where $\nu = 1$. To use both far off-diagonal and near-diagonal asymptotics, we split the integration regions accordingly: $d(x, y) \leq \frac{\log A_p}{\sqrt{A_p}}$ and $d(x, y) \geq \frac{\log A_p}{\sqrt{A_p}}$. Let us start with the simpler condition (ii). For the integral on the far off-diagonal set where $d(x, y) \geq \frac{\log A_p}{\sqrt{A_p}}$, by Theorem 4.2, we have,

$$\lim_{n \to \infty} \sup_{x \in X} \int_{d(x,y) \ge \frac{\log A_p}{\sqrt{A_p}}} \widehat{\mathcal{K}}_p(x,y) d\vartheta_X(y) \le \lim_{p \to \infty} \sup_{x \in X} \int_{d(x,y) \ge \frac{\log A_p}{\sqrt{A_p}}} Ge^{-B\sqrt{A_p}d(x,y)} d\vartheta_X(y) = 0.$$

For the integral over the near-diagonal set where $d(x, y) \leq \frac{\log A_p}{\sqrt{A_p}}$, due to the relations (5.11) and (1.3), we get

$$\lim_{n \to \infty} \sup_{x \in X} \int_{d(x,y) \le \frac{\log A_p}{\sqrt{A_p}}} \widehat{\mathcal{K}}_p(x,y) d\vartheta_X(y) \le \lim_{n \to \infty} \sup_{x \in X} \int_{d(x,y) \le \frac{\log A_p}{\sqrt{A_p}}} 1 \ d\vartheta_X(y) = 0$$

Our next step is to affirm the condition (i). For the integral on the far off-diagonal set where $d(x,y) \geq \frac{\log A_p}{\sqrt{A_p}}$, as $p \to \infty$, the integrand of the numerator approaches zero more rapidly compared to that of the denominator because, by Theorem 4.2, the corresponding decaying orders (to zero) for numerator and denominator are $O(A_p^{-\epsilon})$ and $O(A_p^{-\epsilon/2})$, respectively.

Finally, we verify that the lower limit below will be strictly positive on the near-diagonal set where $\{|v| \leq \frac{\log A_p}{\sqrt{A_p}}\}$:

(5.14)
$$\liminf_{p \to \infty} \frac{\int_X \int_{|v| \le \log A_p} \widehat{\mathcal{K}}_p^2(x, x + \frac{v}{\sqrt{A_p}}) \phi(x) \phi(x + \frac{v}{\sqrt{A_p}}) \, dv \, d\vartheta_X(x)}{\int_{|v| \le \log A_p} \widehat{\mathcal{K}}_p(x, x + \frac{v}{\sqrt{A_p}}) \, dv} > 0$$

Let

(5.15)
$$J(p) := \frac{\int_X \int_{|v| \le \log A_p} \widehat{\mathcal{K}}_p^2(x, x + \frac{v}{\sqrt{A_p}}) \phi(x) \phi(x + \frac{v}{\sqrt{A_p}}) \, dv \, d\vartheta_X(x)}{\int_{|v| \le \log A_p} \widehat{\mathcal{K}}_p(x, x + \frac{v}{\sqrt{A_p}}) \, dv}$$

Let us examine the numerator and the denominator separately. By using the left part of the inequality (4.27) for the denominator and the right part of the same inequality for the numerator, the linearization (4.51) on the neighborhood U_j , where we have the Kähler coordinates at the point x on the polydisk $P^n(x, R) \subset U_j$ provided by Lemma 2.2, we get

(5.16)
$$\int_{X} \int_{|v| \le \log A_p} \widehat{\mathcal{K}}_p^2(x, x + \frac{v}{\sqrt{A_p}}) \phi(x) \phi(x + \frac{v}{\sqrt{A_p}}) \, dv \, d\vartheta_X(x) \\ \ge \int_{X} \int_{|v| \le \log A_p} (\frac{4}{5})^{2n} \frac{|K_p(x, x + \frac{v}{\sqrt{A_p}})|_{h_p}^2 \, \phi(x) \, \phi(x + \frac{v}{\sqrt{A_p}})}{A_p^{2n} \, (1 + D' \eta_p^{2/3})^2} \, dv \, d\vartheta_X(x) := I_1(p).$$

and

(5.17)
$$\int_{|v| \le \log A_p} \widehat{\mathcal{K}}_p(x, x + \frac{v}{\sqrt{A_p}}) \, dv \, \le \int_{|v| \le \log A_p} (\frac{4}{3})^n \frac{|K_p(x, x + \frac{v}{\sqrt{A_p}})|_{h_p}}{A_p^n \left(1 - D' \eta_p^{2/3}\right)} \, dv := I_2(p),$$

which implies, by extracting the weight functions and multiplying and dividing both the integrand of I_1 and I_2 by $e^{-2\sum_{j=1}^n \frac{\lambda_j^p}{A_p} |v_j|^2}$ and $e^{-\sum_{j=1}^n \frac{\lambda_j^p}{A_p} |v_j|^2}$ respectively

(5.18)

$$I_{1}(p) = \int_{X} \int_{|v| \le \log A_{p}} \left(\frac{4}{5}\right)^{2n} \frac{|K_{p}(x, x + \frac{v}{\sqrt{A_{p}}})|^{2} e^{-2\varphi_{p}(x)} e^{-2\varphi_{p}(x + \frac{v}{\sqrt{A_{p}}})} \phi(x) \phi(x + \frac{v}{\sqrt{A_{p}}})}{A_{p}^{2n} \left(1 + D' \eta_{p}^{2/3}\right)^{2} e^{-2\sum_{j=1}^{n} \frac{\lambda_{j}^{p}}{A_{p}} |v_{j}|^{2}}} e^{-2\sum_{j=1}^{n} \frac{\lambda_{j}^{p}}{A_{p}} |v_{j}|^{2}} dv \, d\vartheta_{X}(x)$$

and

(5.19)
$$I_2(p) = \int_{|v| \le \log A_p} \left(\frac{4}{3}\right)^n \frac{|K_p(x, x + \frac{v}{\sqrt{A_p}})|e^{-\varphi_p(x)}e^{-\varphi_p(x + \frac{v}{\sqrt{A_p}})}}{A_p^n \left(1 - D'\eta_p^{2/3}\right)e^{-\sum_{j=1}^n \frac{\lambda_j^p}{A_p}|v_j|^2}} e^{-\sum_{j=1}^n \frac{\lambda_j^p}{A_p}|v_j|^2} dv.$$

By (5.16), (5.17), (5.18) and (5.19), we obtain

(5.20)

$$J(p) \geq \frac{I_{1}(p)}{I_{2}(p)} = \frac{\int_{X} \int_{|v| \leq \log A_{p}} \left(\frac{4}{5}\right)^{2n} \frac{|K_{p}(x, x + \frac{v}{\sqrt{A_{p}}})|^{2} e^{-2\varphi_{p}(x)} e^{-2\varphi_{p}(x + \frac{v}{\sqrt{A_{p}}})} \phi(x) \phi(x + \frac{v}{\sqrt{A_{p}}})}{A_{p}^{2n}(1 + D' \eta_{p}^{2/3})^{2} e^{-2\Sigma_{j=1}^{n}} \frac{\lambda_{j}^{p}}{A_{p}^{j}|v_{j}|^{2}}}{\int_{|v| \leq \log A_{p}} \left(\frac{4}{3}\right)^{n} \frac{|K_{p}(x, x + \frac{v}{\sqrt{A_{p}}})|e^{-\varphi_{p}(x)} e^{-\varphi_{p}(x + \frac{v}{\sqrt{A_{p}}})}}{A_{p}^{n}(1 - D' \eta_{p}^{2/3})e^{-\Sigma_{j=1}^{n}} \frac{\lambda_{j}^{p}}{A_{p}^{p}|v_{j}|^{2}}}{e^{-\Sigma_{j=1}^{n}} \frac{\lambda_{j}^{p}}{A_{p}^{p}|v_{j}|^{2}}} e^{-\Sigma_{j=1}^{n} \frac{\lambda_{j}^{p}}{A_{p}^{p}}|v_{j}|^{2}} dv$$

Since $\phi \in \mathscr{C}^1$, it follows that $\phi(x + \frac{v}{\sqrt{A_p}}) = \phi(x) + O(\frac{v}{\sqrt{A_p}})$. Now, as $p \to \infty$ on the inequality (5.20), and making use of (4.54) and (2.14) yield that

$$\liminf_{p \to \infty} J(p) \ge \frac{\int_X \phi^2(x) d\vartheta_X(x) \int_{\mathbb{C}^n} (\frac{4}{5})^{2n} e^{-\pi \sum_{j=1}^n |v_j|^2} dv}{\int_{\mathbb{C}^n} (\frac{4}{3})^n e^{-\frac{\pi}{2} \sum_{j=1}^n |v_j|^2} dv} = \frac{6^n}{5^{2n}} \int_X \phi^2(x) d\vartheta_X(x) > 0,$$

which ends the proof.

In the papers [CMM] and [BCM], the authors use a more general hypothesis between the first Chern forms and the Kähler form. More precisely, they assume

(5.21)
$$c_1(L_p, h_p) \ge a_p \omega$$
 for all $p \ge 1$ and $a_p > 0$, such that $a_p \to \infty$.

In our setting, the role of a_p is played by A_p (which was defined as $A_p = \int_X c_1(L_p, h_p) \wedge \omega^{n-1}$ in [CMM] and[BCM]), and there are two different limits because of the diophantine approximation relation (2.7) between λ_j^p and A_p . In the case of (5.21), we do not have (2.14) (and consequently (2.15)). However, as Theorem 1.3 in [CMM] shows, there still exists a limit: $\lim_{p\to\infty} \frac{K_p(x)}{\lambda_1^p...\lambda_p^n} = (\frac{2}{\pi})^n$.

Despite the existence of the limit in terms of λ_j^p , we do not know whether the limit $\lim_{p\to\infty} \frac{\lambda_j^p}{A_p}$ exists, which is crucial in the proof of Theorem 5.1. Therefore, the arguments followed in this paper cannot be used to prove a central limit theorem in this framework.

Acknowledgement: We thank Turgay Bayraktar for his interest in and comments on this work.

REFERENCES

- [Bay16] T. Bayraktar. Asymptotic normality of linear statistics of zeros of random polynomials. *Proc. Amer. Math. Soc.*, 145, 2917–2929, 2017
- [BCM] T. Bayraktar, D. Coman, and G. Marinescu. Universality results for zeros of random holomorphic sections. *Trans. Amer. Math. Soc.*, 373(6): 3765-3791(DOI:10.1090/tran/7807), 2020.
- [BCHM18] T. Bayraktar, D. Coman, H. Herrmann, and G. Marinescu. A survey on zeros of random holomorphic sections. *Dolomites Res. Notes Approx.*, 11(4):1–19, 2018.

[Brn] B. Berndtsson. Bergman kernels related to Hermitian line bundles over compact complex manifolds. *Explorations in complex and Riemannian geometry*, Contemp. Math. 332, 1–17, 2003.

- [BBL] E. Bogomolny, O. Bohigas, and P. Leboeuf, *Quantum chaotic dynamics and random polynomials*, J. Statist. Phys. **85**(5-6), 639–679 (1996).
- [BP] A. Bloch and G. Pólya. On the roots of certain algebraic equations. Proc. London Math. Soc., 33:102–114, 1932.

[BG] A. Bojnik, O. Günyüz., Random systems of holomorphic sections of a sequence of line bundles on compact Kähler manifolds. *arXiv:2401.08243*.

- [BSZ] P. Bleher, B. Shiffman, and S. Zelditch. Universality and scaling of correlations between zeros on complex manifolds. *Invent. Math.*, **142**(2):351–395, 2000.
- [CMM] D. Coman, X. Ma, and G. Marinescu. Equidistribution for sequences of line bundles on normal Kähler spaces *Geom. Topol.*, 21(2): 923–962, 2017.
- [CLMM] D. Coman, W. Lu, X. Ma, and G. Marinescu. Bergman kernels and equidistribution for sequences of line bundles on Kähler manifolds. *Adv. Math.*, 414: 923–962, 2017.
- [Dem12] J.-P. Demailly. *Complex analytic and differential geometry*. http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.p 2012.
- [Dem82] J.-P. Demailly. Estimations L^2 pour l'opérateur $\tilde{\partial}$ d'un fibré vectoriel holomorphe semi-positif au-dessus d'une variété Kählérienne complète. Ann. Sci. École Norm. Sup., 15(3): 457-511, 1982.

[DS06] T.-C. Dinh and N. Sibony. Distribution des valeurs de transformations méromorphes et applications. *Comment. Math. Helv.*, 81(1):221–258, 2006.

- [Kli91] M. Klimek. *Pluripotential theory*, volume 6 of *London Mathematical Society Monographs*. *New Series*. The Clarendon Press, Oxford University Press, New York, 1991. Oxford Science Publications.
- [Li] N. Lindholm. Sampling in weighted L^p spaces of entire functions in \mathbb{C}^n and estimates of the Bergman kernel. J. *Funct. Anal.*, 182: 390–426, 2001.
- [MM1] X. Ma and G. Marinescu. *Holomorphic Morse Inequalities and Bergman Kernels*, volume 254 of *Progress in Mathematics*. Birkhäuser, Basel, 2007.
- [Ber] R. Berman. Determinantal point processes and fermions on polarized complex manifolds: Bulk universality *arXiv:* 0811.3341.
- [ET50] P. Erdös and P. Turán. On the distribution of roots of polynomials. Ann. of Math. (2), 51:105–119, 1950.
- [HAM56] J. M. Hammersley. The zeros of a random polynomial. In *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, 1954–1955, vol. II*, pages 89–111, Berkeley and Los Angeles, 1956. University of California Press.
- [Hann] J. H. Hannay Chaotic analytic zero points: exact statistics for those of a random spin state. *Journal of Physics A*, 29: 101–105, 1996.

- [Kac43] M. Kac. On the average number of real roots of a random algebraic equation. *Bull. Amer. Math. Soc.*, 49:314–320, 1943.
- [LO43] J. E. Littlewood and A. C. Offord. On the number of real roots of a random algebraic equation. III. *Rec. Math.* [*Mat. Sbornik*] *N.S.*, 12(54):277–286, 1943.
- [MM1] X. Ma and G. Marinescu. *Holomorphic Morse Inequalities and Bergman Kernels*, volume 254 of *Progress in Mathematics*. Birkhäuser, Basel, 2007.
- [NS] F. Nazarov and M. Sodin. Correlation functions for random complex zeros: strong clustering and local universality. *Comm.Math.Phys.*, 310(1):75–98,2012.
- [NV98] S. Nonnenmacher and A. Voros. Chaotic eigenfunctions in phase space. J. Statist. Phys., 92(3-4):431–518, 1998.
- [SHSM] M. Shub, S. Smale., Complexity of Bezout's theorem. II. Volumes and Probabilities, Computational Algebraic Geometry (Nice, 1992) *Birkhäuser*, Progr. Math. 109: 267–285, 1993.
- [ST] M. Sodin and B. Tsirelson. Random complex zeroes. I. Asymptotic normality Israel J. Math., 144:125–149, 2004. 2, 26
- [SZ99] B. Shiffman and S. Zelditch. Distribution of zeros of random and quantum chaotic sections of positive line bundles. *Comm. Math. Phys.*, 200(3):661–683, 1999.
- [SZ08] B. Shiffman and S. Zelditch. Number variance of random zeros on complex manifolds. *Geom. Funct. Anal.*, 18: 1422–1475, 2008.
- [SZ10] B. Shiffman and S. Zelditch. Number variance of random zeros on complex manifolds, II: Smooth Statistics. *Pure and Applied Mathematics Quarterly*, 6(4): 1145–1167, 2010.
 - FACULTY OF ENGINEERING AND NATURAL SCIENCES, SABANCI UNIVERSITY, İSTANBUL, TURKEY *Email address*: afrimbojnik@sabanciuniv.edu
 - ${\it Email\ address:\ ozangunyuz@alumni.sabanciuniv.edu}$