RECTIFIABLE REIFENBERG AND UNIFORM POSITIVITY UNDER ALMOST CALIBRATIONS

NICK EDELEN, AARON NABER AND DANIELE VALTORTA

ABSTRACT. The Reifenberg theorem [Rei60] tells us that if a set $S \subseteq B_2 \subseteq \mathbb{R}^n$ is uniformly close on all points and scales to a k-dimensional subspace, then S is Hölder homeomorphic to a k-dimensional Euclidean ball. In general this is sharp, for instance such an S may have infinite volume, be fractal in nature, and have no rectifiable structure.

The goal of this note is to show that we can improve upon this for an almost calibrated Reifenberg set, or more generally under a positivity condition in the context of an ϵ -calibration Ω . An ϵ -calibration is very general, the condition holds locally for all continuous k-forms such that $\Omega[L] \leq 1 + \epsilon$ for all k-planes L. We say an oriented k-plane L is α -positive with respect to Ω if $\Omega[L] > \alpha > 0$. If $\Omega[L] > \alpha > 1 - \epsilon$ then we call L an ϵ -calibrated plane.

The main result of this paper is then the following. Assume at all points and scales $B_r(x) \subseteq B_2$ that S is δ -Hausdorff close to a subspace $L_{x,r}$ which is uniformly positive $\Omega[L_{x,r}] > \alpha$ with respect to an ϵ -calibration. Then S is k-rectifiable with uniform volume bounds.

CONTENTS

1. Introduction	1
1.1. Examples and Applications	3
2. Proof of Theorem 1.4	6
2.1. The Family of Approximations S_r	6
2.2. Basic volume bounds on S and S_r	7
2.3. Rectifiability and Volume Control of S	9
References	10

1. INTRODUCTION

For a closed set $S \subset B_2(0) \subseteq \mathbb{R}^m$ and $B_r(x) \subseteq B_2(0)$, following [Tor95] let us define $\theta(x, r)$ by

$$\theta(k;x,r) = \theta(x,r) \equiv r^{-1} \inf_{L^k} d_H \left(S \cap B_r(x), L \cap B_r(x) \right), \tag{1.1}$$

Date: May 7, 2024.

Key words and phrases. Reifenberg theorem, Hausdorff measure, rectifiability, calibrations. N.E. was supported by NSF grant DMS-2204301,

where the inf is taken over all affine k-planes L and d_H is the Hausdorff distance. The classical Reifenberg theorem states that if $\theta(x,r) < \epsilon(n)$ for all $x \in S$ and $B_r(x) \subseteq B_2(0)$, then $S \cap B_1(0)$ is bi-Hölder equivalent to a flat disk $B_1(0) \subset \mathbb{R}^k$. We sometimes say in this case that S satisfies the ϵ -Reifenberg condition. This bi-Hölder equivalence is proved by constructing smooth approximations of S_r at all scales $r \in [0, 1]$, and carefully analyzing their behavior, see Section 2.1.

As is well understood, being ϵ -Reifenberg is not enough to guarantee k-dimensional volume bounds on S or its rectifiability. Many papers have investigated how to obtain these extra results by adding additional constraints on the set S. For instance in [Tor95] it is shown that if one has the L^2 Dini estimate $\int \theta(x, r)^2 \frac{dr}{r} < \epsilon$ for all $x \in S$, then S is bilipschitz to a Euclidean ball. In [NV17] it is shown that if one replaces the pointwise Dini estimate with an average $\int_S \int \theta(x, r)^2 \frac{dr}{r} < \epsilon$, then S is still $W^{1,p}$ equivalent to a Euclidean ball with $p \to \infty$ as $\epsilon \to 0$.

Other interesting extensions are obtained by looking at Jones' β -numbers, which are "one-sided versions" of the quantity θ introduced above, for example:

$$\beta_{\infty}(k;x,r) = \beta_{\infty}(x,r) \equiv r^{-1} \inf_{L^k} \sup_{y \in B_r(x) \cap S} d(y,L).$$

$$(1.2)$$

Notice in particular that any subset S of a k-dimensional plane satisfies $\beta_{\infty}(x,r) = 0$ for all x, r, while $\theta(x,r)$ need not be null if there are holes in S. These quantities are considered in [DT12, NV17, ENV] to prove k-dimensional volume bounds on S and rectifiability. Also, Azzam-Tolsa in [AT15] and Tolsa [Tol15] further investigated the summability properties of beta numbers, and showed that a set is k-rectifiable if and only if $\int \beta^2(x,r) \frac{dr}{r} < \infty$ for a.e. $x \in S$, and similar statements hold for measures. In [ENV], [Nab20] effective statements of this form were proved and then vastly generalized to classify all measures under analogous L^2 Dini estimate assumptions. These sort of estimates play a role in the analysis of singularities, see for instance [NV17, NV18, DLMSV18, EE19, NV20, Zha22].

In this note we take a different approach, and hope to drop the Dini estimate in certain contexts. We show that if a closed set $S \subset B_2(0) \subseteq \mathbb{R}^m$ satisfies an ϵ -Reifenberg condition with respect to k-planes which are uniformly positive with respect to an almost calibration, then $S \cap B_1(0)$ is k-Ahlfors regular and k-rectifiable without the additional Dini assumption. In order to make this precise let us introduce our terminology a little better:

Definition 1.1 (ϵ -Calibration). Let Ω be a smooth k-form over $B_2(0) \subseteq \mathbb{R}^m$. We say that Ω is an ϵ -calibration if

- (1) $|\Omega \Omega_0| \le \epsilon$ for a constant form Ω_0 ,
- (2) for all $x \in \mathbb{R}^m$ and any oriented k-dimensional subspace $L \subseteq \mathbb{R}^m$, we have $\Omega[L] \leq 1 + \epsilon$.

Remark 1.2. If L^k is an oriented subspace then we define $\Omega[L] = \Omega[e_1, \ldots, e_k]$ where e_1, \ldots, e_k is any oriented orthonormal basis of L.

Remark 1.3. Observe that if Ω is any C^0 regular k-form, then we can normalize it by a constant to satisfy (2), and thus after dilating a small ball $B_s(p) \to B_1(0)$ we have that Ω becomes an ϵ -calibration. In particular, the condition on Ω is highly non-restrictive and holds locally for all C^0 k-forms.

Our main result is that if our ϵ -Reifenberg surface is uniformly positive with respect to an almost calibration Ω , then it must be rectifiable with Ahlfor's regularity estimates:

Theorem 1.4 (Rectifiable Reifenberg for Almost Calibrations). Let $S \subset B_2(0) \subseteq \mathbb{R}^n$ be a closed set with $0 \in S$ and let Ω be an ϵ -calibration. Then for all $2\epsilon < \alpha < 1 \exists \delta(n, \epsilon) > 0$ and $A(n, \alpha, \epsilon) > 1$ such that if for all $B_r(x) \subseteq B_2(0)$ there exists an oriented k-dimensional subspace $L = L_{x,r}$ such that

$$d_H\left(S \cap B_r(x), L_{x,r} \cap B_r(x)\right) < \delta r \,, \quad \Omega[L_{x,r}] > \alpha > 0 \,, \tag{1.3}$$

then $S \cap B_1(0)$ is k-rectifiable and for all $x \in S$ with $B_{2r}(x) \subseteq B_2(0)$ we have that

$$(1 - C(n)\delta) \le \frac{\mathcal{H}^k(S \cap B_r(x))}{\omega_k r^k} \le A.$$
(1.4)

Further, $A \to 1$ *as* $\alpha \to 1$ *and* $\epsilon \to 0$.

Remark 1.5. We underline again that in this context there is no need for a summability condition to ensure rectifiability and volume bounds for S, unlike the general Rectifiable Reifenberg case. We simply require the Reifenberg condition, albeit restricted to positively oriented subspaces.

Remark 1.6. Though the theorem is stated in Euclidean space, it is easy to see that it can be applied to a manifold locally by looking at a suitably small chart. Indeed the statements are highly nonsensitive to perturbation.

Remark 1.7. In particular if $\Omega[L_{x,r}] > 1 - \epsilon$, so that S is almost calibrated with respect to Ω , then the volume of all balls $B_r(x)$ on S is close to that of the Euclidean ball.

Remark 1.8. If we view S equipped with the Hausdorff measure as an integral current, oriented by imposing that $\Omega[T_xS] \ge \alpha$, then $\partial S = 0$ as currents in B_2 and S satisfies the following "weak" almost-minimizing property: if S' is any other integral current with S' = S outside $U \subset B_2$ and $\partial S' \cap U = \emptyset$, then $||S||(U) \le C(\epsilon)||S'||(U)$.

The very rough outline is based on the simple observation that one can use the family of Reifenberg approximations S_r as a homotopy to an α -positive k-plane $L = S_0$. Up to some care on boundary terms, one can use this to uniformly control the integral of Ω on S, and hence its volume. We can then take a limit and make a similar conclusion on S itself.

1.1. **Examples and Applications.** Most of the immediate applications arise by considering either calibrations or their associated nonintegrable counterparts. The four examples we will consider are almost complex manifolds, potentially nonintegrable G^2 and Spin(7) manifolds, and almost special Lagrangians:

1.1.1. Almost Complex Manifolds. Let (M^{2n}, g, J) be a manifold with C^0 almost complex structure J and compatible metric g, i.e. g(JX, JY) = g(X, Y) and $J^2 = -Id$. We can write $\omega[X, Y] = g(JX, Y)$ to be the associated continuous 2-form.

In a neighborhood of each point $x \in M$ we can therefore take coordinates $\varphi : B_1(0^{2n}) \to M$ with $\varphi(0) = x$ and whose image is ϵ -small, i.e. after normalizing

$$\left|\varphi^* g_{ij} - \delta_{ij}\right|, \quad \left|\varphi^* J - J_0\right|, \quad \left|\varphi^* \omega - \omega_0\right| < \epsilon, \tag{1.5}$$

where the L^{∞} norm is being measured on $B_1(0^{2n}) \subseteq \mathbb{C}^n$ with J_0 the standard complex structure on \mathbb{C}^n and ω_0 the standard symplectic form on \mathbb{C}^n . In particular, it follows for each k that ω^k is an ϵ -calibration. If we apply Theorem 1.4 to this context we arrive at

Corollary 1.9 (C^0 Almost Complex Manifolds). Let $(B_2(0^{2n}), J)$ be an almost complex structure as in (1.5), so with $\omega = g(J, \cdot)$ we have that ω^k is an ϵ -calibration. Let $S = S^{2k} \subseteq B_2(0^{2n})$ be $\delta(x, \alpha)$ -Reifenberg with respect to α -positive subspaces. That is, assume for each $x \in S$ with $B_r(x) \subseteq B_2$ there exists a subspace $L_{x,r}^{2k} \subseteq \mathbb{R}^{2n}$ with

$$d_H(S \cap B_r(x), L_{x,r} \cap B_r(x)) < \delta, \quad \omega^k[L] > \alpha > 0.$$
(1.6)

Then S is 2k-rectifiable, and for each $x \in S$ with $B_{2r}(x) \subseteq B_2$ we have the Ahlfors regularity

$$(1 - C(n)\delta) \le \frac{\mathcal{H}^k(S \cap B_r(x))}{\omega_k r^k} \le A(n, \alpha).$$
(1.7)

Remark 1.10. If $\alpha \ge 1 - \epsilon$ then we might call S almost complex. In particular, if $L_{x,r}$ are almost-complex subspaces, the theorem holds. Notice also that $A \to 1$ as $\epsilon, \delta \to 0$.

Remark 1.11. If $J = J_0$ is the standard complex structure on \mathbb{C}^n and both $\delta \to 0$ and $\alpha \to 1$ as $r \to 0$ (i.e. *S* is "vanishing complex Reifenberg-flat"), then *S* is in fact a complex subvariety of \mathbb{C}^n . This follows by thinking of *S* as a complex current without boundary and applying the classification result of [Kin71].

1.1.2. *NonIntegrable G2 Structures*. Recall that one defines the associative three form ψ_0 and coassociative four form Ω_0 on \mathbb{R}^7 by

$$\psi_0 = e^{123} - e^{167} - e^{527} - e^{563} - e^{415} - e^{426} - e^{437},$$

$$\Omega_0 = e^{4567} - e^{4523} - e^{4163} - e^{4127} - e^{2637} - e^{1537} - e^{1526}.$$
(1.8)

The definitions are motivated by looking at the imaginary part of the octonian algebra.

Let (M^7, Ω) be a 7-manifold with C^0 regular 3-form ψ . We call ψ a G2-structure if at each $x \in M$ there is a basis $e_1, \ldots, e_7 \in T_x M$ such that

$$\psi(x) = \psi_0 \,. \tag{1.9}$$

Recall [Bry87] that such a basis uniquely defines a metric g on M, and hence by Hodge starring we get a global continuous 4-form Ω such that with respect to the same basis we have $\Omega(x) = \Omega_0$.

In this context we can then find in a neighborhood of each $x \in M$ coordinates $\varphi : B_2(0^7) \subseteq \mathbb{R}^7 \to M$ with $\varphi(0) = x$ and

$$\left|\varphi^*\Omega - \Omega_0\right| < \epsilon \,. \tag{1.10}$$

Our main result is to study Reifenberg sets which are positive with respect to the coassociative form:

Corollary 1.12 (C^0 Almost G2 Structures). Let $(B_2(0^7), \Omega)$ be a G2 structure as in (1.10). Let $S^4 \subseteq B_2(0^7)$ be $\delta(x, \alpha)$ -Reifenberg with respect to α -positive subspaces. That is, assume for each $x \in S$ with $B_r(x) \subseteq B_2$ there exists a subspace $L^4_{x,r} \subseteq \mathbb{R}^7$ with

$$d_H(S \cap B_r(x), L_{x,r} \cap B_r(x)) < \delta, \quad \Omega[L] > \alpha > 0.$$
(1.11)

Then S is 4-rectifiable, and for each $x \in S$ with $B_{2r}(x) \subseteq B_2$ we have the Ahlfors regularity

$$(1 - C(n)\delta) \le \frac{\mathcal{H}^k(S \cap B_r(x))}{\omega_k r^k} \le A(n, \alpha).$$
(1.12)

Remark 1.13. If $\alpha \ge 1 - \epsilon$ then we might call S almost coassociative, and in this case $A \to 1$ as $\epsilon, \delta \to 0$.

1.1.3. Almost Spin(7) Manifolds. Recall that one defines the canonical four form Ω_0 on \mathbb{R}^8 by

$$\Omega_0 = e^{1256} + e^{1278} + e^{3456} + e^{3478} + e^{1357} - e^{1368} - e^{2457}, + e^{2468} - e^{1458} - e^{1467} - e^{2358} - e^{2367} + e^{1234} + e^{5678}.$$
(1.13)

The isomorphism group of this four form is exactly Spin(7). We call a manifold (M^8, Ω) with a C^0 four form Ω an almost Spin(7) manifold if for each $x \in M$ there exists a basis $e_1, \ldots, e_8 \in T_x M$ such that

$$\Omega(x) = \Omega_0 \,. \tag{1.14}$$

In this context we can then find in a neighborhood of each $x \in M$ coordinates $\varphi : B_2(0^8) \subseteq \mathbb{R}^8 \to M$ with $\varphi(0) = x$ and

$$\left|\varphi^*\Omega - \Omega_0\right| < \epsilon \,. \tag{1.15}$$

Our main result is to study Reifenberg sets which are positive with respect to the Spin(7) form:

Corollary 1.14 (C^0 Almost Spin(7) Structures). Let $(B_2(0^8), \Omega)$ be a G2 structure as in (1.15). Let $S^4 \subseteq B_2(0^8)$ be $\delta(x, \alpha)$ -Reifenberg with respect to α -positive subspaces. That is, assume for each $x \in S$ with $B_r(x) \subseteq B_2$ there exists a subspace $L^4_{x,r} \subseteq \mathbb{R}^8$ with

$$d_H(S \cap B_r(x), L_{x,r} \cap B_r(x)) < \delta, \quad \Omega[L] > \alpha > 0.$$
(1.16)

Then S is 4-rectifiable, and for each $x \in S$ with $B_{2r}(x) \subseteq B_2$ we have the Ahlfors regularity

$$(1 - C(n)\delta) \le \frac{\mathcal{H}^k(S \cap B_r(x))}{\omega_k r^k} \le A(n, \alpha).$$
(1.17)

1.1.4. Almost Special Lagrangians. Let (M^{2n}, Ω) be a 2*n*-dimensional manifold with C^0 regular *n*-form Ω . Assume at each $x \in M$ we can find a basis $\frac{\partial}{\partial x^1}, \frac{\partial}{\partial y^1}, \dots, \frac{\partial}{\partial x^n}, \frac{\partial}{\partial y^n} \in T_x M$ such that

$$\Omega(x) = Re\left(dz^1 \wedge \dots \wedge dz^n\right),\tag{1.18}$$

where $dz^j = dx^j + idy^j$ with respect to the dual basis on T_x^*M . Then we call Ω an almost Special Lagrangian structure. Note the above is quite weak, and even on a Kähler manifold the result would be much weaker than a calibration for Special Lagrangians. Indeed it does not require a reduction to SU(n) of the structure group, as the required basis may rotate the complex factor in front of $Re(dz^1 \wedge \cdots \wedge dz^n)$ as x moves.

In this context we can then find in a neighborhood of each $x \in M$ coordinates $\varphi : B_2(0^{2n}) \subseteq \mathbb{C}^n \to M$ with $\varphi(0) = x$ and

$$\left|\varphi^*\Omega - Re\left(dz^1 \wedge \dots \wedge dz^n\right)\right| < \epsilon.$$
(1.19)

We can state the main result, which is that sets S^n which are Reifenberg flat and uniformly positive with respect to an almost calibration for Special Lagrangians are rectifiable:

Corollary 1.15 (C^0 Almost Special Lagrangians). Let $(B_2(0^{2n}), \Omega)$ be an almost calibration for Special Lagrangians as in (1.19). Let $S^n \subseteq B_2(0^{2n})$ be $\delta(x, \alpha)$ -Reifenberg with respect to α -positive subspaces. That is, assume for each $x \in S$ with $B_r(x) \subseteq B_2$ there exists a subspace $L^n_{x,r} \subseteq \mathbb{R}^{2n}$ with

$$d_H(S \cap B_r(x), L_{x,r} \cap B_r(x)) < \delta, \quad \Omega[L] > \alpha > 0.$$
(1.20)

Then S is n-rectifiable, and for each $x \in S$ with $B_{2r}(x) \subseteq B_2$ we have the Ahlfors regularity

$$(1 - C(n)\delta) \le \frac{\mathcal{H}^k(S \cap B_r(x))}{\omega_k r^k} \le A(n, \alpha).$$
(1.21)

Remark 1.16. If $\alpha \ge 1 - \epsilon$ then we might call S an almost Special Lagrangian, and in this case $A \to 1$ as $\epsilon, \delta \to 0$.

2. PROOF OF THEOREM 1.4

2.1. The Family of Approximations S_r . Let $S \subseteq B_2 \subseteq \mathbb{R}^n$ be an δ -Reifenberg set, namely for each $x \in S$ and $B_r(x) \subseteq B_2$ there exists an a k-plane $L_{x,r} \subseteq \mathbb{R}^n$ such that $d_H(S \cap B_r(x), L_{x,r} \cap B_r(x)) < \delta r$. The classical Reifenberg tells us that near B_1 we have that S is homeomorphic to a disk. A key ingredient in this proof is the construction of smooth approximations S_r of S, and their careful analysis. The below is a mild extension of Reifenberg's original construction, which itself follows immediately from his construction:

Lemma 2.1 (Construction of Smooth Approximations S_r). Let $S \subseteq B_{1+\epsilon}(0) \subseteq \mathbb{R}^n$ be a δ -Reifenberg set as above with $\epsilon > 0$ fixed and $\delta < \delta(n, \epsilon)$. For each $r \in (0, \epsilon]$ there exists smooth oriented manifolds $S_r \subseteq \mathbb{R}^n$ which depend smoothly on r such that

(1) $S_{\epsilon} = L_{0,\epsilon}$ is the fixed plane which well approximates S on $B_{1+\epsilon}(0)$.

- (2) On $\mathbb{R}^n \setminus B_{1+\epsilon}$ we have that $S_r = S_{\epsilon} = L_{0,\epsilon}$ is a fixed plane.
- (3) $d_H(S \cap B_1, S_r \cap B_1) < C(n, \epsilon)\delta r$ with second fundamental form bound $r|A_{S_r}| \le C(n, \epsilon)\delta$.
- (4) For each $x \in S_r \cap B_1$ and $y \in S \cap B_r(x)$ we have $d_{Gr}(L_{y,r_y}, T_xS_r) < C(n,\epsilon)\delta$, where $r_y = r_{|y|}$ is piecewise linear with $r_{|y|} = r$ for $|y| \le 1$ and $r_{|y|} = \epsilon$ for $|y| \ge 1 + \epsilon$. In particular we can orient S_r so that $\Omega_0[T_xS_r] > \epsilon/2$ for all $x \in S_r \cap B_1$.
- (5) $\left| \frac{d}{dr} S_r \right| < C(n, \epsilon) \delta$.
- (6) If $S \cap B_{2r}(x)$ is smooth with second fundamental form $r|A_S| \leq \epsilon$, then $S \cap B_r(x) = S_r \cap B_r(x)$.

Remark 2.2. d_{Gr} in (4) is the Grassmann distance between the subspace L_{y,r_y} and the tangent space T_xS_r .

Remark 2.3. $\frac{d}{dr}S_r$ is understood as the normal velocity vector to S_r , so that (5) is the statement that the this normal velocity vector has small norm.

Remark 2.4. Condition (4) is just saying that T_xS_r is always well approximated by a subspace L, which itself well approximates S on the appropriate scale. On B_1 this scale is r and outside of $B_{1+\epsilon}$ this scale is ϵ , with a linear interpolation between.

Rough Proof. As the details are in many references, see for instance [Tor95, Sim, Nab20], and in particular [Nab20, Theorem 4.2] for the construction of a one parameter family of approximating submanifolds S_r , let us just mention a few words on the proof. The idea is simple if somewhat tedious. One picks a Vitali covering $\{B_{r_i}(y_i)\}$ of $S \cap B_{1+\epsilon}$ with $r_i = r_{|y_i|}$ as in (4). Vitali here means that $\{B_{r_i/5}(y_i)\}$ are disjoint. Then choose for each such ball the best approximating plane $L_i = L_{y_i,r_i}$, and then glue them together using estimates provided by the Reifenberg assumption. Condition (5) is not usually stated in this form, instead one often points out that S_{2r} is graphical over S_r with likewise bounds. However, the graphical property immediately implies we can construct as in (5). Indeed, one can build $S_a = S_{r_a}$ for $r_a = 2^{-a}$ and use the graphical property to interpolate and build S_r for $r \in (2^{-a}, 2^{-a+1})$.

Let us state in words the outcome of Lemma 2.1. Beginning with our Reifenberg set S we are building a family of approximations S_r , which by (1) begin at a fixed plane $S_{\epsilon} = L_{0,\epsilon}$ and end at $S_0 = S$, at least on B_1 . This family S_r is independent of r outside of $B_{1+\epsilon}$, and hence equal to the same plane $L_{0,\epsilon}$. By (3) each S_r is δr -close to S on B_1 . The family S_r is moving in a continuous fashion by (5), and the tangent space $T_x S_r$ is always close to some approximating plane of S by (4). Condition (6) tells us that if S is itself scale invariantly smooth on some scale, then the approximation should equation S itself.

2.2. Basic volume bounds on S and S_r . Let us begin with the following, which is under the context of Theorem 1.4 and Lemma 2.1:

Lemma 2.5. Let $S \subseteq B_{1+\epsilon}$ be as in Theorem 1.4. Then for $A = A(n, \alpha, \epsilon)$, $\epsilon < \epsilon(n, \alpha)$ and $\delta < \delta(n, \alpha, \epsilon)$ we have the volume estimate

$$\mathcal{H}^{k}(S_{r} \cap B_{1}) \leq \mathcal{H}^{k}(S_{r} \cap B_{1+\epsilon}) \leq A\omega_{k},$$

$$\mathcal{H}^{k}(S_{r} \cap B_{1}) \geq (1 - C(n)\delta)\omega_{k}$$
(2.1)

independent of r > 0 with $A(n, \alpha, \epsilon) \to 1$ as $\alpha \to 1$ and $\epsilon \to 0$. Moreover, as currents we have

$$(\pi_0)_{\sharp}[S_r] = [\pi_0(S_r)], \quad and \quad \pi_0(S_r) \supset B_{1-C(n)\delta} \cap L_{0,\epsilon},$$

where $\pi_0 : \mathbb{R}^n \to L_{0,\epsilon}$ is the orthogonal projection map.

Proof. Recall that our almost calibration Ω satisfies the L^{∞} control

$$|\Omega[L]| \le 1 + \epsilon, |\Omega - \Omega_0| < \epsilon,$$
(2.2)

where Ω_0 is a constant k-form. It follows from Lemma 2.1.4 and our uniform positivity that for each $x \in S_r$ we have

$$\Omega_0[T_x S_r] \ge \Omega[T_x S_r] - \epsilon \ge \Omega[L_{y,r_y}] - C(n,\epsilon)\delta - \epsilon \ge \alpha - 3\epsilon/2,$$
(2.3)

where by Lemma 2.1.3 $y \in S \cap B_r(x)$ is any choice of point.

On the other hand, $d\Omega_0 = 0$ is a closed k-form trivially as it is constant, and so using Lemma 2.1.1 and Lemma 2.1.5 we have that

$$\int_{S_r \cap B_{1+\epsilon}} \Omega_0 = \int_{L_{0,\epsilon} \cap B_{1+\epsilon}} \Omega_0 \le 1 + C(n)\epsilon.$$
(2.4)

If we then combine this with (2.3) we can turn this into the upper bound

$$\mathcal{H}^{k}(S_{r} \cap B_{1}) = \int_{S_{r} \cap B_{1}} d\mathcal{H}_{S}^{k} \leq \int_{S_{r} \cap B_{1+\epsilon}} d\mathcal{H}_{S}^{k},$$

$$\leq (\alpha - 3\epsilon/2)^{-1} \omega_{k} \int_{S_{r} \cap B_{1+\epsilon}} \Omega_{0} \leq \frac{1 + C(n)\epsilon}{\alpha - 3\epsilon/2} \omega_{k}.$$
 (2.5)

In order to prove the lower bound, we proceed in a manner similar to the general Reifenberg case. Namely consider the projection map $\pi_0 : \mathbb{R}^n \to L_{0,\epsilon}$, which is clearly a Lipschitz map with $|d\pi_0[v]| \le |v|$. Note that

$$\pi_0[S_r \cap \partial B_{1+\epsilon}(0^n)] = L_{0,\epsilon} \cap \partial B_{1+\epsilon}(0^k), \qquad (2.6)$$

independent of r > 0, and that π_0 on $S_{\epsilon} \cap B_{1+\epsilon}(0^n)$ is effectively just the identity map. In particular, for all r > 0 we have that π_0 on $S_r \cap B_{1+\epsilon}(0^n)$ has topological degree one. Hence π_0 must map $S_r \cap B_{1+\epsilon}(0^n)$ onto $B_{1+\epsilon}(0^k)$. On the other hand, it follows from the Hausdorff condition that we then must have

$$\pi_0[S_r \cap B_1(0^n)] \supseteq B_{1-C(n)\delta}(0^k).$$
(2.7)

As π_0 is a submetry, we then get the lower volume estimate

$$\mathcal{H}^k(S_r \cap B_1) \ge \mathcal{H}^k(\pi_0[S_r \cap B_1]) \ge \mathcal{H}^k(B_{1-C(n)\delta}(0^k)) \ge (1-C(n)\delta)\omega_k,$$
(2.8)

as claimed. The assertion $(\pi_0)_{\sharp}[S_r] = [\pi_0(S_r)]$ follows from the constancy theorem and the fact that π_0 has degree one on $S_r \cap B_{1+\epsilon}$.

Let us now use the above to conclude Ahlfor's regularity of our approximating submanifolds:

Lemma 2.6. Let $S \subseteq B_2$ be as in Theorem 1.4, and let S_r be a smooth approximation with $S_r \to S$ on B_1 as in Lemma 2.1. Let $A = A(n, \alpha, \epsilon)$, $\epsilon < \epsilon(n, \alpha)$ and $\delta < \delta(n, \alpha, \epsilon)$, then for all $x \in S_r$ with $B_{2s}(x) \subseteq B_2$ we have the volume estimates

$$(1 - C(n)\delta)\omega_k s^k \le \mathcal{H}^k(S_r \cap B_s(x)) \le A\omega_k s^k.$$
(2.9)

Here $A(n, \alpha, \epsilon) \to 1$ *as* $\alpha \to 1$ *and* $\epsilon \to 0$. *Moreover, when* r < s *then given any* $y \in S \cap B_r(x)$ *we have the identity*

$$(\pi_{y,s})_{\sharp}[S_r \cap B_s(x)] = [\pi_{y,s}(S_r \cap B_s(x))] \quad and \quad \pi_{y,s}(S_r \cap B_s(x)) \supset L_{y,s} \cap B_{(1-C(n)\delta)s}(x).$$
(2.10)

where $\pi_{y,s}$ is the projection onto $L_{y,s}$.

Proof. Take $x \in S_r$ with $B_{2s}(x) \subseteq B_2$. Note that if $s \leq r$ then the result follows due to the smoothness conditions on S_r , indeed one can take $A \approx 1 + C(n)\delta$ using the second fundamental form bounds by Lemma 2.1.3. So we can assume s > r without loss. If we rescale and translate our ball $B_{2s}(x) \to B_2(0)$ with $\tilde{S}_r \subseteq B_2(0)$ the rescaled set, observe that \tilde{S}_r is now itself a Reifenberg set which satisfies the context of Lemma 2.1 and Lemma 2.5. Further the approximations $\tilde{S}_{r,r'}$ of \tilde{S}_r from Lemma 2.1 satisfy $\tilde{S}_{r,r} = \tilde{S}_r$ on B_1 , as in Lemma 2.1.6. Hence we can apply Lemma 2.5 to $\tilde{S}_{r,r'}$ to get the claimed estimates

$$\mathcal{H}^{k}(S_{r} \cap B_{s}(x)) = s^{k} \mathcal{H}^{k}(\tilde{S}_{r} \cap B_{1}(0)) \leq A\omega_{k} s^{k},$$

$$\mathcal{H}^{k}(S_{r} \cap B_{s}(x)) = s^{k} \mathcal{H}^{k}(\tilde{S}_{r} \cap B_{1}(0)) \geq (1 - C(n)\delta)\omega_{k} s^{k}.$$
 (2.11)

The last statement follows in a similar fashion from Lemmas 2.1 and 2.5.

2.3. Rectifiability and Volume Control of S. Consider smooth approximations S_r of S as in Lemma 2.1 such that

$$S_r = L_{0,\epsilon} \text{ outside of } B_{1+\epsilon},$$

$$d_H(S_r \cap B_1, S \cap B_1) < C(n)\delta r.$$
(2.12)

The S_r are oriented k-dimensional surfaces in $B_{1+\epsilon}$ with $\partial S_r = 0$ and uniform bounds $\mathcal{H}^k(S_r \cap B_{1+\epsilon}) \leq A\omega_k$. Given any $r_i \to 0$, we can therefore apply the Feder-Fleming compactness theorem for currents [Sim83] to pass to a subsequence and find an integral k-current T in $B_{1+\epsilon}$ so that $[S_{r_i}] \to T$. The limiting current T has zero boundary and is positively oriented w.r.t. Ω_0 (since the S_{r_i} are).

We claim that $S \cap B_1$ is k-rectifiable, and T = [S] where we orient the tangent spaces of S positively w.r.t. Ω_0 , i.e. $\Omega_0[T_xS] \ge 0$ for \mathcal{H}^k -a.e. x. (Our claim will also imply $[S_r] \to [S]$ in B_1 as $r \to 0$.)

Trivially $\operatorname{spt}(T) \cap B_1 \subset S$, and so in B_1 we can write the mass measure $||T|| = \mathcal{H}^k \llcorner \theta \llcorner S'$ for some $S' \subset S$ and some \mathbb{N} -valued function θ . On the other hand, for any $x \in S \cap B_1$ and $B_{2s}(x) \subset B_2$ we can take the limit of (2.10) to get

$$(\pi_{x,s})_{\sharp}(T \sqcup B_s(x)) = [U_{x,s}] \quad \text{for some } L_{x,s} \cap B_{(1-C(n)\delta)s}(x) \subset U_{x,s} \subset L_{x,s}.$$
(2.13)

Since $\operatorname{Lip}(\pi_{x,s}) \leq 1$, we get

$$||T||(B_s(x)) \ge (1 - C(n)\delta)s^k \omega_k \quad \forall s < 1 - |x|/2,$$
(2.14)

and therefore we deduce S' = S.

So S is k-rectifiable, and for \mathcal{H}^k -a.e. $x \in S \cap B_1 \equiv \operatorname{spt} T \cap B_1$, T has the approximate tangent planewith-multiplicity:

$$(\eta_{x,r})_{\sharp}T \to \theta(x)[T_xS] \quad \text{as } r \to 0,$$

Here $\eta_{x,r}(y) = (y - x)/r$, and $T_x S$ is oriented so that $\Omega_0[T_x S] \ge 0$. In fact since $\Omega_0[T_y S_r] \ge \epsilon/2$ for all $y \in S_r \cap B_1$ we have $\Omega_0[T_x S] \ge \epsilon/2$ also. The identity (2.13) passes to the limit, so we get

$$\theta(x)(\pi_{x,s})_{\sharp}[T_xS] = [L_{x,s}] \quad \forall s < 1 - |x|/2,$$

which implies $\theta(x) = 1$. This proves our claim.

From T = [S], (2.14) and lower-semi-continuity of mass we get the volume estimates

$$(1 - C(n)\delta)\omega_k \le \mathcal{H}^k(S \cap B_1) \le A\omega_k.$$

We can of course now apply the same argument to any ball $B_{2r}(x) \subseteq S$ in order to obtain that S is everywhere rectifiable with the estimates

$$(1 - C(n)\delta)\omega_k r^k \le \mathcal{H}^k(S \cap B_r(x)) \le A\omega_k r^k, \qquad (2.15)$$

finishing the proof of Theorem 1.4.

REFERENCES

- [AT15] J. AZZAM and X. TOLSA, Characterization of n-rectifiability in terms of Jones' square function: Part II, Geom. Funct. Anal. 25 (2015), 1371–1412. MR 3426057. Zbl 06521333. http://dx.doi.org/10.1007/s00039-015-0334-7.
- [Bry87] R. L. BRYANT, Metrics with exceptional holonomy, Ann. of Math. (2) **126** (1987), 525–576. MR 916718. http://dx.doi.org/10.2307/1971360. Available at https://doi.org/10.2307/1971360.
- [DT12] G. DAVID and T. TORO, Reifenberg parameterizations for sets with holes, *Mem. Amer. Math. Soc.* **215** (2012), vi+102. MR 2907827. Zbl 1236.28002. http://dx.doi.org/10.1090/S0065-9266-2011-00629-5.
- [DLMSV18] C. DE LELLIS, A. MARCHESE, E. SPADARO, and D. VALTORTA, Rectifiability and upper Minkowski bounds for singularities of harmonic Q-valued maps, Comment. Math. Helv. 93 (2018), 737–779. MR 3880226. http://dx.doi.org/10.4171/CMH/449. Available at https://doi.org/10.4171/CMH/449.
- [ENV] N. S. EDELEN, A. NABER, and D. VALTORTA, Quantitative Reifenberg theorem for measures, arXiv:1612.08052. Available at https://arxiv.org/abs/1612.08052.
- [EE19] N. EDELEN and M. ENGELSTEIN, Quantitative stratification for some free-boundary problems, *Trans. Amer. Math. Soc.* 371 (2019), 2043–2072. MR 3894044. http://dx.doi.org/10.1090/tran/7401. Available at https://doi.org/10.1090/tran/7401.
- [Kin71] R. KING, The currents defined by analytic varieties, J. Acta Math. 127 (1971).185-220. http://dx.doi.org/10.1007/BF02392053. MR 393550. Available at https://doi.org/10.1007/BF02392053.
- [Nab20] A. NABER, Lecture notes on rectifiable Reifenberg for measures, in *Harmonic analysis and applications*, *IAS/Park City Math. Ser.* 27, Amer. Math. Soc., [Providence], RI, [2020] ©2020, pp. 289–345. MR 4249631.
- [NV17] D. VALTORTA, Rectifiable-Reifenberg and A. NABER and the regularity of stationary harmonic maps, Ann. of Math. (2) 185 (2017), and minimizing 131-227. MR 3583353. Zbl 06686585. http://dx.doi.org/10.4007/annals.2017.185.1.3. Available at https://arxiv.org/abs/1504.02043.
- [NV18] _____, Stratification for the singular set of approximate harmonic maps, Math. Z. 290 (2018), 1415–1455. MR 3856859. http://dx.doi.org/10.1007/s00209-018-2068-3. Available at https://doi.org/10.1007/s00209-018-2068-3.
- [NV20] _____, The singular structure and regularity of stationary varifolds, J. Eur. Math. Soc. (JEMS) 22 (2020), 3305-3382. MR 4153109. http://dx.doi.org/10.4171/jems/987. Available at https://doi.org/10.4171/jems/987.

- [Rei60] E. R. REIFENBERG, Solution of the Plateau Problem for *m*-dimensional surfaces of varying topological type, Acta Math. 104 (1960), 1-92. MR 0114145. Zbl 0099.08503. Available at http://link.springer.com/article/10.1007%2FBF02547186.
- [Sim] L. SIMON, Reifenberg's topological disc theorem. Available at http://www.math.uni-tuebingen.de/ab/analysis/pub/
- [Sim83] _____, Lectures on geometric measure theory, Proceedings of the Centre for Mathematical Analysis, Australian National University 3, Australian National University, Centre for Mathematical Analysis, Canberra, 1983. MR 756417. Zbl 0546.49019.
- [Tol15] X. TOLSA, Characterization of n-rectifiability in terms of Jones' square function: part I, Calc. Var. Partial Differential Equations 54 (2015), 3643–3665. MR 3426090. Zbl 06544048. http://dx.doi.org/10.1007/s00526-015-0917-z.
- [Tor95] T. TORO, Geometric conditions and existence of bi-Lipschitz parameterizations, *Duke Math. J.* **77** (1995), 193–227. MR 1317632. Zbl 0847.42011. http://dx.doi.org/10.1215/S0012-7094-95-07708-4.
- [Zha22] B. ZHANG, Rectifiability and Minkowski bounds for the zero loci of $\mathbb{Z}/2$ harmonic spinors in dimension 4, *Comm. Anal. Geom.* **30** (2022), 1633–1681. MR 4596625.

UNIVERSITY OF NOTRE DAME (USA)

Email address: nedelen@nd.edu

NORTHWESTERN UNIVERSITY (USA) Email address: anaber@math.northwestern.edu

UNIVERSITY OF MILANO-BICOCCA (EU) *Email address*: daniele.valtorta@unimib.it