PROOF OF THE COMPLETE PRESENCE OF A MODULO 4 BIAS FOR THE SEMIPRIMES

Nikola Gyulev, Miroslav Marinov^{*}

Abstract


Dummit, Granville and Kisilevsky have recently shown that the proportion of semiprimes (products of two primes) not exceeding a given x, whose factors are congruent to 3 modulo 4, is more than a quarter when x is sufficiently large. They have also conjectured that this holds from the very beginning, that is, for all $x \ge 9$. We give a proof for $x \ge 10^{21}$ via an explicit approach based on their work. Together with their data for the remaining x, this results in a full proof of the conjecture. Our method consists of techniques with cancellations of sums over primes with different remainders. We also rely on classical estimates for prime counting functions, as well as on very recent explicit improvements by Bennet, Martin, O'Bryant and Rechnitzer, which have wide applications in essentially any setting involving estimations of sums over primes.

1 Introduction

There are many natural questions regarding the structure of the set of prime numbers modulo a given integer. Throughout we denote by $\pi(x)$ the number of primes not exceeding x and by $\pi(x, q, a)$ the number of primes not exceeding x which are congruent to a modulo q. Focusing on modulo 4, with the possible remainders being 1 and 3, we could ask: are the sizes $\pi(x, 4, 1)$ and $\pi(x, 4, 3)$ close for all x; if not, is there dominance from one or the other or does their difference oscillate in a symmetric fashion? A starting point is the Prime Number Theorem for Arithmetic Progressions [3], which gives

$$\pi(x,4,3) \sim \pi(x,4,1) \sim \frac{x}{2\log x}$$

(Throughout we write $f(x) \sim g(x)$ if $g(x) \neq 0$ for sufficiently large x and $\lim_{x\to\infty} \frac{f(x)}{g(x)} = 1$.) Hence it makes sense to conjecture that eventually the two counts will be close throughout or that firstly remainder 3 would dominate, then 1, then 3, etc., roughly symmetrically. Let us now display $\pi(p_n, 4, 3) - \pi(p_n, 4, 1)$, where p_n denotes the *n*-th prime. There is a noticeable imbalance, as remainder 3 seems to be dominating.



This is the so-called *Chebyshev's Bias*, which he first observed in 1853, describing it in a letter to Fuss

^{*}gyulev2718@gmail.com, m.marinov1617@gmail.com

[1]. How might one prove this? It is natural to consider the sum $\sum_{2 , as it precisely shows which residue class is more frequent at each point and by exactly how much. However, the discontinuities of this sum make it impossible to apply any analytic techniques. A possible salvage of this problem is to consider a closely related sum with smooth components. In this manner, Chebyshev stated the following:$

Conjecture (Chebyshev 1856, [1]). As $y \to 0^+$ we have

$$\sum_{p>2} (-1)^{\frac{p-1}{2}} e^{-py} \to -\infty.$$

In 1916 Hardy and Littlewood [5] gave a proof by assuming the Generalized Riemann hypothesis (GRH) for the Dirichlet series $\sum_{m=0}^{\infty} (-1)^m (2m+1)^{-s}$ and Landau [9] proved in 1918 that the two are equivalent. Going back to the plot of the difference, even though the function is mainly positive, there are spots where for a short period remainder 1 catches up, the first two being far away from each other, at $x = p_{2946}$ and $x = p_{50378}$. Surprisingly, it turns out there are infinitely many such exceptional x.

Theorem 1.1 (Littlewood 1914, [10]). There are infinitely many integers x for which

$$\pi(x,4,1) - \pi(x,4,3) \ge \frac{1}{2} \frac{\sqrt{x} \log \log \log x}{\log x}$$

Thus, we cannot blindly substitute y = 0 in Conjecture 1, as it would give $\sum_{2 as <math>x \to \infty$. Hence we may view Conjecture 1 as a test rather than an exact measurement.

In order to measure how often remainder 3 dominates over remainder 1, one explores

$$\frac{\#\{w \le x: \ \pi(w, 4, 3) > \pi(w, 4, 1)\}}{x}$$

and in particular, Knapowski and Turan [8] conjectured that this ratio tends to 1 as $x \to \infty$. However, it turns out to not converge at all, as shown in 1993 by Kaczorowski [7]. A solution was found in 1994 when Rubinstein and Sarnak [16] had an insightful observation – rather than counting 1 for every w for which there are more primes 3 (mod 4), we count $\frac{1}{w}$. This led to the following remarkable result.

Theorem 1.2 (Rubinstein-Sarnak, 1994, [16]). Assume GRH. As $x \to \infty$ we have

$$\frac{1}{\log x} \sum_{w \le x: \ \pi(x,4,3) > \pi(x,4,1)} \frac{1}{w} \to 0,9959...$$

Therefore, with the correct measurement, Chebyshev is right approximately 99.59% of the time. What happens for a general modulo n? Rubinstein and Sarnak [16] show more generally, assuming conjectures related to GRH, that $\pi(x, n, a) > \pi(x, n, b)$ more often, for any quadratic non-residue a and quadratic residue b with gcd(a, n) = gcd(b, n) = 1. Chebyshev's bias naturally leads to similar questions where the main object is slightly different. For example, we can consider the products pq of two primes (also called *semiprimes*) less than a given x. When focusing on modulo 4, the possibilities are $p \equiv q \equiv 3 \pmod{4}$, $p \equiv q \equiv 1 \pmod{4}$ and $p \equiv 3 \pmod{4}$ with $q \equiv 1 \pmod{4}$ or vice-versa. This has recently been explored by Dummit, Granville and Kisilevsky [2] in 2016, whose data suggests strong dominance from remainder 3. The behaviour for general modulo again turns out to depend on comparing quadratic versus non-quadratic residues.

Theorem 1.3 (Dummit-Granville-Kisilevsky, 2016). Let χ be a real non-principal Dirichlet character with conductor d. For $\eta \in \{-1, 1\}$ we have

$$\frac{\#\{m \le x : m = pq \text{ with } \chi(p) = \chi(q) = \eta\}}{\frac{1}{4}\#\{m \le x : m = pq, (m, d) = 1\}} = 1 + \eta \frac{\sum_{p} \frac{\chi(p)}{p} + o(1)}{\log \log x}$$

where p and q represent primes.

(Throughout, for functions f and g such that $g(x) \neq 0$ for sufficiently large x, we write f(x) = o(g(x)) if $\lim_{x\to\infty} f(x)/g(x) = 0$; f(x) = O(g(x)) if $|f(x)| \leq Cg(x)$ for some constant C; $f(x) = \Theta(g(x))$ if f = O(g) and g = O(f).) It follows that the sign of $\sum_{p} \frac{\chi(p)}{p}$ determines the outcome of the bias. For modulo 4 with the character $\chi(p) = (-1)^{\frac{p-1}{2}}$ we have $\sum_{p>2} \frac{(-1)^{\frac{p-1}{2}}}{p} \approx -0.0334$, so for large x the semiprimes $pq \leq x$ with

- $p \equiv q \equiv 3 \pmod{4}$ are approximately $\frac{1}{4} + \frac{0.0835}{\log \log x}$ of all.
- $p \equiv q \equiv 1 \pmod{4}$ are approximately $\frac{1}{4} \frac{0.0835}{\log \log x}$ of all.
- $p \equiv 1 \pmod{4}$, $q \equiv 3 \pmod{4}$ are approximately $\frac{1}{2}$ of all.

An important note is that in this bias from some point on there is full dominance, differing substantially from Chebyshev's where a change of the lead occurs infinitely often. Hence we expect that Chebyshev's bias cannot be used to prove results for products of two primes. Note also that Theorem 1.3 is not enough on its own to justify that the bias for a general modulo d is present, as we also need the following.

Conjecture. For any Dirichlet character χ we have $\sum_{p} \frac{\chi(p)}{p} \neq 0$.

This sum is closely related to the Dirichlet L-function sum $L(1,\chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n}$, for which not much is known besides the fact that it is non-zero. (This is an important step of Dirichlet's proof of the theorem for primes in arithmetic progressions.) A further extension of Theorem 1.3 has been done by Hough [6], who shows that the right-hand side is at least $1 + \frac{\log \log \log x + O(1)}{\log \log x}$ for at least one d with $d \leq x$. The generalization to products of more than two primes has been recently explored by Meng [13]. Finally, Dummit, Granville and Kisilevsky conjecture that the bias mod 4 is present from the very beginning. Our main result is a proof of this conjecture.

Theorem 1.4. For all $x \ge 9$ we have

$$\frac{\#\{m \le x : m = pq \text{ with } p \equiv q \equiv 3 \pmod{4}\}}{\#\{m \le x : m = pq \text{ with } p, q \text{ odd}\}} > \frac{1}{4}.$$

To prove this, one could adapt Dummit, Granville and Kisilevsky's main ideas to statements with explicit bounds. This is precisely what Marinov [12] did in 2020, when he obtained this result for $x \ge 10^{138}$. Our approach is based on his work and it yields a proof for $x \ge 10^{21}$, as it has a different main function, more careful uses of Mertens' estimates, as well as a more precise comparison between $O\left(\frac{\log \log x}{(\log x)^2}\right)$ terms and $O\left(\frac{1}{(\log x)^2}\right)$ terms. This is enough for a full proof, having in mind the experimental data up to 10^{24} of Dummit, Granville and Kisilevsky [2].

2 Strategy and auxiliary results on counting primes

Our approach can be divided into three key steps.

• First, we express the difference between the two counts (with the number of all products of two primes scaled by a factor of $\frac{1}{4}$) in terms of double summations over the primes. Using estimates for prime counting functions, we reduce to a combination of six summations (with a single counter) over subsets of the primes. This is done in a specific way to get rid of double sums and utilize the precise estimate for $\sum_{p\geq 3} \frac{(-1)^{\frac{p-1}{2}}}{p}$, which we arrange to be the coefficient of the dominant term, of order $\Theta\left(\frac{x}{\log x}\right)$.

- We estimate each of the six sums with an elementary function (combination of x and $\log x$). This is done via classical techniques from Analytic Number Theory such as Partial Summation, cancellations with sums over primes with different remainders, as well as estimates for sums over primes. It is worth noting that despite the complexity of some of the integrands, when evaluating them at \sqrt{x} we exhibit a lot of cancellations and the result at the end is simpler than expected. It is possible there are structural reasons behind this, but we cannot completely understand this phenomenon for now.
- Finally, we combine the estimates to derive a lower bound for the initial expression, which turns out to be positive for $x \ge 10^{21}$. The bound is simple enough so that we are able to give a direct proof of the positivity through the derivative.

We now introduce explicit bounds concerning prime counting functions. Firstly, we state an explicit version of the Prime Number Theorem.

Theorem 2.1 (Dusart [4], Rosser-Schoenfeld [15]). We have

$$\frac{x}{\log x} \le \pi(x) \le \frac{x}{\log x} + \frac{x}{(\log x)^2} + \frac{2.51x}{(\log x)^3}$$

where the lower bound holds for $x \ge 17$, and the upper bound is for $x \ge 355991$.

The corresponding version of the Prime Number Theorem for Arithmetic Progressions is as follows.

Theorem 2.2 (Bennet, Martin, O'Bryant, Rechnitzer, Corollary 1.6 in [11] for q = 4). For $x \ge 800$ and $a \in \{1,3\}$ we have

$$\frac{x}{2\log x} < \pi(x,4,a) < \frac{x}{2\log x} + \frac{5x}{4(\log x)^2}.$$

Because of applications of Partial Summation with the sequence with 1/n for prime n, we make use of the following bound.

Theorem 2.3 (Mertens, [14]). For all $x \ge 3$ we have

$$\left|\sum_{p \le x} \frac{1}{p} - \log \log x - M\right| \le \frac{4}{\log(x+1)} + \frac{2}{x \log x}$$

where $M \approx 0.26149$ is the **Meissel-Mertens constant**.

Remark. We shall also make use of the simpler version $\left|\sum_{p\leq x} \frac{1}{p} - \log \log x\right| \leq 1$ for $x \geq 3$. It holds for $x \geq 227$ since $0 \leq M \leq 1$ and $\frac{4}{\log(x+1)} + \frac{2}{x\log x} \leq 1 - M$ (the left-hand side is decreasing). Note that $\log \log x$ is in [0.09, 0.48) for $x \in [3, 5)$, in [0.48, 1.05) for $x \in [5, 17)$ and in [1.05, 1.7] for $x \in [17, 227]$; while $\sum_{p\leq x} \frac{1}{p}$ is $\frac{5}{6}$ for $x \in [3, 5)$, in [0.83, 1.35) for $x \in [5, 17)$ and in [1.35, 1.97] for $x \in [17, 227]$, so indeed the inequality holds for all $x \in [3, 227]$, as well.

We also need bounds which include small x.

Corollary 2.4. For $x \ge 2$ and $a \in \{1,3\}$ we have the following upper bounds

$$\pi(x) \le \frac{x}{\log x} + \frac{x}{(\log x)^2} + \frac{2.54x}{(\log x)^3}, \quad \pi(x, 4, a) \le \frac{x}{2\log x} + \frac{5x}{4(\log x)^2}.$$

We also have the lower bounds

$$\pi(x,4,3) > \frac{x}{2\log x}$$
 for $x \ge 19$ and $\pi(x,4,1) > \frac{x}{2\log x}$ for $x \ge 228$.

Proof. We illustrate only the inequality with $\pi(x)$, the rest are analogous. By Theorem 2.1 it is enough to justify it for $x \leq 355991$. For all integers in [2,355991] a Mathematica computation shows it is correct. Regarding non-integer x, in each interval $(a, a + 1), a \in \mathbb{Z}$, the function

$$h(x) = \frac{x}{\log x} + \frac{x}{(\log x)^2} + \frac{2.54x}{(\log x)^3} - \pi(x), \text{ with } h'(x) = \frac{(\log x)^3 + 0.54\log x - 7.62}{(\log x)^4}$$

is differentiable and the numerator of h'(x) is strictly increasing, with a unique root at $x = x_0 \approx 6.5301$. Hence h(x) is monotonic in each (a, a + 1) except for a = 6, so checking that $h(x_0) > 0$ and knowing that h(x) > 0 when x is an integer completes the proof.

3 Main proof

We are now ready to prove the main theorem.

Proof of Theorem 1.4. We count the products of two primes $pq \leq x$ with $p \equiv q \equiv 3 \pmod{4}$. Any such product can be represented with $p \leq q \leq \frac{x}{p}$ where $p \leq \sqrt{x}$. The count for products $pq \leq x$ with p, q odd is analogous. Hence the desired statement

$$\frac{\#\{m \le x : m = pq \text{ with } p \equiv q \equiv 3 \pmod{4}\}}{\#\{m \le x : m = pq \text{ with } p, q \text{ odd}\}} > \frac{1}{4}$$

is equivalent to

$$W(x) := \sum_{\substack{p \le \sqrt{x} \\ p \equiv 3 \ (4)}} \sum_{\substack{p \le q \le x/p \\ q \equiv 3 \ (4)}} 1 - \frac{1}{4} \sum_{\substack{2 0.}$$

As a consequence of Corollary 2.4, we have

$$\sum_{\substack{p \le q \le x/p \\ q \equiv 3}} 1 = \sum_{\substack{q \le x/p \\ q \equiv 3}} 1 - \sum_{\substack{q$$

Now note the simple observation that $\frac{1-(-1)^{\frac{p-1}{2}}}{2}$ equals 1 if $p \equiv 3 \pmod{4}$ and 0 if $p \equiv 1 \pmod{4}$. Thus,

$$\sum_{\substack{p \le \sqrt{x} \\ p \equiv 3 \, (4)}} \sum_{\substack{p \le q \le x/p \\ q \equiv 3 \, (4)}} 1 \ge \sum_{2$$

with the latter being equal to

$$\frac{1}{4} \sum_{2 2} \frac{(-1)^{\frac{p-1}{2}}}{p} + \frac{x}{4 \log x} \sum_{p > \sqrt{x}} \frac{(-1)^{\frac{p-1}{2}}}{p} - \frac{x}{4 \log x} \sum_{2$$

Next, by Corollary 2.4

$$\sum_{2$$

Hence the main target W(x) is bounded below by

$$-\frac{x}{4\log x}\sum_{p>2}\frac{(-1)^{\frac{p-1}{2}}}{p} + \frac{x}{4\log x}\sum_{p>\sqrt{x}}\frac{(-1)^{\frac{p-1}{2}}}{p} - \frac{x}{4\log x}\sum_{2< p\le\sqrt{x}}\frac{(-1)^{\frac{p-1}{2}}\log p}{p\log\frac{x}{p}} - \sum_{\substack{p\le\sqrt{x}\\p\equiv3(4)}}\left(\frac{p}{2\log p} + \frac{5p}{4(\log p)^2}\right) - \frac{x}{4}\sum_{2< p\le\sqrt{x}}\frac{1}{p(\log\frac{x}{p})^2} - 0.635x\sum_{2< p\le\sqrt{x}}\frac{1}{p(\log\frac{x}{p})^3}.$$

Now we deal with each of the six sums separately.

• Regarding $\sum_{p>2} \frac{(-1)^{\frac{p-1}{2}}}{p}$, the work of Dummit-Granville-Kisilevsky [2] asserts that the decimal expansion of this sum is (-0.334...). For completeness, let us justify that the sum is convergent. Consider $L(s) = \sum_{k=0}^{\infty} (-1)^k (2k+1)^{-s}$ where $s \in \mathbb{C}$. For $\operatorname{Re}(s) > 1$ the Euler Product Formula gives

 $\sum_{k=0}^{\infty} (1) (2k+1) \quad \text{where } b \in \mathbb{C}. \text{ For } \operatorname{Her}(b) \neq 1 \text{ the Dater Frequency formula gives}$

$$\log L(s) = \log \prod_{p \ge 3} \frac{1}{1 - (-1)^{\frac{p-1}{2}} p^{-s}} = -\sum_{p \ge 3} \log \left(1 - (-1)^{\frac{p-1}{2}} p^{-s} \right) = \sum_{p \ge 3} \sum_{k=1}^{\infty} \frac{(-1)^{\frac{-3}{2}}}{k p^{ks}} \tag{1}$$

where in the last step we have applied the Taylor expansion of log(1 - x). Additionally, the Taylor expansion of arctan(x) yields

$$L(1) = \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} = \arctan(1) = \frac{\pi}{4} > 0$$

and hence (1) extends to s = 1. In particular,

$$\sum_{p} \frac{(-1)^{\frac{p-1}{2}}}{p} = \log \frac{\pi}{4} - \sum_{k=2}^{\infty} \sum_{p} \frac{(-1)^{\frac{k(p-1)}{2}}}{kp^k}.$$

It now suffices to justify that the double series converges absolutely, e.g. via the comparison

$$\sum_{k=2}^{\infty} \sum_{p} \left| \frac{(-1)^{\frac{k(p-1)}{2}}}{kp^k} \right| \le \frac{1}{2} \sum_{p} \sum_{k=2}^{\infty} \frac{1}{p^k} = \frac{1}{2} \sum_{p} \frac{1}{p(p-1)} < \frac{1}{2} \sum_{p} \frac{1}{(p-1)^2} < \frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$

and since $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges, the corresponding claim for $\sum_p \frac{(-1)^{\frac{p-1}{2}}}{p}$ follows.

• For $\sum_{p>\sqrt{x}} \frac{(-1)^{\frac{p-1}{2}}}{p}$ we split according to the remainder $b \in \{1,3\}$ modulo 4 of p and then use Partial summation with the function $\frac{1}{t}$ and the indicator sequence of primes with remainder b. This gives

$$\sum_{p>\sqrt{x}} \frac{(-1)^{\frac{p-1}{2}}}{p} = \sum_{b\in\{1,3\}} (-1)^{\frac{b-1}{2}} \sum_{\substack{p>\sqrt{x}\\p\equiv b\ (4)}} \frac{1}{p}$$
$$= \sum_{b\in\{1,3\}} (-1)^{\frac{b-1}{2}} \left(-\frac{\pi(\sqrt{x},4,b)}{\sqrt{x}} + \int_{\sqrt{x}}^{\infty} \frac{\pi(t,4,b)}{t^2} \right)$$
$$= \frac{\pi(\sqrt{x},4,3) - \pi(\sqrt{x},4,1)}{\sqrt{x}} + \int_{\sqrt{x}}^{\infty} \frac{\pi(t,4,1) - \pi(t,4,3)}{t^2} dt$$

By Theorem 2.2 we have for $x \ge 800$

$$|\pi(x,4,3) - \pi(x,4,1)| \le \frac{5x}{4(\log x)^2}$$

Therefore

$$\sum_{p > \sqrt{x}} \frac{(-1)^{\frac{p-1}{2}}}{p} \ge -\frac{5}{(\log x)^2} - \frac{5}{4} \int_{\sqrt{x}}^{\infty} \frac{dt}{t(\log t)^2} = -\frac{5}{2\log x} - \frac{5}{(\log x)^2}.$$

• Using Partial Summation and Corollary 2.4 gives

$$\sum_{\substack{p \le \sqrt{x} \\ p \equiv 3 \, (4)}} \frac{p}{\log p} = \frac{3}{\log 3} + \frac{7}{\log 7} + \frac{11}{\log 11} + \pi(\sqrt{x}, 4, 3) \frac{2\sqrt{x}}{\log x} - \int_{19}^{\sqrt{x}} \pi(t, 4, 3) \frac{\log t - 1}{(\log t)^2} dt$$
$$\leq 10.92 + \frac{2x}{(\log x)^2} + \frac{10x}{(\log x)^3} - \frac{1}{2} \int_{19}^{\sqrt{x}} \frac{(\log x - 1)x}{(\log x)^3} dt$$
$$= 10.92 + \frac{x}{(\log x)^2} + \frac{10x}{(\log x)^3} + \frac{361}{4(\log 19)^2} \le \frac{x}{(\log x)^2} + \frac{10x}{(\log x)^3} + 21.33$$

Similarly, we compute

$$\begin{split} \sum_{\substack{p \leq \sqrt{x} \\ p \equiv 3 \, (4)}} \frac{p}{(\log p)^2} &= \frac{3}{(\log 3)^2} + \frac{7}{(\log 7)^2} + \frac{11}{(\log 11)^2} + \pi(\sqrt{x}, 4, 3) \frac{4\sqrt{x}}{(\log x)^2} - \int_{19}^{\sqrt{x}} \pi(t, 4, 3) \frac{\log t - 2}{(\log t)^3} \, dt \\ &\leq \frac{25}{4} + \frac{4x}{(\log x)^3} + \frac{20x}{(\log x)^4} - \frac{1}{2} \int_{19}^{\sqrt{x}} \frac{t(\log t - 2)}{(\log t)^4} \, dt \\ &= \frac{25}{4} + \frac{4x}{(\log x)^3} + \frac{20x}{(\log x)^4} - \frac{1}{2} \int_{19}^{\sqrt{x}} \frac{t(\log t - \frac{3}{2})}{(\log t)^4} \, dt + \frac{1}{4} \int_{19}^{\sqrt{x}} \frac{t}{(\log t)^4} \, dt \\ &= 9.7855 + \frac{2x}{(\log x)^3} + \frac{20x}{(\log x)^4} + \frac{1}{4} \int_{19}^{\sqrt{x}} \frac{t}{(\log t)^4} \, dt. \end{split}$$

The function $\frac{t}{(\log t)^4}$ is increasing in $[e^4, \infty)$, hence partitioning $[19, \sqrt{x}]$ as $[19, e^4] \cup [e^4, \sqrt{x}]$ and taking the maximum of the integrand in the second interval yields

$$\sum_{\substack{p \le \sqrt{x} \\ p \equiv 3 \, (4)}} \frac{p}{(\log p)^2} \le 11.7844 + \frac{2x}{(\log x)^3} + \frac{20x}{(\log x)^4} + (\sqrt{x} - e^4) \frac{4\sqrt{x}}{(\log x)^4}$$
$$= 11.7844 + \frac{2x}{(\log x)^3} + \frac{24x}{(\log x)^4} - \frac{4e^4\sqrt{x}}{(\log x)^4}.$$

Therefore (dropping the $\sqrt{x}/(\log x)^4$ term)

$$\sum_{\substack{p \le \sqrt{x} \\ p \equiv 3(4)}} \left(\frac{p}{2\log p} + \frac{5p}{4(\log p)^2} \right) \le 25.3955 + \frac{x}{2(\log x)^2} + \frac{15x}{2(\log x)^3} + \frac{30x}{(\log x)^4}.$$

• For a fixed x, we use Partial Summation with the function $1/\left(\log \frac{x}{t}\right)^2$ (on the variable t) and the sequence $\frac{1}{n}$ for n prime. Together with Mertens' estimate in the form of Theorem 2.3, but with the

right-hand side replaced by the slightly larger $\frac{9}{2\log x}$, we obtain

$$\begin{split} \sum_{3 \le p \le \sqrt{x}} \frac{1}{p(\log \frac{x}{p})^2} &= \frac{\sum_{p \le \sqrt{x}} \frac{1}{p} - \frac{1}{2}}{(\log \sqrt{x})^2} - \int_3^{\sqrt{x}} \left(\sum_{p \le t} \frac{1}{p} - \frac{1}{2} \right) \frac{2}{t(\log \frac{x}{t})^3} dt \\ &\le \frac{4\left(\log \log x - 0.23 + \frac{9}{2\log x} \right)}{(\log x)^2} - \int_3^{\sqrt{x}} \left(\log \log t - \frac{9}{2\log t} - 0.24 \right) \frac{2}{t(\log \frac{x}{t})^3} dt \\ &= \frac{4\log \log x}{(\log x)^2} - \frac{0.92}{(\log x)^2} + \frac{18}{(\log x)^3} - \left(\frac{4\log \log x}{(\log x)^2} - \frac{74}{25(\log x)^2} - \frac{36}{(\log x)^3} \right) + A(x) \\ &= \frac{2.04}{(\log x)^2} + \frac{54}{(\log x)^3} + A(x) \end{split}$$

where A(x) is the contribution from the lower limit 3 in the integral. The expression A(x) consists of the sum $\frac{\log \log \frac{x}{3}}{(\log x)^2} + \frac{9\log \log \frac{x}{3}}{(\log x)^2} - \frac{31\log \frac{x}{3}}{25(\log x)^3}$ and a sum of fractions whose denominators are $(\log x)^3$, $(\log \frac{x}{3})(\log x)^3$ and $(\log \frac{x}{3})^2(\log x)^3$ and whose numerators are constants. Having in mind $1.098 < \log 3 < 1.1$ and $0.094 < \log \log 3 < 0.095$, it turns out that these numerators are all negative. Hence we can drop the terms with denominators $(\log \frac{x}{3})(\log x)^3$ and $(\log \frac{x}{3})^2(\log x)^3$, as well as the $\frac{31\log \frac{x}{3}}{25(\log x)^3}$ term, to obtain a simplified upper bound for A(x). Overall, we reach

$$\sum_{3 \le p \le \sqrt{x}} \frac{1}{p(\log \frac{x}{p})^2} \le \frac{\log \log x}{(\log x)^2} + \frac{9\log \log x}{(\log x)^3} + \frac{2.04}{(\log x)^2} + \frac{37}{(\log x)^3}$$

• An analogous approach, with the function $1/\left(\log \frac{x}{t}\right)^3$ (on the variable t), but with the simpler inequality $\left|\sum_{p \le x} \frac{1}{p} - \log \log x\right| \le 1$ in the remark after Theorem 2.3, yields

$$\begin{split} \sum_{3 \le p \le \sqrt{x}} \frac{1}{p \left(\log \frac{x}{p}\right)^3} &= \frac{\sum_{2$$

(We compute the integral and then use $\log \frac{x}{3} \le \log x$ and $\log \log 3 < \frac{3}{2} < \frac{5}{3} < \frac{13}{6}$ for simplicity.)

• For $\sum_{3 \le p \le \sqrt{x}} \frac{(-1)^{\frac{p-1}{2}} \log p}{p \log \frac{x}{p}}$ we firstly justify $\sum_{3 \le p \le 227} \frac{(-1)^{\frac{p-1}{2}} \log p}{p \log \frac{x}{p}} \le 0$ for $x \ge 10^6$. Denote $g(t) = \frac{\log t}{t \log \frac{x}{t}}$ for $t \in [3, 227]$. We have g'(t) < 0 if and only if $\frac{1}{\log t} + \frac{1}{\log \frac{x}{t}} < 1$ and the latter holds for $t \ge 4$ by $\frac{1}{\log 4} + \frac{1}{\log \frac{106}{227}} < 1$ and for $t \in [3, 4]$ by $\frac{1}{\log 3} + \frac{1}{\log \frac{106}{4}} < 1$. Hence g is decreasing and so it suffices to note that g(a) > g(b) for the following pairs (a, b):

(3,5), (7,13), (11,17), (19,29), (23,37), (31,41), (43,53), (47,61), (59,73), (67,89), (71,97), (79,101), (79,100), (79,10), (79,10), (79,10), (79,10), (79,10), (79,10), (79,

(83, 109), (103, 113), (107, 137), (127, 149), (131, 157), (139, 173), (151, 181), (163, 193), (167, 197)and that g(179), g(191), g(199), g(211), g(223) and g(227) are positive. The rest we split into residue classes and use Partial Summation (with $\frac{\log t}{t \log \frac{\pi}{t}}$) to get

$$\sum_{229 \le p \le \sqrt{x}} \frac{(-1)^{\frac{p-1}{2}} \log p}{p \log \frac{x}{p}} = \sum_{b \in \{1,3\}} (-1)^{\frac{b-1}{2}} \sum_{\substack{229 \le p \le \sqrt{x} \\ p \ge b \ (4)}} \frac{\log p}{p \log \frac{x}{p}}$$
$$= \sum_{b \in \{1,3\}} (-1)^{\frac{b-1}{2}} \left[\frac{1}{\sqrt{x}} \sum_{\substack{229 \le p \le \sqrt{x} \\ p \ge b \ (4)}} 1 - \int_{229}^{\sqrt{x}} \frac{\log x - \log t \log(\frac{x}{t})}{t^2 (\log \frac{x}{t})^2} \sum_{\substack{229 \le p \le t \\ p \ge b \ (4)}} 1 dt \right].$$

There are 48 odd primes less than 229, from which 27 are 3 modulo 4 and 21 are 1 modulo 4. Hence the above becomes

$$\sum_{b \in \{1,3\}} (-1)^{\frac{b-1}{2}} \left[\frac{\pi(\sqrt{x},4,b) - \pi(228,4,b)}{\sqrt{x}} + \int_{229}^{\sqrt{x}} \frac{\log t \log(\frac{x}{t}) - \log x}{t^2 (\log \frac{x}{t})^2} (\pi(t,4,b) - \pi(228,4,b)) dt \right]$$
$$= \frac{\pi(\sqrt{x},4,1) - \pi(\sqrt{x},4,3) + 6}{\sqrt{x}} + \int_{229}^{\sqrt{x}} \frac{\log t \log(\frac{x}{t}) - \log x}{t^2 (\log \frac{x}{t})^2} [\pi(t,4,1) - \pi(t,4,3) + 6] dt$$

By Corollary 2.4 we have $\frac{t}{2\log t} < \pi(t,4,b) \le \frac{t}{2\log t} + \frac{5t}{4(\log t)^2}$ for all $t \ge 229$ and $b \in \{1,3\}$. Having in mind $\log t \log \frac{x}{t} \ge \log x$ for $3 \le t \le \sqrt{x}$ and $x \ge 3 \cdot 10^5$ (since $\log t \log \frac{x}{t}$ increases as t increases), we obtain as an upper bound for $\sum_{3 \le p \le \sqrt{x}} \frac{(-1)^{\frac{p-1}{2}\log p}}{p\log \frac{x}{p}}$ the expression

$$\begin{aligned} & \frac{6}{\sqrt{x}} + \frac{5}{(\log x)^2} + \int_{229}^{\sqrt{x}} \frac{\log t \log(\frac{x}{t}) - \log x}{t^2 (\log \frac{x}{t})^2} \left(\frac{5t}{4(\log t)^2} + 6\right) dt \\ &= \frac{6}{\sqrt{x}} + \frac{5}{(\log x)^2} + 6 \int_{229}^{\sqrt{x}} \frac{\log t \log(\frac{x}{t}) - \log x}{t^2 (\log \frac{x}{t})^2} dt + \frac{5}{4} \int_{229}^{\sqrt{x}} \left(\frac{1}{t \log \frac{x}{t} \log t} - \frac{\log x}{t(\log \frac{x}{t})^2 (\log t)^2}\right) dt \\ &= \frac{5 \log \log \frac{x}{229}}{4 \log x} - \frac{5 \log \log \frac{x}{229}}{2(\log x)^2} - \frac{5(\log \log 229 + \frac{1}{\log 229})}{4 \log x} + \frac{6 \log 229}{229 \log \frac{x}{229}} + \frac{25 + 10 \log \log 229}{4(\log x)^2} + \frac{5 \log 229}{4(\log x)^2}. \end{aligned}$$

To simplify the latter, proceed as follows. For the first term use $\log \log \frac{x}{229} < \log \log x$, for the second use $\log \log \frac{x}{229} > \frac{9}{10} \log \log x$ when $x \ge 2 \cdot 10^9$, for the third use $\log \log 229 > 1.692$ and $\frac{1}{\log 229} > 0.184$, for the fourth and sixth one use $\log 229 < 5.44$ and $\log \frac{x}{229} > \frac{3}{4} \log x$ when $x \ge 3 \cdot 10^9$, for the fifth one use $\log \log 229 < 1.696$. Overall, this yields the upper bound

$$\frac{5\log\log x}{4\log x} - \frac{9\log\log x}{4(\log x)^2} - \frac{54}{25\log x} + \frac{21}{2(\log x)^2} + \frac{136}{15(\log x)^3}.$$

Collecting all bounds. In conclusion, the main target W(x) is bounded below by

$$\frac{0.0835x}{\log x} - \frac{9x\log\log x}{16(\log x)^2} - \frac{2.3225x\log\log x}{(\log x)^3} - \frac{1.095x}{(\log x)^2} - \frac{165x}{8(\log x)^3} - \frac{484x}{15(\log x)^4} - 25.3955x$$

which turns out to be positive for $x \ge 2.04 \cdot 10^{20}$. To see this, compute the derivative of the expression and notice (by dropping appropriate positive terms) that its positivity is implied by the inequality $0.0835(\log x)^2 - (0.5625 \log \log x + 1.1785) \log x - (1.1795 \log \log x + 18.9975) \ge 0$, i.e.

$$\frac{0.5625\log\log x + 1.1785}{\log x} + \frac{1.1795\log\log x + 18.9975}{(\log x)^2} \le 0.0835$$

which holds for $x \ge 10^{20}$ as the left-hand side is decreasing. This completes the proof.

Acknowledgments. We would like to thank Angel Raychev for proof-checking particular technical details, as well as Iliyas Noman, Radostin Cholakov, Vladimir Mitankin, and representatives of the American Mathematical Society and Mu Alpha Theta at Regeneron ISEF 2023 for general comments on the paper. This research was supported by the Summer Research School at the High School Students Institute of Mathematics and Informatics (HSSIMI) under an Education with Science grant in Bulgaria and the provided atmosphere for work is greatly appreciated.

References

- P. L. Chebyshev. Lettre de m. le professeur tchébychev à m. fuss sur un nouveaux théorème relatif aux nombres premiers contenus dans les formes 4n + 1 et 4n + 3. Bull. Phys. Acad. Sci. St. Petersburg, 11, 1853.
- [2] A. Granville D. Dummit and H. Kisilevsky. Big biases amongst products of two primes. *Mathematika*, 62(2):502–507, 2016.
- [3] de la Valee Poussin. Recherches analytiques sur la théorie des nombres premiers. Annales de la Société scientifique de Bruxelles, Imprimeur de l'Académie Royale de Belgique, 1896.
- [4] P. Dusart. Autour de la fonction qui compte le nombre de nombres premiers. 1998.
- [5] G. H. Hardy and J. E. Littlewood. Contributions to the theory of the riemann zeta-function and the theory of the distribution of primes. *Acta Mathematica*, 41:119–196, 1916.
- [6] P. Hough. A lower bound for biases amongst products of two primes. Research in Number Theory, 3:1–11, 2017.
- [7] J. Kaczorowski. A contribution to the shanks-rényi race problem. The Quarterly Journal of Mathematics, 44(4):451-458, 1993.
- [8] S. Knapowski and P. Turán. Comparative prime-number theory. i. Acta Mathematica Hungarica, 13(3):299–314, 1962.
- [9] E. Landau. Über einige ältere vermutungen etc. zweite abhandlung. Math. Zeitschr., 1:213–219, 1918.
- [10] J. Littlewood. Sur la distribution des nombres premiers. Comptes Rendus, 1914.
- [11] K. O'Bryant M. A. Bennett, G. Martin and A. Rechnitzer. Explicit bounds for primes in arithmetic progressions. *Illinois Journal of Mathematics*, 62(1-4):427–532, 2018.
- [12] M. Marinov. Biases amongst singleton primes, products of primes and pairs of consecutive primes. Oxford Mathematics MSc Thesis, 2020.
- [13] X. Meng. The distribution of k-free numbers and integers with fixed number of prime factors. 2017.
- [14] F. Mertens. Ein beitrag zur analytischen zahlentheorie. 1874.
- [15] J. B. Rosser and L. Schoenfeld. Approximate formulas for some functions of prime numbers. *Illinois Journal of Mathematics*, 6(1):64–94, 1962.
- [16] M. Rubinstein and P. Sarnak. Chebyshev's bias. *Experimental Mathematics*, 3(3):173–197, 1994.