Nested Instrumental Variables Design: Switcher Average Treatment Effect, Identification, Efficient Estimation and Generalizability

Rui Wang¹, Ying-Qi Zhao², Oliver Dukes³, and Bo Zhang^{* 4}

¹Department of Biostatistics, University of Washington

²Public Health Sciences Division, Fred Hutchinson Cancer Center

³Department of Applied Mathematics, Computer Science and Statistics, Ghent University

⁴Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center

Abstract: Instrumental variables (IV) are a commonly used tool to estimate causal effects from non-randomized data. A prototype of an IV is a randomized trial with non-compliance where the randomized treatment assignment serves as an IV for the non-ignorable treatment received. Under a monotonicity assumption, a valid IV non-parametrically identifies the average treatment effect among a non-identifiable complier subgroup, whose generalizability is often under debate. In many studies, there could exist multiple versions of an IV, for instance, different nudges to take the same treatment in different study sites in a multi-center clinical trial. These different versions of an IV may result in different compliance rates and offer a unique opportunity to study IV estimates' generalizability. In this article, we introduce a novel nested IV assumption and study identification of the average treatment effect among two latent subgroups: always-compliers and switchers, who are defined based on the joint potential treatment received under two versions of a binary IV. We derive the efficient influence function for the SWitcher Average Treatment Effect (SWATE) and propose efficient estimators. We then propose formal statistical tests of the generalizability of IV estimates based on comparing the conditional average treatment effect among the always-compliers and that among the switchers under the nested IV framework. We apply the proposed framework and method to the Prostate, Lung, Colorectal and Ovarian (PLCO) Cancer Screening Trial and study the causal effect of colorectal cancer screening and

^{*}Assistant Professor of Biostatistics, Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center. Email: bzhang3@fredhutch.org

its generalizability.

Keywords: Generalizability; Instrumental variable; Noncompliance; Principal stratification; Semiparametric theory

1 Introduction

1.1 Instrumental variable (IV); complier average treatment effect; generalizability

Scientific queries often involve exploring cause-and-effect relationships. Randomization is a powerful tool for drawing valid causal inference, and randomized controlled trials (RCTs) are considered the gold standard for causal inference. On the other hand, drawing a valid causal conclusion from non-randomized studies is often complicated by many factors, unmeasured confounding being one of the major concerns. Facing potential unmeasured confounding bias, researchers often resort to quasi-experimental devices (Cook et al., 1979). Instrumental variable (IV) methods are among the most popular tools (Angrist et al., 1996; Hernán and Robins, 2006; Baiocchi et al., 2014). A variable is a valid IV if it is correlated with the treatment, does not have a direct effect on the outcome under investigation and is independent of unmeasured treatment-outcome confounders (see, e.g., Baiocchi et al., 2014). Econometricians popularized IV methods under the framework of linear structural equations with omitted variables (see, e.g., Goldberger, 1972). Angrist et al. (1996) first formalized an IV under the potential outcomes (PO) framework (Neyman, 1923; Rubin, 1974), defined the complier average treatment effect (CATE) as a key estimand and studied its nonparametric identification under a monotonicity assumption that states a person's treatment received cannot decrease as the level of IV increases. The concept of identifying and estimating causal effects among a latent subpopulation defined by potential outcomes under both treatment assignments (referred to as "cross-world" potential outcomes) was referred to as principal stratification and generalized to other contexts, including surrogate endpoint, truncation by death, among others (Frangakis and Rubin, 2002; Zhang and Rubin, 2003; Gilbert and Hudgens, 2008).

A key critique towards the CATE framework, and principal stratification in general, is that the CATE only concerns the causal effect on a latent subgroup and cannot be generalized to the entire population without further assumptions; see, e.g., the extensive discussion in Deaton (2009), Heckman and Urzua (2010), Joffe (2011), among others. One strand of work seeks to illuminate the relationship between the CATE and the average treatment effect (ATE) by proposing suitable identification assumptions that point identify the ATE with a valid IV. These assumptions typically impose some homogeneity or no-interaction-type restrictions on the data generating processes (Swanson et al., 2018; Levis et al., 2024). For instance, in the case of binary IV and binary treatment, Robins (1994) proposed the no-current-treatment-interaction assumption, which states that there is no additive interaction between the instrument and covariates in the model for conditional average treatment effect. An alternative assumption is Tchetgen Tchetgen and Vansteelandt's (2013) homogeneous selection bias assumption, which states that the selection bias function doesn't depend on the value of instrument on an additive scale. More recently, Wang and Tchetgen Tchetgen (2018) showed that if there is no additive unmeasured confounder-IV interaction in the model for treatment or additive unmeasured confounder-treatment interaction in the model for potential outcomes, then the ATE can be identified by an IV. These assumptions are typically not empirically verifiable and scale-specific (i.e., only suitable for an additive or multiplicative scale). Under the principal stratification framework, a similar assumption called principal ignorability, which states that there is no difference in the average potential outcomes across principal strata conditional on baseline covariates, is needed for generalizability (Jo and Stuart, 2009). Unfortunately, the principal ignorability assumption is also not empirically verifiable.

1.2 A clinical trial with two versions of IV; nested IV

The archetype of an IV is an RCT with non-compliance, where the treatment assignment in the RCT constitutes a randomized encouragement to receive the treatment and hence an IV for the actual treatment uptake. The Prostate, Lung, Colorectal and Ovarian (PLCO) Cancer Screening Trial is a multiphasic RCT conducted from 1993 to 2011 at 10 US institutions. A total of 154, 897 participants, aged 55 to 74 years, were enrolled and randomized to the intervention or control arm. Participants in the intervention arm were offered cancer screening exams while the control arm participants were advised to receive their usual medical care.

Two methods of consent were used in this trial. Seven centers used a single consent process while the remaining three initially adopted a dual consent process. For centers adopting a single consent process, enrollment and randomization were consented at the same time. For centers adopting a dual consent process, participants were not informed of the study design, including the type of intervention, at enrollment and the first consent; later, enrolled participants who were randomized to the intervention arm were informed of the treatment assignment. Participants would need to consent for administration of screen exams (second consent). Results indicated that the compliance rate among participants under a single consent process was 91.3%, while that among participants under a dual consent process dropped to 71.2%. Because the rate of refusal to receive the screening intervention among dual-consent participants was deemed too high, the three screening centers later switched to the single consent process. In a *post hoc* analysis, Marcus et al. (2014) found that the single versus dual consent process was significantly associated with compliance (odds ratio = 2.2; 95% CI: 2.0 to 2.5), even after adjusting for a rich set of baseline characteristics including age at randomization, race and ethnicity, gender, education, BMI, comorbidity score, among others.

The transition from single to dual consent process appeared to create a natural experiment that *strengthened* the encouragement to comply with the treatment assignment. Given Marcus et al.'s (2014) analysis, it is not unreasonable to assume that a complier under the dual consent process would remain a complier had the person enrolled in the study during the single consent stage. We will refer to the assumption that a complier under one binary IV would remain a complier under a second strengthened IV as a "nested IV" assumption. As another motivating example, consider multiple large-scale HIV prevention trials conducted by the Microbicide Trials Network (MTN) and HIV Prevention Trials Network (HPTN). In different trials testing the same HIV prevention modality (e.g., the daily oral prophylaxis) in similar study populations (e.g., heterosexual women in sub-Saharan Africa), the compliance rate could be rather different due to a variety of reasons (Cohen and Baden, 2012; He et al., 2023), including the method of encouragement to adhere to the prescribed daily pills. A nested IV assumption applied to an integrated analysis of these studies would then state that a participant who was a complier in a trial with a low compliance rate would remain a complier in a trial with a higher compliance rate.

1.3 Implications of a nested IV assumption

While a conventional IV analysis defines latent subpopulations – compliers, always-takers, nevertakers, and defiers – based on potential treatment received under two realizations of one binary IV (Angrist et al., 1996), the nested IV assumption and its associated framework would define subpopulations based on potential treatment received under $2 \times 2 = 4$ realizations of two binary IVs and impose structures on these latent subpopulations in a way analogous to the monotonicity assumption in the original paper by Angrist et al. (1996). Under the nested IV framework. the average treatment effect on two latent subpopulations, referred to as "always-compliers" and "switchers," respectively, can be non-parametrically identified under suitable conditions. "Alwayscompliers" are participants who would be compliers with respect to either IV; for instance, in the PLCO trial, these are participants who would be compliers regardless of the consent process. On the other hand, a non-overlapping latent subpopulation called "switchers" are participants who would be a non-complier (i.e., an always-taker or a never-taker under Angrist et al.'s (1996) monotonicity assumption) under the weaker IV but become a complier under a strengthened IV; in the PLCO trial, these are participants who would not comply under a dual consent process but comply under a single consent process. In addition to identifying and estimating the causal effect among "always-compliers" and "switchers," a comparison of the causal effects on these two disjoint subpopulations, possibly conditional on a set of baseline covariates, helps illuminate the generalizability of IV-based estimates and formally test the no-interaction/principal-ignorability-type assumptions. Such progress towards establishing the generalizability of IV estimates, to the best of our knowledge, is a novel and useful contribution to the literature.

The rest of the article is organized as follows. In Section 2, we describe the nested IV design, define the switcher average treatment effect (SWATE) and present main identification results. In Section 3, we propose estimators for SWATE. In Section 4, we propose formal hypothesis testing procedures that examine treatment effect homogeneity among latent subpopulations under the proposed nested IV framework. Numerical experiments are reported in Section 5. We examine the PLCO study using our proposed method and framework in Section 6 and conclude with a discussion in Section 7.

2 Nested IV design

2.1 Two versions of treatment assignment

Consider a clinical trial center in the PLCO study, say the Henry Ford Health System, that switched from a dual to a single consent process. The clinical trial comprised two stages. Let G = a denote the dual consent stage, and we use $Z \in \{0_a, 1_a\}$ to denote a participant being randomized to the control group $(Z = 0_a)$ or the intervention group $(Z = 1_a)$ in the dual consent stage. In parallel, in the single consent stage, denoted by G = b, we use $Z \in \{0_b, 1_b\}$ to denote a participant being randomized to the control $(Z = 0_b)$ group or the intervention $(Z = 1_b)$ group. Importantly, in either stage, participants randomized to the intervention arm may choose not to take the assigned cancer screening procedure, while participants randomized to the control arm may choose to conduct a cancer screening even if this is not part of their usual medical care. In other words, the randomized treatment assignment is only an encouragement, or a nudge, for participants to receive cancer screening, and participants retain the full right not to adhere to the treatment assignment.

Treatment assignment, or "encouragement to receive the treatment," may contain different elements in two trial stages (G = a or G = b) and hence resulted in different compliance rates; therefore, we use different notations, 1_a and 1_b (or 0_a and 0_b), to denote being assigned to the intervention (or control) arm and encouraged to receive cancer screening (or usual medical care). Our notation acknowledges that there may exist two versions of encouragement and that the Stable Unit Treatment Value Assumption (Rubin, 1974, 1980), or SUTVA, may be violated if one single notation (0 or 1) is used. To further fix ideas, consider a second example in the context of HIV prevention implementation trials; see, e.g., Lambdin et al. (2015). To best leverage the multiple effective HIV prevention modalities in the real world settings, scientists are studying the best implementation strategies to encourage people at elevated HIV risk to use the appropriate HIV prevention tools, like daily oral prophylaxis. To this end, trials with different mechanisms to improve drug uptake (e.g., different drug delivery methods) are often deployed. In this context, one could use G = a or G = b to denote two separate implementation trials, $Z = 1_a$ or $Z = 0_a$ to denote randomization to a first drug delivery method or control, and $Z = 1_b$ or $Z = 0_b$ to denote randomization to a second drug delivery method or control.

2.2 Potential outcomes and identification assumptions

We define potential outcomes and formulate causal estimands under Neyman and Rubin's potential outcomes framework (Neyman, 1923; Rubin, 1974). Suppose we observe n independent and identically distributed samples, $\mathcal{D} = \{O_i = (Z_i, \mathbf{X}_i, D_i, Y_i), i = 1, ..., n\}$, from a target population $P_0 \in \mathcal{P}$, where \mathcal{P} denotes a non-parametric model for the observed data distribution. Throughout the article, we use $\mathbb{E}_P[\cdot]$ to denote taking expectation over some distribution $P \in \mathcal{P}$ and $\mathbb{E}_{P_0}[\cdot]$ taking expectation over the target population. For participant *i*, X_i denotes a set of baseline covariates, Z_i a categorical instrumental variable taking value in $\{0_a, 0_b, 1_a, 1_b\}$, $D_i \in \{0, 1\}$ a binary variable indicating the treatment a participant actually received, and Y_i a real-valued outcome of interest. Let G_i denote a stratification variable. In the PLCO study, G_i indicates the trial stage and hence type of the consent process. We will refer to the tuple $(0_a, 1_a)$ (or $(0_b, 1_b)$) as an IV dose pair (Sun and Wüthrich, 2022). Following Angrist et al. (1996), we say a participant is a complier under the IV dose pair $\{0_g, 1_g\}$ if $\{D(0_g), D(1_g)\} = (0, 1)$, an always-taker if $\{D(0_g), D(1_g)\} = (1, 1)$, a never-taker if $\{D(0_g), D(1_g)\} = (0, 0)$, or a defier if $\{D(0_g), D(1_g)\} = (1, 0)$, $g \in \{a, b\}$. We first describe identification assumptions.

Assumption 1. (Standard IV assumptions).

- (i). (Consistency). D = D(Z = z) if Z = z, Y = Y(D = d, Z = z) if D = d and Z = z, for $d \in \{0, 1\}$ and $z \in \{0_a, 1_a, 0_b, 1_b\}$.
- (ii). (Exclusion restriction). Y(D = d, Z = z) = Y(D = d), for $d \in \{0, 1\}$ and $z \in \{0_a, 1_a, 0_b, 1_b\}$.
- (iii). (IV independence). $Z \perp \{D(Z = z), Y(D = d)\} \mid \mathbf{X} = \mathbf{X}, G = g$, for all d, z and g.
- (iv). (IV relevance). $\mathbb{E}_{P_0}[D \mid Z = 1_g, X] \mathbb{E}_{P_0}[D \mid Z = 0_g, X] \neq 0, g \in \{a, b\}.$

All four parts of Assumption 1 are standard IV identification assumptions and sometimes referred to as *core* IV assumptions (Angrist et al., 1996; Baiocchi et al., 2014). In a clinical trial like the PLCO study, the treatment assignment is most likely affecting clinical outcomes only via its effect on the treatment received, so exclusion restriction (Assumption 1(ii)) is often satisfied. Moreover, IV independence (Assumption 1(iii)) is automatically satisfied by design in a randomized trial, and IV relevance (Assumption 1(iv)) is verifiable and holds in most clinical trials.

In the nested IV framework, the IV dose pair that a participant receives depends on the stratification variable G: a participant will be randomly assigned to 0_a or 1_a if G = a, or 0_b or 1_b if G = b. In addition to the core IV assumptions, we consider the following set of *nested IV assumptions*:

Assumption 2. (Nested IV assumptions).

- (i). (Mean exchangeability over G). $\mathbb{E}_{P_0}[Y(Z = 1_g) Y(Z = 0_g) | G = g, \mathbf{X}] = \mathbb{E}_{P_0}[Y(Z = 1_g) Y(Z = 0_g) | \mathbf{X}]$ and $\mathbb{E}_{P_0}[D(Z = 1_g) D(Z = 0_g) | G = g, \mathbf{X}] = \mathbb{E}_{P_0}[D(Z = 1_g) D(Z = 0_g) | \mathbf{X}]$, $g \in \{a, b\}$.
- (ii). (Partial monotonicity). $D(Z = 1_g) \ge D(Z = 0_g), g \in \{a, b\}.$
- (iii). (Non-equal compliance).

 $\mathbb{E}_{P_0}[D \mid Z = 1_a, X = x] - \mathbb{E}_{P_0}[D \mid Z = 0_a, X = x] \neq \mathbb{E}_{P_0}[D \mid Z = 1_b, X = x] - \mathbb{E}_{P_0}[D \mid Z = 0_b, X = x].$

(iv). (Nested IV). $\{D(Z = 0_a), D(Z = 1_a)\} = (0, 1) \text{ implies } \{D(Z = 0_b), D(Z = 1_b)\} = (0, 1).$

Assumption 2(i) states that, conditional on the observed covariates X, the average effect of IV Z (either version) on D and that on Y can be identified using the corresponding stratum G = a or b. This assumption is satisfied, for example, if participating in the trial does not itself modify the effect of IV on D or Y (Dahabreh et al., 2019). This assumption is also satisfied if the stratification variable G is exogenous. For instance, in the PLCO study, the sudden shift from dual to single consent could be thought of as a happenstance for participants who enrolled right before and after this policy shift, analogous to the so-called "regression discontinuity design" (Thistlethwaite and Campbell, 1960). Assumption 2(ii) states that there is no defier with respect to either version of the IV. Note that we do not impose order structures across IV pairs, that is, we do not assume any ordering among $\{0_a, 0_b\}$, $\{0_a, 1_b\}$, $\{1_a, 0_b\}$, or $\{1_a, 1_b\}$; hence, this partial monotonicity assumption is weaker than the usual monotonicity assumption for an ordinal IV (Imbens and Angrist, 1994). Assumption 2(iii) states that the compliance rates corresponding to G = a and G = b are different; this is likely to hold when the IV is of different versions in two strata. Finally, Assumption 2(iv) states that if a participant is a complier under the IV pair $\{0_a, 1_a\}$, then the participant is also a complier under the IV pair $\{0_b, 1_b\}$. This assumption is key to the identification results and most likely to be satisfied when the binary IV $Z \in \{0_b, 1_b\}$ is stronger than the binary IV $Z \in$ $\{0_a, 1_a\}$. Empirically, the compliance rates can be calculated for both IV pairs, either marginally or conditional on a set of covariates as in Marcus et al. (2014), so the reasonableness of Assumption 2(iv) can often be assessed prior to invoking the nested IV framework.

2.3 Principal strata; causal estimands; identification results

A total of $2^4 = 16$ possible subgroups can be defined based on the joint potential outcomes

$${D(Z = 0_a), D(Z = 1_a), D(Z = 0_b), D(Z = 1_b)};$$

see Table 1. Assumptions 2(ii) (partial monotonicity) and 2(iv) (nested IV) together eliminate 10 out of 16 principal strata and the population then consists of the following 6 latent subgroups:

- 1. Always-Always-Takers (AAT): Always-takers under both IV pairs $\{0_a, 1_a\}$ and $\{0_b, 1_b\}$;
- 2. SWitchers (SW): Always-takers or never-takers under the IV pair $\{0_a, 1_a\}$ but compliers under the IV pair $\{0_b, 1_b\}$;
- 3. Always-Taker-Never-Taker (AT-NT): Always-takers under the IV pair $\{0_a, 1_a\}$ but nevertakers under the IV pair $\{0_b, 1_b\}$;
- 4. Always-COmplier (ACO): Compliers under both IV pairs $\{0_a, 1_a\}$ and $\{0_b, 1_b\}$;
- 5. Never-Taker-Always-Taker (NT-AT): Never-takers under the IV pair $\{0_a, 1_a\}$ but always-takers under the IV pair $\{0_b, 1_b\}$;
- 6. Always-Never-Taker (ANT): Never-takers under both IV pairs $\{0_a, 1_a\}$ and $\{0_b, 1_b\}$.

Remark 1 (One-sided non-compliance). In clinical trials assessing an investigational product, participants assigned placebo often do not have access to the treatment. This is referred to as "onesided non-compliance." In this stylistic case, we necessarily have $D(0_a) = D(0_b) = 0$, and the latent groups NT-AT, AT-NT and AAT in Table 1 no longer exist; moreover, the switcher subgroup now only comprises participants who are never-takers under $(0_a, 1_a)$ but become compliers under $(0_b, 1_b)$.

Let S denote principal stratum. Define the following conditional switcher average treatment effect:

$$\operatorname{SWATE}_{P_0}(\boldsymbol{X}) := \mathbb{E}_{P_0}[Y(D=1) - Y(D=0) \mid S = \operatorname{SW}, \boldsymbol{X}],$$

$D(0_a)$	$D(1_a)$	G = a	$D(0_b)$	$D(1_b)$	G = b	Exists?	Identification Assumption	S
0	0	NT	0	0	NT	Yes		ANT
1	0	DF	0	0	\mathbf{NT}	No	Partial monotonicity	
0	1	CO	0	0	\mathbf{NT}	No	Nested IV	
0	0	\mathbf{NT}	1	0	DF	No	Partial monotonicity	
0	0	\mathbf{NT}	0	1	CO	Yes		SW
1	1	AT	0	0	\mathbf{NT}	Yes		AT-NT
1	0	DF	1	0	DF	No	Partial monotonicity	
1	0	DF	0	1	CO	No	Partial monotonicity	
0	1	CO	1	0	DF	No	Partial monotonicity/Nested IV	
0	1	CO	0	1	CO	Yes		ACO
0	0	\mathbf{NT}	1	1	AT	Yes		NT-AT
1	1	AT	1	0	DF	No	Partial monotonicity	
1	1	AT	0	1	CO	Yes		SW
1	0	DF	1	1	AT	No	Partial monotonicity	
0	1	CO	1	1	AT	No	Nested IV	
1	1	AT	1	1	AT	Yes		AAT

Table 1: Latent populations and reasons for their inclusion/exclusion under Assumption 2(ii) (partial monotonicity) and Assumption 2(iv) (nested IV). NT: never-taker. AT: always-taker. CO: complier. DF: defier. ANT: always-never-taker. SW: switcher. AT-NT: always-taker-never-taker. ACO: always-complier. NT-AT: never-taker-always-taker. AAT: always-taker.

and the conditional always-complier average treatment effect:

$$\operatorname{ACOATE}_{P_0}(\boldsymbol{X}) := \mathbb{E}_{P_0}[Y(D=1) - Y(D=0) \mid S = \operatorname{ACO}, \boldsymbol{X}].$$

Importantly, both $\text{SWATE}_{P_0}(\boldsymbol{X})$ and $\text{ACOATE}_{P_0}(\boldsymbol{X})$ are the average "per-protocol" effect of D on Y, although on different subpopulations, within strata defined by observed covariates \boldsymbol{X} . By averaging $\text{SWATE}_{P_0}(\boldsymbol{X})$ and $\text{ACOATE}_{P_0}(\boldsymbol{X})$ over the target population P_0 , we arrive at the following causal estimands, the SWitcher Average Treatment Effect (SWATE):

$$SWATE_{P_0} := \mathbb{E}_{P_0}[Y(D=1) - Y(D=0) \mid S = SW],$$

and the Always-COmplier Average Treatment Effect (ACOATE):

$$ACOATE_{P_0} := \mathbb{E}_{P_0}[Y(D=1) - Y(D=0) | S = ACO].$$

Theorem 1 is our main identification result and shows that the ACOATE and SWATE can be non-parametrically identified from observed data under identification assumptions in Section 2.2. **Theorem 1.** Under Assumptions 1(i)-(iv) and 2(i)-(iv), $SWATE_{P_0}(\boldsymbol{x})$, $ACOATE_{P_0}(\boldsymbol{x})$, $SWATE_{P_0}(\boldsymbol{x})$, $SWATE_{P_$

$$ACOATE_{P_{0}}(\mathbf{X}) = \theta_{ACO,P_{0}}(\mathbf{X}) := \frac{\delta_{a,P_{0}}(\mathbf{X})}{\eta_{a,P_{0}}(\mathbf{X})}; \quad SWATE_{P_{0}}(\mathbf{X}) = \theta_{SW,P_{0}}(\mathbf{X}) := \frac{\delta_{b,P_{0}}(\mathbf{X}) - \delta_{a,P_{0}}(\mathbf{X})}{\eta_{b,P_{0}}(\mathbf{X}) - \eta_{a,P_{0}}(\mathbf{X})};$$

$$ACOATE_{P_{0}} = \Psi_{ACO,P_{0}} := \frac{\mathbb{E}_{P_{0}}[\delta_{a,P_{0}}(\mathbf{X})]}{\mathbb{E}_{P_{0}}[\eta_{a,P_{0}}(\mathbf{X})]}; \quad SWATE_{P_{0}} = \Psi_{SW,P_{0}} := \frac{\mathbb{E}_{P_{0}}[\delta_{b,P_{0}}(\mathbf{X}) - \delta_{a,P_{0}}(\mathbf{X})]}{\mathbb{E}_{P_{0}}[\eta_{b,P_{0}}(\mathbf{X}) - \eta_{a,P_{0}}(\mathbf{X})]},$$

where, for $P \in \mathcal{P}$,

$$\begin{split} \delta_{a,P}(\boldsymbol{x}) &:= \mathbb{E}_P[Y \mid Z = 1_a, \boldsymbol{X} = \boldsymbol{x}] - \mathbb{E}_P[Y \mid Z = 0_a, \boldsymbol{X} = \boldsymbol{x}];\\ \delta_{b,P}(\boldsymbol{x}) &:= \mathbb{E}_P[Y \mid Z = 1_b, \boldsymbol{X} = \boldsymbol{x}] - \mathbb{E}_P[Y \mid Z = 0_b, \boldsymbol{X} = \boldsymbol{x}];\\ \eta_{a,P}(\boldsymbol{x}) &:= \mathbb{E}_P[D \mid Z = 1_a, \boldsymbol{X} = \boldsymbol{x}] - \mathbb{E}_P[D \mid Z = 0_a, \boldsymbol{X} = \boldsymbol{x}];\\ \eta_{b,P}(\boldsymbol{x}) &:= \mathbb{E}_P[D \mid Z = 1_b, \boldsymbol{X} = \boldsymbol{x}] - \mathbb{E}_P[D \mid Z = 0_b, \boldsymbol{X} = \boldsymbol{x}]. \end{split}$$

2.4 Practical relevance of ACOATE and SWATE: a tool and a goal

Causal estimands ACOATE and SWATE are examples of the so-called principal treatment effects (Frangakis and Rubin, 2002). As discussed in Section 1.1, the practical relevance of any principal stratum effects is always under close scrutiny and often with good reasons (Pearl, 2011; Dawid and Didelez, 2012). Responding to Pearl's (2011) critiques, in our opinion, SWATE and ACOATE are both a useful "tool" in a holistic instrumental variable analysis and a "goal" in itself. First, it is a "tool" towards establishing the generalizability of a causal effect. For instance, researchers can immediately get a crude assessment of the generalizability of the treatment effect by comparing ACOATE to SWATE; we will develop a formal testing procedure for the principal ignorability assumption in Section 4. Moreover, the ACOATE and SWATE are potentially more informative than the standard CATE, by unravelling additional effect heterogeneity between subgroups. Although a study participant cannot be classified into one of the latent subgroups, the distribution of observed covariates among always-compliers and switchers can be characterized. Proposition 1 gives an identification formula for a function of an observed covariate among always-compliers and switchers.

Proposition 1. Let $g(\cdot)$ denote a generic function. The mean of $g(\mathbf{X})$ among switchers and

always-compliers can be identified as follows:

$$\mathbb{E}_{P_0}[g(\boldsymbol{X}) \mid S = SW] = \mathbb{E}_{P_0}\left[g(\boldsymbol{X}) \frac{\eta_{b,P_0}(\boldsymbol{X}) - \eta_{a,P_0}(\boldsymbol{X})}{\mathbb{E}_{P_0}[\eta_{b,P}(\boldsymbol{X}) - \eta_{a,P_0}(\boldsymbol{X})]}\right],$$
$$\mathbb{E}_{P_0}[g(\boldsymbol{X}) \mid S = ACO] = \mathbb{E}_{P_0}\left[g(\boldsymbol{X}) \frac{\eta_{a,P_0}(\boldsymbol{X})}{\mathbb{E}_{P_0}[\eta_{a,P_0}(\boldsymbol{X})]}\right].$$

Based on Proposition 1, the mean and variance of each observed covariate, and more generally certain combinations of observed covariates (e.g., a projected risk score), in the always-complier and switcher subgroups can be characterized and compared to those in the entire study population. For instance, for a binary covariate X, a "prevalence ratio" (Brookhart and Schneeweiss, 2007) may be calculated as P(X = 1 | S = ACO)/P(X = 1) and P(X = 1 | S = SW)/P(X = 1).

3 Estimation and inference

Estimation of the always-complier average treatment effect follows immediately by applying the estimation and inferential methods in the standard IV literature to the binary IV pair $(0_a, 1_a)$; see, e.g., Frölich (2007) and Kennedy (2022). In this section, we will focus on the estimation and inference of switcher average treatment effect.

3.1 Wald-type estimator

When the IV assignment is randomized in each stratum (G = a or b), as in a randomized clinical trial, a classical Wald estimator for ACOATE is $\widehat{\psi}_{ACO}^{Wald} = \frac{\widehat{\delta}_a}{\widehat{\eta}_a}$, where

$$\widehat{\delta}_{a} = \frac{1}{\sum_{i} \mathbb{1}\{Z_{i} = 1_{a}\}} \sum_{Z_{i} = 1_{a}} Y_{i} - \frac{1}{\sum_{i} \mathbb{1}\{Z_{i} = 0_{a}\}} \sum_{Z_{i} = 0_{a}} Y_{i},$$
$$\widehat{\eta}_{a} = \frac{1}{\sum_{i} \mathbb{1}\{Z_{i} = 1_{a}\}} \sum_{Z_{i} = 1_{a}} D_{i} - \frac{1}{\sum_{i} \mathbb{1}\{Z_{i} = 0_{a}\}} \sum_{Z_{i} = 0_{a}} D_{i}.$$

An analogous Wald-type estimator for SWATE can be formulated as follows:

$$\widehat{\psi}_{\mathrm{SW}}^{\mathrm{Wald}} = rac{\widehat{\delta}_b - \widehat{\delta}_a}{\widehat{\eta}_b - \widehat{\eta}_a},$$

where

$$\hat{\delta}_{b} = \frac{1}{\sum_{i} \mathbb{1}\{Z_{i} = 1_{b}\}} \sum_{Z_{i} = 1_{b}} Y_{i} - \frac{1}{\sum_{i} \mathbb{1}\{Z_{i} = 0_{b}\}} \sum_{Z_{i} = 0_{b}} Y_{i},$$
$$\hat{\eta}_{b} = \frac{1}{\sum_{i} \mathbb{1}\{Z_{i} = 1_{b}\}} \sum_{Z_{i} = 1_{b}} D_{i} - \frac{1}{\sum_{i} \mathbb{1}\{Z_{i} = 0_{b}\}} \sum_{Z_{i} = 0_{b}} D_{i}.$$

Theorem 2 summarizes the property of $\widehat{\psi}^{\text{Wald}}_{\text{SW}}.$

Theorem 2. Under regularity conditions specified in Supplemental Material S1.2, we have

$$\sqrt{n}\left(\widehat{\psi}_{SW}^{Wald} - \Psi_{SW,P_0}\right) \rightsquigarrow \mathcal{N}\left(0,\sigma_{Wald}^2\right),$$

where $\sigma_{Wald}^2 = \mathbb{E}_{P_0}[\varphi_{Wald,SW}^2]$, and $\varphi_{Wald,SW}$ is the influence function of $\widehat{\psi}_{SW}^{Wald}$, whose form can be found in Supplemental Material S1.2.

An estimator $\hat{\sigma}_{Wald}^2$ of σ_{Wald}^2 can be obtained using non-parametric bootstrap or a sandwich estimator.

3.2 Non-parametric efficiency bound

We now derive the non-parametric efficiency bound for the target identified parameter Ψ_{SW,P_0} . For a pathwise differentiable parameter, its non-parametric efficiency bound equals the variance of the canonical gradient at P_0 in a non-parametric model. Theorem 3 derives the canonical gradient and hence the non-parametric efficiency bound for the identification functional $\Psi_{SW,P}$.

Theorem 3. The canonical gradient of

$$\Psi_{SW,P} = \frac{\mathbb{E}_{P}[\delta_{b,P}(\boldsymbol{X}) - \delta_{a,P}(\boldsymbol{X})]}{\mathbb{E}_{P}[\eta_{b,P}(\boldsymbol{X}) - \eta_{a,P}(\boldsymbol{X})]}$$

at P_0 is D_{P_0} , where

$$\begin{split} D_{P} &= (z, x, d, y) \mapsto \frac{1}{\mathbb{E}_{P}[\eta_{b,P}(\mathbf{X}) - \eta_{a,P}(\mathbf{X})]} \Biggl\{ \frac{\mathbb{I}\{z = 1_{b}\}}{P(Z = 1_{b} \mid \mathbf{X} = \mathbf{x})} \Bigl[y - \mathbb{E}_{P}[Y \mid Z = 1_{b}, \mathbf{X} = \mathbf{x}] \Bigr] \\ &- \frac{\mathbb{I}\{z = 0_{b}\}}{P(Z = 0_{b} \mid \mathbf{X} = \mathbf{x})} \Bigl[y - \mathbb{E}_{P}[Y \mid Z = 0_{b}, \mathbf{X} = \mathbf{x}] \Bigr] \\ &- \frac{\mathbb{I}\{z = 1_{a}\}}{P(Z = 1_{a} \mid \mathbf{X} = \mathbf{x})} \Bigl[y - \mathbb{E}_{P}[Y \mid Z = 1_{a}, \mathbf{X} = \mathbf{x}] \Bigr] + \frac{\mathbb{I}\{z = 0_{a}\}}{P(Z = 0_{a} \mid \mathbf{X} = \mathbf{x})} \Bigl[y - \mathbb{E}_{P}[Y \mid Z = 0_{a}, \mathbf{X} = \mathbf{x}] \Bigr] \\ &+ \delta_{b_{P}}(\mathbf{x}) - \delta_{a,P}(\mathbf{x}) \Biggr\} - \frac{\Psi_{SW,P}}{\mathbb{E}_{P}[\eta_{b,P}(\mathbf{X}) - \eta_{a,P}(\mathbf{X})]} \Biggl\{ \frac{\mathbb{I}\{z = 1_{b}\}}{P(Z = 1_{b} \mid \mathbf{X} = \mathbf{x})} \Bigl[d - \mathbb{E}_{P}(D \mid Z = 1_{b}, \mathbf{X} = \mathbf{x}) \Bigr] \\ &- \frac{\mathbb{I}\{z = 0_{b}\}}{P(Z = 0_{b} \mid \mathbf{X} = \mathbf{x})} \Bigl[d - \mathbb{E}_{P}(Y \mid Z = 0_{b}, \mathbf{X} = \mathbf{x}) \Bigr] - \frac{\mathbb{I}\{Z = 1_{a}\}}{P(Z = 1_{a} \mid \mathbf{X} = \mathbf{x})} \Bigl[d - \mathbb{E}_{P}(D \mid Z = 1_{a}, \mathbf{X} = \mathbf{x}) \Bigr] \\ &+ \frac{\mathbb{I}\{Z = 0_{a}\}}{P(Z = 0_{a} \mid \mathbf{X} = \mathbf{x})} \Bigl[d - \mathbb{E}_{P}(D \mid Z = 0_{a}, \mathbf{X} = \mathbf{x}) \Bigr] + \eta_{b,P}(\mathbf{x}) - \eta_{a,P}(\mathbf{x}) \Biggr\}. \end{split}$$

Therefore, the non-parametric efficiency bound is $\mathbb{E}_{P_0}[D_{P_0}^2]$.

3.3 Non-parametric efficient estimators

The efficient influence function in Theorem 3 motivates an asymptotically efficient, one-step estimator and an estimating-equation-based estimator. A one-step estimator is based on first order bias correction of a plug-in estimator (Pfanzagl, 1990); an estimating-equation-based estimator is based on solving a efficient-influence-function-based estimating equation (van der Laan and Robins, 2003; Chernozhukov et al., 2018a).

Consider a K-fold random partition $\{I_k\}_{k=1}^K$ of indices $[N] = \{1, ..., K\}$, and define $I_k^c := \{1, ..., n\} - I_k$. For simplicity, we assume I_k 's have the same cardinality. For each $k \in [K]$, we construct estimators $\hat{\pi}_{n,k}$, $\hat{\mu}_{D,n,k}$, and $\hat{\mu}_{Y,n,k}$ of $\pi_{P_0} : (z, x) \mapsto P_0(Z = z \mid \mathbf{X} = \mathbf{x}), \mu_{Y,P_0} : (z, x) \mapsto \mathbb{E}_{P_0}[Y \mid Z = z, \mathbf{X} = \mathbf{x}]$, and $\mu_{D,P_0} : (z, x) \mapsto \mathbb{E}_{P_0}[D \mid Z = z, \mathbf{X} = \mathbf{x}]$, respectively, each based on sample $\{O_i\}_{i\in I_k^c}$. Let $\mathbb{P}_{n,k}$ be the empirical distribution of $\{O_i\}_{i\in I_k}$, and we use $\mathbb{P}_{n,k}\{\cdot\}$ to denote averaging with respect to the empirical measure $\mathbb{P}_{n,k}$. Let $\hat{P}_{n,k}$ be any distribution such that $f(Z \mid \mathbf{X})$ is compatible with $\hat{\pi}_{n,k}$, $f(Y \mid Z, \mathbf{X})$ is compatible with $\hat{\mu}_{Y,n,k}$, $f(D \mid Z, \mathbf{X})$ is compatible with $\hat{\mu}_{n,k}$, and the marginal distribution of $\hat{\mu}_{D,n,k}(z, \mathbf{X}_i)$ is given by its empirical distribution for $z \in \{0_a, 1_a, 0_b, 1_b\}$ in the k-th fold. A one-step estimator $\hat{\psi}_{os}$ can be constructed as follows:

Step I: For each $k \in [K]$, construct the one-step estimator $\widehat{\psi}_{os,k} = \Psi_{\widehat{P}_{n,k}} + \mathbb{P}_{n,k}D_{\widehat{P}_{n,k}}$ for Ψ_{SW,P_0} ; Step II: Construct the final cross-fitted one-step estimator as $\widehat{\psi}_{os} = \frac{1}{K} \sum_{k=1}^{n} \widehat{\psi}_{os,k}$. Alternatively, one may construct an estimator from an estimating equation perspective. Denote $D'(\Psi_{SW,P_0}, \pi_{P_0}, \mu_{Y,P_0}, \mu_{D,P_0}) := D_{P_0}$. Observe that $\mathbb{E}_{P_0}[D'(\Psi_{P_0}, \pi_{P_0}, \mu_{Y,P_0}, \mu_{D,P_0})] = 0$. An estimating-equation-based estimator $\hat{\psi}_{ee}$ can be constructed as follows:

Step I: For each $k \in [K]$, solve the following estimating equation $\mathbb{P}_{n,k} \left\{ D'(\Psi_{P_0}, \widehat{\pi}_{n,k}, \widehat{\mu}_{Y,n,k}, \widehat{\mu}_{D,n,k}) \right\} = 0$. Denote the solution as $\widehat{\psi}_{ee,k}$.

Step II: Construct the final cross-fitted one-step estimator as $\hat{\psi}_{ee} = \frac{1}{K} \sum_{k=1}^{n} \hat{\psi}_{ee,k}$.

Theorem 4 establishes the property of $\widehat{\psi}_{os}$ and $\widehat{\psi}_{ee}$.

Theorem 4. Under regularity conditions specified in Supplemental Material S1.4, we have

$$\sqrt{n}\left(\widehat{\psi} - \Psi_{SW,P_0}\right) = \sqrt{n}\mathbb{P}_n D_{P_0} + o_p(1)$$

for $\hat{\psi} = \hat{\psi}_{os}$ or $\hat{\psi} = \hat{\psi}_{ee}$. That is, the one-step estimator and the estimating-equation-based estimator are asymptotically linear and normal. Furthermore, they are asymptotically equivalent with asymptotic variance achieving the non-parametric efficient bound.

If we further assume $P_0 D_{P_0}^2 < \infty$, then by Central Limit Theorem, we have:

$$\sqrt{n}\left(\widehat{\psi}-\Psi_{\mathrm{SW},P_0}\right)\rightsquigarrow\mathcal{N}\left(0,P_0D_{P_0}^2\right).$$

A sandwich estimator can be used to estimate the asymptotic variance.

Remark 2 (Rates of convergence). For the one-step estimator to be asymptotic linear, regularity conditions require the convergence rates for all nuisance functions to be $o_p(n^{-1/4})$. On the other hand, the estimating-equation-based estimator is rate doubly robust (Rotnitzky et al., 2020), that is, $\hat{\psi}_{ee}$ is asymptotically liner with influence function D_{P_0} when

$$\left\|\pi_{P_0}(z,\cdot) - \pi_{\widehat{P}_{n,k}}(z,\cdot)\right\|_{L^2(P_0)} \left\|\mu_{B,P_0}(z,\cdot) - \mu_{B,\widehat{P}_{n,k}}(z,\cdot)\right\|_{L^2(P_0)} = o_p(1/\sqrt{n}),$$

for $z \in \{0_a, 1_a, 0_b, 1_b\}$, $k \in \{1, ..., K\}$, and B = Y or D.

4 Testable implications of homogeneity-type assumptions in a nested IV design

4.1 Generalizing principal stratum effects

Let $\operatorname{ATE}_{P_0}(\boldsymbol{x}) = \mathbb{E}_{P_0}[Y(D=1) - Y(D=0) \mid \boldsymbol{X} = \boldsymbol{x}]$ denote the conditional average treatment effect among those with covariates $\boldsymbol{X} = \boldsymbol{x}$ and $\operatorname{ATE}_{P_0} = \mathbb{E}_{P_0}[\operatorname{ATE}_{P_0}(\boldsymbol{X})]$. Proposition 2 states two sufficient conditions that identify $\operatorname{ATE}_{P_0}(\boldsymbol{x})$ (and hence $\operatorname{ATE}_{P_0})$ from principal stratum effect $\operatorname{ACOATE}_{P_0}(\boldsymbol{x})$ or $\operatorname{SWATE}_{P_0}(\boldsymbol{x})$.

Proposition 2. Suppose one of the following assumptions holds:

H(i). (Principal ignorability) $\mathbb{E}_{P_0}[Y(D=d) \mid S=s, \mathbf{X}] = \mathbb{E}_{P_0}[Y(D=d) \mid \mathbf{X}]$ for $d \in \{0, 1\}$.

H(ii). (No unmeasured common effect modifier)

$$Cov_{P_0} \Big[\Big(D(Z = 1_a) - D(Z = 0_a) \Big) \Big(Y(D = 1) - Y(D = 0) \Big) \mid \mathbf{X} \Big] = 0,$$

$$Cov_{P_0} \Big[\Big(D(Z = 1_b) - D(Z = 0_b) \Big) \Big(Y(D = 1) - Y(D = 0) \Big) \mid \mathbf{X} \Big] = 0.$$

Then we have $ATE_{P_0}(\boldsymbol{x}) = ACOATE_{P_0}(\boldsymbol{x}) = SWATE_{P_0}(\boldsymbol{x}).$

Remark 3 (One IV setting). Both Assumption H(i) and Assumption H(ii) in Proposition 2 have their counterparts in the usual setting with one IV; see, e.g., Jo and Stuart (2009) and Wang and Tchetgen Tchetgen (2018). With one IV, either assumption implies that the conditional complier average treatment effect with respect to the IV equals the conditional average treatment effect.

Define $\text{COATE}_{P_0}(\mathbf{X}) = \mathbb{E}_{P_0}[Y(D=1) - Y(D=0)|S \in \{\text{SW}, \text{ACO}\}]$, which is the conditional average treatment effect among compliers under IV pair $\{0_b, 1_b\}$. Then under Assumptions 1 and 2, we have $\text{COATE}_{P_0}(\mathbf{X}) = \theta_{\text{CO}, P_0}(\mathbf{X}) := \frac{\delta_{b, P_0}(\mathbf{X})}{\eta_{b, P_0}(\mathbf{X})}$. Observe that

$$\theta_{\mathrm{CO},P_0}(\boldsymbol{X}) \coloneqq \frac{\delta_{b,P_0}(\boldsymbol{X})}{\eta_{b,P_0}(\boldsymbol{X})} = \left\{ \frac{\eta_{b,P_0}(\boldsymbol{X}) - \eta_{a,P_0}(\boldsymbol{X})}{\eta_{b,P_0}(\boldsymbol{X})} \right\} \times \theta_{\mathrm{ACO},P_0}(\boldsymbol{X}) + \left\{ \frac{\eta_{a,P_0}(\boldsymbol{X})}{\eta_{b,P_0}(\boldsymbol{X})} \right\} \times \theta_{\mathrm{SW},P_0}(\boldsymbol{X}).$$

Under the nested IV setting, Proposition 2 implies that Assumption H(i) or H(ii) has the following testable implications: $\theta_{P_0}^{(1)} \equiv \theta_{ACO,P_0}(\mathbf{X}) - \theta_{SW,P_0}(\mathbf{X}) = 0$ a.s. $P_0, \theta_{P_0}^{(2)} \equiv \theta_{ACO,P_0}(\mathbf{X}) - \theta_{CO,P_0}(\mathbf{X}) = 0$ a.s. P_0 , and $\theta_{P_0}^{(3)} \equiv \theta_{SW,P_0}(\mathbf{X}) - \theta_{CO,P_0}(\mathbf{X}) = 0$ a.s. P_0 . In words, under either assumption, we

would expect that, within strata of X, the average treatment effect among always-compliers equals that among the switchers and that among the compliers under the IV pair $\{0_b, 1_b\}$. Next, we propose formal statistical procedures testing the null hypothesis:

$$H_0^{(j)}: \theta_{P_0}^{(j)} = 0 \text{ a.s. } P_0 \text{ vs } H_1^{(j)}: \theta_{P_0}^{(j)} \neq 0, \qquad j = 1, 2, \text{ and } 3.$$
(1)

4.2 Testing the best least squares projection

Consider testing a generic null hypothesis H_0 : $\xi_{P_0} = 0$ a.s. P_0 versus H_1 : $\xi_{P_0} \neq 0$. Here $\xi_{P_0} \in L^2(P_0)$ is a function-valued parameter and it is a function of \mathbf{X} . Let $\Gamma = \{\gamma(\mathbf{x};\beta), \beta \in B\}$ denote a class of functions indexed by β , with $\gamma(\mathbf{x};0) = 0$ for all β and B a subset of some linear space. Assume β is well-identified in the sense that $\beta_1 \neq \beta_2$ implies $\gamma(\cdot;\beta_1) \neq \gamma(\cdot;\beta_2)$. Furthermore, suppose Γ is a closed subspace $L^2(P_0)$. Then the best least squares projection of ξ_{P_0} onto Γ , denoted as $\Pi(\xi_{P_0} \mid \Gamma)$, can be defined as

$$\Pi(\xi_{P_0} \mid \Gamma) = \arg\min_{\gamma \in \Gamma} \|\gamma - \xi_{P_0}\|_{L^2(P_0)}$$

The projection $\Pi(\xi_{P_0} \mid \Gamma)$ has the unique representation as $\gamma(\cdot, \beta_{P_0})$, with β_{P_0} satisfying

$$\beta_0 = \arg\min_{\beta \in B} \mathbb{E}_{P_0} \left[\xi_{P_0}(\boldsymbol{X}) - \gamma(\boldsymbol{X}; \beta_0) \right]^2.$$
(2)

Under the null hypothesis $H_0: \xi_{P_0} = 0$ a.s. P_0 , we necessarily have $\beta_0 = 0$. Therefore, we can test the null hypothesis $H_0: \xi_{P_0} = 0$ a.s. P_0 by first deriving an asymptotically normal estimator and then constructing a Wald-type test for β_0 . Let us now specialize the discussion to the functional parameters $\theta_{P_0}^{(j)}$ and consider testing $H_{P_0}^{(j)}$ in (1). Let $\beta_{P_0}^{(j)}$ denote the projection of $\theta_{P_0}^{(j)}$ as defined in (2). Theorem 5 derives the canonical gradient for $\beta_P^{(j)}$, j = 1, 2, 3.

Theorem 5. Suppose $\partial \gamma(\boldsymbol{x}; \beta) / \partial \beta$ exists and is continuous in β . Under a non-parametric model,

the canonical gradient for $\beta_P^{(j)}$, j = 1, 2, 3 at P_0 , are $\phi_{P_0}^{(j)}$, j = 1, 2, 3. $\phi_P^{(j)}$, j = 1, 2, 3 are defined as

$$\begin{split} \phi_P^{(1)} &: (z, \boldsymbol{x}, d, y) \mapsto C_P^{(1)} \frac{\partial \gamma(\boldsymbol{x}; \beta)}{\partial \beta} \Big|_{\beta = \beta_P} \left[D_P^{(1)}(z, \boldsymbol{x}, d, y) - D_P^{(2)}(z, \boldsymbol{x}, d, y) + \theta_P^{(1)}(\boldsymbol{x}) - \gamma(\boldsymbol{x}; \beta_P) \right], \\ \phi_P^{(2)} &: (z, \boldsymbol{x}, d, y) \mapsto C_P^{(2)} \frac{\partial \gamma(\boldsymbol{x}; \beta)}{\partial \beta} \Big|_{\beta = \beta_P} \left[D_P^{(1)}(z, \boldsymbol{x}, d, y) - D_P^{(3)}(z, \boldsymbol{x}, d, y) + \theta_P^{(2)}(\boldsymbol{x}) - \gamma(\boldsymbol{x}; \beta_P) \right], \\ \phi_P^{(3)} &: (z, \boldsymbol{x}, d, y) \mapsto C_P^{(3)} \frac{\partial \gamma(\boldsymbol{x}; \beta)}{\partial \beta} \Big|_{\beta = \beta_P} \left[D_P^{(2)}(z, \boldsymbol{x}, d, y) - D_P^{(3)}(z, \boldsymbol{x}, d, y) + \theta_P^{(3)}(\boldsymbol{x}) - \gamma(\boldsymbol{x}; \beta_P) \right]. \end{split}$$

where C_P^j are constants, and

$$\begin{split} D_P^{(1)} &= (z, x, d, y) \mapsto \frac{1}{\eta_{a,P}(x)} \Biggl\{ \frac{1\{z = 1_a\}}{P(Z = 1_a \mid \mathbf{X} = x)} \Big[y - \mathbb{E}_P[Y \mid Z = 1_a, \mathbf{X} = x] \Big] \\ &- \frac{1\{z = 0_a\}}{P(Z = 0_a \mid \mathbf{X} = x)} \Big[y - \mathbb{E}_P[Y \mid Z = 0_a, \mathbf{X} = x] \Big] \Biggr\} \\ &- \frac{\delta_{a,P}(x)}{[\eta_{a,P}(x)]^2} \Biggl\{ \frac{1\{Z = 1_a\}}{P(Z = 1_a \mid \mathbf{X} = x)} \Big[d - \mathbb{E}_P[D \mid Z = 1_a, \mathbf{X} = x] \Big] \\ &- \frac{1\{Z = 0_a\}}{P(Z = 0_a \mid \mathbf{X} = x)} \Big[d - \mathbb{E}_P[D \mid Z = 0_a, \mathbf{X} = x] \Big] \Biggr\}, \\ D_P^{(2)} &= (z, x, d, y) \mapsto \frac{1}{\eta_{b,P}(x) - \eta_{a,P}(x)} \Biggl\{ \frac{1\{z = 1_b\}}{P(Z = 1_b \mid \mathbf{X} = x)} \Big[y - \mathbb{E}_P[Y \mid Z = 1_b, x] \Big] \\ &- \frac{1\{z = 0_b\}}{P(Z = 0_b \mid \mathbf{X} = x)} \Big[y - \mathbb{E}_P[Y \mid Z = 0_b, \mathbf{X} = x] \Big] \\ &- \frac{1\{z = 1_a\}}{P(Z = 1_a \mid \mathbf{X} = x)} \Big[y - \mathbb{E}_P[Y \mid Z = 1_a, \mathbf{X} = x] \Big] + \frac{1\{z = 0_a\}}{P(Z = 0_a \mid \mathbf{X} = x)} \Big[y - \mathbb{E}_P[Y \mid Z = 0_a, \mathbf{X} = x] \Big] \Biggr\} \\ &- \frac{\delta_{b,P}(x) - \delta_{a,P}(x)}{[\eta_{b,P}(x) - \eta_{a,P}(x)]^2} \Biggl\{ \frac{1\{z = 1_b\}}{P(Z = 1_b \mid \mathbf{X} = x)} \Big[d - \mathbb{E}_P[D \mid Z = 1_b, \mathbf{X} = x] \Big] \\ &- \frac{1\{z = 0_a\}}{P(Z = 0_b \mid \mathbf{X} = x)} \Big[d - \mathbb{E}_P[Y \mid Z = 0_b, \mathbf{X} = x] \Big] - \frac{1\{Z = 1_a\}}{P(Z = 1_a \mid \mathbf{X} = x)} \Big[d - \mathbb{E}_P[D \mid Z = 1_b, \mathbf{X} = x] \Big] \\ &+ \frac{1\{Z = 0_a\}}{P(Z = 0_a \mid \mathbf{X} = x)} \Big[d - \mathbb{E}_P[D \mid Z = 0_a, \mathbf{X} = x] \Big] \Biggr\}, \end{aligned}$$

$$-\frac{1\{Z=0_b\}}{P(Z=0_b \mid \boldsymbol{X}=\boldsymbol{x})} \Big[d - \mathbb{E}_P[D \mid Z=0_b, \boldsymbol{X}=\boldsymbol{x}] \Big] \bigg\}.$$

When $\gamma(\boldsymbol{x};\beta) = \boldsymbol{x}^T \beta$, we have $C_P^{(1)} = C_P^{(2)} = C_P^{(3)} = \mathbb{E}_P[\boldsymbol{X}\boldsymbol{X}^T]^{-1}$.

Analogous to the development in Section 3.3, we estimate $\beta_{P_0}^{(j)}$ based on one-step estimation with nuisance functions estimated via flexible non-parametric methods and cross-fitting.

Below, we focus on $\gamma(\boldsymbol{x};\beta) = \boldsymbol{x}^T \beta$. In this case, $\beta_P^{(j)}$ admits the following equivalent form:

$$\beta_P^{(j)} = (\mathbb{E}_P[\boldsymbol{X}\boldsymbol{X}^T])^{-1}\mathbb{E}_P[\boldsymbol{X}\boldsymbol{\theta}_P^j(\boldsymbol{X})].$$

For each $\beta_P^{(j)}$, our proposed test statistics can then be constructed analogously as described in Section 3.3 based on the following steps:

Step I: For each $k \in [K]$, construct the one-step estimator $\widehat{\beta}_{os,k}^{(j)} = \beta_{\widehat{P}}^{(j)} + \mathbb{P}_{n,k}\phi_{\widehat{P}_{n,k}}^{(j)}$ for $\beta_{P_0}^{(j)}$. Step II: Construct the final cross-fitted one-step estimator $\widehat{\beta}_{os}^{(j)} = \frac{1}{K} \sum_{k=1}^{n} \widehat{\beta}_{os,k}^{(j)}$. Step III: Estimate the asymptotic variance $\widehat{\Sigma}^{(j)} = \frac{1}{n} \sum_{k=1}^{K} n_k \mathbb{P}_{n,k} \left\{ \phi^{(j)} \widehat{P}_{n,k} \phi_{\widehat{P}_{n,k}}^{(j)} \right\};$

Step IV: Construct the test statistic $W^{(j)} = \widehat{\beta}_{os}^{(j),T} \widehat{\Sigma}^{(j),-1} \widehat{\beta}_{os}^{(j)}$, and define the following test:

$$T_{\text{Wald},\alpha}^{(j)} = \mathbb{1}\left\{W^{(j)} > \chi_{1-\alpha}^2(d_{\boldsymbol{x}})\right\},\,$$

where $d_{\boldsymbol{x}}$ is the dimension of \boldsymbol{x} , and $\chi^2_{1-\alpha}(d_{\boldsymbol{x}})$ is the $1-\alpha$ quantile for Chi-squared distribution with degree of freedom $d_{\boldsymbol{x}}$.

Theorem 6 establishes the property of the proposed estimator for $\beta_{P_0}^{(j)}$ and the test $T_{\text{Wald},\alpha}^{(j)}$.

Theorem 6. Under regularity conditions specified in Supplemental Material S1.7, for j = 1, 2, 3,

1. Estimators $\widehat{\beta}_{os}^{(j)}$ is both consistent and asymptotically normal, with its asymptotic variance achieving the efficiency bound:

$$\sqrt{n}\left(\widehat{\beta}^{(j)} - \beta_{P_0}\right) \rightsquigarrow \mathcal{N}\left(0, \ \mathbb{E}_{P_0}\left[\phi_{P_0}^{(j)T}\phi_{P_0}^{(j)}\right]
ight).$$

2. $T_{Wald,\alpha}^{(j)}$ is an asymptotically size- α test.

Remark 4. The idea of estimating the best least square projection for a function-valued parameter has been discussed in causal inference literature (Chernozhukov et al., 2018b). In our paper, we focus on testing rather than estimation. Under the null hypothesis, the function class Γ can be correctly specified, so that asymptotic type one error can be controlled. We don't need to assume Γ to be correctly specified under the alternative, however, the specification of Γ may affect the asymptotic power of the test.

Remark 5. Similar to the estimation of $SWATE_{P_0}$, one may also construct estimating-equationbased estimator $\beta_P^{(j)}$ by treating $\phi_P^{(j)}$ in Theorem 5 as an estimating function for $\beta_P^{(j)}$. When $\gamma(\mathbf{X}; \beta)$ is linear in \mathbf{X} , one-step estimators of $\beta_{P_0}^{(j)}$ are equivalent to estimating-equation-based estimators.

4.3 Non-parametric Kolmogorov-Smirnov-type test

In this section, we describe a non-parametric Kolmogorov-Smirnov type test motivated by the work of Westling (2022). Again we consider testing a null hypothesis $H_0: \xi_{P_0} = 0$ a.s. P_0 versus $H_1: \xi_{P_0} \neq 0$, where $\xi_{P_0} \in L^2(P_0)$ is a function of \mathbf{X} . Suppose $\mathbf{X} = (\mathbf{X}_1, \mathbf{X}_2)$, where the marginal distribution of \mathbf{X}_1 is absolutely continuous to the counting measure, and the marginal distribution of \mathbf{X}_2 is absolutely continuous to the Lebesgue measure. Let the support of $\mathbf{X} = (\mathbf{X}_1, \mathbf{X}_2)$ be $\mathcal{X} = \mathcal{X}_1 \times \mathcal{X}_2 \subset \mathbb{R}^{d_1} \times \mathbb{R}^{d_2}$. Let $\mathbf{1}_c(x) = \mathbf{1}\{x \in (-\infty, c_1] \times ... \times (-\infty, c_{d_1}] \times ... \times (-\infty, c_{d_1+d_2}]\}$. The construction of the Kolmogorov-Smirnov-type tests is based on the following technical lemma.

Lemma 1. $\theta_{P_0}(x) = 0$ for all $x \in \mathcal{X}_1 \times \mathcal{X}_2$ if and only if $\Omega_{P_0}(c) := \int \xi_{P_0}(x) \mathbb{1}_c(x) dP_0(x) = 0$ for all $c \in \mathcal{X}$ (that is, $\sup_{c \in \mathcal{X}} |\Omega_{P_0}(c)| = 0$).

Therefore, testing the null hypothesis $\xi_{P_0}(x) = 0$ a.s. P_0 is equivalent to testing the null hypothesis that $\Omega_{P_0}(c) = 0$ for any $c \in \mathcal{X}$. One advantage of constructing a test based on $\Omega_{P_0}(c)$ is that $\Omega_{P_0}(c)$ is \sqrt{n} estimable for a fixed c without further parametric assumptions. On the contrary, even when all the nuisance parameters are known, $\theta_{P_0}(x)$ is not \sqrt{n} estimable without further parametric assumptions because the minimax rate is slower than $1/\sqrt{n}$ (Tsybakov, 2008).

The first step of our testing procedure is to construct a RAL estimator of $\Omega_{P_0}(c)$ for each fixed c, denoted as $\widehat{\Omega}(c)$. Similar to the estimation procedure developed in Section 4.2, we can construct a cross-fitted, one-step estimator based on the efficient influence function of $\Omega_P(c)$ at P_0 for any fixed c. Suppose $\widehat{\Omega}_{os}$ admits the following asymptotic expansion:

$$\widehat{\Omega}(c) - \Omega_{P_0}(c) = \mathbb{P}_n D_{P_0}^*(c) + r_n(c)$$

where $D_{P_0}^*(c)$ is the canonical gradient of $\Omega_P(c)$ at P_0 . If we further have $\sup_{c \in \mathcal{X}} |r_n(c)| = o_p(n^{-1/2})$ and $\{D_{P_0}^*(c), c \in \mathcal{X}\}$ is P_0 -Donsker, then the stochastic process $\mathbb{H}_n := \{\sqrt{n}[\widehat{\Omega}(c) - \Omega_{P_0}(c)] : c \in \mathcal{X}\}$ converges weakly in $l^{\infty}(\mathcal{X})$ to a tight, mean-zero Gaussian process \mathbb{H} , with covariance function $\Sigma(s,t) = \mathbb{E}_{P_0}[D_{P_0}^*(s)D_{P_0}^*(t)]$, where $(s,t) \in \mathcal{X} \times \mathcal{X}$. Then by the continuous mapping theorem, $\sup_{c \in \mathcal{X}} |\sqrt{n}\{\widehat{\Omega}_{os}(c) - \Omega_{P_0}(c)\}| \rightsquigarrow ||\mathbb{H}||_{\infty}$. In practice, one may replace $\sup_{c \in \mathcal{X}} |\sqrt{n}\{\widehat{\Omega}(c) - \widehat{\Omega}_{P_0}(c)\}|$ with $\sup_{c \in \mathcal{X}_n} |\sqrt{n}\{\widehat{\Omega}(c) - \Omega_{P_0}(c)\}|$, where $\mathcal{X}_n = \{\mathbf{X}_1, \dots, \mathbf{X}_n\}$. To simulate the α -th quantile of $||\mathbb{H}||_{\infty}$, one may (i) simulate a mean-zero Gaussian process \mathbb{H}_n with covariance function equal to the sample covariance function $\Sigma_n(s,t) = \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left\{ D_{\widehat{P}_{n,k}}^*(s) D_{\widehat{P}_{n,k}}^*(t) \right\}$; (ii) simulate the $1 - \alpha$ -th quantile of $\sup_{c \in \mathcal{X}_n} |\mathbb{H}_n(c)|$, denoted as $Q_{n,\alpha}$; and (iii) construct the test by comparing $Q_{n,\alpha}$ to $\sup_{c \in \mathcal{X}_n} |\sqrt{n}\widehat{\Omega}(c)|$ since $\Omega_{P_0}(c)$ equals 0 under the null.

For j = 1, 2, 3, let ξ_{P_0} be $\theta_{P_0}^{(j)}$, and define $\Omega_{P_0}^{(j)}(c) := \int \theta_{P_0}^{(j)}(x) \mathbf{1}_c(x) dP_0(x)$. Proposition 3 derives the canonical gradient for $\Omega_P^{(j)}(c)$.

Proposition 3. The canonical gradient for $\Omega_P^{(j)}(c)$ at P_0 is $D_{P_0}^{(j)*}(c)$, where

$$\begin{split} D_P^{(1)*}(c) &: o \mapsto D_P^{(1)}(z, x, d, y) \mathbb{1}_c(x) - D_P^{(2)}(g, z, x, d, y) \mathbb{1}_c(x) + \theta_P^{(1)}(x) \mathbb{1}_c(x) - \Omega_P^{(1)}(c), \\ D_P^{(2)*}(c) &: o \mapsto D_P^{(1)}(z, x, d, y) \mathbb{1}_c(x) - D_P^{(3)}(g, z, x, d, y) \mathbb{1}_c(x) + \theta_P^{(2)}(x) \mathbb{1}_c(x) - \Omega_P^{(2)}(c), \\ D_P^{(3)*}(c) &: o \mapsto D_P^{(2)}(z, x, d, y) \mathbb{1}_c(x) - D_P^{(3)}(g, z, x, d, y) \mathbb{1}_c(x) + \theta_P^{(3)}(x) \mathbb{1}_c(x) - \Omega_P^{(3)}(c). \end{split}$$

For $j \in \{1, 2, 3\}$ and any fixed c, the proposed one-step estimator $\widehat{\Omega}_{os}^{(j)}(c)$ of $\Omega_{P_0}^{(j)}(c)$ can then be constructed as follows:

Step I: For each $k \in [K]$, construct the cross-fitted, one-step estimator $\widehat{\Omega}_{os,k}^{(j)}(c) = \Omega_{\widehat{P}_k}^{(j)}(c) + \mathbb{P}_{n,k}D_{\widehat{P}_k}^{(j)*}(c);$

Step II: Construct the final cross-fitted one-step estimator as $\widehat{\Omega}_{os}^{(j)}(c) = \frac{1}{K} \sum_{k=1}^{K} \widehat{\Omega}_{os,k}^{(j)}(c)$.

We summarize our proposed asymptotically size- α test below:

Step I: For each $k \in [K]$ and a fixed c, construct the cross-fitted, one-step estimator $\widehat{\Omega}_{os}^{(j)}(c)$.

- Step II: Define $\mathcal{X}_n = \{ \mathbf{X}_1, \dots, \mathbf{X}_n \}$. Let $Q_{n,\alpha}^{(j)}$ be the 1α quantile of $\max_{c \in \mathcal{X}_n} |\mathbb{H}_n^{(j)}(c)|$, where $(\mathbb{H}_n^{(j)}(\mathbf{X}_1), \dots, \mathbb{H}_n^{(j)}(\mathbf{X}_n))$ is distributed to a mean-zero multivariate normal distribution with covariance matrix given by $\Sigma_n^{(j)}(s,t) = \frac{1}{K} \sum_{k=1}^K \mathbb{P}_{n,k} \left\{ D_{\hat{P}_{n,k}}^{(j)*}(s) D_{\hat{P}_{n,k}}^{(j)*}(t) \right\}$ conditional on (O_1, \dots, O_n) .
- **Step III:** Define the test: $T_{np,\alpha}^{(j)} = \mathbb{1}\{n^{1/2}\sup_i |\widehat{\Omega}^{(j)}(\boldsymbol{X}_i)| > Q_{n,\alpha}^{(j)}\}$.

Theorem 7 summarizes the properties of the proposed test $T_{np,\alpha}^{(j)}$.

Theorem 7. Under regularity conditions specified in Supplemental Material A.1.10,

- 1. $T_{np,\alpha}^{(j)}$ is an asymptotically size α test.
- 2. The asymptotic power of $T_{np,\alpha}^{(j)}$ is 1.
- 3. $T_{np,\alpha}^{(j)}$ has non-trivial power under local alternative distributions P_n :

$$\lim_{n \to \infty} \int \left[n^{1/2} \left(dP_n^{1/2} - dP_0^{1/2} \right) - \frac{1}{2} h dP_0^{1/2} \right] = 0,$$

where h is a score function with $P_0h = 0$ and $P_0h^2 < \infty$.

Remark 6. Unlike $T_{Wald}^{(j)}$, the proposed non-parametric Kolmogorov-Smirnov-type tests are computationally challenging when the dimension of X is moderately large, because one needs to simulate a Gaussian process indexed by a subset of \mathbb{R}^{d_x} . Therefore, we recommend the tests based on the best least-square projection in practice.

5 Simulation study

We have two goals in this simulation study. Section 5.1 investigates the finite-sample performance of the proposed estimators for SWATE. Section 5.2 investigates the level of each linear-projectionbased test for the null hypotheses $H_0^{(j)}$. Because test $T_{\text{Wald},\alpha}^{(j)}$ developed for $H_0^{(j)}$ can each be used to test the same null hypothesis Assumption H(i) or H(ii), Section 5.2 also compares their power.

5.1 Estimation of SWATE

We generate datasets according to the following factorial design:

Sample sizes: N = 1000, 2000, 5000 and 10000.

Baseline covariates: We consider a eight-dimensional covariate $\boldsymbol{X} = (\boldsymbol{X}_1, \boldsymbol{X}_2, \dots, \boldsymbol{X}_8)$, where $(\boldsymbol{X}_1, \boldsymbol{X}_2, \boldsymbol{X}_3) \sim \mathcal{MVN}(\mu, \boldsymbol{\Sigma})$, with $\mu = (0, 1, -0.5)$, $\boldsymbol{\Sigma} = \begin{pmatrix} 1 & 0.2 & -0.3 \\ 0.2 & 1 & 0.1 \\ -0.3 & 0.1 & 1 \end{pmatrix}$, and truncated at -4 and 4, $\boldsymbol{X}_4, \boldsymbol{X}_5, \boldsymbol{X}_6 \sim \text{Bernoulli}(0.5), \boldsymbol{X}_7 \sim \text{Uniform}(-3, 3)$, and $\boldsymbol{X}_8 \sim \text{Binomial}(4, 0.5)$.

Instrumental variable: We generate a binary stratification variable $G \in \{a, b\}$ according to $P(G = a) = 1 - \exp(1 + 0.2X_1 - 0.1X_2 + 0.3X_3)$ and P(G = b) = 1 - P(G = a), where $\exp(x) = \exp(x)/(1 + \exp(x))$ is the inverse of the logit function. Conditioning on G = a or b, we then generate a binary IV $Z \in \{0_a, 1_a\}$ when G = a and $Z \in \{0_b, 1_b\}$ when G = b as follows:

$$P(Z = 1_a \mid G = a) = \exp\{1 + 0.5X_1 - X_2 + 0.7X_3\},\$$
$$P(Z = 1_b \mid G = b) = \exp\{0.5 + 0.6X_1 + 0.3X_2 + 0.4X_3\}.$$

According to this data generating process, the IV assignment is dependent on observed covariates and different for G = a or b.

Unmeasured confounder: We generate an unmeasured confounder $U \sim \mathcal{N}(0, 0.6)$. Scalar representation of unobserved confounding is widely adopted in the literature and referred to as a "principal unobserved covariate" (Rosenbaum and Rubin, 2023).

Principal stratum: The principal stratum status of participant *i*, denoted as S_i , is generated via a multinomial distribution: $P(S_i = j) = \frac{g_j}{\sum_j g_j}, j \in \{\text{ANT}, \text{SW1}, \text{SW2}, \text{AT-NT}, \text{ACO}, \text{NT-AT}, \text{AAT}\}$. The principal strata SW1 and SW2 correspond to participants who convert from always-takers or never-takers to compliers, respectively. We specify g_j as follows:

$$g_{ANT} = \exp\{1 - X_2 + 0.7X_3 + 0.3U\}, \qquad g_{SW1} = \exp\{\alpha_1 + \alpha_2(X_1 + 2X_2 - X_3) + \alpha_3U\},$$

$$g_{SW2} = \exp\{\alpha_1 + \alpha_2(-X_1 - 2X_2) + \alpha_3U\}, \qquad g_{AT-NT} = \exp\{1 + 0.5X_1 + X_2 + 0.5X_3 + 0.5U\},$$

$$g_{ACO} = \exp\{1 + 0.8X_1 - 2X_2 - 2X_3 + 0.5U\}, \qquad g_{NT-AT} = \exp\{1 - 0.5X_1 - 1X_2 - 0.5X_3 - 0.5U\}$$

$$g_{AAT} = \exp\{1 + 2X_1 + 2X_3 - U\},$$

where parameters $(\alpha_1, \alpha_2, \alpha_3)$ control the proportions of switchers. We consider the following 4 choices of $(\alpha_1, \alpha_2, \alpha_3)$: (-0.2, 0.1, 0.0005), (0.5, 0.2, 0.05), (0.3, 0.5, 0.1), and (1, 1, 1), corresponding to 11%, 22%, 32%, and 66% switchers in the population, respectively.

Treatment received: Treatment received D is determined by Z and S.

Outcome: We consider two sets of data generating processes for potential outcomes: For a continuous outcome, we generate $Y(0) = 1 + X_1 + X_2 + X_3 + X_4 + U + \epsilon$ and Y(1) as follows:

$$\begin{split} Y(1) = \mathbb{1} \{ S = \text{ANT or AAT} \} \cdot (1 + X_1 + 2X_2 + 2X_3 + X_4 + U) \\ &+ \mathbb{1} \{ S = \text{AT-NT or NT-AT} \} \cdot (1 + X_1 + X_2 + 2X_3 + X_4 + U) \\ &+ \mathbb{1} \{ S = \text{SWI} \} \cdot (\beta_1 + \beta_2 X_1 + 2X_2 + \beta_3 X_3 + X_4 + U) \\ &+ \mathbb{1} \{ S = \text{ACO} \} \cdot (1 + X_1 + 2X_2 + 0.2X_2^2 + X_3 + X_4 + U) + \epsilon, \end{split}$$

where $(\beta_1, \beta_2, \beta_3) = (2, 2, 2)$ or (4, 4, 4) control the effect size of SWATE, and $\epsilon \sim \mathcal{N}(0, 1)$. For a binary outcome, potential outcomes are generated as follows:

$$\begin{split} P(Y(0) = 1) &= \mathbbm{1}\{S = \text{ANT or AAT}\} \cdot \text{expit}\{1 + X_1 + 2X_2 + 2X_3 + X_4 + X_5 - X_6 - X_7 + X_8 + U\} \\ &+ \mathbbm{1}\{S = \text{AT-NT or NT-AT}\} \cdot \text{expit}\{1 + X_1 + X_2 + 2X_3 + X_4 + X_5 - X_6 - X_7 + X_8 + U\} \\ &+ \mathbbm{1}\{S = \text{SWI}\} \cdot \text{expit}\{\beta_1 + \beta_2 X_1 + 2X_2 + X_3 + \beta_3 X_6 - X_7 + 2X_8 + U\} \\ &+ \mathbbm{1}\{S = \text{ACO}\} \cdot \text{expit}\{1 + X_1 + 2X_2 + 0.2X_2^2 + X_3 + X_4 + X_5 - X_6 - X_7 + X_8 + U\}, \end{split}$$

where $(\beta_1, \beta_2, \beta_3) = (0, 1, -1)$ or (0, 2, -3) control the size of SWATE.

According to the above data generating processes, nuisance functions in the canonical gradient do not have explicit forms. For each simulated dataset, we constructed the proposed estimator $\hat{\psi}_{os}$ with all nuisance functions estimated using the R (R Core Team, 2013) package SuperLearner (Van der Laan et al., 2007). Specifically, we included the random forest and generalized linear models in the SuperLearner library. For each of the $2 \times 4 \times 3 \times 2 = 48$ settings determined by the outcome type, sample size, $(\alpha_1, \alpha_2, \alpha_3)$ and $(\beta_1, \beta_2, \beta_3)$, we repeated the simulation 1000 times.

Figure 1 plots the sampling distributions of estimating equation estimators for various combinations of sample size and switcher proportion when the outcome is continuous and $\beta = (2, 2, 2)$. The true values of SWATE are superimposed as red dash lines. Overall, the sampling distributions appeared normal except in settings where both the proportion of switchers and the sample size are small (e.g., n = 1000 with 11% switchers). For a fixed proportion of switchers, the sampling distributions are more concentrated around the true values as the sample size increases. Similarly, for a fixed sample size, the sampling distribution becomes more concentrated around the true values as the proportion of switchers increases. Table 2 further reports the bias, relative bias and coverage probability for different settings. We have observed come consistent trends. First, when the proportion of switchers is small, say 11%, then a large sample size (e.g., n = 10000) is needed for the proposed estimator to achieve a small relative bias and its 95% confidence intervals to achieve nominal coverage. This phenomenon is known as the weak IV bias (see, e.g., Bound et al., 1995; Andrews et al., 2019). On the other hand, when the proportion of switchers is relatively high (e.g., 66%), then the relative bias is small and the coverage attains approximately its nominal level for a moderately large sample size. For instance, when $\beta = (2, 2, 2)$, the relative bias of the proposed estimator is 2.1% and the coverage of the 95% CI is 94.1% when the sample size is as small as 1000. Table 2 also reports the proportion of switchers were small. For instance, when n = 2000 and the proportion of switchers is 32%, none of the 1000 simulated datasets returned an unstable estimator. On the other hand, the one-step estimator was a bit more unstable compared to the estimating-equation-based estimator; see Table S6 in Supplemental Material S2.

Overall, our simulation studies suggest that when the sample size times the proportion of switchers is greater than 500, the performance of our proposed estimator is favorable with small relative bias and close-to-nominal coverage. Simulation results for a binary outcome are qualitatively similar; see Supplemental Material S2 for details.




Figure 1: Sampling distribution of non-parametric estimator when the outcome is continuous. Simulations are repeated 1000 times. Datasets are generating with $\beta = (2, 2, 2)$. Four proportions of SW are considered. The dashed red lines represents the ground truth SWATE in each setting.

5.2 Testing implications of homogeneity

We generate datasets according to the following factorial design:

Sample sizes: N = 2000, 5000, and 10000.

Baseline covariates: We consider $X = (X_1, X_2)$ from a bivariate normal distribution with $\mu = (0, 0)$ and $\Sigma = I_2$.

Table 2: Simulation results for $\hat{\psi}_{ee}$ when the outcome is continuous. Column **SWATE** reports the true values of SWATE under each setting. Column **Estimate** reports the average point estimates (excluded if the absolute value exceeded 500). Column **SE Est** reports the 90% Winsorized mean of standard error estimates. Column **Acc. Rate** reports the proportion of point estimates not truncated.

β	SW%	SWATE	Sample Size	Estimate	Rel. Bias	Bias	SE Est	Coverage	Acc. Rate
			1000	-1.202	-0.610	-0.559	43.614	0.906	99.8%
			2000	0.765	-2.244	-2.060	10.774	0.923	99.9%
	11%	0.917	5000	1.052	0.147	0.135	1.018	0.941	100%
			10000	1.019	0.110	0.101	0.546	0.938	100%
		1.019	1000	0.799	-0.216	-0.220	4.852	0.909	100%
	22%		2000	1.130	0.109	0.111	0.966	0.949	100%
			5000	1.081	0.061	0.062	0.430	0.961	100%
			10000	1.053	0.033	0.034	0.273	0.954	100%
(2,2,2)		1.377	1000	1.838	0.334	0.461	1.259	0.942	100%
	~		2000	1.486	0.079	0.108	0.606	0.962	100%
	32%		5000	1.383	0.004	0.005	0.304	0.959	100%
			10000	1.378	0.000	0.000	0.202	0.961	100%
		1.557	1000	1.590	0.021	0.033	0.341	0.941	100%
	66%		2000	1.553	-0.003	-0.004	0.210	0.949	100%
			5000	1.560	0.002	0.003	0.122	0.946	100%
			10000	1.560	0.002	0.003	0.083	0.949	100%
	11%	0.906	1000	1.518	0.674	0.612	51.315	0.924	100%
			2000	1.273	3.698	3.354	9.969	0.937	99.9%
			5000	1.168	0.288	0.261	1.216	0.953	100%
			10000	1.111	0.225	0.204	0.607	0.931	100%
	22%	1.146	1000	1.258	0.097	0.111	5.562	0.921	100%
			2000	1.328	0.158	0.181	1.298	0.964	100%
			5000	1.213	0.058	0.066	0.546	0.968	100%
(4,4,4)			10000	1.152	0.004	0.005	0.340	0.953	100%
	32%	1.699	1000	1.814	0.068	0.115	1.721	0.944	100%
			2000	1.966	0.157	0.267	0.853	0.962	100%
			5000	1.741	0.024	0.041	0.425	0.969	100%
			10000	1.735	0.021	0.036	0.278	0.961	100%
		2.315	1000	2.336	0.009	0.020	0.574	0.953	100%
	66%		2000	2.336	0.009	0.021	0.350	0.964	100%
			5000	2.302	-0.006	-0.014	0.202	0.960	100%
			10000	2.306	-0.004	-0.010	0.138	0.973	100%

Instrumental variable: We generate a binary stratification variable $G \in \{a, b\}$ as follows:

$$P(G = a) = \exp\left\{0.1 \cdot \mathbb{1}\{X_1 > 0\} - 0.1 \cdot \mathbb{1}\{X_2 > 0\}\right\}.$$

We generate a binary IV for G = a or b as follows:

$$P(Z = 1_a \mid G = a) = 0.5 + 0.1 \cdot \mathbb{1}\{X_1 > 0\} - 0.1 \cdot \mathbb{1}\{X_1 > 0\},$$

$$P(Z = 1_b \mid G = b) = 0.5 - 0.1 \cdot \mathbb{1}\{X_1 > 0\} + 0.1 \cdot \mathbb{1}\{X_1 > 0\}.$$

Unmeasured confounder: We generate an unmeasured confounder $U \sim \mathcal{N}(-0.3, 0.3)$.

Principal stratum: The principal stratum status of participant i, denoted as S_i , is generated through a multinomial distribution. To this end, we define g_j as follows:

and let $P(S_i = j) = \frac{g_j}{\sum_j g_j}$ for $j \in \{ANT, AT-NT, NT-AT, AAT\}$, $P(S = SW1) = P(S = SW2) = \frac{(1-\alpha) \cdot (g_{SW1}+g_{SW2}+g_{ACO})}{2\sum_j g_j}$, and $P(S = ACO) = \frac{\alpha \cdot (g_{SW1}+g_{SW2}+g_{ACO})}{\sum_j g_j}$, where α controls the proportion of switchers. We let $\alpha = 0.1, 0.2, 0.4, 0.6, 0.8$, and 0.9, which correspond to 9%, 18%, 36%, 54%, 72%, and 81% switchers, respectively. Under this data generating process, the proportion of switchers and always-compliers, i.e., P(S = SW) + P(S = ACO), is fixed at 90%.

Treatment received: Treatment received D is determined by Z and S.


Outcome: We consider the following data generating process for potential outcomes:

$$\begin{split} Y(0) = &1 + X_1 + X_2 + X_3 + X_4 + U + \epsilon, \\ Y(1) = &\mathbb{1}\{S = \text{ANT or AAT}\} \cdot (1 + X_1 + X_2 + U) + &\mathbb{1}\{S = \text{AT-NT or NT-AT}\} \cdot (1 + X_1 + X_2 + U) \\ &+ &\mathbb{1}\{S = \text{SW}\} \cdot (\beta_1 + \beta_2 X_1 + \beta_3 X_2 + U) + &\mathbb{1}\{S = \text{ACO}\} \cdot (1 + 2X_1 + 2X_2 + U) + \epsilon, \end{split}$$

where parameters $(\beta_1, \beta_2, \beta_3)$ control the effect size of SWATE, and $\epsilon \sim N(0, 1)$. We consider $(\beta_1, \beta_2, \beta_3) = (1, 2, 2), (1, 2.5, 2.5), \text{ and } (2, 3, 3)$. When $\beta = (1, 2, 2), \text{ the null hypothesis } H_0^{(j)}$ holds, i.e., $\theta_{P_0}^{(j)} = 0$. Conversely, $\beta = (1, 2.5, 2.5)$ and (2, 3, 3) correspond to $\theta_{P_0}^{(j)} \neq 0$.

For each null hypothesis $H_0^{(j)}$, we constructed the best-linear-projection-based test $T_{\text{Wald},0.05}^{(j)}$ as described in Section 4.2 and evaluated the level and power of each test under different settings specified above based on 1000 simulated datasets.

Figure S.4 in the Supplemental Material S2 plots the nominal Type-I error rate versus the empirical rejection proportion. For $T^{(1)}_{\text{Wald},\alpha}$ that compares $\text{SWATE}_{P_0}(\boldsymbol{X})$ to $\text{ACOATE}_{P_0}(\boldsymbol{X})$, the rejection fraction aligns well with the nominal level when the proportion of switchers is not too high or too low. On the other hand, the approximation of the null distribution is not accurate when the proportion of switchers is small and hence $\text{SWATE}_{P_0}(X)$ is not accurately estimated, or when the proportion of always-compliers is small and hence $\text{ACOATE}_{P_0}(\mathbf{X})$ is not well estimated. Analogously, the null distribution of $T^{(2)}_{\text{Wald},\alpha}$ is best approximated when the proportion of switchers is small and that of $T^{(3)}_{\text{Wald},\alpha}$ is best approximated when the proportion of switchers is large. Importantly, we did not observe any inflation of type-I error for a level-0.05 test in all settings. Figure 2 further compares the power of each test under different settings. In all settings, when the proportion of switchers is small (e.g., 9% and 18%), $T_{\text{Wald},0.05}^{(2)}$ often outperforms $T_{\text{Wald},0.05}^{(1)}$ and $T_{\text{Wald},0.05}^{(3)}$, and $T_{\text{Wald},0.05}^{(3)}$ is the least powerful. When the proportion of switchers is high (e.g., 72% and 81%), $T_{\text{Wald},0.05}^{(3)}$ is the most powerful among three tests. Lastly, when the proportion of switchers is moderate (e.g., 36% and 54%), $T_{\text{Wald},0.05}^{(2)}$ outperforms the other two tests although the three tests have similar performance when the sample size is large. As the sample size increases, the power of all three tests increases. In empirical studies, we would recommend $T_{\text{Wald},0.05}^{(2)}$ when the proportion of always-compliers is high and $T_{Wald,0.05}^{(3)}$ when the proportion of switchers is high.



tests - test 1 - test 2 - test 3

Figure 2: Comparison of power. The red dashed lines correspond to the nominal level. Test 1 compares $\text{SWATE}_{P_0}(\boldsymbol{X})$ to $\text{ACOATE}_{P_0}(\boldsymbol{X})$; test 2 compares $\text{COATE}_{P_0}(\boldsymbol{X})$ to $\text{ACOATE}_{P_0}(\boldsymbol{X})$; test 3 compares $\text{COATE}_{P_0}(\boldsymbol{X})$ to $\text{SWATE}_{P_0}(\boldsymbol{X})$.

6 Revisiting the PLCO study

We applied our developed nested IV framework to re-analyzing the Prostate, Lung, Colorectal and Ovarian (PLCO) Cancer Screening Trial. We focused on the Henry Ford Health System, which underwent a major policy shift in its consent process and switched from dual to single consent in the middle of 1997. Our analysis cohort consisted of study participants who enrolled prior to 1997 (G = a) and after 1997 (G = b). We excluded those who enrolled during 1997. Table 3 summarizes important baseline characteristics and treatment uptake (attending first scheduled cancer screening or not) by stage and treatment assignment. Participants enrolled in the two stages differed moderately in their baseline characteristics; within each stage, participants enrolled in the treatment arm and control arm were comparable by virtue of randomization.

After the study switched to a single consent process, the compliance rate increased. The unadjusted compliance rate was estimated to be 80.1% after 1997 compared to 50.9% before 1997. One might suspect that the shift in the covariate distribution contributed to this marked difference in the compliance rate, so we estimated the covariate-adjusted compliance rate: the covariate-

	Prior to 1	1997 ($G = a$)	After 1997 $(G = b)$		
	Control arm	Treatment arm	Control arm	Treatment arm	
Samlpe size	4,210	4,204	4,970	4,978	
Treatment uptake (%)					
No screening	4210 (100.0)	2063 (49.1)	4970(100.0)	989(19.9)	
Screening	0 (0.0)	2141(50.9)	0 (0.0)	3989(80.1)	
Age	64.75(5.09)	64.69(5.08)	60.40(5.26)	60.42(5.31)	
Sex $(\%)$					
Female	2642(62.8)	2638 (62.7)	2810(56.5)	2798(56.2)	
Male	1568(37.2)	1566 (37.3)	2160(43.5)	2180(43.8)	
Race $(\%)$					
White	3346(79.5)	3289(78.2)	4266(85.8)	4278 (85.9)	
Minority	864 (20.5)	915(21.8)	704 (14.2)	700(14.1)	
Education (%)					
No high school	585(13.9)	616(14.7)	351(7.1)	369(7.4)	
High school	1775(42.2)	1639(39.0)	1586(31.9)	1589(31.9)	
College or above	1850(43.9)	1949 (46.4)	3033(61.0)	$3020 \ (60.7)$	
Smoking Status (%)					
Non-smoker	1803(42.8)	1798 (42.8)	2101 (42.3)	2076 (41.7)	
Current smoker	562(13.3)	$573 \ (13.6)$	700 (14.1)	729(14.6)	
Former smoker	1845 (43.8)	1833 (43.6)	2169(43.6)	2173 (43.7)	
BMI (%)					
< 25	1439(34.2)	1446(34.4)	1438(28.9)	1486 (29.9)	
> 25	2771 (65.8)	2758 (65.6)	3532 (71.1)	3492(70.1)	
Colorectal cancer (%)					
No Confirmed Cancer	4128 (98.1)	4139(98.5)	4905 (98.7)	4920 (98.8)	
Confirmed Cancer	82 (1.9)	65(1.5)	65(1.3)	58 (1.2)	
Following time	12.08(3.87)	11.45(4.47)	9.64(2.31)	9.55(2.47)	
Rate (per 1000 person-yrs)	1.61	1.35	1.36	1.22	

Table 3: Baseline characteristics by study stages and arms. Continuous variables are presented as mean (sd) and categorical variables are presented as count (proportion).

adjusted compliance rate was estimated to be 84.9% after 1997 compared to 52.6% before 1997. These results immediately imply that (1) the moderate shift in the baseline covariates between two stages *cannot* fully explain the significant improvement in the marginal compliance rate; and (2) 84.9% - 52.6% = 32.3% participants were estimated to be "switchers" under the nested IV framework. Who were these switchers that responded to the shift in the consent process? Table 4 summarizes the mean of each baseline covariate in the switcher and always-complier populations using Proposition 1. Interestingly, although always-compliers and switchers were similar in age and smoking status, switchers were more likely to be male, white, and having BMI < 25.

	Switchers	Always-compliers
Age	62.6	62.1
Sex $(\%)$		
Female	50.4%	71.6%
Male	49.6%	28.4%
Race $(\%)$		
White	88.0%	80.8%
Minority	12.0%	19.2%
Education $(\%)$		
No high school	9.3%	10.0%
High school	40.0%	33.1%
College or above	50.7%	56.9%
Smoking Status (%)		
Non-smoker	41.5%	42.8%
Current smoker	15.3%	12.1%
Former smoker	43.2%	45.1%
BMI (%)		
< 25	36.0%	28.7%
> 25	64.0%	71.3%

Table 4: Estimated baseline characteristics of switchers (SW) and always-compliers (ACO)

We then implemented the proposed one-step estimator and estimating-equation-based estimator for the switcher average treatment effect. Specifically, we fitted a logistic regression model for nuisance functions including $P(Z = z \mid G = g, \mathbf{X} = \mathbf{x})$, $P(G = g \mid \mathbf{X} = \mathbf{x})$, and $\mathbb{E}[D \mid Z = z, \mathbf{X} = \mathbf{x}]$, and a Poisson regression model for $\mathbb{E}[Y \mid Z = z, \mathbf{X} = \mathbf{x}]$ with the follow-up time being the offset. The 1000-person-year risk difference was estimated to be 0.083 (95% CI: -0.511 to 0.675) for switchers and -0.049 (95% CI: -0.207 to 0.110) for always-compliers based on the one-step estimator. The results obtained from the estimating-equation-based estimator were similar: -0.041 (95% CI: -0.633 to 0.552) for switchers and -0.021 (95% CI: -0.179 to 0.138) for always-compliers. On the other hand, the Wald estimators with bootstrapped confidence intervals were less accurate, with an estimated 1000-person-year risk difference of 0.614 (95% CI: -2.178 to 3.453) for switchers and -0.499 (95% CI: -1.410 to 0.422) for always-compliers.

Finally, we tested the null hypothesis that the effect of cancer screening on colorectal cancer was the same for all principal strata, within strata defined by 6 observed covariates including age, sex, race, education, smoking status, and BMI. The test statistic equaled 12.35, 5.80 and 12.45 for three best-projection-based tests $T_{\text{Wald},0.05}^{(1)}$, $T_{\text{Wald},0.05}^{(2)}$, and $T_{\text{Wald},0.05}^{(3)}$, respectively. The 95% quantile of a $\chi^2(9)$ distribution is 16.92; hence, results from all three tests did not provide evidence for additional effect heterogeneity among latent principal strata within strata defined by 6 observed covariates.

7 Discussion

In this work, we systematically study the problem with a pair of nested instrumental variables. We show that the switcher average treatment effect, or SWATE, can be non-parametrically identified under a novel nested IV assumption and develop an efficient-influence-function-based estimator for it. We also develop a projection-based test and a non-parametric test that formally assess the effect-homogeneity-type assumptions under our proposed nested-IV framework.

Our paper contributes to the discussion on violation of the SUTVA (Imbens and Rubin, 2015). In the PLCO trial and similar multi-center or multi-phasic trials, there may exist multiple versions of encouragement for treatment uptake, a subtlety that could be easily overlooked. In the presence of multiple versions of an IV, the usual intention-to-treat effect and the Wald estimand do not have their usual causal interpretations. The differentiation between two distinct versions of IV gives us an opportunity to study additional principal causal effects induced by the nested IV structure and test for effect heterogeneity. Interestingly, our identified conditional switcher average treatment has a similar form as the conditional ATE under the so-called instrumented difference-in-differences setting (Ye et al., 2023); however, the SWATE estimand proposed in this article is obtained by integrating over the covariate distribution among switchers, while Ye et al.'s (2023) marginalized estimand is obtained by integrating over the covariate distribution of the entire study population. Furthermore, in our setting, the abundance of principal strata complicates the model for treatment uptake and potential outcomes; hence, the current article focused on non-parametric estimators that permit using flexible non-parametric methods for nuisance functions estimation rather the usual doubly robust estimators that require correct specification for parametric nuisance models (Rotnitzky et al., 2020).

Our methods also provide some insight into testing the equality of two function-valued parameters (Hudson et al., 2021, 2023). We provide a general framework for such testing problems by projecting the function-valued parameter to a subspace of $L^2(P)$. In principle, one can project the parameters onto an infinite-dimensional space; in practice, projection onto a finite dimensional space suffices for type-I error control and appears to have good finite-sample power. Moreover, compared to a non-parametric test, e.g., our generalization of Westling (2022), a projection-based test is computationally more efficient and works smoothly when covariates are of different types.

Switcher average treatment effect is an example of the principal treatment effect. Principal effects are sometimes criticized for not being the most relevant for clinical decision making as they are concerned about the effects among unidentified subgroups; however, SWATE is clinically meaningful when researchers are interested in the practical utility of improving the compliance: a small SWATE effectively conveys the message that the marginal gain of improving compliance may be low for a target population, while a large SWATE could be an encouraging marching order for implementation scientists to investigate how to improve treatment adherence which has potential to translate to improved clinical outcomes.

In general, even a valid IV cannot point identify the marginal and conditional treatment effects without restricting heterogeneity in some sense. Since it was an aim of this work to investigate treatment effect heterogeneity with non-compliance, we did not impose these type of restrictions. It is an interesting question for future work whether alternative non-principal interpretations of the ACOATE and SWATE functionals exist, without imposing homogeneity.

References

Andrews, I., Stock, J. H., and Sun, L. (2019). Weak instruments in instrumental variables regression: Theory and practice. Annual Review of Economics, 11(1):727–753.

Angrist, J. D., Imbens, G. W., and Rubin, D. B. (1996). Identification of causal effects using

instrumental variables. Journal of the American Statistical Association, 91(434):444–455.

- Baiocchi, M., Cheng, J., and Small, D. S. (2014). Instrumental variable methods for causal inference. Statistics in Medicine, 33(13):2297–2340.
- Bound, J., Jaeger, D. A., and Baker, R. M. (1995). Problems with instrumental variables estimation when the correlation between the instruments and the endogenous explanatory variable is weak. *Journal of the American Statistical Association*, 90(430):443–450.
- Brookhart, M. A. and Schneeweiss, S. (2007). Preference-based instrumental variable methods for the estimation of treatment effects: assessing validity and interpreting results. *The international journal of biostatistics*, 3(1).
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., and Newey, W. (2018a). Double/debiased/neyman machine learning of treatment effects. *The Econometrics Journal*, 21(1):C1–C68.
- Chernozhukov, V., Demirer, M., Duflo, E., and Fernandez-Val, I. (2018b). Generic machine learning inference on heterogeneous treatment effects in randomized experiments, with an application to immunization in india. Technical report, National Bureau of Economic Research.
- Cohen, M. S. and Baden, L. R. (2012). Preexposure prophylaxis for HIV—where do we go from here? New England Journal of Medicine, 367(5):459–461.
- Cook, T. D., Campbell, D. T., and Day, A. (1979). *Quasi-experimentation: Design & Analysis Issues for Field Settings*, volume 351. Houghton Mifflin Boston.
- Dahabreh, I. J., Robertson, S. E., Tchetgen, E. J., Stuart, E. A., and Hernán, M. A. (2019). Generalizing causal inferences from individuals in randomized trials to all trial-eligible individuals. *Biometrics*, 75(2):685–694.
- Dawid, P. and Didelez, V. (2012). "imagine a can opener"-the magic of principal stratum analysis. The international journal of biostatistics, 8(1).
- Deaton, A. S. (2009). Instruments of development: Randomization in the tropics, and the search

for the elusive keys to economic development. Technical report, National Bureau of Economic Research.

- Frangakis, C. E. and Rubin, D. B. (2002). Principal stratification in causal inference. *Biometrics*, 58(1):21–29.
- Frölich, M. (2007). Nonparametric iv estimation of local average treatment effects with covariates. Journal of Econometrics, 139(1):35–75. Endogeneity, instruments and identification.
- Gilbert, P. B. and Hudgens, M. G. (2008). Evaluating candidate principal surrogate endpoints. Biometrics, 64(4):1146–1154.
- Goldberger, A. S. (1972). Structural equation methods in the social sciences. *Econometrica: Journal* of the Econometric Society, pages 979–1001.
- He, Q., Gao, F., Dukes, O., Delany-Moretlwe, S., and Zhang, B. (2023). Generalizing the intentionto-treat effect of an active control against placebo from historical placebo-controlled trials to an active-controlled trial: A case study of the efficacy of daily oral TDF/FTC in the HPTN 084 study.
- Heckman, J. J. and Urzua, S. (2010). Comparing iv with structural models: What simple iv can and cannot identify. *Journal of Econometrics*, 156(1):27–37.
- Hernán, M. A. and Robins, J. M. (2006). Instruments for causal inference: an epidemiologist's dream? *Epidemiology*, pages 360–372.
- Hudson, A., Carone, M., and Shojaie, A. (2021). Inference on function-valued parameters using a restricted score test.
- Hudson, A., Geng, E. H., Odeny, T. A., Bukusi, E. A., Petersen, M. L., and van der Laan, M. J. (2023). An approach to nonparametric inference on the causal dose response function.
- Imbens, G. W. and Angrist, J. D. (1994). Identification and estimation of local average treatment effects. *Econometrica*, 62(2):467–475.
- Imbens, G. W. and Rubin, D. B. (2015). Causal Inference for Statistics, Social, and Biomedical Sciences: An Introduction. Cambridge University Press.

- Jo, B. and Stuart, E. A. (2009). On the use of propensity scores in principal causal effect estimation. Statistics in Medicine, 28(23):2857–2875.
- Joffe, M. (2011). Principal stratification and attribution prohibition: good ideas taken too far. *The International Journal of Biostatistics*, 7(1).
- Kennedy, E. H. (2022). Semiparametric doubly robust targeted double machine learning: a review.
- Lambdin, B., Cheng, B., Peter, T., Mbwambo, J., Apollo, T., Dunbar, M., C Udoh, I., Cattamanchi, A., H Geng, E., and Volberding, P. (2015). Implementing implementation science: an approach for hiv prevention, care and treatment programs. *Current HIV Research*, 13(3):244–246.
- Levis, A. W., Kennedy, E. H., and Keele, L. (2024). Nonparametric identification and efficient estimation of causal effects with instrumental variables.
- Marcus, P. M., Ogden, S. L., Gren, L. H., Childs, J. C., Pretzel, S. M., Lamerato, L. E., Walsh, K., Rozjabek, H. M., Mabie, J., Thomas, B., et al. (2014). Non-compliance with the initial screening exam visit in the Prostate, Lung, Colorectal and Ovarian (PLCO) Cancer Screening Trial. *Preventive medicine*, 67:82–88.
- Neyman, J. S. (1923). On the application of probability theory to agricultural experiments. Essay on principles. Section 9. Annals of Agricultural Sciences, 10:1–51.
- Pearl, J. (2011). Principal stratification-a goal or a tool? The International Journal of Biostatistics, 7(1):1–13.
- Pfanzagl, J. (1990). Estimation in semiparametric models, pages 17–22. Springer US, New York, NY.
- R Core Team (2013). R: A language and environment for statistical computing.
- Robins, J. M. (1994). Correcting for non-compliance in randomized trials using structural nested mean models. *Communications in Statistics-Theory and Methods*, 23(8):2379–2412.
- Rosenbaum, P. and Rubin, D. (2023). Propensity scores in the design of observational studies for causal effects. *Biometrika*, 110(1):1–13.

- Rotnitzky, A., Smucler, E., and Robins, J. M. (2020). Characterization of parameters with a mixed bias property. *Biometrika*, 108(1):231–238.
- Rubin, D. (1980). Discussion of "Randomization analysis of experimental data in the Fisher randomization test" by D. Basu. Journal of the American Statistical Association, 75:591–593.
- Rubin, D. B. (1974). Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of Educational Psychology*, 66(5):688.
- Sun, Z. and Wüthrich, K. (2022). Pairwise valid instruments.
- Swanson, S. A., Hernán, M. A., Miller, M., Robins, J. M., and Richardson, T. S. (2018). Partial identification of the average treatment effect using instrumental variables: review of methods for binary instruments, treatments, and outcomes. *Journal of the American Statistical Association*, 113(522):933–947.
- Tchetgen Tchetgen, E. J. and Vansteelandt, S. (2013). Alternative identification and inference for the effect of treatment on the treated with an instrumental variable.
- Thistlethwaite, D. L. and Campbell, D. T. (1960). Regression-discontinuity analysis: An alternative to the expost facto experiment. *Journal of Educational psychology*, 51(6):309.
- Tsybakov, A. (2008). Introduction to Nonparametric Estimation. Springer Series in Statistics. Springer New York.
- Van der Laan, M. J., Polley, E. C., and Hubbard, A. E. (2007). Super learner. Statistical Applications in Genetics and Molecular Biology, 6(1).
- van der Laan, M. J. and Robins, J. M. (2003). Unified Approach for Causal Inference and Censored Data, pages 311–370. Springer New York, New York, NY.
- van der Vaart, A. and Wellner, J. (2023). Weak Convergence and Empirical Processes: With Applications to Statistics. Springer International Publishing.
- Van der Vaart, A. W. (2000). Asymptotic statistics, volume 3. Cambridge university press.

- Wang, L. and Tchetgen Tchetgen, E. (2018). Bounded, efficient and multiply robust estimation of average treatment effects using instrumental variables. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 80(3):531–550.
- Westling, T. (2022). Nonparametric tests of the causal null with nondiscrete exposures. *Journal of the American Statistical Association*, 117(539):1551–1562.
- Ye, T., Ertefaie, A., Flory, J., Hennessy, S., and Small, D. S. (2023). Instrumented difference-indifferences. *Biometrics*, 79(2):569–581.
- Zhang, J. L. and Rubin, D. B. (2003). Estimation of causal effects via principal stratification when some outcomes are truncated by "death". *Journal of Educational and Behavioral Statistics*, 28(4):353–368.

Supporting Information to "Nested Instrumental Variables Design: Switcher Average Treatment Effect, Identification, Efficient Estimation and

Generalizability"

S1 Proofs

S1.1 Proof of Theorem 1

Proof. For SWATE, Since we have

$$\begin{split} & \mathbb{E}_{P_0}[Y|Z = 1_b, X] - \mathbb{E}_{P_0}[Y|Z = 0_b, X] \\ &= \mathbb{E}_{P_0}[Y(Z = 1_b)|Z = 1_b, G = b, X] - \mathbb{E}_{P_0}[Y(Z = 0_b)|Z = b_0, G = b, X] \\ &= \mathbb{E}_{P_0}[Y(Z = 1_b) - Y(Z = 0_b)|G = b, X] \\ &= \mathbb{E}_{P_0}[Y(Z = 1_b) - Y(Z = 0_b)|X] \\ &= \mathbb{E}_{P_0}[Y(D = 1) - Y(D = 0)|S = SW, X]P_0(S = SW|X) \\ &+ \mathbb{E}_{P_0}[Y(D = 1) - Y(D = 0)|S = ACO, X]P_0(S = ACO|X) \end{split}$$

and

$$\mathbb{E}_{P_0}[Y|Z = 1_a, X] - \mathbb{E}_{P_0}[Y|Z = 0_a, X]$$

= $\mathbb{E}_{P_0}[Y(Z = 1_a)|Z = 1_a, G = a, X = x] - \mathbb{E}_{P_0}[Y(Z = 0_a)|Z = 0_a, G = a, X = x]$
= $\mathbb{E}_{P_0}[Y(Z = 1_a) - Y(Z = 0_a)|G = a, X = x]$
= $\mathbb{E}_{P_0}[Y(Z = 1_a) - Y(Z = 0_a)|X = x]$
= $\mathbb{E}_{P_0}[Y(D = 1) - Y(D = 0)|S = ACO, X]P_0(S = ACO|X)$

therefore

$$\mathbb{E}_{P_0}[Y(D=1) - Y(D=0)|S = SWI, X]P_0(S = SWI|X)$$

=(\mathbb{E}_{P_0}[Y|Z = 1_b, X] - \mathbb{E}_{P_0}[Y|Z = 0_b, X]) - (\mathbb{E}_{P_0}[Y|Z = 1_a, X] - \mathbb{E}_{P_0}[Y|Z = 0_a, X])

Similarly, we have

$$\mathbb{E}_{P_0}[D|Z = 1_b, X] - \mathbb{E}_{P_0}[D|Z = 0_b, X]$$

=\mathbb{E}_{P_0}[D|Z = 1_b, S = b, X] - \mathbb{E}_{P_0}[D|Z = 0_b, G = b, X]
=\mathbb{E}_{P_0}[D(Z = 1_b) - D(Z = 0_b)|G = b, X]
=\mathbb{E}_{P_0}[D(Z = 1_b) - D(Z = 0_b)|X]
=\mathbb{P}_0(ACO|X) + P_0(SW|X)

and

$$\mathbb{E}_{P_0}[D|Z = a_1, X] - E[D|Z = a_0, X]$$

= $\mathbb{E}_{P_0}[D|Z = 1_b, G = a, X] - \mathbb{E}_{P_0}[D|Z = 0_b, G = a, X]$
= $\mathbb{E}_{P_0}[D(Z = 1_a) - D(Z = 0_a)|G = a, X]$
= $\mathbb{E}_{P_0}[D(Z = 1_a) - D(Z = 0_a)|X]$
= $P_0(ACO|X)$

Therefore we have

$$(\mathbb{E}_{P_0}[D|Z=1_b, X] - \mathbb{E}_{P_0}[D|Z=0_b, X]) - (\mathbb{E}_{P_0}[D|Z=1_a, X] - \mathbb{E}_{P_0}[D|Z=0_a, X]) = P_0(ACO|X)$$

Therefore,

$$SWATE_{P_0}(X) := \mathbb{E}_{P_0}[Y(D=1) - Y(D=0)|S = SW, X]$$

=
$$\frac{(E[Y|Z=1_b, X] - \mathbb{E}_{P_0}[Y|Z=0_b, X]) - (\mathbb{E}_{P_0}[Y|Z=1_a, X] - \mathbb{E}_{P_0}[Y|Z=0_a, X])}{(\mathbb{E}_{P_0}[D|Z=1_b, X] - \mathbb{E}_{P_0}[D|Z=0_b, X]) - (\mathbb{E}_{P_0}[D|Z=1_a, X] - \mathbb{E}_{P_0}[D|Z=0_a, X])}$$

and

$$SWATE_{P_0} = \int SWATE_{P_0}(x)f_{P_0}(x|S = SW)d\mu(x)$$

= $\int \frac{\delta_{b,P_0}(x) - \delta_{a,P_0}(x)}{\eta_{b,P_0}(x) - \eta_{a,P_0}(x)} \frac{\eta_{b,P_0}(x) - \eta_{a,P_0}(x)}{P_0(S = SW)}f_{P_0}(x)d\mu(x)$
= $\frac{\mathbb{E}_{P_0}[\delta_{b,P_0}(X) - \delta_{a,P_0}(X)]}{P_0(S = SW)}$
= $\frac{\mathbb{E}_{P_0}[\delta_{b,P_0}(X) - \delta_{a,P_0}(X)]}{\mathbb{E}_{P_0}[\eta_{b,P_0}(X) - \eta_{a,P_0}(X)]}$

where $f_{P_0}(x|S = SW)$ and $f_{P_0}(x)$ are conditional density and density function implied by P_0 . Similarly, for ACOATE, we have

$$ACOATE_{P_0}(X) := \mathbb{E}_{P_0}[Y(D=1) - Y(D=0)|S = ACO, X]$$
$$= \frac{\mathbb{E}_{P_0}[Y|Z = 1_a, X] - \mathbb{E}_{P_0}[Y|Z = 0_a, X]}{\mathbb{E}_{P_0}[D|Z = 1_a, X] - \mathbb{E}_{P_0}[D|Z = 0_a, X]}$$

and

$$ACOATE_{P_0} = \frac{\mathbb{E}_{P_0}[\mu_{a,P_0}(X)]}{\mathbb{E}_{P_0}[\eta_{a,P_0}(X)]}$$

S1.2 Proof of Theorem 2

Proof. Let

$$\hat{\mu}_{Y,m} = \frac{1}{n} \sum_{i=1}^{n} Y_i \mathbb{1}\{Z_i = m\}, \quad \mu_{Y,m} = \mathbb{E}_{P_0}[Y_i \mathbb{1}\{Z_i = s\}]$$
$$\hat{\mu}_{D,m} = \frac{1}{n} \sum_{i=1}^{n} D_i \mathbb{1}\{Z_i = m\}, \quad \mu_{D,m} = \mathbb{E}_{P_0}[D_i \mathbb{1}\{Z_i = m\}]$$
$$\hat{\mu}_{Z,m} = \frac{1}{n} \sum_{i=1}^{n} \mathbb{1}\{Z_i = m\}, \quad \mu_{Z,m} = \mathbb{E}_{P_0}[\mathbb{1}\{Z_i = m\}]$$

for $s \in \{a_1, a_0, b_1, b_0\}$. Then it's clear that $\hat{\mu}_{Y,s}, \hat{\mu}_{D,s}, \hat{\mu}_{Z,s}, s \in \{a_1, a_0, b_1, b_0\}$ are asymptotic linear estimators for $\mu_{Y,s}, \mu_{D,s}, \mu_{Z,s}, s \in \{a_1, a_0, b_1, b_0\}$, with influence functions

$$\begin{split} \varphi_{Y,m} &= o \mapsto y \mathbb{1}\{z = m\} - \mu_{Y,m} \\ \varphi_{D,m} &= o \mapsto d\mathbb{1}\{z = m\} - \mu_{D,m} \\ \varphi_{Z,m} &= o \mapsto \mathbb{1}\{z = m\} - \mu_{Z,m} \end{split}$$

Let

$$f(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10}, x_{11}, x_{12}) = \frac{\frac{x_1}{x_2} - \frac{x_3}{x_4} - \frac{x_5}{x_6} + \frac{x_7}{x_8}}{\frac{x_9}{x_2} - \frac{x_{10}}{x_4} - \frac{x_{11}}{x_6} + \frac{x_{12}}{x_8}}$$

Then

$$\hat{\psi}_{SW}^{Wald} = f(\hat{\mu}_{Y,1_b}, \hat{\mu}_{Z,1_b}, \hat{\mu}_{Y,0_b}, \hat{\mu}_{Z,0_b}, \hat{\mu}_{Y,1_a}, \hat{\mu}_{Z,1_a}, \hat{\mu}_{Y,0_a}, \hat{\mu}_{Z,0_a}, \hat{\mu}_{D,1_b}, \hat{\mu}_{D,0_b}, \hat{\mu}_{D,1_a}, \hat{\mu}_{D,0_a})$$

$$\Psi_{SW,P_0} = f(\mu_{Y,1_b}, \mu_{Z,1_b}, \mu_{Y,0_b}, \mu_{Z,0_b}, \mu_{Y,1_a}, \mu_{Z,1_a}, \mu_{Y,0_a}, \mu_{Z,0_a}, \mu_{D,1_b}, \mu_{D,0_b}, \mu_{D,1_a}, \mu_{D,0_a})$$

By delta method (Van der Vaart, 2000), we know $\hat{\psi}_{SW}^{Wald}$ is asymptotically linear estimator of Ψ_{SW,P_0} with influence function

$$\varphi_{Wald,SW}: o \mapsto [\nabla f(\mu_{Y,D,Z})]^T \varphi_{Y,D,Z}(o)$$

where

$$\mu_{Y,D,Z} = (\mu_{Y,1_b}, \mu_{Z,1_b}, \mu_{Y,0_b}, \mu_{Z,0_b}, \mu_{Y,1_a}, \mu_{Z,1_a}, \mu_{Y,0_a}, \mu_{Z,0_a}, \mu_{D,1_b}, \mu_{D,0_b}, \mu_{D,1_a}, \mu_{D,0_a})^T,$$

$$\varphi_{Y,D,Z}(o) = (\varphi_{Y,1_b}, \varphi_{Z,1_b}, \varphi_{Y,0_b}, \varphi_{Z,0_b}, \varphi_{Y,1_a}, \varphi_{Z,1_a}, \varphi_{Y,0_a}, \varphi_{Z,0_a}, \varphi_{D,1_b}, \varphi_{D,0_b}, \varphi_{D,1_a}, \varphi_{D,0_a})^T.$$

Therefore,

$$\sqrt{n} \left(\hat{\psi}_{\mathrm{SW}}^{\mathrm{Wald}} - \Psi_{\mathrm{SW}, P_0} \right) \rightsquigarrow N \left(0, \sigma_{\mathrm{Wald}}^2 \right),$$

where $\sigma_{\text{Wald}}^2 = P_0 \varphi_{Wald,SW}^2$.

S1.3 Proof of Theorem 3

Proof. Let $\{P_t, t \in [0, \delta)\}$ denote a one-dimensional regular submodel of \mathcal{P}_{np} , with $P_{t=0} = P_0$. Then the canonical gradient D(P) at P_0 is the unique function which belongs to $L_0^2(P_0)$ that satisfies

$$\frac{d\Psi(P_t)}{dt}\Big|_{t=0} = P_0(D(P_0)g),$$
(3)

where g is the score function of submodel $\{P_t, t \in [0, \delta)\}$ at t = 0. We observe that

$$\begin{aligned} & \frac{d\Psi(P_t)}{dt}\Big|_{t=0} \\ &= \frac{\frac{d}{dt}\Big|_{t=0}\mathbb{E}_{P_t}[\mathbb{E}_{P_t}[Y|Z=b_1,X] - \mathbb{E}_{P_t}[Y|Z=b_0,X] - \mathbb{E}_{P_t}[Y|Z=a_1,X] + \mathbb{E}_{P_t}[Y|Z=b_0,X]]}{\mathbb{E}_{P_0}[\eta_{b,P_0}(X) - \eta_{a,P_0}(X)]} \\ & + \Psi(P_0)\frac{\frac{d}{dt}\Big|_{t=0}\mathbb{E}_{P_t}[\mathbb{E}_{P_t}[D|Z=b_1,X] - \mathbb{E}_{P_t}[D|Z=b_0,X] - \mathbb{E}_{P_t}[D|Z=a_1,X] + \mathbb{E}_{P_t}[D|Z=b_0,X]]}{\mathbb{E}_{P_0}[\eta_{b,P_0}(X) - \eta_{a,P_0}(X)]} \end{aligned}$$

It's well-known that the canonical gradients for $\mathbb{E}_{P_0}[\mathbb{E}_{P_0}[Y|Z = z, X]]$ and $\mathbb{E}_{P_0}[\mathbb{E}_{P_0}[D|Z = z, X]]$ are

$$T_1(P_0) = E_{P_0}[Y|Z = z, X] + \frac{\mathbb{1}(Z = z)}{P_0(Z = z|X)} \left[Y - \mathbb{E}_{P_0}[Y|Z = z, X] \right] - \mathbb{E}_{P_0} \left[\mathbb{E}_{P_0}[Y|Z = z, X] \right]$$

and

$$T_2(P_0) = \mathbb{E}_{P_0}[D|Z = z, X] + \frac{\mathbb{1}(Z = z)}{P_0(Z = z|X)} \left[D - \mathbb{E}_{P_0}[D|Z = z, X] \right] - \mathbb{E}_{P_0}\left[\mathbb{E}_{P_0}[D|Z = z, X] \right]$$

respectively, therefore

$$\frac{d}{dt} \mathbb{E}_{P_t} \left[\mathbb{E}_{P_t} [Y|Z=z, X] \right]_{t=0} = P_0(T_1(P_0)g)$$
$$\frac{d}{dt} \mathbb{E}_{P_t} \left[\mathbb{E}_{P_t} [Y|Z=z, X] \right]_{t=0} = P_0(T_2(P_0)g)$$

therefore

$$D_{P} = (z, x, d, y) \mapsto \frac{1}{\mathbb{E}_{P}[\eta_{b}(\mathbf{X}) - \eta_{a}(\mathbf{X})]} \left\{ \frac{\mathbb{I}\{z = 1_{b}\}}{P(Z = 1_{b} \mid \mathbf{X} = \mathbf{x})} \left[y - \mathbb{E}_{P}[Y \mid Z = 1_{b}, \mathbf{x}] \right] \\ - \frac{\mathbb{I}\{z = 0_{b}\}}{P(Z = 0_{b} \mid \mathbf{X} = \mathbf{x})} \left[y - \mathbb{E}_{P}[Y \mid Z = 0_{b}, \mathbf{X} = \mathbf{x}] \right] \\ - \frac{\mathbb{I}\{z = 1_{a}\}}{P(Z = 1_{a} \mid \mathbf{X} = \mathbf{x})} \left[y - \mathbb{E}_{P}[Y \mid Z = 1_{a}, \mathbf{X} = \mathbf{x}] \right] + \frac{\mathbb{I}\{z = 0_{a}\}}{P(Z = 0_{a} \mid \mathbf{X} = \mathbf{x})} \left[y - \mathbb{E}_{P}[Y \mid Z = 0_{a}, \mid \mathbf{X} = \mathbf{x}] \right] \\ + \delta_{b,P}(\mathbf{x}) - \delta_{a,P}(\mathbf{x}) \right\} - \frac{\Psi(P)}{\mathbb{E}_{P}[\eta_{b,P}(\mathbf{X}) - \eta_{a,P}(\mathbf{X})]} \left\{ \frac{\mathbb{I}\{z = 1_{b}\}}{P(Z = 1_{b} \mid \mathbf{X} = \mathbf{x})} \left[d - \mathbb{E}_{P}(D \mid Z = 1_{b}, \mid \mathbf{X} = \mathbf{x}) \right] \\ - \frac{\mathbb{I}\{z = 0_{b}\}}{P(Z = 0_{b} \mid \mathbf{X} = \mathbf{x})} \left[d - \mathbb{E}_{P}(Y \mid Z = 0_{b}, \mathbf{X} = \mathbf{x}) \right] - \frac{\mathbb{I}\{Z = 1_{a}\}}{P(Z = 1_{a} \mid \mathbf{X} = \mathbf{x})} \left[d - \mathbb{E}_{P}(D \mid Z = 1_{a}, \mathbf{X} = \mathbf{x}) \right] \\ + \frac{\mathbb{I}\{Z = 0_{a}\}}{P(Z = 0_{a} \mid \mathbf{X} = \mathbf{x})} \left[d - \mathbb{E}_{P}(D \mid Z = 0_{a}, \mathbf{X} = \mathbf{x}) \right] + \eta_{b,P}(\mathbf{x}) - \eta_{a,P}(\mathbf{x}) \right\}.$$

satisfies (3).

S1.4 Regularity conditions and proof for Theorem 4

Suppose that

- 1. There exists $0 < \epsilon < 0.5$ such that $\epsilon < P_0(Z|X) < 1 \epsilon$, $\epsilon < P_0(D|Z,X) < 1 \epsilon$, $\epsilon < \hat{P}_{n,k}(Z|X) < 1 \epsilon$, $\epsilon < \hat{P}_{n,k}(D|Z,X) < 1 \epsilon$ (k = 1,...K), there is a universal constant C such that $|\mathbb{E}_{P_0}[Y|Z,X]| \leq C$ with probability one.
- 2. The nuisance functions estimators are consistent:

$$\begin{aligned} \left\| \pi_{P_0}(z, \cdot) - \pi_{\widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(1) \\ \left\| \mu_{Y, P_0}(z, \cdot) - \mu_{Y, \widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(1) \\ \left\| \mu_{D, P_0}(z, \cdot) - \mu_{D, \widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(1) \end{aligned}$$

for $z \in \{0_a, 1_a, 0_b, 1_b\}$ and $k \in \{1, ..., K\}$

- 3. Rate conditions for nuisance functions estimation:
 - (a) Rate double-robustness:

$$\left\|\pi_{P_0}(z,\cdot) - \pi_{\widehat{P}_{n,k}}(z,\cdot)\right\|_{L^2(P_0)} \left\|\mu_{B,P_0}(z,\cdot) - \mu_{B,\widehat{P}_{n,k}}(z,\cdot)\right\|_{L^2(P_0)} = o_p(1/\sqrt{n})$$

for $z \in \{0_a, 1_a, 0_b, 1_b\}, k \in \{1, ..., K\}, B = Y$ or D.

(b) The convergence rates for all nuisance function estimators are faster then $n^{-1/4}$:

$$\begin{aligned} \left\| \pi_{P_0}(z, \cdot) - \pi_{\widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(n^{-1/4}) \\ \left\| \mu_{Y,P_0}(z, \cdot) - \mu_{Y,\widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(n^{-1/4}) \\ \left\| \mu_{D,P_0}(z, \cdot) - \mu_{D,\widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(n^{-1/4}) \end{aligned}$$

for $z \in \{0_a, 1_a, 0_b, 1_b\}, k \in \{1, ..., K\}.$

For asymptotic linearity of $\hat{\psi}_{ee}$, we need to assume conditions 1, 2 and 3(a) hold, for asymptotic linearity of $\hat{\psi}_{os}$, we need to assume conditions 1, 2 and 3(b) hold.

Proof. For probability measures P and \overline{P} , define

$$R(\overline{P}, P) = \Psi_{SW,\overline{P}} - \Psi_{SW,P} + PD_{\overline{P}}.$$

We will use the notations with overbar to denote functionals defined by \overline{P} , and use the notations without overbar to denote functionals defined by P. The simplied notations are listed below:

$$\begin{split} \overline{\pi}_{z}(x) &= \overline{P}(Z = z | X = x), \pi_{z}(x) = P(Z = z | X = x), \\ \overline{\mu}_{Y,z} &= \mathbb{E}_{\overline{P}}[Y | Z = z, X = x], \mu_{Y,z} = \mathbb{E}_{P}[Y | Z = z, X = x], \\ \overline{\mu}_{D,z} &= \mathbb{E}_{\overline{P}}[D | Z = z, X = x], \mu_{D,z} = \mathbb{E}_{P}[D | Z = z, X = x], \\ \overline{\omega}_{1} &= \mathbb{E}_{\overline{P}}[\delta_{b,\overline{P}}(\mathbf{X}) - \delta_{a,\overline{P}}(\mathbf{X})], \omega_{1} = \mathbb{E}_{P}[\delta_{b,P}(\mathbf{X}) - \delta_{a,P}(\mathbf{X})], \\ \overline{\omega}_{2} &= \mathbb{E}_{\overline{P}}[\eta_{b,\overline{P}}(\mathbf{X}) - \eta_{a,\overline{P}}(\mathbf{X})], \omega_{2} = \mathbb{E}_{P}[\eta_{b,P}(\mathbf{X}) - \eta_{a,P}(\mathbf{X})] \end{split}$$

For $z \in \{1_b, 0_b, 1_a, 0_a\}$, and B is equal to either Y or D, we have (we will use lower case to

denote realization of random variables),

$$\begin{split} & P\left\{\frac{\mathbbm{1}\{z=1_b\}}{\overline{\pi}_z} \Big[b-\overline{\mu}_{B,z}\Big] + \overline{\mu}_{B,z}\right\} \\ &= P\left\{\mathbbm{1}\{z=1_b\} \Big(\frac{1}{\overline{\pi}_z} - \frac{1}{\pi_z} + \frac{1}{\pi_z}\Big) \Big[(b-\mu_{B,z}) + (\mu_{B,z} - \overline{\mu}_{B,z})\Big] + (\overline{\mu}_{B,z} - \mu_{B,z}) + \mu_{B,z}\right\} \\ &= P\left\{\mathbbm{1}\{z=1_b\} \frac{1}{\pi_z} \Big[b-\mu_{B,z}\Big] + \mu_{B,z}\right\} + P\left\{\mathbbm{1}\{z=1_b\} \Big(\frac{1}{\overline{\pi}_z} - \frac{1}{\pi_z}\Big) \Big(\mu_{B,z} - \overline{\mu}_{B,z}\Big)\right\} \\ &+ P\left\{\mathbbm{1}\{z=1_b\} \frac{1}{\pi_z} (\overline{\mu}_{B,z} - \mu_{B,z}) - (\overline{\mu}_{B,z} - \mu_{B,z})\right\} + P\left\{\mathbbm{1}\{z=1_b\} \Big(\frac{1}{\overline{\pi}_z} - \frac{1}{\pi_z}\Big) \Big(y-\mu_{B,z}\Big)\right\} \\ &= P\mu_{B,z} + P\left\{\mathbbm{1}\{z=1_b\} \Big(\frac{1}{\overline{\pi}_z} - \frac{1}{\pi_z}\Big) \Big(\mu_{B,z} - \overline{\mu}_{B,z}\Big)\right\} \end{split}$$

Plug this into $R(\overline{P}, P)$, we have

$$\begin{split} R(\overline{P},P) \\ = & \Psi_{SW,\overline{P}} - \Psi_{SW,P} + PD_{\overline{P}} \\ = & \frac{1}{\overline{\omega}_{2}} \Biggl\{ P \Bigl[\mathbbm{1} \{z = 1_{b} \} \Bigl(\frac{1}{\overline{\pi}_{1_{b}}} - \frac{1}{\pi_{1_{b}}} \Bigr) (\mu_{Y,1_{b}} - \overline{\mu}_{Y,1_{b}}) \Bigr] - P \Bigl[\mathbbm{1} \{z = 0_{b} \} \Bigl(\frac{1}{\overline{\pi}_{0_{b}}} - \frac{1}{\pi_{0_{b}}} \Bigr) (\mu_{Y,0_{b}} - \overline{\mu}_{Y,0_{b}}) \Bigr] \\ & - P \Bigl[\mathbbm{1} \{z = 1_{a} \} \Bigl(\frac{1}{\overline{\pi}_{1_{a}}} - \frac{1}{\pi_{1_{a}}} \Bigr) (\mu_{Y,1_{a}} - \overline{\mu}_{Y,1_{a}}) \Bigr] + P \Bigl[\mathbbm{1} \{z = 0_{a} \} \Bigl(\frac{1}{\overline{\pi}_{0_{a}}} - \frac{1}{\pi_{0_{a}}} \Bigr) (\mu_{Y,0_{a}} - \overline{\mu}_{Y,0_{a}}) \Bigr] \Biggr\} \\ & - \frac{\overline{\omega}_{1}}{\overline{\omega}_{2}^{2}} \Biggl\{ P \Bigl[\mathbbm{1} \{z = 1_{b} \} \Bigl(\frac{1}{\overline{\pi}_{1_{b}}} - \frac{1}{\pi_{1_{b}}} \Bigr) (\mu_{D,1_{b}} - \overline{\mu}_{D,1_{b}}) \Bigr] - P \Bigl[\mathbbm{1} \{z = 0_{b} \} \Bigl(\frac{1}{\overline{\pi}_{0_{b}}} - \frac{1}{\pi_{0_{b}}} \Bigr) (\mu_{D,0_{b}} - \overline{\mu}_{D,0_{b}}) \Bigr] \Biggr\} \end{split}$$

$$-P\Big[\mathbb{1}\{z=1_{a}\}\Big(\frac{1}{\overline{\pi}_{1_{a}}}-\frac{1}{\pi_{1_{a}}}\Big)(\mu_{D,1_{a}}-\overline{\mu}_{D,1_{a}})\Big]+P\Big[\mathbb{1}\{z=0_{a}\}\Big(\frac{1}{\overline{\pi}_{0_{a}}}-\frac{1}{\pi_{0_{a}}}\Big)(\mu_{D,0_{a}}-\overline{\mu}_{D,0_{a}})\Big]\Big\}$$
$$+\frac{1}{\overline{\omega}_{2}^{2}}(\omega_{2}-\overline{\omega}_{2})(\overline{\omega}_{1}-\omega_{1})+\frac{\omega_{1}}{\overline{\omega}_{2}^{2}}(\omega_{2}-\overline{\omega}_{2})^{2}$$

Then for k = 1, ..., K,

$$\begin{aligned} \hat{\psi}_{os,k} &- \Psi_{SW,P_0} \\ &= \Psi_{SW,\hat{P}_{n,k}} + \mathbb{P}_{n,k} D_{\hat{P}_n} - \Psi_{SW,P_0} \\ &= (\mathbb{P}_{n,k} - P_0) D_{P_0} + (\mathbb{P}_{n,k} - P_0) (D_{\hat{P}_{n,k}} - D_{P_0}) + R(\hat{P}_{n,k}, P_0) \end{aligned}$$

We will show that $\sqrt{n}R(\hat{P}_{n,k}, P_0) = o_p(1)$. Now we define new notations

$$\begin{split} \hat{\pi}_{z,k}(x) &= \hat{P}_{n,k}(Z = z | X = x), \\ \pi_{0,z}(x) &= P_0(Z = z | X = x), \\ \hat{\mu}_{Y,z,k} &= \mathbb{E}_{\hat{P}_{n,k}}[Y | Z = z, X = x], \\ \mu_{0,Y,z} &= \mathbb{E}_{P_0}[Y | Z = z, X = x], \\ \hat{\mu}_{D,z,k} &= \mathbb{E}_{\hat{P}_{n,k}}[D | Z = z, X = x], \\ \mu_{0,D,z} &= \mathbb{E}_{P_0}[D | Z = z, X = x], \\ \hat{\omega}_{1,k} &= \mathbb{E}_{\hat{P}_{n,k}}[\delta_{b,\hat{P}_{n,k}}(\mathbf{X}) - \delta_{a,\hat{P}_{n,k}}(\mathbf{X})], \\ \omega_{0,1} &= \mathbb{E}_{P_0}[\delta_{b,P_0}(\mathbf{X}) - \delta_{a,P_0}(\mathbf{X})], \\ \hat{\omega}_{2,k} &= \mathbb{E}_{\hat{P}_{n,k}}[\eta_{b,\hat{P}_{n,k}}(\mathbf{X}) - \eta_{a,\hat{P}_{n,k}}(\mathbf{X})], \\ \omega_{0,2} &= \mathbb{E}_{P_0}[\eta_{b,P_0}(\mathbf{X}) - \eta_{a,P_0}(\mathbf{X})]. \end{split}$$

Then with probability approaching one, we know there exists a constant C such that

$$\Big|\frac{1}{\hat{\omega}_{2,k}}\Big| \leq C, \Big|\frac{1}{\hat{\omega}_{2,k}^2}\Big| \leq C, \Big|\frac{\hat{\omega}_{1,k}}{\hat{\omega}_{2,k}^2}\Big| \leq C, \Big|\frac{\hat{\omega}_{1,k}}{\hat{\omega}_{2,k}^2\omega_{0,2}}\Big| \leq C,$$

Under our assumptions for nuisance parameter estimation, we have

$$P_0\Big[\mathbbm{1}\{z=z\}\Big(\frac{1}{\hat{\pi}_{z,k}}-\frac{1}{\pi_z}\Big)(\mu_{B,z}-\hat{\mu}_{B,z,k})\Big] \le C\Big\|\frac{1}{\hat{\pi}_{z,k}}-\frac{1}{\pi_z}\Big\|_{L^2(P_0)}\Big\|\mu_{B,z}-\hat{\mu}_{B,z,k}\Big\|_{L^2(P_0)} = o_p(n^{-1/2})$$

for $z \in \{1_b, 0_b, 1_a, 0_a\}$. Similarly, by the fact we used cross-fitting and Holder's inequality, we have

$$\begin{aligned} |\omega_{0,2} - \hat{\omega}_{2,k}| |\omega_{0,1} - \hat{\omega}_{1,k}| &= o_p(n^{-1/2}), \\ |\omega_{0,2} - \hat{\omega}_{2,k}|^2 &= o_p(n^{-1/2}) \end{aligned}$$

Therefore, we have proved that $R(\hat{P}_{n,k}, P_0) = o_p(n^{-1/2}).$

Now we discuss how to handle $(\mathbb{P}_{n,k} - P_0)(D_{\hat{P}_{n,k}} - D_{P_0})$. Since we used cross-fitting, we have

for fixed t > 0,

$$\begin{split} & P_0 \left\{ \left| [\mathbb{P}_{n,k} - P_0] [D_{\hat{P}_{n,k}} - D_{P_0}] \right| > (n/K)^{-1/2} t \middle| \ k\text{-th subsample} \right\} \\ & \leq \min \left\{ 1, \frac{\operatorname{Var}\{ [D_{\hat{P}_{n,k}} - D_{P_0}] | \ k\text{-th subsample}\}}{t^2} \right\} \\ & = \min \left\{ 1, \frac{P_0\{ [D_{\hat{P}_{n,k}} - D_{P_0}]^2 | \ k\text{-th subsample}\}}{t^2} \right\} \\ & = \min \left\{ 1, \frac{P_0[D_{\hat{P}_{n,k}} - D_{P_0}]^2}{t^2} \right\} \\ & = \min \left\{ 1, o_p(1) \right\} \end{split}$$

Taking an expectation of both sides and applying the Dominated convergence theorem shows that

$$P_0 \left\{ \left| [\mathbb{P}_{n,k} - P_0] [D_{\hat{P}_{n,k}} - D_{P_0}] \right| > (n/K)^{-1/2} t \right\}$$

= $E[\min 1, o_p(1)]$
 $\to 0$

As t > 0 was arbitrary, we have

$$(\mathbb{P}_{n,k} - P_0)(D_{\hat{P}_{n,k}} - D_{P_0}) = o_p(n^{-1/2}).$$

To sum up,

$$\begin{split} &\sqrt{n}(\hat{\psi}_{os} - \Psi_{SW,P_0}) \\ = &\sqrt{n}(\frac{1}{K} \sum_{k=1}^{K} (\hat{\psi}_{os,k} - \Psi_{SW,P_0})) \\ = &\sqrt{n}(\frac{1}{K} \sum_{k=1}^{K} (\mathbb{P}_{n,k} - P_0) D_{P_0}) + o_p(1) \\ = &\sqrt{n}(\mathbb{P}_n - P_0) D_{P_0} \\ &\rightsquigarrow N(0, P_0 D_{P_0}^2) \end{split}$$

where the last line holds if we further assume $P_0 D_{P_0}^2 < \infty$.

For $\widehat{\psi}_{ee}$, we notice that

$$\widehat{\psi}_{ee,k} = \frac{\mathbb{P}_{n,k}A_{n,k}}{\mathbb{P}_{n,k}B_{n,k}}$$

where

$$\begin{split} A_{n,k} &= \frac{\mathbb{1}\{z = 1_b\}}{\hat{P}_{n,k}(Z = 1_b \mid \mathbf{X} = \mathbf{x})} \Big[y - \mathbb{E}_{\hat{P}_{n,k}}[Y \mid Z = 1_b, \mathbf{X} = \mathbf{x}] \Big] - \frac{\mathbb{1}\{z = 0_b\}}{\hat{P}_{n,k}(Z = 0_b \mid \mathbf{X} = \mathbf{x})} \Big[y - \mathbb{E}_{\hat{P}_{n,k}}[Y \mid Z = 0_b, \mathbf{X} = \mathbf{x}] \Big] \\ &- \frac{\mathbb{1}\{z = 1_a\}}{\hat{P}_{n,k}(Z = 1_a \mid \mathbf{X} = \mathbf{x})} \Big[y - \mathbb{E}_{\hat{P}_{n,k}}[Y \mid Z = 1_a, \mathbf{X} = \mathbf{x}] \Big] + \frac{\mathbb{1}\{z = 0_a\}}{\hat{P}_{n,k}(Z = 0_a \mid \mathbf{X} = \mathbf{x})} \Big[y - \mathbb{E}_{\hat{P}_{n,k}}[Y \mid Z = 0_a, \mathbf{X} = \mathbf{x}] \Big] \\ &+ \delta_{b,\hat{P}_{n,k}}(\mathbf{x}) - \delta_{a,\hat{P}_{n,k}}(\mathbf{x}) \end{split}$$

$$B_{n,k} = \frac{\mathbb{1}\{z = 1_b\}}{\hat{P}_{n,k}(Z = 1_b \mid \boldsymbol{X} = \boldsymbol{x})} \Big[d - \mathbb{E}_{\hat{P}_{n,k}}[D \mid Z = 1_b, \boldsymbol{X} = \boldsymbol{x}] \Big] - \frac{\mathbb{1}\{z = 0_b\}}{\hat{P}_{n,k}(Z = 0_b \mid \boldsymbol{X} = \boldsymbol{x})} \Big[d - \mathbb{E}_{\hat{P}_{n,k}}[D \mid Z = 0_b, \boldsymbol{X} = \boldsymbol{x}] \Big] \\ - \frac{\mathbb{1}\{z = 1_a\}}{\hat{P}_{n,k}(Z = 1_a \mid \boldsymbol{X} = \boldsymbol{x})} \Big[d - \mathbb{E}_{\hat{P}_{n,k}}[D \mid Z = 1_a, \boldsymbol{X} = \boldsymbol{x}] \Big] + \frac{\mathbb{1}\{z = 0_a\}}{\hat{P}_{n,k}(Z = 0_a \mid \boldsymbol{X} = \boldsymbol{x})} \Big[d - \mathbb{E}_{\hat{P}_{n,k}}[D \mid Z = 0_a, \boldsymbol{X} = \boldsymbol{x}] \Big] \\ + \eta_{b,\hat{P}_{n,k}}(\boldsymbol{x}) - \eta_{a,\hat{P}_{n,k}}(\boldsymbol{x})$$

Since under the given conditions, $\mathbb{P}_{n,k}A_{n,k}$ and $\mathbb{P}_{n,k}B_{b,k}$ are asymptotically linear estimators for $E_{P_0}[\delta_{b,P_0} - \delta_{a,P_0}]$ and $E_{P_0}[\eta_{b,P_0} - \eta_{a,P_0}]$, respectively, the influence functions (denoted as D_{A,P_0} and D_{B,P_0}) at P_0 are

$$\begin{aligned} D_{A,P_0} : o \mapsto E_{P_0}[Y|Z = 1_b, X = x] + \frac{\mathbb{1}(z = 1_b)}{P_0(Z = 1_b|X = x)} \big[y - \mathbb{E}_{P_0}[Y|Z = z, X = x] \big] \\ &- E_{P_0}[Y|Z = 0_b, X = x] - \frac{\mathbb{1}(z = 0_b)}{P_0(Z = 0_b|X = x)} \big[y - \mathbb{E}_{P_0}[Y|Z = z, X = x] \big] \\ &- E_{P_0}[Y|Z = 1_a, X = x] - \frac{\mathbb{1}(z = 1_a)}{P_0(Z = 1_a|X = x)} \big[y - \mathbb{E}_{P_0}[Y|Z = z, X = x] \big] \\ &+ E_{P_0}[Y|Z = o_a, X = x] + \frac{\mathbb{1}(z = 0_a)}{P_0(Z = 0_a|X = x)} \big[y - \mathbb{E}_{P_0}[Y|Z = z, X = x] \big] \\ &- E_{P_0}[\delta_{b,P_0} - \delta_{a,P_0}] \end{aligned}$$

and

$$\begin{split} D_{B,P_0} &: o \mapsto E_{P_0}[D|Z = 1_b, X = x] + \frac{\mathbb{1}(z = 1_b)}{P_0(Z = 1_b|X = x)} \Big[d - \mathbb{E}_{P_0}[D|Z = z, X = x] \Big] \\ &- E_{P_0}[D|Z = 0_b, X = x] - \frac{\mathbb{1}(z = 0_b)}{P_0(Z = 0_b|X = x)} \Big[d - \mathbb{E}_{P_0}[D|Z = z, X = x] \Big] \\ &- E_{P_0}[D|Z = 1_a, X = x] - \frac{\mathbb{1}(z = 1_a)}{P_0(Z = 1_a|X = x)} \Big[d - \mathbb{E}_{P_0}[D|Z = z, X = x] \Big] \\ &+ E_{P_0}[D|Z = o_a, X = x] + \frac{\mathbb{1}(z = 0_a)}{P_0(Z = 0_a|X = x)} \Big[d - \mathbb{E}_{P_0}[D|Z = z, X = x] \Big] \\ &- E_{P_0}[\eta_{b,P_0} - \eta_{a,P_0}]. \end{split}$$

Let $f(x_1, x_2) = x_1/x_2$, then by delta method Van der Vaart (2000), $\widehat{\psi}_{ee,k}$ should also be asymptotically linear, where the influence function should be

$$o \mapsto (\nabla f(E_{P_0}[\delta_{b,P_0} - \delta_{a,P_0}], E_{P_0}[\eta_{b,P_0} - \eta_{a,P_0}])^T (D_{A,P_0}(o), D_{A,P_0}(o))^T$$

which is exactly D_{P_0} . Therefore, we also have

$$\sqrt{n/K}(\hat{\psi}_{ee,k} - \Psi_{SW,P_0}) = \sqrt{n/K}(\mathbb{P}_{n,k} - P_0)D_{P_0} + o_p(1/\sqrt{n})$$

Therefore,

$$\sqrt{n}(\hat{\psi}_{ee} - \Psi_{SW,P_0}) = \sqrt{n}(\mathbb{P}_n - P_0)D_{P_0} + o_p(1/\sqrt{n}).$$

S1.5 Proof of Theorem 5

Proof. β_P satisfies the following equation:

$$\int q(x;\beta_P)(\theta_P(x) - \gamma(x;\beta_{P_t}))dP(x) = 0,$$

where $q(x; \beta_P) = \frac{\gamma(x;\beta)}{\partial \beta}|_{\beta = \beta_P}$

Now we pick a one-dimensional parametric submodel $\{P_t : t \in [0, \epsilon)\}$, then we have

$$\int q(x;\beta_{P_t})(\theta_{P_t}(x) - \gamma(x;\beta_{P_t}))dP_t(x) = 0$$

Take derivative with respect to t, then evaluate the derivative at t = 0, we have

$$\int \frac{\partial q(x;\beta)}{\partial \beta} \Big|_{\beta=\beta_{P_0}} \frac{d\beta_{P_t}}{dt} \Big|_{t=0} [\theta_{P_0}(x) - \gamma(x;\beta_{P_0})] dP_0(x) + \int q(x;\beta_{P_0}) \Big[\frac{\theta_{P_t}}{\partial t} \Big|_{t=0} - q^T(x;\beta_{P_0}) \frac{\partial \beta_{P_t}}{\partial t} \Big|_{t=0} \Big] dP_0(x) \int q(x;\beta_{P_0}) [\theta_{P_0}(x) - \gamma(x;\beta_{P_0})] s_0(x) dP_0(x) = 0$$

Therefore

$$C\frac{\partial\beta_{P_t}}{\partial t} = \int q(x;\beta_{P_0})\frac{\theta_{P_t}(x)}{\partial t}\Big|_{t=0} dP_0(x) + E_{P_0}\Big[q(X;\beta_{P_0})[\theta_0(X) - \gamma(X;\beta_{P_0})]s_0(O)\Big],$$

where C is a constant, when $\gamma(x;\beta) = x^T\beta$, we have $\frac{\partial q(x;\beta)}{\partial \beta} = 0$, $q(x;\beta) = x$ and therefore $C = \int x^T x dP_0$. Note that $q(X;\beta_{P_0})[\theta_0(X) - \gamma(X;\beta_{P_0})]$ is already mean zero, therefore we only need to arrange the first term. Since $\theta_{P_t}(x) = \theta_{1,P_t}(x) - \theta_{2,P_t}(x)$. We first handle $\frac{\partial \theta_{1,P_t}(x)}{\partial t}$:

$$\begin{split} & \frac{\partial \theta_{1,P_{t}}(x)}{\partial t} \\ &= \frac{\frac{\partial \delta_{a,P_{t}}(x)}{\partial t}|_{t=0}\eta_{a,P_{0}}(x) - \delta_{a,P_{0}}(x)\frac{\partial \eta_{a,P_{t}}(x)}{\partial t}|_{t=0}}{[\eta_{a,P_{0}}(x)]^{2}} \\ &= \frac{1}{\eta_{a,P_{0}}(x)} \Big[\frac{\partial \delta_{a,P_{t}}(x)}{\partial t}\Big|_{t=0} - \theta_{1,P_{0}}(x)\frac{\partial \eta_{a,P_{t}}(x)}{\partial t}\Big|_{t=0}\Big] \end{split}$$

It remains to arrange

$$\int q(x;\beta_{P_0}) \frac{1}{\eta_{a,P_0}(x)} \frac{\delta_{a,P_t}(x)}{\partial t} \Big|_{t=0} dP_0(x) - \int q(x;\beta_{P_0}) \frac{\theta_{1,P_0}(x)}{\eta_{a,P_0}(x)} \frac{\partial \eta_{a,P_t}(x)}{\partial t} \Big|_{t=0} dP_0(x)$$

For any h(x), we calculate $\mathbb{E}_{P_0}[h(X)\frac{\partial}{\partial t}E_{P_t}[C|Z=z,X]|_{t=0}]$, where C is equal to either Y or D.

$$\begin{split} & \mathbb{E}_{P_0} \left[h(X) \frac{\partial}{\partial t} E_{P_t}[C|Z = z, X] \Big|_{t=0} \right] \\ = & \mathbb{E}_{P_0} \left[h(X) E_{P_0}[CS_{C|Z=z,X}(C,X)|Z = z, X] \right] \\ = & \mathbb{E}_{P_0} \left[h(X) \mathbb{E}_{P_0}[(C - \mathbb{E}_{P_0}(C|Z = z, X)S_{C|Z=z,X}(C,X)|Z = z, X] \right] \\ = & \mathbb{E}_{P_0} \left[\frac{h(X)}{P_0(Z = z|X)} \mathbb{E}_{P_0}[1\{Z = z\}(C - E_{P_0}(C|Z,X)S_{C|Z=z,X}(C,X)|X] \right] \\ = & \mathbb{E}_{P_0} \left[\frac{h(X)\mathbb{E}_{P_0}[1\{Z = z\}(C - \mathbb{E}_{P_0}[C|Z,X])S_{C|Z,X}(C,Z,X)|X]}{P_0(Z = z|X)} \right] \\ = & \mathbb{E}_{P_0} \left[\frac{h(X)[1\{Z = z\}(C - \mathbb{E}_{P_0}[C|Z,X])}{P_0(Z = z|X)} S_{C|Z,X}(C,Z,X) \right] \end{split}$$

Furthermore,

$$\mathbb{E}_{P_0}\left[\frac{h(X)[1\{Z=z\}(C-\mathbb{E}_{P_0}[C|Z,X])}{P_0(Z=z|X)}S_{Z,X}(Z,X)\right] = 0$$

Therefore we have

$$\mathbb{E}_{P_0}\left[\frac{h(X)[1\{Z=z\}(C-\mathbb{E}_{P_0}[C|Z,X])}{P_0(Z=z|X)}S_O(O)\right] = 0$$

Therefore,

$$\begin{split} &\int q(x;\beta_{P_0}) \frac{1}{\eta_{a,P_0}(x)} \frac{\delta_{a,P_t}(x)}{\partial t} \Big|_{t=0} dP_0(x) \\ = & \mathbb{E}_{P_0} \Bigg[\frac{q(X;\beta_0)}{\eta_{a,P_0}(X)} \Bigg[\frac{1\{Z=1_a\}}{P_0(Z=1_a|X)} (Y - \mathbb{E}_{P_0}(Y|Z=1_a,X)) \\ &- \frac{1\{Z=0_a\}}{P_0(Z=0_a|X)} (Y - \mathbb{E}_{P_0}(Y|Z=0_a,X)) \Bigg] S_O(O) \Bigg] \end{split}$$

Similarly,

$$\int q(x; \beta_{P_0}) \frac{\theta_{1,P_0}(x)}{\eta_{a,P_0}(x)} \frac{\eta_{a,P_t}(x)}{\partial t} \Big|_{t=0} dP_0(x)$$

= $\mathbb{E}_{P_0} \left[\frac{q(X; \beta_0) \theta_{1,P_0}(X)}{\eta_{a,P_0}(X)} \left[\frac{1\{Z = 1_a\}}{P_0(Z = 1_a|X)} (D - \mathbb{E}_{P_0}(D|Z = 1_a, X)) - \frac{1\{Z = 0_a\}}{P_0(Z = 0_a|X)} (D - \mathbb{E}_{P_0}(D|Z = 0_a, X)) \right] S_O(O) \right]$

We repeat the similar calculation for the term involving θ_{2,P_t} , then we can obtain the canonical gradient as stated in theorem 5.

S1.6 Proof of Proposition 2

Proof. Suppose principal ignorability holds, then by definition of principal ignorability, we have

$$ATE_{P_0}(x) = SWATE_{P_0}(x) = ACOATE_{P_0}(x).$$

Suppose no unmeasured common effect modifier holds, then

$$(\mathbb{E}_{P_0}[Y|X, Z = 1_a] - \mathbb{E}_{P_0}[Y|X, Z = 0_a]) - (\mathbb{E}_{P_0}[Y|X, Z = 1_b] - \mathbb{E}_{P_0}[Y|X, Z = 0_b])$$
$$= (\mathbb{E}_{P_0}[Y(Z = 1_a)|X] - \mathbb{E}_{P_0}[Y(Z = 0_a)|X]) - (\mathbb{E}_{P_0}[Y(Z = 1_b)|X] - \mathbb{E}_{P_0}[Y(Z = 0_b)|X])$$

Notice that

$$\begin{split} & \mathbb{E}_{P_0}[Y(Z=1_a)|X] - \mathbb{E}_{P_0}[Y(Z=0_a)|X] \\ = & \mathbb{E}_{P_0}[D(Z=1_a)Y(D=1) + (1 - D(Z=1_a))Y(D=0)|X] - \\ & \mathbb{E}_{P_0}[D(Z=0_a)Y(D=1) + (1 - D(Z=0_a))Y(D=0)|X] \\ = & \mathbb{E}_{P_0}[(D(Z=1_a) - D(Z=0_a))(Y(D=1) - Y(D=0))|X] \\ = & \mathbb{E}_{P_0}[D(Z=1_a) - D(Z=0_a)|X] \mathbb{E}_{P_0}[Y(D=1) - Y(D=0)|X] \end{split}$$

similarly we have

$$\mathbb{E}_{P_0}[Y(Z=1_b)|X] - \mathbb{E}_{P_0}[Y(Z=0_b)|X] = \mathbb{E}_{P_0}[D(Z=1_b) - D(Z=0_b)|X] \mathbb{E}_{P_0}[Y(D=1) - Y(D=0)|X]$$

Therefore,

$$(\mathbb{E}_{P_0}[Y|X, Z = 1_a] - \mathbb{E}_{P_0}[Y|X, Z = 0_a]) - (\mathbb{E}_{P_0}[Y|X, Z = 1_b] - \mathbb{E}_{P_0}[Y|X, Z = 0_b])$$

= $(\mathbb{E}_{P_0}[D(Z = 1_a) - D(Z = 0_a)|X] - \mathbb{E}_{P_0}[D(Z = 1_b) - D(Z = 0_b)|X])\mathbb{E}_{P_0}[Y(D = 1) - Y(D = 0)|X]$
= $[(\mathbb{E}_{P_0}[D|X, Z = 1_a] - \mathbb{E}_{P_0}[D|X, Z = 0_a]) - (\mathbb{E}_{P_0}[D|X, Z = 1_b] - \mathbb{E}_{P_0}[D|X, Z = 0_b])]ATE_{P_0}(X)$

Therefore, $ATE_{P_0}(X) = SWATE_{P_0}(X)$. Similarly, we have $ATE_{P_0}(X) = ACOATE_{P_0}(X)$. \Box

S1.7 Proof of Theorem 6

Suppose that

- 1. There exists $0 < \epsilon < 0.5$ such that $\epsilon < P_0(Z|X) < 1 \epsilon$, $\epsilon < P_0(D|Z,X) < 1 \epsilon$, $\epsilon < \widehat{P}_{n,k}(Z|X) < 1 \epsilon$, $\epsilon < \widehat{P}_{n,k}(D|Z,X) < 1 \epsilon$ (k = 1,...K), there is a universal constant C such that $|\mathbb{E}_{P_0}[Y|Z,X]| \leq C$ with probability one.
- 2. $||X||_{\infty} < C, \lambda_{\min}(\frac{1}{n}\sum_{i=1}^{n}X_{i}^{T}X_{i}) > 0$ with probability one. $\lambda_{\min}E[X_{i}^{T}X_{i}] > 0$. Here for any matrix $H, \lambda_{\min}H$ denoted the smallest eigenvalue of H.
- 3. Rate conditions for nuisance functions estimation: The convergence rates for all nuisance function estimators are faster then $n^{-1/4}$:

$$\begin{aligned} \left\| \pi_{P_0}(z, \cdot) - \pi_{\widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(n^{-1/4}) \\ \left\| \mu_{Y,P_0}(z, \cdot) - \mu_{Y,\widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(n^{-1/4}) \\ \left\| \mu_{D,P_0}(z, \cdot) - \mu_{D,\widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(n^{-1/4}) \end{aligned}$$

for $z \in \{0_a, 1_a, 0_b, 1_b\}, k \in \{1, ..., K\}.$

Proof. The proof is very similar to the proof of theorem 4. We will only state the proof for test

 $T^{(1)}_{Wald,\alpha}.$ We define

$$R(\overline{P}, P) = \beta_{\overline{P}}^{(1)} - \beta_{P}^{(1)} + P\phi_{\overline{P}}^{(1)}.$$

Similar to the proof of Theorem 4, we define the following notations:

$$\begin{split} \overline{\pi}_z(x) &= \overline{P}(Z = z | X = x), \pi_z(x) = P(Z = z | X = x), \\ \overline{\mu}_{Y,z} &= \mathbb{E}_{\overline{P}}[Y | Z = z, X = x], \mu_{Y,z} = \mathbb{E}_P[Y | Z = z, X = x], \\ \overline{\mu}_{D,z} &= \mathbb{E}_{\overline{P}}[D | Z = z, X = x], \mu_{D,z} = \mathbb{E}_P[D | Z = z, X = x], \\ \overline{\delta}_g &= \delta_{g,\overline{P}}, \delta_g = \delta_{g,P}, \overline{\eta}_g = \eta_{g,\overline{P}}, \eta_g = \eta_{g,P} \end{split}$$

for $z \in \{1_b, 0_b, 1_a, 0_a\}$ and $g \in \{a, b\}$.

We first study for fixed $z' \in \{1_b, 0_b, 1_a, 0_a\}$, direct calculation yields

$$\begin{split} &P\left[\frac{1}{\bar{\eta}_g}\left(\frac{\mathbbm{1}\{z=z'\}}{\bar{\pi}_{z'}}(y-\bar{\mu}_{Y,z'})\right)\right]\\ =&P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\left(\frac{\mathbbm{1}\{z=z'\}}{\pi_{z'}}(y-\mu_{Y,z'})\right)\right] + P\left[\frac{1}{\eta_g}\left(\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(y-\mu_{Y,z'})\right)\right]\\ &+P\left[\frac{1}{\eta_g}\left(\mathbbm{1}\{z=z'\}\frac{1}{\pi_{z'}}(\bar{\mu}_{Y,z'}-\mu_{Y,z'})\right)\right] + P\left[\frac{1}{\eta_g}\frac{\mathbbm{1}\{z=z'\}}{\pi_{z'}}(y-\mu_{Y,z'})\right]\\ &+P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\frac{1}{\pi_{z'}}(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] + P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(y-\mu_{Y,z'})\right]\\ &+P\left[\frac{1}{\eta_g}\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right]\\ &+P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right]\\ &=P\left[\frac{1}{\eta_g}(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] + P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] + P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] \\ &+P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] + P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] \\ &+P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] + P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'}\right)\right] \\ &+P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] + P\left[\frac{1}{\eta_g}\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'}\right)\right] \\ &+P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] + P\left[\frac{1}{\eta_g}\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] \\ &+P\left[\left(\frac{1}{\bar{\eta}_g}-\frac{1}{\eta_g}\right)\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] + P\left[\frac{1}{\eta_g}\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] \\ &+P\left[\frac{1}{\eta_g}\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\right] \\ &+P\left[\frac{1}{\eta_g}\mathbbm{1}\{z=z'\}\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y,z'})\left(\frac{1}{\bar{\pi}_{z'}}-\frac{1}{\pi_{z'}}\right)(\mu_{Y,z'}-\bar{\mu}_{Y$$

Then

$$\begin{split} &\beta_{\overline{P}}^{(1)} - \beta_{P}^{(1)} + P\phi_{\overline{P}}^{(1)} \\ = &(\overline{P}xx^{T})^{-1}\overline{P} \left[x \left(\frac{\overline{\delta}_{a}}{\overline{\eta}_{a}} - \frac{\overline{\delta}_{b} - \overline{\delta}_{a}}{\overline{\eta}_{b} - \overline{\eta}_{a}} \right) \right] - (Pxx^{T})^{-1}P \left[x \left(\frac{\delta_{a}}{\eta_{a}} - \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}} \right) \right] \\ &+ (\overline{P}xx^{T})^{-1}P \left[x \left(D_{\overline{P}}^{1} - D_{\overline{P}}^{2} + \frac{\overline{\delta}_{a}}{\overline{\eta}_{a}} - \frac{\overline{\delta}_{b} - \overline{\delta}_{a}}{\overline{\eta}_{b} - \overline{\eta}_{a}} - x^{T}(\overline{P}xx^{T})^{-1}\overline{P} \left[x \left(\frac{\overline{\delta}_{a}}{\overline{\eta}_{a}} - \frac{\overline{\delta}_{b} - \overline{\delta}_{a}}{\overline{\eta}_{b} - \overline{\eta}_{a}} \right) \right] \right) \right] \\ &= \left\{ (\overline{P}xx^{T})^{-1}\overline{P} \left[x \frac{\overline{\delta}_{a}}{\overline{\eta}_{a}} \right] - (Pxx^{T})^{-1}P \left[x \frac{\delta_{a}}{\eta_{a}} \right] + (\overline{P}xx^{T})^{-1}P \left[x \left(D_{\overline{P}}^{1} + \frac{\overline{\delta}_{a}}{\overline{\eta}_{a}} - x^{T}(\overline{P}xx^{T})^{-1}\overline{P} \left[x \frac{\overline{\delta}_{a}}{\overline{\eta}_{a}} \right] \right) \right] \right\} \\ &- \left\{ (\overline{P}xx^{T})^{-1}\overline{P} \left[x \frac{\overline{\delta}_{b} - \overline{\delta}_{a}}{\overline{\eta}_{b} - \overline{\eta}_{a}} \right] - (Pxx^{T})^{-1}P \left[x \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}} \right] \\ &+ (\overline{P}xx^{T})^{-1}P \left[x \left(D_{\overline{P}}^{2} + \frac{\overline{\delta}_{b} - \overline{\delta}_{a}}{\overline{\eta}_{b} - \overline{\eta}_{a}} - x^{T}(\overline{P}xx^{T})^{-1}\overline{P} \left[x \frac{\overline{\delta}_{b} - \overline{\delta}_{a}}{\overline{\eta}_{b} - \overline{\eta}_{a}} \right] \right) \right] \right\} \end{split}$$

Let

$$\begin{split} A &:= (\overline{P}xx^T)^{-1}\overline{P}\left[x\frac{\overline{\delta}_a}{\overline{\eta}_a}\right] - (Pxx^T)^{-1}P\left[x\frac{\delta_a}{\eta_a}\right] + (\overline{P}xx^T)^{-1}P\left[x\left(D_{\overline{P}}^1 + \frac{\overline{\delta}_a}{\overline{\eta}_a} - x^T(\overline{P}xx^T)^{-1}\overline{P}\left[x\frac{\overline{\delta}_a}{\overline{\eta}_a}\right]\right)\right] \\ B &:= (\overline{P}xx^T)^{-1}\overline{P}\left[x\frac{\overline{\delta}_b - \overline{\delta}_a}{\overline{\eta}_b - \overline{\eta}_a}\right] - (Pxx^T)^{-1}P\left[x\frac{\delta_b - \delta_a}{\eta_b - \eta_a}\right] \\ &+ (\overline{P}xx^T)^{-1}P\left[x\left(D_{\overline{P}}^2 + \frac{\overline{\delta}_b - \overline{\delta}_a}{\overline{\eta}_b - \overline{\eta}_a} - x^T(\overline{P}xx^T)^{-1}\overline{P}\left[x\frac{\overline{\delta}_b - \overline{\delta}_a}{\overline{\eta}_b - \overline{\eta}_a}\right]\right)\right] \end{split}$$

Direct calculation yields

$$\begin{split} &A = (\overline{P}xx^{T})^{-1}P(xx^{T})\Big(P(xx^{T})^{-1} - \overline{P}(xx^{T})^{-1}\Big)\left(\overline{P}x\frac{\overline{\delta}_{a}}{\overline{\eta}_{a}} - Px\frac{\delta_{a}}{\eta_{a}}\right) \\ &+ ((\overline{P}xx^{T})^{-1}P(xx^{T}) - I)\Big((Pxx^{T})^{-1} - (\overline{P}xx^{T})^{-1}\Big)P\Big(x\frac{\delta_{a}}{\eta_{a}}\Big) \\ &+ P\left[x\left(\frac{1}{\overline{\eta}_{a}} - \frac{1}{\eta_{a}}\right)(\mu_{Y,1a} - \overline{\mu}_{Y,1a})\right] + P\left[x\frac{1}{\eta_{a}}\mathbbm{1}\{z = 1_{a}\}\left(\frac{1}{\overline{\pi}_{1a}} - \frac{1}{\pi_{1a}}\right)(\mu_{Y,1a} - \overline{\mu}_{Y,1a})\right] \\ &- P\left[x\left(\frac{1}{\overline{\eta}_{a}} - \frac{1}{\eta_{a}}\right)(\mu_{Y,0a} - \overline{\mu}_{Y,0a})\right] - P\left[x\frac{1}{\eta_{a}}\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{0a}} - \frac{1}{\pi_{0a}}\right)(\mu_{D,1a} - \overline{\mu}_{D,1a})\right] \\ &- P\left[x\left(\frac{\overline{\delta}_{a}}{\overline{\eta}_{a}^{2}} - \frac{\delta_{a}}{\eta_{a}^{2}}\right)(\mu_{D,1a} - \overline{\mu}_{D,1a})\right] + P\left[x\frac{\delta_{a}}{\eta_{a}^{2}}\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{0a}} - \frac{1}{\pi_{1a}}\right)(\mu_{D,1a} - \overline{\mu}_{D,1a})\right] \\ &+ P\left[x\left(\frac{\overline{\delta}_{a}}{\overline{\eta}_{a}^{2}} - \frac{\delta_{a}}{\eta_{a}^{2}}\right)(\mu_{D,0a} - \overline{\mu}_{D,0a})\right] + P\left[x\frac{\delta_{a}}{\eta_{a}^{2}}\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{0a}} - \frac{1}{\pi_{1a}}\right)(\mu_{D,0a} - \overline{\mu}_{D,0a})\right] \\ &+ P\left[x\left(\frac{1}{\overline{\eta}_{a}} - \frac{1}{\eta_{a}}\right)\mathbbm{1}\{z = 1_{a}\}\left(\frac{1}{\overline{\pi}_{1a}} - \frac{1}{\pi_{0a}}\right)(\mu_{Y,1a} - \overline{\mu}_{Y,1a})\right] \\ &- P\left[x\left(\frac{1}{\overline{\eta}_{a}} - \frac{1}{\eta_{a}}\right)\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{1a}} - \frac{1}{\pi_{0a}}\right)(\mu_{Y,0a} - \overline{\mu}_{Y,0a})\right] \\ &- P\left[x\left(\frac{1}{\overline{\eta}_{a}} - \frac{1}{\eta_{a}}\right)\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{1a}} - \frac{1}{\pi_{0a}}\right)(\mu_{D,1a} - \overline{\mu}_{D,1a})\right] \\ &+ P\left[x\left(\frac{1}{\overline{\eta}_{a}} - \frac{1}{\eta_{a}^{2}}\right)\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{1a}} - \frac{1}{\pi_{0a}}\right)(\mu_{D,0a} - \overline{\mu}_{D,0a})\right] \\ &+ P\left[x\left(\frac{1}{\overline{\eta}_{a}^{2}} - \frac{\delta_{a}}{\eta_{a}^{2}}\right)\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{1a}} - \frac{1}{\pi_{0a}}\right)(\mu_{D,0a} - \overline{\mu}_{D,0a})\right] \\ &+ P\left[x\left(\frac{1}{\overline{\eta}_{a}^{2}} - \frac{\delta_{a}}{\eta_{a}^{2}}\right)\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{0a}} - \frac{1}{\pi_{0a}}\right)(\mu_{D,0a} - \overline{\mu}_{D,0a})\right] \\ &+ P\left[x\left(\frac{1}{\overline{\eta}_{a}^{2}} - \frac{\delta_{a}}{\eta_{a}^{2}}\right)\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{0a}} - \frac{1}{\pi_{0a}}\right)(\mu_{D,0a} - \overline{\mu}_{D,0a})\right] \\ &+ P\left[x\left(\frac{1}{\overline{\eta}_{a}^{2}} - \frac{\delta_{a}}{\eta_{a}^{2}}\right)\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{0a}} - \frac{1}{\pi_{0a}}\right)(\mu_{D,0a} - \overline{\mu}_{D,0a})\right] \\ &+ P\left[x\left(\frac{1}{\overline{\eta}_{a}^{2}} - \frac{\delta_{a}}{\eta_{a}^{2}}\right)\mathbbm{1}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{0a}} - \frac{1}{\pi_{0a}}\right)(\mu_{D,0a} - \overline{\mu}_{D,0a})$$

Similarly, for B, direct calculation yields

$$\begin{split} B &= (\overline{P}xx^{T})^{-1}P(xx^{T})(P(xx^{T})^{-1} - \overline{P}(xx^{T})^{-1})\left(\overline{P}x\frac{\overline{\delta}_{0} - \overline{\delta}_{a}}{\overline{\eta}_{b} - \eta_{a}} - Px\frac{\overline{\delta}_{b} - \overline{\delta}_{a}}{\overline{\eta}_{b} - \eta_{a}}\right) \\ &+ ((\overline{P}x^{T})^{-1}P(xx^{T}) - I)\left((Pxx^{T})^{-1} - (\overline{P}xx^{T})^{-1}\right)P\left(x\frac{\overline{\delta}_{b} - \overline{\delta}_{a}}{\overline{\eta}_{b} - \eta_{a}}\right) \\ &+ P\left[x\left(\frac{1}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{1}{\eta_{b} - \eta_{a}}\right)(\mu_{Y,1_{a}} - \overline{\mu}_{Y,1_{b}})\right] + P\left[x\frac{1}{\eta_{b} - \eta_{a}}\mathbb{I}\{z = 1_{b}\}\left(\frac{1}{\overline{\pi}_{1_{b}}} - \frac{1}{\pi_{1_{b}}}\right)(\mu_{Y,0_{b}} - \overline{\mu}_{Y,0_{b}})\right] \\ &- P\left[x\left(\frac{1}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{1}{\eta_{b} - \eta_{a}}\right)(\mu_{Y,0_{a}} - \overline{\mu}_{Y,1_{a}})\right] - P\left[x\frac{1}{\eta_{b} - \eta_{a}}\mathbb{I}\{z = 0_{b}\}\left(\frac{1}{\overline{\pi}_{1_{a}}} - \frac{1}{\pi_{1_{a}}}\right)(\mu_{Y,1_{a}} - \overline{\mu}_{Y,1_{a}})\right] \\ &+ P\left[x\left(\frac{1}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{1}{\eta_{b} - \eta_{a}}\right)(\mu_{Y,0_{a}} - \overline{\mu}_{Y,0_{a}})\right] + P\left[x\frac{1}{\eta_{b} - \eta_{a}}\mathbb{I}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{1_{a}}} - \frac{1}{\pi_{1_{a}}}\right)(\mu_{Y,1_{a}} - \overline{\mu}_{Y,1_{a}})\right] \\ &- P\left[x\left(\frac{1}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{1}{\eta_{b} - \eta_{a}}\right)(\mu_{Y,0_{a}} - \overline{\mu}_{Y,0_{a}})\right] + P\left[x\frac{1}{\eta_{b} - \eta_{a}}\mathbb{I}\{z = 0_{a}\}\left(\frac{1}{\overline{\pi}_{1_{a}}} - \frac{1}{\pi_{1_{a}}}\right)(\mu_{Y,0_{a}} - \overline{\mu}_{Y,0_{a}})\right] \\ &- P\left[x\left(\frac{1}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{1}{\eta_{b} - \eta_{a}}\right)(\mu_{Y,0_{a}} - \overline{\mu}_{Y,0_{a}})\right] + P\left[x\frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}}\mathbb{I}\{z = 0_{b}\}\left(\frac{1}{\overline{\pi}_{1_{a}}} - \frac{1}{\pi_{1_{a}}}\right)(\mu_{D,1_{a}} - \overline{\mu}_{Y,0_{a}})\right] \\ &+ P\left[x\left(\frac{\delta_{b} - \overline{\delta}_{a}}{\overline{\eta_{b} - \overline{\eta}_{a}} - \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}}\right)(\mu_{D,0_{a}} - \overline{\mu}_{D,0_{b}})\right] + P\left[x\frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}}\mathbb{I}\{z = 0_{b}\}\left(\frac{1}{\overline{\pi}_{1_{a}}} - \frac{1}{\pi_{1_{a}}}\right)(\mu_{D,1_{a}} - \overline{\mu}_{D,1_{b}})\right] \\ &+ P\left[x\left(\frac{\delta_{b} - \overline{\delta}_{a}}{\overline{\eta_{b} - \overline{\eta}_{a}} - \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}}\right)(\mu_{D,0_{a}} - \overline{\mu}_{D,0_{b}})\right] - P\left[x\frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}}\mathbb{I}\{z = 0_{b}\}\left(\frac{1}{\overline{\pi}_{1_{a}}} - \frac{1}{\pi_{0_{a}}}\right)(\mu_{D,0_{a}} - \overline{\mu}_{D,1_{b}}\right)\right] \\ &+ P\left[x\left(\frac{\delta_{b} - \overline{\delta}_{a}}{\overline{\eta_{b} - \overline{\eta}_{a}}}\right)\mathbb{I}\{z = 0_{b}\}\left(\frac{1}{\pi_{1_{a}}} - \frac{1}{\pi_{0}}}\right)(\mu_{D,0_{a}} - \overline{\mu}_{D,0_{b}}\right)\right] \\ &+ P\left[x\left(\frac{\delta_{b} - \overline{\delta}_{a}}{\overline{\eta_{b} - \eta_{a}}}\right)\mathbb{I}\{z = 0_{b}\}\left(\frac{1}{\overline{\pi}_{1_{a}}} - \frac{1}$$

Now we show how to bound $R(\widehat{P}_{n,k}, P_0)$.

$$\begin{split} & (\mathbb{P}_{n,k}xx^{T})^{-1}P_{0}(xx^{T})\Big((P_{0}xx^{T})^{-1} - \mathbb{P}_{n,k}(xx^{T})^{-1}\Big)\Bigg(\mathbb{P}_{n,k}x\frac{\widehat{\delta}_{a,k}}{\widehat{\eta}_{a,k}} - P_{0}x\frac{\delta_{a,0}}{\eta_{a,0}}\Bigg) \\ & \leq \|(\mathbb{P}_{n,k}xx^{T})^{-1}\|\|P_{0}(xx^{T})\|\Big\|\Big((P_{0}xx^{T})^{-1} - (\mathbb{P}_{n,k}xx^{T})^{-1}\Big)\Big\|\Big\|\Bigg(\mathbb{P}_{n,k}x\frac{\widehat{\delta}_{a,k}}{\widehat{\eta}_{a,k}} - P_{0}x\frac{\delta_{a,0}}{\eta_{a,0}}\Big)\Big\| \\ & \lesssim \Big\|\Big((Pxx^{T})^{-1} - (\mathbb{P}_{n,k}xx^{T})^{-1}\Big\|_{F}\Bigg\|\Bigg(\mathbb{P}_{n,k}x\frac{\widehat{\delta}_{a,k}}{\widehat{\eta}_{a,k}} - P_{0}x\frac{\delta_{a,0}}{\eta_{a,0}}\Bigg)\Bigg\| \\ & \lesssim O_{p}(1/\sqrt{n})o_{p}(1) \\ & = o_{p}(1/\sqrt{n}) \end{split}$$

Similarly, we can show that

$$((\mathbb{P}_{n,k}xx^T)^{-1}P_0(xx^T) - I)\Big((P_0xx^T)^{-1} - (\mathbb{P}_{n,k}xx^T)^{-1}\Big)P_0\Big(x\frac{\delta_{a,0}}{\eta_{a,0}}\Big) = o_p(1/\sqrt{n})$$

Also,

$$\begin{split} &P_0 \left[x \left(\frac{\widehat{\delta}_{a,k}}{\widehat{\eta}_{a,k}^2} - \frac{\delta_{a,0}}{\eta_{a,0}^2} \right) (\mu_{D,z,0} - \widehat{\mu}_{D,z,k}) \right] \\ \lesssim &P_0 \left[\left| \left(\frac{\widehat{\delta}_{a,k}}{\widehat{\eta}_{a,k}^2} - \frac{\delta_{a,0}}{\eta_{a,0}^2} \right) (\mu_{D,z,0} - \widehat{\mu}_{D,z,k}) \right| \right] \\ \lesssim &P_0 \left[|\widehat{\delta}_{a,k} - \delta_{a,0}| |\mu_{D,z,0} - \widehat{\mu}_{D,z,k}| \right] + P_0 \left[|\widehat{\eta}_{a,k} - \eta_{a,0}| |\mu_{D,z,0} - \widehat{\mu}_{D,z,k}| \right] \\ \leq & \left(\left\| \widehat{\delta}_{a,k} - \delta_{a,0} \right\|_{L^2(P_0)} + \left\| \widehat{\eta}_{a,k} - \eta_{a,0} \right\|_{L^2(P_0)} \right) \left\| \mu_{D,z,0} - \widehat{\mu}_{D,z,k} \right\| \\ = &o_p (1/\sqrt{n}) \end{split}$$

holds for $z \in \{0_a, 1_a, 0_b, 1_b\}$. Similarly, we can show

$$\begin{split} &P_0 \left[x \left(\frac{1}{\hat{\eta}_{a,k}} - \frac{1}{\eta_{a,0}} \right) (\mu_{Y,z,0} - \hat{\mu}_{Y,z,k}) \right] = o_p(1/\sqrt{n}) \\ &P_0 \left[x \frac{1}{\eta_{a,0}} \mathbbm{1}\{z = z'\} \left(\frac{1}{\hat{\pi}_{z,k}} - \frac{1}{\pi_{z',0}} \right) (\mu_{Y,z',0} - \hat{\mu}_{Y,z',k}) \right] = o_p(1/\sqrt{n}) \\ &P_0 \left[x \frac{\delta_{a,0}}{\eta_{a,0}^2} \mathbbm{1}\{z = z'\} \left(\frac{1}{\hat{\pi}_{z',k}} - \frac{1}{\pi_{z',0}} \right) (\mu_{D,z',0} - \hat{\mu}_{D,z',0}) \right] = o_p(1/\sqrt{n}) \\ &P_0 \left[x \left(\frac{1}{\hat{\eta}_{a,k}} - \frac{1}{\eta_{a,0}} \right) \mathbbm{1}\{z = z'\} \left(\frac{1}{\hat{\pi}_{z',k}} - \frac{1}{\pi_{z',0}} \right) (\mu_{Y,z',0} - \overline{\mu}_{Y,z',0}) \right] = o_p(1/\sqrt{n}) \\ &P_0 \left[x \left(\frac{\hat{\delta}_{a,k}}{\hat{\eta}_{a,k}^2} - \frac{\delta_{a,0}}{\eta_{a,0}^2} \right) \mathbbm{1}\{z = z'\} \left(\frac{1}{\hat{\pi}_{z',k}} - \frac{1}{\pi_{z',0}} \right) (\mu_{D,z',0} - \hat{\mu}_{D,z',0}) \right] = o_p(1/\sqrt{n}) \\ &P_0 \left[\frac{x \delta_{a,0}}{\eta_{a,0}^2 \hat{\eta}_{a,k}} \left[\delta_{a,0} (\hat{\eta}_{a,k} - \eta_{a,0})^2 + \eta_{a,0} (\hat{\eta}_{a,k} - \eta_{a,0}) (\delta_{a,0} - \hat{\delta}_{a,k}) \right] \right] = o_p(1/\sqrt{n}) \end{split}$$

To sum up, $A = o_p(1/\sqrt{n})$. Similarly, we can prove $B = o_p(1/\sqrt{n})$. Therefore, $R(\hat{P}_{n,k}, P_0) = o_p(1/\sqrt{n})$.

Following the same steps in Theorem 4, we know

$$\sqrt{n}(\widehat{\beta}_{os}^{(1)} - \beta_{P_0}^{(1)}) = \mathbb{P}_n \phi_{P_0}^{(1)} + o_p(1)$$

Therefore,

$$\sqrt{n}(\widehat{\beta}_{os}^{(1)} - \beta_{P_0}) \rightsquigarrow \mathcal{N}\left(0, \ \mathbb{E}_{P_0}\left[\phi_{P_0}^{(j)T}\phi_{P_0}^{(j)}\right]\right),$$

and

$$W^{(j)} \rightsquigarrow \chi^2(d_x)$$

Therefore, $T_{Wald}^{(j)}$ is an asymptotically size- α test.

S1.8 Proof of Lemma 1

Proof. \Rightarrow is obvious. We only need to show \Leftarrow .

For \Leftarrow , it is equivalent to show $A \Rightarrow B$, where A is

There exists an x^* , such that $\xi_{P_0}(x^*) > 0$.

and \boldsymbol{B} is

There exists an
$$c$$
, such that $\Omega_{P_0}(c) > 0$.

We prove this by contradiction. That is, the statement that there exists an x^* , such that $\xi_{P_0}(x^*) > 0$ contradicts with the statement that $\Omega_{P_0}(c) = 0$ for all c.

Let $x^* = (x_1^*, ..., x_{d_1}^*, ..., x_{d_1+d_2}^*)$ such that $x^* \in \mathcal{X}_1 \times \mathcal{X}_2$. Then there exists a set, and $C := (x_1^*, ..., x_{d_1}^*, ..., (a_{d_1+1}, b_{d_1+1}), ..., (a_{d_1+d_2}, b_{d_1+d_2}))$ such that

$$P_0(C) > 0$$

 $|a_i < x_i^* < b_i|, |b_i - a_i|$ small enough such that $\xi_{P_0}(x) > 0$ for $x \in C$. We assume $(-\infty, x_1^*] \times ... \times (-\infty, x_{d_1}^*] \cap \mathcal{X}_1$ is not empty, the case when it is empty can be handled similarly. Then there must exist a point $(a_1, ..., a_{d_1}) \in \mathcal{X}_1$ such that $\mathcal{X}_1 \cap (a_1, x_1^*] \times ... (a_{d_1}, x_{d_1}^*] = (x_1^*, ..., x_{d_1}^*)$. Then

$$\int \xi_{P_0}(x) \mathbb{1}\{x \in (a_1, x_1^*] \times \dots \times (a_{d_1}, x_{d_1}^*] \times (a_{d_1+1}, b_{d_1+1}) \times \dots \times (a_{d_2}, b_{d_2})\} dP_0(x) > 0$$

Since X_2 is absolutely continuous, then

$$\int \theta_{P_0}(x) 1\{x \in (a_1, x_1^*] \times \dots \times (a_{d_1}, x_{d_1}^*] \times (a_{d_1+1}, b_{d_1+1}] \times \dots \times (a_{d_2}, b_{d_2}]\} dP_0(x) > 0$$
(4)

Let $b_i = x_i^*$ for $i \leq d_1$, define

$$C = \{c = (c_1, ..., c_n) : c_i \in \{a_i, b_i\}, i = 1, ..., d_1 + d_2\}$$

and let $n(c) = \#\{i : c_i = a_i\}$. Then if $\Omega_{P_0}(c) = 0$ for all $c \in \mathbb{R}^{d_x}$,

$$0 = \sum_{c \in C} (-1)^{n(c)} \Omega(c)$$

= $\int \xi_{P_0}(x) \mathbf{1}_{(a_1, b_1](x) \times \cdots (a_{d_1+d_2}, b_{d_1+d_2}]} dP_0(x),$

which contradicts with (4).

S1.9 Proof of Proposition 3

Proof. We will pick $\Omega_P^{(j)}(c)$ as example. We pick a one-dimensional parametric submodel $\{P_t : t \in [0, \epsilon)\}$, our goal is to calculate

$$\frac{d\Omega_{P_t}^{(1)}(c)}{dt}\Big|_{t=0} = \frac{d\mathbb{E}_{P_t}[\theta_{ACO,P_t}(X)\mathbb{1}_c(X)]}{dt}\Big|_{t=0} - \frac{d\mathbb{E}_{P_t}[\theta_{SW,P_t}(X)\mathbb{1}_c(X)]}{dt}\Big|_{t=0}$$

Similar to the proof of theorem 5, one can show that

$$\frac{d\mathbb{E}_{P_t}[\theta_{ACO,P_t}(X) - \mathbb{1}_c(X)]}{dt}\Big|_{t=0}$$
$$= \mathbb{E}_{P_0}\left[(\theta_{ACO,P_t}(X) - \mathbb{E}_{P_0}[\theta_{ACO,P_t}(X)] + D_P^{(1)})\mathbb{1}_c(X)S(O)\right]$$

and

$$\frac{d\mathbb{E}_{P_t}[\theta_{SW,P_t}(X) - \mathbb{1}_c(X)]}{dt}\Big|_{t=0}$$
$$= \mathbb{E}_{P_0}\left[(\theta_{SW,P_t}(X) - \mathbb{E}_{P_0}[\theta_{SW,P_t}(X)] + D_P^{(2)})\mathbb{1}_c(X)S(O) \right]$$

so that

$$D_P^{(1)*}(c) = o \mapsto D_P^{(1)}(z, x, d, y) \mathbb{1}_c(x) - D_P^{(2)}(g, z, x, d, y) \mathbb{1}_c(x) + \theta_P^{(1)}(x) \mathbb{1}_c(x) - \Omega_P^{(1)}(c)$$

S1.10 Proof of Theorem 7

Suppose that

- 1. There exists $0 < \epsilon < 0.5$ such that $\epsilon < P_0(Z|X) < 1 \epsilon$, $\epsilon < P_0(D|Z,X) < 1 \epsilon$, $\epsilon < \widehat{P}_{n,k}(Z|X) < 1 \epsilon$, $\epsilon < \widehat{P}_{n,k}(D|Z,X) < 1 \epsilon$ (k = 1,...K), there is a universal constant C such that $|\mathbb{E}_{P_0}[Y|Z,X]| \leq C$ and $|\mathbb{E}_{\widehat{P}_{n,k}}[Y|Z,X]| \leq C$ with probability one.
- 2. $\{D_{P_0}^{(j)*}(c) : c \in \mathcal{X}\}$ is P_0 -Donsker for $j \in \{1, 2, 3\}$.
- 3. Rate conditions for nuisance functions estimation: The convergence rates for all nuisance function estimators are faster then $n^{-1/4}$:

$$\begin{aligned} \left\| \pi_{P_0}(z, \cdot) - \pi_{\widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(n^{-1/4}) \\ \left\| \mu_{Y,P_0}(z, \cdot) - \mu_{Y,\widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(n^{-1/4}) \\ \left\| \mu_{D,P_0}(z, \cdot) - \mu_{D,\widehat{P}_{n,k}}(z, \cdot) \right\|_{L^2(P_0)} &= o_p(n^{-1/4}) \end{aligned}$$

for
$$z \in \{0_a, 1_a, 0_b, 1_b\}, k \in \{1, ..., K\}.$$

Proof. First we will establish asymptotic linearity of $\widehat{\Omega}_{os,k}^{(1)}(c)$ for a given c, so that we can establish asymptotic linearity of $\widehat{\Omega}_{os}^{(1)}(c)$.

We firstly define

$$R(\overline{P}, P; c) = \Omega_{\overline{P}}^{(1)}(c) - \Omega_{P}^{(1)}(c) + PD_{\overline{P}}^{(1)*}(c).$$

Similar to the proof of theorem 6, $R(\overline{P}, P; c) = A + B$, where

$$\begin{split} A =& P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}}\right)(\mu_{Y,1_{a}}-\overline{\mu}_{Y,1_{a}})\right] + P\left[\mathbbm{1}_{c}\frac{1}{\eta_{a}}\mathbbm{1}\{z=1_{a}\}\left(\frac{1}{\overline{\pi}_{1_{a}}}-\frac{1}{\pi_{1_{a}}}\right)(\mu_{Y,1_{a}}-\overline{\mu}_{Y,1_{a}})\right] \\ &- P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}}\right)(\mu_{Y,0_{a}}-\overline{\mu}_{Y,0_{a}})\right] - P\left[\mathbbm{1}_{c}\frac{1}{\eta_{a}}\mathbbm{1}\{z=0_{a}\}\left(\frac{1}{\overline{\pi}_{0_{a}}}-\frac{1}{\pi_{0_{a}}}\right)(\mu_{Y,0_{a}}-\overline{\mu}_{Y,0_{a}})\right] \\ &- P\left[\mathbbm{1}_{c}\left(\frac{\overline{\delta}_{a}}{\overline{\eta}_{a}^{2}}-\frac{\delta_{a}}{\eta_{a}^{2}}\right)(\mu_{D,1_{a}}-\overline{\mu}_{D,1_{a}})\right] + P\left[\mathbbm{1}_{c}\frac{\delta_{a}}{\eta_{a}^{2}}\mathbbm{1}\{z=1_{a}\}\left(\frac{1}{\overline{\pi}_{1_{a}}}-\frac{1}{\pi_{1_{a}}}\right)(\mu_{D,1_{a}}-\overline{\mu}_{D,1_{a}})\right] \\ &+ P\left[\mathbbm{1}_{c}\left(\frac{\overline{\delta}_{a}}{\overline{\eta}_{a}^{2}}-\frac{\delta_{a}}{\eta_{a}^{2}}\right)(\mu_{D,0_{a}}-\overline{\mu}_{D,0_{a}})\right] + P\left[\mathbbm{1}_{c}\frac{\delta_{a}}{\eta_{a}^{2}}\mathbbm{1}\{z=0_{a}\}\left(\frac{1}{\overline{\pi}_{0_{a}}}-\frac{1}{\pi_{1_{a}}}\right)(\mu_{D,0_{a}}-\overline{\mu}_{D,0_{a}})\right] \\ &+ P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}}\right)\mathbbm{1}\{z=1_{a}\}\left(\frac{1}{\overline{\pi}_{1_{a}}}-\frac{1}{\pi_{1_{a}}}\right)(\mu_{Y,1_{a}}-\overline{\mu}_{Y,1_{a}})\right] \\ &- P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}}\right)\mathbbm{1}\{z=0_{a}\}\left(\frac{1}{\overline{\pi}_{0_{a}}}-\frac{1}{\pi_{0_{a}}}\right)(\mu_{Y,0_{a}}-\overline{\mu}_{Y,0_{a}})\right] \\ &- P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}}\right)\mathbbm{1}\{z=0_{a}\}\left(\frac{1}{\overline{\pi}_{1_{a}}}-\frac{1}{\pi_{0_{a}}}\right)(\mu_{D,1_{a}}-\overline{\mu}_{D,1_{a}})\right] \\ &+ P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}^{2}}\right)\mathbbm{1}\{z=0_{a}\}\left(\frac{1}{\overline{\pi}_{1_{a}}}-\frac{1}{\pi_{0_{a}}}\right)(\mu_{D,0_{a}}-\overline{\mu}_{D,0_{a}})\right] \\ &+ P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}^{2}}\right)\mathbbm{1}\{z=0_{a}\}\left(\frac{1}{\overline{\pi}_{0_{a}}}-\frac{1}{\pi_{0_{a}}}\right)(\mu_{D,0_{a}}-\overline{\mu}_{D,0_{a}})\right] \\ &+ P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}^{2}}\right)\mathbbm{1}\{z=0_{a}\}\left(\frac{1}{\overline{\pi}_{0_{a}}}-\frac{1}{\pi_{0_{a}}}\right)(\mu_{D,0_{a}}-\overline{\mu}_{D,0_{a}})\right] \\ &+ P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}^{2}}\right)\mathbbm{1}\{z=0_{a}\}\left(\frac{1}{\overline{\pi}_{0_{a}}}-\frac{1}{\pi_{0_{a}}}\right)(\mu_{D,0_{a}}-\overline{\mu}_{D,0_{a}})\right] \\ &+ P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}^{2}}\right)\mathbbm{1}\{z=0_{a}\}\left(\frac{1}{\overline{\pi}_{0_{a}}}-\frac{1}{\pi_{0_{a}}}\right)(\mu_{D,0_{a}}-\overline{\mu}_{D,0_{a}})\right] \\ &+ P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}^{2}}\right)\mathbbm{1}\{z=0_{a}\}\left(\frac{1}{\overline{\pi}_{0_{a}}}-\frac{1}{\overline{\pi}_{0_{a}}}\right)(\mu_{D,0_{a}}-\overline{\mu}_{D,0_{a}})\right) \\ &+ P\left[\mathbbm{1}_{c}\left(\frac{1}{\overline{\eta}_{a}}-\frac{1}{\eta_{a}^{2$$

Similarly, for B, direct calculation yields

$$\begin{split} B = & P \left[\mathbbm{1}_{c} \left(\frac{1}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{1}{\eta_{b} - \eta_{a}} \right) (\mu_{Y,1b} - \overline{\mu}_{Y,1b}) \right] + P \left[\mathbbm{1}_{c} \frac{1}{\eta_{b} - \eta_{a}} \mathbbm{1}_{z} \mathbbm{1}_{z} \mathbbm{1}_{b} \left(\frac{1}{\overline{\pi}_{1b}} - \frac{1}{\pi_{1b}} \right) (\mu_{Y,1b} - \overline{\mu}_{Y,1b}) \right] \\ & - P \left[\mathbbm{1}_{c} \left(\frac{1}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{1}{\eta_{b} - \eta_{a}} \right) (\mu_{Y,0c} - \overline{\mu}_{Y,0c}) \right] - P \left[\mathbbm{1}_{c} \frac{1}{\eta_{b} - \eta_{a}} \mathbbm{1}_{z} \mathbbm{1}_{a} \mathbbm{1}_{z} \mathbbm{1}_{a} \right) \left(\frac{1}{\overline{\pi}_{1a}} - \frac{1}{\pi_{1a}} \right) (\mu_{Y,1c} - \overline{\mu}_{Y,1c}) \right] \\ & - P \left[\mathbbm{1}_{c} \left(\frac{1}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{1}{\eta_{b} - \eta_{a}} \right) (\mu_{Y,1c} - \overline{\mu}_{Y,1c}) \right] + P \left[\mathbbm{1}_{c} \frac{1}{\eta_{b} - \eta_{a}} \mathbbm{1}_{z} \mathbbm{1}_{a} \mathbbm{1}_{z} \mathbbm{1}_{a} \right) \left(\frac{1}{\overline{\pi}_{1a}} - \frac{1}{\pi_{1a}} \right) (\mu_{Y,1c} - \overline{\mu}_{Y,1c}) \right] \\ & + P \left[\mathbbm{1}_{c} \left(\frac{1}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}} \right) (\mu_{D,1c} - \overline{\mu}_{D,1c}) \right] + P \left[\mathbbm{1}_{c} \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}} \mathbbm{1}_{z} \mathbbm{1}_{a} \mathbbm{1}_{a} \mathbbm{1}_{a} \mathbbm{1}_{a} \right) (\mu_{D,1c} - \overline{\mu}_{D,1c}) \right] \\ & + P \left[\mathbbm{1}_{c} \left(\frac{\delta_{b} - \delta_{a}}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}} \right) (\mu_{D,0c} - \overline{\mu}_{D,0c}) \right] + P \left[\mathbbm{1}_{c} \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}} \mathbbm{1}_{z} \mathbbm{1}_{a} \mathbbm{1}_{a} \mathbbm{1}_{a} \right) (\mu_{D,0c} - \overline{\mu}_{D,0c}) \right] \\ & + P \left[\mathbbm{1}_{c} \left(\frac{\delta_{b} - \delta_{a}}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}} \right) (\mu_{D,0c} - \overline{\mu}_{D,0c}) \right] + P \left[\mathbbm{1}_{c} \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}} \mathbbm{1}_{z} \mathbbm{1}_{z} \mathbbm{1}_{z} \right] \left(\frac{1}{\overline{\pi}_{1a}} - \frac{1}{\pi_{1a}} \right) (\mu_{D,0c} - \overline{\mu}_{D,0c}) \right] \\ & - P \left[\mathbbm{1}_{c} \left(\frac{\delta_{b} - \delta_{a}}{\overline{\eta}_{b} - \overline{\eta}_{a}} - \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}} \right) (\mu_{D,0c} - \overline{\mu}_{D,0c}) \right] - P \left[\mathbbm{1}_{c} \frac{\delta_{b} - \delta_{a}}{\eta_{b} - \eta_{a}} \mathbbm{1}_{z} \mathbbm{1}_{z} \right] (\mu_{D,0c} - \overline{\mu}_{D,0c}) \right] \\ & + P \left[\mathbbm{1}_{c} \left(\frac{\delta_{b} - \delta_{a}}{\overline{\eta}_{b} - \overline{\eta}_{a}} \right) \mathbbm{1}_{z} \mathbbm{1}_{z} \mathbbm{1}_{z} \left(\frac{1}{\pi_{1a}} - \frac{1}{\pi_{1a}} \right) (\mu_{D,1c} - \overline{\mu}_{D,1c}) \right] \\ & + P \left[\mathbbm{1}_{c} \left(\frac{\delta_{b} - \delta_{a}}{\overline{\eta}_{b} - \eta_{a}} \right) \mathbbm{1}_{z} \mathbbm{1}_{z} \left(\frac{1}{\overline{\pi}_{1a}} - \frac{1}{\pi_{1a}} \right) (\mu_{D,0c} - \overline{\mu}_{D,0c}) \right] \\ & + P \left[\mathbbm{$$

By similar argument in Theorem 6 and $\mathbb{1}_c \leq 1$, we know $\sup_c |R(\hat{P}_{n,k}, P_0)(c)| = o_p(1/\sqrt{n})$. Now we consider the empirical process term

$$\mathbb{G}_{n,k}(D_{\widehat{P}_{n,k}}^{(1)*}(c) - D_{P_0}^{(1)*}(c)) = \sqrt{n_k}(\mathbb{P}_{n,k} - P_0)(D_{\widehat{P}_{n,k}}^{(1)*}(c) - D_{P_0}^{(1)*}(c))$$

Define $\mathcal{F} := \{D_{\widehat{P}_{n,k}}^{(1)*}(c) - D_{P_0}^{(1)*}(c) : c \in \mathcal{X}\}$, then $\sup_c |\mathbb{G}_{n,k}(D_{\widehat{P}_{n,k}}^{(1)*}(c) - D_{P_0}^{(1)*}(c))|$ can be written as $\sup_{f \in \mathcal{F}} |\mathbb{G}_{n,k}f|$. To show the supreme of this empirical process is $o_p(1)$, we will bound

$$\mathbb{E}_{P_0}\left[\sup_{f\in\mathcal{F}_{n,k}}\left|\mathbb{G}_{n,k}f\right|\right] = \mathbb{E}_{P_0}\left[\mathbb{E}_{P_0}\left[\sup_{f\in\mathcal{F}_{n,k}}\left|\mathbb{G}_{n,k}f\right|\middle|O_i, i\in I_k\right]\right].$$

After condition on $\{O_i, i \in I_k^c\}$, the estimated nuisance functions can be treated as fixed. Let $F_{n,k}(o)$ be an envelop function of $\mathcal{F}_{n,k}$. Then by Theorem 2.14.1 of van der Vaart and Wellner (2023). It holds that

$$\mathbb{E}_{P_0}\left[\sup_{f\in\mathcal{F}_{n,k}} |\mathbb{G}_{n,k}f| \left| O_i, i\in I_k \right] \lesssim \sqrt{\mathbb{E}_{P_0}\left[F_{n,k}(O)^2 \left| O_i, i\in I_k^c \right] J(1,\mathcal{F}_{n,k}|F_{n,k},L^2) \right]} \right]$$

where $J(1, \mathcal{F}_{n,k}|F_{n,k}, L^2)$ is the uniform entropy integral. Our goal is to show that

- $J(1, \mathcal{F}_{n,k} | F_{n,k}, L^2) = O(1)$
- There exists an $F_{n,k}(o)$ such that $\sqrt{\mathbb{E}_{P_0}\left[F_{n,k}(O)^2 \middle| O_i, i \in I_k^c\right]} = o(1).$

To show $J(1, \mathcal{F}_{n,k} | F_{n,k}, L^2) = O(1)$, we notice that $\mathcal{F}_{n,k}$ is a linear combination of elements from the following three class

$$\mathcal{A}_{1} := \{ [D_{\hat{P}_{n,k}}^{(1)*}(o) - D_{P_{0}}^{(1)*}(o)] \mathbb{1}_{c} : c \in \mathcal{C} \}$$
$$\mathcal{A}_{2} := \left\{ \left[\frac{\delta_{a,\hat{P}_{n,k}}}{\eta_{a,\hat{P}_{n,k}}} - \frac{\delta_{a,P_{0}}}{\eta_{a,P_{0}}} \right] \mathbb{1}_{c} : c \in \mathcal{C} \right\}$$
$$\mathcal{A}_{3} := \left\{ \left[\mathbb{P}_{n,k} \frac{\delta_{a,\hat{P}_{n,k}}}{\eta_{a,\hat{P}_{n,k}}} \mathbb{1}_{c} - P_{0} \frac{\delta_{a,P_{0}}}{\eta_{a,P_{0}}} \mathbb{1}_{c} \right] : c \in \mathcal{C} \right\}$$

Since $\mathcal{A}_4 := \{\mathbb{1}_c : c \in \mathcal{X}\}$ is a VC class with VC index $d_x + 1$, by Theorem 2.6.7 of van der Vaart and Wellner (2023), the covering number satisfies

$$N(\epsilon \|F\|_{Q,2}, \mathcal{A}_4, L^2(Q)) \le K\left(\frac{1}{\epsilon}\right)^{2d_x}$$

By the similar reasoning from the proof of Lemma 2 in Westling (2022), We know $J(1, \mathcal{F}_{n,k} | F_{n,k}, L^2) = O(1)$.

Next we derive an envelope function $F_{n,k}$, and verify it satisfies $\sqrt{\mathbb{E}_{P_0}\left[F_{n,k}(O)^2 \left| O_i, i \in I_k^c \right]\right]} = o(1).$

$$D_{\hat{P}_{n,k}}^{(1)*} \mathbb{1}_{c} - D_{P_{0}}^{(1)*}(c) = \underbrace{\left[D_{\hat{P}_{n,k}}^{(1)} \mathbb{1}_{c} + \theta_{ACO,\hat{P}_{n,k}} \mathbb{1}_{c} - \int \mathbb{1}_{c} \theta_{ACO,\hat{P}_{n,k}} dP_{n,k}\right] - \left[D_{P_{0}}^{(1)} \mathbb{1}_{c} + \theta_{ACO,P_{0}} - \int \mathbb{1}_{c} \theta_{ACO,P_{0}} dP_{0}\right]}_{A} + \underbrace{\left[D_{\hat{P}_{n,k}}^{(2)} \mathbb{1}_{c} + \theta_{SW,\hat{P}_{n,k}} - \int \mathbb{1}_{c} \theta_{SW,\hat{P}_{n,k}} dP_{n,k}\right] - \left[D_{P_{0}}^{(2)} \mathbb{1}_{c} + \theta_{SW,P_{0}} - \int \mathbb{1}_{c} \theta_{SW,P_{0}} dP_{0}\right]}_{B}}_{B}$$

We firstly consider term A. For $z' \in \{o_a, 1_a, 0_b, 1_b\}$,

$$\begin{split} & \left| \frac{\mathbbm{1}_c}{\eta_{a,\hat{P}_{n,k}}} \left[\frac{\mathbbm{1}\{z=z'\}}{\hat{\pi}_{z',k}} [y - \hat{\mu}_{Y,z',k}] \right] - \frac{\mathbbm{1}_c}{\eta_{a,P_0}} \left[\frac{\mathbbm{1}\{z=z'\}}{\pi_{z',0}} [y - \mu_{Y,z',0}] \right] \right| \\ & \leq \left| \left(\frac{1}{\eta_{a,\hat{P}_{n,k}}} - \frac{1}{\eta_{a,P_0}} \right) \left[\frac{\mathbbm{1}\{z=z'\}}{\hat{\pi}_{z',k}} [y - \hat{\mu}_{Y,z',k}] \right] + \frac{1}{\eta_{a,P_0}} \left[\mathbbm{1}\{z=z'\} \left(\frac{1}{\hat{\pi}_{z',k}} - \frac{1}{\pi_{z',0}} \right) [y - \hat{\mu}_{Y,z',k}] \right] \right| \\ & + \frac{1}{\eta_{a,P_0}} \left[\mathbbm{1}\{z=z'\} \frac{1}{\pi_{z',0}} [(\mu_{Y,z',0} - \hat{\mu}_{Y,z',k})] \right] \right| \\ & \leq C \left| \eta_{a,\hat{P}_{n,k}} - \eta_{a,P_0} \right| (|y| + C) + C \left| \hat{\pi}_{z',k} - \pi_{z',0} \right| (|y| + C) + C (|\hat{\mu}_{Y,z',k} - \mu_{Y,z',0}|) \\ & \leq C \left| \max_{z \in \{0_a,1_a\}} \left| \hat{\mu}_{D,z,k} - \mu_{D,z,0} \right| (|y| + C) + C \left| \hat{\pi}_{z',k} - \pi_{z',0} \right| (|y| + C) + C |\hat{\mu}_{Y,z',k} - \mu_{Y,z',0}| \end{split}$$

Similarly,

$$\begin{aligned} &\left|\frac{\mathbb{1}_{c}\delta_{a,\widehat{P}_{n,k}}}{\eta_{a,\widehat{P}_{n,k}}^{2}} \left[\frac{\mathbb{1}\{z=z'\}}{\widehat{\pi}_{z',k}} [d-\widehat{\mu}_{D,z',k}]\right] - \frac{\mathbb{1}_{c}\delta_{a,P_{0}}}{\eta_{a,P_{0}}^{2}} \left[\frac{\mathbb{1}\{z=z'\}}{\pi_{z',0}} [d-\mu_{D,z',0}]\right]\right| \\ \leq & C \max_{z\in\{0_{a},1_{a}\},B\in\{Y,D\}} \left|\widehat{\mu}_{B,z,k} - \mu_{B,z,0}\right| + C \left|\widehat{\pi}_{z',k} - \pi_{z',0}\right| + C |\widehat{\mu}_{D,z',k} - \mu_{D,z',0}| \end{aligned}$$

Furthermore, we have

$$\begin{aligned} &|\theta_{ACO,\hat{P}_{n,k}} \mathbb{1}_{c} - \theta_{ACO,P_{0}} \mathbb{1}_{c}| \leq C \max_{z \in \{0_{a},1_{a}\}, B \in \{Y,D\}} \left| \hat{\mu}_{B,z,k} - \mu_{B,z,0} \right| \\ &\int \mathbb{1}_{c} \theta_{ACO,\hat{P}_{n,k}} dP_{n,k} - \int \mathbb{1}_{c} \theta_{ACO,P_{0}} dP_{0} \leq \sup_{c \in \mathcal{C}} \left| (\mathbb{P}_{n,k} - P_{0}) \mathbb{1}_{c} \theta_{ACO,\hat{P}_{n,k}} \right| + C \max_{z \in \{0_{a},1_{a}\}, B \in \{Y,D\}} P_{0} \left| \hat{\mu}_{B,z,k} - \mu_{B,z,0} \right| \end{aligned}$$

Therefore,

$$\begin{split} |A| \leq & \left| D_{\hat{P}_{n,k}}^{(1)} \mathbbm{1}_{c} - D_{P_{0}}^{(1)} \mathbbm{1}_{c} \right| + \left| \theta_{ACO,\hat{P}_{n,k}} \mathbbm{1}_{c} - \theta_{ACO,P_{0}} \mathbbm{1}_{c} \right| + \left| \int \mathbbm{1}_{c} \theta_{ACO,\hat{P}_{n,k}} d\mathbbm{1}_{n,k} - \int \mathbbm{1}_{c} \theta_{ACO,P_{0}} dP_{0} \right] \\ \leq & C \max_{z \in \{0_{a},1_{a}\}, B \in \{Y,D\}} \left| \hat{\mu}_{B,z,k} - \mu_{B,z,0} \right| (|y| + C) + C \sup_{z \in \{0_{a},1_{a}\}} \left| \hat{\pi}_{z,k} - \pi_{z,0} \right| (|y| + C) \\ & + \sup_{c \in \mathcal{C}} \left| (\mathbbm{1}_{n,k} - P_{0}) \mathbbm{1}_{c} \theta_{ACO,\hat{P}_{n,k}} \right| + C \max_{z \in \{0_{a},1_{a}\}, B \in \{Y,D\}} P_{0} \left| \hat{\mu}_{B,z,k} - \mu_{B,z,0} \right| \end{split}$$

We can obtain a similar bound for B, to sum up, we can take

$$\begin{split} F_{n,k} \\ = & C \max_{z \in \{0_a, 1_a, 0_b, 1_b\}, B \in \{Y, D\}} \left| \hat{\mu}_{B, z, k} - \mu_{B, z, 0} \right| (|y| + C) + C \sup_{z \in \{0_a, 1_a, 0_b, 1_b\}} \left| \hat{\pi}_{z, k} - \pi_{z, 0} \right| (|y| + C) \\ & + \sup_{c \in \mathcal{C}} |(\mathbb{P}_{n, k} - P_0) \mathbb{1}_c \theta_{SW, \widehat{P}_{n, k}}| + \sup_{c \in \mathcal{C}} |(\mathbb{P}_{n, k} - P_0) \mathbb{1}_c \theta_{ACO, \widehat{P}_{n, k}}| \\ & + C \max_{z \in \{0_a, 1_a, 0_b, 1_b\}, B \in \{Y, D\}} P_0 \Big| \hat{\mu}_{B, z, k} - \mu_{B, z, 0} \Big| \end{split}$$

and our assumption equipped with triangle inequality implies that $||F_{n,k}||_{P_{0,2}} \xrightarrow{P_{0}} 0$, uniform boundedness of $||F_{n,k}||_{P_{0,2}}$ implies $E[||F_{n,k}||_{P_{0,2}}] \to 0$ therefore,

$$\begin{split} & \mathbb{E}_{P_0} \left[\sup_{f \in \mathcal{F}_{n,k}} |\mathbb{G}_{n,k}f| \right] \\ = & \mathbb{E}_{P_0} \left[\mathbb{E}_{P_0} \left[\sup_{f \in \mathcal{F}_{n,k}} |\mathbb{G}_{n,k}f| \Big| O_i, i \in I_k \right] \right] \\ & \lesssim & \mathbb{E}_{P_0} \left\{ \sqrt{\mathbb{E}_{P_0} \left[F_{n,k}(O)^2 \Big| O_i, i \in I_k^c \right]} \right\} J(1, \mathcal{F}_{n,k}|F_{n,k}, L^2) \\ = & o(1)O(1) = o(1) \end{split}$$

To sum up, we have already shown that

$$\max_{k} \sup_{c} |\mathbb{G}_{n,k}(D_{\widehat{P}_{n,k}}^{(1)*}(c) - D_{P_0}^{(1)*}(c))| = o_p(1)$$

and the one-step estimator has the following expansion:

$$\widehat{\Omega}_{os}^{(1)}(c) - \Omega_{P_0}^{(1)}(c) = \frac{1}{n} \sum_{i=1}^n D_{P_0}^{(1)*}(c)(O_i) + r_n(c).$$

with $\sup_{c \in \mathcal{X}} |r_n(c)| = o_p(1/\sqrt{n}).$

The rest of the proof is similar to that of Westling (2022).

S2 Additional simulation results




Figure S.1: Sampling distribution of estimating equation estimator when the outcome is binary. Simulations are repeated 1000 times. Datasets are generating with $\beta = (0, 1, -1)$. Four proportions of SW are considered. The dashed red lines represents the ground truth SWATE in each setting.




Figure S.2: Sampling distribution of one-step estimator when the outcome is continuous. Simulations are repeated 1000 times. Datasets are generating with $\beta = (0, 1, -1)$. Four proportions of SW are considered. The dashed red lines represents the ground truth SWATE in each setting.




Figure S.3: Sampling distribution of one-step estimator when the outcome is binary. Simulations are repeated 1000 times. Datasets are generating with $\beta = (0, 1, -1)$. Four proportions of SW are considered. The dashed red lines represents the ground truth SWATE in each setting.

β	SWI $\%$	SWATE	Sample size	Estimate	Relative bias	Bias	Se est	Coverage	Acceptance rate
			1000	0.383	0.955	3.704	17.015	0.912	0.999
			2000	-0.070	-0.328	-1.272	4.664	0.956	1.000
	11%	0.257	5000	0.246	-0.012	-0.045	0.337	0.966	1.000
			10000	0.247	-0.011	-0.043	0.171	0.949	1.000
			1000	0.192	-0.048	-0.200	1.232	0.937	1.000
	220	0.044	2000	0.236	-0.005	-0.021	0.304	0.963	1.000
	22%	0.241	5000	0.248	0.008	0.032	0.136	0.962	1.000
			10000	0.247	0.006	0.026	0.087	0.944	1.000
(0,1-1)			1000	0.279	0.067	0.313	0.395	0.939	1.000
	220	0.010	2000	0.226	0.014	0.064	0.178	0.976	1.000
	32%	0.212	5000	0.210	-0.003	-0.012	0.095	0.963	1.000
			10000	0.213	0.001	0.003	0.063	0.964	1.000
		0.208	1000	0.224	0.016	0.075	0.121	0.934	1.000
	66%		2000	0.204	-0.004	-0.018	0.073	0.951	1.000
			5000	0.206	-0.003	-0.012	0.043	0.956	1.000
			10000	0.205	-0.003	-0.016	0.029	0.956	1.000
			1000	1.091	0.531	1.013	20.669	0.922	0.998
			2000	0.576	0.051	0.097	2.751	0.935	1.000
	11%	0.524	5000	0.581	0.057	0.109	0.321	0.961	1.000
			10000	0.541	0.016	0.031	0.174	0.952	1.000
		0.518	1000	0.805	0.287	0.555	1.293	0.924	1.000
			2000	0.555	-1.261	-2.434	0.311	0.971	0.999
	22%		5000	0.530	0.012	0.023	0.140	0.968	1.000
			10000	0.521	0.003	0.006	0.088	0.940	1.000
(0,2,-3)		0.511	1000	0.510	-0.002	-0.003	0.376	0.960	1.000
			2000	0.520	0.009	0.018	0.188	0.970	1.000
	32%		5000	0.512	0.001	0.002	0.098	0.974	1.000
			10000	0.512	0.001	0.001	0.064	0.968	1.000
			1000	0.522	0.012	0.023	0.126	0.935	1.000
			2000	0.513	0.003	0.006	0.077	0.950	1.000
	66%	0.510	5000	0.512	0.001	0.003	0.044	0.955	1.000
			10000	0.510	0.000	-0.001	0.030	0.954	1.000

Table S5: Simulation results for $\widehat{\psi}_{ee}$ when the outcome is binary

¹ True values of SWATE under each setting.
 ² Truncated means of point estimates. Point estimates are truncated if absolute value are bigger than 500.
 ³ 90% Winsorized mean of standard error estimates.

⁴ Proportion of point estimates which are not truncated.

β	SWI $\%$	SWATE	Sample size	Estimate	Relative bias	Bias	Se est	Coverage	Acceptance rate
			1000	-3.122	-4.040	-4.402	43.614	0.863	0.947
	~		2000	-5.087	-6.004	-6.543	10.774	0.864	0.976
	11%	0.917	5000	0.744	-0.174	-0.190	1.018	0.928	0.997
			10000	0.902	-0.015	-0.017	0.546	0.932	1.000
			1000	-7.244	-8.263	-8.109	4.852	0.849	0.987
	~		2000	0.597	-0.422	-0.415	0.966	0.914	0.998
	22%	1.019	5000	1.008	-0.011	-0.011	0.430	0.943	1.000
			10000	1.025	0.006	0.006	0.273	0.953	1.000
(2,2,2)			1000	-0.404	-1.781	-1.293	1.259	0.885	0.996
			2000	1.232	-0.146	-0.106	0.606	0.931	1.000
	32%	1.377	5000	1.334	-0.044	-0.032	0.304	0.936	1.000
			10000	1.364	-0.014	-0.010	0.202	0.946	1.000
			1000	1.531	-0.026	-0.017	0.341	0.938	1.000
	66%	1.557	2000	1.522	-0.036	-0.023	0.210	0.934	1.000
			5000	1.550	-0.007	-0.005	0.122	0.938	1.000
			10000	1.557	-0.001	-0.001	0.083	0.951	1.000
			1000	-4.529	-5.436	-5.994	51.315	0.859	0.946
			2000	-1.545	-2.452	-2.704	9.969	0.856	0.971
	11%	0.906	5000	0.090	-0.817	-0.901	1.216	0.910	0.992
			10000	0.941	0.035	0.038	0.607	0.935	1.000
			1000	-2.536	-3.682	-3.211	5.562	0.865	0.978
	22% 1.146		2000	-0.637	-1.783	-1.555	1.298	0.906	0.995
		1.146	5000	1.080	-0.067	-0.058	0.546	0.953	1.000
			10000	1.108	-0.038	-0.033	0.340	0.942	1.000
(4, 4, 4)			1000	-0.306	-2.006	-1.180	1.721	0.907	0.998
			2000	1.412	-0.287	-0.169	0.853	0.945	1.000
	32%	1.699	5000	1.615	-0.084	-0.050	0.425	0.955	1.000
			10000	1.688	-0.011	-0.006	0.278	0.953	1.000
			1000	2.199	-0.117	-0.050	0.574	0.943	1.000
			2000	2.286	-0.029	-0.013	0.350	0.952	1.000
	66%	2.315	5000	2.284	-0.032	-0.014	0.202	0.954	1.000
			10000	2.298	-0.018	-0.008	0.138	0.967	1.000

Table S6: Simulation results for $\widehat{\psi}_{os}$ when the outcome is continuous

¹ True values of SWATE under each setting.
² Truncated means of point estimates. Point estimates are truncated if absolute value are bigger than 500.
³ 90% Winsorized mean of standard error estimates.

⁴ Proportion of point estimates which are not truncated.

β	SWI $\%$	SWATE	Sample size	Estimate	Relative bias	Bias	Se est	Coverage	Acceptance rate
			1000	0.769	0.512	1.984	2.679	0.914	0.790
	~		2000	0.404	0.146	0.568	1.090	0.917	0.884
	11%	0.257	5000	0.284	0.026	0.100	0.370	0.922	0.988
			10000	0.255	-0.003	-0.012	0.172	0.937	1.000
			1000	0.444	0.204	0.846	0.859	0.915	0.931
	220	0.041	2000	0.219	-0.022	-0.090	0.324	0.924	0.998
	22%	0.241	5000	0.242	0.002	0.007	0.136	0.953	1.000
			10000	0.243	0.002	0.010	0.087	0.956	1.000
(0,1-1)			1000	0.381	0.168	0.792	0.414	0.927	0.984
	220	0.010	2000	0.233	0.021	0.097	0.186	0.955	1.000
	32%	0.212	5000	0.215	0.002	0.011	0.095	0.944	1.000
			10000	0.210	-0.002	-0.011	0.063	0.956	1.000
		0.208	1000	0.214	0.006	0.029	0.120	0.954	1.000
	66%		2000	0.209	0.000	0.002	0.073	0.945	1.000
			5000	0.205	-0.003	-0.013	0.043	0.951	1.000
			10000	0.205	-0.003	-0.015	0.029	0.952	1.000
			1000	0.506	-0.019	-0.036	2.304	0.917	0.796
			2000	0.495	-0.029	-0.056	1.725	0.919	0.894
	11%	0.524	5000	0.461	-0.064	-0.122	0.363	0.930	0.994
			10000	0.513	-0.011	-0.022	0.179	0.931	0.999
		0.518	1000	0.505	-0.013	-0.026	0.853	0.904	0.923
	22%		2000	0.440	-0.078	-0.151	0.331	0.930	0.987
			5000	0.508	-0.010	-0.020	0.139	0.963	1.000
			10000	0.521	0.003	0.005	0.089	0.953	1.000
(0,2,-3)			1000	0.445	-0.067	-0.131	0.397	0.911	0.989
		0.511	2000	0.491	-0.020	-0.040	0.197	0.953	1.000
	32%		5000	0.504	-0.007	-0.014	0.098	0.950	1.000
			10000	0.509	-0.002	-0.004	0.065	0.965	1.000
			1000	0.514	0.004	0.008	0.130	0.939	1.000
			2000	0.508	-0.003	-0.005	0.077	0.951	1.000
	66%	0.51	5000	0.507	-0.004	-0.007	0.045	0.956	1.000
			10000	0.507	-0.003	-0.006	0.030	0.950	1.000

Table S7: Simulation results for $\widehat{\psi}_{os}$ when the outcome is binary

¹ True values of SWATE under each setting.
 ² Truncated means of point estimates. Point estimates are truncated if absolute value are bigger than 500.
 ³ 90% Winsorized mean of standard error estimates.

⁴ Proportion of point estimates which are not truncated.




Figure S.4: Simulation results for type-one-error control. The red dash line is the theoretical asymptotic size for each under under different levels.