#### Abstract

A novel approach to accounting for the influence of initial system-bath correlations on the dynamics of an open quantum system, based on the conventional projection operator technique, is suggested. To avoid the difficulties of treating the initial correlations, the conventional Nakajima-Zwanzig inhomogeneous generalized master equations (GMEs) for a system's reduced statistical operator and correlation function are exactly converted into the homogeneous GMEs (HGMEs), which take into account the initial correlations in the kernel governing the evolution of these HGMEs. In the second order (Born) approximation in the systembath interaction, the obtained HGMEs are local in time and valid at all timescales. They are further specialized for a realistic equilibrium Gibbs initial (at  $t = t_0$ ) system+bath state (for a system reduced statistical operator an external force at  $t > t_0$  is applied) and then for a bath of oscillators (Boson field). As an example, the evolution of a selected quantum oscillator (a localized mode) interacting with a Boson field (Fano-like model) is considered at different timescales. It is shown explicitly how the initial correlations influence the oscillator evolution process. In particular, it is shown that the equilibrium system's correlation function acquires at the large timescale the additional constant phase factor conditioned by survived initial system-bath correlations.

# Influence of initial correlations on evolution over time of an open quantum system

Victor F. Los

Institute of Magnetism, Nat. Acad. Sci. and Min. Edu. Sci. of Ukraine, 36-b Vernadsky Blvd., 03142, Kiev, Ukraine

May 14, 2024

#### 1 Introduction

Rigorous derivation of the reliable, tractable and effective evolution equations for the measurable values (statistical expectations) characterizing a nonequilibrium state of a many-particle system remains a principal task of statistical physics. The derivation is conventionally started from the Liouville-von Neumann equation for a statistical operator (distribution function) of the whole system, which depends on the enormous number of variables and thus is practically useless. Fortunately, for practical purposes, one needs the equations only for reduced statistical operators (distribution functions) for a system of interest depending on a much smaller number of variables and which can be obtained by integration off the environment (irrelevant) variables. It is expected that these equations, although obtained from underlying reversible microscopic many-particle dynamics (Liouville-von Neumann equation), should generally be the equations converting into irreversible kinetic ones on some timescale.

In order to derive the equation for reduced statistical operator, several assumptions are usually made. One of them is related to the correlations between a selected system under consideration and its environment in the initial state of a full system (initial correlations). For example, all derivations of the Boltzmann equation for a many-particle system or the Lindblad equation for evolution of a system interacting with a bath are mainly based on either factorizing-type initial conditions (random phase approximation (RPA) or "molecular chaos") corresponding to uncorrelated initial state or on Bogoliubov's principle of weakening of initial correlations [1]. The uncorrelated initial state is not very realistic [2] and the Bogoliubov principle is not always applicable, e.g., when the correlations do not damp with time. Such a situation is realized, e.g., in a quantum many-body system, where the quantum correlations, caused by the particles' statistics, do not damp with time [3], or for an equilibrium initial state of the whole system [2, 4]. More generally, the disregarding of the initial (at the initial time moment  $t_0$  correlation (irrelevant) term, e.g., in the inhomogeneous Nakajima-Zwanzig generalized master equation (GME) [5, 6] for the relevant (reduced) part of a statistical operator of interest, implies, in fact, the "propagation of chaos", i.e. the absence of correlations also at  $t > t_0$  (see, e.g., [2, 7]). The latter has not been proved yet [8]. Since there are no convincing arguments for neglecting the initial correlations [2, 7], it would be desirable to effectively include initial correlations into consideration and thus to have a completely closed (homogeneous) and valid on all timescales evolution equation for a reduced statistical operator. The approaches, like the Bogoliubov principle of weakening of initial correlations, result in the closed homogeneous equations for a system of interest describing the evolution on the large enough timescale, when correlations might damp, and, therefore, are not able to describe the influence of initial correlations on the entire evolution process. The effective accounting in the evolution equation for initial correlations assumes that such an equation at the large timescale would not lead to the appearance of the so called "secular" terms growing with time as, e.g., it is the case when one is trying to solve the inhomogeneous equations of the BBGKY (after the names of Bogoliubov, Born, Green, Kirkwood and Yvon) chain [1]. Generally, the homogeneous equations have a much wider range of applicability with regard to the timescale than the inhomogeneous ones.

Of a special interest is the situation when a system interacts with a stationary environment (a bath), i.e., an open quantum system. Open quantum systems have become an active area of research, owing to its potential applications in many different fields such as quantum gases, quantum optics, quantum information processing, quantum computing (to mention a few), and the general problems of statistical physics [9]. In this area, the evolution of a system is conventionally described by the Lindblad equation (see, [10], [11] and [9]), which is considered now a cornerstone of the theory of open quantum systems. Actually, this equation follows from the Redfield equation [12, 13], which is a local in time equation for a system S density matrix  $\rho_S(t)$  obtained from the von-Neumann equation of motion for the combined (system+environment) statistical operator in the second (Born) approximation in a weak system-environment interaction with additional assumptions that the total system-bath density matrix  $\rho(t)$  can be factorized for any time t, i.e.,  $\rho(t) = \rho_S(t)\rho_B$ , where  $\rho_B$  is a bath density matrix, and that the system density matrix at any retarded time  $\rho_S(t')$  (t' < t) can be replaced by that at the present time  $\rho_S(t)$  (Markovian approximation). To make the Redfield master equation fully Markovian one, the Born-Markov approximation is further applied, which is justified when the bath correlation time  $\tau_B$  is small compared with the relaxation time of a system  $\tau_S$ , i.e.,  $\tau_B \ll \tau_S$ . The Born-Markov approximation means that we consider the system evolution at a large timescale  $t \sim \tau_S \gg \tau_B$ . Generally, the Markovian Redfield equation does not guarantee the dynamics complete positivity, and in order to achieve this goal, the additional approximation, which involves an averaging over the rapidly oscillating terms in the master equation (the rotating wave approximation), is needed. This procedure, which eliminates the very rapidly oscillating during the time  $\tau_S$  terms, means that the evolution equation is valid for a large timescale  $t \sim \tau_S$ . The final quantum master equation, which results from the mentioned approximations, is the Lindblad equation preserving all properties of the density matrix (including complete positivity). Therefore, the Lindblad equation (like the Redfield equation) has its fundamental limitations (see also [14]).

The essential conventional assumption, made at the derivation of the Redfield (and the Lindblad) equation, is that the system-environment correlations (including the correlations at the initial moment of time  $t_0$ ) are ignored. There are several approaches for including the correlations either for specific Hamiltonians (see, e.g., [15]), or in the frameworks of completely positive trace preserving (CPTP) maps [16] and of the Lindblad-like equations ([17]).

In this paper, we suggest the different approach to the initial correlation problem based on the modified standard time-independent projection operator formalism allowing for exact transformation of the conventional Nakajima-Zwanzig-like inhomogeneous generalized master equations (GMEs) for the relevant parts of a system statistical operator or correlation function, containing initial correlations in the inhomogeneous (irrelevant) initial condition term (a source), into the homogeneous generalized master equations (HGMEs) with no irrelevant source term. These exact HGMEs contain initial correlations in the modified flow and collision terms. The parameter of initial correlations is defined by a series in the product of irrelevant parts of statistical operator or correlation function and the inverse of the corresponding relevant parts. In the second order approximation in the system-environment interaction, the obtained HGMEs are local in time and contain general initial system-bath interaction in the kernel governing the evolution of these equations. They exactly describe in the Born approximation the system evolution on any timescale (no Bogoliubov's principle of weakening of initial correlations or the Markov approximation is applied) and have an additional terms defined by initial correlations. These terms are considered for an interesting (and quite realistic) case of the system+bath equilibrium Gibbs initial state, when initial correlations do not damp with time. The equations for  $\rho_S(t)$  (at  $t > t_0$ , when an external force is applied) and system's correlation function are further specialized for the bath of oscillators (Boson field) and then for a quantum oscillator as a system of interest (Fano-like system). These equations are valid on any timescale and describe the evolution of a localized Bose mode. The solutions of equations show that the equilibrium initial correlations influence the selected system's evolution process. For short times,  $\overline{\omega t} \ll 1$  $(\overline{\omega}$  is a bath characteristic frequency), the relaxation process is quadratic in time, and for long times,  $\overline{\omega t} \gg 1$ , the evolution exhibits the standard behavior, but with additional phase shift for the equilibrium system's correlation function conditioned by survived equilibrium initial correlations. Consequences of such a behavioral for the observables are discussed.

# 2 Time-independent projection operator formalism with no "molecular chaos"-type approximation

#### 2.1 Homogeneous equation for the relevant statistical operator

We start with the von-Neumann equation for a statistical operator F(t) of N  $(N \gg 1)$  quantum particles

$$\frac{\partial}{\partial t}F(t) = L(t)F(t). \tag{1}$$

Here, F(t) satisfies the normalization condition

$$TrF(t) = 1, (2)$$

and L is the Liouville superoperator acting on an arbitrary operator A(t) as

$$L(t)A(t) = \frac{1}{i\hbar}[H(t), A(t)], \exp[L(t)]A(t) = \exp[H(t)/i\hbar]A(t)\exp[-H(t)/i\hbar],$$
(3)

where [,] is a commutator and H(t) is the system's Hamiltonian, generally dependent on time.

The formal solution to Eq. (1) is

$$F(t) = U(t, t_0) F(t_0), U(t, t_0) = T \exp\left[\int_{t_0}^t ds L(s)\right],$$
(4)

where T denotes the chronological time-ordering operator, which orders the product of time-dependent operators such that their time-arguments increase from right to left, and  $F(t_0)$  is the statistical operator at some initial moment of time  $t_0$  (initial condition).

We employ first the standard projection operator technique [5], [6], [18]. By applying the time-independent projection operators  $P = P^2$  and  $Q = Q^2 =$ 1 - P (QP = PQ = 0) to Eq. (1), it is easy to obtain the equations for the relevant  $f_r(t) = PF(t)$  and irrelevant  $f_i(t) = QF(t)$  parts of F(t)

$$\frac{\partial}{\partial t}f_r(t) = PL(t)[f_r(t) + f_i(t)],$$
  
$$\frac{\partial}{\partial t}f_i(t) = QL(t)[f_r(t) + f_i(t)].$$
(5)

A relevant part of F(t) is defined in a way permitting the calculation of the average values (observables) of the operators depending on the much smaller number  $s \ll N$  of relevant variables than that of the total system. This is typically achieved by selecting a projector P which integrates off the irrelevant variables excessive for the calculation of observables of interest.

A formal solution to the second Eq. (5) has the form

$$f_i(t) = \int_{t_0}^t dt' S(t, t') Q L(t') f_r(t') + S(t, t_0) f_i(t_0),$$
(6)

where

$$S(t,t_0) = T \exp\left[\int_{t_0}^t ds QL(s)\right].$$
(7)

Inserting this solution into the first Eq. (5), we obtain the conventional exact time-convolution generalized master equation (TC-GME) known as the Nakajima-Zwanzig equation for the relevant part of the statistical operator (see also [9])

$$\frac{\partial f_r(t)}{\partial t} = PL(t)[f_r(t) + \int_{t_0}^t dt' S(t,t')QL(t')f_r(t') + S(t,t_0)f_i(t_0)].$$
(8)

Serving as a basis for many applications, Eq. (8), nevertheless, contains the undesirable and in general non-negligible inhomogeneous term (the last term in the right hand side of (8)), which depends (via  $f_i(t_0)$ ) on the same large number of variables as the statistical operator  $F(t_0)$  at the initial instant  $t_0$ . Therefore, Eq. (8) does not provide for a complete reduced description of a multiparticle system in terms of relevant (reduced) statistical operator. Applying Bogoliubov's principle of weakening of initial correlations [1] allowing to eliminate the influence of  $f_i(t_0)$  on the large enough timescale  $t - t_0 \gg t_{cor}$  ( $t_{cor}$  is the correlation time caused by inter-particle interaction) or using a factorized initial condition, when  $f_i(t_0) = QF(t_0) = 0$ , one can achieve the above-mentioned goal and obtain the homogeneous GME for  $f_r(t)$ , i.e. Eq. (8) with no initial condition term. However, obtained in such a way homogeneous GME is either approximate and valid only on a large enough time scale (when all initial correlations vanish) or applicable only for a rather artificial (actually not realistic), as pointed in [2], initial conditions (no correlations at an initial instant of time). As to our understanding, there is no satisfactory way for eliminating the irrelevant initial condition term (see, e.g., [7]).

In order to obtain an exact homogeneous equation for the relevant part of a statistical operator, one can try to transfer the inhomogeneous initial correlations term in the right hand side of Eq. (8) to the (super)operator acting on the relevant part  $f_r(t)$ . To achieve this goal, we suggest to present the initial (irrelevant) term  $f_i(t_0) = QF(t_0)$  as a following exact identity

$$f_{i}(t_{0}) = f_{i}(t_{0})F^{-1}(t_{0})U^{-1}(t,t_{0})(P+Q)U(t,t_{0})F(t_{0})$$
  
=  $K_{0}U^{-1}(t,t_{0})[f_{r}(t) + f_{i}(t)],$   
 $K_{0} = f_{i}(t_{0})F^{-1}(t_{0}), U^{-1}(t,t_{0}) = T_{-}\exp[-\int_{t_{0}}^{t} dsL(s)],$  (9)

where  $U^{-1}(t, t_0)$  is the backward-in-time evolution operator,  $T_{-}$  is the antichronological time-ordering operator arranging the time-dependent operators L(s) in such a way that the time arguments increase from left to right,  $U^{-1}(t, t_0)U(t, t_0) = 1$ ,  $F^{-1}(t_0)$  is inverse of  $F(t_0)$ ,  $F^{-1}(t_0)F(t_0) = 1$ , and P + Q = 1. Hence, additional identity (9) is obtained by multiplying the irrelevant part by unity  $F^{-1}(t_0)F(t_0)$  and inserting  $U^{-1}(t, t_0)U(t, t_0) = 1$  and P(t) + Q(t) = 1.

In (9), we introduce the parameter of initial correlations

$$K_{0} = f_{i}(t_{0})F^{-1}(t_{0}) = f_{i}(t_{0})[f_{r}(t_{0}) + f_{i}(t_{0})]^{-1}$$
  
=  $f_{i}(t_{0})f_{r}^{-1}(t_{0})[1 + f_{i}(t_{0})f_{r}^{-1}(t_{0})]^{-1}$   
=  $(1 - K_{0})f_{i}(t_{0})f_{r}^{-1}(t_{0}),$  (10)

which is a series in  $f_i(t_0)f_r^{-1}(t_0)$ .

We now have two equations, (6) and (9), relating  $f_i(t)$  to  $f_i(t_0)$ . Finding  $f_i(t_0)$  from these equations as a function of  $f_r(t)$  and substituting it in (8), we obtain the equation

$$\frac{\partial f_r(t)}{\partial t} = PL(t)R(t, t_0)[f_r(t) + \int_{t_0}^t dt' S(t, t')QL(t')f_r(t')],$$
(11)

where operator  $R(t, t_0)$  is defined as

$$R(t,t_0) = 1 + K(t,t_0),$$
  

$$K(t,t_0) = S(t,t_0) \left[1 - K_0(t,t_0)\right]^{-1} K_0 U^{-1}(t,t_0),$$
  

$$K_0(t,t_0) = K_0 U^{-1}(t,t_0) S(t,t_0).$$
(12)

Thus, we have obtained the exact integro-differential Eq. (11), which is the homogeneous one (in contrast to Eq. (8)) and takes into account the initial correlations and their dynamics on an equal footing with collisions via the modification of the (super)operator (memory kernel) of GME (8) acting on the relevant part of the statistical operator  $f_r(t)$ . The obtained exact kernel of Eq. (11) can serve as a starting point for consecutive perturbation expansions. In many cases such expansions of the homogeneous equations (like (11)) have much broader range of validity than that for the inhomogeneous ones (like (8)).

#### 2.2 Homogeneous equation for system's correlation function

Another useful approach to studying the reduced dynamics of a many-particle system is the correlation functions approach. We divide the total system of N particle into a system of interest S with a number of particles  $s \ll N$  and its environment (the rest of particles). Let us consider the following two-time correlation function for the time-dependent operators  $\alpha_S(t) = e^{-Lt}\alpha_S(0)$  and  $\beta_S(t) = e^{-Lt}\beta_S(0)$  related to the system S

$$\langle \alpha_S(0)\beta_S(t)\rangle = Tr_S[\beta_S(0)\rho_S^{\alpha}(t)], \rho_S^{\alpha}(t) = Tr_BF_{\alpha}(t)], F_{\alpha}(t) = e^{Lt}F_{\alpha}(0), F_{\alpha}(0) = \rho_{tot}(0)\alpha_S(0),$$
(13)

where the superoperator L is given by (3) (with time-independent L), and the averaging symbol  $\langle ... \rangle = Tr_{S+B}[...\rho_{tot}(0)]$  is defined as the trace  $Tr_{S+B}$  over the total S+B system degrees of freedom with the time-independent statistical operator  $\rho_{tot}(0) = \rho_{tot}$  at the initial time moment  $t_0 = 0$  (the Heisenberg picture), which generally is not a statistical operator for S+B system in an equilibrium state. We also used the invariance of trace under the cyclic permutation of operators. The superoperator  $F_{\alpha}(t)$  is subject to the following equation

$$\frac{\partial}{\partial t}F_{\alpha}(t) = LF_{\alpha}(t), \qquad (14)$$

which defines the evolution of the correlation function (13) in time .

As earlier, with the help of the projection operators P and Q = 1 - Pand taking into account that in Eq. (14) L does not depend on time, we can write down the Nakajima-Zwanzig equation (8) for the relevant part of the superoperator  $F_{\alpha}(t)$ ,  $f_r^{\alpha}(t) = PF_{\alpha}(t)$ , as

$$\frac{\partial f_r^{\alpha}(t)}{\partial t} = PL[f_r^{\alpha}(t) + \int_0^t dt' e^{QL(t-t')} QLf_r^{\alpha}(t') + e^{QLt}f_i^{\alpha}(0)], \qquad (15)$$

where the irrelevant part of  $F_{\alpha}(t)$  is  $f_{i}^{\alpha}(t) = QF_{\alpha}(t)$ . As it is seen from Eq. (15), the projector P selects the relevant part of  $F_{\alpha}(t)$ , which is supposed to be sufficient for obtaining the closed equation for the evolution of correlation function for system's S operators, i.e., an operator P projects the large-dimensional Hilbert space of the total system on the smaller dimensional space of the system of interest S. But Eq. (15) is the inhomogeneous one which contains the irrelevant initial condition term  $f_{i}^{\alpha}(0) = (\rho_{tot} - P\rho_{tot})\alpha_{S}(0)$ .

To make this equation homogeneous (completely closed), the above described procedure can be employed. As a result, we arrive at the following homogeneous equation

$$\frac{\partial f_r^{\alpha}(t)}{\partial t} = PLR^{\alpha}(t)[f_r^{\alpha}(t) + \int_0^t dt' e^{QL(t-t')}QLf_r^{\alpha}(t')], \qquad (16)$$

where operator  $R^{\alpha}(t)$  is defined as

$$R^{\alpha}(t) = 1 + K^{\alpha}(t),$$

$$K^{\alpha}(t) = e^{QLt} \left[1 - K_{0}^{\alpha}(t)\right]^{-1} K_{0}^{\alpha} e^{-Lt},$$

$$K_{0}^{\alpha}(t) = K_{0}^{\alpha} e^{-Lt} e^{QLt},$$

$$K_{0}^{\alpha} = f_{i}^{\alpha}(0) F_{\alpha}^{-1}(0) = (1 - K_{0}^{\alpha}) f_{i}^{\alpha}(0) [f_{r}^{\alpha}(0)]^{-1}.$$
(17)

#### 3 System in a bath

For the case under consideration, when a system of interest S interacts with another quantum stationary system B, called bath, the Hamiltonian of the whole S + B system can be presented as

$$H(t) = H_S(t) + H_B + H_{SB},$$
(18)

where  $H_S(t)$  is related to a system S and can depend on time through an applied external force,  $H_B$  and  $H_{SB}$  are the Hamiltonians of a bath and of a system-bath interaction, respectively. The corresponding Liouville superoperator is

$$L(t) = L_S(t) + L_B + L_{SB}.$$
 (19)

We are interested in finding the evolution equation for statistical operator of the system  ${\cal S}$ 

$$\rho_S(t) = Tr_B F(t) \tag{20}$$

from Eq. (11) ( $Tr_B$  is the partial trace of the bath degrees of freedom). To this end, it is convenient to introduce the following projection operators

$$P = \rho_B T r_B, Q = 1 - P, T r_B \rho_B = 1,$$
(21)

where  $\rho_B$  stands for the normalized statistical operator of a bath. Then the relevant and irrelevant parts of the statistical operator F(t) are

$$f_r(t) = PF(t) = \rho_B \rho_S(t), f_i(t) = QF(t) = F(t) - \rho_B \rho_S(t).$$
(22)

Then, Eq. (11) can be rewritten in a more specific and simple form for the Liouvillian (19) and projection operators (21) if we take into consideration the following properties

$$PL_S(t)Q = QL_S(t)P = 0, PL_B = L_BP = 0, PL_BQ = QL_BP = 0.$$
 (23)

where we took into account that  $[H_B, \rho_B] = 0$ . We also assume that

$$\langle H_{SB} \rangle_B = T r_B H_{SB} \rho_B = 0 \tag{24}$$

and, therefore,

$$PL_{SB}P = 0, (25)$$

which is a rather typical situation (see below).

Now, Eq. (11) can be rewritten as

$$\frac{\partial f_r(t)}{\partial t} = L_S(t)f_r(t) + PL_{SB}\overline{K}(t,t_0)f_r(t) + PL_{SB}[1+\overline{K}(t,t_0)]\int_{t_0}^t dt'\overline{S}(t,t')L_{SB}f_r(t'),$$
(26)

where  $\overline{K}(t, t_0)$  is defined by (12) with S(t, t') replaced by

$$\overline{S}(t,t') = T \exp\left\{\int_{t'}^{t} ds [L_0(s) + QL_{SB}]\right\},$$

$$L_0(s) = L_S(s) + L_B$$
(27)

Taking  $Tr_B$  from both sides of Eq. (26) and using definitions (21) and (22), we obtain the following exact equation for a system's statistical operator

$$\frac{\partial \rho_S(t)}{\partial t} = L_S(t)\rho_S(t) + Tr_B L_{SB}\overline{K}(t,t_0)\rho_B\rho_S(t) + Tr_B L_{SB}[1+\overline{K}(t,t_0)] \int_{t_0}^t dt'\overline{S}(t,t')L_{SB}\rho_B\rho_S(t').$$
(28)

This equation differs from the standard form of such an equation (see, e.g., [9]) by the additional terms in the kernel containing  $\overline{K}(t, t_0)$ , which account for the influence of initial system-bath correlations on the evolution of the system. We also note, that conventionally used "molecular chaos"-type approximation,  $f_r(t') = \rho_B \rho_S(t')$  (including the initial time moment  $t' = t_0$ ), is not used when going from Eq. (26) to Eq. (28).

In the correlation function approach, when we have a time-independent  $L_S$   $(H_S)$ , the relevant and irrelevant parts of  $F_{\alpha}(t)$  selected by the projector (21) are

$$f_r^{\alpha}(t) = PF_{\alpha}(t) = \rho_B \rho_S^{\alpha}(t), f_i^{\alpha}(t) = QF_{\alpha}(t) = F_{\alpha}(t) - \rho_B \rho_S^{\alpha}(t),$$
  

$$\rho_S^{\alpha}(t) = Tr_B F_{\alpha}(t).$$
(29)

Then, using (23) and (25), Eq. (16) can be simplified as

$$\frac{\partial f_r^{\alpha}(t)}{\partial t} = L_S f_r^{\alpha}(t) + P L_{SB} \overline{K}^{\alpha}(t) f_r^{\alpha}(t) + P L_{SB} [1 + \overline{K}^{\alpha}(t)] \int_0^t dt' e^{(L_0 + Q L_{SB})(t - t')} L_{SB} f_r^{\alpha}(t'), \quad (30)$$

where  $\overline{K}^{\alpha}(t)$  is defined by (17) with the substitution  $e^{QLt} \to \exp(L_0 + QL_{SB})$ . We remind, that for the correlation function case, the Hamiltonian is time-independent and  $L_0 = L_S + L_B$ .

Again, by applying  $Tr_B$  to Eq. (30) from the left, we obtain the following exact equation for  $\rho_S^{\alpha}(t)$ 

$$\frac{\partial \rho_{S}^{\alpha}(t)}{\partial t} = L_{S} \rho_{S}^{\alpha}(t) + Tr_{B} L_{SB} \overline{K}^{\alpha}(t) \rho_{B} \rho_{S}^{\alpha}(t) + Tr_{B} L_{SB} [1 + \overline{K}^{\alpha}(t)] \int_{0}^{t} dt' e^{(L_{0} + QL_{SB})(t-t')} L_{SB} \rho_{B} \rho_{S}^{\alpha}(t'), (31)$$

which defines the evolution of the correlation function (13) according to

$$\frac{\partial}{\partial t} \langle \alpha_S(0)\beta_S(t) \rangle = Tr_S[\beta_S(0)\frac{\partial}{\partial t}\rho_S^{\alpha}(t)].$$
(32)

Obtained Eqs. (28) and (31) are the homogeneous exact evolution equations for a system's reduced statistical operator and correlation function in the case of a system interacting with a bath which exactly account for initial correlations.

#### 4 The Born approximation

Equations (28) and (31) are exact but very complicated. Let us consider the case of a weak system-bath interaction when  $H_{SB}$  is proportional to some small parameter  $\varepsilon$ . The collision terms (third in the r.h.s. of Eq.(28) and Eq. (31)) are already proportional to  $\varepsilon^2$ . Then, we will restrict ourselves by the second order in the system-bath interaction (the Born approximation). So, we need to estimate (with regard to the power of  $\varepsilon$ ) the initial correlations parameters  $\overline{K}(t, t_0)$  and  $\overline{K}^{\alpha}(t)$ . In the zero approximation in  $\varepsilon$ ,

$$\overline{S}(t,t_0) = U_0(t,t_0) = T \exp[\int_{t_0}^t ds L_0(s)],$$

$$U_0^{-1}(t,t_0) = T_- \exp[-\int_{t_0}^t ds L_0(s)],$$

$$\exp[(L_0 + QL_{SB})t] = \exp(L_0 t), e^{-Lt} = e^{-L_0 t}.,$$
(33)

and, therefore,  $\overline{K_0}(t, t_0) = K_0$  and  $\overline{K}_0^{\alpha} = K_0^{\alpha}$  (see (12) and (17)). Therefore,  $\overline{K}(t, t_0) = \overline{K_1}(t, t_0) = U_0(t, t_0)(1 - K_0)^{-1}K_0U_0^{-1}(t, t_0)$ ,  $\overline{K}^{\alpha}(t) = \overline{K}_1^{\alpha}(t) = e^{L_0t}(1 - K_0^{\alpha})^{-1}K_0^{\alpha}e^{-L_0t}$ , and because  $(1 - K_0)^{-1}K_0 = f_i(t_0)f_r^{-1}(t_0)$ ,  $(1 - K_0^{\alpha})^{-1}K_0^{\alpha} = f_i^{\alpha}(0)[f_r^{\alpha}(0)]^{-1}$ , as it follows from (10) and (17), we finally have in the adopted approximation

$$\overline{K}_{1}(t,t_{0}) = G_{SB}(t,t_{0}) = U_{0}(t,t_{0})f_{i}(t_{0})f_{r}^{-1}(t_{0})U_{0}^{-1}(t,t_{0}), 
\overline{K}_{1}^{\alpha}(t) = G_{SB}^{t}(t) = e^{L_{0}t}f_{i}^{\alpha}(0)[f_{r}^{\alpha}(0)]^{-1}e^{-L_{0}t},$$
(34)

where

$$\begin{aligned}
f_i(t_0)f_r^{-1}(t_0) &= [F(t_0) - \rho_B \rho_S(t_0)][\rho_B \rho_S(t_0)]^{-1}, \rho_S(t_0) = Tr_B F(t_0), \\
f_i^{\alpha}(0)[f_r^{\alpha}(0)]^{-1} &= (\rho_{tot} - \rho_B \rho_S^t) \alpha_S(0)[\rho_B \rho_S^t \alpha_S(0)]^{-1} \\
&= (\rho_{tot} - \rho_B \rho_S^t)[\rho_B \rho_S^t]^{-1}, \rho_S^t = Tr_B \rho_{tot}
\end{aligned}$$
(35)

(see (22)) and (29). Note, that  $G_{SB}^t(t)$  (34) does not depend on  $\alpha_S(0)$  (see (35)). Estimating  $[F(t_0) - \rho_B \rho_S(t_0)] \sim \varepsilon$  and  $(\rho_{tot} - \rho_B \rho_S^t) \sim \varepsilon$ , we obtain that both functions (34)  $\sim \varepsilon$ .

Thus, in the second order in  $\varepsilon$ , we have from Eqs. (28) and (31) the following equations for  $\rho_S(t)$  and  $\rho_S^{\alpha}(t)$  after the change of integration variable  $t - t' = \tau$ 

$$\frac{\partial \rho_S(t)}{\partial t} = L_S(t)\rho_S(t) + Tr_B L_{SB} G_{SB}(t,t_0)\rho_B \rho_S(t) 
+ Tr_B L_{SB} \int_0^{t-t_0} d\tau U_0(t,t-\tau) L_{SB} \rho_B \rho_S(t-\tau), 
\frac{\partial \rho_S^{\alpha}(t)}{\partial t} = L_S \rho_S^{\alpha}(t) + Tr_B L_{SB} G_{SB}^t(t) \rho_B \rho_S^{\alpha}(t) 
+ Tr_B L_{SB} \int_0^t d\tau e^{L_0 \tau} L_{SB} \rho_B \rho_S^{\alpha}(t-\tau).$$
(36)

The obtained Eqs. (36) contain (as compared to the standard case) the additional terms in the kernels governing the evolution of  $\rho_S(t)$  and  $\rho_S^{\alpha}(t)$ , which account for initial correlations (terms with  $G_{SB}(t, t_0)$  and  $G_{SB}^t(t)$ ).

However, Eqs. (36) are still a time-convolution and non-Markovian ones. To make them time local, we will employ an approach used for obtaining a time-convolutionless GME ([19, 20]). It follows from Eqs. (4) and (13), that

$$F(t-\tau) = U^{-1}(t,t-\tau)F(t) = U^{-1}(t,t-\tau)[f_r(t)+f_i(t)],$$
  

$$F_{\alpha}(t-\tau) = e^{-L\tau}F_{\alpha}(t) = e^{-L\tau}[f_r^{\alpha}(t)+f_i^{\alpha}(t)].$$
(37)

By applying the projector P(21) to this relation, using the definitions (22), (29) and that  $f_i(t) \sim \varepsilon$ ,  $f_i^{\alpha}(t) \sim \varepsilon$ , we obtain in the zero approximation in  $\varepsilon$ ,

$$\rho_{S}(t-\tau) = U_{S}^{-1}(t,t-\tau)\rho_{S}(t), \rho_{S}^{\alpha}(t-\tau) = e^{-L_{S}\tau}\rho_{S}^{\alpha}(t)$$
$$U_{S}^{-1}(t,t-\tau) = T_{-}\exp[-\int_{t-\tau}^{t}d\lambda L_{S}(\lambda)], \qquad (38)$$

where we have also used that  $Tr_B \exp(-L_B \tau)\rho_B = 1$ .

~

Now, remaining in the Born (second in  $\varepsilon$ ) approximation, Eqs. (36) can be rewritten in the time-local form as

$$\frac{\partial \rho_S(t)}{\partial t} = L_S(t)\rho_S(t) + Tr_B L_{SB} G_{SB}(t,t_0)\rho_B \rho_S(t) 
+ Tr_B L_{SB} \int_0^{t-t_0} d\tau U_0(t,t-\tau) L_{SB} \rho_B U_S^{-1}(t,t-\tau)\rho_S(t), 
\frac{\partial \rho_S^{\alpha}(t)}{\partial t} = L_S \rho_S^{\alpha}(t) + Tr_B L_{SB} G_{SB}^t(t) \rho_B \rho_S^{\alpha}(t) 
+ Tr_B L_{SB} \int_0^t d\tau e^{L_0 \tau} L_{SB} \rho_B e^{-L_S \tau} \rho_S^{\alpha}(t).$$
(39)

Equations (39) represent the central result of this section. They are exact in the second approximation in the system-bath interaction and homogeneous (completely closed) time-local evolution equations for the system's statistical operator and correlation function, respectively. No "molecular chaos"-type or Bogoliubov's principle of weakening of initial correlations approximation has been used. These equations effectively account for the influence in time of initial correlations on the system's evolution process via the functions  $G_{SB}(t, t_0)$  or  $G_{SB}^{t}(t)$  (34) defined for an arbitrary initial statistical operator  $F(t_0)$  or  $\rho_{tot}(0)$ , respectively.

#### 4.1An equilibrium Gibbs initial state

Let us suppose, that up to the moment of time  $t_0$  the total system is in an equilibrium state with the Gibbs statistical operator but just after  $t_0$  (at t > $t_0$ ) an external (generally time-dependent) force (described by the Hamiltonian  $H_{ext}(t)$  is applied to a system S driving it from an initial state, i.e.,

$$F(t \leq t_0) = \rho_{eq} = Z^{-1} \exp(-\beta H), H = H_S + H_B + H_{SB}, t \leq t_0,$$
  

$$\beta = 1/k_B T, Z = Tr_{S+B} \exp(-\beta H),$$
  

$$H(t) = H_S(t) + H_B + H_{SB}, H_S(t) = H_S + H_{ext}(t), t > t_0.$$
(40)

In this case, the evolution of the system's statistical operator can be described by Eq. (39) for  $\rho_S(t)$  with the equilibrium initial statistical operator for the total system  $F(t_0) = Z^{-1} \exp(-\beta H)$ . For this quite realistic initial equilibrium state we can explicitly find the initial condition function  $G_{SB}(t, t_0)$  (34) by making use of the following exact identity

$$e^{-\beta H} = e^{-\beta H_0} - \int_0^\beta d\lambda e^{-\lambda H_0} H_{SB} e^{\lambda H} e^{-\beta H},$$
  

$$H_0 = H_S + H_B.$$
(41)

We also select

$$\rho_B = \rho_B^{eq} = \frac{e^{-\beta H_B}}{Tr_B e^{-\beta H_B}} \tag{42}$$

in the definition (21) for the projection operator.

Then, in the linear approximation in  $H_{SB}$ ,

$$\rho_{S}(t_{0}) = Tr_{B}F(t_{0}) = Tr_{B}e^{-\beta H}/Z$$

$$\approx Tr_{B}[e^{-\beta H_{0}} - \int_{0}^{\beta} d\lambda e^{-\lambda H_{0}}H_{SB}e^{\lambda H_{0}}e^{-\beta H_{0}}]/Tr_{S+B}[e^{-\beta H_{0}} - \int_{0}^{\beta} d\lambda e^{-\lambda H_{0}}H_{S}e^{-\beta H_{0}}]/Tr_{S+B}[e^{-\beta H_{0}} - \int_{0}^{\beta}$$

It is not difficult to see that

$$Tr_B \int_0^\beta d\lambda e^{-\lambda H_0} H_{SB} e^{\lambda H_0} e^{-\beta H_0} = 0, \qquad (44)$$

and. therefore, in this approximation,

$$\rho_S(t_0) = e^{-\beta H_S} / Tr_S e^{-\beta H_S}.$$
(45)

where we have used the condition (24).

Thus, in the first approximation in  $H_{SB}$ 

$$[F(t_0) - \rho_B \rho_S(t_0)] [\rho_B \rho_S(t_0)]^{-1} = e^{-\beta H} [\rho_B^{eq} \rho_S(t_0)]^{-1} - 1$$
$$= -\int_0^\beta d\lambda e^{-\lambda H_0} H_{SB} e^{\lambda H_0}, \qquad (46)$$

and, finally, the function defining the influence of initial correlations on the evolution of the system's statistical operator, given by Eq. (39), is

$$G_{SB}(t,t_0) = -U_0(t,t_0) \int_0^\beta d\lambda e^{-\lambda H_0} H_{SB} e^{\lambda H_0} U_0^{-1}(t,t_0).$$
(47)

Likewise, selecting  $\rho_{tot}(0) = Z^{-1} \exp(-\beta H)$  and  $\rho_B = \rho_B^{eq}$  (42), we obtain for function  $G_{SB}^t(t)$  (34)

$$G_{SB}^{t}(t) = -e^{L_{0}t} \int_{0}^{\beta} d\lambda e^{-\lambda H_{0}} H_{SB} e^{\lambda H_{0}} e^{-L_{0}t}.$$
(48)

This function describes the influence of initial correlation on the evolution of the equilibrium correlation function (13) (defined with the  $\rho_{tot}(0) = Z^{-1} \exp(-\beta H)$ ) according to the second Eq. (39).

It is interesting to note, that the terms describing the influence of initial equilibrium correlations, given by Eqs. (47) and (48), coincide with those obtained in works [22, 21] (dealing with the polaron mobility) by different method of converting the inhomogeneous Nakajima-Zwanzig equation into the homogeneous one based on the identity (41), which, however, is applicable only for initial Gibbs state (40) for the total system. The method suggested in this work (see Sec 2) is applicable for any initial state.

#### 4.2 A system interacting with the Boson field

Let us consider the case when a system interacts with the Boson field which acts as the bath, i.e., we assume that

$$H_{B} = \sum_{k} \hbar \omega_{k} b_{k}^{+} b_{k},$$
  

$$H_{SB} = \sum_{k} [C_{k}(S)b_{k} + C_{k}^{+}(S)b_{k}^{+}],$$
(49)

where  $\hbar \omega_k$  is the energy of the field quantum characterized by the set of quantum numbers  $k, b_k^+, b_k$  are the Bose-operators of creation and annihilation of the field quantum, and  $C_k(S)$  is an operator acting on a system S.

For simplicity, we assume that an external force (contributing to  $L_S(t)$ ) in Eq. (39) for system's statistical operator  $\rho_S(t)$  is weak and thus disregard it in the initial correlation and collision terms (linear response regime). In this case (and putting  $t_0 = 0$ ), the mentioned terms in Eqs. (39) for  $\rho_S(t)$  and  $\rho_S^{\alpha}(t)$  become formally identical.

Then, the term of initial correlations in Eq. (39) for  $\rho_S(t)$  with the use of

(47), (48) (equilibrium initial state), and (49), can be written as

$$I_{S}(t,t_{0}) = Tr_{B}L_{SB}G_{SB}(t,t_{0})\rho_{B}\rho_{S}(t) = -PL_{SB}e^{L_{0}t} \int_{0}^{\beta} d\lambda e^{-\lambda H_{0}}H_{SB}e^{\lambda H_{0}}e^{-L_{0}t}\rho_{B}^{eq}\rho_{S}(t)$$
$$= i\int_{0}^{\beta} d\lambda \sum_{k} \{e^{-i\omega_{k}(t-i\lambda)}(1+N_{k})[C_{k}(S), C_{k}^{+}(S_{t,\lambda})\rho_{S}(t)] + e^{i\omega_{k}(t-i\lambda)}N_{k}[C_{k}^{+}(S), C_{k}(S_{t,\lambda})\rho_{S}(t)]\},$$
(50)

where we put  $\hbar = 1$  and

$$C_k(S_{t,\lambda}) = e^{-iH_S t} e^{-\lambda H_S} C_k(S) e^{iH_S t} e^{\lambda H_S}, C_k^+(S_{t,\lambda}) = e^{-iH_S t} e^{-\lambda H_S} C_k^+(S) e^{iH_S t} e^{\lambda H_S}.$$
(51)

Here we also used, that for  $H_B$ , given by (49),

$$e^{-iH_{B}t}b_{\mathbf{k}}e^{iH_{B}t} = e^{i\omega_{\mathbf{k}}t}b_{\mathbf{k}}, e^{-iH_{B}t}b_{\mathbf{k}}^{+}e^{iH_{B}t} = e^{-i\omega_{\mathbf{k}}t}b_{\mathbf{k}}^{+},$$

$$e^{-\lambda H_{B}}b_{k}e^{\lambda H_{B}} = e^{\lambda\omega_{k}}b_{k}, e^{-\lambda H_{B}}b_{k}^{+}e^{\lambda H_{B}} = e^{-\lambda\omega_{k}}b_{k}^{+}$$

$$< b_{k}b_{k_{1}} >_{B} = 0, < b_{k}^{+}b_{k_{1}}^{+} >_{B} = 0,$$

$$< b_{k}b_{k_{1}}^{+} >_{B} = (1+N_{k})\delta_{kk_{1}}, < b_{k}^{+}b_{k_{1}} >_{B} = N_{k}\delta_{kk_{1}},$$

$$< \dots >_{B} = Tr_{\Sigma}(\dots\rho_{B}^{eq}), N_{k} = [\exp(\beta\omega_{k}) - 1]^{-1}$$
(52)

Likewise, we have for the term of Eq. (39), which can be identified with a collision integral,

$$C_{S}(t,t_{0}) = Tr_{B}L_{SB}\int_{0}^{t} d\tau e^{L_{0}\tau}L_{SB}\rho_{B}^{eq}e^{-L_{S}\tau}\rho_{S}(t)$$

$$= -\int_{0}^{t} d\tau \sum_{k} \{e^{-i\omega_{k}\tau}(1+N_{k})[C_{k}(S), C_{k}^{+}(S_{\tau})\rho_{S}(t)]$$

$$+e^{i\omega_{k}\tau}N_{k}[C_{k}^{+}(S), C_{k}(S_{\tau})\rho_{S}(t)] + h.c\}.$$
(53)

where

$$C_k(S_{\tau}) = e^{-iH_S\tau} C_k(S) e^{iH_S\tau}, C_k^+(S_{\tau}) = e^{-iH_S\tau} C_k^+(S) e^{iH_S\tau}.$$
 (54)

The same expressions with the substitution  $\rho_S(t) \rightarrow \rho_S^{\alpha}(t)$  are valid for the initial correlations and collision terms of Eq. (39) for system's correlation function (see (32)). We note, that in the adopted approximation, the collision integral (53) is valid for any initial condition (not only for the equilibrium Gibbs state (40)).

## 5 Quantum oscillator (Fano-like model)

The results, given by Eqs. (50) and (53), define the evolution equations (39) for the system's density matrix and correlaton function with account for initial

correlations. These results hold for any system's Hamiltonian  $H_S(t)$ , any small parameter of the system-Boson field interaction  $C_k(S)$ , and equilibrium initial state of the total system.

Let us further specify the system and consider the following Hamiltonian for a single-mode cavity system (could be a nanocavity in nanostructures or photonic crystals) interacting with a Boson reservoir

$$H(t = 0) = \omega_c a^+ a + \sum_k \omega_k b_k^+ b_k + \sum_k V_k (ab_k^+ + a^+ b_k),$$
  

$$H(t > 0) = H_{ext}^S(t) + H(t = 0),$$
(55)

where the first term in H(t = 0) is the Hamiltonian of the cavity field with frequency  $\omega_c$  (corresponds to the system Hamiltonian  $H_S$ ),  $a^+$  and a are the creation and annihilation operators of the cavity field. The second term (corresponds to  $H_B$ ) describes the environment as the reservoir of the infinite Boson modes, and the third term  $(H_{SB})$  is the the system-bath interaction with the coupling strength  $V_k$ . The Hamiltonian (55) represents a Fano-type model of a localized state coupled with a continuum [23]. Thus, we can now calculate explicitly the initial correlation term (50) and the collision term (53) for Eqs. (39) by putting in these equations

$$C_{k}(S) = V_{k}a^{+}, C_{k}^{+}(S) = V_{k}a,$$

$$C_{k}(S_{\tau}) = e^{-iH_{S}\tau}C_{k}(S)e^{iH_{S}\tau} = V_{k}e^{-i\omega_{c}t}a^{+},$$

$$C_{k}^{+}(S_{\tau}) = e^{-iH_{S}\tau}C_{k}^{+}(S)e^{iH_{S}\tau} = V_{k}e^{i\omega_{c}t}a,$$

$$C_{k}(S_{t,\lambda}) = e^{-iH_{S}t}e^{-\lambda H_{S}}C_{k}(S)e^{iH_{S}t}e^{\lambda H_{S}} = V_{k}e^{-i\omega_{c}(t-i\lambda)}a^{+}$$

$$C_{k}^{+}(S_{t,\lambda}) = e^{-iH_{S}t}e^{-\lambda H_{S}}C_{k}^{+}(S)e^{iH_{S}t}e^{\lambda H_{S}} = V_{k}e^{i\omega_{c}(t-i\lambda)}a.$$
(56)

For initial correlation term the result is

$$I_{S}(t) = i \int_{0}^{\beta} d\lambda \sum_{k} V_{k}^{2} \{ e^{-i(\omega_{k} - \omega_{c})(t - i\lambda)} (1 + N_{k}) [a^{+}, a\rho_{S}(t)] \} + e^{i(\omega_{k} - \omega_{c})(t - i\lambda)} N_{k} [a, a^{+}\rho_{S}(t)] \} = i \sum_{k} V_{k}^{2} \{ e^{-i(\omega_{k} - \omega_{c})t} (1 + N_{k}) \frac{e^{(\omega_{c} - \omega_{k})\beta} - 1}{\omega_{c} - \omega_{k}} [a^{+}, a\rho_{S}(t)] + e^{i(\omega_{k} - \omega_{c})t} N_{k} \frac{e^{(\omega_{k} - \omega_{c})\beta} - 1}{\omega_{k} - \omega_{c}} [a, a^{+}\rho_{S}(t)] \},$$
(57)

and the collision term is given by

$$C_{S}(t) = -\int_{0}^{t} d\tau \sum_{k} V_{k}^{2} \{ e^{-i(\omega_{k}-\omega_{c})\tau} (1+N_{k})[a^{+}, a\rho_{S}(t)] + e^{i(\omega_{k}-\omega_{c})\tau} N_{k}[a, a^{+}\rho_{S}(t)] + e^{i(\omega_{k}-\omega_{c})\tau} (1+N_{k})[\rho_{S}(t)a^{+}, a] e^{-i(\omega_{k}-\omega_{c})\tau} N_{k}[\rho_{S}(t)a, a^{+}] \}$$

$$= -\int_{0}^{t} d\tau \sum_{k} V_{k}^{2} \{ e^{-i(\omega_{k}-\omega_{c})\tau} (1+N_{k})[a^{+}, a\rho_{S}(t)] + e^{i(\omega_{k}-\omega_{c})\tau} N_{k}[a, a^{+}\rho_{S}(t)] + h.c \}$$
(58)

Thus, we have the following equations for system's statistical operator and equilibrium correlation function in the case of the Hamiltonian (55)

$$\frac{\partial \rho_S(t)}{\partial t} = L_S(t)\rho_S(t) + I_S(t) + C_S(t),$$
  

$$\frac{\partial \rho_S^{\alpha}(t)}{\partial t} = L_S\rho_S^{\alpha}(t) + I_S^{\alpha}(t) + C_S^{\alpha}(t),$$
(59)

where  $I_S^{\alpha}(t)$  and  $C_S^{\alpha}(t)$  are defined by Eqs. (57), (58) with the substitution  $\rho_S(t) \to \rho_S^{\alpha}(t)$  for the second Eq. (59), and  $L_S(t)$  differs from  $L_S$  by the term conditioned by an external driving force.

It is instructive to consider Eqs. (59) neglecting the initial condition terms. Then, let us assume that the integration over  $\tau$  in (58) can be extended to infinity, i.e., that we are interested in the evolution on the relaxation timescale  $t \sim \tau_{rel} \gg |\omega_k - \omega_c|^{-1}$  (the Markov approximation). We also define the integrals over  $\tau$  as

$$\int_{0}^{\infty} d\tau e^{\pm i(\omega_k - \omega_c)\tau} = \lim_{\eta \to +0} \int_{0}^{\infty} d\tau e^{\pm i(\omega_k - \omega_c)\tau - \eta\tau} = \pi \delta(\omega_k - \omega_c) \pm iP \frac{1}{\omega_k - \omega_c},$$
(60)

where P stands for the integral principal value. As a result, we obtain the following expression for the collision term

$$C_{S}(t) = -i\Delta\omega_{c}[a^{+}a,\rho_{S}(t)] + J(\omega_{c})(1+N_{c})[a\rho_{S}(t)a^{+} - \frac{1}{2}\{a^{+}a,\rho_{S}(t)\}] + J(\omega_{c})N_{c}[a^{+}\rho_{S}(t)a - \frac{1}{2}\{aa^{+},\rho_{S}(t)\}], t \sim \tau_{rel} \gg |\omega_{k} - \omega_{c}|^{-1}, \Delta\omega_{c} = P \int_{0}^{\infty} \frac{d\omega}{2\pi} \frac{J(\omega)}{\omega_{c} - \omega}, J(\omega) = 2\pi \sum_{k} V_{k}^{2}\delta(\omega_{k} - \omega), N_{c} = N(\omega_{c}), \{A, B\} = AB + BA.$$
(61)

This is the standard Lindblad form for a collision term of a quantum oscillator (see, e.g. [9]), where  $\Delta \omega_c$  is a shift of a frequency  $\omega_c$  due to interaction with a bath and  $J(\omega)$  is the bath spectral density.

Let us now consider Eqs. (59) with the contributions of the initial correlation. We will try to obtain the solution of Eqs. (59) for any timescale and, therefore, will not use the Markov approximation for these equations (Eq. (61) is only applicable for a large timescale). Considering, e.g., the equations for  $\langle a \rangle_S^t = Tr_S[a\rho_S(t)]$  and  $\langle a^+ \rangle_S^t = Tr_S[a^+\rho_S(t)]$  and using (57) and (58), we easily obtain

$$\langle aI_{S}(t)\rangle_{S} = Tr_{S}[aI_{S}(t)] = i\gamma_{i}(t)\langle a\rangle_{S}^{t}, \gamma_{i}(t) = \sum_{k} V_{k}^{2} e^{-i(\omega_{k}-\omega_{c})t}(1+N_{k})\frac{e^{(\omega_{c}-\omega_{k})\beta}-1}{\omega_{c}-\omega_{k}},$$

$$\langle a^{+}I_{S}(t)\rangle_{S} = -i\gamma_{i}^{+}(t)\langle a^{+}\rangle_{S}^{t}, \gamma_{i}^{+}(t) = \sum_{k} V_{k}^{2} e^{i(\omega_{k}-\omega_{c})t}N_{k}\frac{e^{(\omega_{k}-\omega_{c})\beta}-1}{\omega_{k}-\omega_{c}},$$

$$\langle aC_{S}(t)\rangle_{S} = -\gamma_{c}(t)\langle a\rangle_{S}^{t}, \gamma_{c}(t) = \int_{0}^{t} d\tau \sum_{k} V_{k}^{2} e^{-i(\omega_{k}-\omega_{c})\tau},$$

$$\langle a^{+}C_{S}(t)\rangle_{S} = -\gamma_{c}^{+}(t)\langle a^{+}\rangle_{S}^{t}, \gamma_{c}^{+}(t) = \gamma_{c}^{*}(t) = \int_{0}^{t} d\tau \sum_{k} V_{k}^{2} e^{i(\omega_{k}-\omega_{c})\tau},$$
(62)

Thus, we have from Eqs. (59) with account for initial correlations

$$\frac{\partial \langle a \rangle_{S}^{t}}{\partial t} = -i\omega_{c} \langle a \rangle_{S}^{t} - iTr_{s} \{ [a, H_{ext}^{S}(t)]\rho_{S}(t)] \} + i\gamma_{i}(t) \langle a \rangle_{S}^{t} - \gamma_{c}(t) \langle a \rangle_{S}^{t},$$

$$\frac{\partial \langle a^{+} \rangle_{S}^{t}}{\partial t} = i\omega_{c} \langle a^{+} \rangle_{S}^{t} - iTr_{s} \{ [a^{+}, H_{ext}^{S}(t)]\rho_{S}(t)] \} - i\gamma_{i}^{+}(t) \langle a^{+} \rangle_{S}^{t} - \gamma_{c}^{+}(t) \langle a^{\dagger} \langle \mathfrak{H}_{S}^{t} \rangle \}$$

In order to consider Eqs. (63), we should introduce the external force in the Hamiltonian (55), e.g., as

$$H_{ext}^{S}(t) = E_0 \exp(i\omega_d t)a + E_0 \exp(-i\omega_d t)a^+,$$
(64)

where  $E_0$  is the strength of the external driving field with frequency  $\omega_d$ . Then, the inhomogeneous terms in Eqs. (63) are

$$iTr_{s}\{[a, H_{ext}^{S}(t)]\rho_{S}(t)]\} = iE_{0}\exp(-i\omega_{d}t),$$
  
$$iTr_{s}\{[a^{+}, H_{ext}^{S}(t)]\rho_{S}(t)]\} = -iE_{0}\exp(i\omega_{d}t).$$
 (65)

For an equilibrium two-time correlation function  $\langle \alpha_S(0)a(t)\rangle$  and  $\langle \alpha_S(0)a^+(t)\rangle$ (see (13)) with  $\rho_{tot} = \rho_{eq}$  (40) we have

$$\langle \alpha_S(0)a(t) \rangle_{eq} = Tr_S\{a[\rho_S^{\alpha}(t)]_{eq}\}, \langle \alpha_S(0)a^+(t) \rangle_{eq} = Tr_S\{a^+[\rho_S^{\alpha}(t)]_{eq}\},$$
  
$$[\rho_S^{\alpha}(t)]_{eq} = Tr_B e^{Lt} \rho_{eq} \alpha_S(0).$$
(66)

In this correlation functions case, Eqs. (62) hold with the substitution  $\langle a \rangle_S^t \rightarrow \langle \alpha_S(0)a(t) \rangle, \langle a^+ \rangle_S^t \rightarrow \langle \alpha_S(0)a^+(t) \rangle$ . Therefore, the equations for these correlation functions are

$$\frac{\partial \langle \alpha_S(0)a(t) \rangle_{eq}}{\partial t} = \gamma(t) \langle \alpha_S(0)a(t) \rangle_{eq}, \gamma(t) = -i\omega_c + i\gamma_i(t) - \gamma_c(t),$$
  
$$\frac{\partial \langle \alpha_S(0)a^+(t) \rangle_{eq}}{\partial t} = \gamma^+(t) \langle \alpha_S(0)a^+(t) \rangle_{eq}, \gamma^+(t) = i\omega_c - i\gamma_i^+(t) - \gamma_c^+(t)67)$$

It is worth noting, that Eqs. (63) are inhomogeneous due to the driving Hamiltonian  $H_{ext}^{S}(t)$ , whereas Eqs. (67) for correlations functions are the homogeneous ones.

In order to illustrate the influence of initial correlations on the evolution process, let us consider the more simple homogeneous Eqs. (67), which describe the dynamics of the cavity field fluctuations. They can be easily solved and the result is

$$\langle \alpha_{S}(0)a(t) \rangle_{eq} = \exp[\Gamma(t)] \langle \alpha_{S}(0)a(0) \rangle_{eq}, \Gamma(t) = \int_{0}^{t} dt' \gamma(t') = -i\omega_{c}t + i\Gamma_{i}(t) - \Gamma_{c}(t),$$

$$\langle \alpha_{S}(0)a^{+}(t) \rangle_{eq} = \exp[\Gamma^{+}(t)] \langle \alpha_{S}(0)a^{+}(0) \rangle_{eq}, \Gamma^{+}(t) = \int_{0}^{t} dt' \gamma^{+}(t') = i\omega_{c}t - i\Gamma_{i}^{+}(t) - \Gamma_{c}^{+}(t),$$

$$\Gamma_{i}(t) = \int_{0}^{t} dt' \gamma_{i}(t'), \Gamma_{i}^{+}(t) = \int_{0}^{t} dt' \gamma_{i}^{+}(t'), \Gamma_{c}(t) = \int_{0}^{t} dt' \gamma_{c}(t'), \Gamma_{c}^{+}(t) = \int_{0}^{t} dt' \gamma_{c}^{+}(68).$$

where the relaxation functions in Eqs. (68) after integration over t' acquire, as it follows from (62) and (67), the following form

$$\Gamma_{i}(t) = i \sum_{k} V_{k}^{2} (1+N_{k}) [e^{(\omega_{c}-\omega_{k})\beta} - 1] \\ \times \frac{1 - \cos(\omega_{k} - \omega_{c})t + i\sin(\omega_{k} - \omega_{c})t}{(\omega_{k} - \omega_{c})^{2}},$$

$$\Gamma_{i}^{+}(t) = i \sum_{k} V_{k}^{2} N_{k} [e^{(\omega_{k}-\omega_{c})\beta} - 1] \\ \times \frac{1 - \cos(\omega_{k} - \omega_{c})t - i\sin(\omega_{k} - \omega_{c})t}{(\omega_{k} - \omega_{c})^{2}},$$

$$\Gamma_{c}(t) = \sum_{k} V_{k}^{2} [\frac{1 - \cos(\omega_{k} - \omega_{c})t + i\sin(\omega_{k} - \omega_{c})t}{(\omega_{k} - \omega_{c})^{2}} + \frac{t}{i(\omega_{k} - \omega_{c})}],$$

$$\Gamma_{c}^{+}(t) = \sum_{k} V_{k}^{2} [\frac{1 - \cos(\omega_{k} - \omega_{c})t - i\sin(\omega_{k} - \omega_{c})t}{(\omega_{k} - \omega_{c})^{2}} - \frac{t}{i(\omega_{k} - \omega_{c})}]. (69)$$

Using the spectral density of the Bosonic reservoir  $J(\omega)$  (61), the relaxation

functions (69) can be rewritten as

$$\Gamma_{i}(t) = \frac{1}{\hbar^{2}} \frac{i}{2\pi} \int_{-\omega_{c}}^{\infty} d\omega J(\omega_{c} + \omega) [1 + N(\hbar\omega_{c} + \hbar\omega)](e^{-\hbar\omega\beta} - 1) \\
\times \frac{1 - \cos(\omega t) + i\sin(\omega t)}{\omega^{2}},$$

$$\Gamma_{i}^{+}(t) = \frac{1}{\hbar^{2}} \frac{i}{2\pi} \int_{-\omega_{c}}^{\infty} d\omega J(\omega_{c} + \omega) N(\hbar\omega_{c} + \hbar\omega)(e^{\hbar\omega\beta} - 1) \\
\times \frac{1 - \cos(\omega t) - i\sin(\omega t)}{\omega^{2}},$$

$$\Gamma_{c}(t) = \frac{1}{\hbar^{2}} \frac{1}{2\pi} \int_{-\omega_{c}}^{\infty} d\omega J(\omega_{c} + \omega) [\frac{1 - \cos(\omega t) + i\sin(\omega)t}{\omega^{2}} + \frac{t}{i\omega}],$$

$$\Gamma_{c}^{+}(t) = \frac{1}{\hbar^{2}} \frac{1}{2\pi} \int_{-\omega_{c}}^{\infty} d\omega J(\omega_{c} + \omega) [\frac{1 - \cos(\omega t) - i\sin(\omega t)}{\omega^{2}} - \frac{t}{i\omega}],$$

$$N(\hbar\omega) = [\exp(\beta\hbar\omega) - 1]^{-1},$$
(70)

where we recover the Planck constant.

The formulae (68), (70) give the exact in the second order (Born) approximation solution to the system (cavity) equilibrium correlation functions accounting for initial correlations and valid on any timescale (in contrast to Eq. (61)). The evolution of these correlations functions is time-reversible (invariant to the  $t \to -t$ ,  $i \to -i$  replacement).

It is interesting to follow the evolution of the correlation functions with time in more detail. To this end, we introduce the following timescales

$$\overline{\omega}t \ll 1, \overline{\omega}t \gg 1,\tag{71}$$

where  $\overline{\omega}$  is the characteristic bath frequency, e.g.,  $\hbar \overline{\omega} \sim k_B T$  (see also the definition of the relaxation time after Eq. (59)), or introducing the coherence time  $t_{coh}$ ,

$$t \ll t_{coh}, t \gg t_{coh}, t_{coh} \sim \hbar/k_B T \tag{72}$$

Let us first consider the case of the small timescale  $\overline{\omega}t \ll 1$ . In this case, we can approximate  $\cos(\omega t) = 1 - \frac{1}{2}(\omega t)^2$  and  $\sin(\omega t) = \omega t$ . As a result we obtain

for the correlation functions (68)

$$\langle \alpha_{S}(0)a(t) \rangle_{eq} = \exp[\Gamma_{<}(t)] \langle \alpha_{S}(0)a(0) \rangle_{eq}, \overline{\omega}t \ll 1,$$

$$\Gamma_{<}(t) = -i(\omega_{c} + \Delta_{i}\omega_{c})t - \gamma t^{2},$$

$$\Delta_{i}\omega_{c} = \frac{1}{\hbar^{2}} \int_{-\omega_{c}}^{\infty} \frac{d\omega}{2\pi} \frac{J(\omega_{c} + \omega)}{\omega} [1 + N(\hbar\omega_{c} + \hbar\omega)](e^{-\hbar\omega\beta} - 1),$$

$$\gamma = \frac{1}{2\hbar^{2}} \int_{-\omega_{c}}^{\infty} \frac{d\omega}{2\pi} J(\omega_{c} + \omega) \{1 + [1 + N(\hbar\omega_{c} + \hbar\omega)](e^{-\hbar\omega\beta} - 1)\},$$

$$\langle \alpha_{S}(0)a^{+}(t) \rangle_{eq} = \exp[\Gamma_{<}^{+}(t)] \langle \alpha_{S}(0)a^{+}(0) \rangle_{eq}, \overline{\omega}t \ll 1,$$

$$\Gamma_{<}^{+}(t) = i[\omega_{c} - (\Delta_{i}\omega_{c})^{+}]t - \gamma^{+}t^{2},$$

$$(\Delta_{i}\omega_{c})^{+} = \frac{1}{\hbar^{2}} \int_{-\omega_{c}}^{\infty} \frac{d\omega}{2\pi} \frac{J(\omega_{c} + \omega)}{\omega} N(\hbar\omega_{c} + \hbar\omega)(e^{\hbar\omega\beta} - 1),$$

$$\gamma^{+} = \frac{1}{2\hbar^{2}} \int_{-\omega_{c}}^{\infty} \frac{d\omega}{2\pi} J(\omega_{c} + \omega)[1 - N(\hbar\omega_{c} + \hbar\omega)(e^{\hbar\omega\beta} - 1)]$$

$$(73)$$

which shows quadratic in time evolution caused not only by collisions (first terms in  $\gamma$ ,  $\gamma^+$ ) but also by initial correlations (second terms). There are also the cavity frequency shifts  $\Delta_i \omega_c$ ,  $(\Delta_i \omega_c)^+$  conditioned exclusively by initial correlations. We note, that due to the contribution of initial correlations, these quantities depend on temperature.

A more interesting is the situation on the kinetic timescale, which we define as

$$t \gtrsim \tau_{rel} \gg t_{coh} \sim \frac{1}{\overline{\omega}} \sim \frac{\hbar}{k_B T},$$
(74)

where the coherence time  $t_{coh}$  is defined by (72). The time hierarchy (74) is supposed to be realized in the considered case of a weak system-environment interaction. On the kinetic timescale we can take a limit  $t \to \infty$  and use in Eqs. (70) the following representations of  $\delta$ - function

$$\lim_{t \to \infty} \frac{1 - \cos(\omega t)}{\omega^2} = \pi \delta(\omega) |t|, \lim_{t \to \infty} \frac{\sin(\omega t)}{\omega} = \pi \delta(\omega)$$
(75)

Then, the relaxation functions (70) become

$$\Gamma_{i}(t) = \Phi(\beta) = \tau_{rel}^{-1} \hbar [1 + N(\hbar\omega_{c})]\beta, \tau_{rel}^{-1} = \frac{J(\omega_{c})}{2\hbar^{2}},$$

$$\Gamma_{i}^{+}(t) = \Phi^{+}(\beta) = \tau_{rel}^{-1} \hbar N(\hbar\omega_{c})\beta,$$

$$\Gamma_{c}(t) = \tau_{rel}^{-1} |t| + i\Delta\omega_{c}t, \Gamma_{c}^{+}(t) = \tau_{rel}^{-1} |t| - i\Delta\omega_{c}t,$$

$$\Delta\omega_{c} = -\frac{1}{2\pi\hbar^{2}} P \int_{-\omega_{c}}^{\infty} d\omega \frac{J(\omega_{c} + \omega)}{\omega},$$

$$t \gg 1/\overline{\omega},$$
(76)

where we observed that  $\int_{-\omega_c}^{\infty} d\omega J(\omega_c) \frac{\delta(\omega)}{\omega} = \int_{-\infty}^{\infty} d\omega J(\omega_c) \frac{\delta(\omega)}{\omega} = 0$  and assumed that the integral over  $\omega$  is meant as its principal value. Note, that the frequency shift here  $\Delta \omega_c$  coincides with that defined in (61) (where  $\hbar = 1$ ).

Thus, the time evolution of the system correlation functions on the large timescale is given by the following expressions (see (68))

$$\langle \alpha_S(0)a(t)\rangle_{eq} = \exp[i\Phi(\beta)]\exp[-i\widetilde{\omega}_c t - \tau_{rel}^{-1}|t|]\langle \alpha_S(0)a(0)\rangle_{eq} \langle \alpha_S(0)a^+(t)\rangle_{eq} = \exp[-i\Phi^+(\beta)]\exp[i\widetilde{\omega}_c t - \tau_{rel}^{-1}|t|]\langle \alpha_S(0)a^+(0)\rangle_{eq}, \widetilde{\omega}_c = \omega_c + \Delta\omega_c, t \gg 1/\overline{\omega}.$$

$$(77)$$

This is the main result of this section: Equations (77) differ from the standard ones (see, e.g., [24]) by the extra phase factors  $\exp[i\Phi(\beta)]$  and  $\exp[-i\Phi^+(\beta)]$ which emerge due to initial correlations (it is easy to verify, that using the Lindblad equation (61) in the absence of initial correlations, we can obtain the result (77) with no extra phase factors). These extra factors can influence the observables, in the considered example associated with the cavity field fluctuations. The result (77) also shows, that although the initial correlations influence the relaxation with time process (see (68) and (70)), on the large timescale they cease to do that ( $\Phi(\beta)$  does not depend on time) but can contribute to the observables. It is also interesting to note, that Eqs. (77) are not time-reversible (although Eqs. (68) are) and this is due to initial correlations. If we disregard initial correlations, Eqs. (77) become time-reversible. Thus, irreversibility on the large (asymptotic) timescale ( $t \to \infty$ ) emerges here due to initial correlations.

Generally, the system correlation function determines the response of the system on the applied driving force, and, if the initial correlations survive on the kinetic timescale as in the considered above case, they can influence the kinetic coefficients. For example, the additional phase factor (of the type given by (77)) influences the polaron mobility, as it was demonstrated earlier in work [21] treating the dynamics of an electron interacting with a bath of phonons in the polar crystals.

#### 6 Summary

We have presented a novel approach to the dynamics of a system coupled with a bath. The approach is based on the exact homogeneous (completely closed) generalized master equations for the relevant parts of a system's statistical operator and correlation function (Eqs. (11) and (16)). These equations exactly follow from the inhomogeneous Nakajima-Zwanzig GME without any "molecular chaos"-type approximations and account for initial correlations in the kernel governing their evolution in time. For a system in a bath, the obtained equations are equivalent to the homogeneous equations for a system of interest statistical operator and correlation function (Eqs. (28) and (31)). In the second (Born) approximation in the system-bath interaction these equations reduce to the time-local ones (39) which describe the evolution of a system interacting with a bath at any timescale and any initial state of the entire system (system +bath).

Equations (39) are the instruments for investigation of a dynamics of a specific system in a bath. They have been further specialized for the important and quite realistic initial Gibbs equilibrium state for the whole system and the bath as a Boson field. The corresponding initial correlation and collision terms are given by (50) and (53).

As an application, the quantum oscillator (cavity mode) interacting with a Boson field (Fano-like model) has been considered. The equations for an oscillator statistical operator and correlation function have been obtained (Eqs. (59)) with the initial correlations and collision terms given by Eqs. (57) and (58). The solution (68) for the cavity field fluctuations (correlation functions), where the generators are defined by Eqs. (70), has been obtained. This solution describes the evolution of the fluctuations of localized cavity mode at any timescale and shows the influence of initial correlations on the evolution process. For small times, the evolution is quadratic in time, while at the large timescale (see Eq. (77)), when the Markov approximation is applicable (actually at  $t \to \infty$ ), the initial correlations cease to influence the dynamics of the cavity mode fluctuations, but the initial correlations terms survive and result in the additional phase factor in the correlation functions, which distinguishes this result from the standard one obtained from the corresponding Lindblad equation,

### References

- N. N. Bogoliubov, Problems of Dynamical Theory in Statistical Physics (Gostekhizdat, Moscow, 1946, in Russian); English transl.: Stud. Statist. Mech. 1 (North-Holland, Amsterdam, 1962).
- [2] N. G. van Kampen, J. Stat. Phys. **115**, 1057 (2004).
- [3] V. F. Los, J. Stat. Mech. 013107 (2024).
- [4] V.F. Los, J Stat. Mech. 1, 013211 (2022).
- [5] S. Nakajima, Progress of Theoretical Physics **20**, 948 (1958).
- [6] R. Zwanzig, J. Chem. Phys. 33, 1338 (1960).
- [7] D. Wallace, Probability and irreversibility in modern statistical mechanics: Classical and quantum, 2021, arXiv: 2104.11223v1.
- [8] M. Kac, Probability and Related Topics in Physical Sciences (Interscience, New York, 1959).
- [9] H. -P. Breuer, F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, New York, 2002).

- [10] G. Lindblad, Commun. Math. Phys. 48, 119 (1976).
- [11] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, J. Math. Phys. 17, 821 (1976).
- [12] A. G. Redfield, IBM J. Res. Dev. 1, 19 (1957).
- [13] K. Blum, Density Matrix Theory and Applications (Plenum Press, New York, London, 1981).
- [14] Devashish Tupkary, Abhishek Dhar, Manas Kulkarni, and Archak Purkayastha, Phys. Rev. A 105, 032208 (2022).
- [15] Hua-Tang Tan and Wei-Min Zhang, Phys. Rev. A 83, 032102 (2011).
- [16] Gerardo A. Paz-Silva, Michael J. W. Hall, and Howard M. Wiseman, Phys. Rev. A 100, 042120 (2019).
- [17] S. Alipour, A. T. Rezakhani, A. P. Babu, K. Mølmer, M. Möttönen, and T. Ala-Nissila, Phys. Rev. X 10, 041024 (2020).
- [18] I. Prigogine, Non-Equilibrium Statistical Mechanics (Interscience Publishers, New York, 1962).
- [19] F. Shibata, Y. Takahashi, and N. Hashitsume, J. Stat. Phys. 17, 171 (1977).
- [20] F. Shibata and T. Arimitsu, J. Phys. Soc. Jap. 49, 891 (1980).
- [21] V. F. Los, J. Stat. Phys. 168, 857 (2017).
- [22] V. F. Los, Physica A **503**, 476 (2018).
- [23] U. Fano, Phys. Rev. **124**, 1866 (1961).
- [24] Chan U Lei and Wei-Min Zhang, arXiv: 1011.4570v2.