NOTES ON HYPERBOLIC BRANCHING BROWNIAN MOTION

WOLFGANG WOESS

ABSTRACT. Euclidean branching Brownian motion (BBM) has been intensively studied during many decades by renowned researchers. BBM on hyperbolic space has received less attention. A profound study of Lalley and Sellke (1997) provided insight on the recurrent, resp. transient regimes of BBM on the Poincare' disk. In particular, they determined the Hausdorff dimension of the limit set on the boundary circle in dependance on the fission rate of the branching particles. In the present notes, some further features are exhibited, such as the rate of the maximal hyperbolic distance to the starting point and the behaviour of the empiricial distributions of the branching population, as time goes to infinity.

1. INTRODUCTION

This is ongoing work; these notes present its state of mid 2024.

Euclidean branching Brownian motion on \mathbb{R} has been an intensively studied topic. A good reference for the earlier developments is the comprehensive monograph by ATHREYA AND NEY [1]. In short, a particle performs Brownian motion for an exponentially distributed time and then fissions in two new particles, each of which continues its own Brownian motion independently of the other for exponential time, then in turn fissions in two, and so on. Primary object of studies has been the evolution of the random population at time t, as $t \to \infty$. One of the many interesting features concerns the position of the rightmost particle at time t. Famous work of MCKEAN [24] showed that the distribution of the rightmost particle is directly linked with the travelling wave equation studied in the famous work of KOLMOGOROV, PETROVSKI AND PISCOUNOV [21].

Similarly, there is the study of branching random walk (BRW) in discrete time, where particles evolve according to a Galton-Watson process: at the points of fission, the new particles perform independent steps according to a given random walk. Branching random walk on \mathbb{R} appeared, for example, in the monograph by HARRIS [15, §III.16]. Both models have evolved significantly, and both are examples of tree-indexed Markov processes in the sense of BENJAMINI AND PERES [3], where the linear time is replaced by a - possibly random - tree (discrete or continuous).

Here, the focus is on branching Brownian motion (BBM) on the hyperbolic disk. That is, the motion of the particles follows the continuous time Markov process whose infinitesimal generator is $\frac{1}{2}\mathfrak{L}$, where \mathfrak{L} is the Laplace-Beltrami operator of the unit disk equipped with the Poincaré metric (or equivalently, the upper half plane model). It has not been as widely considered as several other branching random processes. A very significant in-depth study is due to LALLEY AND SELLKE [23]. They considered hyperbolic BBM where the random

Date: May 12, 2024.

Key words and phrases. Hyperbolic disk, hyperbolic Laplacian, branching Brownian motion, maximal distance, empirical distributions, boundary convergence.

times between successive fissions are exponentially distributed with parameter $\lambda > 0$. There is a phase transition: for $\lambda \leq 1/8$ (transient phase), each compact subset of \mathbb{D} is no more visited by the branching population from some random time onwards, while for $\lambda > 1/8$ (recurrent phase) each open subset of \mathbb{D} continues to be visited by the population. The main results of [23] concern the random *limit set* Λ , that is, the set of accumulation points of the trace of the population on the boundary $\partial \mathbb{D}$, the unit circle. It is the full circle when $\lambda > 1/8$, while it is a perfect set with *Hausdorff dimension* $(1 - \sqrt{1 - 8\lambda})/2$ when $\lambda \leq 1/8$.

An analogous phenomenon was exhibited for branching random walk on regular trees, resp. free groups, by LIGGETT [22] and HUETER AND LALLEY [16]. This was extended to BRW on free products of groups by CANDELLERO, GILCH AND MÜLLER [8], and very recently to BRW on hyperbolic groups by SIDORAVICIUS, WANG AND XIANG [26] and subsequently to BRW on relatively hyperbolic groups by DUSSAULE, WANG AND YANG [12]. For BRW on finitely generated groups, resp. transitive graphs, this study has been accompanied by the investigation of the *trace*, that is, the subgraph spanned by all points visited by the BRW. See BENJAMINI AND MÜLLER [2], CANDELLERO AND ROBERTS, [10], GILCH AND MÜLLER [13], HUTCHCROFT [17] as well as [26] and [12].

Returning to hyperbolic BBM, in the present notes some results are added to the profound study of [23]. For the maximal hyperbolic distance from the origin of a particle at time t, we show that its rate is the same as for the comparison process which is Euclidean BBM where the underlying one-dimensional BM has drift 1/2. Then we consider the empirical distributions of hyperbolic BBM: these are the normalised occupation measures of the population at the times t > 0.

Following a suggestion by V. A. Kaimanovich, the asymptotic behaviour of these random distributions was recently studied for BRW on graphs in parallel work of KAIMANOVICH AND WOESS [18] and CANDELLERO AND HUTCHCROFT [9]. In the Euclidean setting of BRW and BBM, they had been studied since the mid 1960ies, see e.g. NEY [25], STAM [27], KAPLAN AND ASMUSSEN [19], UCHIYAMA [28], BIGGINS [4], while [18] and [9] are more relevant in a "non-amenable" setting.

Here, variants of CLT-type asymptotics of the empricial distributions of hyperbolic BBM are derived once more via the Euclidean comparison process. Then then the average rate of escape, that is, the rate of the average distance from the origin of the particles at time t, is shown to coincide with the one of (non-branching) hyperbolic BM. It is proved that the empirical distributions converge weakly (on the closed disk) to a random limit distribution on the boundary, that is, the unit circle. Some properties of the limit distribution are derived, and various open questions are posed.

In these notes, the author has chosen to present the initial parts in a rather broad way. This concerns, in paricular, the construction of the underlying time tree (Yule tree) and the chosen notation, which differs a bit from the previous mainstream.

Acknowledgements. The author acknowledges helpful email exchange with John Biggins, Maury Bramson, Nicola Kistler, Steve Lalley, Sebastian Müller, Enzo Orsingher, Yuichi Shiozawa, Anton Wakolbinger (personal contact) and Ofer Zeitouni.

2. The Yule tree

The introduction of the continuous random population tree described in this section goes back to YULE [29].

Consider the binary tree $\{\mathfrak{l}, \mathfrak{r}\}^*$ consisting of all binary sequences (words) $\mathfrak{v} = \mathfrak{s}_1 \cdots \mathfrak{s}_n$ with $n \geq 0$ and $\mathfrak{s}_k \in \{\mathfrak{l}, \mathfrak{r}\}$.¹ For n = 0, this is the empty sequence ϵ . If \mathfrak{v} has the form $\mathfrak{v} = \mathfrak{u}\mathfrak{s}$ with $\mathfrak{s} \in \{\mathfrak{l}, \mathfrak{r}\}$, then the *predecessor* of \mathfrak{v} is $\mathfrak{v}' = \mathfrak{u}$, and \mathfrak{v} is one of the two successors of \mathfrak{u} . The edges of the tree are all $[\mathfrak{v}', \mathfrak{v}]$, where $\mathfrak{v} \neq \epsilon$. We augment this tree by an additional vertex α which is only connected to ϵ , so that $\epsilon' = \alpha$. Now consider a sequence of i.i.d. random variables $\ell_{\mathfrak{v}}, \mathfrak{v} \in \{\mathfrak{l}, \mathfrak{r}\}^*$ having exponential distribution with parameter $\lambda > 0$. For the moment, we can realise it for example on the product space

$$\left(\Omega^{\mathrm{Yule}}, \mathcal{A}^{\mathrm{Yule}}, \mathbb{P}^{\mathrm{Yule}}\right) = \bigotimes_{\mathfrak{v} \in \{\mathfrak{l}, \mathfrak{r}\}^*} \left(\mathbb{R}_+, \mathcal{B}_{\mathbb{R}^+}, Exp_\lambda\right)_{\mathfrak{v}},$$

where each factor is a copy of the probability space consisting of the positive real half-line with the Borel sigma-algebra and the exponential distribution with parameter λ , so that $\ell_{\mathfrak{v}}$ is the projection on the *v*-coordinate. Later on, it will be embedded into a bigger probability space.

We then get a random tree \mathcal{T} , a 1-complex where each edge $[\mathfrak{v}', \mathfrak{v}]$ is an interval with the random length $\ell_{\mathfrak{v}}$, and we write the edge as

(1)
$$[\mathfrak{v}',\mathfrak{v}] = \{\tau = \mathfrak{v}_s : 0 \le s \le \ell_{\mathfrak{v}}\},\$$

where v_s is the point in the interval at distance s from \mathfrak{v}' . Then the length (= distance from α) of a vertex \mathfrak{v} is defined recursively by $|\alpha| = 0$ and $|\mathfrak{v}| = |\mathfrak{v}'| + \ell_{\mathfrak{v}}$, and the length of $\tau = v_s \in [v^-, v]$ is $|\tau| = |v^-| + s$.

This is the Yule tree. It is interpreted as a genealogical tree, where α is the "ancestor" at time 0, an thinking of it as a particle, after time ℓ_{ϵ} it fissions in two particles. The timelines of the new particles are the edges $[\epsilon, \mathfrak{l}]$ and $[\epsilon, \mathfrak{r}]$, respectively, and after times $\ell_{\mathfrak{l}}$, resp. $\ell_{\mathfrak{r}}$, each of them fissions again in 2 particles. Recursively, a particle at the end of its timeline $[\mathfrak{v}', \mathfrak{v}]$ fissions in two, whose new timelines are the edges $[\mathfrak{v}, \mathfrak{v}\mathfrak{l}]$ and $[\mathfrak{v}, \mathfrak{v}\mathfrak{r}]$, respectively.

The population at time $t \ge 0$ is the set $\mathcal{T}(t) = \mathcal{T}_{\epsilon}(t)$ of all particles (\equiv vertices or interior points on the edges) at distance t from α . We write

$$N(t) = N_{\epsilon}(t) = |\mathcal{T}(t)|$$

for their number. Note that by continuity of the exponential distribution, at any time $t \ge 0$, with probability 1 there is at most one vertex v of \mathcal{T} with |v| = t. That is, no two fissions take place simultaneously.

For any $\mathfrak{u} \in {\mathfrak{l}, \mathfrak{r}}^*$ let $\mathcal{T}_{\mathfrak{u}}$ be the subtree of \mathcal{T} in which \mathfrak{u} has the role of ϵ in the above description and \mathfrak{u}' the role of α . In other words, it is spanned by the vertex set

$$\{\mathfrak{u}'\} \cup \{\mathfrak{u}\mathfrak{v} : \mathfrak{v} \in \{\mathfrak{l},\mathfrak{r}\}^*\}.$$

¹To avoid confusion with numbers, we avoid the binary symbols 0, 1. The symbols l, \mathfrak{r} and \mathfrak{s} stand for "left", "right" and "side", respectively.

We write $\mathcal{T}_{\mathfrak{u}}(t)$ for the associated part of the population, that is, the set of elements of $\mathcal{T}_{\mathfrak{u}}$ at distance t from u^- , and $N_u(t)$ for their number.

For any subset $U \subset {\{\mathfrak{l}, \mathfrak{r}\}}^*$ of vertices which is prefix-free (that is, no element of U is an initial part of another element in U), then the trees $\mathcal{T}_{\mathfrak{u}}$, $\mathfrak{u} \in U$ are i.i.d. In particular, all the generation sizes $N_u(t)$, $\mathfrak{u} \in {\{\mathfrak{l}, \mathfrak{r}\}}^*$ have the same distribution on \mathbb{N} . The following is well-known (even for more general models); we provide a short proof.

Lemma 2.1. For any $t \ge 0$, the population size at time t has geometric distribution:

$$\mathbb{P}[N(t) = n] = e^{-\lambda t} \left(1 - e^{-\lambda t}\right)^{n-1}, \quad n \in \mathbb{N}.$$

In particular, the expected population size is $\mathbb{E}(N(t)) = e^{\lambda t}$.

Proof. We have

$$N_{\epsilon}(t) = \begin{cases} 1, & \text{if } \ell_{\epsilon} \ge t \\ N_{\mathfrak{l}}(t - \ell_{\epsilon}) + N_{\mathfrak{r}}(t - \ell_{\epsilon}), & \text{if } \ell_{\epsilon} < t. \end{cases}$$

Therefore the characteristic function (in the variable x) is

$$\varphi_{N(t)}(x) = \mathbb{E}\Big(\exp\big(\mathfrak{i}\,x\,N_{\epsilon}(t)\big)\Big) = e^{\mathfrak{i}\,x}\,\mathbb{P}[\ell_{\epsilon} \ge t] + \mathbb{E}\Big(\exp\big(\mathfrak{i}\,x\,N_{\mathfrak{l}}(t-\ell_{\epsilon})+\mathfrak{i}\,x\,N_{\mathfrak{r}}(t-\ell_{\epsilon})\big)\,\mathbf{1}_{[\ell_{\epsilon}< t]}\Big)$$
$$= e^{\mathfrak{i}\,x-\lambda t} + \int_{0}^{t}\varphi_{N(t-s)}(x)^{2}\,\lambda e^{-\lambda s}\,ds\,.$$

Just for the purpose of this proof, set $f(t) = e^{\lambda t} \varphi_{N(t)}(x)$. Then the above equation transforms into the integral equation

$$f(t) = e^{ix} + \lambda \int_0^t f(s)^2 e^{-\lambda s} ds$$

Thus, $f(t) = 1/(e^{-ix} + e^{-\lambda t} - 1)$, and

$$\varphi_{N(t)}(x) = \frac{e^{ix} e^{-\lambda t}}{1 - e^{ix}(1 - e^{-\lambda t})},$$

which we recognise as the characteristic function of the geometric distribution with parameter (success probability) $p = e^{-\lambda t}$.

The population $\mathcal{T}(t)$ at time t can be viewed as a cross-section of \mathcal{T} at level t. For the tree viewed as a spatial generalisation of time, it can be interpreted to have a role analogous to the one of a stopping time. If we cut the tree \mathcal{T} along that cross-section, we obtain the subtree $\mathcal{T}(\prec t)$ of all elements τ with $|\tau| \leq t$. We let $V(\mathcal{T}(\prec t)) = {\mathfrak{l}, \mathfrak{r}}^* \cap \mathcal{T}(\prec t)$ be the random set consisting of those vertices of the original binary tree which are part of $\mathcal{T}(\prec t)$. By Lemma 2.1, $V(\mathcal{T}(\prec t))$ is finite with probability 1.

The sigma-algebra $\mathcal{F}_t^{\text{Yule}}$ comprising the information inherent to the Yule tree up to and including time t is generated by all random variables ℓ_v which intervene in the construction of $\mathcal{T}(\prec t)$. The following is well known.

Proposition 2.2. The process $(N(t) e^{-\lambda t})_{t\geq 0}$ is a martingale with respect to the filtration $(\mathcal{F}_t^{Yule})_{t\geq 0}$, and there exists an almost surely positive random variable W such that

$$\lim_{t \to \infty} N(t) e^{-\lambda t} = W \quad almost \ surrely \ and \ in \ L^1.$$

3. Hyperbolic disk and Brownian motion

A standard model of two-dimensional hyperbolic space is the *Poincaré disk* with the hyperbolic length element and resulting metric

(2)
$$ds = \frac{2\sqrt{dx^2 + dy^2}}{1 - |z|^2} \quad \text{and} \quad d(z, w) = \log \frac{|1 - z\bar{w}| + |z - w|}{|1 - z\bar{w}| - |z - w|}.$$

Its orientation preserving isometry group consists of all Möbius transformations of the form

(3)
$$gz = \frac{az+c}{\bar{c}z+\bar{a}}, \quad a,c \in \mathbb{C}, \ |a|^2 - |c|^2 = 1.$$

Together with the reflection $z \mapsto -\bar{z}$, it generates the full isometry group. Of course, these transformations also act on the boundary of \mathbb{D} , the unit circle $\partial \mathbb{D}$. We shall use the group \mathcal{A} of all transformations as in (3) which fix the boundary point 1, that is, $a + c \in \mathbb{R}$. It acts simply transitively on \mathbb{D} . In particular, For each $z_0 \in \mathbb{D}$, there is a unique element $g_{z_0} \in \mathcal{A}$ which maps 0 to z_0 . It is given by

(4)
$$g_{z_0}z = \frac{(1-z_0)z + (z_0 - |z_0|^2)}{(\bar{z}_0 - |z_0|^2)z + (1-\bar{z}_0)}.$$

The hyperbolic Laplace (or Laplace-Beltrami) operator in the variable z = x + i y is

(5)
$$\mathfrak{L} = \frac{(1-|z|^2)^2}{4} \left(\partial_x^2 + \partial_y^2\right).$$

It is self-adjoint on $L^2(\mathbb{D}, \mathsf{m})$, where $\mathsf{m} = \mathsf{m}_{\mathbb{D}}$ is the hyperbolic measure,

(6)
$$d\mathbf{m}(z) = \frac{4dz}{(1-|z|^2)^2}.$$

The infinitesimal generator of hyperbolic Brownian motion $(B_t)_{t\geq 0}$ is $\frac{1}{2}\mathfrak{L}$. (We remark that the factor $\frac{1}{2}$ is used because in the analogous Euclidean setting we want that Brownian motion at time t = 1 has standard normal distribution.)

One can see the latter as a version of the standard two-dimensional Euclidean Brownian motion slowed down as it gets close to the unit circle. Responsible for slowing down is the factor $\frac{(1-|z|^2)^2}{4}$ which only depends on the Euclidean distance of the current position z from the circle.

Hyperbolic Brownian motion (BM) has independent increments and continuous trajectories (almost surely). Since the hyperbolic Laplacian commutes with all hyperbolic isometries, hyperbolic BM is invariant under the latter: if g is as in (3) and (B_t) is (a version of) hyperbolic BM starting at $z_0 \in \mathbb{D}$ then (gB_t) is (a version of) hyperbolic BM starting at gz_0 . Equivalently, the heat kernel $p_t(\cdot, \cdot)$ with respect to **m** associated with $\frac{1}{2}\mathfrak{L}$ is invariant under the diagonal actions of every hyperbolic isometry. We write P_t for the associated transition

operator of hyperbolic BM: for any measurable set $K \subset \mathbb{D}$ and $z \in \mathbb{D}$, resp., measurable function $f : \mathbb{D} \to \mathbb{R}$,

(7)
$$P_t(z,K) = \int_K p_t(z,w) \, d\mathbf{m}(w) \quad \text{and} \quad P_t f(z) = \int_{\mathbb{D}} p_t(z,w) f(w) \, d\mathbf{m}(w) \, ,$$

whenever that integral is well defined.

Now let $(\Omega_b, \mathcal{A}_b, \mathbb{P}_b)$ be a suitable probability space on which hyperbolic BM starting at the origin is defined. With starting point 0, as $t \to \infty$, the process converges almost surely to a $\partial \mathbb{D}$ -valued random variable B_{∞} . The distribution of B_{∞} , being rotation invariant, is equidistribution on the unit circle: $d\xi = \frac{1}{2\pi} e^{i\phi} d\phi$, where $d\phi$ is the Lebesgue measure on $[-\pi, \pi]$. (The elements of $\partial \mathbb{D}$ are denoted ξ, η , etc.)

When the starting point is z_0 then, working on the same probability space, (a model of) the limit random variable is $g_{z_0}B_{\infty}$. The density of its distribution ν_{z_0} with respect to $d\xi$ is the *Poisson kernel*

(8)
$$\Pi(z_0,\xi) = \frac{1-|z_0|^2}{|\xi-z_0|^2} \qquad (z_0 \in \mathbb{D}, \ \xi = e^{i\phi} \in \partial \mathbb{D})$$

Since $d(B_t, o) \to \infty$ almost surely, we are also interested in the speed. It is linear, and to understand it, it may be useful to pass to two other models of hyperbolic space.

The second one, besides the disk, is the upper half plane model $\mathbb{H} = \{u + \mathfrak{i}v : u, v \in \mathbb{R}, v > 0\}$. The metric $d = d_{\mathbb{D}}$ of (2) is transported to the metric $d_{\mathbb{H}}$ via the Möbius map from \mathbb{D} to \mathbb{H}

(9)
$$z \mapsto \mathfrak{i} \frac{1+z}{1-z}.$$

It maps 0 to i, and the boundary points -1 to 0 and 1 to $i\infty$, and

(10)
$$d_{\mathbb{H}}(z,w) = \log \frac{|z-\bar{w}| + |z-w|}{|z-\bar{w}| - |z-w|}$$

We shall often switch back and forth between \mathbb{D} and \mathbb{H} , and will mostly use unified notation o for our origin, that is, o = 0 in \mathbb{D} and $o = \mathfrak{i}$ in \mathbb{H} . We also remark here that in the upper half plane model, the group \mathfrak{A} of all transformations that fix the boundary point 1 in \mathbb{D} (that is, $a + c \in \mathbb{R}$ for a, c as in (3)) becomes, via conjugation with the map (9), the *affine group* af all transformations

$$z \mapsto az + b$$
, $a > 0$, $b \in \mathbb{R}$ $(z \in \mathbb{H})$.

(The a here is not the same as in (3).)

In the coordinates $(u, v) \in \mathbb{H}$, the hyperbolic Laplacian becomes $v^2(\partial^2 u + \partial^2 v)$. Then we make one more change of variables, setting $w = \log v$ to obtain the *logarithmic model*, where now $(u, w) \in \mathbb{R}^2$ and the hyperbolic Laplacian becomes

(11)
$$\mathfrak{L} = e^{2w}\partial_u^2 + \partial_w^2 - \partial_w.$$

Its projection on the vertical coordinate w is $\partial_w^2 - \partial_w$, so that $\frac{1}{2}(\partial_w^2 - \partial_w)$ is the infinitesimal generator of one-dimensional Euclidean Brownian motion with drift -1/2. Tracing this

back to the upper half plane and disk models, writing $B_t^{\mathbb{H}}$ and $B_t^{\mathbb{D}}$ for hyperbolic Brownian motion in the respective coordinates, when $B_0^{\mathbb{H}} = \mathfrak{i}$, resp. $B_0^{\mathbb{D}} = 0$,

(12)
$$\log(\Im B_t^{\mathbb{H}}) = \log \frac{1 - |B_t^{\mathbb{D}}|^2}{|1 - B_t^{\mathbb{D}}|^2} \sim N\left(-\frac{1}{2}t, t\right),$$

where (as usual) \Im denotes the imaginary part, ~ means "has distribution" and $N(a, s^2)$ is the normal distribution with mean a and variance s^2 . From this we get the following, where we can omit the superscript referring to the respective model.

Lemma 3.1. The following central limit theorem and rate of escape results hold, as $t \to \infty$:

$$\frac{d(B_t, B_0) - \frac{1}{2}t}{\sqrt{t}} \to \mathcal{N}(0, 1) \quad in \ law, \ and \quad \frac{d(B_t, B_0)}{t} \to \frac{1}{2} \quad almost \ surely.$$

Proof. As stated above (8), in terms of the disk model, it is well known that $B_t^{\mathbb{D}}$ converges almost surely to a limit random variable $B_{\infty}^{\mathbb{D}}$ with continuous distribution. Taking this to the upper half plane model, we get $B_{\infty}^{\mathbb{H}} \in \mathbb{R}$ and $\Im B_t^{\mathbb{H}} \to 0$ almost surely (since $\mathbb{P}[B_{\infty}^{\mathbb{H}} = \mathfrak{i} \infty] = 0$). Now, if $z = x + \mathfrak{i} y \in \mathbb{H}$ then an easy computation with the hyperbolic metric in \mathbb{H} as in (10) shows that

(13)
$$d_{\mathbb{H}}(x+\mathfrak{i}\,y,\mathfrak{i}\,) + \log y = \log \frac{1}{2} \Big(1+|z|^2 + \sqrt{(1+|z|^2)^2 - 4y^2} \Big)$$

We get that

$$d_{\mathbb{H}}(B_t, \mathfrak{i}) + \log(\Im B_t^{\mathbb{H}}) \to \log(1 + (B_{\infty}^{\mathbb{H}})^2)$$
 almost surely, as $t \to \infty$,

and the limit is almost surely finite. Combining this with (12) completes the proof.

We shall need estimates of the tail behaviour of the random variables $d(B_t, o), t > 0$. The heat kernel $p_t(o, z), z \in \mathbb{D}$ (resp., $\in \mathbb{H}$) associated with $\frac{1}{2}\mathfrak{L}$ only depends on R = d(z, o), and it has a uniform estimate in space and time, see DAVIES AND MANDOUVALOS [11, Thm. 3.1]². We state here the upper bound in a way which is suited for our purpose:

(14)

$$p_t(z,w) \le \frac{Const}{\sqrt{1+R}} \Psi\left(\frac{1+R}{t}\right) \exp\left(-\frac{(R+\frac{t}{2})^2}{2t}\right) \quad \text{for } R, t > 0, \text{ where}$$

$$R = d(z,w) \quad \text{and} \quad \Psi(x) = \begin{cases} x^{3/2}, & 0 \le x \le 1, \\ x^{1/2}, & x \ge 1. \end{cases}$$

For the following proposition, the author acknowledges a suggestion by Yuichi Shiozawa.

²Concerning the heat kernel, different normalisations of the Laplacian are an ongoing source of small confusion. Analysist typically use the heat kernel for \mathfrak{L} as given in (5). Some of them also omit the factor 1/4. In probability, we want that Euclidean BM at time t = 1 has variance 1, so that we use the heat kernel for $\frac{1}{2}\mathfrak{L}$. Although not explicitly stated, [11] uses the first of these three options, so that here we had to replace their t by t/2.

Proposition 3.2. Let $\mathcal{M} = \max\{d(B_t, B_0) : t \leq 1\}$. Then there is K > 0 such that for any c > 0

$$\mathbb{P}[\mathcal{M} \ge c] \le 2 \mathbb{P}[d(B_1, B_0) \ge c] \le K \exp\left(-\frac{(c - \frac{1}{2})^2}{2}\right).$$

Proof. We work in the upper half plane model, and we may suppose without loss of generality that $B_0 = B_0^{\mathbb{H}} = o \, (= \mathfrak{i})$. Here, we shall omit the sub- and superscripts \mathbb{H} . Recall from (14) that in any model, $(B_t)_{t\geq 0}$ is istropic, that is, its transition kernel only depends on time and hyperbolic distance: $(d(B_t, o))_{t\geq 0}$ is a Markov process on the state space $[0, \infty)$ with continuous trajectories. In \mathbb{H} , the point e^{-c} is at hyperbolic distance c from $o = \mathfrak{i}$. Thus, using (13), for s, t > 0,

(15)
$$\mathbb{P}[d(B_{s+t}, o) \ge c \mid d(B_s, o) = c] = \mathbb{P}[d(B_t, o) \ge c \mid B_0 = e^{-c}] \\ \ge \mathbb{P}[-\log(\Im B_t) \ge c \mid -\log(\Im B_0) = c] \ge 1/2$$

because by (12), if $-\log(\Im B_0) = c$ then $-\log(\Im B_t) \sim N(c + \frac{1}{2}t, t)$.

Now consider the a.s. finite stopping time

$$T_c = \inf\{t > 0 : d(B_t, o) = c\}$$

Then, using the strong Markov property, the fact that $d(B_{T_c}, o) = c$, and isotropy of hyperbolic BM,

$$\mathbb{P}[\mathcal{M} \ge c] = \mathbb{P}_0[T_c \le 1] \\ = \mathbb{P}[d(B_1, o) \ge c, T_c \le 1] + \mathbb{P}[d(B_1, o) < c, T_c \le 1] \\ = \mathbb{P}[d(B_1, o) \ge c] + \int_0^1 \mathbb{P}[d(B_1, o) < c \mid d(B_s, o) = c] \ d \mathbb{P}[T_c = s]$$

now applying (15)

$$\leq \mathbb{P}[d(B_1, o) \geq c] + \int_0^1 \mathbb{P}[d(B_1, o) \geq c \mid d(B_s, o) = c] \ d\mathbb{P}[T_c = s]$$
$$= 2\mathbb{P}[d(B_1, o) \geq c].$$

This proves the first of the two inequalities. The second is going to be derived from (14). Note that for t = 1, we have to use $\Psi(x) = x^{1/2}$ in that heat kernel estimate.

We return to the disk model. We can express the hyperbolic measure of (6) first in terms of Euclidean polar coordinates (r, φ) with r < 1 and then replace r = |z| by $R = d(z, o) = \log \frac{1+r}{1-r}$. In the coordinates (R, φ) , we get

(16)
$$d\mathbf{m}(z) = \sinh R \ dR \ d\varphi \quad \text{in } \mathbb{D}.$$

Then

$$\mathbb{P}[d(B_1, o) \ge c] = \int_{\{z \in \mathbb{D} : d(z, o) \ge c\}} p_1(0, z) \, d\mathsf{m}(z)$$

$$\le Const \int_{-\pi}^{\pi} \int_c^{\infty} \exp\left(-\frac{(R + \frac{1}{2})^2}{2}\right) \frac{e^R - e^{-R}}{2} \, dR \, d\varphi$$

$$\le Const' \int_c^{\infty} \exp\left(-\frac{(R - \frac{1}{2})^2}{2}\right) \, dR$$

$$\le Const'' \exp\left(-\frac{(c - \frac{1}{2})^2}{2}\right)$$

for a suitable constant Const''.

4. Hyperbolic branching Brownian motion

We now construct hyperbolic branching Brownian motion. Whenever it is suitable, we can switch between the different models of hyperbolic plane, but primarily we have the disk model in mind.

We need a probability space $(\Omega, \mathcal{A}, \mathbb{P})$ on which one can realise countably many i.i.d. random variables $\ell_{\mathfrak{v}}$, $\mathfrak{v} \in {\mathfrak{l}, \mathfrak{r}}^*$ and, independently of the latter, countably many independent hyperbolic Brownian motions $(B_t^{\mathfrak{v}})$, $\mathfrak{v} \in {\mathfrak{l}, \mathfrak{r}}^*$, each one starting at 0. If $(\Omega_b, \mathcal{B}_b, \mathbb{P}_b)$ is one probability space on which hyperbolic BM can be defined, then we can take the product space

$$(\Omega, \mathcal{A}, \mathbb{P}) = (\Omega^{\text{Yule}}, \mathcal{A}^{\text{Yule}}, \mathbb{P}^{\text{Yule}}) \otimes (\Omega_B, \mathcal{A}_B, \mathbb{P}_B), \text{ where}$$
$$(\Omega_B, \mathcal{A}_B, \mathbb{P}_B) = \bigotimes_{\mathfrak{v} \in \{\mathfrak{l}, \mathfrak{r}\}^*} (\Omega_b, \mathcal{B}_b, \mathbb{P}_b)_{\mathfrak{v}},$$

along with the corresponding projections. (It may also be realised on the Lebesgue space = the unit interval with the Lebesgue measure).

With each fission point $\mathbf{v} \in {\{\mathbf{l}, \mathbf{r}\}}^*$, we associate a random element $\mathbf{g}_{\mathbf{v}} \in \mathfrak{A}$, where \mathfrak{A} is the affine group, defined below (3) for the disk model: if $z_{\mathbf{v}} = B_{\ell_{\mathbf{v}}}^{\mathbf{v}} o$ then $\mathbf{g}_{\mathbf{v}} = g_{z_{\mathbf{v}}}$ as defined by (4). The $\mathbf{g}_{\mathbf{v}}, \mathbf{v} \in {\{\mathbf{l}, \mathbf{r}\}}^*$, are i.i.d.

Furthermore, we take a starting point $z_0 \in \mathbb{D}$. We shall often write \mathbb{P}_{z_0} for the probability measure and \mathbb{E}_{z_0} for the corresponding expectation in order to remember the starting point.

Then hyperbolic branching Brownian motion assigns a random variable B_{τ} to every element τ of the Yule tree \mathcal{T} (vertex or interior element of some edge) as follows, using the notation of (1):

(A) If the path from α to $\mathfrak{v} \in {\{\mathfrak{l}, \mathfrak{r}\}}^*$ in our tree has the vertices $\alpha = \mathfrak{v}_0, \mathfrak{v}_1, \ldots, \mathfrak{v}_k = \mathfrak{v}$ (with $\mathfrak{v}_i^- = \mathfrak{v}_{j-1}$) then for $\tau = \mathfrak{v}_s \in [\mathfrak{v}', \mathfrak{v}]$,

$$\boldsymbol{B}_{\tau} = g_{z_0} \mathbf{g}_{\boldsymbol{\mathfrak{v}}_1} \cdots \mathbf{g}_{\boldsymbol{\mathfrak{v}}_{k-1}} B_s^{\boldsymbol{\mathfrak{v}}} \, .$$

This can also be described by the following recursive construction:

(B1) At the "ancestor" α ,

$$B_{lpha}=z_{0}$$
 .

(B2) If for a vertex $\mathfrak{v} \in {\mathfrak{l}, \mathfrak{r}}^*$, we already have $B_{\mathfrak{v}'} = z \in \mathbb{D}$, then on the edge $[\mathfrak{v}', \mathfrak{v}]$, we continue with

$$\boldsymbol{B}_{\boldsymbol{\mathfrak{v}}_s} = g_z B_s^{\boldsymbol{\mathfrak{v}}} \,, \quad 0 \le s \le \ell_{\boldsymbol{\mathfrak{v}}} \,,$$

where the random element $g_z \in \mathfrak{A}$ is given by (4), with z in the place of z_0 .

In particular, in our construction, hyperbolic BBM starting at z_0 is the image under g_{z_0} of hyperbolic BBM starting at 0.

Remarks 4.1. (a) In principle, we could choose any initial family of Möbius transformations such that for each $z_0 \in \mathbb{D}$ there is precisely one g_{z_0} mapping o to z_0 . For example (suggestion by Steve Lalley) one could take the map

$$z\mapsto \frac{z+z_0}{\bar{z}_0z+1}\,,$$

which is as in (3) with $a = 1/(1 - |z_0|^2)^{1/2}$ and $c = z_0/(1 - |z_0|^2)^{1/2}$. When $z_0 = 0$ it is the identity map, while otherwise it fixes the diameter segment of the unit disk through the origin and z_0 .

(b) However, the equivalence between the above two constructions (A) and (B1)+(B2) relies on the fact that the group \mathfrak{A} acts simply transitively, while this is not the case for the semigroup generated by the mappings of (a).

A feature of the construction (A) is that in this way, hyperbolic BBM is governed by a branching random walk on the group \mathfrak{A} . The underlying Galton-Watson tree is not random, but the binary tree $\{\mathfrak{l}, \mathfrak{r}\}^*$. That is, each member of the corresponding population fissions in precisely 2 children. The branching random walk starting at g_{z_0} is then $(g_{z_0}\mathbf{G}_{\mathfrak{v}})_{\mathfrak{v}\in\{\mathfrak{l},\mathfrak{r}\}^*}$, where

(17)
$$\mathbf{G}_{\mathfrak{v}} = \mathbf{g}_{\mathfrak{v}_1} \cdots \mathbf{g}_{\mathfrak{v}_k}$$

when the path from ϵ to \mathfrak{v} is $[\epsilon = \mathfrak{v}_1, \mathfrak{v}_2, \dots, \mathfrak{v}_k = \mathfrak{v}]$.

An infinite ray π in the Yule tree \mathcal{T} is a line isometric with $[0, \infty)$ which is spanned by a sequence of vertices $[\alpha, \epsilon, ...]$ where every vertex $\mathfrak{v} \in \pi$ has precisely one successor in π . Along π , the process $(\mathbf{B}_{\tau})_{\tau \in \pi}$ is a hyperbolic BM starting at z_0 whose element at time t is \mathbf{B}_{τ} with $\tau = v_s$, where $v \in \pi$ is a vertex and $|v_s| = t$ as defined below (1). If we have two distinct rays π and π' then their confluent $\pi \wedge \pi'$ is the furthest vertex from α shared by the two. Then the two hyperbolic Brownian motions along π and π' coincide up to $\pi \wedge \pi'$ and thereafter continue independently.

5. The maximal distance

We first consider an issue which has been intensively studied for Euclidean BBM since the pioneering work of KOLMOGOROV, PETROVSKI AND PISCOUNOV [21], namely, the behaviour of the maximal hyperbolic distance from the starting point of the population at time t. We only consider the basic result on the rate of escape. For details in the Euclidean case, see e.g. BRAMSON [7] and, in higher dimension, KIM, LUBETZKY AND ZEITOUNI [20].

Theorem 5.1. Set $Max_t = \max\{d(B_{\tau}, o) : \tau \in \mathcal{T}(t)\}$. Then

$$\frac{\textit{Max}_t}{t} \to r^* = \frac{1}{2} + \sqrt{2\lambda}$$

almost surely, as $t \to \infty$.

Proof. In the upper half plane model, the family $(-\log \Im B^{\mathbb{H}}_{\tau})_{\tau \in \mathcal{T}}$ is one-dimensional Euclidean branching BM with drift 1/2. It is well known that

(18)
$$\frac{1}{t} \max\{-\log \Im \boldsymbol{B}_{\tau} : \tau \in \mathcal{T}(t)\} \to r^* \text{ almost surely, as } t \to \infty$$

for r^* as given above, see e.g. [24] and Remark 5.2 below.

Furthermore, again in the upper half plane model, $d(z, o) \ge -\log \Im z$ by (13), which we have used already. This and (18) imply that

$$\liminf_{t \to \infty} \frac{\mathsf{Max}_t}{t} \ge r^* \quad \text{almost surely.}$$

For what follows, it is preferable to return to the disk model. Note that the particles' positions at a given time t, that is, the random variables

$$\boldsymbol{B}_{\tau}, \ \tau \in \mathcal{T}(t)$$

have the same distribution (but are not independent) as the ordinary hyperbolic random variable B_t . Let us assume, without loss of generality, that the starting point is o.

We now choose a constant $C > r^*$. If N(t) = n then write $\mathcal{T}(t) = \{\tau_1, \ldots, \tau_n\}$. Then

(19)

$$\mathbb{P}[\mathsf{Max}_{t} \geq Ct, \ N(t) = n] = \mathbb{P}\left(\bigcup_{j=1}^{n} [d(\boldsymbol{B}_{\tau_{j}}, o) \geq Ct, \ N(t) = n]\right)$$

$$\leq \sum_{j=1}^{n} \mathbb{P}[d(\boldsymbol{B}_{\tau_{j}}, o) \geq Ct, \ N(t) = n]$$

$$= n \mathbb{P}[d(B_{t}, o) \geq Ct] \mathbb{P}[N(t) = n],$$

since the position of B_{τ_j} does not depend on the value of N(t).

Therefore

$$\begin{split} \mathbb{P}[\mathsf{Max}_t \geq Ct] &= \sum_{n=1}^{\infty} \mathbb{P}[\mathsf{Max}_t \geq Ct \,, \, N(t) = n] \\ &= \sum_{n=1}^{\infty} n \mathbb{P}[d(B_t, o) \geq Ct] \, \mathbb{P}[N(t) = n] \\ &= \mathbb{P}[d(B_t, o) \geq Ct] \, \mathbb{E}\big(N(t)\big) = \mathbb{P}[d(B_t, o) \geq Ct] \, e^{\lambda t}. \end{split}$$

We need to estimate $\mathbb{P}[d(B_t, o) \ge Ct]$. This works similarly as in the last part of the proof of Proposition 3.2. We use once more (14). Up to a change of the leading constant, we can also use $\Psi(x) = x^{1/2}$ for $x \ge 1/2$ (instead of $x \ge 1$). This applies to the range which we

are now considering, namely $R \ge Ct > r^*t \ge t/2$. We perform the following estimate for (non-branching) hyperbolic BM in \mathbb{D} starting at o.

$$\mathbb{P}[d(B_t, o) \ge C t] = \int_{\{z:d(z,o) \ge Ct\}} p_t(o, z) \, d\mathsf{m}(z)$$

$$\le Const \int_{-\pi}^{\pi} \int_{Ct}^{\infty} \frac{\sqrt{1+R}}{t} \exp\left(-\frac{(R+\frac{t}{2})^2}{2t}\right) \frac{e^R - e^{-R}}{2} \, dR \, d\varphi$$

$$\le Const' \int_{Ct}^{\infty} \frac{\sqrt{1+R}}{t} \exp\left(-\frac{(R-\frac{t}{2})^2}{2t}\right) dR$$

substituting $s = (R - \frac{t}{2})/\sqrt{t}$

$$= \operatorname{Const'} \int_{(C-\frac{1}{2})\sqrt{t}}^{\infty} \sqrt{\frac{1}{t} + \frac{1}{2} + \frac{s}{\sqrt{t}}} \exp\left(-\frac{s^2}{2}\right) ds$$

since $\frac{s}{\sqrt{t}} \ge C - \frac{1}{2} > \sqrt{2\lambda}$

$$\leq \operatorname{Const}'' \int_{(C-\frac{1}{2})\sqrt{t}}^{\infty} \frac{s}{\sqrt{t}} \exp\left(-\frac{s^2}{2}\right) ds = \operatorname{Const}'' \frac{1}{\sqrt{t}} \exp\left(-\frac{(C-\frac{1}{2})^2 t}{2}\right),$$

(The constants are not assumed to be the same as in the proof of Proposition 3.2. Also, here Const'' depends on λ .)

Combining our computations, we get

$$\mathbb{P}[\mathsf{Max}_t \ge Ct] \le Const'' \frac{1}{\sqrt{t}} \exp\left(-\frac{\left((C - \frac{1}{2})^2 - 2\lambda\right)t}{2}\right)$$

Since $C > r^*$, we have $(C - \frac{1}{2})^2 - 2\lambda > 0$. Consequently, at integer times

$$\sum_{n=1} \mathbb{P}[\mathsf{Max}_n \geq Cn] < \infty \,,$$

and by the Borel-Cantelli Lemma,

$$\limsup_{n \to \infty} \frac{\mathsf{Max}_n}{n} \le C \quad \text{almost surely.}$$

This holds for every $C > r^*$, whence in view of the lower bound

$$\lim_{n \to \infty} \frac{\mathsf{Max}_n}{n} = r^* \quad \text{almost surely.}$$

We now need to fill in the "gaps" for real $t \in (n - 1, n)$, where n runs through the positive integers. For this purpose, we first define

$$\overline{\mathcal{M}} = \max\{d(\boldsymbol{B}_{\tau}, o) : \tau \in \mathcal{T}(\prec 1)\}$$

Claim. For any c > 0,

where \mathcal{M} is as in Proposition 3.2, that is, the maximal distance of ordinary hyperbolic BM from the starting point within the time interval [0, 1].

Proof of the Claim. Suppose that N(1) = k, and just within this proof, let $\mathcal{T}(1) = \{\tau_1, \ldots, \tau_k\}$. For each j, consider the geodesic path in \mathcal{T} from the root α to τ_j . Along this timeline, we see an ordinary hyperbolic BM starting at 0, runnig up to time 1. Let \mathcal{M}_j be the maximal distance from 0 of this BM. The random variables $\mathcal{M}_1, \ldots, \mathcal{M}_k$ are not independent, but they all have the same distribution as \mathcal{M} . Furthermore,

$$\overline{\mathcal{M}} = \max\{\mathcal{M}_1, \ldots, \mathcal{M}_k\}.$$

The proof of the Claim is now completed in the same way as in (19).

Note that the event $[|v| \notin \mathbb{N}$ for all $\mathfrak{v} \in {\mathfrak{l}, \mathfrak{r}}^*]$ has probability 1. We work on that event. Now let t > 0 and $n = \lfloor t \rfloor$. Each $\tau \in \mathcal{T}(n)$ is the root of a single subtree $\mathcal{T}_{\tau} = \mathcal{T}_{\tau}(\prec 1)$ of height 1 within \mathcal{T} , whose end-vertices belong to $\mathcal{T}(n+1)$. Let

$$\overline{\mathcal{M}}_{\tau} = \max\{d(\boldsymbol{B}_{\theta}, \boldsymbol{B}_{\tau}) : \theta \in \mathcal{T}_{\tau}\}$$

The random variables $\overline{\mathcal{M}}_{\tau}, \tau \in \mathcal{T}(n)$, are independent and (since the exponential distribution is memoryless) have the same distribution as the RV $\overline{\mathcal{M}}$ of the Claim. We clearly have

$$\operatorname{Max}_t \leq \operatorname{Max}_n + \max\{\overline{\mathcal{M}}_\tau : \tau \in \mathcal{T}(n)\}$$

The proof of the theorem will be complete if we can show that

(20)
$$\lim_{n \to \infty} \frac{1}{n} \max\{\overline{\mathcal{M}}_{\tau} : \tau \in \mathcal{T}(n)\} = 0 \quad \text{almost surely.}$$

Let c > 0. Then, once more in the same way as in (19), ad using the Claim,

$$\mathbb{P}\left[\max\{\overline{\mathcal{M}}_{\tau}: \tau \in \mathcal{T}(n)\} > cn\right] \leq \mathbb{P}\left[\overline{\mathcal{M}} > cn\right] e^{\lambda n}$$
$$\leq \mathbb{P}[\mathcal{M} > cn] e^{\lambda(n+1)}$$
$$\leq K \exp\left(-\frac{(cn - \frac{1}{2})^2 - 2\lambda(n+1)}{2}\right)$$

by Proposition 3.2. Summing over all n, the resulting series converges for every c > 0, and once more, the Borel-Cantelli Lemma yields (20).

Remark 5.2. Regarding the value of r^* , the literature deals with Euclidean BBM where the underlying BM is standard with drift 0. Also, usually only the case $\lambda = 1$ is considered for the fissioning rate (i.e., the parameter of the exponential distribution of the edge lengths of the Yule tree). The $\frac{1}{2}$ in our formula for r^* comes from the drift $\frac{1}{2}$ of our underlying Brownian motion. Then, in the drift-free case with $\lambda = 1$, on finds e.g. in [24] – see also BOVIER [6, Lemma 3.4] – that the maximum at time t of Euclidean BBM has rate $\sqrt{2}$. Now let \mathcal{T} be the Yule tree with parameter λ as in §2, and let $(\boldsymbol{\beta}_{\tau})_{\tau \in \mathcal{T}}$ be the associated standard Euclidean BBM (where the underlying BM is drift-free). Write $\lambda \mathcal{T}$ for the random tree where all the edge lengths are multiplied by λ . This is the Yule tree with fissioning rate 1. Then by the scale-invariance of BM, $(\frac{1}{\sqrt{\lambda}} \boldsymbol{\beta}_{\tau})_{\tau \in \mathcal{T}}$ is standard Euclidean BBM. From

here, one obtains the general formula. This is of course well-known and clear to the experts, but the hints may be useful for the novice.

6. The empirical distributions

We can interpret hyperbolic BBM as a Markov process on the space of *populations* in \mathbb{D} , where a population is a finitely supported measure of the form

$$\mathbf{M} = \sum_{j=1}^{n} \delta_{z_j}, \quad n \in \mathbb{N}, \ z_j \in \mathbb{D} \text{ not necessarily distinct.}$$

(Another interpretation is to see this as a multiset; see [18].)

Definition 6.1. Starting with one particle at position z_0 , the *occupation measure* of hyperbolic BBM at time $t \ge 0$ is the population

$$\mathbf{M}_t = \mathbf{M}_t^{z_0} = \sum_{\tau \in \mathcal{T}: |\tau| = t} \delta_{B_\tau}$$

and the associated *empirical distribution* is

$$\mu_t = \mu_t^{z_0} = \frac{1}{N(t)} \mathbf{M}_t \,.$$

Thus, μ_t is a finitely supported probability measure on the disk. Both \mathbf{M}_t and μ_t depend on the starting point z_0 , and in our construction, with g_{z_0} as in (4), $\mathbf{M}_t^{z_0} = g_{z_0}\mathbf{M}_t^o$ is the image of \mathbf{M}_t^o under the mapping $z \mapsto g_{z_0}z$. (That is, for a measure μ on $\overline{\mathbb{D}}$ and an isometry g of \mathbb{D} , we have $g\mu(K) = \mu(g^{-1}K)$.) We shall often omit the starting point in the notation.

Instead of starting BBM at a single point, we may start with an arbitrary initial population \mathbf{M}_0 . Then the population at time t is

(21)
$$\mathbf{M}_t = \sum_{j=1}^n \mathbf{M}_t^{z_j}, \quad \text{if} \quad \mathbf{M}_0 = \sum_{j=1}^n \delta_{z_j},$$

where the occupation measures $\mathbf{M}_{t}^{z_{i}}$ are independent even when some of the z_{i} coincide.

We start with an obvious consequence of the continuity of the distributions of the random variables B_{τ} .

Fact 6.2. For distinct $\tau, \tau' \in \mathcal{T}$, one has $\mathbb{P}[B_{\tau} = B_{\tau'}] = 0$.

In particular, for any fixed $t \ge 0$, with probability 1 the measure μ_t is equidistribution on N(t) distinct points.

Of course this does not imply that it never happens that μ_t is other than equidistributed. However, the following is also quite clear.

Lemma 6.3. Let π and π' be two distinct rays in the Yule tree. Then, with probability 1, there is a random $t_0 \geq 0$ such that $\mathbf{B}_{\tau} \neq \mathbf{B}_{\tau'}$ for all $\tau \in \pi$ and $\tau' \in \pi'$ with $|\tau|, |\tau'| > t_0$.

Proof. Let $v = \pi \wedge \pi'$. (Compare with the last lines of §4.) Conditionally upon the location (value) $\mathbf{B}_{v} = z_{0}$, beyond v the two Brownian motions along π and π' are independent replicas of hyperbolic BM starting at z_{0} . Each of them has an a.s. random limit ξ , resp. $\xi' \in \partial \mathbb{D}$. Both are distributed according to $\nu_{z_{0}}$; see (24) and the preceding lines. Continuity of $\nu_{z_{0}}$ yields that $\xi \neq \xi'$ almost surely. This implies the statement of the lemma.

Again, this does not necessarily mean that from some random time onwards, all μ_t are equidistributed. But if we restrict to all μ_n , $n \in \mathbb{N}$, then with probability 1 all of them *are* equidistributed.

For a stronger result than Lemma 6.3 concerning branching random walks on finitely generated groups, see HUTCHCROFT [17].

Question 6.4. Can one prove a result analogous to [17] that in the regime $\lambda \leq 1/8$, two independent copies of hyperbolic BBM meet at most finitely often? It appears to be true at integer times for any λ .

In the place of μ_t , it will also be useful to use the discrete random measure

$$\sigma_t = \sigma_t^{z_0} = \frac{1}{e^{\lambda t}} \mathbf{M}_t^{z_0} \,,$$

the image of σ_t^0 under $z \mapsto g_{z_0} z$. It is not a probability measure, but its expectation is a deterministic probability measure on \mathbb{D} , as the following Lemma shows.

Lemma 6.5. Let $K \subset \mathbb{D}$ be compact. Then the expected number of particles present within K at time t is

$$\mathbb{E}_{z_0}(\mathbf{M}_t(K)) = e^{\lambda t} \mathbb{P}_{z_0}[B_t \in K],$$

where B_t is hyperbolic Brownian motion started at z_0 . Thus,

$$\mathbb{E}_{z_0}\left(\int_0^\infty \sigma_t(K)\,dt\right)<\infty\,,$$

and therefore

$$\lim_{t \to \infty} \mu_t(K) = 0 \quad almost \ surely.$$

Proof. Once more, if N(t) = n then write $\mathcal{T}(t) = \{\tau_1, \ldots, \tau_n\}$. Then, similarly to (19),

$$\mathbb{E}_{z_0} \left(\mathbf{M}_t(K) \right) = \sum_{n=1}^{\infty} \mathbb{E}_{z_0} \left(\sum_{j=1}^n \mathbf{1}_K (\mathbf{B}_{\tau_j} \mid N(t) = n \right) \mathbb{P}[N(t) = n]$$
$$= \sum_{n=1}^{\infty} n \mathbb{P}_{z_0}[B_t \in K] \mathbb{P}[N(t) = n] = \mathbb{E} \left(N(t) \right) \mathbb{P}_{z_0}[B_t \in K].$$

This shows the first identity, and finiteness of $\mathbb{E}_{z_0}(\int_0^\infty \sigma_t(K) dt)$ follows from trancience of hyperbolic BM. It implies that the random variable $\int_0^\infty \sigma_t(K) dt$ is a.s. finite, whence

$$\lim_{t \to \infty} \mu_t(K) = \frac{1}{W} \lim_{t \to \infty} \sigma_t(K) = 0 \quad \text{almost surely,}$$

where W is the martingale limit of Proposition 2.2.

Thus, the mass of the empirical distributions μ_t "disappears at infinity" in the hyperbolic metric, resp. topology. (More precisely, the μ_t tend to 0 vaguely within the hyperbolic disk.)

We can express the first formula of Lemma 2.1 in terms of the transition operator (7) of (non-branching) hyperbolic BM: for a measurable set $K \subset \mathbb{D}$ and starting point z_0 ,

(22)
$$\mathbb{E}_{z_0}(\sigma_t(K)) = P_t(z, K).$$

Thus, for a measurable function $f : \mathbb{D} \to \mathbb{R}$

(23)
$$\mathbb{E}_{z_0}\left(\int_{\mathbb{D}} f \, d\mathbf{M}_t\right) = e^{\lambda t} P_t f(z_0) \quad \text{and} \quad \mathbb{E}_{z_0}\left(\int_{\mathbb{D}} f \, d\sigma_t\right) = P_t f(z_0) \,,$$

as long as the involved integral is well defined. A harmonic function is a C^2 -function h on \mathbb{D} which satisfies $\mathfrak{L}h \equiv 0$. It is well known that every harmonic function satisfies

$$P_t h = h$$
 for every $t > 0$.

Now let $\mathcal{F}_t = \mathcal{F}_t^{\text{BBM}}$ be the sigma-algebra generated by the information of hyperbolic BBM up to and including time t. (It projects naturally onto $\mathcal{F}_t^{\text{Yule}}$.)

Proposition 6.6. If h is a bounded or positive harmonic function, then the family of random variables

$$\int h \, d\sigma_t = e^{-\lambda t} \sum_{\tau \in \mathcal{T}: |\tau| = t} h(\boldsymbol{B}_{\tau}) \,, \quad t \ge 0$$

is a martingale with respect to the filtration $(\mathcal{F}_t)_{t\geq 0}$.

Proof. For any bounded or positive measurable function $f : \mathbb{D} \to \mathbb{C}$, consider its lift \tilde{f} to the space of populations:

$$\widetilde{f}(\mathbf{M}) = \int_{\mathbb{D}} f \, d\mathbf{M} = \sum_{z \in \mathbb{D}} f(z) \, \mathbf{M}(\{z\}) \,,$$

a finite sum. Let $(\widetilde{P}_t)_{t\geq 0}$ be the family of transition operators (i.e. the transition semigroup) of hyperbolic BBM, that is, of the Markov process $(\mathbf{M}_t)_{t\geq 0}$. By (21), for any population $\mathbf{M}_0 = \sum_{j=1}^n \delta_{z_j}$,

$$\widetilde{P}_{t}\widetilde{f}(\mathbf{M}_{0}) = \mathbb{E}_{\mathbf{M}_{0}}\left(\widetilde{f}(\mathbf{M}_{t})\right) = \mathbb{E}_{\mathbf{M}_{0}}\left(\sum_{z} f(z) \mathbf{M}_{t}(\{z\})\right)$$
$$= \mathbb{E}_{\mathbf{M}_{0}}\left(\sum_{z} f(z) \sum_{j=1}^{n} \mathbf{M}_{t}^{z_{j}}(z)\right) = \sum_{j=1}^{n} \mathbf{M}_{0}(z_{j}) \mathbb{E}_{z_{j}}\left(\sum_{z} f(z) \mathbf{M}_{t}^{z_{j}}(z)\right)$$
$$= \sum_{j=1}^{n} \mathbf{M}_{0}(z_{j}) \mathbb{E}_{z_{j}}\left(\int f d\mathbf{M}_{t}^{z_{j}}(z)\right) = \sum_{j=1}^{n} \mathbf{M}_{0}(z_{j}) e^{\lambda t} P_{t}f(z_{j})$$
$$= e^{\lambda t} \widetilde{P_{t}f}(\mathbf{M}_{0})$$

by (23). In particular, if h is a bounded or positive harmonic function on \mathbb{D} then $\widetilde{P}_t \widetilde{h} = e^{\lambda t} \widetilde{h}$ for every $t \ge 0$. This yields that $\int h \, d\sigma_t = e^{-\lambda t} \int h \, d\mathbf{M}_t$ is a martingale. \Box

Theorem 6.7. With probability 1, as $t \to \infty$, the measures σ_t converge weakly to a Borel measure σ_{∞} on $\partial \mathbb{D}$, and the probability measures μ_t converge weakly to a probability measure μ_{∞} on $\partial \mathbb{D}$. With W as in Proposition 2.2, we have

$$\sigma_{\infty} = W \mu_{\infty}, \quad and \quad \mathbb{E}_{z_0}(\sigma_{\infty}) = \nu_{z_0},$$

the measure on $\partial \mathbb{D}$ whose density with respect to the normalised Lebesgue measure on the circle is the Poisson kernel (8).

In particular, for every $\xi \in \partial \mathbb{D}$, we have $\mu_{\infty}(\{\xi\}) = 0$ almost surely.

Proof. We use a potential theoretic argument. Note that the harmonic functions for \mathfrak{L} are the same as the harmonic functions for the Euclidean Laplacian on \mathbb{D} . The corresponding Dirichlet problem is solvable: given a continuous function φ on $\partial \mathbb{D}$, there is a unique harmonic function h_{φ} on \mathbb{D} which provides a continuous extension of φ to the interior of the disk. Indeed, h_{φ} is the Poisson transform of φ ,

(24)
$$h_{\varphi}(z) = \int_{\partial \mathbb{D}} \Pi(z,\xi) \varphi(\xi) \, d\xi \, .$$

Now let f be any continuous function on the closed disk $\overline{\mathbb{D}}$. Considering σ_t as a measure on $\overline{\mathbb{D}}$, we have to show that $\int_{\mathbb{D}} f \, d\sigma_t$ converges almost surely. Let $\varphi = f|_{\partial \mathbb{D}}$ be the restriction of f to the unit circle, and let h_{φ} be the associated solution of the Dirichlet problem. This is a bounded harmonic function, so that

$$\int_{\mathbb{D}} h_{\varphi} \, d\sigma_t$$

converges almost surely by Proposition 6.6. On the other hand,

$$\lim_{|z| \to 1} f(z) - h_{\varphi}(z) = 0 \quad \text{uniformly in } z.$$

Given $\varepsilon > 0$, take $r \in (0, 1)$ such that $|f(z) - h_{\varphi}(z)| < \varepsilon$ for $|z| \ge r$. Let $K = \{z \in \mathbb{D} : |z| \le r\}$. By Proposition 2.2 and Lemma 6.5,

$$\int_{K} (f - h_{\varphi}) d\mu_{t} = \frac{e^{\lambda t}}{N(t)} \int_{K} (f - h_{\varphi}) d\sigma_{t} \to 0 \quad \text{almost surely,}$$

and since μ_t is a probability measure,

$$\left|\int_{\mathbb{D}\setminus K} (f-h_{\varphi}) \, d\mu_t\right| < \varepsilon \, .$$

This shows weak convergence. The identity $\sigma_{\infty} = W \mu_{\infty}$ is immediate from Proposition 2.2. Finally, if $f \in C(\overline{\mathbb{D}})$ then by (23) and dominated convergence

$$\mathbb{E}_{z_0}\left(\int_{\mathbb{D}} f \, d\sigma_{\infty}\right) = \lim_{t \to \infty} \mathbb{E}_{z_0}\left(\int_{\mathbb{D}} f \, d\sigma_t\right) = \lim_{t \to \infty} P_t f(z_0) = \lim_{t \to \infty} \mathbb{E}_{z_0}(f(B_t)) = \mathbb{E}_{z_0}(f(B_\infty))$$
$$= \int_{\partial \mathbb{D}} f \, d\nu_{z_0} \, .$$

In particular, for any $\xi \in \partial \mathbb{D}$, $\mathbb{E}_{z_0}(\sigma_{\infty}(\{\xi\})) = \nu_{z_0}(\{\xi\})$, so that $\sigma_{\infty}(\{\xi\}) = 0$ almost surely.

Note that the latter does not necessarily imply that almost surely, the limit measure carries no point mass.

The radial projection of \mathbb{D} to the boundary circle $\partial \mathbb{D}$ is the mapping

$$rad(re^{i\psi}) = e^{i\psi}$$
, if $0 < r < 1$, and $rad(0) = 1$

(The value rad(0) is of no specific importance and might be chosen arbitrarily.) We can consider the radial projection of hyperbolic Brownian motion, and the image of μ_t under the mapping rad.

Corollary 6.8. With probability 1, the probability measures

$$\mu_t^{rad} = \frac{1}{N(t)} \sum_{\tau \in \mathcal{T}: |\tau| = t} \delta_{rad(B_{\tau})}$$

on $\partial \mathbb{D}$ converge weakly to μ_{∞} , as $t \to \infty$.

Proof. Let $\varphi \in C(\mathbb{D})$, and let h_{φ} be its Poisson transform (24), providing the continuous extension of φ to $\overline{\mathbb{D}}$ which is harmonic on \mathbb{D} . To "hide" the discontinuity of $rad(\cdot)$ at 0, let

$$f(z) = \begin{cases} \left(h_{\varphi}(z) - \varphi(rad(z))\right) \min\{2|z|, 1\} & \text{if } z \in \mathbb{D}, \\ 0 & \text{if } z \in \partial \mathbb{D} \end{cases}$$

Then $f \in C(\overline{\mathbb{D}})$. We get that with probability 1,

$$\lim_{t \to \infty} \int_{\overline{\mathbb{D}}} f \, d\mu_t = 0$$

On the other hand,

$$\lim_{t \to \infty} \int_{\overline{\mathbb{D}}} h_{\varphi} \, d\mu_t = \int_{\partial \mathbb{D}} \varphi \, d\mu_{\infty}$$

This concludes the proof.

In analogy with Corollary 6.8, we have the following for the real parts of hyperbolic BBM in the upper half plane model. We omit the very similar proof.

Corollary 6.9. Let

$$\mu_t^{\Re} = \frac{1}{N(t)} \sum_{\tau \in \mathcal{T} : |\tau| = t} \delta_{\Re B_{\tau}^{\mathbb{H}}}.$$

18

Then, with probability one, μ_t^\Re converges weakly to $\mu_\infty^\mathbb{H}$, as $t\to\infty$.

Another feature of the "disappearance" of the population at infinity is the average rate of escape.

Definition 6.10. The *(empirical) distance distribution* of hyperbolic BBM at time $t \ge 0$ is the finitely supported random measure μ_t^d on \mathbb{R}_+ which is the image of μ_t under the mapping $\mathbb{D} \ni z \mapsto d(z, o)$, that is,

$$\mu_t^d = \frac{1}{N(t)} \sum_{\tau \in \mathcal{T}: |\tau| = t} \delta_{d(\boldsymbol{B}_{\tau}, o)}.$$

Theorem 6.11. With probability one, μ_t^d is asymptotically normal with mean t/2 and variance t. That is, its distribution function satisfies

$$\mu_t^d \left(-\infty, \frac{1}{2}t + x\sqrt{t}\right] \to \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-s^2/2} \, ds \quad \text{for every } x \in \mathbb{R},$$

as $t \to \infty$.

Before the proof, we make a detour to the upper half plane model and write $(\boldsymbol{B}_{\tau}^{\mathbb{H}})_{\tau \in \mathcal{T}}$ for the resulting model of hyperbolic BBM: it is the image of $(\boldsymbol{B}_{\tau})_{\tau \in \mathcal{T}}$ as costructed on \mathbb{D} under the mapping (9). Considering the logarithmic model, we see the following.

Fact 6.12. The vertical projection

$$oldsymbol{B}^{ ext{vert}}_{ au} = -\log \Im oldsymbol{B}^{\mathbb{H}}_{ au}\,, \quad au \in \mathcal{T},$$

is one-dimensional branching Brownian motion on \mathbb{R} ; the underlying Euclidean Brownian motion at time t has drift $\frac{1}{2}t$ and variance is t; see (11).

The reason for choosing the minus sign is that this Euclidean BBM has the same asymptotic drift and variance as the one proposed in Theorem 6.11, and we will use comparison of the two processes.

For the following, see NEY [25, Thm. 2], KAPLAN AND ASMUSSEN [19, Thm. 3] as well as BIGGINS [5].

Proposition 6.13. Let

$$\mu_t^{vert} = \frac{1}{N(t)} \sum_{\tau \in \mathcal{T}: |\tau| = t} \delta_{B_{\tau}^{vert}}.$$

With probability 1, as $t \to \infty$, the family of discrete probability distributions μ_t^{vert} on \mathbb{R} is asymptotically normal with mean t/2 and variance t.

[25] only has convergence in mean square, whence in probability. [19] has almost sure convergence when the drift of the base Brownian motion is 0 with obvious extension to non-zero drift; see also the last section of [5].

Proof of Theorem 6.7. We work with the upper half plane model and indicate this by the superscript \mathbb{H} . Let Ω_0 be the event on which $\mu_t^{\mathbb{H}}$ converges weakly and $\mu_{\infty}^{\mathbb{H}}(\mathfrak{i}\infty) = 0$. On Ω_0 , the latter measure assigns mass 0 to the boundary point $\mathfrak{i}\infty$, so that it lives on the lower boundary line \mathbb{R} . Let $f \in C^{\infty}(\mathbb{R})$ be uniformly continuous and set

$$f_t(x) = f\left(\frac{x - \frac{1}{2}t}{\sqrt{t}}\right).$$

By the Portmanteau Theorem, we need to show that on Ω_0 ,

$$\lim_{t \to \infty} \int_{\mathbb{R}} f_t(x) \, d\mu_t^d(x) = \int_{\mathbb{R}} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} f(x) \, dx \, .$$

Given $\varepsilon > 0$ there is a random bound $M < \infty$ such that

$$\mu_{\infty}^{\mathbb{H}}([-M, M]) \ge 1 - \varepsilon.$$

Consider the closed set $U_M = \{z \in \overline{\mathbb{H}} : |z| \leq M, \Im z \leq 1\} \subset \mathbb{H}$, where $\overline{\mathbb{H}} = \mathbb{H} \cup \mathbb{R} \cup \{\mathfrak{i} \infty\}$ is the geometric compactification of the hyperbolic upper half plane. Then

(25)
$$\limsup_{t \to \infty} \mu_t^{\mathbb{H}}(\overline{\mathbb{H}} \setminus U_M) \le \varepsilon$$

We decompose

$$\int_{\mathbb{R}} f_t(x) \, d\mu_t^d(x) = \int_{\overline{\mathbb{H}}} f_t\big(d_{\mathbb{H}}(z,o)\big) \, d\mu_t^{\mathbb{H}}(z)$$

into $\int_{U_M} + \int_{\overline{\mathbb{H}} \setminus U_M}$. Regarding the second part, we have by (25)

$$\limsup_{t \to \infty} \left| \int_{\overline{\mathbb{H}} \setminus U_M} f_t \big(d_{\mathbb{H}}(z, o) \big) \, d\mu_t^{\mathbb{H}}(z) \right| \le \varepsilon \, \|f\|_{\infty}$$

We now apply the same decomposition to

$$\int_{\mathbb{R}} f_t(x) \, d\mu_t^{\text{vert}}(x) = \int_{\overline{\mathbb{H}}} f_t\left(-\log\Im z\right) \, d\mu_t^{\mathbb{H}}(z)$$

and get

$$\limsup_{t \to \infty} \left| \int_{\mathbb{H} \setminus U_M} f_t \left(-\log \Im z \right) d\mu_t^{\mathbb{H}}(z) \right| \le \varepsilon \, \|f\|_{\infty}$$

We now consider the difference of the integrals over U_M , recalling that for $\Im z \leq 1$ one has $d_{\mathbb{H}}(\Im z, o) = -\log \Im z$:

$$\begin{split} \left| \int_{U_M} f_t \big(d_{\mathbb{H}}(z, o) \big) \, d\mu_t^{\mathbb{H}}(z) - \int_{U_M} f_t \big(-\log \Im z \big) \, d\mu_t^{\mathbb{H}}(z) \right| \\ & \leq \int_{U_M} \left| f_t \big(d_{\mathbb{H}}(z, o) \big) - f_t \big(d_{\mathbb{H}}(\Im z, o) \big) \right| \, d\mu_t^{\mathbb{H}}(z) \, . \end{split}$$

By (13), we have $|d_{\mathbb{H}}(z, o) - d_{\mathbb{H}}(\Im z, o)| \le \log(1 + M^2)$ on U_M . Therefore

$$\left|\frac{d_{\mathbb{H}}(z,o) - \frac{1}{2}t}{\sqrt{t}} - \frac{d_{\mathbb{H}}(\Im z,o) - \frac{1}{2}t}{\sqrt{t}}\right| \le \frac{\log(1+M^2)}{\sqrt{t}},$$

and by uniform continuity of f,

$$\lim_{t \to \infty} \left| f_t \big(d_{\mathbb{H}}(z, o) \big) - f_t \big(d_{\mathbb{H}}(\Im z, o) \big) \right| = 0 \quad \text{uniformly on } U_M.$$

We infer that on Ω_0

$$\lim_{t \to \infty} \int_{\mathbb{R}} f_t(x) \, d\mu_t^d(x) = \lim_{t \to \infty} \int_{\mathbb{R}} f_t(x) \, d\mu_t^{\text{vert}}(x) \, ,$$

and Proposition 6.13 yields the proposed asymptotic behaviour.

The average displacement of hyperbolic BBM is

(26)
$$\int_{\mathbb{D}} d(z,o) \, d\mu_t(z) = \frac{1}{N(t)} \sum_{\tau \in \mathcal{T}: |\tau|=t} d(\boldsymbol{B}_{\tau},o)$$

Theorem 6.14. With probability 1, the average displacement of hyperbolic BBM has rate of escape

$$\lim_{t \to \infty} \frac{1}{t} \int_{\mathbb{D}} d(z, o) \, d\mu_t(z) = \frac{1}{2} \, .$$

Proof. In the sample space Ω of hyperbolic BBM, let Ω_1 be the set on which μ_t^d is asymptotically normal as in Theorem 6.11, and

$$\frac{1}{\mathsf{Max}_t} \to r^*$$

as in Theorem 5.1, where $\mathsf{Max}_t = \max\{d(B_{\tau}, o) : \tau \in \mathcal{T}, |\tau| = t\}$. We can consider this as providing a random environment for the family of probability distributions μ_t^d , and consider one quenched case, i.e., a fixed $\omega \in \Omega_1$. We can think of the associated deterministic family μ_t^d , t > 0, as the distributions of a family of random variables X_t . By asymptotic normality,

$$\lim_{t \to \infty} \frac{X_t}{t} = \frac{1}{2} \quad \text{in probability.}$$

On the other hand,

$$\frac{|X_t|}{t} \le \frac{\mathsf{Max}_t}{t} \,,$$

which is bounded. We can apply the Dominated Convergence Theorem, and denoting by \mathbb{E}_{ω} the quenched expectation, we get

$$\lim_{t \to \infty} \frac{\mathbb{E}_{\omega}(X_t)}{t} = \frac{1}{2}.$$

Now

$$\mathbb{E}_{\omega}(X_t) = \int_{\mathbb{R}} x \, d\mu_t^d(x) = \int_{\mathbb{D}} d(z, o) \, d\mu_t(z) \, d\mu_t$$

and this proves the claim.

21

7. Some properties of the random limit distributions

Here, we shall collect a few properties of the measures $\mu_{\infty}^{z_0}$ plus related open questions. Before that, we start with a general fact with a simple proof, comunicated to the author by V. A. Kaimanovich during the work on the paper [18].

Proposition 7.1. Let \mathcal{X} be a separable compact space and $\mathcal{P}(\mathcal{X})$ be the space of Borel probability measures on \mathcal{X} . With the weak*topology, it is again separable and compact. Now let μ and μ' be independent random probability measures on \mathcal{X} , with distributions $\boldsymbol{\nu}$ and $\boldsymbol{\nu}'$, respectively. That is, the latter are probability measures on $\mathcal{P}(\mathcal{X})$, and the distribution of (μ, μ') on \mathcal{X}^2 is $\boldsymbol{\nu} \otimes \boldsymbol{\nu}'$.

Suppose that the expectation of μ (barycentre of ν), that is, the deterministic probability measure in \mathcal{X} given by

$$\overline{\mu}(K) = \int_{\mathcal{P}(\mathcal{X})} \mu(K) \, d\boldsymbol{\nu}(\mu) \quad (K \subset \mathcal{X} \text{ compact})$$

on \mathcal{X} is purely non-atomic. Then, almost surely, μ' and μ share no atoms.

("Almost surely" refers to the probability measure $\boldsymbol{\nu} \otimes \boldsymbol{\nu}'$.)

Proof. Step 1. Since $\overline{\mu}$ is non-atomic, for any $x \in \mathcal{X}$

 $\mu(x) = 0$ for $\boldsymbol{\nu}$ - almost every $\mu \in \mathcal{P}(\mathcal{X})$.

Step 2. Let $\mu' \in \mathcal{P}(\mathcal{X})$ be fixed (deterministic). Then ν – almost every $\mu \in \mathcal{P}(\mathcal{X})$ has no common atoms with μ' , that is,

$$\int_{\mathcal{P}(\mathcal{X})} \mu(x)\mu'(x) \, d\boldsymbol{\nu}(\mu) = 0 \quad \text{for all } x \in \mathcal{X} \, .$$

Proof. Let $x_i, i \in I$ be the finitely or countably many atoms of μ' (if any). Then the statement of Step 1 holds for each x_i .

Conclusion. Step 2 holds for every probability measure ν' on $\mathcal{P}(\mathcal{X})$. Therefore

$$\iint_{\mathcal{P}(\mathcal{X})\times\mathcal{P}(\mathcal{X})}\mu(x)\mu'(x)\,d\boldsymbol{\nu}(\mu)\,d\boldsymbol{\nu}'(\mu')=0\quad\text{for all }x\in\mathcal{X}\,,$$

which is the proposed result.

For any vertex $\mathfrak{u} \in {\mathfrak{l}, \mathfrak{r}}^*$, consider the subtree $\mathcal{T}_{\mathfrak{u}}$ as defined in §2. Its distribution is the same as the one of \mathcal{T} . Recall that u^- has the role of the ancestor α in $\mathcal{T}_{\mathfrak{u}}$. For the associated martingale according to Proposition 2.2, we write

(27)
$$W_{\mathfrak{u}} = \lim_{t \to \infty} N_{\mathfrak{u}}(t) e^{-\lambda t}$$

for its almost surely existing and nondegenerate limit. If $B_{\mathfrak{u}'} = z_{\mathfrak{u}'} = z$ then $(g_z^{-1}B_\tau)_{\tau\in\mathcal{T}_{\mathfrak{u}}}$ is hyperbolic BBM starting at o, whose pieces along the edges come from the same construction on our probability space as for the process starting with the ancestor α . We have $g_z = \mathbf{G}_{\mathfrak{u}'}$ as defined by (17). (Simple transitivity of the group \mathfrak{A} is useful here, since in this way we can avoid the need to handle uncountably many different versions.) In particular, all the empirical distributions of $(g_z^{-1} \mathbf{B}_\tau)_{\tau \in \mathcal{T}_u}$ converge almost surely, and by continuity of the action of g_z , we have almost surely all the limits

(28)
$$\mu_{\mathfrak{u},\infty}^{z} = \lim_{t \to \infty} \mu_{\mathfrak{u},t}^{z}$$
, where $z = z_{\mathfrak{u}'} = B_{\mathfrak{u}'}$ and $\mu_{\mathfrak{u},t}^{z} = \frac{1}{N_{\mathfrak{u}}(t)} \sum_{\tau \in \mathcal{T}_{\mathfrak{u}} : |\tau| = t + |u^{-}|} B_{\tau}$

for for every z and for each of the countably many vertices of \mathcal{T} . Along with $\mu_{\mathfrak{u},t}^z$ and $\mu_{\mathfrak{u},\infty}^z$, we also have

$$\sigma_{\mathfrak{u},t}^{z} = N_{\mathfrak{u}}(t)e^{-\lambda t}\,\mu_{\mathfrak{u},t}^{z} \quad \text{and} \quad \sigma_{\mathfrak{u},\infty}^{z} = W_{\mathfrak{u}}\,\mu_{\mathfrak{u},\infty}^{z}$$

with $W_{\mathfrak{u}}$ as in (27).

Lemma 7.2. Let $\mathfrak{u}, \mathfrak{v} \in {\{\mathfrak{l}, \mathfrak{r}\}}^*$ be such that none of the two is an ancestor (predecessor) of the other. Consider hyperbolic BBM indexed by the time trees $\mathcal{T}_{\mathfrak{u}}$ and $\mathcal{T}_{\mathfrak{v}}$, respectively, and starting at o. Then with probability 1, the limit distributions $\mu_{\mathfrak{u},\infty}^o$ and $\mu_{\mathfrak{v},\infty}^o$ share no atoms.

Proof. The two random measures are independent. Furthermore,

$$\mu^o_{\mathfrak{u},\infty} = \frac{1}{W_{\mathfrak{u}}} \sigma^o_{\mathfrak{u},\infty}.$$

Now we know from Theorem 6.7 that the expectation of the random measure $\sigma_{u,\infty}^o$ is ν_o , the limit distribution on \mathbb{D} of hyperbolic Brownian motion starting at o. This is normalised Lebesgue measure, whence it has no atoms. Since W_u is almost surely finite and positive, also the expectation of $\mu_{u,\infty}^o$ has no atoms. The lemma now follows from Proposition 7.1. \Box

Now we can deduce a first interesting property.

Theorem 7.3. With probability 1, the support of the random limit distribution $\mu_{\infty}^{o} = \mu_{\epsilon,\infty}^{o}$ is infinite.

Proof. For any measure μ on $\partial \mathbb{D}$, we shall write $\operatorname{atoms}(\mu)$ for the set of atoms of μ , and $\operatorname{supp}(\mu)$ for its support.

Let $\mathfrak{u} \in {\mathfrak{l}, \mathfrak{r}}^*$ and consider hyperbolic BBM according to our construction of §4 with time tree $\mathcal{T}_{\mathfrak{u}}$ and starting point $\mathbf{B}_{\mathfrak{u}'} = o$. The two children of u are $u\mathfrak{l}$ and $u\mathfrak{r}$. For each of the two independent subtrees $\mathcal{T}_{\mathfrak{u}\mathfrak{l}}$ and $\mathcal{T}_{\mathfrak{u}\mathfrak{r}}$, the vertex u has the role that the ancestor α has in \mathcal{T} . Write $z_1 = \mathbf{B}_{\mathfrak{u}}$, so that $g_{z_1} = \mathbf{g}_{\mathfrak{u}}$, the random element of \mathfrak{A} as described in §4. Denote by $\mathbf{M}_{\mathfrak{u}\mathfrak{l},t}^{z_1}$ and $\mathbf{M}_{\mathfrak{u}\mathfrak{r},t}^{z_1}$ the occupation measures at time t (that is, distance t from u in $\mathcal{T}_{\mathfrak{l}}$ and $\mathcal{T}_{\mathfrak{r}}$), respectively. Then

$$\mathbf{M}_{\mathfrak{u},t}^{o} = \mathbf{M}_{\mathfrak{ul},t-\ell_{\mathfrak{u}}}^{z_{1}} + \mathbf{M}_{\mathfrak{ur},t-\ell_{\mathfrak{u}}}^{z_{1}}.$$

Dividing by $N_{\epsilon}(t)$ and letting $t \to \infty$,

$$\mu_{\mathfrak{u},\infty}^{o} = \mathbf{c}_{\mathfrak{u}\mathfrak{l}}\,\mu_{\mathfrak{u}\mathfrak{l},\infty}^{z_{1}} + \mathbf{c}_{\mathfrak{u}\mathfrak{r}}\,\mu_{\mathfrak{u}\mathfrak{r},\infty}^{z_{1}}, \quad \text{where} \quad \mathbf{c}_{\mathfrak{u}\mathfrak{l}} = \frac{W_{\mathfrak{u}\mathfrak{l}}}{W_{\mathfrak{u}}}\,e^{-\lambda\ell_{\mathfrak{u}}} \text{ and } \mathbf{c}_{\mathfrak{u}\mathfrak{r}} = \frac{W_{\mathfrak{u}\mathfrak{r}}}{W_{\mathfrak{u}}}\,e^{-\lambda\ell_{\mathfrak{u}}},$$

a non-trivial convex combination of two probability measures. We can also rewrite this as

$$\mathbf{g}_{\mathfrak{u}}^{-1}\mu_{\mathfrak{u},\infty}^{o}=\mathbf{c}_{\mathfrak{u}\mathfrak{l}}\,\mu_{\mathfrak{u}\mathfrak{l},\infty}^{o}+\mathbf{c}_{\mathfrak{u}\mathfrak{r}}\,\mu_{\mathfrak{u}\mathfrak{r},\infty}^{o}$$
 .

From this identity and Lemma 7.2 we infer that

 $\mathsf{atoms}(\mu^o_{\mathfrak{u},\infty}) = \mathbf{g}_\mathfrak{u} \mathsf{atoms}(\mu^o_{\mathfrak{u}\mathfrak{l},\infty}) \mathrel{\dot{\cup}} \mathbf{g}_\mathfrak{u} \mathsf{atoms}(\mu^o_{\mathfrak{u}\mathfrak{l},\infty})$

is almost surely a disjoint union.

Recursively, we get the following convex combination for each n:

(29)
$$\mu^{o}_{\epsilon,\infty} = \sum_{\mathfrak{v} \in \{\mathfrak{l},\mathfrak{r}\}^{n}} \mathbf{c}^{(n)}_{\mathfrak{v}} \mathbf{G}_{\mathfrak{v}'} \mu^{o}_{\mathfrak{v},\infty}$$

with positive random constants $\mathbf{c}_{\boldsymbol{v}}^{(n)}$ and $\mathbf{G}_{v'}$ as defined by (17), and

$$\operatorname{atoms}(\mu^o_{\mathfrak{u},\infty}) = \bigcup_{\mathfrak{v} \in \{\mathfrak{l},\mathfrak{r}\}^n} \mathbf{G}_{\mathfrak{v}'} \operatorname{atoms}(\mu^o_{\mathfrak{v},\infty}) \quad \text{almost surely.}$$

If $\operatorname{supp}(\mu_{\mathfrak{u},\infty}^o)$ is finite then the support of the measure coincides with the set of its atoms, and there must be \mathfrak{v} such that $\mu_{\mathfrak{v},\infty}^o$ has no atoms. But along with $\mu_{\epsilon,\infty}^o$, by (29) all $\mu_{\mathfrak{v},\infty}^o$ have finite support consisting only of atoms, a contradiction.

Definition 7.4. The *limit set* of hyperbolic BBM is the random set Λ of accumulation points of $(B_{\tau})_{\tau \in \mathcal{T}}$ on the unit circle $\partial \mathbb{D}$.

LALLEY AND SELLKE [23] have proved the following significant result.

Theorem 7.5. [23] In the unit circle with the arclength measure, there is the following dichotomy.

(i) For $\lambda \leq 1/8$, with probability 1, Λ is a Cantor set (totally disconnected and perfect), and its Hausdorff dimension is $(1 - \sqrt{1 - 8\lambda})/2$. Furthermore, Λ is contained in a proper sub-arc of \mathbb{D} .

(ii) For $\lambda > 1/8$, with probability 1, $\Lambda = \partial \mathbb{D}$.

It is clear that

$$\operatorname{supp}(\mu_{\infty}^{o}) \subseteq \Lambda$$
.

At this point several interesting questions arise to which at present the author does not know the anser.

Questions 7.6. (a) Is $supp(\mu_{\infty}^{o}) \subset \Lambda$ properly?

(b) What is the Hausdorff dimension of μ_{∞}^{o} (i.e. the smallest Hausdorff dimension of a set with full μ_{∞}^{o} -measure)? Is it strictly smaller than the Hausdorff dimension of Λ ?

(c) Is μ_{∞}^{o} purely non-atomic? Is it absolutely continuous with respect to Lebesgue (arc) measure in the supercritical regime $\lambda > 1/8$?

All answers may well depend on the regime $\lambda \leq 1/8$, resp. > 1/8.

References

- Athreya, K. B. and Ney, P. E.: Branching Processes. Grundlehren der mathematischen Wissenschaften 196. Springer, New York, 1972.
- [2] Benjamini, I. and Müller, S.: On the trace of branching random walks. Groups Geom. Dyn. 6 (2012) 231-247.
- [3] Benjamini, I. and Peres, Y.: Markov chains indexed by trees. Ann. Probab. 22 (1994) 219–243.
- [4] Biggins, J. D.: The central limit theorem for the supercritical branching random walk, and related results. Stochastic Process. Appl. 34 (1990) 255–274.
- [5] Biggins, J. D.: Uniform convergence of martingales in the branching random walk. Ann. Probab. 20 (1992) 137–151.
- [6] Bovier, Anton(D-BONN-AM).: From spin glasses to branching Brownian motion and back? In Random Walks, Random Fields, and Disordered Systems, pp. 1–64. Lecture Notes in Math. 2144, Springer, Cham, 2015.
- Bramson, M.: Convergence of solutions of the Kolmogorov equation to travelling waves. Mem. Amer. Math. Soc. 44 (1983) no. 285.
- [8] Candellero, E., Gilch, L. and Miller, S.: Branching random walks on free products of groups. Proc. London Math. Soc. 104 (2012) 1085–1120.
- [9] Candellero, E. and Hutchcroft, T.: On the boundary at infinity for branching random walk. Electronic Commun. Probab. 28 (2023) 1–12.
- [10] Candellero, E. and Roberts, M. I.: The number of ends of critical branching random walks. ALEA Lat. Am. J. Probab. Math. Stat. 12 (2015) 55–67.
- [11] Davies, E. B. and Mandouvalos, N.: Heat kernel bounds on hyperbolic space and Kleinian groups. Proc. London Math. Soc. (3) 57 (1988) 182–208.
- [12] Dussaule, M., Wang, L. and Yang, W.: Branching random walks on relatively hyperbolic groups. Preprint (2022), arXiv:2211.07213.
- [13] Gantert, N. and Müller, S.: The critical branching Markov chain is transient. Markov Process. Related Fields 12 (2006) 805–814.
- [14] Gilch, L. A. and Müller, S.: Ends of branching random walks on planar hyperbolic Cayley graphs. In Groups, Graphs and Random Walks, London Math. Soc. Lecture Note Ser. 436, pp. 205–214, Cambridge Univ. Press, Cambridge, 2017.
- [15] Harris, T.: The Theory of Branching Processes. Grundlehren der mathematischen Wissenschaften 119. Springer, Berlin 1963.
- [16] Hueter, I. and Lalley, S. P.: Anisotropic branching random walks on homogeneous trees. Probab. Theory Rel. Fields 116 (2000) 57–88.
- [17] Hutchcroft, T.: Non-intersection of transient branching random walks. Probab. Theory Rel. Fields 178 (2020) 1–23.
- [18] Kaimanovich, V. A. and Woess, W.: Limit distributions of branching Markov chains. Ann. Inst. H. Poincaré Probab. Stat. 59 (2023) 1951–1983.
- [19] Kaplan, N. and Asmussen, S.: Branching random walks II. Stochastic Process. Appl. 4 (1976) 15–31.
- [20] Kim, Y. H., Lubetzky, E. and Zeitouni, O.:The maximum of branching Brownian motion in \mathbb{R}^d . Ann. Appl. Probab. **33** (2023) 1315–1368.
- [21] Kolmogorov, A. N., Petrovski, I. and Piscounov, N.: Étude de l'équation de la diffusion avec croissance de la quantité de matiére et son application à un problème biologique. Moscow Univ. Math. Bull. 1 (1937) 1–25.
- [22] Liggett, T. M.: Branching random walks and contact processes on homogeneous trees. Probab. Theory Related Fields 106 (1996) 495–519.
- [23] Lalley, S. P. and Sellke, T.: Hyperbolic branching Brownian motion. Probab. Theory Rel. Fields. 108 (1997) 171–192.
- [24] McKean, H. P.: Application of Brownian motion to the equation of Kolmogorov-Petrovskii-Piskunov. Comm. Pure Appl. Math. 28 (1975)323–331.

- [25] Ney, P. M.: The convergence of a random distribution function associated with a branching process. J. Math. Anal. Appl. 12 (1965), 316–327.
- [26] Sidoravicius, V., Wang, L. and Xiang, K.: Limit set of branching random walks on hyperbolic groups. Comm. Pure Appl. Math. 76 (2023) 2765–2803.
- [27] Stam, A. J.: On a conjecture by Harris. Z. Wahrscheinlichkeitstheorie verw. Gebiete 5 (1966) 202–206.
- [28] Uchiyama, K.: Spatial growth of a branching process of particles living in \mathbb{R}^d . Ann. Probab. 10 (1982)896–918.
- [29] Yule, G.U.: A mathematical theory of evolution, based on the conclusions of Dr. J. C. Willis. Philos. Trans. Roy. Soc. London Ser. B 213 (1924) 21–87.

Institut für Diskrete Mathematik, Technische Universität Graz, Steyrergasse 30, A-8010 Graz, Austria

Email address: woess@tugraz.at