

A characterization of the existence of zeros for operators with Lipschitzian derivative and closed range

BIAGIO RICCERI

ABSTRACT. Let H be a real Hilbert space and $\Phi : H \rightarrow H$ be a C^1 operator with Lipschitzian derivative and closed range. We prove that $\Phi^{-1}(0) \neq \emptyset$ if and only if, for each $\epsilon > 0$, there exist a convex set $X \subset H$ and a convex function $\psi : X \rightarrow \mathbf{R}$ such that $\sup_{x \in X}(\|x\|^2 + \psi(x)) - \inf_{x \in X}(\|x\|^2 + \psi(x)) < \epsilon$ and $0 \in \overline{\text{conv}}(\Phi(X))$.

2020 Mathematics Subject Classification. Primary 47J05.

Key words. Nonlinear operator, Lipschitzian derivative, minimax theorem.

The aim of this very short note is to establish the following result:

THEOREM 1. - *Let H be a real Hilbert space, let $\Omega \subseteq H$ be an open set and let $\Phi : \Omega \rightarrow H$ be a C^1 operator with Lipschitzian derivative.*

Then, the following assertions are equivalent:

- (a) $\inf_{x \in \Omega} \|\Phi(x)\| = 0$;
- (b) *for every $\epsilon > 0$, there exist a convex set $X \subset \Omega$ and a convex function $\psi : X \rightarrow \mathbf{R}$ such that*

$$\sup_{x \in X}(\|x\|^2 + \psi(x)) - \inf_{x \in X}(\|x\|^2 + \psi(x)) < \epsilon$$

and

$$0 \in \overline{\text{conv}}(\Phi(X)) .$$

More precisely, the key result of this note is Theorem 2 below. Theorem 1 then follows as a by-product of it.

THEOREM 2. - *Let H be a real Hilbert space, let $\Omega \subseteq H$ be an open set and let $\Phi : \Omega \rightarrow H$ be a C^1 operator such that Φ' is Lipschitzian, with Lipschitz constant L . Moreover, assume that*

$$\delta := \inf_{x \in \Omega} \|\Phi(x)\| > 0 .$$

Then, for each convex set $X \subset \Omega$ such that

$$\sup_{x \in X}(\|x\|^2 + \psi(x)) - \inf_{x \in X}(\|x\|^2 + \psi(x)) < \frac{2\delta}{L}$$

for some convex function $\psi : X \rightarrow \mathbf{R}$, one has

$$0 \notin \overline{\text{conv}}(\Phi(X)) .$$

PROOF. Fix any convex set $X \subset \Omega$ and any convex function $\psi : X \rightarrow \mathbf{R}$ satisfying

$$\sup_{x \in X}(\|x\|^2 + \psi(x)) - \inf_{x \in X}(\|x\|^2 + \psi(x)) < \frac{2\delta}{L} . \tag{1}$$

Set

$$Y = \{x \in H : \|x\| \leq 1\}$$

and consider the functions $\varphi : X \rightarrow \mathbf{R}$ and $f, g : X \times Y \rightarrow \mathbf{R}$ defined by

$$\varphi(x) = \frac{L}{2}(\|x\|^2 + \psi(x)) ,$$

$$f(x, y) = \langle \Phi(x), y \rangle$$

and

$$g(x, y) = f(x, y) + \varphi(x)$$

for all $x \in X, y \in Y$. We claim that

$$\inf_X \sup_Y f - \sup_Y \inf_X f \leq \sup_X \varphi - \inf_X \varphi . \quad (2)$$

Arguing by contradiction, assume that

$$\inf_X \sup_Y f - \sup_Y \inf_X f > \sup_X \varphi - \inf_X \varphi .$$

We then would have

$$\sup_Y \inf_X g \leq \sup_Y \inf_X f + \sup_X \varphi < \inf_X \sup_Y f + \inf_X \varphi \leq \inf_X \sup_Y g . \quad (3)$$

For each $y \in Y$, the function $f(\cdot, y)$ is C^1 and one has

$$f'_x(x, y)(u) = \langle \Phi'(x)(u), y \rangle$$

for all $x \in \Omega, u \in H$. Also, for each $v, w \in \Omega$, we have

$$\begin{aligned} \|f'_x(v, y) - f'_x(w, y)\| &= \sup_{u \in Y} |\langle \Phi'(v)(u) - \Phi'(w)(u), y \rangle| \leq \sup_{u \in Y} \|\Phi'(v)(u) - \Phi'(w)(u)\| \\ &= \|\Phi'(v) - \Phi'(w)\|_{\mathcal{L}(H)} \leq L\|v - w\| . \end{aligned} \quad (4)$$

In view of (4), for each $y \in Y$, the function $g(\cdot, y)$ is convex ([2], Corollary 42.7). But then, thanks to a standard minimax theorem ([1]), we would have

$$\sup_Y \inf_X g = \inf_X \sup_Y g$$

contradicting (3). So, (2) does hold. Notice that

$$\inf_X \sup_Y f = \inf_{x \in X} \|\Phi(x)\| .$$

Therefore, from (2) we infer that

$$\frac{L}{2} \left(\inf_{x \in X} (\|x\|^2 + \psi(x)) - \sup_{x \in X} (\|x\|^2 + \psi(x)) \right) + \inf_{x \in X} \|\Phi(x)\| \leq \sup_{y \in Y} \inf_{x \in X} \langle \Phi(x), y \rangle$$

and hence, in view of (1),

$$0 < \sup_{y \in Y} \inf_{x \in X} \langle \Phi(x), y \rangle .$$

Fix $\gamma > 0$ and $\tilde{y} \in Y$ so that

$$\inf_{x \in X} \langle \Phi(x), \tilde{y} \rangle \geq \gamma .$$

Thus, if we set

$$C = \{u \in H : \langle u, \tilde{y} \rangle \geq \gamma\} ,$$

we have

$$\overline{\text{conv}}(\Phi(X)) \subset C$$

while $0 \notin C$ and the proof is complete. \triangle

Proof of Theorem 1. The implication (a) \rightarrow (b) is immediate. So, assume that (b) holds. We have to prove (a). Arguing by contradiction, suppose that $\inf_{x \in \Omega} \|\Phi(x)\| > 0$. By assumption, there are a convex set $X \subset \Omega$ and a convex function $\psi : X \rightarrow \mathbf{R}$ such that

$$\sup_{x \in X} (\|x\|^2 + \psi(x)) - \inf_{x \in X} (\|x\|^2 + \psi(x)) < 2 \frac{\inf_{x \in \Omega} \|\Phi(x)\|}{L}$$

and

$$0 \in \overline{\text{conv}}(\Phi(X)) ,$$

against Theorem 2. \triangle

REMARK 1. - Notice that, in general, Theorem 1 does not hold if Φ is only C^1 . To see this, consider a C^1 function $h : \mathbf{R} \rightarrow \mathbf{R}$ with the following property: there exist two sequences in $]0, +\infty[$ $\{\alpha_n\}, \{\beta_n\}$ such that $\lim_{n \rightarrow \infty} \alpha_n = +\infty$, $\lim_{n \rightarrow \infty} (\beta_n^2 - \alpha_n^2) = 0$, $\alpha_n < \beta_n$, $h(\alpha_n) = 0$, $h(\beta_n) = -\pi$ for all $n \in \mathbf{N}$. Then, consider the function $\Phi : \mathbf{R}^2 \rightarrow \mathbf{R}^2$ defined by

$$\Phi(x, y) = (\sin(h(x)), \cos(h(x)))$$

for all $(x, y) \in \mathbf{R}^2$. So, Φ is C^1 and $\inf_{(x, y) \in \mathbf{R}^2} \|\Phi(x, y)\| = 1$. However, for each $\epsilon > 0$, if n is large enough, we have $\beta_n^2 - \alpha_n^2 < \epsilon$ and $0 \in \text{conv}(\Phi([\alpha_n, \beta_n] \times \{0\}))$.

REMARK 2. - We are not aware of known results close to Theorem 1 so that a proper comparison can be made.

Acknowledgements: The author has been supported by PRIN 2022BCFHN2 "Advanced theoretical aspects in PDEs and their applications", by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) and by the Università degli Studi di Catania, PIACERI 2020-2022, Linea di intervento 2, Progetto MAFANE.

References

- [1] M. SION *On general minimax theorems*, Pacific J. Math., **8** (1958), 171-176.
- [2] E. ZEIDLER, *Nonlinear functional analysis and its applications*, vol. III, Springer-Verlag, 1985.

Department of Mathematics and Informatics
University of Catania
Viale A. Doria 6
95125 Catania, Italy
e-mail address: ricceri@dmf.unict.it