

Measurement-induced phase transitions in systems with diffusive dynamics

Hyunsoo Ha,¹ Akshat Pandey,² Sarang Gopalakrishnan,³ and David A. Huse¹

¹*Department of Physics, Princeton University, Princeton, NJ 08544, USA*

²*Department of Physics, Stanford University, Stanford, CA 94305, USA*

³*Department of Electrical and Computer Engineering,
Princeton University, Princeton, NJ 08544, USA*

(Dated: May 16, 2024)

The competition between scrambling and projective measurements can lead to measurement-induced entanglement phase transitions (MIPT). In this work, we show that the universality class of the MIPT is drastically altered when the system is coupled to a diffusing conserved density. Specifically, we consider a 1+1d random Clifford circuit locally monitored by classically diffusing particles (“measurers”). The resulting diffusive correlations in the measurement density are a relevant perturbation to the usual space-time random MIPT critical point, producing a new universality class for this phase transition. We find “Griffiths-like” effects due to rare space-time regions where, e.g., the diffusive measurers have a low or high density, but these are considerably weaker than the Griffiths effects that occur with quenched randomness that produce rare spatial regions with infinite lifetime.

Measurement-induced phase transitions (MIPTs) are a recently discovered class of phase transitions in quantum dynamical systems subject to repeated measurements. MIPTs are transitions in the entanglement structure of typical individual *quantum trajectories*—i.e., in the state of the system conditional on a set of measurement outcomes [1–3]. They separate a “mixed” or “volume-law” phase in which the dynamics generate volume-law entanglement that is able to preserve some quantum information about the initial state, from a “pure” or “area-law” phase in which measurements rapidly extract all quantum information and prevent the buildup of entanglement [4, 5]. The MIPT in spatially local one-dimensional random quantum circuits, with local unitary gates and single-site projective measurements, has been the best-studied example [6–12]; however, MIPTs have also been explored in other geometries [13–17]. Experimental studies of MIPTs remain challenging because studying individual trajectories requires procedures (such as post-selection or classical simulations) for which the resource requirements grow exponentially with system size; nevertheless, signatures of MIPTs have been seen in recent experiments with trapped ions and superconducting qubits [18–21], and recent theoretical proposals indicate how MIPTs may be realized in larger-scale experiments [22–24].

The critical properties of MIPTs depend on the spatial dimension and range of the gates [15, 25–27]. In addition, typical trajectories are inherently random in space and time—since the measurement outcomes are random—and the critical properties are sensitive to the correlations in this randomness. The “standard” MIPT occurs in one-dimensional circuits where the unitary gates and measurement locations are chosen at random with no space-time correlations, and the measurement outcomes are set by the Born rule. A number of circuit-averaged observables at this standard MIPT seem to be governed by a

nonunitary conformal field theory [9], which, in the limit of infinite on-site Hilbert space dimension, can be identified as percolation. When the measurement locations or the gate parameters have strong enough long-range correlations, the universality class changes. For example, if there is quenched spatial randomness in the measurement probabilities, the transition is instead governed by an infinite-randomness fixed point with infinitely anisotropic spacetime scaling [28, 29].

In the standard MIPT, the dynamics is not subject to any conservation laws. Even in the absence of measurements, conservation laws constrain entanglement growth in quantum circuits [30]. In this paper we explore how the presence of a diffusing conserved charge (conserved also by the measurements) affects the MIPT in 1+1 dimensions. Generically, the measurement rate at the transition will depend linearly on small changes in the overall charge density [31]. When this is true, as we argue below, the coupling to the diffusing charge is relevant at the critical point, and thus changes the universality class of this phase transition [32]. Note that, in addition to the MIPT (which is an *entanglement* transition), circuits with a conserved charge also exhibit a “charge-sharpening” phase transition [23, 33, 34]. The MIPT takes place in the charge-sharp phase, in which measurements have collapsed the charge distribution, so an individual quantum trajectory has a well-defined spacetime charge density profile at large length scales.

The MIPT can be studied efficiently for large systems composed of Clifford gates and Pauli measurements [3, 5, 8–10, 35]. Unfortunately, if one restricts such a random Clifford circuit to have a conserved $U(1)$ charge (e.g. \hat{Z}_i summed over all qubits i) this eliminates the volume-law entangled phase and thus also the phase transition we wish to study. However, in more general models this phase transition occurs within the charge-sharp phase, in which charge fluctuations are essentially classical [33].

Thus we introduce a new type of model in order to efficiently numerically study this transition. In this model, a diffusing conserved *classical* charge is separate from the quantum Clifford random circuit, the circuit alone has no conservation laws, and the measurement rate at a given space-time point is set by the (classical) charge at that point. This allows us to simulate large systems and thus investigate the resulting new universality class of the MIPT. In the absence of the conserved charge, the universality class of the transition does differ in some details between Clifford and non-Clifford (e.g., Haar-random) systems, although in many respects they are qualitatively similar [8, 10]. With the conserved charge, we can again expect differences between non-Clifford and Clifford universality classes; here, we study the latter because it is much more accessible numerically by using the new type of model that we introduce. Our main results are to locate and characterize the critical point of the MIPT in this model. We also provide numerical evidence and analytical arguments to show that diffusive correlations affect the properties of the phases on either side of the MIPT.

Our model has classical particles diffusing via the symmetric simple exclusion process (SSEP) on the same one-dimensional lattice that the qubits occupy. The particle density is equal to one particle per two qubits. These classical particles are the “measurers”: at the time of measurements, each qubit that is at a site occupied by a particle is measured with probability p ; see Fig. 1. This imparts diffusive correlations in space-time to the locations of the measurements. In all data presented here, the measurers have diffusivity $D = 1$, and we use periodic boundary conditions except where otherwise stated.

Generalized Harris argument— We begin by giving a generalized Harris argument [36, 37] for the relevance of diffusive correlations in the measurement rate. The standard MIPT (with uncorrelated measurements) has correlation length exponent ν and dynamical exponent $z = 1$, so for p near the critical point p_c the correlation region has space-time volume $\sim \xi^{(d+1)}(p) \sim |p - p_c|^{-(d+1)\nu}$ for $d + 1$ dimensions. Space-time correlations in the measurements will be relevant if they cause the measurement rate averaged over a space-time correlation volume to vary (between different correlation volumes of the same system) by an amount δp that vanishes more slowly than $|p - p_c| \sim \xi^{-1/\nu}$ as $p \rightarrow p_c$.

If diffusive measurers are added perturbatively to the standard uncorrelated model, the number of measurers in a correlation volume does not change significantly within one correlation time; the measurer number is strongly correlated along the time direction. The variation in the number of measurers is proportional to the square root of the spatial volume, resulting in a variation of measurer density within a correlation volume proportional to $\xi^{-d/2}$. Therefore, $\delta p \sim \xi^{-d/2}(p)$, which is relevant when $\nu < 2/d$, as in the usual Harris argument [38]. In partic-




FIG. 1. Random Clifford circuit monitored by diffusing classical particles. (a) Illustration of classical SSEP rules governing measurer dynamics, where hopping to nearest-neighbor sites occurs as a continuous-time Poisson process with a specified rate and is forbidden if the adjacent site is already occupied. (b) An example of measurer trajectories. (c) A circuit with measurer locations as in (b). Two-qubit random Clifford gates are applied in a brickwork manner. At each time between layers of gates, each measurer makes a \hat{Z} projective measurement of the local qubit with probability p .

ular, for the 1+1-dimensional case that we consider here, the uncorrelated critical point has $\nu \cong 1.3 < 2/d = 2$. Therefore, diffusive correlations are indeed a relevant perturbation for the uncorrelated MIPT. Note that this relevance criterion does not change as long as the measurers move slower than ballistically (so that the number of measurers within a correlation volume does not change significantly in one correlation time in the $p \rightarrow p_c$ limit).

Critical point: numerical results— The generalized Harris argument establishes that diffusive correlations drive the system away from the spacetime-uncorrelated critical point. To study the new critical point that *does* occur, we turn to numerical simulations of these Clifford circuits. We begin by estimating the critical measurement rate p_c based on the peak of the antipodal mutual information (AMI) for subsystems of size $N/8$. The AMI vanishes as $N \rightarrow \infty$ in both the volume-law and area-law phases and is expected to exhibit a peak at p_c under fairly general conditions [3]. As shown in Fig. 2(a), the peak in the AMI for accessible sizes suggests $p_c \approx 0.33$. Interpreted conservatively, our data on the AMI constrains p_c to lie in the interval $(0.31, 0.35)$: for p within this interval, the AMI grows with L for all accessible sizes, so any value in this range could be the critical point. The estimated p_c for this model is approximately twice that in the standard random-Clifford MIPT. This factor of roughly two is plausible because the number of measurers is $N_p = N/2$, so the average measurement rate over all spacetime is $p/2$. In [39], we attempt to extract the correlation-length exponent ν at




FIG. 2. Entanglement and purification dynamics at and near the critical point. (a) Antipodal mutual information in the long-time steady state shows that the critical point p_c is located near $p = 0.33$. (b) Purification time τ_P . The median over samples is presented as a representative of a typical sample. The gray dotted line is the relaxation time scale for the diffusive measurers with $z = 2$. The inset shows the apparent activated dynamical scaling at $p = 0.33$, with the horizontal axis re-scaled to $N^{0.44}$.

this critical point by collapsing the AMI as well as a different observable, the tripartite mutual information. This gives us a rough estimate of $\nu = 1.6(5)$. The relatively poor quality of the collapse indicates that despite the large system sizes we are considering, finite-size effects are still severe.

To explore dynamical scaling at the critical point, we turn to the purification dynamics of an initially maximally mixed state for measurement rates p around this critical window. At the critical point, we expect that the timescale for purifying a system of size N scales as $\tau_P(N) \sim N^z$, where z is the dynamical critical exponent. For the standard MIPT, $z = 1$, and for the case of quenched randomness, $z = \infty$ [28]. In the diffusive case, our numerical results [Fig. 2(b)] clearly indicate that $z \geq 2$. Assuming that $N = 256$ is larger than any relevant microscopic scale, the slope $d(\log \tau_P)/d(\log N)$ at $p = 0.35$ lower bounds the value of z . This lower bound is close to the diffusive value $z = 2$, which, as we will discuss, is the largest value that is theoretically well-motivated. If we instead focus on the apparent p_c —given by the peak in the AMI—we find a slope that is clearly drifting to larger values with increasing N . This would imply $z > 3.5$ and would be most consistent with a formally infinite z , with power-law dynamical scaling replaced by some form of activated scaling, as at the infinite-randomness critical point. Indeed, as shown in the inset to Fig. 2(b), an activated scaling ansatz with scaling $\log \tau_P \sim N^\psi$ with $\psi \cong 0.44$ is consistent with the purification data.

Thus our numerical results are consistent with at least two possibilities for the critical point. One possible scenario, suggested by a straightforward analysis of the numerics, gives $p_c \approx 0.33$ and $z > 3.5$. Although consistent with the numerics, this scenario seems implausible, by

the following logic: Imagine running a renormalization-group scheme that gives an exponent $z > 2$. After many steps of this scheme, the system will have $O(1)$ remaining degrees of freedom per $\ell \times \ell^z$ patch for some large ℓ . The remaining disorder at this scale is spatially uncorrelated, so if the renormalized description of the circuit resembles the original microscopic description, the flow beyond this scale will be governed by the standard MIPT and z will decrease. We emphasize that this argument makes some nontrivial assumptions about the nature of the flow, and does not rule out the $z > 2$ scenario. A simpler possibility, however, is that the AMI is showing strong finite-size effects and the true critical point lies near the upper end of the allowed range (0.31, 0.35) and has $z = 2$, so the apparent values of $z > 2$ are actually in the volume-law phase. We present some tentative numerical support for this possibility in [39].

Rare region effects— We now turn to the phases on either side of this new MIPT. As in the infinite-randomness case, we expect that some properties of these phases are dominated by rare spacetime regions that are locally “in the other phase”. To influence dynamics on a timescale t , such a rare configuration must persist for time t . In the standard MIPT, the measurement probabilities are drawn independently at each time, so this probability decays exponentially in time. However, in the diffusive case, a rare region of size $\ell \gtrsim \sqrt{Dt}$ typically persists for time t . The probability “cost” of such a region is stretched-exponential in t , so rare-region effects can dominate late-time dynamics of certain quantities whose typical behavior is exponential decay. The quantities that show these effects are different in the two phases:

In the volume-law phase, we find that regions with anomalously high density of measurers behave as bottlenecks that suppress *entanglement growth* due to the resulting high density of measurements. We prepare a random pure product initial state, evolve with open boundary conditions until time t , bipartition the system with a single cut at position x near the midpoint ($x = N/2$) and obtain the entanglement entropy that we label $S_{N/2}(t)$. We define the total probability $P_{S_{N/2}}^<(t)$ at time t of having $S_{N/2}(t) < \bar{S}_{N/2}(t)/2$: the probability of the entanglement entropy for a single instance being suppressed by at least a factor of two from its mean over many instances at time t .

In the usual MIPT, the most probable way to produce such a large suppression is due to a space-time path of time duration $\sim t$ that has substantially suppressed entanglement growth along the entire path. Therefore one expects $-\log P_{S_{N/2}}^< \sim t$. In our diffusive circuit, the tail is due to the rare regions of spatial width $O(\sqrt{Dt})$ such that they can survive until time t . Therefore, we expect $-\log P_{S_{N/2}}^< \sim \sqrt{Dt}$. With time-independent quenched randomness, the rare region is a bottleneck of the entan-




FIG. 3. Rare region effects in the volume-law phase in models with (a) diffusing measurers, (b) space-time uncorrelated measurements, and (c) immobile measurers having quenched randomness. Note, the horizontal axis is different in each case. The presented measurement rates are $p \cong p_c/2$, thus well into the volume-law phase. The system size is $N = 1024$, and cuts are sampled within a distance of 64 sites of the center of the chain.

glement growth with a rate exponentially decaying with its length. In this case the most probable way to suppress the entanglement is due to a nearby bottleneck that is of spatial width $O(\log t)$, leading to the Griffiths power law: $-\log P_{S_{N/2}}^< \sim \log t$ [28]. We numerically verified the results are consistent with expectations for all three models (Fig. 3). This probability also oscillates in time as it decays because entanglement in Clifford circuits is integer-valued [39].

In the area-law phase, rare regions with low measurement density slow down *purification* from the maximally mixed initial state. We focus on two quantities: the mean total entropy $S(t)$ and the tail distribution of the purification time τ_P , denoted as $P_{\tau_P}^>(t) \equiv P(\tau_P \geq t)$. These two quantities behave similarly at very late times when $S(t) < 1$, as both diagnose the last bit of entropy to be purified.

In the intermediate time regime of $1 \ll Dt < O(N^2)$, the entropy remains in rare regions with a lower density of measurers that were already present at time zero. For these to remain until time t they must be of spatial length $\ell \gtrsim \sqrt{Dt}$. The probability of having a rare region decays exponentially in its length ℓ , and once the probability is rare, there are approximately N/ℓ independent locations where such rare regions may occur. Therefore, we expect the following intermediate-time scaling:

$$\log\left(\frac{S(t)}{N/\sqrt{Dt}}\right) \sim -\sqrt{Dt}, \quad \log\left(\frac{P_{\tau_P}^>(t)}{N/\sqrt{Dt}}\right) \sim -\sqrt{Dt}, \quad (1)$$

which are clearly observed in Fig. 4(a,b). The typical purification time diverges with system size as $\sim (\log N)^2/D$: The largest rare region length is typically $O(\log N)$, and the typical purification time is determined by the diffusive lifetime ($z = 2$) of this rare region.




FIG. 4. Rare region effects in the area-law phase. (a,b) Numerical demonstrations of Eqs. (1) at intermediate times, where the longest-living rare region determines the decay of the total entropy and the tail of the purification time distribution. (c,d) At later times $S(t)$ and $P_{\tau_P}^>(t)$ decay exponentially with t/N . Color scale indicates N .

Beyond the diffusive time scale $t \gg O(N^2/D)$, almost all the rare samples in the area-law phase that have not fully purified have entropy one bit. This logical operator has survived in spite of the strong measurements due to a sequence of rare spacetime regions. The most probable way for a logical to survive for such long times is for there to be a rare region of size of order N : the density of measurers is low over some fraction of order one half of the full system. Such a pattern occurs with a probability that is exponentially small in N . This rare pattern would normally disappear after a time $\sim N^2/D$, so for it to persist even longer to time $t \gg N^2/D$ means it did not decay away of order Dt/N^2 times. Thus the probability of the system not purifying to these late times is

$$\log P_{\tau_P}^>(t) \cong \log S(t) \sim (-N) \times \frac{Dt}{N^2} = -\frac{Dt}{N}. \quad (2)$$

This is consistent with Fig. 4(c,d), where for large t/N both $S(t)$ and $P_{\tau_P}^>(t)$ decay exponentially with t/N (orange dotted lines are parallel).

Conclusion— We have shown that diffusive correlations due to a conservation law change the universality class of the MIPT, as well as the long-time asymptotics of the phases on either side of it—specifically, entanglement dynamics in the volume-law phase and purification dynamics in the area-law phase. We have provided analytic arguments to show that the universality class of the transition must change, and to estimate the “Griffiths-like” effects from diffusive correlations in the phases. To isolate the effect of diffusive correlations while still reaching large system sizes, we numerically explored a model in which the diffusive correlations are put in by hand. However, our analytic arguments do not rely on the specifics

of this model, and we expect that our main conclusions are general, although critical exponents might differ between Clifford and generic circuits.

If $z > 2$, the correlation length exponent has a weaker lower bound, as the single correlation volume lasts for ξ^{z-2} correlation times of the measurers on scale ξ . Hence, the rms variation in the measurement rate between such space-time correlation volumes is $\delta p \sim \xi^{(2-d-z)/2}$, matching to the $\sim \xi^{d/2}$ for $z \leq 2$. This gives a generalized Chayes inequality $\nu \geq 2/(z+d-2) = 2/(z-1)$ for 1+1 dimension and $z \geq 2$. Therefore, if the system is behaving on some scale with apparent $z > 2$, the apparent ν we extract may only obey the generalized inequality and “explain” our apparent $\nu < 2$ (although it has a large uncertainty).

The critical point we have explored is one of a large family of new critical points that arise naturally in the dynamics of quantum and/or classical information. The sharpening transition is another example in this vein; in one dimension disorder is irrelevant at this transition, but in higher dimensions (or for symmetry groups other than $U(1)$) it might be a relevant perturbation. The dynamics of out-of-time-order correlators in classical diffusive spin chains is another closely related example [37]. It was also recently shown that the coupling between quantum and classical circuits leads to nontrivial entanglement growth [40]. These critical points involve an information-theoretic quantity, such as entanglement, charge fluctuations, or an out-of-time-order correlator, coupled nonreciprocally to one or more hydrodynamic modes. Developing a framework for describing these new phase transitions is an important topic for future work.

H.H. thanks Grace Sommers and Su-un Lee for advice on numerics, Zack Weinstein, Ahana Chakraborty, Vir Bulchandani, and Rhine Samajdar for fruitful discussions. A.P. is grateful to Aditya Cowsik and Nicholas O’Dea for discussions. S.G. and D.A.H. thank Jed Pixley, Drew Potter, and Romain Vasseur for helpful discussions and collaborations on related topics. D.A.H. was supported in part by NSF QLCI grant OMA-2120757. This material is based upon work supported by the U.S. Department of Energy, Office of Science, National Quantum Information Science Research Centers, Co-design Center for Quantum Advantage (C2QA) under contract number DE-SC0012704 (theoretical analysis performed by S.G.). Simulations presented in this work were performed on computational resources managed and supported by Princeton Research Computing.

[1] Yaodong Li, Xiao Chen, and Matthew P. A. Fisher. Quantum zeno effect and the many-body entanglement transition. *Phys. Rev. B*, 98:205136, Nov 2018. doi:10.1103/PhysRevB.98.205136. URL <https://link.aps.org/doi/10.1103/PhysRevB.98.205136>.

- [2] Brian Skinner, Jonathan Ruhman, and Adam Nahum. Measurement-induced phase transitions in the dynamics of entanglement. *Phys. Rev. X*, 9:031009, Jul 2019. doi:10.1103/PhysRevX.9.031009. URL <https://link.aps.org/doi/10.1103/PhysRevX.9.031009>.
- [3] Yaodong Li, Xiao Chen, and Matthew P. A. Fisher. Measurement-driven entanglement transition in hybrid quantum circuits. *Phys. Rev. B*, 100:134306, Oct 2019. doi:10.1103/PhysRevB.100.134306. URL <https://link.aps.org/doi/10.1103/PhysRevB.100.134306>.
- [4] Soonwon Choi, Yimu Bao, Xiao-Liang Qi, and Ehud Altman. Quantum error correction in scrambling dynamics and measurement-induced phase transition. *Phys. Rev. Lett.*, 125:030505, Jul 2020. doi:10.1103/PhysRevLett.125.030505. URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.030505>.
- [5] Michael J. Gullans and David A. Huse. Dynamical purification phase transition induced by quantum measurements. *Phys. Rev. X*, 10:041020, Oct 2020. doi:10.1103/PhysRevX.10.041020. URL <https://link.aps.org/doi/10.1103/PhysRevX.10.041020>.
- [6] Romain Vasseur, Andrew C. Potter, Yi-Zhuang You, and Andreas W. W. Ludwig. Entanglement transitions from holographic random tensor networks. *Phys. Rev. B*, 100:134203, Oct 2019. doi:10.1103/PhysRevB.100.134203. URL <https://link.aps.org/doi/10.1103/PhysRevB.100.134203>.
- [7] Yimu Bao, Soonwon Choi, and Ehud Altman. Theory of the phase transition in random unitary circuits with measurements. *Phys. Rev. B*, 101:104301, Mar 2020. doi:10.1103/PhysRevB.101.104301. URL <https://link.aps.org/doi/10.1103/PhysRevB.101.104301>.
- [8] Aidan Zabalo, Michael J. Gullans, Justin H. Wilson, Sarang Gopalakrishnan, David A. Huse, and J. H. Pixley. Critical properties of the measurement-induced transition in random quantum circuits. *Phys. Rev. B*, 101:060301, Feb 2020. doi:10.1103/PhysRevB.101.060301. URL <https://link.aps.org/doi/10.1103/PhysRevB.101.060301>.
- [9] Yaodong Li, Xiao Chen, Andreas W. W. Ludwig, and Matthew P. A. Fisher. Conformal invariance and quantum nonlocality in critical hybrid circuits. *Phys. Rev. B*, 104:104305, Sep 2021. doi:10.1103/PhysRevB.104.104305. URL <https://link.aps.org/doi/10.1103/PhysRevB.104.104305>.
- [10] A. Zabalo, M. J. Gullans, J. H. Wilson, R. Vasseur, A. W. W. Ludwig, S. Gopalakrishnan, David A. Huse, and J. H. Pixley. Operator scaling dimensions and multifractality at measurement-induced transitions. *Phys. Rev. Lett.*, 128:050602, Feb 2022. doi:10.1103/PhysRevLett.128.050602. URL <https://link.aps.org/doi/10.1103/PhysRevLett.128.050602>.
- [11] Andrew C. Potter and Romain Vasseur. *Entanglement Dynamics in Hybrid Quantum Circuits*, pages 211–249. Springer International Publishing, Cham, 2022. ISBN 978-3-031-03998-0. doi:10.1007/978-3-031-03998-0_9. URL https://doi.org/10.1007/978-3-031-03998-0_9.
- [12] Matthew P.A. Fisher, Vedika Khemani, Adam Nahum, and Sagar Vijay. Random quantum circuits. *Annual Review of Condensed Matter Physics*, 14(Volume 14, 2023):335–379, 2023. ISSN 1947-5462. doi:https://doi.org/10.1146/annurev-conmatphys-031720-030658. URL <https://www.>

- annualreviews.org/content/journals/10.1146/annurev-conmatphys-031720-030658.
- [13] Javier Lopez-Piqueres, Brayden Ware, and Romain Vasseur. Mean-field entanglement transitions in random tree tensor networks. *Phys. Rev. B*, 102:064202, Aug 2020. doi:10.1103/PhysRevB.102.064202. URL <https://link.aps.org/doi/10.1103/PhysRevB.102.064202>.
- [14] Sagar Vijay. Measurement-driven phase transition within a volume-law entangled phase. *arXiv preprint arXiv:2005.03052*, 2020. URL <https://arxiv.org/abs/2005.03052>.
- [15] Oliver Lunt, Marcin Sznyszewski, and Arijeet Pal. Measurement-induced criticality and entanglement clusters: A study of one-dimensional and two-dimensional clifford circuits. *Phys. Rev. B*, 104:155111, Oct 2021. doi:10.1103/PhysRevB.104.155111. URL <https://link.aps.org/doi/10.1103/PhysRevB.104.155111>.
- [16] Adam Nahum, Shitadhi Roy, Brian Skinner, and Jonathan Ruhman. Measurement and entanglement phase transitions in all-to-all quantum circuits, on quantum trees, and in landau-ginsburg theory. *PRX Quantum*, 2:010352, Mar 2021. doi:10.1103/PRXQuantum.2.010352. URL <https://link.aps.org/doi/10.1103/PRXQuantum.2.010352>.
- [17] Xiaozhou Feng, Brian Skinner, and Adam Nahum. Measurement-induced phase transitions on dynamical quantum trees. *PRX Quantum*, 4:030333, Sep 2023. doi:10.1103/PRXQuantum.4.030333. URL <https://link.aps.org/doi/10.1103/PRXQuantum.4.030333>.
- [18] Crystal Noel, Pradeep Niroula, Daiwei Zhu, Andrew Risinger, Laird Egan, Debopriyo Biswas, Marko Cetina, Alexey V Gorshkov, Michael J Gullans, David A Huse, et al. Measurement-induced quantum phases realized in a trapped-ion quantum computer. *Nature Physics*, 18(7):760–764, 2022. URL <https://www.nature.com/articles/s41567-022-01619-7>.
- [19] Jin Ming Koh, Shi-Ning Sun, Mario Motta, and Austin J Minnich. Measurement-induced entanglement phase transition on a superconducting quantum processor with mid-circuit readout. *Nature Physics*, 19(9):1314–1319, 2023. URL <https://www.nature.com/articles/s41567-023-02076-6>.
- [20] Google Quantum AI and Collaborators. Measurement-induced entanglement and teleportation on a noisy quantum processor. *Nature*, 2023. doi:10.1038/s41586-023-06505-7. URL <https://doi.org/10.1038/s41586-023-06505-7>.
- [21] Hirsh Kamakari, Jiace Sun, Yaodong Li, Jonathan J. Thio, Tanvi P. Gujarati, Matthew P. A. Fisher, Mario Motta, and Austin J. Minnich. Experimental demonstration of scalable cross-entropy benchmarking to detect measurement-induced phase transitions on a superconducting quantum processor. 2024. URL <https://arxiv.org/abs/2403.00938>.
- [22] Yaodong Li, Yijian Zou, Paolo Glorioso, Ehud Altman, and Matthew P. A. Fisher. Cross entropy benchmark for measurement-induced phase transitions. *Phys. Rev. Lett.*, 130:220404, Jun 2023. doi:10.1103/PhysRevLett.130.220404. URL <https://link.aps.org/doi/10.1103/PhysRevLett.130.220404>.
- [23] Fergus Barratt, Utkarsh Agrawal, Andrew C. Potter, Sarang Gopalakrishnan, and Romain Vasseur. Transitions in the learnability of global charges from local measurements. *Phys. Rev. Lett.*, 129:200602, Nov 2022. doi:10.1103/PhysRevLett.129.200602. URL <https://link.aps.org/doi/10.1103/PhysRevLett.129.200602>.
- [24] Samuel J Garratt and Ehud Altman. Probing post-measurement entanglement without post-selection. *arXiv preprint arXiv:2305.20092*, 2023. URL <https://arxiv.org/abs/2305.20092>.
- [25] Takaaki Minato, Koudai Sugimoto, Tomotaka Kuwahara, and Keiji Saito. Fate of measurement-induced phase transition in long-range interactions. *Phys. Rev. Lett.*, 128:010603, Jan 2022. doi:10.1103/PhysRevLett.128.010603. URL <https://link.aps.org/doi/10.1103/PhysRevLett.128.010603>.
- [26] Maxwell Block, Yimu Bao, Soonwon Choi, Ehud Altman, and Norman Y. Yao. Measurement-induced transition in long-range interacting quantum circuits. *Phys. Rev. Lett.*, 128:010604, Jan 2022. doi:10.1103/PhysRevLett.128.010604. URL <https://link.aps.org/doi/10.1103/PhysRevLett.128.010604>.
- [27] T. Müller, S. Diehl, and M. Buchhold. Measurement-induced dark state phase transitions in long-ranged fermion systems. *Phys. Rev. Lett.*, 128:010605, Jan 2022. doi:10.1103/PhysRevLett.128.010605. URL <https://link.aps.org/doi/10.1103/PhysRevLett.128.010605>.
- [28] Aidan Zabalo, Justin H. Wilson, Michael J. Gullans, Romain Vasseur, Sarang Gopalakrishnan, David A. Huse, and J. H. Pixley. Infinite-randomness criticality in monitored quantum dynamics with static disorder. *Phys. Rev. B*, 107:L220204, Jun 2023. doi:10.1103/PhysRevB.107.L220204. URL <https://link.aps.org/doi/10.1103/PhysRevB.107.L220204>.
- [29] Gal Shkolnik, Aidan Zabalo, Romain Vasseur, David A. Huse, J. H. Pixley, and Snir Gazit. Measurement induced criticality in quasiperiodic modulated random hybrid circuits. *Phys. Rev. B*, 108:184204, Nov 2023. doi:10.1103/PhysRevB.108.184204. URL <https://link.aps.org/doi/10.1103/PhysRevB.108.184204>.
- [30] Tibor Rakovszky, Frank Pollmann, and C. W. von Keyserlingk. Sub-ballistic growth of rényi entropies due to diffusion. *Phys. Rev. Lett.*, 122:250602, Jun 2019. doi:10.1103/PhysRevLett.122.250602. URL <https://link.aps.org/doi/10.1103/PhysRevLett.122.250602>.
- [31] Ahana Chakraborty, Kun Chen, Aidan Zabalo, Justin H. Wilson, and J. H. Pixley. Charge and entanglement criticality in a u(1)-symmetric hybrid circuit of qubits. *arXiv:2307.13038*, 2023. URL <https://arxiv.org/abs/2307.13038>.
- [32] Note1. At special points where the critical measurement rate does not depend linearly on the density, the coupling to diffusion may be irrelevant. We will not consider this case in the present work.
- [33] Utkarsh Agrawal, Aidan Zabalo, Kun Chen, Justin H. Wilson, Andrew C. Potter, J. H. Pixley, Sarang Gopalakrishnan, and Romain Vasseur. Entanglement and charge-sharpening transitions in u(1) symmetric monitored quantum circuits. *Phys. Rev. X*, 12:041002, Oct 2022. doi:10.1103/PhysRevX.12.041002. URL <https://link.aps.org/doi/10.1103/PhysRevX.12.041002>.
- [34] Fergus Barratt, Utkarsh Agrawal, Sarang Gopalakrishnan, David A. Huse, Romain Vasseur, and Andrew C. Potter. Field theory of charge sharpening in symmetric monitored quantum circuits. *Phys. Rev. Lett.*, 129:120604, Sep 2022. doi:10.1103/PhysRevLett.129.120604. URL <https://link.aps.org/doi/10.1103/PhysRevLett.129.120604>.

- [aps.org/doi/10.1103/PhysRevLett.129.120604](https://doi.org/10.1103/PhysRevLett.129.120604).
- [35] Michael J. Gullans and David A. Huse. Scalable probes of measurement-induced criticality. *Phys. Rev. Lett.*, 125:070606, Aug 2020. doi:10.1103/PhysRevLett.125.070606. URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.070606>.
- [36] Thomas Vojta and Ronald Dickman. Spatiotemporal generalization of the harris criterion and its application to diffusive disorder. *Phys. Rev. E*, 93:032143, Mar 2016. doi:10.1103/PhysRevE.93.032143. URL <https://link.aps.org/doi/10.1103/PhysRevE.93.032143>.
- [37] Henry Ando and David A. Huse. Conserved quantities in models of classical chaos. *arXiv:2307.11733*, 2023. URL <https://arxiv.org/abs/2307.11733>.
- [38] A B Harris. Effect of random defects on the critical behaviour of ising models. *Journal of Physics C: Solid State Physics*, 7(9):1671, may 1974. doi:10.1088/0022-3719/7/9/009. URL <https://dx.doi.org/10.1088/0022-3719/7/9/009>.
- [39] See the Supplemental Material.
- [40] Kai Klocke, Joel E. Moore, and Michael Buchhold. Power-law entanglement and hilbert space fragmentation in non-reciprocal quantum circuits. 2024. URL <https://arxiv.org/abs/2405.06021>.

Supplemental Material for “Measurement-induced phase transitions in systems with diffusive dynamics”

Hyunsoo Ha,¹ Akshat Pandey,² Sarang Gopalakrishnan,³ and David A. Huse¹

¹*Department of Physics, Princeton University, Princeton, New Jersey 08544, USA*

²*Department of Physics, Stanford University, Stanford, California 94305, USA*

³*Princeton University, Department of Electrical and Computer Engineering, Princeton, NJ 08544, USA*

(Dated: May 16, 2024)

In this Supplemental Material, we (i) overview the diagnostics we used to probe the transition and the phases, (ii) present further results near the critical point, (iii) investigate the code length during the purification process, and (iv) discuss the periodic structure seen in $P_{S_{N/2}}^<(t)$ in the volume-law phase.

S1. Diagnostics Overview

A. Entanglement probes

We use entanglement probes to investigate the phase diagram of our model. Starting from a pure state (mostly using product initial states, but we can also start from random pure states), we characterize distributions of several quantities over the course of the dynamics, as well as in the long-time steady state. Specifically, we consider the (i) half-chain entanglement entropy $S_{N/2}$, (ii) antipodal mutual information: AMI, and (iii) tripartite mutual information \mathcal{I}_3 , defined as follows:

$$S_{N/2} \equiv S_A; \tag{S1}$$

$$\text{AMI} \equiv S_A + S_B - S_{A \cup B}; \tag{S2}$$

$$\mathcal{I}_3 \equiv S_A + S_B + S_C - S_{A \cup B} - S_{B \cup C} - S_{A \cup C} + S_{A \cup B \cup C}. \tag{S3}$$




Figure S1. Illustrations of partitions into subsystems involved in various entanglement probes. (Left) Half-chain bipartite entanglement $S_{N/2}$ from the periodic and open boundary condition. (Middle) Antipodal mutual information (AMI) between subregions A and B with length $N/8$ and centers $N/2$ apart. (Right) Tripartite mutual information \mathcal{I}_3 where $|A| = |B| = |C| = N/4$.

The bipartite entanglement $S_{N/2}$ is the simplest probe diagnosing the volume-law and the area-law phase, characterized by $S_{N/2} \sim N$ and $S_{N/2} \sim O(1)$, respectively, in the steady state. Also, with a product initial state in the volume-law phase, $S_{N/2}(t)$ increases linearly in t before it saturates to an order- N value. We additionally investigate $S_{N/2}$ with open boundary conditions while exploring the Griffith-like effects because now, a single cut defines a bipartition, and it better captures the rare-region effect. However, $S_{N/2}$ is sensitive to entanglement on all length scales. AMI and \mathcal{I}_3 are, in some situations, nicer objects to work with, since they are sensitive only to long-range entanglement.

In particular, we use the antipodal mutual information AMI to estimate the location of the critical point.¹ We expect $\mathcal{I}_{AB} \rightarrow 0$ in both volume-law and area-law phases in the thermodynamic limit, so for finite systems AMI has a peak near the critical point, which can be found without making any scaling assumption.

We also investigated the tripartite mutual information \mathcal{I}_3 , a probe that has been found to be useful for precisely characterizing the spacetime-uncorrelated conventional MIPT.² It is asymptotically zero in the area-law phase and is

negative and proportional to N at the volume-law phase. However, the value of \mathcal{I}_3 is sensitive to the specific circuit realization, and the probability distribution $\mathcal{P}[\mathcal{I}_3]$ exhibits a fat tail under strong space-time correlation, i.e., quenched randomness.^{3,4} In such scenarios, the average value of \mathcal{I}_3 is not representative of the distribution, and it may be more informative to study the quantity $\mathcal{P}[\mathcal{I}_3 = 0]$, which we also explore.

B. Purification dynamics

Together with the entanglement probes, we investigate purification dynamics,⁵ establishing a direct connection between space and time. Specifically, we prepare a fully mixed initial state $\hat{\rho}_i = \hat{\mathbb{I}}/2^N$ and track the von Neumann entropy $S(t) = -\text{tr}(\hat{\rho}_t \log_2 \hat{\rho}_t)$ over time. The purification time τ_P is when the system is completely purified, so that $S(t) = 0$ for all $t \geq \tau_P$. The volume-law to area-law phase transition in the conventional MIPT is also a transition from a “mixed” phase with exponentially long purification times $\log \tau_P \sim N$, to a “purifying” phase where τ_P grows only as $\log N$. At the critical point, $\tau_P \sim N^z$ with $z = 1$ for spacetime-uncorrelated measurements, while $\log \tau_P \sim N^\psi$ (so that $z = \infty$) for quenched spatial randomness.

C. Contiguous code length

The contiguous code length⁵⁻⁷ shows where the logical operator associated with the state is located. In this work, we calculate the contiguous code length when there is a single logical pair (or, equivalently, $(N - 1)$ stabilizers). A single logical pair can be prepared in the purification protocol starting from the maximally mixed state, waiting until exactly one bit remains unpurified. Calling the logical operators $\hat{\ell}_X$, $\hat{\ell}_Y$, and $\hat{\ell}_Z$, where the logicals follow the same algebra as Pauli \hat{X} , \hat{Y} , and \hat{Z} operators respectively, we can find the shortest contiguous interval that supports each logical operator, with the respective lengths d_α defined as

$$d_\alpha = \min_A \{|A| : \exists \hat{\ell}_\alpha \text{ supported in } A\}, \quad \alpha \in \{X, Y, Z\}. \quad (\text{S4})$$

where A denotes an interval within the chain.

Initially, we entangle the logical pair with a reference ancilla (labeled a) to form a Bell pair so that $\hat{X}_a \otimes \hat{\ell}_X$ and $\hat{Z}_a \otimes \hat{\ell}_Z$ are the stabilizer generators of the ancilla-extended system. We then run the circuit solely on the system, so the time-evolved logicals are $\hat{X}_a \otimes \hat{\ell}_X(t)$, $\hat{Y}_a \otimes \hat{\ell}_Y(t)$, and $\hat{Z}_a \otimes \hat{\ell}_Z(t)$. The system gets purified at time τ_P once the measurements coincide with the logical operator (assuming $\hat{\ell}_X(\tau_P)$ is measured), leading to the disentanglement of the ancilla from the system. The stabilizers at the purification time are $\hat{\mathbb{I}}_a \otimes \hat{\ell}_X(\tau_P)$ and $\hat{X}_a \otimes \hat{\mathbb{I}}$.

To independently capture the code length for each logical component, we adopt the so-called clipped gauge.^{1,8} Each stabilizer generator has a left and a right “end” of nontrivial content, so a pure state of length L has a total of $2L$ ends. The clipped gauge imposes the condition that exactly two ends, originating from either different generators or the same one, are positioned at every site. Such a representation allows efficient calculation S_A of the contiguous interval A by counting the string of generators with one end inside the interval and the other outside.

When applying the clipped gauge to a system with a single logical pair coupled to an ancilla as a Bell pair (so the total length is $N + 1$), we find two stabilizer generators with nontrivial content starting from the first site (the ancilla’s location) and ending at $r_1 + 1$ and $r_2 + 1$, respectively ($r_1 \leq r_2$). The mutual information between the ancilla and the system within an interval from 1 to r (denoted as $A_{[1, \dots, r]}$) is determined as follows:

$$\mathcal{I}(\text{ancilla}; A_{[1, \dots, r]}) = \begin{cases} 0 & (r < r_1), \\ 1 & (r_1 \leq r < r_2), \\ 2 & (r_2 \leq r). \end{cases} \quad (\text{S5})$$

If the shorter stabilizer generator has Pauli operator $\hat{\alpha}_a$ at the ancilla, then $A_{[1, \dots, r]}$ with $r_1 \leq r < r_2$ supports the logical operator $\hat{\ell}_\alpha$, while the other two are absent. For intervals longer than r_2 , they include all three ($\hat{\ell}_X(t)$, $\hat{\ell}_Y(t)$, and $\hat{\ell}_Z(t)$) logical operators:

$$A_{[1, \dots, r]} \text{ includes } \begin{cases} \text{no logicals} & (r < r_1), \\ \hat{\ell}_\alpha & (r_1 \leq r < r_2), \\ \hat{\ell}_X, \hat{\ell}_Y, \text{ and } \hat{\ell}_Z & (r_2 \leq r). \end{cases} \quad (\text{S6})$$

To determine the code length of logicals in a periodic chain, we repeatedly compute in the clipped gauge while keeping the ancilla position fixed, cyclically permuting the system. During each iteration, we obtain r_1 and r_2 , and identify which logical component is supported on length r_1 (flagged with \hat{a}_a). Subsequently, we designate length r_1 to α and r_2 to $\{X, Y, Z\} \setminus \{\alpha\}$, comparing these values across iterations to identify the shortest length and its corresponding interval for all X, Y , and Z logical components.

We note that the shortest interval supporting each logical may not be unique, but the shortest length is well defined. Furthermore, since logicals with different components anticommute, the intervals supporting different logicals should always overlap on at least one site.

S2. Results near the critical point

A. Determining the critical point




Figure S2. (a) Re-scaled AMI for large system sizes ($N \geq 128$) (b) $\log_{10} Q_{\text{AMI}}$, an indicator of the quality of the one-parameter scaling ansatz Eqn. S7 for different values of β and ν . Smaller values represent the better collapse, with $\log_{10} Q_{\text{AMI}} = 0$ indicating the best result. (c) $\mathcal{P}[I_3 = 0]$ at the steady value for various measurement rates and system sizes. Inset shows the finite-size scaling of $\mathcal{P}[I_3 = 0]$ for large system sizes ($N \geq 128$). (d) $\log_{10} Q_{I_3}$ shows the quality of the scaling ansatz Eqn. S8 for different values of ν and p_c .

As explained in the main text, we rely on the steady-state antipodal mutual information to determine the critical measurement rate, constrained to be in between the interval $p_c \in (0.31, 0.35)$. Solely focusing on the peak of AMI suggests $p_c \approx 0.33$. The increasing peak for larger system sizes is a notable difference from the conventional space-time uniform MIPT, where AMI has a constant value at the critical point. The evident increasing peak indicates the absence of conformal invariance in our model.

For each sample, circuits are applied until time $t = 5t_{\text{classical}} + 1000$, and the entanglement probes for each sample are obtained by averaging over the last 1000 time steps. Here, $t_{\text{classical}} = (N/2\pi)^2$ represents the relaxation time of the classical SSEP particles. We carefully checked that all the probes were well saturated to the steady-state value after $5t_{\text{classical}}$ (within our accessible system sizes/times).

Adopting the peak to be p_c , next, we attempt to extract the correlation-length exponent ν from the following scaling ansatz

$$\text{AMI} \sim N^\beta f(N^{1/\nu}(p - p_c)) \quad (\text{S7})$$

where f is the scaling function. As shown in Fig. S2(a), we use $N=128, 256$, and 512 to avoid being distracted by the smaller system sizes, which are unlikely to scale properly, and find $\nu = 1.6$. However, the data collapse is poor, as we quantified in Fig. S2(b) with the objective function Q_{AMI} that determines the quality of the scaling ansatz for different values of β and ν .^{2,9}

We additionally present results on the tripartite mutual entanglement $\mathcal{P}[I_3 = 0]$ in Fig. S2(c). The pronounced drift towards larger p_c for larger system sizes becomes evident as we focus on the intersection of the two graphs, indicating significant finite-size effects. We again attempt to extract the correlation length exponent from the following ansatz

$$P[I_3 = 0] \sim g(N^{1/\nu}(p - p_c)) \quad (\text{S8})$$

with a scaling function g , and get $\nu = 1.5$ as shown in the inset of Fig. S2(c). Fig. S2(d) shows the objective function for the scaling ansatz Eqn. S8. From both AMI and $P[I_3 = 0]$ observables, we extract $\nu = 1.6(5)$; however, the large

uncertainties presented in Fig. S2(b,d) and the strong finite-size effects shown in Fig. S2(c) that everything may still strongly drift hinder the accurate characterization of the scaling relations.

B. Entanglement growth at the critical point




Figure S3. Entanglement probes at $p = 0.33$. (a) $S_2(t)$ shows a logarithmic trend. (b,c) Steady-state properties are presented with (b) S_2 and (c) AMI, averaged over samples, showing an algebraic increase over sample sizes.

Once the critical point is identified, we investigate the entanglement growth and explicitly present the result for $p = 0.33$ in Fig. S3, indicating a logarithmic trend [Fig. S3(a)]. A notable distinction is observed in the steady-state entanglement, as depicted in Fig. S3(b,c), where the relation between S_2 and AMI over various system sizes demonstrates an algebraic relation $\bar{S} \sim N^{\psi'}$ with $\psi' \approx 0.4$. This behavior resembles the quenched randomness critical point, which has $\bar{S} \sim N^{1/2}$, and the power-law scaling was also observed in the system with quantum and classical circuits coupled together.¹⁰

C. Purification dynamics at the critical point




Figure S4. (a) Purification time τ_P . The median over samples is presented as a representative of the typical samples. The inset shows activated dynamical scaling at $p = 0.33$, with the x-axis rescaled to $N^{0.44}$. (b) The lifetime of the last logical bit τ_1 . The inset shows $p = 0.33$ with the x-axis rescaled to $N^{0.39}$. The measurement rates (p) are indicated in both graphs with numbers on each plot with different colors. The gray dotted line is the relaxation time scale for the diffusive measurers with $z = 2$.




Figure S5. Entropy from the purification dynamics, presented over re-scaled time t/N^2 .

As explained in the main text, we explore the purification dynamics to find a direct connection between space and time at the critical point. As shown in Fig. S4 (a), it is clear that the new critical point has $z > 1$, indicating a different universality class from the conventional MIPT. From the constrain of $p_c \in (0.31, 0.35)$ based on the AMI results, we can bound the dynamical exponent $z \geq 2$, and there is a signature of diverging dynamical exponent $z \rightarrow \infty$ in the thermodynamic limit if $p_c = 0.33$.

The long-lived behavior of the last logical pair also captures the extraordinarily long timescale. Starting from the fully mixed state with entropy $S(0) = N$, the entropy monotonically decreases over time. Furthermore, the entropy is always an integer, due to the Clifford character of our circuits. Therefore, the moment when the entropy remains precisely one bit — the “last logical” — is well defined. The probability distribution of the last logical’s survival time turns out to have an exponential tail near the critical point. One can extract the last logical’s lifetime τ_1 by fitting the exponential tail. This lifetime has roughly the same behavior as the purification times, as shown in Fig. S4(b).

We also present how the entropy $S(t)$ decays over time (scaled as t/N^2) at different measurement rates [Fig. S5]. We investigate the possibility of the dynamical exponent being the same as that of the measurers’ SSEP viz. $z = 2$. For $p = 0.33$, the entropy develops a plateau for larger system sizes, clearly showing $z > 2$, while for the slightly higher measurement rate ($p = 0.35$), the plots seem to collapse for various system sizes. Collapsing plots near $p \approx 0.35$ with re-scaled time t/N^2 suggests a scenario, discussed in the main text, where p_c is larger than the $p \approx 0.33$ which has heretofore been the focus of our discussion, with $z = 2$ dynamical scaling at criticality. Even at these rather large system sizes, we believe our model suffers from significant finite-size corrections to scaling. We leave further investigation to characterize the critical behavior for future work.

S3. Purification of the last logical bit

In this section, we focus on the $z \rightarrow \infty$ scenario and provide a qualitative argument focusing on the code length of the last logical bit. Naively, if one considers a rare region of length ℓ with a relatively low density of measurers. The reduced density in this rare region leads to a lower measurement rate, allowing logical operators to remain secure inside. However, due to the diffusive nature of measurers, the rare region has a finite lifetime of $O(\ell^2)$, and the logicals will be exposed to a higher measurement rate afterward, suggesting that the dynamical exponent should be bounded by $z \leq 2$.

Contrary to expectations, however, the logical can hop between rare regions, as illustrated in Section S5. This occurs when one logical operator (denoted as \hat{X}_L) increases in size while the other logical operator (\hat{Z}_L) maintains its length. The shorter logical appears to tunnel between rare regions, taking advantage of the sufficiently long pair. This mechanism may be necessary so that the logical can persist even after the rare region dissipates, allowing the possibility of $z > 2$.

A. Contiguous code length of the last logical




Figure S6. (a,b) Contiguous code length of the logical before it gets purified. (a) Standard MIPT in Clifford circuits and (b) our diffusive measurer model at $p = 0.33$ are presented. The x-axis displays $\tau_P - t$, the remaining time before being purified. All calculations are performed in the system length $N = 128$. (c) Probability distribution of the contiguous code length. Both standard MIPT¹ at the critical point and diffusive model at $p = 0.33$ are presented. The system size increases as the plot gets darker. The code length scales proportional to the system size.

Fig. S6(a,b) illustrates the contiguous code length of the last logical pair for the standard MIPT in Clifford circuits¹ at the critical point, and in our model with $p = 0.33$. The x-axis denotes $\tau_P - t$, the time until purification. The code length of three nontrivial logical operators \hat{X}_L , \hat{Z}_L , and $\hat{Y}_L = i\hat{Z}_L\hat{X}_L$ was calculated, and we averaged the shortest, longest, and mean values over samples.

The shortest logical operator consistently shrinks to zero (single-site occupation) preceding purification in both models. However, the distinction is clear when one focuses on the longest logical operator. In contrast to the standard model, the diffusive MIPT model exhibits an increase in the length of the longest logical operator. This observation supports our previously addressed mechanism.

The long-lived behavior is also reflected in the contiguous code length probability distribution, shown in Fig. S6(c). In the conventional MIPT, the probability is finite at the zero-length, while the diffusive model produces a clear peak near $0.1N$, with the distribution vanishing at zero code length. This implies the code length remains extensive most of the time and has a much smaller chance of going to zero.

B. Hyperuniformity of measurers when the last logical purifies

From the observations on code length above, we qualitatively and quantitatively analyzed that (1) the contiguous code distance is proportional to the system size for the typical times, and (2) even very short logical operators with $O(1)$ length may be secured by the rare region with a lower density of measurers, and can grow longer. Based on these observations, we propose that the purification lifetime is controlled by events when the spatial rare regions




Figure S7. (a) Schematic of the height function $W(x)$. (b) Probability distribution of the $W(x)$ variance sampled when the ancilla is disentangled from the system. Compared to the control group (blue) from randomly distributed measurers, the probability distribution of $\text{var}(W)$ gets sharper, and the peak shifts towards zero. (c) Hyperuniformity of the measurers with wandering exponent $\beta = 0.415$. The sample-averaged standard deviation of W and the system sizes have clear algebraic relations, as shown in the log-log plot.

are suppressed system-wide. This leads to ultra-slow dynamics with purification time characterized by stretched exponential of system size.

As a measure of the uniformity of classical particles, we define a height function $W(x)$:

$$W(x) = \sum_{r=1}^x (-1)^{n(r)}, \quad (\text{S9})$$

with $W(0) = 0$, and $n(r)$ being the occupation number of measurers at r . We have $W(N) = W(0)$, since the number of measurers is exactly half the system size. If measurers are randomly distributed, which is true for the typical distribution of SSEP particles, the standard deviation of the height function scales $\sigma(W) \sim N^{1/2}$. However, the measurer locations are not random at the moment when the system gets totally purified, and hyperuniformity,^{11–14} characterized by the wandering exponent β , is observed at this particular time: $\sigma(W) \sim N^\beta$, with $\beta = 0.415$ [Fig. S7(c)]. Fig. S7(b) also shows the distribution $\text{var}(W)$ at the purification time is clearly different from the distribution for randomly distributed measurers (blue line). The probabilistic chance of measurers to locate hyperuniformly is approximately $\exp(N^{1-2\beta})$ (ignoring constants). This heuristic mechanism supports a picture of ultra-slow dynamics with $\psi \approx 1 - 2\beta$.

S4. In-periodic structure of $P_{S_2}^<(t)$

The tail distribution of entanglement growth effectively captures Griffiths-like effects within the volume-law phase, arising from rare regions of low measurer density with finite lifetimes due to particle diffusion. We define $P_{S_2}^<(t)$ as the probability of bipartite entanglement being suppressed by more than half the average over samples for a single cut. In our analysis, we observed distinct scaling behaviors for $P_{S_2}^<(t)$ across three classes of models with different spacetime correlations of measurements.

However, alongside the anticipated scaling relations, we observed the emergence of a nearly periodic structure over time (see Fig. 3 in the main text). This periodicity is a characteristic feature of the Clifford circuit, where all $S_2(x, t)$ values are integers for each run. Unlike individual samples of S_2 , the mean value \bar{S}_2 , and consequently $\bar{S}_2/2$, are not necessarily integers. As \bar{S}_2 increases by 2 units, the range of integers considered for $P_{S_2}^<(t)$ expands as a step function over time.

This characteristic is readily apparent in the spacetime-uncorrelated MIPT,¹ where entanglement grows ballistically, i.e., $\bar{S}_2(N/2, t) = v_E t$. Consequently, the range of counting for $P_{S_2}^<(t)$ increases every $2/v_E$ periods, leading to the observed periodic structure over time. Figure S8 illustrates this feature for measurement rates $p = 0.08$ and $p = 0.04$, where the period is indeed determined by $2/v_E$.




Figure S8. (a,b) Results from an open-boundary chain of length $N = 1024$ and measurement rate $p = 0.08$. The measurements are spacetime-independent and identically distributed, and the measurement rates are low enough to be well inside the volume-law phase. In (a), we observe ballistic growth of bipartite entanglement from a single cut at the middle, indicating a velocity of entanglement propagation of approximately $v_E \sim 0.15$. In (b), the nearly periodic structure in $P_{S_2}^<(t)$ has a period of approximately $T \sim 13$, which aligns with the expected period of $2/v_E$, as explained in the text. (c,d) Here, we present another demonstration with different measurement rates, $p = 0.04$, and find $v_E \sim 0.25$ and $T \sim 8$, consistent with the anticipated relationship.

S5. Appendix: visualizing the last remaining logical bit

We here visualize the last remaining logical bit for $p = 0.33$ with the contiguous code length and the corresponding interval described in Section S1 C. In Fig. S9, we show the shortest(longest) logical operator among $\hat{\ell}_X$, $\hat{\ell}_Y$, and $\hat{\ell}_Z$ at individual time with yellow(green). The background gray shows the space-time location of the measurements.

To make a stark contrast in spatial measurer density to apparently visualize the rare regions, we next run the dimer model, where the measurers form a pair as a dimer, and each dimer evolves under SSEP rules. Mesurer particle individually measures the qubit with probability p . We picked the number of dimers to be $N_d = N/8$, so the total number of particles is $N/4$. In this model, the AMI value peaked at $p = 0.65$; hence, we picked $p = 0.65$ in Fig. S10.

-
- ¹ Yaodong Li, Xiao Chen, and Matthew P. A. Fisher, “Measurement-driven entanglement transition in hybrid quantum circuits,” *Phys. Rev. B* **100**, 134306 (2019).
 - ² Aidan Zabalo, Michael J. Gullans, Justin H. Wilson, Sarang Gopalakrishnan, David A. Huse, and J. H. Pixley, “Critical properties of the measurement-induced transition in random quantum circuits,” *Phys. Rev. B* **101**, 060301 (2020).
 - ³ Aidan Zabalo, Justin H. Wilson, Michael J. Gullans, Romain Vasseur, Sarang Gopalakrishnan, David A. Huse, and J. H. Pixley, “Infinite-randomness criticality in monitored quantum dynamics with static disorder,” *Phys. Rev. B* **107**, L220204 (2023).
 - ⁴ Gal Shkolnik, Aidan Zabalo, Romain Vasseur, David A. Huse, J. H. Pixley, and Snir Gazit, “Measurement induced criticality in quasiperiodic modulated random hybrid circuits,” *Phys. Rev. B* **108**, 184204 (2023).
 - ⁵ Michael J. Gullans and David A. Huse, “Dynamical purification phase transition induced by quantum measurements,” *Phys. Rev. X* **10**, 041020 (2020).
 - ⁶ Sergey Bravyi and Barbara Terhal, “A no-go theorem for a two-dimensional self-correcting quantum memory based on stabilizer codes,” *New Journal of Physics* **11**, 043029 (2009).
 - ⁷ Matteo Ippoliti, Michael J. Gullans, Sarang Gopalakrishnan, David A. Huse, and Vedika Khemani, “Entanglement phase transitions in measurement-only dynamics,” *Phys. Rev. X* **11**, 011030 (2021).
 - ⁸ Adam Nahum, Jonathan Ruhman, Sagar Vijay, and Jeongwan Haah, “Quantum entanglement growth under random unitary dynamics,” *Phys. Rev. X* **7**, 031016 (2017).
 - ⁹ Naoki Kawashima and Nobuyasu Ito, “Critical behavior of the three-dimensional $\pm j$ model in a magnetic field,” *Journal of the Physical Society of Japan* **62**, 435–438 (1993), <https://doi.org/10.1143/JPSJ.62.435>.
 - ¹⁰ Kai Klocke, Joel E. Moore, and Michael Buchhold, “Power-law entanglement and hilbert space fragmentation in non-reciprocal quantum circuits,” (2024), [arXiv:2405.06021 \[cond-mat.stat-mech\]](https://arxiv.org/abs/2405.06021).
 - ¹¹ Salvatore Torquato and Frank H. Stillinger, “Local density fluctuations, hyperuniformity, and order metrics,” *Phys. Rev. E* **68**, 041113 (2003).
 - ¹² Salvatore Torquato, “Hyperuniformity and its generalizations,” *Phys. Rev. E* **94**, 022122 (2016).
 - ¹³ Zheng Ma and Salvatore Torquato, “Random scalar fields and hyperuniformity,” *Journal of Applied Physics* **121**, 244904 (2017).

- (2017), https://pubs.aip.org/aip/jap/article-pdf/doi/10.1063/1.4989492/19773224/244904_1_online.pdf.
¹⁴ Salvatore Torquato, "Hyperuniform states of matter," *Physics Reports* **745**, 1–95 (2018), hyperuniform States of Matter.




Figure S9. The shortest logical operator (left) and both the longest and the shortest logical pair (right) in our diffusive model.




Figure S10. The shortest logical operator (left) and both the longest and the shortest logical pair (right) in the dimer model.