ERGODICITY OF SOME STOCHASTIC FOKKER-PLANCK EQUATIONS WITH ADDITIVE COMMON NOISE

FRANÇOIS DELARUE, RAPHAËL MAILLET, ETIENNE TANRÉ

ABSTRACT. In this paper we consider stochastic Fokker-Planck Partial Differential Equations (PDEs), obtained as the mean-field limit (i.e., as the number of particles tends to ∞) of weakly interacting particle systems subjected to both independent (or idiosyncratic) and common Brownian noises. We provide sufficient conditions under which the deterministic counterpart of the Fokker-Planck equation, which corresponds to particle systems that are just subjected to independent noises, has several invariant measures, but for which the stochastic version admits a unique invariant measure under the presence of the additive common noise. The very difficulty comes from the fact that the common noise is just of finite dimension while the state variable, which should be seen as the conditional marginal law of the system given the common noise, lives in a space of infinite dimension. In this context, our result holds true if, in addition to standard confining properties, the mean field interaction term forces the system to be attracted by its conditional mean given the common noise and the intensity of the idiosyncratic noise is small.

Mathematics Subject Classification (2020): 37A50, 35Q70, 60H15, 60J25

Keywords: Non-linear Fokker-Planck equations , McKean-Vlasov equations with common noise, Stochastic Partial Differential Equations, Common noise, Asymptotic stability

Contents

1. Introduction	2
1.1. Deterministic Fokker-Planck equations	2
1.2. Common noise	3
1.3. Our contribution	4
1.4. Further connections with the literature on mean field models	5
1.5. Organisation of the paper	5
1.6. Notation	5
2. Main result: uniqueness recovery thanks to common noise	6
2.1. Stochastic Fokker-Planck PDE and Conditional McKean-Vlasov SDE	6
2.2. Assumptions, existence and uniqueness of solutions	7
2.3. Restoration of uniqueness	9
2.4. Discussions and examples	14
3. Proof of Theorem 8	19
3.1. First preliminary estimate: conditional variance	19
3.2. Second preliminary estimate: conditional exponential moments	20
3.3. Completion of the proof of Theorem 8	22

Received by the editors May 17, 2024.

F. DELARUE ; R. MAILLET ; E. TANRÉ

4. Proof of Proposition 7	22
4.1. Ansatz	22
4.2. Contraction for classical diffusion processes	23
4.3. Proof of Proposition 7	25
Acknowledgments	34
References	34
Appendix A.	36
A.1. Proof of Equation (38)	36
A.2. Proof of Equation (55)	37

1. INTRODUCTION

This paper addresses the long-term behavior of solutions to a class of non-linear Stochastic Partial Differential Equations (SPDEs) of the form

(1)
$$d_t m_t = \nabla \cdot \left(\frac{\sigma^2 + \sigma_0^2}{2} \nabla m_t - m_t b(\cdot, m_t)\right) dt - \sigma_0 \nabla m_t \cdot dB_t^0, \quad \text{on } [0, +\infty) \times \mathbb{R}^d,$$

where $B^0 = (B_t^0)_{t \ge 0}$ is a *d*-dimensional Brownian motion, called *common noise*. Above, the unknown $(m_t)_{t \ge 0}$ is regarded as a stochastic process with values in the space $\mathcal{P}(\mathbb{R}^d)$ of probability measures on \mathbb{R}^d . Accordingly, the coefficient *b* is a function from $\mathbb{R}^d \times \mathcal{P}(\mathbb{R}^d)$ to \mathbb{R}^d depending on both space and measure arguments. The two parameters σ and σ_0 are non-negative scalars: the former is the intensity of the so-called idiosyncratic noise and the latter is the intensity of the common noise.

Equation (1) is in fact intended to describe the flow of conditional marginal laws of the solution to the conditional McKean-Vlasov equation:

(2)
$$dX_t = b(X_t, \mathcal{L}(X_t|B^0))dt + \sigma dB_t + \sigma_0 \nabla m_t \cdot dB_t^0, \quad t \ge 0,$$

where *B* is another Brownian motion, independent of B^0 , and $\mathcal{L}(\cdot|B^0)$ stands for the conditional law given the realization of the common noise. Formally, m_t in (1) is expected to coincide with $\mathcal{L}(X_t|B^0)$ in (2). The rigorous connection between (1) and (2) is addressed more carefully in the core of the paper.

1.1. **Deterministic Fokker-Planck equations.** When $\sigma_0 = 0$, i.e., in absence of common noise, Equation (1) boils down to a standard Fokker-Planck equation describing the evolution of the marginal laws of the solutions to the standard McKean-Vlasov equation (2) obtained by replacing $\mathcal{L}(X_t|B^0)$ by the law of X_t . In this setting, the long-time analysis of the solutions has been a notoriously challenging problem for over twenty years, giving rise to numerous contributions, many of them still recent. While the problem is of mathematical interest in its own right, it also finds some application to calculus of variation since certain equations like (1) can be interpreted as gradient flows on the space of probability measures, see the seminal works [1, 15, 37, 49, 50]. For instance, such a gradient structure occurs when $b(x, \mu) = -\nabla V(x) - \nabla W * \mu(x)$, for two differentiable real-valued functions V and W on \mathbb{R}^d , W being symmetric, where * stands for the

2

convolution operator. In this case, the potential lying above the dynamics writes

(3)
$$\mu \in \mathcal{P}(\mathbb{R}^d) \mapsto \int_{\mathbb{R}^d} V(x)\mu(\mathrm{d}x) + \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} W(x-y)\mu(\mathrm{d}x)\mu(\mathrm{d}y) + \frac{\sigma^2}{2} \int_{\mathbb{R}^d} \ln(\frac{\mathrm{d}\mu}{\mathrm{d}x}(x))\mu(\mathrm{d}x).$$

In fact, due to the mean field interaction (whether it derives from a potential or not), the sole presence of the noise B in the dynamics (2) (with $\sigma_0 = 0$) does not suffice to guarantee the uniqueness of an invariant measure (or equivalently of a stationary solution to (1)) under quite general conditions on b. This is in contrast with the long run behavior of Stochastic Differential Equations (SDEs) (or equivalently linear Fokker-Planck equations), for which mere confining properties of the drift suffice to produce ergodicity (provided $\sigma > 0$). In the mean field setting, some further structural conditions are necessary. In the potential example (3), a standard condition is to demand V and W to be (strongly) convex, which forces in fact a form of convexity of the potential (3) on the space of probability measures. We refer for instance to the earlier works [7, 15, 45, 46], which also include additional convergence results to the stationary regime. For a tiny list of variants, in which *perturbations* of the convex potential case are addressed, we refer to [6, 8, 10, 21, 25]. We refrain from detailing all the possible extensions of the previous cases under which the invariant measure remains unique. Let us just say, for our purposes, that the above list of references includes cases where the lack of convexity is compensated for by the presence of a strong enough diffusion coefficient. This large noise regime is compared with our own setting in Remark 9 below. At this point, we would especially like to stress that there are known explicit simple cases in which uniqueness does not hold, which observation supports our previous claim: the presence of a nonlinear mean field term may easily lead to the existence of multiple invariant measures, even in situations where the mass stays confined under the dynamics (1). Specifically, it has been proven in [35] (see also [19]) that, when V is only uniformly convex outside of a ball but admits a double-well, W is quadratic (but non-zero) and the diffusion coefficient σ is sufficiently small, there exist several stationary solutions to (1) when it derives from the potential (3) (and $\sigma_0 = 0$).

1.2. **Common noise.** Very basically, our primary objective in this article is to revisit the class of examples addressed in [35], but in presence of a common noise, i.e., $\sigma_0 > 0$. In this regard, it is worth noting that, when $b(x, \mu) = -\nabla V(x) - \nabla W * \mu(x)$, with *V* being uniformly convex and *W* being convex, the common noise does not change the picture recalled in the previous paragraph. Indeed, the second author has shown in [44] that (1) has a unique invariant measure, which is consistent with the results obtained earlier without common noise with the slight subtlety that the invariant measure is then understood as a probability measure on $\mathcal{P}(\mathbb{R}^d)$.

Of course, our goal is to go one step beyond and prove that the common noise can change the picture positively, meaning that it can force uniqueness of the invariant measure even though the deterministic analogue of (1) (i.e., with $\sigma_0 = 0$) has several stationary solutions. Actually, this type of result, if it holds true, must be part of the very broad theory of ergodic Markov processes, with the specific feature that the state space here is $\mathcal{P}(\mathbb{R}^d)$. It is worth observing that, from the same point of view, the deterministic and linear analogue of (2) (i.e., *b* only depends on *x* and $\sigma_0 = 0$) should be regarded as a Fokker-Planck equation obtained by forcing a standard ordinary differential equation by a Brownian motion of intensity σ . This is just to say that, in the Euclidean setting, the same program is very well-understood and just consists in addressing the ergodic properties of a non-degenerate SDE. For sure, the very difficulty in our setting comes from the fact that the unknown in (1) lives in a space of infinite dimension (once again, the space of probability measures) while the noise (i.e., B^0) is just finite-dimensional and thus *completely* degenerate. This says that, at best, noise can be expected to restore uniqueness of the invariant measure only in specific cases. And, precisely, this is our objective to identify a class of cases that would become ergodic under the action of the common noise and that would include some of the examples addressed in [35].

In fact, this question was already addressed by the second author in [44] when $\sigma = 0$, leaving open setting with $\sigma > 0$. Apart from this, the idea of using the averaging properties of the common noise has been used in several contexts, for instance in the analysis of existence and uniqueness of equilibria to mean field games (see for instance [23, 29] for games with a finite dimensional common noise and [22] for games with an infinite dimensional common noise). More recently, the first author has just released an article, [24], in which the ergodic properties of (1) (or, more precisely, (2)) are studied when *b* is general but the common noise is infinite dimensional and the dimension *d* is 1. Part of the challenge then precisely lies in the construction of the noise, which is a question different from the one addressed here. A similar problem was addressed in the prior work [3], for another type of infinite dimensional common noise that may not preserve the space of probability measures.

1.3. Our contribution. We here focus on drifts of the type

$$b(x,\mu) = G(x) + F\left(x - \int_{\mathbb{R}^d} x \,\mu(\mathrm{d}x)\right), \quad x \in \mathbb{R}^d, \, \mu \in \mathcal{P}(\mathbb{R}^d),$$

where *F* and *G* are functions from \mathbb{R}^d into itself. The function *F* is assumed to be strictly decreasing, which occurs for instance if $F = -\nabla W$ for a strictly convex function. More interestingly, *G* is just assumed to be confining in a mild sense that is detailed in Assumption 1 below. In particular, *G* is not required to be strictly decreasing and, when it derives from a potential *V*, *V* may not be convex. Therefore, this framework allows us to choose $G = -\nabla V$ for a non-convex potential *V* and $F = -\nabla W$, for $W(x) = \alpha x^2$ for some $\alpha > 0$, and thus contains the case treated by [35] (up to an additional common noise), since $b(x, \mu)$ rewrites $-\nabla V(x) - \nabla W * \mu(x)$.

Our main contribution is to establish that, for a small diffusion coefficient σ and for an interaction force F that is sufficiently decreasing, the system has a unique invariant measure, which attracts exponentially fast any other initial condition (with appropriate integrability properties). Our strategy of proof is based on the key idea that the long run behavior of (2) should be dictated by the long time dynamics of the conditional mean $\mathbb{E}[X_t|B^0]$ given the common noise. However, this intuition is not completely correct, as the distance between X_t and its conditional mean $\mathbb{E}[X_t|B^0]$ does not tend to 0 in long time, at least when $\sigma > 0$. In fact, some residual fluctuation persists due to the presence of the idiosyncratic noise and this is one of our main achievement here to explain how to handle this residual fluctuation in long time. This is precisely the point where we need σ to be small and the interaction force F to be sufficiently decreasing. As for the function G, it mainly plays a role in the long time analysis of the conditional mean $\mathbb{E}[X_t|B^0]$ itself, which is shown to share many similarities with the long time analysis of the (standard) SDE driven by the drift G and the noise $\sigma_0 B^0$. In particular, we draw heavily on previous works on coupling methods for long time analysis of SDEs, especially the approach developed in [26] on couplings by reflection (inspired from [41]), which plays a key role in our study. Notice that our result complements the earlier one [44] dedicated to the case $\sigma = 0$.

1.4. Further connections with the literature on mean field models. Stochastic PDEs like (1) and related conditional McKean-Vlasov equations of type (2) were introduced in [20, 39]. Further, they were studied in a series of works [42, 43, 32, 30, 33, 28] in connection with stochastic scalar conservations laws, in which case σ_0 is typically required to depend on the local value at point x of dm_t/dx . The more recent contribution [17] addresses uniqueness to (1) under weaker conditions than in [39]. Equation (1) has also become very popular in mean field game theory, see for instance the book [12]. A variant, including a reflection term, has been studied recently in [9]. As for (2), we refer to [34] for a general existence and uniqueness result of weak solutions.

Last but not least, it is worth recalling that the connection between nonlinear Fokker-Planck equations and McKean-Vlasov equations (without common noise) goes back to the pioneering works of [38, 47, 31]. In this context, a key question concerns the particular approximation of the solutions, usually referred to as *propagation of chaos*, see for instance [52, 48]. In connection with the existence of stationary solutions to the Fokker-Planck equation (1), it is in general a difficult question to wonder whether propagation of chaos holds uniformly in time or not. We feel better not to give a list of references in this direction but we quote that the question appears for instance in the earlier contribution [16]. Obviously, this would be very interesting to address the same problem but in presence of a common noise for the model studied here.

1.5. **Organisation of the paper.** In Section 2, we list the assumptions and give the main statements of the paper. We also address some examples and compare in particular our results with those from [35] (when $\sigma_0 = 0$ and uniqueness does not hold). The proof of the main theorem (Theorem 8) is split in two parts. A first step is to derive Theorem 8 from a key auxiliary estimate (Proposition 7) on the contraction properties of the semigroup induced by the solution of (1). This is done in Section 3. A second step, which is in fact the core of the paper, is to prove this key estimate. We do so in Section 4, using coupling arguments. Auxiliary results are proven in Appendix.

1.6. **Notation.** Throughout the paper, for a Polish space E, $\mathcal{P}(E)$ stands for the space of Borel probability measures on E equipped with the topology of weak convergence and the corresponding Borel σ -algebra (which is induced by the mappings $m \in \mathcal{P}(E) \mapsto m(A)$ for any Borel subset A of E). Whenever there exists a distance d so that (E, d) is a metric space, we call $\mathcal{P}_p(E)$, for any p > 0, the collection of elements $\mu \in \mathcal{P}_p(E)$ such that

$$\exists x_0 \in E : \int_E d(x_0, x)^p \mu(\mathrm{d}x) < +\infty.$$

In fact, the integral above is finite or not, whatever the choice of x_0 . We then define for any $p \ge 1$, the *p*-Wasserstein distance on $\mathcal{P}_p(E)$ as

$$\mathbf{d}_p^E(\mu,\nu) := \inf_{\pi \in \Pi(\mu,\nu)} \left(\int_{E \times E} d(x,y)^p \, \pi(\mathbf{d} x,\mathbf{d} y) \right)^{1/p}, \quad \mu,\nu \in \mathcal{P}_1(E),$$

where $\Pi(\mu, \nu)$ stands for the set of all couplings of μ and ν (i.e., all the joint probability measures on $E \times E$ with μ and ν as first and second marginals).

Also, for any measurable $\varphi : E \to \mathbb{R}$ and any probability measure $m \in \mathcal{P}(E)$, we define the duality product $\langle \cdot ; \cdot \rangle_E$ as

$$\langle m; \varphi \rangle_E = \int_E \varphi(x) m(\mathrm{d}x),$$

whenever the integral in the right-hand side does exit (for instance so is the case if φ is bounded or φ is at most of linear growth and $m \in \mathcal{P}_1(E)$ and so forth...). When $E = \mathbb{R}^d$, we define

$$\mu_1(m) := \int_{\mathbb{R}^d} x \, m(\mathrm{d}x), \quad m \in \mathcal{P}_1(\mathbb{R}^d);$$

$$\mu_2(m) := \int_{\mathbb{R}^d} |x|^2 \, m(\mathrm{d}x), \quad m \in \mathcal{P}_2(\mathbb{R}^d);$$

$$v(m) := \int_{\mathbb{R}^d} |x - \mu_1(m)|^2 \, m(\mathrm{d}x), \quad m \in \mathcal{P}_2(\mathbb{R}^d);$$

which stand respectively for the expectation, the moment of order two and the variance of m. It is important to pay attention to the fact that despite the similarities in the notation, the objects $\mu_1(m) \in \mathbb{R}^d$ and $\mu_2(m) \in \mathbb{R}$ are of different dimensions.

When *m* is random, say is a measurable mapping from $(\Omega_0, \mathcal{F}^0, \mathbb{P}_0)$ into $\mathcal{P}(E)$ (or $\mathcal{P}_1(E)$), its law *P* is an element of $\mathcal{P}(\mathcal{P}(E))$ (or $\mathcal{P}(\mathcal{P}_1(E))$), in the sense that, for any bounded measurable function $\phi : \mathcal{P}(E) \to \mathbb{R}$, $\mathbb{E}_0[\phi(m)] = \langle P; \phi \rangle_{\mathcal{P}(E)}$ (and similarly, when working on $\mathcal{P}_1(E)$), where \mathbb{E}_0 is the expectation associated with \mathbb{P}_0 . Repeatedly in this paper, we also consider continuous stochastic processes $(m_t)_{t\geq 0}$, typically constructed on $(\Omega_0, \mathcal{F}^0, \mathbb{P}_0, \mathbb{P}_0)$, with values in the space $\mathcal{P}_1(\mathbb{R}^d)$ equipped with $d_1^{\mathbb{R}^d}$ (and thus also in $\mathcal{P}(\mathbb{R}^d)$, equipped with the weak convergence topology). When $(m_t)_{t\geq 0}$ is Markovian, we define its semigroup $(\mathscr{P}_t)_{t\geq 0}$ by letting, for any function $\phi : \mathcal{P}(\mathbb{R}^d) \to \mathbb{R}$ and any $t \geq 0$,

$$\mathscr{P}_t \phi: \begin{array}{c} \mathscr{P}(\mathbb{R}^d) \to \mathbb{R} \\ m \mapsto \mathbb{E}_0[\phi(m_t)|m_0 = m] \end{array}$$

Moreover, for any twice differentiable function $f : \mathbb{R}^d \to \mathbb{R}$, ∇f , Δf and $\nabla^2 f$ respectively stand for the gradient, Laplacian and Hessian matrix of f. For any differentiable function $F : \mathbb{R}^d \to \mathbb{R}^d$, we denote by $DF : \mathbb{R}^d \to \mathcal{M}_d(\mathbb{R})$ its Jacobian matrix.

For any real squared matrix $A \in \mathcal{M}_d(\mathbb{R})$, we denote the Euclidian norm of A by

$$||A||| = \sqrt{\operatorname{Tr}(A^{\top}A)},$$

with A^{\top} and Tr respectively standing for the transpose matrix of A and the Trace operator on $\mathcal{M}_d(\mathbb{R})$. Lastly, the Euclidean norm on \mathbb{R}^d is denoted $|\cdot|$ and the inner product \cdot .

2. MAIN RESULT: UNIQUENESS RECOVERY THANKS TO COMMON NOISE

2.1. Stochastic Fokker-Planck PDE and Conditional McKean-Vlasov SDE. In this paper, we focus on a mean field model with common noise in which the nonlinearity in the dynamics occurs through the mean (i.e., the expectation). To make it clear, on a filtered probability space $(\Omega_0, \mathcal{F}^0, \mathbb{P}^0, \mathbb{P}_0)$ (satisfying the usual conditions) equipped with a *d*-dimensional \mathbb{F}^0 -Brownian motion B^0 and for $F, G : \mathbb{R}^d \to \mathbb{R}^d$ two interaction and confinement forces, we are interested in the

solutions of the following Stochastic Partial Differential Equation (SPDE):

(4)
$$d_t m_t = \nabla \cdot \left(\frac{\sigma^2 + \sigma_0^2}{2} \nabla m_t - m_t \left(G + F(\cdot - \mu_1(m_t))\right)\right) dt - \sigma_0 \nabla m_t \cdot dB_t^0, \quad t \ge 0.$$

which is understood the weak sense, i.e., the process $(m_t)_{t \ge 0}$ satisfies for all $t \ge 0$ and $\varphi \in C_c^{\infty}(\mathbb{R}^d)$,

$$\mathrm{d}\langle m_t, \varphi \rangle = \langle m_t, \mathscr{L}_{m_t} \varphi \rangle \,\mathrm{d}t + \sigma_0 \langle m_t, (\nabla \varphi)^\top \rangle \,\mathrm{d}B_t^0,$$

where for any probability measure $m \in \mathcal{P}_1(\mathbb{R}^d)$, the operator \mathscr{L}_m acts on a smooth function φ with compact support in the following manner:

$$\mathscr{L}_m \varphi = -\left(G + F(\cdot - \mu_1(m)) \cdot \nabla \varphi + \frac{\sigma_0^2 + \sigma^2}{2} \Delta \varphi\right).$$

Equation (4) admits a Lagrangian representation in the form of conditional McKean-Vlasov equation. Throughout the article, this McKean-Vlasov equation is constructed on a product probability space, obtained by tensorizing the filtered probability space $(\Omega_0, \mathcal{F}^0, \mathbb{P}^0, \mathbb{P}_0)$ (on which Equation (4) is defined) with another filtered probability space $(\Omega_1, \mathcal{F}^1, \mathbb{P}^1, \mathbb{P}_1)$. The product structure is denoted

$$(\Omega := \Omega_0 \times \Omega_1, \ \mathcal{F}, \ \mathbb{F}, \ \mathbb{P}),$$

where $(\mathcal{F}, \mathbb{P})$ is the completion of $(\mathcal{F}^0 \otimes \mathcal{F}^1, \mathbb{P}_0 \otimes \mathbb{P}_1)$ and \mathbb{F} is the right continuous augmentation of $(\mathcal{F}^0_t \otimes \mathcal{F}^1_t)_{t \ge 0}$. In this paper we denote by \mathbb{E}_0 , \mathbb{E}_1 and \mathbb{E} the expectations on $(\Omega_0, \mathcal{F}^0, \mathbb{P}^0, \mathbb{P}_0)$, $(\Omega_1, \mathcal{F}^1, \mathbb{F}^1, \mathbb{P}_1)$ and $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ respectively. Analogously, for any random variable *X* defined on $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, we denote by $\mathcal{L}_0(X)$ and $\mathcal{L}_1(X)$ the conditional laws of the random variable *X* given \mathcal{F}^1 and \mathcal{F}^0 respectively. Of course, $\mathcal{L}(X)$ stands for the law of *X* on the whole product space.

Next, the initial law of (4) may be random and then distributed according to a probability measure $P_0 \in \mathcal{P}(\mathcal{P}_1(\mathbb{R}^d))$. The initial condition m_0 is then defined as an \mathcal{P}_0^0 -measurable random variable with values in $\mathcal{P}_1(\mathbb{R}^d)$ such that $\mathcal{L}(m_0) = P_0$. We can now define on the whole probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ a random variable X_0 such that $\mathcal{L}_1(X_0) = m_0$ almost surely. Precisely, by Lemma 2.4 in [14], for \mathbb{P}^0 -a.e. $\omega_0 \in \Omega_0$, $X_0(\omega_0, \cdot)$ is a random variable on $(\Omega_1, \mathcal{F}^1, \mathbb{P}_1)$ and the conditional law of X_0 given \mathcal{F}^0 , $\mathcal{L}_1 : \Omega_0 \ni \omega_0 \mapsto \mathcal{L}(X(\omega_0, \cdot))$, defines a random variable from $(\Omega_0, \mathcal{F}^0, \mathbb{P}_0)$ into $\mathcal{P}(\mathbb{R}^d)$, which can be taken equal to m_0 almost surely.

In addition to B^0 , we consider a *d*-dimensional \mathbb{F}^1 -Brownian motion B supported by the space $(\Omega_1, \mathcal{F}^1, \mathbb{P}_1)$. The Lagrangian formulation of (4) then reads in the form of the following conditional McKean-Vlasov equation, set on $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$:

(5)
$$dX_t = G(X_t)dt + F(X_t - \mathbb{E}_1[X_t])dt + \sigma dB_t + \sigma_0 dB_t^0, \quad t \ge 0,$$

with X_0 as initial condition.

2.2. Assumptions, existence and uniqueness of solutions. In the sequel, we make the following assumptions on the coefficients *F* and *G*. We start with *G*:

Assumption 1. The drift $G : \mathbb{R}^d \to \mathbb{R}^d$ is differentiable in \mathbb{R}^d . Moreover, (A1.1) *G* is confining in the sense that there exists a function $\kappa : [0; +\infty) \to \mathbb{R}$ such that

$$\forall x, y \in \mathbb{R}^d, \quad (G(x) - G(y)) \cdot (x - y) \leq -\frac{\sigma_0^2}{2} \kappa(|x - y|) |x - y|^2$$

with

$$\lim \sup_{r \to +\infty} \kappa(r) > 0 \quad \text{and} \quad \int_0^1 r \kappa(r)^- \mathrm{d} r < \infty,$$

where $\kappa^- = \max(0, -\kappa)$.

(A1.2) *G* is Lipschitz-continuous on \mathbb{R}^d , meaning that there exists a constant $L_G > 0$ such that

$$\forall x, y \in \mathbb{R}^d, \quad |G(x) - G(y)| \le L_G |x - y|.$$

(A1.3) *G* is differentiable and its Jacobian matrix *DG* is Lipschitz-continuous on \mathbb{R}^d , meaning that there exists a constant $C_G > 0$ such that for any $x, y \in \mathbb{R}^d$,

$$||DG(x) - DG(y)|| \le C_G |x - y|.$$

In order to state properly the assumption on *F*, we introduce the following definition:

Definition 1. For a pair $(\alpha, C) \in \mathbb{R} \times [0, +\infty)$, we call $\mathcal{S}(\alpha, C, L)$ the collection of L-Lipschitz-continuous and differentiable functions H from \mathbb{R}^d into itself that are α -decreasing and whose derivative is C Lipschitz-continuous, i.e., for all $(x, y) \in \mathbb{R}^{2d}$,

- (i) $(x-y) \cdot (H(x) H(y)) \leq -\alpha |x-y|^2$;
- $(ii) \quad \|DH(x) DH(y)\| \leq C|x y|.$

Given Definition 1, we make the following assumption on *F*:

Assumption 2. $F : \mathbb{R}^d \to \mathbb{R}^d$ satisfies F(0) = 0. Moreover, there exist $\alpha_F > 0$ and $C_F \ge 0$ such that $F \in S(\alpha_F, C_F, C_F)$.

About Assumption **1***.* The following remarks about Assumption **1** are in order:

- Assumption 1 is quite intuitive and consistent with the ideas presented in [35]. Essentially, this assumption guarantees that *G* is confining and behaves linearly at infinity. Also, it is worth noting that our approach can accommodate a force that derives from a non-convex potential, which, around the origin, behaves for example like $x \mapsto |x|^4 |x|^2$.
- When $\sigma_0 > 0$, there exists a canonical choice of κ , given by

$$\kappa(r) := \inf \left\{ -\frac{2}{\sigma_0^2} \frac{(x-y) \cdot (G(x) - G(y))}{|x-y|^2}; \quad x, y \in \mathbb{R}^d \text{ s.t. } |x-y| = r \right\}, \quad r > 0.$$

Assumption 1 says that κ is necessarily positive outside of a ball. This implies in particular that $\sigma_0^2 \kappa^-/2$ is bounded from below by some constant $m_G \in \mathbb{R}$, which gives

$$(x-y)\cdot (G(x)-G(y)) \leqslant -m_G |x-y|^2, \quad x,y \in \mathbb{R}^d.$$

This implies in particular that $G \in \mathcal{S}(m_G, C_G, L_G)$.

When $m_G > 0$ we say that the drift is (strictly) decreasing. This case is easier to study under similar conditions on F because the monotone structure of G ensures that two solutions of the SDE (5) (with different initial conditions) get closer in large time. In this context, uniqueness of the invariant measure of the solutions of Equation (4) is covered in [44, Section 2], but it is fair to say that the common noise then plays little role because the invariant measure is also unique when $\sigma_0 = 0$. Notice that in the specific case when Gderives from a potential, this potential is strictly convex under the condition $m_G > 0$.

8

(6)

About Assumption 2. Assumption 2 is somewhat surprising, especially as we are asking below the coefficient α_F to be large (see the statement of Theorem 8). Intuitively, one might indeed expect that uniqueness of the invariant measure becomes less likely if the interaction is strong. At least, this is exactly what happens in absence of common noise, the extreme case being $F \equiv 0$ in which the dynamics become a mere diffusion equation (the long time analysis of which is much easier to study). However, the situation is different here, thanks to the peculiar structure of the dynamics and to the presence of the common noise ($\sigma_0 > 0$). In brief, F has a contracting effect, which forces the process to be attracted by its conditional expectation (given the common noise), with the latter being a nearly solution of an ergodic SDE. Even though this picture is not exactly correct due to some minor errors caused by the idiosyncratic noise, this is indeed a key step in our analysis to quantify the accuracy of this approximation and to derive from it uniqueness of the invariant measure. We refer to Section 3.1.

Throughout, we work under Assumptions 1 and 2. In this context, we claim

Proposition 2. Under Assumptions 1 and 2 and for an initial condition X_0 satisfying $\mathbb{E}[|X_0|] < \infty$ (or equivalently $\mathbb{E}_0 \int_{\mathbb{R}^d} |x| m_0(dx) < \infty$ for $m_0 := \mathcal{L}_0(X_0)$), the conditional McKean-Vlasov equation (5) has a unique \mathbb{F} -progressively measurable solution $(X_t)_{t \ge 0}$, with continuous trajectories, such that, for all T > 0, $\mathbb{E}[\sup_{0 \le t \le T} |X_t|] < \infty$.

Proof. The proof consists in a straightforward fixed point argument, using the Lipschitz properties of the two coefficients F and G. We refer to [14, Chap. 2].

By the superposition principle for conditional McKean-Vlasov equations (see [40]), we deduce that existence and uniqueness also hold true for the stochastic Fokker-Planck equation:

Proposition 3. Under Assumptions 1 and 2 and for an initial condition m_0 satisfying $\mathbb{E}_0 \int_{\mathbb{R}^d} |x| m_0(dx) < \infty$, the stochastic Fokker-Planck equation (4) has a unique solution $(m_t)_{t\geq 0}$ in the space of \mathbb{F}^0 -progressively measurable processes with values in $\mathcal{P}_1(\mathbb{R}^d)$ satisfying $\mathbb{E}_0[\sup_{0\leq t\leq T}\int_{\mathbb{R}^d} |x| m_t(dx)] < \infty$.

Proof. Existence of a solution is a straightforward consequence of the existence part in the statement of Proposition 2. This is Proposition 1.2 in [40]. Uniqueness is more difficult to obtain. We invoke Theorem 1.3 in [40]. In brief, any solution $(\tilde{m}_t)_{t\geq 0}$ (in the weak sense) to (4) induces a weak solution to the McKean-Vlasov equation $\tilde{X} = (\tilde{X}_t)_{t\geq 0}$ to (5), weak in the sense that the private (or idiosyncratic) noise, say \tilde{B} , becomes part of the solution. It holds $\tilde{m}_t = \mathcal{L}(\tilde{X}_t | \mathcal{F}_T^0)$. In fact, by strong uniqueness to (5), a relevant form of Yamada-Watanabe theorem applies to the current setting and implies that $\mathcal{L}(\tilde{X}_t | \mathcal{F}_T^0)$ is necessarily equal to $\mathcal{L}_0(X_t)$, with $X = (X_t)_{t\geq 0}$ being the solution given by Proposition 2.

2.3. **Restoration of uniqueness.** We now collect our main results about the long time behaviour of the process $(m_t)_{t \ge 0}$, solution of Equation (4). Before we do so, we first state the precise definition of an invariant measure for such a stochastic process, when seen as a process with values in the space of probability measures:

Definition 4 (Invariant measure). We say that $\overline{P} \in \mathcal{P}_2(\mathcal{P}_1(\mathbb{R}^d))$ is an invariant measure for the process $(m_t)_{t\geq 0}$ when the latter is regarded as taking values in $\mathcal{P}_1(\mathbb{R}^d)$, if the law of m_t is independent of t, when

 m_0 is distributed according to \overline{P} , i.e., for all continuous and bounded function $\phi \in \mathcal{C}_b(\mathcal{P}_1(\mathbb{R}^d))$

$$\mathbb{E}_0[\phi(m_t)] = \mathbb{E}_0[\phi(m_0)] = \int_{\mathcal{P}_1(\mathbb{R}^d)} \phi(m)\bar{P}(\mathrm{d}m), \quad \forall t > 0$$

Of course, if \overline{P} is an invariant probability distribution, then the process $(m_t)_{t\geq 0}$ with initial condition \overline{P} , has the same law as the process $(m_{t+T})_{t\geq 0}$ for any T > 0. This follows directly from the fact that m_0 and m_T have the same law combined with the weak Markov property. Moreover, it is worth observing that, under the condition $\overline{P} \in \mathcal{P}_2(\mathcal{P}_1(\mathbb{R}^d))$ imposed in the statement, the function $m \mapsto d_1^{\mathbb{R}^d}(\delta_0, m) = \int_{\mathbb{R}^d} |x| m(dx)$ is square integrable. This constraint may seem rather artificial. In fact, given the form of the dynamics in Equation (4), it is natural to require the invariant measures to be supported by $\mathcal{P}_1(\mathbb{R}^d)$. Then, the aforementioned integrability condition provides an additional growth property that makes easier the identification of those invariant measures, see the proof of Proposition 5 right below.

Let us begin with the existence of an invariant measure P:

Proposition 5. Under Assumptions 1 and 2, there exists at least one invariant measure \overline{P} for the process defined by Equation (4). Moreover, under the condition $\alpha > L_G$, any invariant measure belongs to $\mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$.

The proof of Proposition 5 makes use of the notion of derivatives on the space of measures. Following [11] (see also [13, Chap. 5] and [14, Chap. 4]), we recall the following definition:

Definition 6. Let us define $C_b^2(\mathcal{P}_2(\mathbb{R}^d))$ as the collection of continuous and bounded functions Φ : $\mathcal{P}_2(\mathbb{R}^d) \to \mathbb{R}$ with the following properties (throughout the definition, $\mathcal{P}_2(\mathbb{R}^d)$ is equipped with $d_2^{\mathbb{R}^d}$):

• There exists a unique jointly continuous function $\partial_m \Phi : (m, x) \in \mathcal{P}(\mathbb{R}^d) \times \mathbb{R}^d \to \partial_m \Phi(m, x) \in \mathbb{R}^d$, at most of quadratic growth in x for any m, such that

$$\lim_{h \to 0} \frac{\Phi(m + h(m' - m)) - \Phi(m)}{h} = \int_{\mathbb{R}^d} \partial_m \Phi(m, v)(m' - m)(\mathrm{d}v),$$

for all $m, m' \in \mathcal{P}(\mathbb{R}^d)$ and

$$\int_{\mathbb{R}^d} \partial_m \Phi(m, v) m(\mathrm{d}v) = 0, \quad m \in \mathcal{P}(\mathbb{R}^d);$$

- For any $m \in \mathcal{P}_2(\mathbb{R}^d)$, the mapping $x \mapsto \partial_m \Phi(m, x)$ is differentiable, with the gradient being denoted $D_m \Phi(m, x)$; the mapping $(m, x) \mapsto D_m \Phi(m, x)$ is jointly continuous in (m, x) and at most of linear growth in x uniformly in m in bounded subsets of $\mathcal{P}_2(\mathbb{R}^d)$;
- For any $x \in \mathbb{R}^d$, every component of the \mathbb{R}^d -valued function $m \mapsto D_m \Phi(m, x)$ satisfies the same conditions as in the first bullet point, resulting in a mapping $(m, x, y) \in \mathcal{P}_2(\mathbb{R}^d) \times \mathbb{R}^d \times \mathbb{R}^d \mapsto D_m^2 \Phi(m, x, y) \in \mathbb{R}^{d \times d}$, which is jointly continuous (in the three arguments) and at most of quadratic growth in (x, y), uniformly in m in bounded subsets of $\mathcal{P}_2(\mathbb{R}^d)$;
- For any m ∈ P₂(ℝ^d), the function x ∈ ℝ^d → D_mΦ(m, x) is differentiable; the Jacobian, denoted (m, x) → D²_{xm}Φ(m, x), is jointly continuous in (m, x) and at most of linear growth in x, uniformly in m in bounded subsets of P₂(ℝ^d).

Proof of Proposition 5. Step 1. The proof of the existence of an invariant measure closely follows the approach taken in Proposition 2 of [44], using the results of Section 1.2 of [40]. The key step

(4), which can be achieved with relative ease in our context. Details are left to the reader.

Step 2. We now address the second part of the statement (any invariant measure in $\mathcal{P}_2(\mathcal{P}_1(\mathbb{R}^d))$) is in fact in $\mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$). Let us thus consider an invariant measure $\bar{P} \in \mathcal{P}_2(\mathcal{P}_1(\mathbb{R}^d))$ and call m_0 an \mathcal{F}_0 -valued random variable with values in $\mathcal{P}_1(\mathbb{R}^d)$ such that $\mathcal{L}_0(m_0) = \bar{P}$. We denote by $(m_t)_{t \ge 0}$ the solution to Equation (4) with m_0 as initial condition. We also introduce for any R > 0, a smooth function $\chi_R^d : \mathbb{R}^d \to \mathbb{R}^d$ that coincides with the identity on the ball $B_{\mathbb{R}^d}(0, R)$ and that is equal to 0 outside $B_{\mathbb{R}^d}(0, 2R)$. We can assume the Lipschitz constant of χ_R^d to be bounded by 1. For any R > 0, we denote by P_0^R the law of $\chi_R^d \# m_0$ (where # is the pushforward operator) and construct $(m_t^R)_{t\ge 0}$, the solution to Equation (4) with $\chi_R^d \# m_0$ as initial condition. Letting $(P_t^R)_{t\ge 0} := (\mathcal{L}_0(m_t^R))_{t\ge 0}$, we deduce from the truncation performed on the initial condition and from Equation (4) that, for any R > 0 and any $t \ge 0$, $P_t^R(\mathcal{P}_2(\mathbb{R}^d)) = 1$. We obtain from the chain rule proven in [13, Chap. 5] and [14, Chap. 4] (see [40] for the form that is retained below) that for any function $\Phi \in \mathcal{C}_b^2(\mathcal{P}_2(\mathbb{R}^d))$, for any $t \ge 0$,

(7)
$$\langle P_t^R - P_0^R; \Phi \rangle = \int_0^t \langle P_s^R; \mathcal{M}\Phi \rangle \mathrm{d}s$$

where for any $m \in \mathcal{P}_2(\mathbb{R}^d)$,

$$\begin{split} \mathcal{M}\Phi(m) &:= \int_{\mathbb{R}^d} \left[D_m \Phi(m,x) \cdot (G(x) + F(x - \mu_1(m)) + \frac{\sigma^2 + \sigma_0^2}{2} \operatorname{Tr} \left[D_{xm}^2 \Phi(m,x) \right] \right] m(\mathrm{d}x) \\ &+ \frac{\sigma_0^2}{2} \int_{\mathbb{R}^{2d}} \operatorname{Tr} \left[D_{mm}^2 \Phi(m,x,y) \right] m(\mathrm{d}x) m(\mathrm{d}y). \end{split}$$

Let us now consider $\Phi(m) := \int_{\mathbb{R}^d} \phi(x - \mu_1(m))m(dx)$, for a function $\phi : \mathbb{R}^d \to \mathbb{R}$ which we assume to be bounded and smooth, with bounded derivatives of order 1 and 2. In particular, Φ is bounded. Now, $m \mapsto \Phi(m) \in \mathcal{C}^2_b(\mathcal{P}_2(\mathbb{R}^d))$ and we can write, for any $m \in \mathcal{P}_2(\mathbb{R}^d)$ and $x, y \in \mathbb{R}^d$,

$$\begin{split} D_m \Phi(m,x) &= \nabla \phi(x-\mu_1(m)) - \int_{\mathbb{R}^d} \nabla \phi(y-\mu_1(m)) m(\mathrm{d}y), \\ D_{xm}^2 \Phi(m,x) &= \nabla^2 \phi(x-\mu_1(m)), \\ D_{mm}^2 \Phi(m,x,y) &= -\nabla^2 \phi(x-\mu_1(m)) - \nabla^2 \phi(y-\mu_1(m)) + \int_{\mathbb{R}^d} \nabla^2 \phi(z-\mu_1(m)) m(\mathrm{d}z). \end{split}$$

Plugging this into Equation (7), we obtain

$$\begin{split} \int_{\mathcal{P}_{2}(\mathbb{R}^{d})} \Phi(m)(P_{t}^{R} - P_{0}^{R})(\mathrm{d}m) \\ &= \int_{0}^{t} \left[\int_{\mathcal{P}_{2}(\mathbb{R}^{d})} \int_{\mathbb{R}^{d}} \nabla \phi(x - \mu_{1}(m)) \cdot (G(x) + F(x - \mu_{1}(m)))m(\mathrm{d}x)P_{s}^{R}(\mathrm{d}m) \\ &- \int_{\mathcal{P}_{2}(\mathbb{R}^{d})} \left(\int_{\mathbb{R}^{d}} \nabla \phi(y - \mu_{1}(m))m(\mathrm{d}y) \right) \cdot \left(\int_{\mathbb{R}^{d}} (G(x) + F(x - \mu_{1}(m)))m(\mathrm{d}x) \right) P_{s}^{R}(\mathrm{d}m) \\ &+ \frac{1}{2} \left(\sigma^{2} + \sigma_{0}^{2} \right) \int_{\mathbb{R}^{d}} \mathrm{Tr} \left[\nabla^{2} \phi(z - \mu_{1}(m)) \right] m(\mathrm{d}z) - \frac{1}{2} \sigma_{0}^{2} \int_{\mathbb{R}^{d}} \mathrm{Tr} \left[\nabla^{2} \phi(z - \mu_{1}(m)) \right] m(\mathrm{d}z) \right] \mathrm{d}s, \end{split}$$

which gives (by simplifying the last line)

$$\int_{\mathcal{P}_{2}(\mathbb{R}^{d})} \Phi(m)(P_{t}^{R} - P_{0}^{R})(\mathrm{d}m)$$

$$= \int_{0}^{t} \left[\int_{\mathcal{P}_{2}(\mathbb{R}^{d})} \int_{\mathbb{R}^{d}} \nabla \phi(x - \mu_{1}(m)) \cdot (G(x) + F(x - \mu_{1}(m)))m(\mathrm{d}x)P_{s}^{R}(\mathrm{d}m) - \int_{\mathcal{P}_{2}(\mathbb{R}^{d})} \left(\int_{\mathbb{R}^{d}} \nabla \phi(y - \mu_{1}(m))m(\mathrm{d}y) \right) \cdot \left(\int_{\mathbb{R}^{d}} (G(x) + F(x - \mu_{1}(m)))m(\mathrm{d}x) \right) P_{s}^{R}(\mathrm{d}m)$$

$$+ \frac{1}{2}\sigma^{2} \int_{\mathbb{R}^{d}} \operatorname{Tr} \left[\nabla^{2} \phi(z - \mu_{1}(m)) \right]m(\mathrm{d}z) \right] \mathrm{d}s.$$

Step 3. We now want to let $R \to \infty$. Let us begin with the term on the left-hand side

$$\int_{\mathcal{P}(\mathbb{R}^d)} \Phi(m)(P_t^R - P_0^R)(\mathrm{d}m) = \int_{\mathcal{P}(\mathbb{R}^d)} \Phi(m)(P_t^R - \bar{P})(\mathrm{d}m) + \int_{\mathcal{P}(\mathbb{R}^d)} \Phi(m)(\bar{P} - P_0^R)(\mathrm{d}m).$$

Using the Lipschitz-continuity of *F* and *G*, we can show that, for any $T \ge 0$, there exists a constant $C_T \ge 0$, such that, for any $t \in [0, T]$,

(9)
$$d_2^{\mathcal{P}_1(\mathbb{R}^d)}(P_t^R,\bar{P}) \leqslant C_T d_2^{\mathcal{P}_1(\mathbb{R}^d)}(P_0^R,\bar{P}).$$

Moreover, we can write

$$\mathbf{d}_{2}^{\mathcal{P}_{1}(\mathbb{R}^{d})}(P_{0}^{R},\bar{P})^{2} \leqslant \int_{\mathcal{P}(\mathbb{R}^{d})} \left(\int_{\mathbb{R}^{d}} |\chi_{R}^{d}(x) - x| m(\mathrm{d}x) \right)^{2} \bar{P}(\mathrm{d}m).$$

Since χ_R^d is equal to 0 in 0 and is 1-Lipschitz continuous and since $\overline{P} \in \mathcal{P}_2(\mathcal{P}_1(\mathbb{R}^d))$, the above right-hand side tends to 0 as R tends to ∞ , that is

(10)
$$\lim_{R \to +\infty} \mathrm{d}_2^{\mathcal{P}_1(\mathbb{R}^d)}(P_0^R, \bar{P}) = 0.$$

Equations (9) and (10) say that for all $T \ge 0$,

(11)
$$\lim_{R \to +\infty} \sup_{0 \le t \le T} \mathrm{d}_2^{\mathcal{P}_1(\mathbb{R}^d)}(P_t^R, \bar{P}) = 0$$

Back to (8), we observe that all the functions of m that are integrated with respect to \overline{P} therein are continuous in m with respect to $d_1^{\mathbb{R}^d}$. Moreover, they are all at most of linear growth with respect to $\int_{\mathbb{R}^d} |x| m(\mathrm{d}x)$. This makes it possible to let R tend to $+\infty$. We get

$$\begin{split} &\int_{\mathcal{P}_1(\mathbb{R}^d)} \int_{\mathbb{R}^d} \nabla \phi(x - \mu_1(m)) \cdot (G(x) + F(x - \mu_1(m))) m(\mathrm{d}x) \bar{P}(\mathrm{d}m) \\ &- \int_{\mathcal{P}_1(\mathbb{R}^d)} \left(\int_{\mathbb{R}^d} \nabla \phi(y - \mu_1(m)) m(\mathrm{d}y) \right) \cdot \left(\int_{\mathbb{R}^d} (G(x) + F(x - \mu_1(m))) m(\mathrm{d}x) \right) \bar{P}(\mathrm{d}m) \\ &+ \frac{1}{2} \sigma^2 \int_{\mathcal{P}_1(\mathbb{R}^d)} \int_{\mathbb{R}^d} \mathrm{Tr} \big[\nabla^2 \phi(z - \mu_1(m)) \big] m(\mathrm{d}z) \bar{P}(\mathrm{d}m) = 0. \end{split}$$

Step 4. Now, we take $\phi(x)$ in the form $\vartheta(|x|)$ where ϑ is a smooth function from $[0, +\infty)$ into itself, that is equal to the identity on [0, A] and that is equal to 3A/2 on $[2A, +\infty)$, for some A > 0. We assume ϑ' to be non-negative and bounded by the identity and ϑ'' to take values in [0, 2].

Under this choice, $\nabla \phi(x)$ rewrites $[\vartheta'(|x|)/|x|]x$ and $\operatorname{Tr}[\nabla^2 \phi(x)]$ is bounded by a constant c_d only depending on the dimension d. Then, we can find a constant C, only depending on G (and whose value is allowed to change from line to line) such that, for any $m \in \mathcal{P}_2(\mathbb{R}^d)$,

$$\begin{split} & \left| \int_{\mathbb{R}^{d}} \nabla \phi(x - \mu_{1}(m)) \cdot G(x)m(\mathrm{d}x) \right| \\ & \leq \left| \int_{\mathbb{R}^{d}} \nabla \phi(x - \mu_{1}(m)) \cdot \left[G(x) - G(\mu_{1}(m)) \right] m(\mathrm{d}x) \right| + \left| \int_{\mathbb{R}^{d}} \nabla \phi(x - \mu_{1}(m)) \cdot G(\mu_{1}(m))m(\mathrm{d}x) \right| \\ & \leq L_{G} \int_{\mathbb{R}^{d}} \vartheta'(|x - \mu_{1}(m)|)|x - \mu_{1}(m)|m(\mathrm{d}x) + C \int_{\mathbb{R}^{d}} \vartheta'(|x - \mu_{1}(m)|) (1 + |\mu_{1}(m)|)m(\mathrm{d}x) \\ & \leq L_{G} \int_{\mathbb{R}^{d}} \vartheta'(|x - \mu_{1}(m)|)|x - \mu_{1}(m)|m(\mathrm{d}x) + C \left(1 + \mathrm{d}_{1}^{\mathbb{R}^{d}} (\delta_{0}, m)^{2} \right). \end{split}$$

We insist on the fact that C is independent of ϑ . Now using the assumption on F (see Assumption 2), we obtain

$$\int_{\mathbb{R}^d} \nabla \phi(x - \mu_1(m)) \cdot F(x - \mu_1(m)) m(\mathrm{d}x) \leq -\alpha \int_{\mathbb{R}^d} \vartheta'(|x - \mu_1(m)|) |x - \mu_1(m)| m(\mathrm{d}x).$$

Rewriting the conclusion of the third step gives

$$(\alpha - L_G) \int_{\mathcal{P}_1(\mathbb{R}^d)} \int_{\mathbb{R}^d} \vartheta'(|x - \mu_1(m)|) |x - \mu_1(m)| m(\mathrm{d}x) \bar{P}(\mathrm{d}m)$$

$$\leqslant c_d \sigma^2 + C \left(1 + \int_{\mathcal{P}_1(\mathbb{R}^d)} \mathrm{d}_1^{\mathbb{R}^d} (\delta_0, m)^2 \bar{P}(\mathrm{d}m) \right).$$

Now, we can choose ϑ' along a non-decreasing sequence converging pointwise to the identity function on $[0, +\infty)$. By monotone convergence, we obtain

$$(\alpha - L_G) \int_{\mathcal{P}_1(\mathbb{R}^d)} \int_{\mathbb{R}^d} |x - \mu_1(m)|^2 m(\mathrm{d}x) \bar{P}(\mathrm{d}m) \leqslant c_d \sigma^2 + C \left(1 + \int_{\mathcal{P}_1(\mathbb{R}^d)} \mathrm{d}_1^{\mathbb{R}^d} (\delta_0, m)^2 \bar{P}(\mathrm{d}m) \right),$$
 nich completes the proof. \Box

which completes the proof.

(12)

Uniqueness is much more challenging to establish. When the intensity of the idiosyncratic noise σ is positive, this question has not been addressed yet. The following proposition is one key step in this regard and demonstrates that, under certain assumptions, the initial condition is forgotten in long time.

Proposition 7. For any $\alpha_F > \max(L_G, m_G)$ in Assumption 2, for any $\sigma_0 > 0$, there exists $\bar{\sigma} > 0$, depending on F, G, σ_0 and d, such that for all $\sigma \leq \overline{\sigma}$, for any 1-Lipschitz continuous function $\phi : \mathfrak{P}_1(\mathbb{R}^d) \to \mathfrak{P}_1(\mathbb{R}^d)$ \mathbb{R} (with $\mathcal{P}_1(\mathbb{R}^d)$ being equipped with $d_1^{\mathbb{R}^d}$) and any $m, \widetilde{m} \in \mathcal{P}_2(\mathbb{R}^d)$,

$$|\mathscr{P}_t\phi(m) - \mathscr{P}_t\phi(\widetilde{m})| \leq \mathfrak{K}(m,\widetilde{m})e^{-\mathfrak{c}t}, \quad t \ge 0,$$

for some $\mathfrak{c} := c(G, F, \sigma, \sigma_0) > 0$ and some $\mathcal{K}(m, \widetilde{m})$, which is described next.

Indeed, given the choice of α_F , there exist a constant $c_1^{\alpha_F} > 0$, depending on G and F, and a constant C_d , depending on d (and thus independent of ϕ , m and \widetilde{m}), such that

$$\mathcal{K}(m,\widetilde{m}) := C_d(\mu_2(m)^{1/2} + \mu_2(\widetilde{m})^{1/2}) \exp\left(\frac{c_1^{\alpha_F}}{c_{\alpha_F}}\left(v(m)^{1/2} + v(\widetilde{m})^{1/2}\right)\right),$$

(0)-

where $c_{\alpha_F} := \alpha_F - L_G$. We recall that μ_1, μ_2 and v are defined in Section 1.6.

Proposition 7 shows that we forget the initial condition at an exponential rate for any sufficiently integrable initial conditions m and \tilde{m} . This is a key result of the paper, the proof of which is postponed to Section 4.

As a consequence of Proposition 7, we get uniqueness of the invariant measure and deduce that the system converges exponentially fast toward the stationary regime, both facts being stated in the main statement below.

Theorem 8. For any $\alpha_F > \max(L_G, m_G)$ in Assumption 2, for any $\sigma_0 > 0$, there exists $\overline{\sigma} > 0$, depending on F, G, σ_0 and d, such that for all $\sigma \leq \overline{\sigma}$, there is a unique invariant measure $\overline{P} \in \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$ for the process $(m_t)_{t\geq 0}$, solution of Equation (4). More precisely, there exists a constant C > 0, such that for any probability measure $m \in \mathcal{P}(\mathbb{R}^d)$ and any 1-Lipschitz continuous function $\phi : \mathcal{P}_1(\mathbb{R}^d) \to \mathbb{R}$, (with $\mathcal{P}_1(\mathbb{R}^d)$ being equipped with $d_1^{\mathbb{R}^d}$), we have

$$\left|\mathscr{P}_t\phi(m) - \int_{\mathscr{P}(\mathbb{R}^d)} \mathscr{P}_t\phi(\widetilde{m})\overline{P}(\mathrm{d}\widetilde{m})\right| \leq C\left(1 + \mu_2(m)^{1/2}\right)\exp(\lambda v(m)^{1/2})e^{-\mathfrak{c}t}, \quad t \ge 0,$$

with $\lambda := c_1^{\alpha_F}/c_{\alpha_F} > 0$, and for c, c_{α_F} and $c_1^{\alpha_F}$ as in the statement of Proposition 7.

Remark 9. Let us comment here on the choice of parameters in the above statement. First, it is important to note that the type and strength of the interaction play a central role in our framework. The interaction, in both its form and intensity ($\alpha_F \ge \overline{\alpha}$), forces the long-time behaviour of the entire process $(m_t)_{t\ge0}$ to be dictated by its mean $(\mu_1(m_t))_{t\ge0}$, which is the cornerstone of the proof. Asking the interaction to be strong, as we require in the condition $\alpha_F \ge \overline{\alpha}$, may seem surprising at first sight. Indeed, this is fundamentally different from the standard picture that exists for models without common noise, where strong interactions are typically expected to give rise to several invariant measures. In our case, the presence of the common noise, even of a possibly small intensity, induces a phase transition. For any positive value of σ_0 and for a sufficiently large interaction (independently of the value of σ_0), we can indeed find $\sigma > 0$ such that uniqueness of the invariant measure is ensured in presence of a common noise of intensity σ_0 but is lost when the common noise is disabled. This makes a conceptual difference with the previous results obtained on the subject in [36], where uniqueness is restored when the global contribution of the two noises is large enough with respect to the strength of the interaction. In comparison, σ and σ_0 have opposite roles to each other in our approach.

2.4. **Discussions and examples.** In this section, we present two explicit examples in which common noise forces uniqueness of the invariant measure. The first example fits explicitly the framework of this work and the second one may be seen as a variant of (4). At last, we also provide a counter-example where uniqueness does not hold despite the presence of the common noise.

A prototype: dynamics driven by a confining potential and a linear interaction. Let us consider the case $G(x) = -\nabla V(x), x \in \mathbb{R}^d$, for a differentiable function $V : \mathbb{R}^d \to \mathbb{R}$, and $F(x) = -\alpha x, x \in \mathbb{R}^d$, for a certain $\alpha > 0$. Then, (4) becomes

(13)
$$d_t m_t = \frac{\sigma^2 + \sigma_0^2}{2} \Delta m_t dt + \nabla \cdot \left[m_t \left(\nabla V + \alpha \left(\cdot - \mu_1(m_t) \right) \right) \right] dt - \sigma_0 \nabla m_t \cdot dB_t^0, \quad t \ge 0.$$

When $\sigma_0 = 0$, this equation reduces to

(14)
$$\partial_t m_t = \frac{\sigma^2}{2} \Delta m_t + \nabla \cdot \left[m_t \left(\nabla V + \alpha \left(\cdot - \mu_1(m_t) \right) \right) \right]$$

When *V* is a double well potential, the latter equation may have several stationary solutions. For instance, if $\sigma = 0$ (in addition to the assumption $\sigma_0 = 0$), any critical point x_0 of *V* induces a stationary solution, concentrated at x_0 . In the standard example where d = 1 and $V(x) = |x|^4/4 - |x|^2/2$ (see for instance [19, 35]), those stationary solutions are the Dirac masses δ_1 , δ_0 and δ_{-1} . Non-uniqueness persists when σ is strictly positive but small: with the same potential *V* and for σ less than a certain threshold $\bar{\sigma}$, there are three invariant measures, respectively centered around 1, 0 and -1, to the McKean-Vlasov SDE

$$dX_t = -V'(X_t)dt - \alpha(X_t - \mathbb{E}_1[X_t])dt + \sigma dB_t, \quad t \ge 0.$$

Lack of uniqueness can be explained as follows. There is a competition between the noise *B* (weighted by the intensity factor σ) and the interaction term function $(x, m) \mapsto -\alpha(x - \mu_1(m))$. On the one hand, the process *X* is attracted by the minimizers of *V* and by its expectation. On the other hand, it is subjected to the diffusive effect of the noise. Whenever the coefficient σ is too small, the interaction term dominates. Any invariant measure has a small variance and must be concentrated around a minimizer of *V*, hence forcing the multiplicity of the stationary solutions to (14).

When $\sigma_0 > 0$ but $\sigma = 0$ (and under conditions similar to Assumption 1), whenever α is assumed large enough, the second author has shown in [44] that the process $(m_t)_{t\geq 0}$ admits a unique invariant measure $P_0 \in \mathcal{P}(\mathcal{P}(\mathbb{R}^d))$, and thus established restoration of uniqueness in this situation. In this particular case, the invariant measure is supported by Dirac masses:

$$\bar{P}_0(\mathrm{d}m) = \int_{\mathbb{R}^d} \delta_{\delta_a}(\mathrm{d}m) m^*(\mathrm{d}a),$$

where m^* is solution of the stationary Fokker-Planck equation $(\sigma_0^2/2)\Delta m^* + \nabla \cdot (m^*\nabla V) = 0$. Moreover, as consequence of the contracting properties of *F*, it can be shown that

$$v(m_t) \to 0,$$

when $t \to +\infty$, which implies that, asymptotically (in time), the solution X to (5) coincides with its conditional expectation given the common noise, hence justifying the shape of the measure \bar{P}_0 .

In the case $\sigma_0 > 0$ and σ small but (strictly) positive, our result extends the analysis carried out in [44]. However, the long-time behaviour of $(v(m_t))_{t \ge 0}$ is different and the latter does not tend to 0 as t tends to ∞ . This difference is substantial and makes the proof significantly more challenging in our analysis. The argument is explained in the next two sections. In short, one shows that the long-time behavior of the conditional mean $(\mu_1(m_t))_{t\ge 0}$ coincides with that one of the standard SDE $dY_t = -\nabla V(Y_t)dt + \sigma_0 dB_t^0$, $t \ge 0$ up to an error that is small with σ and $1/\alpha$. Although the fluctuation $(X_t - \mathbb{E}_1(X_t) = X_t - \mu_1(m_t))_{t\ge 0}$ does not vanish in long time, which is the main mathematical difficulty here, the regime $0 < \sigma \ll 1$ shares many conceptual similarities with the case $\sigma = 0$. In particular, the long-time behavior of $(m_t)_{t\ge 0}$ is mostly dictated by the ergodic properties of the standard SDE obtained by putting $F \equiv 0$ and $\sigma = 0$ in (5). This example is typical of our study. When the value of σ is significantly large and α is relatively small, it is straightforward to establish the uniqueness of the invariant measure using coupling arguments on the idiosyncratic noise, as in [25]. Still, it is fair to say that we do not have, at this stage, a good understanding of the behaviour of the model between these two regimes.

A variant: dynamics driven by a quadratic potential and a nonlinear first order mean field term. In this paragraph, we cook up a variant of the model covered by Theorem 8 that does not exactly fit the form postulated in (4) but that can be studied in a similar manner. This example is built up as a perturbation of an Ornstein-Uhlenbeck process. Precisely, we consider the stochastic Fokker-Planck equation

(15)
$$d_t m_t = \frac{\sigma^2 + \sigma_0^2}{2} \Delta m_t dt + \nabla \cdot \left[m_t \left(ax - f(\mu_1(m_t)) \right) \right] dt - \sigma_0 \nabla m_t \cdot dB_t^0, \quad t \ge 0,$$

whose probabilistic counterpart writes

$$\mathrm{d}X_t = -aX_t\mathrm{d}t + f(\mathbb{E}_1[X_t])\mathrm{d}t + \sigma\mathrm{d}B_t + \sigma_0\mathrm{d}B_t^0, \quad t \ge 0,$$

for some a > 0 and $f : \mathbb{R}^d \to \mathbb{R}^d$. Equivalently, the conditional law $(\mathcal{L}_1(X_t))_{t \ge 0}$ of X given the common noise $(\mathcal{L}_1(X_t))_t$ is a solution to (15).

Obviously, the mean field term $f(\mathbb{E}_1[X_t])$ cannot be put in the form $F(X_t - \mathbb{E}_1[X_t])$, from which we see that this example is outside the scope of Theorem 8. That said, it is in fact not that far from the framework addressed in this article and, in particular, it obeys phenomena similar to those underpinning the proof of Theorem 8.

Actually, the long-time analysis of equation (15) is rather straightforward. When $\sigma_0 = 0$, any invariant measure *m* is solution to

$$\frac{\sigma^2}{2}\Delta m - \nabla \cdot \left[m\left(f(\mu_1(m)) - ax\right)\right] = 0, \quad x \in \mathbb{R}^d.$$

By integrating with respect to x, the previous equation leads to the condition

(16)
$$f(\mu_1(m)) = a\mu_1(m),$$

which is in fact sufficient in the following sense: once $\mu_1(m)$ has been found, the entire measure m can be defined as the invariant measure of the Ornstein-Uhlenbeck dynamics $dX_t = -a(X_t - \mu_1(m))dt + \sigma dB_t$, $t \ge 0$. Therefore, Equation (16) says that there exist several invariant measures as soon as f/a admits several fixed points in \mathbb{R}^d .

In presence of a common noise (i.e., $\sigma_0 > 0$), the conditional mean $(\mu_1(m_t))_{t \ge 0}$ solves the SDE

(17)
$$d\mu_1(m_t) = -a\mu_1(m_t)dt + f(\mu_1(m_t))dt + \sigma_0 dB_t^0, \quad t \ge 0,$$

and under appropriate confining conditions on the *effective* drift $x \mapsto -ax + f(x)$ (which may be compatible with the fact that (16) has several fixed points), (17) admits a unique invariant measure, say m^* (on \mathbb{R}^d), which implies that any two invariant measures on $\mathcal{P}_1(\mathbb{R}^d)$ (in the sense of Definition 4) must have the same marginal law by the projection $m \mapsto \mu_1(m)$. As above, the entire invariant measure (on $\mathcal{P}_1(\mathbb{R}^d)$) can be recovered by observing that $d(X_t - \mathbb{E}_1(X_t)) =$ $-a(X_t - \mathbb{E}_1(X_t))dt + \sigma dB_t, t \ge 0$. This shows that the invariant measure is

(18)
$$\bar{P}(\mathrm{d}m) = \int_{\mathbb{R}^d} \delta_{\mathcal{N}_d(\theta, [\sigma^2/(2a)]I_d)}(\mathrm{d}m) m^*(\mathrm{d}\theta),$$

where $\mathcal{N}_d(\theta, [\sigma^2/(2a)]I_d)$ is the *d*-dimensional Gaussian law with θ as mean and $\sigma^2/(2a)I_d$ as covariance (I_d standing for the *d*-dimensional identity matrix). In particular, there is a unique invariant measure even though (16) has several fixed points.

The spirit of this new example is clear: similar to the prototype addressed in the previous paragraph, its long-time analysis is in fact governed by the simpler Fokker-Planck equation (17) for the sole conditional mean. Here the situation is even simpler because the residual fluctuation $(X_t - \mathbb{E}_1(X_t))_{t\geq 0}$ has a trivial behavior (whilst the analysis of the residual is non-trivial in the prototype example).

Combining the two examples. For sure, the reader may wonder about a global framework that would cover both the prototype and the variant examples. While this would be indeed possible to extend in such a way the current setting, we have decided not to go up to this level of generality in order to keep the presentation and the notation at a reasonable level. In order to guess what the more general form of (4) should be, one first needs to understand which of the two coefficients *F* and *G* (in (4)) correspond respectively to $x \mapsto -ax$ and $m \mapsto f(\mu_1(m))$ in (15). At first sight, one may be tempted to regard $x \mapsto -ax$ as a specific example of *G*, but this is the wrong choice. In fact, the correct answer is to write -ax as $-a(x - \mu_1(m))$, which prompts us to associate the coefficient *a* in this model with the coefficient α in the prototype example. Next, we should see the remaining coefficient $m \mapsto -a\mu_1(m) + f(\mu_1(m))$ as a *new G*, with the subtlety that, in our global framework, *G* must be allowed to depend on both *x* and $\mu_1(m)$. In clear, the new version of (4) should be

(19)
$$d_t m_t = \nabla \cdot \left(\frac{\sigma^2 + \sigma_0^2}{2} \nabla m_t - m_t \left[G(\mu_1(m_t), \cdot) + F(\cdot - \mu_1(m_t)) \right] \right) dt - \sigma_0 \nabla m_t \cdot dB_t^0,$$

for a function $G : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}^d$. Back to the prototype example addressed in the previous paragraph, this says that the long-time behavior of $(m_t)_{t\geq 0}$ should now be compared with those of the standard SDE $dY_t = G(Y_t, Y_t)dt + \sigma_0 dB_t^0$, up to some fluctuation terms that one may expect to control properly if F is sufficiently decreasing. In this context, what truly matters are the confining properties of the *doubled* mapping $(x, x) \mapsto G(x, x)$.

We strongly believe that Theorem 8 could be extended to this setting, with a similar analysis. From a technical point of view, the gain would be limited: the difficulty would not come from the variable $\mu_1(m)$ in *G* but from the variable *x* (precisely because the strategy is to replace in the end the argument X_t by its conditional expectation $\mathbb{E}_1(X_t)$ in (5)). This difficulty is already present in the prototype example (13) and, in contrast, it is clear from the example (15) that the presence of the variable $\mu_1(m)$ in *G* does not raise any substantial difficulty. This explains how choice to work on (4) (and not on (19)).

An example without uniqueness recovery. As highlighted earlier, the finite-dimensional nature of the common noise constitutes a significant limitation to obtain ergodic properties on the process $(m_t)_{t\geq0}$ (see in contrast the recent work [24] by the first author for a 1d case with an infinite dimensional noise). In particular, this is the thrust of our work to identify one class of mean field dynamics for which the common noise really helps in this matter. However, it is clear that there is no chance to get uniqueness of the invariant measure for a generic class of stochastic Fokker-Planck SPDEs forced by a *simple* noise like B^0 .

A counter-example in dimension d = 1 is

$$\mathbf{d}_t m_t = \frac{\sigma(v(m_t))^2 + \sigma_0^2}{2} \partial_{xx}^2 m_t \mathbf{d}t + \partial_x(xm_t) \mathbf{d}t - \sigma_0 \partial_x m_t \mathbf{d}B_t^0,$$

for some function $\sigma : [0, +\infty) \to (0, +\infty)$. Whenever $\sigma_0 = 0$ and the function $\sigma^2(\cdot)/4$ admits several fixed points, the stationary solutions to the above equation cannot be unique. In this case, $(m_t)_{t\geq 0}$ is indeed solution of

(20)
$$\partial_t m_t = \frac{\sigma(v(m_t))^2}{2} \partial_{xx}^2(m_t) + \partial_x(xm_t), \quad t \ge 0$$

Then, similar to the stationary solutions found in the previous example, the stationary solutions (equivalently, the invariant measures of (5)) are here solutions of

$$\frac{\mathrm{d}m}{\mathrm{d}x}(x) = \exp\left(-2\frac{x^2}{\sigma^2(v(m))}\right), \quad x \in \mathbb{R}.$$

Then, one can show that, if the function $x \mapsto \sigma^2(x)/4$ admits several fixed points, then there are several stationary solutions to (20).

When adding common noise, the situation does not get better. For instance, assuming that the initial condition $P_0 = \delta_{\delta_0}$, one can show that for all t > 0, \mathbb{P}_0 -almost surely, m_t is a (random) Gaussian probability measure with parameters

$$\begin{cases} \mu_1(m_t) = \sigma_0 \int_0^t e^{s-t} dB_s^0, \\ v(m_t) = \int_0^t e^{2(s-t)} \sigma^2(v(m_s)) ds. \end{cases}$$

Then, it is not difficult to see that $(\mu_1(m_t))_{t \ge 0}$ is an ergodic process (it is in fact an Ornstein-Uhlenbeck process with a negative mean-reverting parameter). As for the dynamics of the variance $(v(m_t))_{t \ge 0}$, we get

$$\partial_t v(m_t) = -2v(m_t) + \frac{1}{2}\sigma^2(v(m_t)), \quad t \ge 0.$$

Obviously, the above equation has several stationary solutions (say σ^*) if the function $x \mapsto \sigma^2(x)/4$ admits several fixed points. In the latter case, $(m_t)_{t\geq 0}$ has several invariant measures, whose form is similar to (18) except that σ therein is now replaced by any σ^* . In this specific example, it is noteworthy that the common noise has no influence on the mean field term $(v(m_t))_{t\geq 0}$ (which is completely deterministic). Clearly, this is a consequence of the additive structure of the noise, the effect of which is just to shift (or to translate) the measures $(m_t)_{t\geq 0}$.

Conclusion. All these examples illustrate that the form of the mean field interaction in (4), based on the sole (conditional) mean state is key in the derivation of Theorem 8. In particular, this structure makes it possible to transmit the noise from the dynamics of the state variable $(X_t)_{t\geq0}$ in (5) to its conditional mean $(\mathbb{E}_1(X_t))_{t\geq0}$. As suggested in the discussion on the prototype example, the next step is to prove that the long run behavior of $(\mathbb{E}_1(X_t))_{t\geq0}$ is dictated by the ergodic properties of the SDE $dY_t = G(Y_t)dt + \sigma_0 dB_t^0$, at least if α is large and σ is small. This is the main line of the proof that is presented in the next two sections.

18

3. PROOF OF THEOREM 8

In this section, we explain how to deduce Theorem 8 from Proposition 7, but the proof of the latter is postponed to Section 4.

Before moving on to the proof of Theorem 8, we need two preliminary estimates: the first one addresses the conditional variance of the solution to (4), and the second one the conditional exponential moments of the solution.

3.1. First preliminary estimate: conditional variance. Given a solution *X* to Equation (5), we first focus on the conditional expectation given the common noise $(\mathbb{E}_1[X_t])_{t \ge 0}$:

Proposition 10. Let us consider a d-dimensional Brownian motion β^0 defined on $(\Omega_0, \mathcal{F}^0, \mathbb{P}_0)$ and adapted to the filtration \mathbb{F}^0 . Let Assumptions 1 and 2 be in force with the additional constraint $c_{\alpha_F} := \alpha_F - L_G > 0$. Let us consider $P_0 \in \mathcal{P}(\mathcal{P}_1(\mathbb{R}^d))$ and X_0 such that $\mathcal{L}_1(X_0) = m_0$ a.s., for some random measure m_0 satisfying $\mathcal{L}_0(m_0) = P_0$. Then, for X with dynamic given by

$$\begin{cases} dX_t = G(X_t)dt + F(X_t - \mathbb{E}_1[X_t])dt + \sigma dB_t + \sigma_0 d\beta_t^0\\ X_{t|t=0} = X_0, \end{cases}$$

we have for all $t \ge 0$,

(21)
$$\mathbb{E}_{1}\left[|X_{t} - \mathbb{E}_{1}[X_{t}]|^{2}\right] \leq \mathbb{E}_{1}\left[|X_{0} - \mathbb{E}_{1}[X_{0}]|^{2}\right]e^{-2c_{\alpha_{F}}t} + \frac{d\sigma^{2}}{c_{\alpha_{F}}},$$

where the previous inequality holds $\mathbb{P}_0 - a.s.$

Remark 11. (1) Inequality (21) does not depend on the choice of the Brownian motion $(\beta_t^0)_{t\geq 0}$ in the dynamics of X. Accordingly, it can be recast in terms of the solution (4) in the form

(22)
$$v(m_t) \leqslant v(m_0)e^{-2c_{\alpha_F}t} + \frac{d\sigma^2}{c_{\alpha_F}}, \quad t \ge 0.$$

(2) The result can be interpreted as follows. When $\sigma = 0$, the result is consistent with [44] and says that the process $(X_t)_{t\geq 0}$ is attracted by its conditional expectation in the long run. When $\sigma > 0$, the result provides a sharp estimate of the residual conditional variance.

(3) Back to Theorem 8, the challenge is precisely to prove that the long-time behaviour of X is dictated by the long-time behaviour of $\mathbb{E}_1[X]$ even though the residual term is not zero.

Proof of Proposition **10***.* Using the dynamics of the process *X*, one can write

$$d(X_t - \mathbb{E}_1[X_t]) = (G(X_t) - \mathbb{E}_1[G(X_t)])dt + (F(X_t - \mathbb{E}_1[X_t]) - \mathbb{E}_1[F(X_t - \mathbb{E}_1[X_t])])dt + \sigma dB_t$$

Then, applying Itô's formula, it comes

$$d|X_t - \mathbb{E}_1[X_t]|^2 = 2(X_t - \mathbb{E}_1[X_t]) \cdot (G(X_t) - \mathbb{E}_1[G(X_t)])dt$$

+ 2(X_t - \mathbb{E}_1[X_t]) \cdot (F(X_t - \mathbb{E}_1[X_t]) - \mathbb{E}_1[F(X_t - \mathbb{E}_1[X_t])])dt
+ 2\sigma(X_t - \mathbb{E}_1[X_t]) \cdot dB_t
+ d\sigma^2 dt

Using the fact that G is Lipschitz continuous and taking expectation, we get that

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}_{1}\left[|X_{t} - \mathbb{E}_{1}[X_{t}]|^{2}\right] \leq 2L_{G}\mathbb{E}_{1}\left[|X_{t} - \mathbb{E}_{1}[X_{t}]|^{2}\right] + 2\mathbb{E}_{1}\left[(X_{t} - \mathbb{E}_{1}[X_{t}]) \cdot F(X_{t} - \mathbb{E}_{1}[X_{t}])\right] + d\sigma^{2}.$$

Moreover, we can write

$$\mathbb{E}_{1}\left[(X_{t} - \mathbb{E}_{1}[X_{t}]) \cdot F(X_{t} - \mathbb{E}_{1}[X_{t}]) \right] = \mathbb{E}_{1}\left[(X_{t} - \mathbb{E}_{1}[X_{t}]) \cdot (F(X_{t} - \mathbb{E}_{1}[X_{t}]) - F(0)) \right] \\ \leqslant -\alpha_{F} \mathbb{E}_{1}\left[|X_{t} - \mathbb{E}_{1}[X_{t}]|^{2} \right],$$

where we recall that $\alpha_F > 0$ is defined in Assumption 2. Then,

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}_1[|X_t - \mathbb{E}_1[X_t]|^2] \leq -2(\alpha_F - L_G)\mathbb{E}_1[|X_t - \mathbb{E}_1[X_t]|^2] + d\sigma^2$$

Finally, we get the result using Grönwall's Lemma, as soon as $\alpha_F > L_G$.

3.2. Second preliminary estimate: conditional exponential moments. Let us state the following lemma which allows us to control the exponential moments of the process $(m_t)_{t\geq 0}$ under one invariant measure \overline{P} (which will be in the end 'the' invariant measure):

Lemma 12. Under Assumptions 1 and 2, assuming that $\alpha_F > L_G$, there exists an invariant measure $\overline{P} \in \mathfrak{P}_2(\mathfrak{P}_2(\mathbb{R}^d))$ for the process $(m_t)_{t \ge 0}$, such that

(23)
$$\int_{\mathcal{P}(\mathbb{R}^d)} \exp\left(2c_2 v(\widetilde{m})^{1/2}\right) \bar{P}(\mathrm{d}\widetilde{m}) < +\infty$$

where $c_2 := c_1^{\alpha_F}/c_{\alpha_F}$.

Remark 13. The result presented in Lemma 12 aligns with the global picture we gave for the behavior of the model. In particular, it is natural (and in fact well-expected) in the regime $\sigma \ll 1$ (which corresponds to the framework of Proposition 7). Indeed, we already know from [44] (see also Section 2.4) that, if $\sigma = 0$, the invariant measure \overline{P}^0 is supported by Dirac masses; therefore, the conclusion of Lemma 12 becomes straightforward when replacing \overline{P} by \overline{P}^0 in the statement. The thrust of Lemma 12 is thus to extend the result to positive values of σ . Although our proof relies on a direct computation, it is worth noticing that any invariant measure \overline{P} (for $\sigma > 0$) should satisfy

$$\mathbf{d}_1^{\mathcal{P}(\mathbb{R}^d)}(\bar{P}, \bar{P}^0) \leqslant C\sigma^2,$$

for some constant C independent of σ . In particular, any invariant \overline{P} is close to \overline{P}^0 when σ is small, hence justifying our intuition that Lemma 12 is a perturbation of the regime $\sigma = 0$.

Proof of Lemma 12. Consider a probability measure $P_0 \in \mathcal{P}(\mathcal{P}(\mathbb{R}^d))$ with $P_0(\mathcal{P}_2(\mathbb{R}^d)) = 1$ and the measure valued stochastic process $(m_t)_{t \ge 0}$ solving (4) with $\mathcal{L}_0(m_0) = P_0$ as initial condition. Then, letting $P_t = \mathcal{L}_0(m_t)$, $\forall t \ge 0$, we can easily show with arguments similar to those used in the proof of Proposition 2 in [44] that the sequence $(\mathcal{Q}_T)_{T \ge 0}$ defined by

$$\mathcal{Q}_T := \frac{1}{T} \int_0^T P_t \mathrm{d}t$$

admits a converging subsequence, still denoted by $(\Omega_T)_{T \ge 0}$, that converges to an invariant measure \overline{P} . For any T > 0,

$$\int_{\mathcal{P}(\mathbb{R}^d)} \exp\left(2c_2 v(\widetilde{m})^{1/2}\right) \mathcal{Q}_T(\mathrm{d}\widetilde{m}) = \frac{1}{T} \int_0^T \int_{\mathcal{P}(\mathbb{R}^d)} \exp\left(2c_2 v(\widetilde{m})^{1/2}\right) P_t(\mathrm{d}\widetilde{m})$$
$$= \frac{1}{T} \int_0^T \mathbb{E}_0\left[\exp\left(2c_2 v(m_t)^{1/2}\right)\right] \mathrm{d}t.$$

20

Moreover, (22) gives

$$\int_{\mathcal{P}(\mathbb{R}^d)} \exp\left(2c_2 v(\tilde{m})^{1/2}\right) \mathcal{Q}_T(\mathrm{d}\tilde{m}) \leq \frac{1}{T} \int_0^T \mathbb{E}_0 \left[\exp\left(2c_2 v(m_0)^{1/2} + \frac{2c_2\sigma}{\sqrt{c_{\alpha_F}}}\right) \right] \mathrm{d}t$$
$$\leq \mathbb{E}_0 \left[\exp\left(2c_2 v(m_0)^{1/2} + \frac{2c_2\sigma}{\sqrt{c_{\alpha_F}}}\right) \right],$$

which can be assumed to be finite by choosing an appropriate initial condition. Recall indeed that the choice of P_0 in the construction of the sequence $(\mathfrak{Q}_T)_{T \ge 0}$ is free. Therefore, assuming that

(24)
$$\int_{\mathcal{P}(\mathbb{R}^d)} \exp\left(2c_2 v(\widetilde{m})^{1/2}\right) P_0(\mathrm{d}\widetilde{m}) < +\infty,$$

and using lower-semi-continuity of the function $P \mapsto \langle \exp(2c_2 v(\cdot)^{1/2}); P \rangle_{\mathcal{P}(\mathbb{R}^d)}$, we get that

(25)
$$\int_{\mathcal{P}(\mathbb{R}^d)} \exp\left(2c_2 v(\widetilde{m})^{1/2}\right) \bar{P}(\mathrm{d}\widetilde{m}) \leq \lim \inf_{T \to +\infty} \int_{\mathcal{P}(\mathbb{R}^d)} \exp\left(2c_2 v(\widetilde{m})^{1/2}\right) \mathcal{Q}_T(\mathrm{d}\widetilde{m}) < +\infty,$$

which is (23).

In order to prove that $\overline{P} \in \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$, we use the same approach and get for any T > 0,

(26)
$$\int_{\mathcal{P}(\mathbb{R}^d)} \mu_2(\widetilde{m}) \mathcal{Q}_T(\mathrm{d}\widetilde{m}) = \frac{1}{T} \int_0^T \int_{\mathcal{P}(\mathbb{R}^d)} \mu_2(\widetilde{m}) P_t(\mathrm{d}\widetilde{m}) \mathrm{d}t$$
$$= \frac{1}{T} \int_0^T \mathbb{E}_0[\mu_2(m_t)] \mathrm{d}t.$$

Now, we claim that, under (24) and Assumptions 1 and 2,

$$\sup_{t \ge 0} \mathbb{E}_0[\mu_2(m_t)] < +\infty$$

Indeed, there exist two positive constants $b_1, b_2 > 0$, such that for all $(x, y) \in \mathbb{R}^d$,

$$(x-y) \cdot (G(x) - G(y)) \leq -b_1 |x-y|^2 + b_2.$$

Then, we get

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E}_0[\mu_2(m_t)] &\leqslant -2b_1 \mathbb{E}_0[\mu_2(m_t)] + 2b_2 + (\sigma^2 + \sigma_0^2) \\ &+ 2|G(0)|\mathbb{E}_0[\mu_2(m_t)]^{1/2} + 2C_F \mathbb{E}_0\left[\mu_2(m_t)^{1/2} v(m_t)^{1/2}\right]. \end{aligned}$$

Using once again (22) and Jensen's inequality to control the last term, we obtain

$$\frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E}_{0}[\mu_{2}(m_{t})] \leq -2b_{1}\mathbb{E}_{0}[\mu_{2}(m_{t})] + 2\left(|G(0)| + C_{F}v(m_{0})^{1/2} + \frac{C_{F}\sigma}{\sqrt{c_{\alpha_{F}}}}\right) \mathbb{E}_{0}\left[\mu_{2}(m_{t})\right]^{1/2}
+ 2b_{2} + d(\sigma^{2} + \sigma_{0}^{2})
\leq -b_{1}\mathbb{E}_{0}[\mu_{2}(m_{t})] + b_{1}^{-1}\left(|G(0)| + C_{F}v(m_{0})^{1/2} + \frac{C_{F}\sigma}{\sqrt{c_{\alpha_{F}}}}\right)^{2} + 2b_{2} + d(\sigma^{2} + \sigma_{0}^{2}),$$

and conclude using Grönwall's Lemma that $\sup_{t\geq 0} \mathbb{E}_0[\mu_2(m_t)] < +\infty$. Finally, using the same lower-semi-continuity argument as in (25) and returning to (26), we get

(27)
$$\int_{\mathcal{P}(\mathbb{R}^d)} \mu_2(\widetilde{m}) \overline{P}(\mathrm{d}\widetilde{m}) \leq \lim_{T \to +\infty} \int_{\mathcal{P}(\mathbb{R}^d)} \mu_2(\widetilde{m}) \mathcal{Q}_T(\mathrm{d}\widetilde{m}) < +\infty.$$

The latter implies that $\overline{P} \in \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$.

3.3. **Completion of the proof of Theorem 8.** We are now ready for the proof of Theorem 8 which is a straightforward combination of Proposition 7 and Lemma 12.

Proof of Theorem 8. We know from Proposition 7 that for any $\alpha_F \ge \max(L_G, m_G)$ and any $\sigma_0 > 0$, there exist $\overline{\sigma} > 0$ (depending on F, G, σ_0 and d) such that for all $\sigma \le \overline{\sigma}$ and for any 1-Lipschitz continuous function $\phi : \mathcal{P}_1(\mathbb{R}^d) \to \mathbb{R}$ and all $m, \widetilde{m} \in \mathcal{P}_2(\mathbb{R}^d)$,

$$|\mathscr{P}_t\phi(m) - \mathscr{P}_t\phi(\widetilde{m})| \leq \mathcal{K}(m,\widetilde{m})e^{-\mathfrak{c}t}, \quad t \ge 0,$$

for some $\mathfrak{c} > 0$ independent of m and \tilde{m} . Let us now consider an invariant measure $\bar{P} \in \mathcal{P}_2(\mathcal{P}_2(\mathbb{R}^d))$, with a finite conditional exponential moments, whose existence is ensured by Lemma 12. Then,

(28)
$$\left| \mathscr{P}_{t}\phi(m) - \int_{\mathscr{P}(\mathbb{R}^{d})} \mathscr{P}_{t}\phi(\widetilde{m})\overline{P}(\mathrm{d}\widetilde{m}) \right| \leq \int_{\mathscr{P}(\mathbb{R}^{d})} |\mathscr{P}_{t}\phi(m) - \mathscr{P}_{t}\phi(\widetilde{m})|\overline{P}(\mathrm{d}\widetilde{m})| \leq e^{-\mathfrak{c}t} \int_{\mathscr{P}(\mathbb{R}^{d})} \mathscr{K}(m,\widetilde{m})\overline{P}(\mathrm{d}\widetilde{m}).$$

Now, coming back to Proposition 7 for the shape of $\mathcal{K}(m, \tilde{m})$, we obtain that

$$\begin{split} \int_{\mathcal{P}(\mathbb{R}^d)} \mathcal{K}(m, \widetilde{m}) \bar{P}(\mathrm{d}\widetilde{m}) &\leqslant C_d \mu_2(m)^{1/2} \exp\left(c_2 v(m)^{1/2}\right) \int_{\mathcal{P}(\mathbb{R}^d)} \exp\left(c_2 v(m)^{1/2}\right) \bar{P}(\mathrm{d}\widetilde{m}) \\ &+ C_d \exp\left(c_2 v(\widetilde{m})^{1/2}\right) \int_{\mathcal{P}(\mathbb{R}^d)} \mu_2(\widetilde{m})^{1/2} \exp\left(c_2 v(\widetilde{m})^{1/2}\right) \bar{P}(\mathrm{d}\widetilde{m}), \end{split}$$

with $c_2 := c_1^{\alpha_F} / c_{\alpha_F}$.

Using Hölder inequality and leveraging on Lemma 12 we have

$$\int_{\mathcal{P}(\mathbb{R}^d)} \mathcal{K}(m, \widetilde{m}) \bar{P}(\mathrm{d}\widetilde{m}) \leqslant C_d \left(1 + \mu_2(m)^{1/2}\right) \exp\left(c_2 v(m)^{1/2}\right).$$

Substituting in (28), we complete the proof.

4. PROOF OF PROPOSITION 7

This section is dedicated to the proof of Proposition 7.

4.1. **Ansatz.** The strategy of proof of Proposition 7 relies on the auxiliary Proposition 10. This latter result says that the residual conditional variance of *X* (solution to Equation (5)) is small with σ . In turn, it prompts us to focus on the process $\mathbb{E}_1[X] = (\mathbb{E}_1[X_t])_{t \ge 0}$. The latter has the following dynamics:

$$d\mathbb{E}_1[X_t] = \mathbb{E}_1[G(X_t)]dt + \mathbb{E}_1[F(X_t - \mathbb{E}_1[X_t])]dt + \sigma_0 dB_t^0, \quad t \ge 0.$$

Whilst this is not a closed equation (in $\mathbb{E}_1[X_t]$), our approach is to write it as a perturbation of a finite-dimensional stochastic differential equation. Heuristically, we have (thanks to Proposition 10) that on a large time scale

(29)
$$\mathbb{E}_1[F(X_t - \mathbb{E}_1[X_t])] \approx \sigma^2 / c_{\alpha_F},$$

(30)
$$\mathbb{E}_1[G(X_t)] = G(\mathbb{E}_1[X_t]) + [\mathbb{E}_1[G(X_t)] - G(\mathbb{E}_1[X_t])] \approx G(\mathbb{E}_1[X_t]) + \sigma^2 / c_{\alpha_F}$$

Equations (29) and (30) are rather informal, and we refrain from giving here a rigorous meaning to the symbol \approx in both of them. However, these two expansions suggest that the dynamics of $(\mathbb{E}_1[X_t])_{t\geq 0}$ indeed resembles to the one of the diffusion process *Y*, constructed on $(\Omega_0, \mathbb{F}^0, \mathbb{P}_0)$ as the solution of

(31)
$$dY_t = G(Y_t) dt + \sigma_0 dB_t^0, \quad t \ge 0; \quad Y_0 = \mathbb{E}_1(X_0).$$

Here, it is worth emphasizing that, even obvious, (31) is not a McKean-Vlasov but a mere diffusion equation. In particular, the long time analysis of (31) falls within a much wider literature, since the study of the long time behaviour of diffusion processes has been an important topic of interest, see [2, 4, 5] to name just a few. Below, we make use of coupling arguments, in the spirit of Eberle [26], who stated contraction properties for the law of the solution of (31) by building on ideas developed earlier in [41]. The next section gives some details on this approach and the contraction results available in this framework.

4.2. **Contraction for classical diffusion processes.** Following [26], we introduce the following quantities :

$$\begin{aligned} R_0 &:= \inf\{R \ge 0 : \kappa(r) \ge 0, \quad \forall r \ge R\}, \\ R_1 &:= \inf\{R \ge R_0 : \kappa(r)R(R - R_0) \ge 8, \quad \forall r \ge R\}, \end{aligned}$$

so that $R_0 \leq R_1$. Thanks to Assumption 1, R_0 and R_1 are finite. Moreover, let

$$\begin{aligned} \varphi(r) &:= \exp\left(-\frac{1}{4}\int_0^r s\kappa(s)^- \mathrm{d}s\right), \quad \Phi(r) &:= \int_0^r \varphi(s)\mathrm{d}s, \\ g(r) &:= 1 - \left(\int_0^{r \wedge R_1} \Phi(s)\varphi(s)^{-1}\mathrm{d}s\right) \left(2\int_0^{R_1} \Phi(s)\varphi(s)^{-1}\mathrm{d}s\right)^{-1}, \end{aligned}$$

and (32)

$$f(r) := \int_0^r \varphi(s) g(s) \mathrm{d}s.$$

Then, we have that

- φ is non-increasing, $\varphi(0) = 1$, and $\varphi(r) = \varphi(R_0)$ for any $r \ge R_0$;
- *g* is non-increasing, g(0) = 1, and g(r) = 1/2 for any $r \ge R_1$;
- f' is bounded on $[0, +\infty)$;

• f is concave, f(0) = 0, f'(0) = 1, and

$$\Phi(r)/2 \leq f(r) \leq \Phi(r)$$
 for any $r \geq 0$.

The last display combined with the fact that for all $r \ge 0$, $\Phi(r)/2 \ge \varphi(R_0)r/2$ and $\Phi(r) \le r$, implies

(33)
$$\kappa_1 r \leqslant f(r) \leqslant r,$$

with $\kappa_1 := \varphi(R_0)/2$. This allows us to define the equivalent deformation of 1-Wasserstein distance

$$\mathbf{d}_f^{\mathbb{R}^d}(\mu,\nu) \coloneqq \inf_{\pi \in \Pi(\mu,\nu)} \int f(|x-y|) \pi(\,\mathrm{d} x,\,\mathrm{d} y),$$

for any two probability measures μ , ν on \mathbb{R}^d , where the infimum is taken over all couplings π of μ and ν . Last but not least, the function f we constructed satisfies

(34)
$$f''(r) - \frac{1}{4}r\kappa(r)f'(r) \le -\frac{c}{2\sigma_0^2}f(r) \quad \text{for all } r > 0,$$

and for some c > 0. Such concentration property for the function f is classical, and the proof of inequality (34) can be found in [18, 25, 26]. Leveraging on the contraction property of f given by Equation (34), we get the following result (as a direct consequence of [26]):

Theorem 14 (Corollary 2 in [26]). Under Assumption 1 and with $\sigma_0 > 0$, there exists a constant $c := c_{\sigma_0} > 0$, such that for any pair (Y, \tilde{Y}) of solutions to Equation (31) with different initial conditions, we have

(35)
$$d_f^{\mathbb{R}^d}\left(\mathcal{L}_1(Y_t), \mathcal{L}_1(\widetilde{Y}_t)\right) \leq d_f^{\mathbb{R}^d}\left(\mathcal{L}_1(Y_0), \mathcal{L}_1(\widetilde{Y}_0)\right) e^{-ct}, \quad t \ge 0.$$

Remark 15. Thanks to (33), we get that inequality (35) holds (up to some constant) when replacing $d_f^{\mathbb{R}^d}$ by the classical $d_1^{\mathbb{R}^d}$ Wasserstein distance.

The idea of the proof of Theorem 14 is to use the reflection coupling introduced in [41] and widely used in the literature [18, 25, 26, 27]. The classical reflection coupling is constructed by multiplying the noise by a symmetry matrix mapping instantaneously $Y_t - \tilde{Y}_t$ on its opposite, see [41, Section 3] for details. In this framework, the distance between the two processes Y and \tilde{Y} evolves according to an Itô process with constant noise intensity. The hope is thus to benefit from the presence of the noise to obtain further recurrent properties that prevent the processes from staying away from one another. For sure, one cannot expect this intuition to hold true for any type of drift; in order to guarantee a relevant form of recurrence, some further confining properties are necessary.

When dealing with the stochastic differential equation (31), the condition $\limsup_{r\to+\infty} \kappa(r) > 0$ required in Assumption 1 is key in the verification of the latter confining properties. In itself, this is not an obvious result: one must indeed keep in mind that *G* is not strictly decreasing, meaning that κ may be negative on some part of the space. Clearly, things become even more subtle in presence of the conditional McKean-Vlasov interaction. Still, one would like to settle down a similar reflection argument for Equation (5), which raises preliminary questions of measurability. Briefly, in order to preserve the form of the common noise, one must consider a reflection matrix that is measurable with respect to the latter (and not to the idiosyncratic noise). The idea developed in

24

the next subsection is to introduce a coupling (X, \tilde{X}) of solutions to (5) in which the reflection is achieved by reflecting with respect to $\mathbb{E}_1[X_t] - \mathbb{E}_1[\tilde{X}_t]$. However, while reflecting with respect to the conditional expectations may seem appropriate to our objective, it cannot suffice on its own to get a proof of Proposition 7 that would be a mere copy and paste of the argument of [26]. The difficulty that one meets when adapting [26] to (5) is clear: the processes that solve (5) are not those that are used in the reflection. This is where Proposition 10 comes in and this is one of the main innovation of our work: we use the fact that the distance between the solutions to (5) and their conditional expectations (which are also the processes entering the reflection) is small with σ . Mathematically, the difficulty is to revisit the whole machinery of [26] by using the fact that, although it is not zero, the latter distance is small.

4.3. **Proof of Proposition 7.** To ease the reading of this subsection, we briefly outline the strategy of proof:

Proof Outline.

- Step 1. We construct a coupling (X, X) inspired by the reflection coupling introduced in [41] (as explained in subsection 4.2) and adapted to the presence of a common noise.
- *Steps 2, 3 & 4.* We prove that this coupling allows the conditional expectations $\mathbb{E}_1[X]$ and $\mathbb{E}_1[\widetilde{X}]$ to get closer in time, up to some residual error that depends on the distance between the centred processes $X \mathbb{E}_1[X]$ and $\widetilde{X} \mathbb{E}_1[\widetilde{X}]$. This is established thanks to the confining property of *G* guaranteed by Assumption 1.
- Step 5. In parallel (the key point is to run the two arguments at the same time), we prove the mirror result, namely we show that the centred processes X − E₁[X] and X̃ − E₁[X̃] get closer in time, up to some residual error that depends on the distance between the conditional expectations E₁[X] and E₁[X̃]. This is established thanks to the strictly contracting properties of the interaction term F guaranteed by Assumption 2.
- *Step 6*. We combine together the results of Step 4 and Step 5 to conclude.

Proof. Step 1. Let us consider a bounded and $d_1^{\mathbb{R}^d}$ -Lipschitz continuous function $\phi : \mathcal{P}_1(\mathbb{R}^d) \to \mathbb{R}$. In particular there exists $\|\phi\|_{\text{lip}} > 0$ such that for any $m, \tilde{m} \in \mathcal{P}_1(\mathbb{R}^d)$,

$$|\phi(m) - \phi(\widetilde{m})| \leq \|\phi\|_{\operatorname{lip}} \mathrm{d}_1^{\mathbb{R}^a}(m, \widetilde{m}).$$

Let us then fix $m, \tilde{m} \in \mathcal{P}_2(\mathbb{R}^d)$ together with two independent random variables X_0, \tilde{X}_0 , defined on $(\Omega_1, \mathbb{F}^1, \mathbb{P}_1)$ such that $\mathcal{L}_1(X_0) = m$ and $\mathcal{L}_1(\tilde{X}_0) = \tilde{m}$. Then, we extend X_0 and \tilde{X}_0 to the product space $(\Omega, \mathbb{F}, \mathbb{P})$ in a natural manner.

Remark 16. It is important to notice here that, even if the initial conditions X_0 , \tilde{X}_0 are defined on the product space $(\Omega, \mathbb{F}, \mathbb{P})$, they are independent of \mathcal{F}^0 .

For a given $\delta > 0$, let us now introduce the following coupling of two solutions of the SDE (5) with initial conditions X_0 and \tilde{X}_0 :

$$(36) \quad \begin{cases} \mathrm{d}X_t^{\delta} = G(X_t^{\delta})\mathrm{d}t + F\left(X_t^{\delta} - \mathbb{E}_1[X_t^{\delta}]\right)\mathrm{d}t + \sigma\mathrm{d}B_t + \sigma_0\left\{\pi_{\delta}(E_t^{\delta})\mathrm{d}B_t^0 + \lambda_{\delta}(E_t^{\delta})\mathrm{d}\widetilde{B}_t^0\right\}\\ \mathrm{d}\widetilde{X}_t^{\delta} = G(\widetilde{X}_t^{\delta})\mathrm{d}t + F\left(\widetilde{X}_t^{\delta} - \mathbb{E}_1[\widetilde{X}_t^{\delta}]\right)\mathrm{d}t + \sigma\mathrm{d}B_t\\ + \sigma_0\left\{\pi_{\delta}(E_t^{\delta})\left(\mathrm{Id} - 2e_t^{\delta}\left(e_t^{\delta}\right)^{\mathsf{T}}\right)\mathrm{d}B_t^0 + \lambda_{\delta}(E_t^{\delta})\mathrm{d}\widetilde{B}_t^0\right\},\end{cases}$$

where

- the initial conditions are given by X^δ₀ = X₀ and X̃^δ₀ = X̃₀;
- \widetilde{B}^0 is a *d*-dimensional Brownian motion defined on $(\Omega_0, \mathcal{F}^0, \mathbb{P}_0)$ adapted to the filtration \mathbb{F}^0 and independent of B^0 ;
- for all $t \ge 0$, $E_t^{\delta} := \mathbb{E}_1[X_t^{\delta}] \mathbb{E}_1[\widetilde{X}_t^{\delta}]$, and

$$e_t^{\delta} := \begin{cases} E_t^{\delta} / |E_t^{\delta}| & \text{if } |E_t^{\delta}| \neq 0\\ 0 & \text{otherwise;} \end{cases}$$

• the function π_{δ} is Lipschitz continuous on \mathbb{R}^d with value in [0, 1], such that for any $x \in \mathbb{R}^d$,

$$\pi_{\delta}(x) = \begin{cases} 0 & \text{if} \quad |x| \leq \delta/2 \\ 1 & \text{if} \quad |x| \geq \delta. \end{cases}$$

• the function λ_{δ} is Lipschitz-continuous and satisfies $\lambda_{\delta}^2(x) = 1 - \pi_{\delta}^2(x)$, for all $x \in \mathbb{R}^d$.

The first thing to notice is that under the Lipschitz continuity assumptions on *G* and *F*, and the integrability properties of the initial conditions, we have existence and uniqueness of a weak solution of (36). Moreover, thanks to the choice of π_{δ} and λ_{δ} , the pair $(X^{\delta}, \tilde{X}^{\delta})$ is a coupling of (Z, \tilde{Z}) where for *Z* and \tilde{Z} admits the same dynamic,

(37)
$$dZ_t = G(Z_t)dt + F(Z_t - \mathbb{E}_1[Z_t])dt + \sigma dB_t + \sigma_0 dB_t^0,$$

with initial condition $Z_0 = X_0$ and $\tilde{Z}_0 = \tilde{X}_0$. This is mainly due to Levy's characterisation of the Brownian motion which is applicable here as for all $x \in \mathbb{R}^d$, $\lambda_{\delta}^2(x) + \pi_{\delta}^2(x) = 1$. Details on the coupling method are given in subsection 4.2. Below, we focus on the long-time behavior of

$$\mathbb{E}_0\big[\mathbb{E}_1\big[|X_t^\delta - \widetilde{X}_t^\delta|\big]\big],$$

for a given $\delta > 0$.

Step 2. Letting
$$A_t^{\delta} := X_t^{\delta} - \mathbb{E}_1[X_t^{\delta}]$$
 and $\widetilde{A}_t^{\delta} := \widetilde{X}_t^{\delta} - \mathbb{E}_1[\widetilde{X}_t^{\delta}]$, we get that for all $t \ge 0$,
 $|X_t^{\delta} - \widetilde{X}_t^{\delta}| \le |\mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}]| + |A_t^{\delta} - \widetilde{A}_t^{\delta}|.$

The idea of the proof is to control both $|\mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}]|$ and $|A_t^{\delta} - \widetilde{A}_t^{\delta}|$ simultaneously. This will allow us to prove that both terms converge in fact to 0 (in some sense). Let us begin with the difference of the conditional expectations:

$$d\left(\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right) = \mathbb{E}_{1}[G(X_{t}^{\delta}) - G(\widetilde{X}_{t}^{\delta})]dt + \mathbb{E}_{1}[F(A_{t}^{\delta}) - F(\widetilde{A}_{t}^{\delta})]dt + 2\sigma_{0}\pi_{\delta}(E_{t}^{\delta})e_{t}^{\delta}e_{t}^{\delta} \cdot dB_{t}^{0}.$$

26

Using Itô's formula and recalling that $E_t^{\delta} = \mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}]$ and $e_t^{\delta} \cdot E_t^{\delta} = |E_t^{\delta}|$, we obtain

$$d\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|^{2} = 2\left(\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right) \cdot \left(\mathbb{E}_{1}[G(X_{t}^{\delta})] - \mathbb{E}_{1}[G(\widetilde{X}_{t}^{\delta})]\right) dt + 2\left(\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right) \cdot \left(\mathbb{E}_{1}[F(A_{t}^{\delta})] - \mathbb{E}_{1}[F(\widetilde{A}_{t}^{\delta})]\right) dt + 4\sigma_{0}\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right| \pi_{\delta}(E_{t}^{\delta})e_{t}^{\delta} \cdot dB_{t}^{0} + 2\sigma_{0}^{2}\pi_{\delta}^{2}(E_{t}^{\delta})dt.$$

This combined with the particular shape of the reflection, and noting that $(\int_0^t e_s^{\delta} \cdot dB_s^0)_t$ is a 1-dimensional Brownian motion tank's to Levy's characterisation allows us to write

(38)

$$d \left| \mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}] \right| = e_{t}^{\delta} \cdot \left(\mathbb{E}_{1}[G(X_{t}^{\delta})] - \mathbb{E}_{1}[G(\widetilde{X}_{t}^{\delta})] \right) dt$$

$$+ e_{t}^{\delta} \cdot \left(\mathbb{E}_{1}[F(A_{t}^{\delta})] - \mathbb{E}_{1}[F(\widetilde{A}_{t}^{\delta})] \right) dt$$

$$+ 2\sigma_{0}\pi_{\delta}(E_{t}^{\delta})e_{t}^{\delta} \cdot dB_{t}^{0}, \quad t \ge 0.$$

Deriving Equation (38) is not straightforward and can be done using similar techniques as in [25]. The underlying concept is that the presence of π_{δ} keeps the process E^{δ} from lingering near zero. The complete proof of (38) can be found in Appendix A. We now recall (see Assumption 1) that κ is a continuous function on $(0, \infty)$ satisfying

(39)
$$\liminf_{r \to \infty} \kappa(r) > 0 \quad \text{and} \quad \int_0^1 r \kappa(r)^- \mathrm{d}r < \infty.$$

Then, using Itô's formula once again, we obtain

$$df\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) = f'\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) e_{t}^{\delta} \cdot \left(\mathbb{E}_{1}[G(X_{t}^{\delta})] - \mathbb{E}_{1}[G(\widetilde{X}_{t}^{\delta})]\right) dt \\ + f'\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) e_{t}^{\delta} \cdot \left(\mathbb{E}_{1}[F(A_{t}^{\delta})] - \mathbb{E}_{1}[F(\widetilde{A}_{t}^{\delta})]\right) dt \\ + 2\sigma_{0}f'\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) \pi_{\delta}(E_{t}^{\delta})e_{t}^{\delta} \cdot dB_{t}^{0} \\ + 2\sigma_{0}^{2}f''\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) \pi_{\delta}^{2}(E_{t}^{\delta}) dt.$$

Then, we rewrite

(40)

$$df\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) = f'\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) e_{t}^{\delta} \cdot \left(G(\mathbb{E}_{1}[X_{t}^{\delta}]) - G(\mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}])\right) dt + 2\sigma_{0}^{2}f''\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) \pi_{\delta}^{2}(E_{t}^{\delta}) dt + 2\sigma_{0}f'\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) \pi_{\delta}(E_{t}^{\delta})e_{t}^{\delta} \cdot dB_{t}^{0} + r_{t}^{1}dt,$$

where for all $t \ge 0$, $r_t^1 := \mathbf{I}_1(t) + \mathbf{II}_1(t) + \mathbf{III}_1(t)$, with

$$\mathbf{I}_{1}(t) = f'\left(\left|E_{t}^{\delta}\right|\right)e_{t}^{\delta} \cdot \mathbb{E}_{1}\left[\int_{0}^{1} \{DG(\rho X_{t}^{\delta} + (1-\rho)\widetilde{X}_{t}^{\delta}) - DG(\rho \mathbb{E}_{1}[X_{t}^{\delta}] + (1-\rho)\mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}])\} d\rho \\ \left(\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right)\right];$$

(41)
$$\mathbf{I}_{1}(t) = f'\left(\left|E_{t}^{\delta}\right|\right)e_{t}^{\delta} \cdot \mathbb{E}_{1}\left[\int_{0}^{1}\left\{DG(\rho X_{t}^{\delta} + (1-\rho)\widetilde{X}_{t}^{\delta}) - DG(\rho\mathbb{E}_{1}[X_{t}^{\delta}] + (1-\rho)\mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}])\right\}d\rho$$
$$\left(A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}\right)\right];$$

$$\mathbb{III}_{1}(t) = f'\left(\left|E_{t}^{\delta}\right|\right)e_{t}^{\delta} \cdot \mathbb{E}_{1}\left[\int_{0}^{1} DF(\rho A_{t}^{\delta} + (1-\rho)\widetilde{A}_{t}^{\delta}) d\rho \left(A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}\right)\right],$$

where we used the following two identities: $\mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}] + A_t^{\delta} - \widetilde{A}_t^{\delta} = X_t^{\delta} - \widetilde{X}_t^{\delta}$ and $\mathbb{E}_1[A_t^{\delta}] - \mathbb{E}_1[\widetilde{A}_t^{\delta}] = 0.$

Step 3. Recall (34):

$$f''(r) - \frac{1}{4}r\kappa(r)f'(r) \leqslant -\frac{c}{2\sigma_0^2}f(r) \quad \text{ for all } r > 0.$$

Combining the above equation with the fact that, from Assumption 1,

$$(x-y)\cdot (G(x)-G(y)) \leqslant -\frac{2}{\sigma_0^2}\kappa(|x-y|)|x-y|^2, \quad x,y \in \mathbb{R}^d,$$

we obtain

$$f'\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) e_{t}^{\delta} \cdot \left(G(\mathbb{E}_{1}[X_{t}^{\delta}]) - G(\mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}])\right) + 2\sigma_{0}^{2}f''\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) \pi_{\delta}^{2}(E_{t}^{\delta})$$

$$(42) \qquad \leqslant -\frac{\sigma_0}{2} f'\left(\left|E_t^{\delta}\right|\right) \left|E_t^{\delta}\right| \kappa\left(\left|E_t^{\delta}\right|\right) + 2\sigma_0^2 f''\left(\left|E_t^{\delta}\right|\right) \pi_\delta^2(E_t^{\delta}) \leqslant 2\sigma_0^2 \left(-\frac{1}{4} f'\left(\left|E_t^{\delta}\right|\right) \left|E_t^{\delta}\right| \kappa\left(\left|E_t^{\delta}\right|\right) + f''\left(\left|E_t^{\delta}\right|\right)\right) \pi_\delta^2(E_t^{\delta}) - \frac{\sigma_0^2}{2} f'\left(\left|E_t^{\delta}\right|\right) \left|E_t^{\delta}\right| \kappa\left(\left|E_t^{\delta}\right|\right) \lambda_\delta^2(E_t^{\delta}) \leqslant -cf\left(\left|\mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}]\right|\right) + h(\delta),$$

where

$$h(\delta) := \frac{\sigma_0^2}{2} f'\left(\left|E_t^{\delta}\right|\right) |E_t^{\delta}| \kappa\left(\left|E_t^{\delta}\right|\right) \lambda_{\delta}^2(E_t^{\delta}).$$

Now, recalling that $\lambda_{\delta}(x) = 0$ as soon as $|x| \ge \delta$, we obtain for $\delta \in [0, 1]$,

(43)
$$|h(\delta)| \leq \frac{\sigma_0^2}{2} \|f'\|_{\infty} \delta \kappa_2,$$

where $\kappa_2 = \sup_{0 \le r \le 1} \kappa(r)$, from which we deduce that $h(\delta) \to 0$ with δ . Then, back to (40), we can write

$$df\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right) \leq -cf\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right)dt + |r_{t}^{1}|dt + h(\delta)dt$$

+
$$2\sigma_0 f'\left(\left|\mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}]\right|\right) \pi_{\delta}(E_t^{\delta})e_t^{\delta} \cdot \mathrm{d}B_t^0,$$

and

$$\frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E}_0 \left[f\left(\left| \mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}] \right| \right) \right] \leqslant -c \, \mathbb{E}_0 \left[f\left(\left| \mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}] \right| \right) \right] + \mathbb{E}_0[|r_t^1|] + \mathbb{E}_0[|h(\delta)|]$$

Using Grönwall's Lemma, we obtain

(44)
$$\mathbb{E}_{0}\left[f\left(\left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right)\right] \leqslant \mathbb{E}_{0}\left[f\left(\left|\mathbb{E}_{1}[X_{0}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{0}^{\delta}]\right|\right)\right]e^{-ct} + \int_{0}^{t}e^{-c(t-s)}\mathbb{E}_{0}[|r_{s}^{1}|]\mathrm{d}s + \frac{h(\delta)}{c}.$$

Step 4. It remains to control the three parts in the expansion of the residual term r_{1}^{1} , see (41). For the two first ones, we use the Lipschitz continuity of DG with respect to the norm $\|\cdot\|$ and write

$$|\mathbf{I}_{1}(t)| \leq C_{G} \mathbb{E}_{1} \left[|X_{t}^{\delta} - \mathbb{E}_{1}[X_{t}^{\delta}]|^{2} + |\widetilde{X}_{t}^{\delta} - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]|^{2} \right]^{1/2} \left| \mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}] \right| ;$$

$$|\mathbf{I}_{1}(t)| \leq C_{G} \mathbb{E}_{1} \left[|X_{t}^{\delta} - \mathbb{E}_{1}[X_{t}^{\delta}]|^{2} + |\widetilde{X}_{t}^{\delta} - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]|^{2} \right]^{1/2} \mathbb{E}_{1} \left[|A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}|^{2} \right]^{1/2} .$$

Letting $\eta_t^{\delta} = \mathbb{E}_1 \left[|X_t^{\delta} - \mathbb{E}_1[X_t^{\delta}]|^2 + |\widetilde{X}_t^{\delta} - \mathbb{E}_1[\widetilde{X}_t^{\delta}]|^2 \right]^{1/2}$, for all $t \ge 0$, and recalling from Proposition 10 that

$$\eta_t^\delta \leqslant \eta_0^\delta e^{-c_{\alpha_F}t} + 2\sigma c_{\alpha_F}^{-1/2}\sqrt{d},$$

we obtain

(45)
$$\begin{aligned} |\mathbf{I}_{1}(t)| + |\mathbf{II}_{1}(t)| \\ \leqslant C_{G} \left(\eta_{0}^{\delta} e^{-c_{\alpha_{F}}t} + 2\sigma c_{\alpha_{F}}^{-1/2}\sqrt{d} \right) \left(\mathbb{E}_{1} \left[|A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}|^{2} \right]^{1/2} + \left| \mathbb{E}_{1} [X_{t}^{\delta}] - \mathbb{E}_{1} [\widetilde{X}_{t}^{\delta}] \right| \right). \end{aligned}$$

For the term $\mathbf{III}_1(t)$, we use the fact that $\mathbb{E}_1[A_t^{\delta}] = \mathbb{E}_1[\widetilde{A}_t^{\delta}] = 0$ and write

(46)
$$|\mathbf{III}_{1}(t)| = \left| \mathbb{E}_{1} \left[\int_{0}^{1} \{ DF(\rho A_{t}^{\delta} + (1-\rho)\widetilde{A}_{t}^{\delta}) - DF(0) \} \mathrm{d}\rho \cdot (A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}) \right] \right|$$
$$\leq C_{F} \left(\eta_{0}^{\delta} e^{-c_{\alpha_{F}}t} + 2\sigma c_{\alpha_{F}}^{-1/2} \sqrt{d} \right) \mathbb{E}_{1} \left[|A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}|^{2} \right]^{1/2},$$

where we used once again Proposition 10 together with the Lipschitz continuity of F to get the last line. Then, combining (40), (45) and (46), we have

$$\mathbb{E}_{0}[|r_{t}^{1}|] \leq \mathbb{E}_{0}[|\mathbf{I}_{1}(t)| + |\mathbf{\Pi}_{1}(t)| + |\mathbf{\Pi}_{1}(t)|]$$

$$\leq (C_{G} + C_{F}) \left(\eta_{0}^{\delta} e^{-c_{\alpha_{F}}t} + 2\sigma c_{\alpha_{F}}^{-1/2}\sqrt{d}\right) \mathbb{E}_{0} \left[\mathbb{E}_{1}\left[|A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}|^{2}\right]^{1/2} + \left|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]\right|\right].$$

Here, we recall that η_0^{δ} is deterministic because m and \tilde{m} are deterministic, see Remark 16. Finally, using inequalities (33) and (44), we obtain

(47)

$$\kappa_{1}\mathbb{E}_{0}\left[|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]|\right] \leq e^{-ct} \left|\mathbb{E}_{1}[X_{0}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{0}^{\delta}]\right| + \frac{h(\delta)}{c} \\
+ C_{b} \int_{0}^{t} e^{-c(t-s)} \left(\eta_{0}^{\delta}e^{-c_{\alpha_{F}}s} + 2\sigma c_{\alpha_{F}}^{-1/2}\sqrt{d}\right) \mathbb{E}_{0} \left[\mathbb{E}_{1}\left[|A_{s}^{\delta} - \widetilde{A}_{s}^{\delta}|^{2}\right]^{1/2} + |\mathbb{E}_{1}[X_{s}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{s}^{\delta}]|\right] ds,$$
where $C_{b} := C_{G} + C_{F}.$

Step 5. We now focus on the (normed) difference process $(|A_t^{\delta} - \widetilde{A}_t^{\delta}|)_{t \ge 0}$. Thanks to the properties of F, as stated in Assumption 2, we are here able to show that it decreases exponentially fast to zero.

Recalling from Step 2 that $A_t^{\delta} = X_t^{\delta} - \mathbb{E}_1[X_t^{\delta}]$ for $t \ge 0$, we have

$$dA_t^{\delta} = \left(G(X_t^{\delta}) - \mathbb{E}_1[G(X_t^{\delta})]\right) dt + \left(F(X_t^{\delta} - \mathbb{E}_1[X_t^{\delta}]) dt - \mathbb{E}_1[F(X_t^{\delta} - \mathbb{E}_1[X_t^{\delta}]]\right) dt + \sigma dB_t,$$

from which we obtain

$$\frac{\mathrm{d}}{\mathrm{d}t}(A_t^{\delta} - \widetilde{A}_t^{\delta}) = \left[G(X_t^{\delta}) - G(\widetilde{X}_t^{\delta})\right] + \left[F(A_t^{\delta}) - F(\widetilde{A}_t^{\delta})\right] \\ - \left[\mathbb{E}_1[G(X_t^{\delta})] - \mathbb{E}_1[G(\widetilde{X}_t^{\delta})]\right] - \left[\mathbb{E}_1[F(A_t^{\delta})] - \mathbb{E}_1[F(\widetilde{A}_t^{\delta})]\right].$$

Then,

(48)

$$\frac{\mathrm{d}}{\mathrm{d}t}|A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}|^{2} = 2(A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}) \cdot (G(X_{t}^{\delta}) - G(\widetilde{X}_{t}^{\delta})) \\
+ 2(A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}) \cdot (F(A_{t}^{\delta}) - F(\widetilde{A}_{t}^{\delta})) \\
- 2(A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}) \cdot (\mathbb{E}_{1}[G(X_{t}^{\delta})] - \mathbb{E}_{1}[G(\widetilde{X}_{t}^{\delta})]) \\
- 2(A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}) \cdot (\mathbb{E}_{1}[F(A_{t}^{\delta})] - \mathbb{E}_{1}[F(\widetilde{A}_{t}^{\delta})])$$

We can immediately leverage on Assumption 2 to control the second term on the right hand side: ${}^{*\delta} = \widetilde{A}{}^{\delta}_{\iota}|^2.$ δ $\widetilde{\delta}$ $\widetilde{\delta}$ $\widetilde{\delta}$ $\widetilde{\delta}$

(49)
$$2(A_t^{\delta} - \tilde{A}_t^{\delta}) \cdot (F(A_t^{\delta}) - F(\tilde{A}_t^{\delta})) \leqslant -2\alpha_F |A_t^{\delta} - \tilde{A}_t^{\delta}|^2$$

Let us now focus on the terms involving G. We have

$$\begin{split} & (A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot \left(G(X_t^{\delta}) - G(\widetilde{X}_t^{\delta}) - \mathbb{E}_1[G(X_t^{\delta})] + \mathbb{E}_1[G(\widetilde{X}_t^{\delta})] \right) \\ &= (A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot \left[\left(\int_0^1 \left\{ DG(\rho X_t^{\delta} + (1 - \rho) \widetilde{X}_t^{\delta}) - DG(\rho \mathbb{E}_1[X_t^{\delta}] + (1 - \rho) \mathbb{E}_1[\widetilde{X}_t^{\delta}]) \right\} d\rho \right) (X_t^{\delta} - \widetilde{X}_t^{\delta}) \right] \\ &+ (A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot \left[\left(\int_0^1 DG(\rho \mathbb{E}_1[X_t^{\delta}] + (1 - \rho) \mathbb{E}_1[\widetilde{X}_t^{\delta}]) d\rho \right) (X_t^{\delta} - \widetilde{X}_t^{\delta}) \right] \\ &- (A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot \mathbb{E}_1 \left[\left(\int_0^1 \left\{ DG(\rho X_t^{\delta} + (1 - \rho) \widetilde{X}_t^{\delta}) - DG(\rho \mathbb{E}_1[X_t^{\delta}] + (1 - \rho) \mathbb{E}_1[\widetilde{X}_t^{\delta}]) \right\} d\rho \right) (X_t^{\delta} - \widetilde{X}_t^{\delta}) \right] \\ &- (A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot \left[\left(\int_0^1 DG(\rho \mathbb{E}_1[X_t^{\delta}] + (1 - \rho) \mathbb{E}_1[\widetilde{X}_t^{\delta}]) d\rho \right) (\mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}]) \right]. \end{split}$$

.

By combining the second and fourth term in the right hand side, we obtain

$$\begin{split} & (A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot \left(G(X_t^{\delta}) - G(\widetilde{X}_t^{\delta})) - \mathbb{E}_1[G(X_t^{\delta})] + \mathbb{E}_1[G(\widetilde{X}_t^{\delta})] \right) \\ = & (A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot \left[\left(\int_0^1 \left\{ DG(\rho X_t^{\delta} + (1 - \rho) \widetilde{X}_t^{\delta}) - DG(\rho \mathbb{E}_1[X_t^{\delta}] + (1 - \rho) \mathbb{E}_1[\widetilde{X}_t^{\delta}]) \right\} \mathrm{d}\rho \right) (X_t^{\delta} - \widetilde{X}_t^{\delta}) \right] \\ & + & (A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot \left(\int_0^1 DG(\rho \mathbb{E}_1[X_t^{\delta}] + (1 - \rho) \mathbb{E}_1[\widetilde{X}_t^{\delta}]) \mathrm{d}\rho \right) (A_t^{\delta} - \widetilde{A}_t^{\delta}) \\ & - & (A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot \mathbb{E}_1 \left[\left(\int_0^1 \left\{ DG(\rho X_t^{\delta} + (1 - \rho) \widetilde{X}_t^{\delta}) - DG(\rho \mathbb{E}_1[X_t^{\delta}] + (1 - \rho) \mathbb{E}_1[\widetilde{X}_t^{\delta}]) \right\} \mathrm{d}\rho \right) (X_t^{\delta} - \widetilde{X}_t^{\delta}) \right] \\ = : & \overline{\Delta}_1(t) + \overline{\Delta}_2(t) + \overline{\Delta}_3(t). \end{split}$$

Then, if we perform the following decomposition of the first term on the right-hand side, we obtain

(50)
$$\overline{\mathbf{A}}_1(t) = \overline{\mathbf{A}}\overline{\mathbf{A}}_1(t) + \overline{\mathbf{A}}\overline{\mathbf{A}}_2(t),$$

with

$$\begin{split} \overline{\mathbf{A}}\overline{\mathbf{A}}_{2}(t) &:= (A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}) \cdot \Big(\int_{0}^{1} \Big\{ DG(\rho X_{t}^{\delta} + (1-\rho)\widetilde{X}_{t}^{\delta}) \\ &- DG(\rho \mathbb{E}_{1}[X_{t}^{\delta}] + (1-\rho)\mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]) \Big\} \mathrm{d}\rho \Big) (A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}). \end{split}$$

Then, combining the terms $\overline{\Delta}_2(t)$ and $\overline{\Delta}\overline{\Delta}_2(t)$, we get

(51)

$$(A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot \left(G(X_t^{\delta}) - G(\widetilde{X}_t^{\delta}) - \mathbb{E}_1[G(X_t^{\delta})] + \mathbb{E}_1[G(\widetilde{X}_t^{\delta})] \right)$$

$$= \overline{\Delta \Delta}_1(t) + [\overline{\Delta}_2(t) + \overline{\Delta}_2(t)] + \overline{\Delta}_3(t)$$

$$=: \overline{\Delta \Delta}_1(t) + \overline{\mathbf{I}}_2(t) + \overline{\Delta}_3(t),$$

where $\overline{AA}_1(t)$ and $\overline{A}_3(t)$ have been already defined and $\underline{II}_2(t)$ is defined by

$$\mathbf{I}_{2}(t) := (A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}) \cdot \left(\int_{0}^{1} DG(\rho X_{t}^{\delta} + (1-\rho)\widetilde{X}_{t}^{\delta}) \mathrm{d}\rho \right) (A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}).$$

We now control $\underline{\Pi}_2(t)$ as we handled Equation (49). We get

(52)
$$\mathbf{I}_2(t) \leqslant m_G |A_t^{\delta} - \widetilde{A}_t^{\delta}|^2,$$

where we recall from (6) that $m_G \in \mathbb{R}$ stands for the upper bound of the function $r \mapsto [-\sigma_0^2 \kappa(r)/2]$ introduced in Assumption 1. Indeed, as a consequence of 1, one can show that for any $x, y \in \mathbb{R}^d$,

$$y \cdot (DG(x)y) \leq m_G |y|^2.$$

Then, inserting (49), (51) and (52) into (48), we obtain

$$\frac{\mathrm{d}}{\mathrm{d}t} |A_t^{\delta} - \widetilde{A}_t^{\delta}|^2 \leq -2(\alpha_F - m_G)|A_t^{\delta} - \widetilde{A}_t^{\delta}|^2 + \overline{\Delta} \overline{\Delta}_1(t) + \overline{\Delta}_3(t) + 2(A_t^{\delta} - \widetilde{A}_t^{\delta}) \cdot (\mathbb{E}_1[F(A_t^{\delta})] - \mathbb{E}_1[F(\widetilde{A}_t^{\delta})]).$$

By taking expectation under \mathbb{P}_1 and by using the fact that $\mathbb{E}_1[A_t^{\delta}] = \mathbb{E}_1[\widetilde{A}_t^{\delta}] = 0$, we get rid the last two terms in the above right-hand side. We obtain

(53)
$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}_{1}\left[|A_{t}^{\delta}-\widetilde{A}_{t}^{\delta}|^{2}\right] \leqslant -2(\alpha_{F}-m_{G})\mathbb{E}_{1}\left[|A_{t}^{\delta}-\widetilde{A}_{t}^{\delta}|^{2}\right]+2\mathbb{E}_{1}\left[\overline{\mathbf{A}}\mathbf{A}_{1}(t)\right]$$

In order to control $\overline{\Delta\Delta}_1(t)$, we come back to (50) and use the fact that $\mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}]$ is measurable with respect to the common noise. By Cauchy-Schwarz' inequality, we obtain

$$\mathbb{E}_1\left[\overline{\mathbf{A}}\overline{\mathbf{A}}_1(t)\right] \leqslant C_G \mathbb{E}_1[|A_t^{\delta} - \widetilde{A}_t^{\delta}|^2]^{1/2} \eta_t^{\delta} |\mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}]|,$$

where we recall that $\eta_t^{\delta} = \mathbb{E}_1[|A_t^{\delta}|^2 + |\widetilde{A}_t^{\delta}|^2]^{1/2}$. Then, from Proposition 10, we get

$$\mathbb{E}_1\left[\overline{\mathbf{A}}\overline{\mathbf{A}}_1(t)\right] \leqslant \mathbb{E}_1\left[|A_t^{\delta} - \widetilde{A}_t^{\delta}|^2\right]^{1/2} \left(\eta_0^{\delta} e^{-c_{\alpha_F}t} + \sigma c_{\alpha_F}^{-1/2}\sqrt{d}\right) |\mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}]|,$$

and (53) rewrites

(54)
$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}_{1}\left[|A_{t}^{\delta}-\widetilde{A}_{t}^{\delta}|^{2}\right] \leqslant -2(\alpha_{F}-m_{G})\mathbb{E}_{1}\left[|A_{t}^{\delta}-\widetilde{A}_{t}^{\delta}|^{2}\right] +2\mathbb{E}_{1}[|A_{t}^{\delta}-\widetilde{A}_{t}^{\delta}|^{2}]^{1/2}\left(\eta_{0}^{\delta}e^{-c_{\alpha_{F}}t}+\sigma c_{\alpha_{F}}^{-1/2}\sqrt{d}\right)|\mathbb{E}_{1}[X_{t}^{\delta}]-\mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]|.$$

Then, we get

(55)
$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}_{1}\left[|A_{t}^{\delta}-\widetilde{A}_{t}^{\delta}|^{2}\right]^{1/2} \leq -(\alpha_{F}-m_{G})\mathbb{E}_{1}\left[|A_{t}^{\delta}-\widetilde{A}_{t}^{\delta}|^{2}\right]^{1/2} + C_{G}(\eta_{0}^{\delta}e^{-c_{\alpha_{F}}t}+\sigma c_{\alpha_{F}}^{-1/2}\sqrt{d})|\mathbb{E}_{1}[X_{t}^{\delta}]-\mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]|,$$

see Appendix A.2 for details. Using Grönwall's Lemma, we obtain

$$\begin{split} \mathbb{E}_{1}[|A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}|^{2}]^{1/2} &\leqslant \mathbb{E}_{1}[|A_{0}^{\delta} - \widetilde{A}_{0}^{\delta}|^{2}]^{1/2} \exp\left(-(\alpha_{F} - m_{G})t\right) \\ &+ C_{G} \int_{0}^{t} e^{-(\alpha_{F} - m_{G})(t-s)} (\eta_{0}^{\delta} e^{-c_{\alpha_{F}}s} + \sigma c_{\alpha_{F}}^{-1/2} \sqrt{d}) |\mathbb{E}_{1}[X_{s}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{s}^{\delta}] |\mathrm{d}s. \end{split}$$

Finally, taking again expectation \mathbb{E}_0 ,

(56)
$$\mathbb{E}_{0}\left[\mathbb{E}_{1}\left[|A_{t}^{\delta}-\widetilde{A}_{t}^{\delta}|^{2}\right]^{1/2}\right] \leq \exp(-(\alpha_{F}-m_{G})t)\mathbb{E}_{1}\left[|A_{0}^{\delta}-\widetilde{A}_{0}^{\delta}|^{2}\right]^{1/2} + C_{G}\int_{0}^{t}e^{-(\alpha_{F}-m_{G})(t-s)}(\eta_{0}^{\delta}e^{-c_{\alpha_{F}}s} + \sigma c_{\alpha_{F}}^{-1/2}\sqrt{d})\mathbb{E}_{0}\left[|\mathbb{E}_{1}\left[X_{s}^{\delta}\right]-\mathbb{E}_{1}\left[\widetilde{X}_{s}^{\delta}\right]|\right]\mathrm{d}s$$

32

Step 6. Now, as (47) remains true replacing the rate c by $\tilde{c} \leq c$, we can assume without loss of generality that $\alpha_F > m_G + c$. We can then combine Equations (47) and (56) to obtain, for all $t \ge 0$,

$$\mathbb{E}_{0} \left[\mathbb{E}_{1} \left[|A_{t}^{\delta} - \widetilde{A}_{t}^{\delta}|^{2} \right]^{1/2} + |\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]| \right] \\$$
(57)
$$\leq \left(\mathbb{E}_{1} \left[|A_{0}^{\delta} - \widetilde{A}_{0}^{\delta}|^{2} \right]^{1/2} + |\mathbb{E}_{1}[X_{0}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{0}^{\delta}]| \right) e^{-ct} + \frac{h(\delta)}{c} \\
+ c_{3}^{b} \int_{0}^{t} e^{-c(t-s)} \left(\eta_{0}^{\delta} e^{-c_{\alpha_{F}}s} + 2\sigma c_{\alpha_{F}}^{-1/2} \sqrt{d} \right) \mathbb{E}_{0} \left[\mathbb{E}_{1} \left[|A_{s}^{\delta} - \widetilde{A}_{s}^{\delta}|^{2} \right]^{1/2} + |\mathbb{E}_{1}[X_{s}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{s}^{\delta}]| \right] ds,$$

for some constant $c_3^b > 0$ that depends on F and G. With the notation $\Theta_t^{\delta} := \mathbb{E}_1 \left[|A_t^{\delta} - \widetilde{A}_t^{\delta}|^2 \right]^{1/2} + |\mathbb{E}_1[X_t^{\delta}] - \mathbb{E}_1[\widetilde{X}_t^{\delta}]|$, Equation (57) reads

$$\mathbb{E}_{0}[\Theta_{t}^{\delta}] \leqslant \Theta_{0}^{\delta} e^{-ct} + \frac{h(\delta)}{c} + c_{3}^{b} \int_{0}^{t} e^{-c(t-s)} \left(\eta_{0}^{\delta} e^{-c_{\alpha_{F}}s} + 2\sigma c_{\alpha_{F}}^{-1/2}\sqrt{d}\right) \mathbb{E}_{0}\left[\Theta_{s}^{\delta}\right] \mathrm{d}s.$$

Then, using Grönwall Lemma, we obtain

(58)
$$e^{ct} \mathbb{E}_{0} \left[\Theta_{t}^{\delta} \right] \leq \left(\Theta_{0} + \frac{h(\delta)}{c} \right) \exp \left(\frac{c_{3}^{b}}{c_{\alpha_{F}}} \eta_{0} + \frac{c_{3}^{b} \sigma \sqrt{d}}{\sqrt{c_{\alpha_{F}}}} t \right) + h(\delta) \int_{0}^{t} e^{cs} \exp \left(\frac{c_{3}^{b}}{c_{\alpha_{F}}} \eta_{0} + \frac{c_{3}^{b} \sigma \sqrt{d}}{\sqrt{c_{\alpha_{F}}}} (t-s) \right) \mathrm{d}s$$

We emphasize here that we write Θ_0 and η_0 without reference to δ because these quantities are predetermined and independent of δ . Now, let us notice that creating the couplage we defined two Brownian motions adapted to the filtration \mathbb{F}_0 :

$$\beta_t^{\delta} = \int_0^t \pi_{\delta}(E_s^{\delta}) (\mathrm{Id} - 2e_s^{\delta}(e_s^{\delta})^{\mathsf{T}}) \mathrm{d}B_s^0 + \int_0^t \lambda_{\delta}(E_s^{\delta}) \mathrm{d}\widetilde{B}_s^0$$
$$\widetilde{\beta}_t^{\delta} = \int_0^t \pi_{\delta}(E_s^{\delta}) \mathrm{d}B_s^0 + \int_0^t \lambda_{\delta}(E_s^{\delta}) \mathrm{d}\widetilde{B}_s^0.$$

Setting $m_t^{\delta} = \mathcal{L}_1(X_t^{\delta})$ et $\widetilde{m}_t^{\delta} = \mathcal{L}_1(\widetilde{X}_t^{\delta})$, we get

$$d_t m_t^{\delta} = \nabla \cdot \left(\frac{\sigma^2 + \sigma_0^2}{2} \nabla m_t^{\delta} - m_t^{\delta} \left(G + F(\cdot - \mu_1(m_t^{\delta}))\right)\right) dt - \sigma_0 \nabla m_t^{\delta} \cdot d\beta_t^{\delta},$$

$$d_t \widetilde{m}_t^{\delta} = \nabla \cdot \left(\frac{\sigma^2 + \sigma_0^2}{2} \nabla \widetilde{m}_t^{\delta} - \widetilde{m}_t^{\delta} \left(G + F(\cdot - \mu_1(\widetilde{m}_t^{\delta}))\right)\right) dt - \sigma_0 \nabla \widetilde{m}_t^{\delta} \cdot d\widetilde{\beta}_t^{\delta}.$$

Then, one has $\mathscr{P}_t \phi(m) = \mathbb{E}_0 \left[\phi(m_t^{\delta}) | m_0^{\delta} = m \right]$ and $\mathscr{P}_t \phi(\widetilde{m}) = \mathbb{E}_0 \left[\phi(\widetilde{m}_t^{\delta}) | \widetilde{m}_0^{\delta} = \widetilde{m} \right]$. Then, we obtain that for any $\delta > 0$,

(59)
$$\|\mathscr{P}_t\phi(m) - \mathscr{P}_t\phi(\widetilde{m})\| \leq \|\phi\|_{\operatorname{lip}} \mathbb{E}_0\left[\mathbb{E}_1\left[\left|X_t^{\delta} - \widetilde{X}_t^{\delta}\right|\right]\right] \\ \leq \|\phi\|_{\operatorname{lip}} \mathbb{E}_0[\Theta_t^{\delta}]$$

Plugging the estimate given by Equation (58) into Equation (59) gives

$$\begin{split} |\mathscr{P}_t \phi(m) - \mathscr{P}_t \phi(\widetilde{m})| \leq & \|\phi\|_{\mathrm{lip}} \left(\Theta_0 + \frac{h(\delta)}{c}\right) \exp\left(\frac{c_3^b}{c_{\alpha_F}}\eta_0 + \left(\frac{c_3^b \sigma \sqrt{d}}{\sqrt{c_{\alpha_F}}} - c\right)t\right) \\ & + \|\phi\|_{\mathrm{lip}} h(\delta) \int_0^t e^{-c(t-s)} \exp\left(\frac{c_3^b}{c_{\alpha_F}}\eta_0 + \frac{c_3^b \sigma \sqrt{d}}{\sqrt{c_{\alpha_F}}}(t-s)\right) \mathrm{d}s. \end{split}$$

The latter being true for all $\delta > 0$, we can consider the limit $\delta \to 0$ to finally obtain

$$|\mathscr{P}_t\phi(m) - \mathscr{P}_t\phi(\widetilde{m})| \leq \|\phi\|_{\mathrm{lip}}\Theta_0 \exp\left(\frac{c_3^b}{c_{\alpha_F}}\eta_0 + \frac{c_3^b\sigma\sqrt{d}}{\sqrt{c_{\alpha_F}}}t - ct\right)$$

Then, for σ and α satisfying $\mathfrak{c} := \sigma c_3^b c_{\alpha_F}^{-1/2} - c < 0$, we get the expected exponential rate of convergence. Moreover, writing

$$\Theta_{0} \exp\left(\frac{c_{3}^{b}}{c_{\alpha_{F}}}\eta_{0}\right) = \left(\mathbb{E}_{1}\left[|A_{0}-\widetilde{A}_{0}|^{2}\right]^{1/2} + |\mathbb{E}_{1}[X_{0}] - \mathbb{E}_{1}[\widetilde{X}_{0}]|\right) \exp\left(\frac{c_{3}^{b}}{c_{\alpha_{F}}}\eta_{0}\right)$$
$$\leq C_{2}\left(\mu_{2}(m)^{1/2} + \mu_{2}(\widetilde{m})^{1/2}\right) \exp\left(\frac{c_{3}^{b}}{c_{\alpha_{F}}}(v(m)^{1/2} + v(\widetilde{m})^{1/2})\right),$$

for a new constant positive C_2 , we complete the proof of Proposition 7.

Acknowledgments. François Delarue acknowledges the financial support of the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (AdG ELISA project, Grant agreement No. 101054746). Raphaël Maillet wishes to thank the Université Côte d'Azur for its hospitality during the preparation of this work, along with Pierre Cardaliaguet for fruitful discussions.

REFERENCES

- L. Ambrosio, N. Gigli, and G. Savaré, <u>Gradient flows in metric spaces and in the space of probability measures</u>, second ed., Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 2008. MR-2401600
- [2] C. Ané, S. Blachère, D. Chafaï, P. Fougères, I. Gentil, F. Malrieu, C. Roberto, and G. Scheffer, <u>Sur les inégalités de Sobolev logarithmiques</u>, With a preface by D. Bakry and M. Ledoux, Panoramas et Synthèses [Panoramas and Syntheses], vol. 10, Société Mathématique de France, Paris, 2000. MR-1845806
- [3] L. Angeli, J. Barré, M. Kolodziejczyk, and M. Ottobre, <u>Well-posedness and stationary solutions of McKean-Vlasov</u> (S)PDEs, J. Math. Anal. Appl. 526 (2023), no. 2, Paper No. 127301, 49, doi. MR-4579326
- [4] D. Bakry, F. Barthe, P. Cattiaux, and A. Guillin, A simple proof of the Poincaré inequality for a large class of probability measures including the log-concave case, Electron. Commun. Probab. 13 (2008), 60–66, doi. MR-2386063
- [5] D. Bakry and Michel Émery, <u>Diffusions hypercontractives</u>, Séminaire de probabilités, XIX, 1983/84, Lecture Notes in Math., vol. 1123, Springer, Berlin, 1985, pp. 177–206, doi. MR-889476
- [6] K. Bashiri, On the long-time behaviour of McKean-Vlasov paths, Electron. Commun. Probab. 25 (2020), Paper No. 52, 14, doi. MR-4129676
- [7] S. Benachour, B. Roynette, and P. Vallois, <u>Nonlinear self-stabilizing processes</u>. II. Convergence to invariant probability, Stochastic Process. Appl. 75 (1998), no. 2, 203–224, doi. MR-1632197
- [8] F. Bolley, I. Gentil, and A. Guillin, Uniform convergence to equilibrium for granular media, Arch. Ration. Mech. Anal. 208 (2013), no. 2, 429–445, doi. MR-3035983
- [9] P. Briand, P. Cardaliaguet, P.-E. Chaudru de Raynal, and Y. Hu, Forward and backward stochastic differential equations with normal constraints in law, Stochastic Process. Appl. 130 (2020), no. 12, 7021–7097, doi. MR-4167200

- [10] O. A. Butkovsky, On ergodic properties of nonlinear Markov chains and stochastic McKean-Vlasov equations, Theory Probab. Appl. 58 (2014), no. 4, 661–674, doi. MR-3403022
- [11] P. Cardaliaguet, Notes on mean field games, Tech. report, Technical report, 2010.
- [12] P. Cardaliaguet, F. Delarue, J.-M. Lasry, and P.-L. Lions, <u>The master equation and the convergence problem in mean field games</u>, Annals of Mathematics Studies, vol. 201, Princeton University Press, Princeton, NJ, 2019, doi. <u>MR-3967062</u>
- [13] R. Carmona and F. Delarue, <u>Probabilistic theory of mean field games with applications.</u> I, Mean field FBSDEs, control, and games, Probability Theory and Stochastic Modelling, vol. 83, Springer, Cham, 2018. MR-3752669
- [14] _____, Probabilistic theory of mean field games with applications. II, Mean field games with common noise and master equations, Probability Theory and Stochastic Modelling, vol. 84, Springer, Cham, 2018. MR-3753660
- [15] J.A. Carrillo, R.J. McCann, and C. Villani, <u>Kinetic equilibration rates for granular media and related equations:</u> entropy dissipation and mass transportation estimates, Rev. Mat. Iberoamericana **19** (2003), no. 3, 971–1018, doi. MR-2053570
- [16] P. Cattiaux, A. Guillin, and F. Malrieu, Probabilistic approach for granular media equations in the non-uniformly convex case, Probab. Theory Related Fields 140 (2008), no. 1-2, 19–40, doi. MR-2357669
- [17] M. Coghi and B. Gess, <u>Stochastic nonlinear Fokker-Planck equations</u>, Nonlinear Anal. 187 (2019), 259–278, doi. MR-3954095
- [18] G. Conforti, Coupling by reflection for controlled diffusion processes: turnpike property and large time behavior of Hamilton-Jacobi-Bellman equations, Ann. Appl. Probab. 33 (2023), no. 6A, 4608–4644, doi. MR-4674060
- [19] D.A. Dawson, <u>Critical dynamics and fluctuations for a mean-field model of cooperative behavior</u>, J. Statist. Phys. **31** (1983), no. 1, 29–85, doi. MR-711469
- [20] D.A Dawson and J. Vaillancourt, Stochastic McKean-Vlasov equations, NoDEA Nonlinear Differential Equations Appl. 2 (1995), no. 2, 199–229, doi. MR-1328577
- [21] P. Del Moral and J. Tugaut, <u>Uniform propagation of chaos and creation of chaos for a class of nonlinear diffusions</u>, Stoch. Anal. Appl. **37** (2019), no. 6, 909–935, doi. MR-4020054
- [22] F. Delarue, <u>Restoring uniqueness to mean-field games by randomizing the equilibria</u>, Stoch. Partial Differ. Equ. Anal. Comput. 7 (2019), no. 4, 598–678, doi. MR-4022284
- [23] F. Delarue and R. Foguen-Tchuendom, Selection of equilibria in a linear quadratic mean-field game, Stochastic Process. Appl. 130 (2020), no. 2, 1000–1040, doi. MR-4046528
- [24] F. Delarue and W. Hammersley, <u>Rearranged stochastic heat equation: Ergodicity and related gradient descent on the</u> space of probability measures, arXiv 2403.16140 (2024).
- [25] A. Durmus, A. Eberle, A. Guillin, and R. Zimmer, <u>An elementary approach to uniform in time propagation of chaos</u>, Proc. Amer. Math. Soc. 148 (2020), no. 12, 5387–5398, doi. MR-4163850
- [26] A. Eberle, <u>Reflection couplings and contraction rates for diffusions</u>, Probab. Theory Related Fields 166 (2016), no. 3-4, 851–886, doi. MR-3568041
- [27] A. Eberle and R. Zimmer, Sticky couplings of multidimensional diffusions with different drifts, Ann. Inst. Henri Poincaré Probab. Stat. 55 (2019), no. 4, 2370–2394, doi. MR-4029157
- [28] B. Fehrman and B. Gess, <u>Well-posedness of nonlinear diffusion equations with nonlinear, conservative noise</u>, Arch. Ration. Mech. Anal. 233 (2019), no. 1, 249–322, doi. MR-3974641
- [29] R. Foguen-Tchuendom, Uniqueness for linear-quadratic mean field games with common noise, Dyn. Games Appl. 8 (2018), no. 1, 199–210, doi. MR-3764697
- [30] P.K. Friz and B. Gess, <u>Stochastic scalar conservation laws driven by rough paths</u>, Ann. Inst. H. Poincaré C Anal. Non Linéaire 33 (2016), no. 4, 933–963, doi. MR-3519527
- [31] T. Funaki, <u>A certain class of diffusion processes associated with nonlinear parabolic equations</u>, Z. Wahrsch. Verw. Gebiete 67 (1984), no. 3, 331–348, doi. MR-762085
- [32] B. Gess and P.E. Souganidis, <u>Scalar conservation laws with multiple rough fluxes</u>, Commun. Math. Sci. 13 (2015), no. 6, 1569–1597, doi. MR-3351442
- [33] _____, Stochastic non-isotropic degenerate parabolic-hyperbolic equations, Stochastic Process. Appl. 127 (2017), no. 9, 2961–3004, doi. MR-3682120
- [34] W. Hammersley, D. Šiška, and Ł. Szpruch, <u>Weak existence and uniqueness for McKean-Vlasov SDEs with common</u> noise, Ann. Probab. 49 (2021), no. 2, 527–555, doi. MR-4255126

F. DELARUE ; R. MAILLET ; E. TANRÉ

- [35] S. Herrmann and J. Tugaut, <u>Non-uniqueness of stationary measures for self-stabilizing processes</u>, Stochastic Process. Appl. **120** (2010), no. 7, 1215–1246, doi. MR-2639745
- [36] B. Jianhai and W. Jian, Long time behavior of one-dimensional mckean-vlasov sdes with common noise, arXiv:2401.07665 (2024).
- [37] R. Jordan, D. Kinderlehrer, and F. Otto, The variational formulation of the Fokker-Planck equation, SIAM J. Math. Anal. 29 (1998), no. 1, 1–17, doi. MR-1617171
- [38] M. Kac, Foundations of kinetic theory, Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, 1954–1955, vol. III, Univ. California Press, Berkeley-Los Angeles, Calif., 1956, pp. 171–197. MR-84985
- [39] T.G. Kurtz and J. Xiong, Particle representations for a class of nonlinear SPDEs, Stochastic Process. Appl. 83 (1999), no. 1, 103–126, doi. MR-1705602
- [40] D. Lacker, M. Shkolnikov, and J. Zhang, <u>Superposition and mimicking theorems for conditional McKean-Vlasov</u> equations, J. Eur. Math. Soc. (JEMS) 25 (2023), no. 8, 3229–3288, doi. MR-4612111
- [41] T. Lindvall and L. C. G. Rogers, Coupling of multidimensional diffusions by reflection, Ann. Probab. 14 (1986), no. 3, 860–872, https://www.jstor.org/stable/pdf/2244138.pdf. MR-841588
- [42] P.-L. Lions, B. Perthame, and P.E. Souganidis, <u>Scalar conservation laws with rough (stochastic) fluxes</u>, Stoch. Partial Differ. Equ. Anal. Comput. 1 (2013), no. 4, 664–686, doi. MR-3327520
- [43] P.-L Lions, B. Perthame, and P.E. Souganidis, <u>Scalar conservation laws with rough (stochastic) fluxes: the spatially dependent case</u>, Stoch. Partial Differ. Equ. Anal. Comput. 2 (2014), no. 4, 517–538, doi. MR-3274890
- [44] R. Maillet, <u>A note on the Long-Time behaviour of Stochastic McKean-Vlasov Equations with common noise</u>, arXiv:2306.16130 (2023).
- [45] F. Malrieu, Logarithmic Sobolev inequalities for some nonlinear PDE's, Stochastic Process. Appl. 95 (2001), no. 1, 109–132, doi. MR-1847094
- [46] _____, Convergence to equilibrium for granular media equations and their Euler schemes, Ann. Appl. Probab. 13 (2003), no. 2, 540–560, doi. MR-1970276
- [47] H. P. McKean, <u>A class of Markov processes associated with nonlinear parabolic equations</u>, Proc. Nat. Acad. Sci. U.S.A. 56 (1966), 1907–1911, doi. MR-221595
- [48] S. Méléard, Asymptotic behaviour of some interacting particle systems; McKean-Vlasov and Boltzmann models, Probabilistic models for nonlinear partial differential equations (Montecatini Terme, 1995), Lecture Notes in Math., vol. 1627, Springer, Berlin, 1996, pp. 42–95, doi. MR-1431299
- [49] F. Otto, Evolution of microstructure in unstable porous media flow: a relaxational approach, Comm. Pure Appl. Math. 52 (1999), no. 7, 873–915, doi. MR-1682800
- [50] _____, The geometry of dissipative evolution equations: the porous medium equation, Comm. Partial Differential Equations 26 (2001), no. 1-2, 101–174, doi. MR-1842429
- [51] D. Revuz and M. Yor, <u>Continuous martingales and brownian motion</u>, vol. 293, Springer Science & Business Media, 2013.
- [52] A.-S. Sznitman, <u>Topics in propagation of chaos</u>, École d'Été de Probabilités de Saint-Flour XIX—1989, Lecture Notes in Math., vol. 1464, Springer, Berlin, 1991, pp. 165–251, doi. MR-1108185

APPENDIX A.

A.1. Proof of Equation (38).

Proof. Using Itô's formula, and recalling that $E^{\delta} = X^{\delta} - \widetilde{X}^{\delta}$, we get that for any $\delta > 0$,

$$\begin{aligned} \mathbf{d} |\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]|^{2} &= 2(\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]) \cdot \left(\mathbb{E}_{1}[G(X_{t}^{\delta})] - \mathbb{E}_{1}[G(\widetilde{X}_{t}^{\delta})]\right) \mathbf{d}t \\ &+ 2(\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]) \cdot \left(\mathbb{E}_{1}[F(X_{t}^{\delta} - \mathbb{E}_{1}[X_{t}^{\delta}])] - \mathbb{E}_{1}[G(F(\widetilde{X}_{t}^{\delta} - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]))]\right) \mathbf{d}t \\ &+ 4\sigma_{0}\pi_{\delta}(E_{t}^{\delta})((\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]) \cdot e_{t}^{\delta})e_{t}^{\delta} \cdot \mathbf{d}B_{t}^{0} \\ &+ 4\sigma_{0}^{2}\pi_{\delta}^{2}(E_{t}^{\delta})\mathbf{d}t.\end{aligned}$$

For any $\varepsilon > 0$, let us introduce $\psi_{\varepsilon} : [0, +\infty] \in r \mapsto (r + \varepsilon)^{1/2}$. Since this function is twice continuously differentiable, we can write

$$d\psi_{\varepsilon} \left(|\mathbb{E}_{1}[X_{t}^{\delta}] - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta}]|^{2} \right) = 2\psi_{\varepsilon}'(|E_{t}^{\delta}|^{2})E_{t}^{\delta} \cdot \left(\mathbb{E}_{1}[G(X_{t}^{\delta})] - \mathbb{E}_{1}[G(\widetilde{X}_{t}^{\delta})]\right) dt + 2\psi_{\varepsilon}'(|E_{t}^{\delta}|^{2})E_{t}^{\delta} \cdot \left(\mathbb{E}_{1}[F(X_{t}^{\delta} - \mathbb{E}_{1}[X_{t}^{\delta})] - \mathbb{E}_{1}[F(\widetilde{X}_{t}^{\delta} - \mathbb{E}_{1}[\widetilde{X}_{t}^{\delta})]\right) dt + 4\sigma_{0}\psi_{\varepsilon}'(|E_{t}^{\delta}|^{2})\pi_{\delta}(E_{t}^{\delta})(E_{t}^{\delta} \cdot e_{t}^{\delta})e_{t}^{\delta} \cdot dB_{t}^{0} + 4\sigma_{0}^{2}\psi_{\varepsilon}'(|E_{t}^{\delta}|^{2})\pi_{\delta}^{2}(E_{t}^{\delta}) dt + 8\sigma_{0}^{2}\psi_{\varepsilon}''(|E_{t}^{\delta}|^{2})|E_{t}^{\delta}|^{2}\pi_{\delta}(E_{t}^{\delta}) dt.$$

We now want to take the limit $\varepsilon \to 0$. Using dominated convergence theorem and stochastic dominated convergence theorem as stated in [51, Theorem 2.12], combined with the bound $4r\psi'_{\varepsilon}(r^2) \leq 1$, we can deal with the first two lines and get that for all $t \geq 0$,

$$(61) \lim_{\varepsilon \to 0} \int_0^t 2\psi_{\varepsilon}'(|E_s^{\delta}|^2) E_s^{\delta} \cdot \left(\mathbb{E}_1[G(X_s^{\delta})] - \mathbb{E}_1[G(\widetilde{X}_s^{\delta})] \right) \mathrm{d}s = \int_0^t e_s^{\delta} \cdot \left(\mathbb{E}_1[G(X_s^{\delta})] - \mathbb{E}_1[G(\widetilde{X}_s^{\delta})] \right) \mathrm{d}s,$$
$$\lim_{\varepsilon \to 0} \int_0^t 2\psi_{\varepsilon}'(|E_s^{\delta}|^2) E_s^{\delta} \cdot \left(\mathbb{E}_1[F(X_t^{\delta} - \mathbb{E}_1[X_t^{\delta}])] - \mathbb{E}_1[F(\widetilde{X}_t^{\delta} - \mathbb{E}_1[\widetilde{X}_t^{\delta}])] \right) \mathrm{d}s$$

(62)
$$\lim_{\varepsilon \to 0} \int_{0} 2\psi_{\varepsilon}'(|E_{s}^{\delta}|^{2})E_{s}^{\delta} \cdot \left(\mathbb{E}_{1}[F(X_{t}^{\delta} - \mathbb{E}_{1}[X_{t}^{\delta}])] - \mathbb{E}_{1}[F(\tilde{X}_{t}^{\delta} - \mathbb{E}_{1}[\tilde{X}_{t}^{\delta}])]\right) ds$$
$$= \int_{0}^{t} e_{s}^{\delta} \cdot \left(\mathbb{E}_{1}[F(X_{t}^{\delta} - \mathbb{E}_{1}[X_{t}^{\delta}])] - \mathbb{E}_{1}[F(\tilde{X}_{t}^{\delta} - \mathbb{E}_{1}[\tilde{X}_{t}^{\delta}])]\right) ds,$$

and

(63)
$$\lim_{\varepsilon \to 0} \int_0^t 4\sigma_0 \psi_{\varepsilon}'(|E_s^{\delta}|^2) \pi_{\delta}(E_s^{\delta}) (E_s^{\delta} \cdot e_s^{\delta}) e_s^{\delta} \cdot \mathrm{d}B_s^0 = \int_0^t 2\sigma_0 \pi_{\delta}(E_s^{\delta}) e_s^{\delta} \cdot \mathrm{d}B_s^0,$$

almost surely. For the last two terms in Equation (60), let us remark that for all $r \ge 0$,

(64)
$$\psi_{\varepsilon}'(r) + 2\psi''(r) = \frac{1}{2(r+\varepsilon)^{1/2}} - \frac{r}{2(r+\varepsilon)^{3/2}} \to 0,$$

when ε tends to 0. In order to we need to properly take the limit $\varepsilon \to 0$ in Equation (60), we need to take advantage of the presence of the function π_{δ} for $\delta > 0$. In Indeed, we have

$$\begin{aligned} \left| 4\sigma_0^2 \psi_{\varepsilon}'(|E_t^{\delta}|^2) \pi_{\delta}^2(E_t^{\delta}) + 8\sigma_0^2 \psi_{\varepsilon}''(|E_t^{\delta}|^2) |E_t^{\delta}|^2 \pi_{\delta}(E_t^{\delta}) \right| \\ \leqslant \left| \pi_{\delta}^2(E_t^{\delta}) \sigma_0^2 \left(4\psi_{\varepsilon}'(|E_t^{\delta}|^2) + 8\psi_{\varepsilon}''(|E_t^{\delta}|^2) |E_t^{\delta}|^2 \right) \right|. \end{aligned}$$

Moreover, we know that $\psi'_{\varepsilon}(r^2) + 2\psi''_{\varepsilon}(r^2)r^2 \leq r^{-3}$, for all r > 0 and $\varepsilon \leq 1$. Using the presence of π_{δ} , we have that the integrand is null near 0. Then, we can once again apply dominated convergence theorem and obtain

(65)
$$\lim_{\varepsilon \to 0} \int_0^{\iota} 4\left\{\sigma_0^2 \psi_{\varepsilon}'(|E_s^{\delta}|^2) \pi_{\delta}(E_s^{\delta})^2 + 8\sigma_0^2 \psi_{\varepsilon}''(|E_s^{\delta}|^2) |E_s^{\delta}|^2 \psi_{\delta}(E_s^{\delta})\right\} \mathrm{d}s = 0.$$

Finally, combining Equations (61), (62) (63) and (65), we get the expected result.

A.2. Proof of Equation (55).

Proof. To prove Equation, let us introduce the following technical Lemma:

Lemma 17. For any positive and differentiable function $u : [0, +\infty) \rightarrow [0, +\infty)$ such that

$$\frac{\mathrm{d}}{\mathrm{d}t}u(t) \leqslant 2k(t)\sqrt{u(t)},$$

for some function $k : [0, +\infty) \rightarrow \mathbb{R}$, we have

$$\frac{\mathrm{d}}{\mathrm{d}t}\sqrt{u(t)} \leqslant k(t).$$

Proof of Lemma 17. We consider for any $\varepsilon > 0$ the function $\psi_{\varepsilon} : [0, +\infty) \ni r \mapsto (r + \varepsilon)^{1/2}$. The function ψ_{ε} being differentiable with positive derivative, we get

$$\frac{\mathrm{d}}{\mathrm{d}t}\psi_{\varepsilon}\left(u(t)\right) \leqslant 2k(t)\sqrt{u(t)}\psi_{\varepsilon}'\left(u(t)\right).$$

Using the fact that $2r\psi_{\varepsilon}'(r^2) = r(r^2 + \varepsilon)^{-1/2} \leq 1$, we use dominated convergence theorem to conclude the proof.

Such a lemma is classical and applying it to Equation (54) concludes the proof of Equation (55). \Box

FRANÇOIS DELARUE, UNIVERSITÉ CÔTE D'AZUR, LABORATOIRE J.A.DIEUDONNÉ, 06108 NICE, FRANCE *Email address*: francois.delarue@univ-cotedazur.fr

RAPHAËL MAILLET, UNIVERSITÉ PARIS-DAUPHINE & PSL, CNRS, CEREMADE, 75016 PARIS, FRANCE *Email address*: maillet@ceremade.dauphine.fr

ETIENNE TANRÉ, UNIVERSITÉ CÔTE D'AZUR, INRIA, CNRS, LJAD, FRANCE *Email address*: Etienne.Tanre@inria.fr