

Quantum Chromodynamics Resolution of the ATOMKI Anomaly in ^4He Nuclear Transitions

Valery Kubarovsky

Thomas Jefferson National Accelerator Laboratory, Newport News, VA 23606, USA

Jennifer Rittenhouse West

*Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA and
EIC Center at Thomas Jefferson National Accelerator Laboratory, Newport News, VA 23606, USA*

Stanley J. Brodsky

SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94309, USA

Observations of anomalous angular correlations in electron-positron pairs produced from excited states of ^4He , ^8Be and ^{12}C nuclei have been suggested as due to the creation and subsequent decay of a new light particle of mass ~ 17 MeV. In this work, we investigate the possibility that the source of the observed signals is a set of new excitation channels created by the 12-quark hidden-color Fock state within the ^4He nuclear wavefunction dubbed the “hexadiquark.” We calculate the invariant e^+e^- mass spectrum for the electromagnetic transition from a new excitation of ^4He , estimating its differential and total decay width. We find that we can fit the shape of the anomalous signal with the QCD Fock state at excitation energy $E^* = 17.9 \pm 1$ MeV and a Gaussian form factor for the electromagnetic decay. We address the physical issues with the fit parameters using properties of the hexadiquark state, in particular the three weakly repulsive 6_C interactions of $\text{SU}(3)_C$ between diquark pairs. Experimental tests of our model are described in detail. In light of this work, we emphasize the need for independent experimental confirmation or refutation of the ATOMKI results as well as dedicated experiments to search for the proposed new excitations of ^4He and other α -cluster nuclei.

I. Introduction

Observations by the ATOMKI collaboration of anomalous angular correlations in electron-positron pairs produced in the nuclear decays $^4\text{He}^* \rightarrow ^4\text{He} + e^+e^-$ [1, 2], $^8\text{Be}^* \rightarrow ^8\text{Be} + e^+e^-$ [3] and $^{12}\text{C}^* \rightarrow ^{12}\text{C} + e^+e^-$ [4] have been attributed to the creation and subsequent decay of a new light particle of mass of ~ 17 MeV, dubbed the X17. The same group has reported observations of the lepton pair in the off-resonance region of $^7\text{Li}(p, e^+e^-)^8\text{Be}$ direct proton-capture reactions [5]. The signals have generated a great deal of theoretical interest in both the particle and nuclear physics communities [6–29]. In this work, we focus on the ^4He experiment in which the observed invariant mass $m_{e^+e^-}$ of the lepton pair was found to be $m_X = 16.94 \pm 0.12(\text{stat.}) \pm 0.21(\text{syst.})$ MeV. We show that excitation and decay of the recently proposed QCD hidden-color Fock state within the ^4He nuclear wavefunction [30] provides a viable and compelling explanation of the ATOMKI phenomenon as well as a clear set of experimental predictions. The model acts in all $A \geq 4$ nuclei and explains why the X17 signal has only been observed in α -nuclei.

The light-front Fock state expansion of QCD has led to new perspectives for the nonperturbative eigenstructure of hadrons. The $[ud]$ scalar diquark in the $\bar{3}_C$ representation of $\text{SU}(3)_C$, for example, is a configuration that appears to play a fundamental role in hadron spectroscopy such as the $|u[ud]\rangle$ Fock state in the proton wavefunction and baryonic Regge trajectories [31, 32]. Tetraquarks are very likely bound states of diquarks and anti-diquarks [33–36]. Quark-antiquark, quark-diquark, and diquark-antidiquark bound states all have an identical $3_C - \bar{3}_C$ color-confining interaction and therefore matching spectroscopy in QCD [37].

The 12-quark (6-diquark) hidden-color state $[[ud][ud][ud][ud][ud][ud]]$ dubbed the “hexadiquark” is a novel color-singlet of QCD [30]. Such hidden-color states are predictions of the $\text{SU}(3)_C$ group theory basis of QCD and have been studied for over four decades [38–43]. In general, a hadronic or nuclear eigenstate of the QCD Hamiltonian is a sum over all color-singlet Fock states which match its quantum numbers. This has a special impact for the ^4He nucleus, since the hexadiquark Fock state is an unusually low mass state with the same quantum numbers as the nuclear Fock state $|nnpp\rangle$. Nuclear wavefunctions are typically dominated by the neutron and proton Fock state, i.e., the multiple (A) 3-quark color-singlet Fock state. In the ^4He nuclear wavefunction, this is the $|nnpp\rangle$ state containing A=4 color singlets (2 protons and 2 neutrons). In addition to this state, every 12-quark combination with the same quantum numbers of ^4He is also in the wavefunction - including the single color-singlet hexadiquark. The hexadiquark is a special hidden-color state because it is an unusually low mass combination of quarks - all spin-0, isospin-0 and S-wave combinations - that obeys the spin-statistics theorem at every stage of the build. This means it has the opportunity to have a larger effect on nuclear physics than is typical. The six scalar diquarks within the hexadiquark hidden-color

state have predicted experimental signatures in diffractive dissociation of ^4He nuclei, as recently published [44].

QCD also predicts orbital and radial excitations between the $[ud]$ diquark components of the hexadiquark and therefore new excitations of ^4He beyond the standard excitations predicted by nucleonic degrees of freedom. The excitation energy of the hexadiquark can be below the energy required to produce nuclear decays such as $^4\text{He} \rightarrow p + ^3\text{H}$ and have evaded detection thus far while containing new experimental predictions as will be demonstrated. The first hints of an excited state of ^4He were discovered over 80 years ago [45, 46] and the question of how a new subdominant excited state could have been missed must be addressed. Hidden-color states cannot decay through conventional nuclear channels as all of the constituents have color charge (specifically quarks, diquarks and duo-diquarks). The weakly repulsive QCD bonds between diquarks, utilizing the symmetric sextet 6_C channel of $\text{SU}(3)_C$, allow for low (nuclear) energy excitations and require dedicated searches for their signatures. An exhaustive list of QCD decay channels and experimental signatures is presented in Section IV.

II. Fitting the nuclear decay amplitude for $^4\text{He}^* \rightarrow ^4\text{He} + e^+e^-$

The signal template we use to match the ATOMKI e^+e^- angular correlation is a narrow Gaussian peak in the electron-positron invariant mass with a mean value ~ 17 MeV and width $\sigma \sim 0.7$ MeV (see Figure 1) based on unpublished work by the ATOMKI collaboration [2] and cross-checked with the NA64 X17 invariant mass simulation [26]. We calculate the invariant mass $m_{e^+e^-}$ spectrum in the electromagnetic transition of an excited state of ^4He , $^4\text{He}^*(E^*) \rightarrow ^4\text{He} + e^+e^-$, where E^* is the excited state energy. The mass difference between the first known excited state and the ground state of the helium nucleus is 20.21 MeV [47] and E^* is initially fixed to this value. Its decay to the ^4He ground state via a virtual photon cannot explain the narrow width of the observed X17 signal. The value of E^* is subsequently allowed to float to a different value corresponding to the proposed excitation of the 12-quark color-singlet hexadiquark Fock state in the ^4He nuclear wavefunction [30].




FIG. 1. Fit to the ATOMKI data (in red) by a Gaussian function and background distribution (in black) of the $m_{e^+e^-}$ invariant mass distribution based on [2] and cross-checked with [26]. The mean value of the fitted mass is ~ 17 MeV with width $\sigma \sim 0.7$ MeV. The signal alone is shown in blue.

The invariant mass distribution of the lepton pair $m_{e^+e^-}$ from the decay of excited states of ^4He has not been calculated until now. The result of our calculation shows that decay due to an excitation of the hexadiquark Fock state in the eigensolution of ^4He can describe the signal and its narrow width; in fact, this signal is to be expected from the existence of the new hidden-color QCD state.

We begin with the square of the amplitude for the electromagnetic decay $^4\text{He}^* \rightarrow ^4\text{He} + e^+e^-$, given by [15]

$$|\mathcal{M}|^2 = \sum_{\text{spin}} |M|^2 = \frac{e^4 C_{E0}^2}{\Lambda^4} [2(p_+ \cdot p_0)(p_- \cdot p_0) - m_0^2(p_+ \cdot p_-) - m_e^2 m_0^2], \quad (1)$$

where p_+ and p_- are the 4-momenta of the positron and electron respectively, p_0 and m_0 are the 4-momentum and

mass of the ground state of ^4He and m_e is the electron mass. Final state spins have been summed over. The definitions of Λ as the nuclear energy scale and C_{E0}^2 as the Wilson coefficient of the decay operator are contained in [15]. These constants are not relevant for the analysis carried out in this work as they affect only the normalization which we match to the signal.

The matrix element depends on only two kinematical variables, q^2 and $\cos \theta^*$, where $q = p_+ + p_-$ is the 4-momentum of the virtual photon, $q^2 = m_{e^+e^-}^2$ is the e^+e^- invariant mass squared and θ^* is the electron angle in the virtual photon center of mass system. Rewriting Eq.1 in terms of these variables gives

$$|\mathcal{M}(q^2, \cos \theta^*)|^2 = \frac{\alpha^2 C_{E0}^2 (4\pi)^2}{\Lambda^4 8q^2} [q^2 - \cos^2 \theta^* (q^2 - 4m_e^2)] [m^4 + (m_0^2 - q^2)^2 - 2m(m_0^2 + q^2)] \quad (2)$$

where m is the mass of the excited $^4\text{He}^*$ state.

The differential width $d\Gamma$ for the three body decay is given by

$$d\Gamma = \frac{1}{(2\pi)^5} \frac{1}{16m^2} |\mathcal{M}|^2 |p_e^*| |p_0| dm_{e^+e^-} d\Omega_e^* d\Omega_0 \quad (3)$$

where $d\Omega = d\phi d\cos \theta$, $(|p_e^*|, \Omega_e^*)$ is the electron momentum in the center of mass of the electron-positron pair, $(|p_0|, \Omega_0)$ is the ^4He momentum in the rest frame of the $^4\text{He}^*$. The minimum value of the $m_{e^+e^-}$ invariant mass is $2m_e$ and maximum value is determined by the mass difference between the excited and ground ^4He states, $\Delta m = 20.21$ MeV.

The integration of the matrix element $|\mathcal{M}(q^2, \cos \theta^*)|^2$ over angles $d\Omega_e^* d\Omega_0$ gives

$$|\mathcal{M}_q(q^2)|^2 = \int |\mathcal{M}(q^2, \cos \theta^*)|^2 d\Omega_e^* d\Omega_0 = \frac{\alpha^2 C_{E0}^2 (4\pi)^4}{\Lambda^4 12q^2} (q^2 + 2m_e^2) [m^4 + (m_0^2 - q^2)^2 - 2m^2(m_0^2 + q^2)]. \quad (4)$$

The matrix element $|\mathcal{M}_q(q^2)|^2$ simplifies in the limit $m_e = 0$, to

$$|\mathcal{M}_q(q^2)|^2 = \frac{\alpha^2 C_{E0}^2 (4\pi)^4}{\Lambda^4 12} [m^4 + (m_0^2 - q^2)^2 - 2m^2(m_0^2 + q^2)]. \quad (5)$$

There is only one variable left, the invariant electron-positron mass $m_{e^+e^-} = \sqrt{q^2} = \sqrt{(p_+ + p_-)^2}$, and we find

$$d\Gamma = \frac{1}{(2\pi)^5} \frac{1}{16m^2} |\mathcal{M}_q|^2 |p_e^*| |p_0| dm_{e^+e^-}. \quad (6)$$

The numerical integration over the three-body phase space is

$$\Gamma_{E0} = \int_{2m_e}^{m-m_0} dm_{e^+e^-} |\mathcal{M}_q(q^2)|^2 |p_e^*| |p_{He}| = \frac{\alpha^2 C_{E0}^2}{\Lambda^4} \frac{\Delta m^5}{60\pi} 0.982, \quad (7)$$

which coincides with the approximate solution (Eq. 9, below) to an accuracy of 2%.

In the approximation $q^2/m^2 < \Delta m^2/m^2 = 3 \cdot 10^{-5} \ll 1$ and negligible electron mass, the differential width $d\Gamma/dm_{e^+e^-}$ is

$$\frac{d\Gamma}{dm_{e^+e^-}} = \frac{\alpha^2 C_{E0}^2}{12\pi\Lambda^4} m_{e^+e^-} (\Delta m^2 - m_{e^+e^-}^2)^{\frac{3}{2}}. \quad (8)$$

The full width has an analytic solution in this case,

$$\begin{aligned} \Gamma_{E0} &= \frac{\alpha^2 C_{E0}^2}{12\pi\Lambda^4} \int_{2m_e}^{m-m_0} m_{e^+e^-} (\Delta m^2 - m_{e^+e^-}^2)^{\frac{3}{2}} dm_{e^+e^-} \\ &= \frac{\alpha^2 C_{E0}^2}{\Lambda^4} \frac{(\Delta m^2 - 4m_e^2)^{\frac{5}{2}}}{60\pi} \approx \frac{\alpha^2 C_{E0}^2}{\Lambda^4} \frac{\Delta m^5}{60\pi} \end{aligned} \quad (9)$$

that with 2% accuracy corresponds to the exact solution of Eq. 7.

The differential width $d\Gamma/dm_{e^+e^-}$ from Eq. 2, integrated over $\cos\theta^*$ and normalized to one (shown as the blue curve on the left panel of Figure 2) is very smooth from the minimum value of $m_{e^+e^-} = 2m_e$ up to the maximum of $\Delta m = 20.21$ MeV. The ATOMKI signal cannot arise from the straightforward analysis carried out in this section.

III. Electromagnetic transition form factors $F(q^2)$ in the nuclear decay ${}^4\text{He}^* \rightarrow {}^4\text{He} + e^+e^-$

The analysis carried out in the previous section implicitly assumed a form factor $F(q^2)$ equal to unity. A more realistic form factor significantly changes the $m_{e^+e^-}$ invariant mass distribution in the nuclear decay ${}^4\text{He}^* \rightarrow {}^4\text{He} + e^+e^-$

$$\frac{d\Gamma}{dm_{e^+e^-}} = \frac{\alpha^2 C_{E0}^2}{12\pi\Lambda^4} m_{e^+e^-} (\Delta m^2 - m_{e^+e^-}^2)^{\frac{3}{2}} |F(q^2)|^2. \quad (10)$$

We show the best fit form factor and its failure to describe the signal in this section, as well as the successful fit for a new excitation energy near 18 MeV.

The experimental $m_{e^+e^-}$ distribution is not given in the ATOMKI work. In order to compare the theoretical analysis in this section to the signal, we first extract the ATOMKI data from [2], graph them and confirm that we can replicate their plot. We then use this as a template to fit our signal, noting that it is based on the unpublished ATOMKI work [2] and cross-checked with the invariant mass plot in the NA64 Collaboration X17 simulations [26]. In the next step, we use the extracted signal and background distributions from the template and make a fit with two functions to describe both distributions. Figure 1 presents our fit of ATOMKI data (shown in red). The background (shown in black) was fitted independently because the experiment measured the background distribution with much higher accuracy than the signal. We used a third-degree polynomial to fit the background. The fit result is $f_{\text{bg}}(x) = 1584 - 306x + 20x^2 - 0.434x^3$ (where $x = m_{e^+e^-}$) with $\chi^2/ndf = 0.99$. The same function was used in all fits in Figure 1, Figure 2 and Figure 3. The signal in Figure 1 was described by a Gaussian function with 3 parameters: the mass value of the peak position (16.8 ± 0.23 MeV), the width ($\sigma = 0.70 \pm 0.17$ MeV) and overall normalization. This fit is very close to the signal template. We now begin to model the results theoretically.

Several realistic form factor models are tested in our attempt to model the data template. We compare the shape of the differential width $d\Gamma/dm_{e^+e^-}$ for the electromagnetic transition with form factor $F(q^2) \propto \exp(q^2/\lambda^2)$ with $\lambda = 14.5$ MeV (see Eq. 10) with a maximum near 17 MeV and ATOMKI anomaly shown on the left panel of Figure 2 and including $F(q^2) = 1$ for comparison. We find the width of the distribution to be too large to obtain a satisfactory description of the experimental distribution. The signal template is too narrow in comparison with the theoretical curve as shown in Figure 2, comparing the red curve (theoretical curve) and black curve (signal template). On the right panel of Figure 2 we present the attempt to fit the signal template, again with electromagnetic transition form factors $F(q^2) \propto \exp(q^2/\lambda^2)$. The background function is the same as in Figure 1. This fit has only two parameters, λ and the normalization of the theoretical function Eq. 10. The fit is not satisfactory. A form factor of $F(q^2) \propto q^2$ also failed to describe the signal due to its large width.

We now introduce a new parameter, the mass of a new excited ${}^4\text{He}^*$ state, into our model. We motivate this state physically by the 12-quark color singlet Fock state model of ${}^4\text{He}$ nucleus in which the ${}^4\text{He}$ wavefunction is dominated by a linear combination of $|nnpp\rangle$ and the 12-quark hexadiquark QCD state [30]. The result of the fit with form factor $F(q^2) \propto \exp(q^2/\lambda^2)$ is presented in Figure 3. There are three parameters of the fit, the excitation energy of the new ${}^4\text{He}^*$ state $E^* = 17.9 \pm 1$ MeV ($= \Delta m$ in Eq. 10), the form factor parameter $\lambda = 6.2 \pm 0.5$ MeV and the overall normalization. The model describes the data template very well. Our parameter λ is close to the momentum scale found in earlier work on nuclear transitions [16].

We have reproduced the ATOMKI signal with the introduction of a new excitation of ${}^4\text{He}$ but there are serious physical issues that must be addressed. First, the energy of the electron-positron pair in this case has to be close to 17.9 MeV. This is at the lower limit of the selection of the positrons and electrons with total energy in the region (18, 22) MeV and 2.6σ from the central excitation energy 20.49 MeV created by the 900 keV proton beam. There are no error bars given by ATOMKI but a reasonable visual estimate is 0.9 MeV which would reduce our σ values. These are not prohibitively large deviations and we do not discuss them further but they must be noted. More importantly, this is an excitation energy that has never been observed in a decay to a conventional hadronic or nuclear final state. In addition, the form factor parameter λ is an order of magnitude smaller than expected for a typical nuclear transition. We address both of the latter points in the following section.




FIG. 2. *Left panel:* Differential width $d\Gamma/dm_{e^+e^-}$ (in arbitrary units) for the transition ${}^4\text{He}^*(20.21) \rightarrow {}^4\text{He} + e^+e^-$ as a function of the electron-positron invariant mass $m_{e^+e^-}$. Blue curve has form factor $F(q^2) = 1$ (see Eq. 10). The black curve is the Gaussian distribution of the ATOMKI anomaly with $m = 17$ MeV and $\sigma = 0.7$ MeV. Red curve has form factor $F(q^2) \propto \exp(q^2/\lambda^2)$ with $\lambda = 14.5$ MeV. With this value of the λ parameter, the distribution maximum is at $m = 17$ MeV. The functions are normalized such that the peak values are the same for all curves. *Right panel:* Fit to signal template (event counts vs. invariant mass distribution) for the transition ${}^4\text{He}^*(20.21) \rightarrow {}^4\text{He} + e^+e^-$. The fit is an electromagnetic transition with form factor $F(q^2) \propto \exp(q^2/\lambda^2)$ (in red). The background distribution is shown in black, the signal alone is shown in blue. The fit is not satisfactory.




FIG. 3. Fit to signal template (event counts vs. invariant mass distribution) for the electromagnetic transition $\text{HdQ}^*(17.9) \rightarrow {}^4\text{He} + (e^+e^-)$. The fit assumes a form factor $F(q^2) \propto \exp(q^2/\lambda^2)$ with the form factor parameter $\lambda = 6.2$ MeV (in red). The background distribution is shown in black, and the signal alone is shown in blue (see Eq. 10 with $\Delta m = 17.9$ MeV).

IV. QCD Hexadiquark excitations

The hexadiquark (HdQ) is a hidden-color QCD state [38–43] in the ${}^4\text{He}$ nuclear wavefunction consisting of six scalar $[ud]$ diquarks [48, 49] in a color singlet configuration [30]. Excitation of a subset of diquarks, either radial or angular momentum excitations, gives rise to nuclear transitions with unconventional decays. In conventional nuclear physics, for example, the electron scattering experiment $e + {}^4\text{He} \rightarrow e' + X$ will produce hadronic excitations of ${}^4\text{He}$. In contrast, if an electron (or proton) scatters on the hexadiquark Fock state of ${}^4\text{He}$, it can produce orbital or radial excitations between any two of the $[ud]$ diquarks. Typically such excitations would be at energies near 300 MeV above the ground state, in analogy to the N^* and Δ excited states of nucleons. Excitations of single diquarks from spin-0

isospin-0 to spin-1 isospin-1 states have energies of approximately 200 MeV [48, 49].

However, the HdQ is an unusual QCD eigenstate because, in addition to being a low mass spin-0 isospin-0 S-wave hidden-color state, the wavefunction contains three symmetric and *repulsive* $3_C \times 3_C \rightarrow \bar{6}_C$ interactions. These interactions are required by the spin-statistics constraints upon clustering pairs of $[ud]$ diquarks into each of the three $[ud][ud]$ duo-diquarks contained within the 12-quark color-singlet wavefunction. While the final construction of the three duo-diquarks into the HdQ is strongly bound with color factor $C_F = -5$, the QCD interaction between diquarks in a duo-diquark is repulsive which significantly lowers the excitation energies of the HdQ and increases the size of the HdQ.

There are two ways to excite the 6_C bond between diquarks: an orbital angular momentum $L = 1$ excitation and an $L = 0$ radial excitation. These are color-singlet excitations, they do not break the strongly bound HdQ state apart, and therefore cannot decay to a conventional nuclear physics state such as $p + {}^3\text{H}$ or $n + {}^3\text{He}$. However, since the $L = 1$ excitation has $J^P = 1^-$, the HdQ excitation can decay to the ${}^4\text{He}$ ground state by the emission of a virtual photon which makes an e^+e^- pair: ${}^4\text{He}^*(17.9) \rightarrow {}^4\text{He} + \gamma^* \rightarrow {}^4\text{He} + e^+e^-$. This state can also emit a 17.9 ± 1 MeV real photon. The $L = 0$ radial excitation between diquarks has $J^P = 0^+$ and can only decay via virtual photon which creates the e^+e^- pair. Thus the three decay modes of the HdQ predict an e^+e^- pair of invariant mass 17.9 ± 1 MeV and/or a 17.9 ± 1 MeV gamma ray photon.

In the case of an excitation of the ${}^4\text{He}$ nucleus to a known excited state, there are two additional decay modes with distinct signatures. In the first scenario, the transition is described by the 2-stage process

$$p + {}^3\text{H} \rightarrow {}^4\text{He}^*(20.21) \text{ MeV} \rightarrow {}^4\text{He}^*(17.9) \text{ MeV} + \gamma(2.31 \pm 1 \text{ MeV}) \rightarrow {}^4\text{He} + (e^+e^-) + \gamma(2.31 \pm 1 \text{ MeV}) \quad (11)$$

for the first excited state of ${}^4\text{He}$ at 20.21 MeV ($\Gamma = 0.50$ MeV). The second scenario occurs for the second excited state of ${}^4\text{He}$ at 21.01 MeV ($\Gamma = 0.84$ MeV),

$$p + {}^3\text{H} \rightarrow {}^4\text{He}^*(21.01) \text{ MeV} \rightarrow {}^4\text{He}^*(17.9) \text{ MeV} + \gamma(3.11 \pm 1 \text{ MeV}) \rightarrow {}^4\text{He} + (e^+e^-) + \gamma(3.11 \pm 1 \text{ MeV}) \quad (12)$$

It is important to note that multiple processes can excite the HdQ state, including direct proton capture in the initial $p + {}^3\text{H}$ collision creating $\gamma + \text{HdQ}^*$ and then decaying down to $e^+e^- + {}^4\text{He}$, *i.e.*,

$$p + {}^3\text{H} \rightarrow \gamma + {}^4\text{He}^*(17.9) \rightarrow {}^4\text{He} + e^+e^- + \gamma. \quad (13)$$

A low energy photon will also be present in the final state, however, the signature of this decay is the e^+e^- pair of invariant mass 17.9 ± 1 MeV. In addition to the decay signatures outlined above (summarized in Tables I and II), a key method for confirmation of the hexadiquark solution to the ATOMKI puzzle is an investigation of the first excited state of ${}^4\text{He}$ to identify a secondary “shoulder” peak within the range of 17.9 ± 1 MeV. Since the hexadiquark contains three sets of duo-diquarks, the 17.9 ± 1 MeV excitation will be accompanied by excitations at ~ 36 MeV and ~ 54 MeV. Larger α -nuclei, including ${}^8\text{Be}$ and ${}^{12}\text{C}$, will have even higher multiples of the 17.9 ± 1 MeV excitation for each additional hexadiquark state within the nuclear wavefunction.

We propose that the HdQ is the origin of a previously missed subdominant ${}^4\text{He}^*(17.9)$ excited state. It can be observed in the interaction $e + {}^4\text{He} \rightarrow e' + {}^4\text{He}^*(17.9) \rightarrow e' + {}^4\text{He} + [e^+e^-]$. The characteristic mass scale of the transition form factor $F(q^2)$ producing the timelike lepton pair in the $\text{HdQ}^* \rightarrow \text{HdQ} + \gamma^*$ process can be significantly lighter than the standard QCD mass scale because it reflects the large size structure of the HdQ due to the repulsive interactions within each of the three duo-diquarks. This can cause a λ parameter of order 10 MeV in the transition $\text{HdQ}^* \rightarrow \text{HdQ}$ form factor as required by our fit. The conclusion from these arguments is that an excitation of the HdQ Fock state of ${}^4\text{He}$ can account for the excellent fit and parameters in Figure 3 to the ATOMKI signal template.

We note that the hexadiquark state is predicted to exist as a hidden-color QCD Fock state within every nucleus containing an alpha particle. Each such nucleus will also have HdQ hidden color excitations; the resulting HdQ^* decays should therefore also be observed in lepton pair decays from excitations of ${}^8\text{Be}$ and ${}^{12}\text{C}$. All nuclei containing at least one α -particle will have this excitation and the strength of the signal is predicted to increase with each additional α -particle. HdQ transitions cannot occur in $A = 3$ nuclei, in deuterium, or in nucleons (all of which contain less than 12 quarks) and therefore will not be observed in experiments that use these targets. However, other hidden-color Fock states in $A < 4$ nuclei that use the repulsive 6_C representation of $\text{SU}(3)_C$ also allow for new lower energy excitations with unconventional decays.

There are important complementary tests for the hexadiquark solution. The HdQ component of ${}^4\text{He}$ should be observed as a peak in the missing mass in $e + {}^4\text{He} \rightarrow e' + X$. The diquark composition of the ${}^4\text{He}$ nucleus may be observed via $e + {}^4\text{He} \rightarrow e' + [ud] + X$ where the $[ud]$ -diquark jet [50] is produced opposite to the scattered electron.

New Hexadiquark excitation ${}^4\text{He}^*(17.9 \pm 1)$	Decay from ${}^4\text{He}^*(17.9 \pm 1) \rightarrow$	Experimental signature
L=0 radial excitations $[ud] \leftrightarrow [ud]$	${}^4\text{He} + \gamma^* \rightarrow {}^4\text{He} + e^+e^-$	$e^+e^- (17.9 \pm 1)$
L=1 orbital excitations $[ud] \odot [ud]$	${}^4\text{He} + \gamma$ ${}^4\text{He} + \gamma \rightarrow {}^4\text{He} + e^+e^-$	$\gamma(17.9 \pm 1)$ $e^+e^- (17.9 \pm 1)$

TABLE I. Experimental signatures for direct excitations of the QCD hexadiquark Fock state in units of MeV. Each hexadiquark contains 3 diquark-diquark correlations and therefore subdominant nuclear excitations near 36 MeV and 54 MeV are predicted in ${}^4\text{He}$, ${}^8\text{Be}$, ${}^{12}\text{C}$ and all higher A α -cluster nuclei will have additional excitations at multiples of 17.9 ± 1 MeV, exactly 3 additional excitations per α .

Excitation energy of ${}^4\text{He}$	Decay channel ${}^4\text{He}^*(E^*) \rightarrow$	Experimental signature
${}^4\text{He}^*(20.21)$	${}^4\text{He}^*(17.9 \pm 1) + \gamma(2.31 \pm 1) \rightarrow {}^4\text{He} + \gamma(2.31 \pm 1) + e^+e^-$ ${}^4\text{He}^*(17.9 \pm 1) + \gamma(2.31 \pm 1) \rightarrow {}^4\text{He} + \gamma(17.9 \pm 1)$	$e^+e^- (17.9 \pm 1), \gamma(2.31 \pm 1)$ $\gamma(17.9 \pm 1), \gamma(2.31 \pm 1)$
${}^4\text{He}^*(21.01)$	${}^4\text{He}^*(17.9 \pm 1) + \gamma(3.11 \pm 1) \rightarrow {}^4\text{He} + \gamma(3.11 \pm 1) + e^+e^-$ ${}^4\text{He}^*(17.9 \pm 1) + \gamma(3.11 \pm 1) \rightarrow {}^4\text{He} + \gamma(3.11 \pm 1) + \gamma(17.9 \pm 1)$	$e^+e^- (17.9 \pm 1), \gamma(3.11 \pm 1)$ $\gamma(17.9 \pm 1), \gamma(3.11)$

TABLE II. Experimental signatures for 2-stage excitations in which excitation of a known ${}^4\text{He}$ excited state is followed by a decay to the proposed new hexadiquark excitation, ${}^4\text{He}^* \rightarrow \text{He}^*(17.9) \rightarrow X$, in units of MeV.

V. Summary

We have presented a QCD-based solution to the ATOMKI anomalies that utilizes the predicted hexadiquark hidden-color Fock state in the ${}^4\text{He}$ nuclear wavefunction [30]. The hexadiquark state mixes with the conventional $|nnpp\rangle$ nuclear physics state of ${}^4\text{He}$ and offers a hidden-color explanation of its anomalously large binding energy. Our calculation of the $m_{e^+e^-}$ invariant mass spectrum in the electromagnetic transition ${}^4\text{He}^*(E^*) \rightarrow {}^4\text{He} + e^+e^-$ as well as the observed differential and total width of the decay are fit to an electromagnetic decay of a new excitation of ${}^4\text{He}$ of energy $E^* = 17.9 \pm 1$ MeV above the ground state. The cause of ${}^4\text{He}^*(17.9)$ is proposed to be low energy diquark-diquark pair excitations within the hidden-color hexadiquark component of the nuclear wavefunction, enabled by the novel spin-statistics mandated use of weakly repulsive 6_C interactions of $\text{SU}(3)_C$ between diquark pairs.

We have shown that it is possible to reproduce the ATOMKI signal with $J^P = 0^+$ and $J^P = 1^-$ excitations between the diquark constituents of the hexadiquark Fock state. In one production scenario, the proton + tritium collision first produces the conventional first nuclear excited state ${}^4\text{He}^*$ which decays by photon emission to an excitation of the QCD Fock state. The hexadiquark excitation may be produced by direct proton capture as well. All experimental signatures and decay channels are summarized in Tables I and II.

This work is both timely and important due to the negative searches for the new X17 boson carried out thus far [26, 51, 52] and planned searches at Jefferson Lab [29], in Italy with the Positron Annihilation to Dark Matter Experiment (PADME [53, 54]) and elsewhere. Our QCD-based solution of the ATOMKI results, which must occur in all $A \geq 4$ nuclei including the ${}^4\text{He}$, ${}^8\text{Be}$, ${}^{12}\text{C}$ nuclei in the ATOMKI experiments, is in contrast to the new particle explanation.

Acknowledgements

J.R.W. acknowledges support by the LDRD program of LBNL, the EIC Center at Jefferson Lab and the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under contract number DE-AC02-05CH11231. V.K. acknowledges support by the U.S. Department of Energy, Office of Science, and Office of Nuclear Physics under contracts DE-AC05-06OR23177. JLAB-PHY-22-3641. The work of S.J.B. was supported in part by the Department of Energy under Contract No. DE-AC02-76SF00515. SLAC-PUB-17683.

-
- [1] A. J. Krasznahorkay, M. Csatlós, L. Csige, J. Gulyás, A. Krasznahorkay, B. M. Nyakó, I. Rajta, J. Timár, I. Vajda, N. J. Sas, New anomaly observed in He4 supports the existence of the hypothetical X17 particle, Phys. Rev. C 104 (2021) 044003. doi:10.1103/PhysRevC.104.044003. [arXiv:2104.10075](#).
 - [2] A. J. Krasznahorkay, et al., New evidence supporting the existence of the hypothetic X17 particle (2019). [arXiv:1910.10459](#).

- [3] A. J. Krasznahorkay, et al., Observation of Anomalous Internal Pair Creation in Be8 : A Possible Indication of a Light, Neutral Boson, *Phys. Rev. Lett.* 116 (2016) 042501. doi:doi:10.1103/PhysRevLett.116.042501. [arXiv:1504.01527](#).
- [4] A. J. Krasznahorkay, et al., New anomaly observed in C12 supports the existence and the vector character of the hypothetical X17 boson, *Phys. Rev. C* 106 (2022) L061601. doi:doi:10.1103/PhysRevC.106.L061601. [arXiv:2209.10795](#).
- [5] N. J. Sas, et al., Observation of the X17 anomaly in the ${}^7\text{Li}(p, e^+e^-){}^8\text{Be}$ direct proton-capture reaction (2022). [arXiv:2205.07744](#).
- [6] M. Battaglieri, et al., US Cosmic Visions: New Ideas in Dark Matter 2017: Community Report, in: U.S. Cosmic Visions: New Ideas in Dark Matter, 2017. [arXiv:1707.04591](#).
- [7] J. Backens, M. Vanderhaeghen, X17 Discovery Potential in the $\gamma\text{N}\rightarrow\text{e}^+\text{e}^-\text{N}$ Process at Electron Scattering Facilities, *Phys. Rev. Lett.* 128 (2022) 091802. doi:doi:10.1103/PhysRevLett.128.091802. [arXiv:2110.06055](#).
- [8] B. Fornal, Is There a Sign of New Physics in Beryllium Transitions?, *Int. J. Mod. Phys. A* 32 (2017) 1730020. doi:doi:10.1142/S0217751X17300204. [arXiv:1707.09749](#).
- [9] J. L. Feng, B. Fornal, I. Galon, S. Gardner, J. Smolinsky, T. M. P. Tait, P. Tanedo, Particle physics models for the 17 MeV anomaly in beryllium nuclear decays, *Phys. Rev. D* 95 (2017) 035017. doi:doi:10.1103/PhysRevD.95.035017. [arXiv:1608.03591](#).
- [10] M. Viviani, E. Filandri, L. Girlanda, C. Gustavino, A. Kievsky, L. E. Marcucci, R. Schiavilla, X17 boson and the $\text{H3}(p, e^+e^-)\text{He4}$ and $\text{He3}(n, e^+e^-)\text{He4}$ processes: A theoretical analysis, *Phys. Rev. C* 105 (2022) 014001. doi:doi:10.1103/PhysRevC.105.014001. [arXiv:2104.07808](#).
- [11] P. Agrawal, et al., Feebly-interacting particles: FIPs 2020 workshop report, *Eur. Phys. J. C* 81 (2021) 1015. doi:doi:10.1140/epjc/s10052-021-09703-7. [arXiv:2102.12143](#).
- [12] J. Alexander, et al., Dark Sectors 2016 Workshop: Community Report, 2016. [arXiv:1608.08632](#).
- [13] J. Kozaczuk, D. E. Morrissey, S. R. Stroberg, Light axial vector bosons, nuclear transitions, and the ${}^8\text{Be}$ anomaly, *Phys. Rev. D* 95 (2017) 115024. doi:doi:10.1103/PhysRevD.95.115024. [arXiv:1612.01525](#).
- [14] T. Kitahara, Y. Yamamoto, Protophobic Light Vector Boson as a Mediator to the Dark Sector, *Phys. Rev. D* 95 (2017) 015008. doi:doi:10.1103/PhysRevD.95.015008. [arXiv:1609.01605](#).
- [15] J. L. Feng, T. M. P. Tait, C. B. Verhaaren, Dynamical Evidence For a Fifth Force Explanation of the ATOMKI Nuclear Anomalies, *Phys. Rev. D* 102 (2020) 036016. doi:doi:10.1103/PhysRevD.102.036016. [arXiv:2006.01151](#).
- [16] X. Zhang, G. A. Miller, Can nuclear physics explain the anomaly observed in the internal pair production in the Beryllium-8 nucleus?, *Phys. Lett. B* 773 (2017) 159–165. doi:doi:10.1016/j.physletb.2017.08.013. [arXiv:1703.04588](#).
- [17] X. Zhang, G. A. Miller, Can a protophobic vector boson explain the ATOMKI anomaly?, *Phys. Lett. B* 813 (2021) 136061. doi:doi:10.1016/j.physletb.2021.136061. [arXiv:2008.11288](#).
- [18] C. Frugiuele, E. Fuchs, G. Perez, M. Schlaffer, Constraining New Physics Models with Isotope Shift Spectroscopy, *Phys. Rev. D* 96 (2017) 015011. doi:doi:10.1103/PhysRevD.96.015011. [arXiv:1602.04822](#).
- [19] D. Banerjee, et al. (NA64), Search for a Hypothetical 16.7 MeV Gauge Boson and Dark Photons in the NA64 Experiment at CERN, *Phys. Rev. Lett.* 120 (2018) 231802. doi:doi:10.1103/PhysRevLett.120.231802. [arXiv:1803.07748](#).
- [20] G. L. Castro, N. Quintero, Tests of the Atomki anomaly in lepton pair decays of heavy mesons, *Phys. Rev. D* 103 (2021) 093002. doi:doi:10.1103/PhysRevD.103.093002. [arXiv:2101.01865](#).
- [21] J. A. Dror, R. Lasenby, M. Pospelov, New constraints on light vectors coupled to anomalous currents, *Phys. Rev. Lett.* 119 (2017) 141803. doi:doi:10.1103/PhysRevLett.119.141803. [arXiv:1705.06726](#).
- [22] D. S. M. Alves, N. Weiner, A viable QCD axion in the MeV mass range, *JHEP* 07 (2018) 092. doi:doi:10.1007/JHEP07(2018)092. [arXiv:1710.03764](#).
- [23] A. Ariga, et al. (FASER), FASER’s physics reach for long-lived particles, *Phys. Rev. D* 99 (2019) 095011. doi:doi:10.1103/PhysRevD.99.095011. [arXiv:1811.12522](#).
- [24] L. Delle Rose, S. Khalil, S. J. D. King, S. Moretti, New Physics Suggested by Atomki Anomaly, *Front. in Phys.* 7 (2019) 73. doi:doi:10.3389/fphy.2019.00073. [arXiv:1812.05497](#).
- [25] A. C. Hayes, J. L. Friar, G. M. Hale, G. T. Garvey, Angular correlations in the e^+e^- decay of excited states in Be8, *Phys. Rev. C* 105 (2022) 055502. doi:doi:10.1103/PhysRevC.105.055502. [arXiv:2106.06834](#).
- [26] E. Depero, et al. (NA64), Hunting down the X17 boson at the CERN SPS, *Eur. Phys. J. C* 80 (2020) 1159. doi:doi:10.1140/epjc/s10052-020-08725-x. [arXiv:2009.02756](#).
- [27] O. Seto, T. Shimomura, Atomki anomaly and dark matter in a radiative seesaw model with gauged $B - L$ symmetry, *Phys. Rev. D* 95 (2017) 095032. doi:doi:10.1103/PhysRevD.95.095032. [arXiv:1610.08112](#).
- [28] O. Seto, T. Shimomura, Atomki anomaly in gauged $U(1)_R$ symmetric model, *JHEP* 04 (2021) 025. doi:doi:10.1007/JHEP04(2021)025. [arXiv:2006.05497](#).
- [29] A. Ahmidouch, et al., A Direct Detection Search for Hidden Sector New Particles in the 3-60 MeV Mass Range (2021). [arXiv:2108.13276](#).
- [30] J. Rittenhouse West, S. J. Brodsky, G. F. de Teramond, A. S. Goldhaber, I. Schmidt, QCD hidden-color hexadiquark in the core of nuclei, *Nucl. Phys. A* 1007 (2021) 122134. doi:doi:10.1016/j.nuclphysa.2020.122134. [arXiv:2004.14659](#).
- [31] H. G. Dosch, G. F. de Teramond, S. J. Brodsky, Superconformal Baryon-Meson Symmetry and Light-Front Holographic QCD, *Phys. Rev. D* 91 (2015) 085016. doi:doi:10.1103/PhysRevD.91.085016. [arXiv:1501.00959](#).
- [32] G. F. de Teramond, H. G. Dosch, S. J. Brodsky, Baryon Spectrum from Superconformal Quantum Mechanics and its Light-Front Holographic Embedding, *Phys. Rev. D* 91 (2015) 045040. doi:doi:10.1103/PhysRevD.91.045040. [arXiv:1411.5243](#).
- [33] G. ’t Hooft, G. Isidori, L. Maiani, A. D. Polosa, V. Riquer, A Theory of Scalar Mesons, *Phys. Lett. B* 662 (2008) 424–430. doi:doi:10.1016/j.physletb.2008.03.036. [arXiv:0801.2288](#).

- [34] P. Masjuan, E. Ruiz Arriola, Regge trajectories of Excited Baryons, quark-diquark models and quark-hadron duality, *Phys. Rev. D* 96 (2017) 054006. doi:doi:10.1103/PhysRevD.96.054006. [arXiv:1707.05650](#).
- [35] M. Anselmino, E. Predazzi, S. Ekelin, S. Fredriksson, D. B. Lichtenberg, Diquarks, *Rev. Mod. Phys.* 65 (1993) 1199–1234. doi:doi:10.1103/RevModPhys.65.1199.
- [36] M. Nielsen, S. J. Brodsky, Hadronic superpartners from a superconformal and supersymmetric algebra, *Phys. Rev. D* 97 (2018) 114001. doi:doi:10.1103/PhysRevD.97.114001. [arXiv:1802.09652](#).
- [37] S. J. Brodsky, G. F. de Teramond, H. G. Dosch, J. Erlich, Light-Front Holographic QCD and Emerging Confinement, *Phys. Rept.* 584 (2015) 1–105. doi:doi:10.1016/j.physrep.2015.05.001. [arXiv:1407.8131](#).
- [38] S. J. Brodsky, B. T. Chertok, The Asymptotic Form-Factors of Hadrons and Nuclei and the Continuity of Particle and Nuclear Dynamics, *Phys. Rev. D* 14 (1976) 3003–3020. doi:doi:10.1103/PhysRevD.14.3003.
- [39] M. Harvey, Effective nuclear forces in the quark model with Delta and hidden color channel coupling, *Nucl. Phys. A* 352 (1981) 326–342. doi:doi:10.1016/0375-9474(81)90413-9.
- [40] S. J. Brodsky, C.-R. Ji, G. P. Lepage, Quantum Chromodynamic Predictions for the Deuteron Form-Factor, *Phys. Rev. Lett.* 51 (1983) 83. doi:doi:10.1103/PhysRevLett.51.83.
- [41] M. Harvey, On the Fractional Parentage Expansions of Color Singlet Six Quark States in a Cluster Model, *Nucl. Phys. A* 352 (1981) 301. doi:doi:10.1016/0375-9474(81)90412-7, [Erratum: *Nucl. Phys. A* 481, 834 (1988)].
- [42] M. Bashkanov, S. J. Brodsky, H. Clement, Novel Six-Quark Hidden-Color Dibaryon States in QCD, *Phys. Lett. B* 727 (2013) 438–442. doi:doi:10.1016/j.physletb.2013.10.059. [arXiv:1308.6404](#).
- [43] G. A. Miller, Pionic and Hidden-Color, Six-Quark Contributions to the Deuteron b1 Structure Function, *Phys. Rev. C* 89 (2014) 045203. doi:doi:10.1103/PhysRevC.89.045203. [arXiv:1311.4561](#).
- [44] J. Rittenhouse West, S. J. Brodsky, G. F. de Téramond, I. Schmidt, Diffractive dissociation of alpha particles as a test of isophobic short-range correlations inside nuclei, *Phys. Lett. B* 805 (2020) 135423. doi:doi:10.1016/j.physletb.2020.135423. [arXiv:1912.11288](#).
- [45] N. Jarmie, R. C. Allen, T (p, p) T Scattering near the T (p, n) He-3 Threshold, *Phys. Rev.* 114 (1959) 176–178. doi:doi:10.1103/PhysRev.114.176.
- [46] N. Jarmie, M. G. Silbert, D. B. Smith, J. S. Loos, Proton-Triton Elastic Scattering below 1 MeV, *Phys. Rev.* 130 (1963) 1987–1989. doi:doi:10.1103/PhysRev.130.1987.
- [47] T. Walcher, Excitation of 4 He by inelastic electron scattering at low momentum transfer, *Phys. Lett. B* 31 (1970) 442–444. doi:doi:10.1016/0370-2693(70)90148-6.
- [48] R. L. Jaffe, Exotica, *Phys. Rept.* 409 (2005) 1–45. doi:doi:10.1016/j.physrep.2004.11.005. [arXiv:hep-ph/0409065](#).
- [49] J. R. West, Diquark Induced Short-Range Nucleon-Nucleon Correlations & the EMC Effect, *Nuc. Phys. A* (2023). [arXiv:2009.06968](#).
- [50] E.-M. Ilgenfritz, J. Kripfganz, A. Schiller, Fragmentation of Quark and Diquark Jets, *Acta Phys. Polon. B* 9 (1978) 881.
- [51] J. R. Batley, et al. (NA48/2), Search for the dark photon in π^0 decays, *Phys. Lett. B* 746 (2015) 178–185. doi:doi:10.1016/j.physletb.2015.04.068. [arXiv:1504.00607](#).
- [52] D. Banerjee, et al. (NA64), Improved limits on a hypothetical X(16.7) boson and a dark photon decaying into e^+e^- pairs, *Phys. Rev. D* 101 (2020) 071101. doi:doi:10.1103/PhysRevD.101.071101. [arXiv:1912.11389](#).
- [53] F. Bossi, et al. (PADME), Cross-section measurement of two-photon in-flight annihilation of positrons at $s=20$ MeV with the PADME detector, *Phys. Rev. D* 107 (2023) 012008. doi:doi:10.1103/PhysRevD.107.012008. [arXiv:2210.14603](#).
- [54] L. Darmé, M. Mancini, E. Nardi, M. Raggi, Resonant search for the X17 boson at PADME, *Phys. Rev. D* 106 (2022) 115036. doi:doi:10.1103/PhysRevD.106.115036. [arXiv:2209.09261](#).