NMSSM with correct relic density and an additional 95 GeV Higgs boson

Ulrich Ellwanger^{*a*} and Cyril Hugonie^{*b*}

^a IJCLab, CNRS/IN2P3, University Paris-Saclay, 91405 Orsay, France ulrich.ellwanger@ijclab.in2p3.fr

^b LUPM, UMR 5299, CNRS/IN2P3, Université de Montpellier, 34095 Montpellier, France cyril.huqonie@umontpellier.fr

Abstract

We investigate whether it is possible within the NMSSM to describe simultaneously a correct dark matter relic density complying with the latest null results from the LZ experiment, an extra Higgs boson with a mass of ~ 95 GeV visible in the bb channel at LEP and the diphoton channel at the LHC, and the deviation of the anomalous magnetic moment of the muon from its value within the Standard Model. We find that this is still possible for a variety of dark matter annihilation mechanisms, for singlino-like but also bino-like lightest supersymmetric particles. We show the signal rates of the extra Higgs boson and the dark matter detection rates as function of the dark matter mass and annihilation mechanism. The dark matter direct detection cross section may well fall below the neutrino floor. The masses of the lightest electroweakly interacting supersymmetric particles are typically not far above 100 GeV, but not excluded due to unconventional decays.

1 Introduction

Supersymmetric extensions of the Standard Model may explain the dark matter of the universe. The lightest supersymmetric partner particle is stable, provided R-parity is conserved. It can well be neutral and freeze out into the desired relic density $\Omega_{DM}h^2 = 0.1187$ (we add $\pm 10\%$ theoretical uncertainty) in agreement with WMAP/Planck results [1,2]. However, it also has to comply with the absence of signals in direct detection experiments in spin-independent channels by the CRESST [3], PandaX-II [4], LUX [5], XENON1T [6,8], DarkSide [7] and LZ [9] collaborations, and in spin-dependent channels by the PICO-2L [10], LUX [11] and PandaX-II [12], Xenon1T [6,8] and LZ [9] collaborations.

Within the Minimal Supersymmetric extension of the Standard Model (MSSM) it is no longer easy to comply with these constraints [13]; in [14] it has been argued that additional contributions to the dark matter relic density would be required, but co-annihilation of higgsinos with charginos, with masses beyond 1 TeV, may still be a possibility [15]. In [16] it has been observed that co-annihilation with charginos or sleptons can allow for dark matter made out of a Lightest Supersymmetric Particle (LSP) consisting of a bino/wino-mixture, however at the price of some fine tuning [17–19] (see also [20–32]). Staus degenerate with the LSP seem also possible [33]. Given the lower bounds on chargino and slepton masses already from LEP, the mass of the LSP has then to be above 100 GeV. In [34] it is shown that, considering the constraints from LZ [9] and the LHC, even if additional contributions to the dark matter relic density are allowed the MSSM requires heavy higgsinos, or narrow regions in parameter space with light higgsinos or light staus.

Such constraints are alleviated within the Next-to-Minimal extension of the Standard Model (NMSSM) [35] where the LSP can have at least a strong singlino-component and correspond to the fermionic superpartner of the gauge singlet superfield. A pure singlino has no couplings to quarks and leptons, hence such an LSP has very small direct detection cross sections which can fall below the projections for the exclusion reach of the XENON-nT [36] and LZ [37], or even below the neutrino floor [38]. Hence the singlino is a quite attractive dark matter candidate [39–97] which can remain consistent with the absence of signals in direct detection experiments.

Actually hints exist for additional Higgs bosons as present in the NMSSM. The combination of searches for the SM Higgs boson by the ALEPH, DELPHI, L3 and OPAL experiments at LEP [98] showed some mild excess of events in the $Z^* \rightarrow Z + b\bar{b}$ channel in the mass region 95 – 100 GeV. Searches for Beyond-the-Standard Model (BSM) Higgs bosons at the LHC in the diphoton channel were performed by CMS and ATLAS. A search at run 1 by CMS showed a ~ 2 σ excess at 97 GeV [99], which was confirmed by CMS later in [100] and in [101] for a mass hypothesis of 95.4 GeV. A somewhat less sensitive search in the diphoton channel by ATLAS in [102] lead to an upper limit on the fiducial cross section which did not contradict the possible excess observed by CMS, a recent analysis by ATLAS in the diphoton channel in [103] showed a mild excess of 1.7σ at 95 GeV. A search for BSM Higgs bosons in the di-tau channel by CMS in [104] showed an excess of 2.6σ (local) for a mass of 95 – 100 GeV.

The ~ 2σ excess at LEP was quantified in [105]. Let us denote the extra (lighter) Higgs boson by H_{95} , with a reduced coupling to vector bosons W^{\pm}, Z (relative to the coupling of a SM-like Higgs boson of corresponding mass) given by $C_V(H_{95})$. Then the authors in [105] define

$$\mu_{bb}^{LEP} \equiv C_V(H_{95})^2 \times BR(H_{95} \to b\bar{b}) / BR(H_{SM}^{95} \to b\bar{b}) = 0.117 \pm 0.057$$
(1.1)

where H_{SM}^{95} denotes a fictitious SM-like Higgs boson with a mass of 95 GeV.

The best fits for a diphoton signal of H_{95} in CMS and ATLAS were combined in [106]. The authors in [106] obtain

$$\mu_{\gamma\gamma}^{LHC} = \frac{\sigma(gg \to H_{95} \to \gamma\gamma)}{\sigma(gg \to H_{SM}^{95} \to \gamma\gamma)} = 0.24^{+0.09}_{-0.08} . \tag{1.2}$$

Again, H_{SM}^{95} denotes a SM-like Higgs boson with a mass of ~ 95 GeV.

The best fit for the excess in the di-tau channel at 95 GeV observed by CMS in [104] corresponds to a cross section times branching fraction

$$\sigma(gg \to H_{95} \to \tau\tau) = 7.8^{+3.9}_{-3.1} \text{pb} .$$
 (1.3)

Relative to H_{SM}^{95} we obtain

$$\mu_{\tau\tau}^{LHC} = \frac{\sigma(gg \to H_{95} \to \tau\tau)}{\sigma(gg \to H_{SM}^{95} \to \tau\tau)} = 1.38^{+0.69}_{-0.55} \,. \tag{1.4}$$

Such an additional Higgs boson with a mass of ~ 95 GeV has been discussed within the NMSSM in [105–121]; mostly, however, allowing for a too small dark matter relic density (assuming additional contributions), and/or before the consideration of constraints from LZ [9]. It is the aim of the present paper to verify whether attractive scenarios within the NMSSM exist with a correct relic density, satisfying the LZ-constraints, and an additional Higgs boson with a mass of ~ 95 GeV. To this end we scan the parameter space over the NMSSM-parameters λ , κ , A_{λ} , A_{κ} , μ_{eff} , tan β in the Higgs sector and non universal soft supersymmetry breaking terms in the gaugino/sfermion sector; for their definitions we refer to review in [35]. We employ the codes NMSSMTools-6.0.2 [122–124] and micrOMEGAs_3 [125].

We impose constraints from b-physics and constraints from searches for BSM Higgs bosons by ATLAS and CMS as implemented in NMSSMTools-6.0.2. The references to constraints from BSM Higgs-boson searches and b-physics (of little relevance here) are listed on the web page https://www.lupm.in2p3.fr/users/nmssm/history.html. Constraints from the absence of a Landau singularity for the Yukawa couplings below the GUT scale confine values of the NMSSM-specific coupling λ to $\lambda \leq 0.7$ (see [35] for the the meaning of NMSSM specific parameters).

Constraints from the anomalous magnetic moment of the muon a_{μ} [126, 127] as in [128] can always be satisfied by choosing the soft supersymmetry breaking smuon, bino and/or wino masses small enough, and the trilinear coupling A_{μ} large enough. These parameters have little impact on the quantities of relevance here which are shown in the Figures in the next Section, hence constraints from a_{μ} are not imposed on most of the points in parameter space. However, for the benchmark points in the next Section, constraints from the a_{μ} are satisfied. The constraint on M_W as applied in [128] is not used since it relies on a single experimental result which differs significantly from many others. All soft supersymmetry breaking terms are taken below 5 TeV.

Constraints on the sparticle spectrum from the absence of signals at the LHC are taken into account using the code SModels-2.2.0 [129–132]. Constraints on light electroweakly

interacting sparticles (in the $\sim 100 - 200$ GeV range) from the LHC rely on production cross sections for higgsinos or winos, and their decays into W- or Z-bosons, see e.g. the recent ATLAS combination in [133]. In the NMSSM, both production cross sections of light electroweakly interacting sparticles as well as corresponding decay rates can be suppressed and the constraints from the LHC become weak. Moreover, in the presence of several light electroweakly interacting sparticles, heavy sparticles (squarks, gluino) can undergo several distinct cascade decays such that the probability for each of them is reduced. As a consequence such scenarios are difficult to rule out.

We require that the singlet-like scalar has a mass in the range 95.4 ± 3 GeV (allowing for a theoretical uncertainty of 3 GeV), μ_{bb}^{LEP} in the 2σ range of (1.1), and $\mu_{\gamma\gamma}^{LHC}$ in the 2σ range of (1.2). For the SM-like Higgs boson we require a mass within 125.2 ± 3 GeV (allowing for theoretical uncertainties), and that the couplings in the κ -framework satisfy combined limits of CMS [134] and ATLAS [135]. These limits require that the reduced couplings $C_V(H_{SM})$ of the SM-like Higgs boson to gauge fields satisfy $C_V(H_{SM}) \gtrsim 0.96$, hence the mixing angle between H_{95} and H_{SM} is limited such that the reduced coupling of H_{95} to $\tau\tau$ is bounded to below 25%. Then the (reduced) cross section × BR($H_{95} \rightarrow \tau\tau$) can hardly exceed 0.1 in contrast to what is desired within the 2σ range for the di-tau excess $\mu_{\tau\tau}^{LHC}$ in (1.4); its description within the 2σ range is therefore left aside since impossible for the type II Yukawa couplings present in the (N)MSSM.

In the next Section we present our results in the form of Figures and benchmark points, Section 3 is devoted to conclusions.

2 Results

Various mechanisms exist within the NMSSM for a sufficiently fast annihilation of dark matter in the form of a singlino-like LSP:

- (a) Annihilation via a pseudoscalar in the s-channel. Then M_{A_1} should be about $2 \times M_{\chi_1^0}$ such that the annihilation cross section is enhanced by the s-channel pole. We find that this annihilation mechanism is possible in the NMSSM for $M_{\chi_1^0} \leq 24$ GeV, i.e. $M_{A_1} \leq 50$ GeV. A_1 is always mostly singlet-like, and decays into $b\bar{b}$ for $M_{A_1} \gtrsim 9$ GeV, into $\tau\tau$ otherwise.
- (b) Annihilation via the Standard Model-like Higgs boson H_{SM} or the mostly singlet-like Higgs boson H_{95} in the s-channel if the singlino mass is about half the corresponding mass. (With $M_{H_{95}} \sim 95$ GeV, the Z-boson plays little role in the s-channel since the couplings of an LSP with $M_{LSP} \sim M_Z/2$ to H_{95} are typically much stronger.)
- (c) Co-annihilation with charginos, sleptons, or the next-to-lightest neutralinos. (Subsequently we include staus and their sneutrinos in the notion sleptons.) Often several of these sparticles intervene simultaneously with different strengths; then these sparticles must be close in mass (within ~ 10 GeV) to the singlino. In most cases, the lightest charged or electroweakly interacting sparticle must be heavier than ~ 100 GeV from constraints from LEP. Then a lower bound of ~ 90 GeV holds for the singlino, and the properties of the valid points shown in the figures below depend little on the nature of the

sparticle(s) which intervene in co-annihilation. An exception are sneutrinos which can be lighter, as light as 55 GeV if they annihilate through the H_{SM} funnel.

(d) A bino-like LSP as in the MSSM under the latest constraints [34] (or in the NMSSM e.g. in [136, 137]). For a practically pure bino, LEP and LHC constraints allow for masses up to 330 GeV down to ~ 20 GeV. Compared to singlino-like LSPs, pair-annihilation of binos via a (pseudo-)scalar in the s-channel is relatively less important. The reason is that singlino-singlino-Higgs Yukawa couplings ~ λ exist in the NMSSM for the singlet-like (pseudo-)scalar at tree level, and other scalars through mixing. However, bino-bino-Higgs Yukawa couplings do not exist for pure binos since the bino is a gaugino of an abelian gauge symmetry. The dominant annihilation channelis therefore co-annihilation with sleptons.

All mechanisms allow for a dark matter relic density in agreement with the WMAP/Planck result $\Omega_{DM}h^2 = 0.1187 \pm 10\%$.

It is interesting to compare the possible values of the signal strengths μ_{bb}^{LEP} , $\mu_{\gamma\gamma}^{LHC}$ and $\mu_{\tau\tau}^{LHC}$ as well as the dark matter cross sections (spin independent on protons σ_p^{SI} , and spin dependent on neutrons σ_n^{SD}) for the various possible dark matter masses associated to various annihilation processes, as function of the mass of the LSP.

In Figs. 1 we show in how far points in the NMSSM can accomodate an additional Higgs boson near 95 GeV with signal rates μ_{bb}^{LEP} at LEP as in eq. (1.1). (Only points within the 2σ range of (1.1) are shown.) Here and in the subsequent figures, points on the left hand side refer to a singlino-like LSP. Points corresponding to annihilation via a pseudoscalar A_1 (A1 funnel) are shown in violet, annihilation via H_{95} or H_{SM} in orange, co-annihilation with charginos, sleptons or next-to-lightest neutralinos in blue. Note that, for a LSP mass near $H_{SM}/2$ or $H_{95}/2$, the dominant dark matter annihilation mechanism can be either via the corresponding Higgs funnel or via co-annihilation. Points with a bino-like LSP are shown on the right hand side in black. The red dots denote benchmark points whose details are given in the Table 2-4 below. Comparing the right hand side and the left hand side of Figs. 1 one observes that the possible values of μ_{bb}^{LEP} are generally somewhat larger for bino LSPs compared to singlino LSPs.

The contributing range of parameters depends somewhat on the annihilation mechanism. In Table 1 we give these ranges for the NMSSM specific parameters and the soft supersymmetry breaking masses M_1 , M_2 and M_3 where the latter refer to the bino, wino and gluino masses, respectively. The soft supersymmetry breaking mass terms for scalars vary from 83 GeV (for sleptons) and 320 GeV (for squarks) up to 5 TeV, the soft supersymmetry breaking trilinear couplings from -5 TeV to 5 TeV. (The upper limits of 5 TeV are imposed by hand.) Given the possibly complicated squark decay cascades, squarks can be relatively light without contradicting constraints from searches at the LHC. In any case, due to radiative corrections the pole squark masses can differ considerably from the soft supersymmetry breaking mass terms.

In Fig. 2 we show $\mu_{\gamma\gamma}^{LHC}$ for points within the 2σ range of the signal rate (1.2) for an additional Higgs boson near 95 GeV in the $\gamma\gamma$ channel at the LHC. The colors denote the dark matter annihilation mechanisms as in Fig. 1.

Fig. 3 shows the signal rate $\mu_{\tau\tau}^{LHC}$ for an additional Higgs boson near 95 GeV for points in the NMSSM within the 2σ ranges of the signal rates (1.1) and (1.2), and all other constraints. Actually Fig. 3 looks similar to Fig. 1, the reason being that the couplings of H_{95} to b-quarks



Figure 1: μ_{bb}^{LEP} for points in the NMSSM satisfying all constraints. Left hand side: Points with a singlino-like LSP. Points corresponding to annihilation via a pseudoscalar A_1 (A1 funnel) are shown in violet, annihilation via H_{95} or H_{SM} in orange, co-annihilation with charginos, sleptons, next-to-lightest neutralinos in blue. Right hand side: Points with a bino-like LSP. The red dots denote benchmark points whose details are given in the Tables 2-4 below.

	A_1 -funnel	H_{95} - or H_{SM} -funnels	co-annihilation	bino LSP
λ	0.17 - 0.52	0.18 - 0.32	0.019 - 0.34	0.019 - 0.22
κ	$-(6.3-0.36)\times10^{-3}$	$-(1.48-0.59)\times10^{-2}$	-0.011 - 0.024	0.0019 - 0.020
A_{λ}	2862 - 5000	3486 - 5000	-5000 - 4320	-(5000-1503)
A_{κ}	12 - 65	-122 - 175	-3.4-551	-1.0 - 585
μ_{eff}	350 - 1388	575 - 946	-1326 - 136	-(1226-361)
$\tan\beta$	3.3-11	4.0 - 7.7	3.2-14	3.8 - 19
M_1	15 - 5000	-227 - 5000	56 - 5000	25 - 329
M_2	89-5000	91-5000	-5000 - 680	-(5000-89)
M_3	746 - 5000	729 - 5000	722-5000	721 - 5000

Table 1: Range of NMSSM specific input parameters and bino (M_1) , wino (M_2) and gluino (M_3) masses for the various dark matter annihilation mechanisms. All dimensionful parameters are given in GeV.

and τ -leptons are related in the NMSSM with its type 2 Yukawa couplings of H_{SM} ; the Yukawa couplings of H_{95} are inherited from those of H_{SM} through mixing. A larger mixing angle would enhance the couplings of H_{95} , but reduce simultaneously those of H_{SM} . The latter are limited from below by the latest measurements of ATLAS [135] and CMS [134]. Hence it is difficult in the NMSSM to obtain large couplings of H_{95} , i.e. larger signal rates than those shown in Fig. 3 which are below the 2 σ range of 0.28 for $\mu_{\tau\tau}^{LHC}$ in (1.4).

In Fig. 4 we show the spin-independent (SI) and spin-dependent (SD) direct detection cross section. We also show the present exclusion of the LZ experiment [9], projections for the LZ



Figure 2: $\mu_{\gamma\gamma}^{LHC}$ for an additional Higgs boson near 95 GeV for points in the NMSSM satisfying all constraints. The colors denote the dark matter annihilation mechanisms as in Fig. 1.



Figure 3: $\mu_{\tau\tau}^{LHC}$ for an additional Higgs boson near 95 GeV for points in the NMSSM satisfying all constraints. The colors denote the dark matter annihilation mechanisms as in Fig. 1.

exclusion reach from [37], and the neutrino floor from [38]. We see that the SI direct detection cross section can well fall below the neutrino floor, in which case direct detection experiments are not able to verify this form of dark matter.

In Fig. 5 we show the mass M_{A1} of the mostly singlet-like pseudoscalar, which is $\sim 2 \times M_{LSP}$ for points in the A_1 -funnel. For $M_{A1} \leq 60$ GeV, A_1 may be visible in decays of H_{SM} , see the latest study by CMS in [138]. Accordingly A_1 may be visible (but not necessarily) for all dark matter annihilation mechanisms.

In Tables 2-4 we show the properties of some benchmark points (BPs) with distinct dark matter annihilation processes: Input parameters in Table 2, some Higgs masses and the nature



Figure 4: σ_p^{SI} and σ_n^{SD} for points in the NMSSM with the correct relic density and satisfying all constraints. The colors denote the dark matter annihilation mechanisms as in Fig. 1. The full green line indicates the present upper limits from LZ [9], the dashed green line the future prospects on upper limits from LZ from [37], the dotted green line the neutrino floor from [38].

of light electroweakinos in Table 3, and the properties shown in Figs. 1-5 in Table 4.

BP1 corresponds to dark matter annihilation via the A_1 funnel, accordingly the mass of the singlino-like LSP is close to $M_{A1}/2$. BP2 and BP3 correspond to dark matter annihilation via the H_{95} and H_{SM} funnels, respectively. Accordingly the masses of the singlino-like LSP are close to $M_{H_{95}}/2$ and $M_{H_{SM}}/2$, respectively. BP4 corresponds to co-annihilation with a slepton and/or the second-lightest bino-like neutralino; in fact several possibilities are included here. σ_p^{SI} for BP4 is below the neutrino floor. Finally BP5 corresponds to a bino-like LSP which is possible in the NMSSM as well as in the MSSM.



Figure 5: M_{A1} for points in the NMSSM satisfying all constraints. The colors denote the dark matter annihilation mechanisms as in Fig. 1.

	BP1	BP2	BP3	BP4	BP5
λ	0.458	0.242	0.205	0.0223	0.0590
κ	-5.74×10^{-3}	-7.26×10^{-3}	-8.06×10^{-3}	9.63×10^{-3}	7.75×10^{-3}
A_{λ}	3970	4999	4544	-3390	-4995
A_{κ}	34.9	11.0	11.0	183	252
$\mu_{ ext{eff}}$	404	736	794	-179	-681
an eta	9.76	6.25	5.13	7.01	8.94
M_1	163	2065	76.6	196	45.2
M_2	111	100	96.7	-502	-106
M_3	2735	4429	1201	3505	825

Table 2: NMSSM specific input parameters and bino (M_1) , wino (M_2) and gluino (M_3) masses for the five benchmark points. All dimensionful parameters are given in GeV.

3 Conclusions

The NMSSM allows to describe simultaneously three phenomena related to physics beyond the Standard Model: A dark matter relic density of the amount observed in cosmology, but compatible with the absence of direct detection signals; a deviation of the muon anomalous magnetic moment by 4.2σ from its value in the Standard Model; anomalies which indicate the presence of an extra scalar of a mass near 95 GeV in the *bb* channel at LEP, and in $\gamma\gamma$ and $\tau\tau$ channels at the LHC. The latter would be impossible to describe within the MSSM. The LSP may have a mass well below 100 GeV, and at least part of the electroweak sparticle sector (sleptons, winos, bino) is typically relatively light. Such sparticles are difficult to rule out at the LHC, and generate complicated decay cascades also for the heavier strongly interacting sparticles. Their verification may then be a demanding task for the future.

	BP1	BP2	BP3	BP4	BP5
M_{H3}	3966	4852	4367	2069	5592
M_{A1}	21	18	31	206	260
LSP	singl.	singl.	singl.	singl.	bino
$M_{\rm LSP}$	9.0	43.9	62.9	157	44.1
NLSP	$wino^{\pm}$	wino ^{\pm}	bino	bino	wino ⁰
$M_{\rm NLSP}$	115	107	74.8	164	113
Slepton	$\tilde{\nu}_{ au}$	$\tilde{\nu}_e$	$\tilde{e}_{R.h.}$	$\tilde{\nu}_e$	$\tilde{\nu}_{ au}$
$M_{\rm Slepton}$	140	171	255	165	93.4

Table 3: Some BSM Higgs masses, the nature of the LSP, NLSP, the lightest slepton, and the corresponding masses for five BMpoints. All dimensionful parameters are given in GeV.

	BP1	BP2	BP3	BP4	BP5
μ_{bb}^{LEP}	1.18×10^{-2}	0.115	0.106	0.105	0.116
$\mu_{\gamma\gamma}^{LHC}$	0.331	8.01×10^{-2}	8.01×10^{-2}	8.75×10^{-2}	0.130
$\mu_{ au au}^{LHC}$	1.28×10^{-2}	0.112	0.102	9.34×10^{-2}	0.117
σ_n^{SI}	3.43×10^{-11}	4.98×10^{-12}	1.59×10^{-12}	1.49×10^{-13}	8.90×10^{-12}
σ_n^{SD}	3.66×10^{-5}	3.12×10^{-7}	1.15×10^{-7}	1.06×10^{-7}	4.40×10^{-7}

Table 4: μ_{bb}^{LEP} , $\mu_{\gamma\gamma}^{LHC}$, $\mu_{\tau\tau}^{SI}$, σ_p^{SI} and σ_n^{SD} (the latter in pb) for the five benchmark points. σ_p^{SI} for BP4 is below the neutrino floor.

References

- G. Hinshaw *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **208** (2013) 19 [arXiv:1212.5226 [astro-ph.CO]].
- [2] P. A. R. Ade *et al.* [Planck Collaboration], Astron. Astrophys. 571 (2014) A16 [arXiv:1303.5076 [astro-ph.CO]].
- [3] G. Angloher *et al.* [CRESST], Eur. Phys. J. C 76 (2016) no.1, 25 [arXiv:1509.01515 [astroph.CO]].
- [4] A. Tan *et al.* [PandaX-II Collaboration], Phys. Rev. Lett. **117** (2016) no.12, 121303 [arXiv:1607.07400 [hep-ex]].
- [5] D. S. Akerib *et al.* [LUX Collaboration], Phys. Rev. Lett. **118** (2017) no.2, 021303 [arXiv:1608.07648 [astro-ph.CO]].
- [6] E. Aprile *et al.* [XENON Collaboration], Phys. Rev. Lett. **119** (2017) no.18, 181301 [arXiv:1705.06655 [astro-ph.CO]].
- [7] P. Agnes *et al.* [DarkSide], Phys. Rev. Lett. **121** (2018) no.8, 081307 [arXiv:1802.06994 [astro-ph.HE]].
- [8] E. Aprile *et al.* [XENON Collaboration], arXiv:1805.12562 [astro-ph.CO].
- [9] J. Aalbers *et al.* [LZ], Phys. Rev. Lett. **131** (2023) no.4, 041002 [arXiv:2207.03764 [hep-ex]].
- [10] C. Amole *et al.* [PICO Collaboration], Phys. Rev. D **93** (2016) no.6, 061101
 [arXiv:1601.03729 [astro-ph.CO]].
- [11] D. S. Akerib *et al.* [LUX Collaboration], Phys. Rev. Lett. **116** (2016) no.16, 161302
 [arXiv:1602.03489 [hep-ex]].
- [12] C. Fu *et al.* [PandaX-II Collaboration], Phys. Rev. Lett. **118** (2017) no.7, 071301 Erratum:
 [Phys. Rev. Lett. **120** (2018) no.4, 049902] [arXiv:1611.06553 [hep-ex]].
- [13] H. Baer, V. Barger and H. Serce, Phys. Rev. D 94 (2016) no.11, 115019 [arXiv:1609.06735 [hep-ph]].
- [14] H. Baer, V. Barger, D. Sengupta and X. Tata, Eur. Phys. J. C 78 (2018) no.10, 838 doi:10.1140/epjc/s10052-018-6306-y [arXiv:1803.11210 [hep-ph]].
- [15] A. Delgado and M. Quirós, Phys. Rev. D 103 (2021) no.1, 015024 [arXiv:2008.00954 [hepph]].
- [16] M. Chakraborti, S. Heinemeyer, I. Saha and C. Schappacher, Eur. Phys. J. C 82 (2022) no.5, 483 [arXiv:2112.01389 [hep-ph]].
- [17] F. Wang, W. Wang, J. Yang, Y. Zhang and B. Zhu, Universe 8 (2022) no.3, 178 [arXiv:2201.00156 [hep-ph]].

- [18] M. E. Gomez, Q. Shafi, A. Tiwari and C. S. Un, Eur. Phys. J. C 82 (2022) no.6, 561 [arXiv:2202.06419 [hep-ph]].
- [19] J. M. Yang and Y. Zhang, Sci. Bull. 67 (2022) no.14, 1430-1436 [arXiv:2204.04202 [hep-ph]].
- [20] M. Chakraborti, S. Heinemeyer and I. Saha, Eur. Phys. J. C 81 (2021) no.12, 1069 [arXiv:2103.13403 [hep-ph]].
- [21] S. Iwamoto, T. T. Yanagida and N. Yokozaki, Phys. Lett. B 823 (2021), 136768 doi:10.1016/j.physletb.2021.136768 [arXiv:2104.03223 [hep-ph]].
- [22] M. Van Beekveld, W. Beenakker, M. Schutten and J. De Wit, SciPost Phys. 11 (2021) no.3, 049 [arXiv:2104.03245 [hep-ph]].
- [23] P. Cox, C. Han and T. T. Yanagida, Phys. Rev. D 104 (2021) no.7, 075035 [arXiv:2104.03290 [hep-ph]].
- [24] M. Chakraborti, L. Roszkowski and S. Trojanowski, JHEP 05 (2021), 252 [arXiv:2104.04458 [hep-ph]].
- [25] H. Baer, V. Barger and H. Serce, Phys. Lett. B 820 (2021), 136480 [arXiv:2104.07597 [hep-ph]].
- [26] Q. Shafi and C. S. Un, [arXiv:2107.04563 [hep-ph]].
- [27] A. K. Forster and S. F. King, Nucl. Phys. B 976 (2022), 115700 [arXiv:2109.10802 [hepph]].
- [28] K. Agashe, M. Ekhterachian, Z. Liu and R. Sundrum, JHEP 09 (2022), 142 [arXiv:2203.01796 [hep-ph]].
- [29] J. Zhao, J. Zhu, P. Zhu and R. Zhu, Phys. Rev. D 107 (2023) no.5, 055030 [arXiv:2211.14587 [hep-ph]].
- [30] S. Baum, M. Carena, T. Ou, D. Rocha, N. R. Shah and C. E. M. Wagner, JHEP 11 (2023), 037 doi:10.1007/JHEP11(2023)037 [arXiv:2303.01523 [hep-ph]].
- [31] Y. He, L. Meng, Y. Yue and D. Zhang, Phys. Rev. D 108 (2023) no.11, 115010 [arXiv:2303.02360 [hep-ph]].
- [32] S. Bisal, A. Chatterjee, D. Das and S. A. Pasha, [arXiv:2311.09938 [hep-ph]].
- [33] M. Chakraborti, S. Heinemeyer and I. Saha, Eur. Phys. J. C 84 (2024) no.2, 165 [arXiv:2308.05723 [hep-ph]].
- [34] R. K. Barman, G. Bélanger, B. Bhattacherjee, R. Godbole and R. Sengupta, "Current status of the light neutralino thermal dark matter in the phenomenological MSSM," [arXiv:2402.07991 [hep-ph]].

- [35] U. Ellwanger, C. Hugonie and A. M. Teixeira, Phys. Rept. 496 (2010) 1 [arXiv:0910.1785 [hep-ph]].
- [36] E. Aprile et al. [XENON], JCAP 11 (2020), 031 [arXiv:2007.08796 [physics.ins-det]].
- [37] D. S. Akerib *et al.* [LZ], Phys. Rev. D **101** (2020) no.5, 052002 [arXiv:1802.06039 [astroph.IM]].
- [38] J. Billard, L. Strigari and E. Figueroa-Feliciano, Phys. Rev. D 89 (2014) no.2, 023524 [arXiv:1307.5458 [hep-ph]].
- [39] D. G. Cerdeno, C. Hugonie, D. E. Lopez-Fogliani, C. Munoz and A. M. Teixeira, JHEP 0412 (2004) 048 [hep-ph/0408102].
- [40] G. Belanger, F. Boudjema, C. Hugonie, A. Pukhov and A. Semenov, JCAP 0509 (2005) 001 [hep-ph/0505142].
- [41] D. G. Cerdeno, E. Gabrielli, D. E. Lopez-Fogliani, C. Munoz and A. M. Teixeira, JCAP 0706 (2007) 008 [hep-ph/0701271 [HEP-PH]].
- [42] V. Barger, P. Langacker, I. Lewis, M. McCaskey, G. Shaughnessy and B. Yencho, Phys. Rev. D 75 (2007) 115002 [hep-ph/0702036 [HEP-PH]].
- [43] G. Belanger, C. Hugonie and A. Pukhov, JCAP 0901 (2009) 023 [arXiv:0811.3224 [hepph]].
- [44] D. Albornoz Vasquez, G. Belanger, C. Boehm, A. Pukhov and J. Silk, Phys. Rev. D 82, 115027 (2010) [arXiv:1009.4380 [hep-ph]].
- [45] M. Perelstein and B. Shakya, Phys. Rev. D 88 (2013) no.7, 075003 [arXiv:1208.0833 [hepph]].
- [46] J. Kozaczuk and S. Profumo, Phys. Rev. D 89 (2014) no.9, 095012 [arXiv:1308.5705 [hepph]].
- [47] J. Cao, C. Han, L. Wu, P. Wu and J. M. Yang, JHEP 1405 (2014) 056 [arXiv:1311.0678 [hep-ph]].
- [48] J. S. Kim and T. S. Ray, Eur. Phys. J. C **75** (2015) 40 [arXiv:1405.3700 [hep-ph]].
- [49] U. Ellwanger and C. Hugonie, JHEP **1408** (2014) 046 [arXiv:1405.6647 [hep-ph]].
- [50] K. Ishikawa, T. Kitahara and M. Takimoto, Phys. Rev. Lett. **113** (2014) no.13, 131801 [arXiv:1405.7371 [hep-ph]].
- [51] T. Han, Z. Liu and S. Su, JHEP **1408** (2014) 093 [arXiv:1406.1181 [hep-ph]].
- [52] C. Cheung, M. Papucci, D. Sanford, N. R. Shah and K. M. Zurek, Phys. Rev. D 90 (2014) no.7, 075011 [arXiv:1406.6372 [hep-ph]].

- [53] J. Huang, T. Liu, L. T. Wang and F. Yu, Phys. Rev. D 90 (2014) no.11, 115006 [arXiv:1407.0038 [hep-ph]].
- [54] M. Cahill-Rowley, J. Gainer, J. Hewett and T. Rizzo, JHEP 1502 (2015) 057 [arXiv:1409.1573 [hep-ph]].
- [55] J. Guo, J. Li, T. Li and A. G. Williams, Phys. Rev. D 91 (2015) no.9, 095003 [arXiv:1409.7864 [hep-ph]].
- [56] J. Cao, L. Shang, P. Wu, J. M. Yang and Y. Zhang, Phys. Rev. D 91 (2015) no.5, 055005 [arXiv:1410.3239 [hep-ph]].
- [57] T. Gherghetta, B. von Harling, A. D. Medina, M. A. Schmidt and T. Trott, Phys. Rev. D 91 (2015) 105004 [arXiv:1502.07173 [hep-ph]].
- [58] X. J. Bi, L. Bian, W. Huang, J. Shu and P. F. Yin, Phys. Rev. D 92 (2015) 023507 [arXiv:1503.03749 [hep-ph]].
- [59] C. Han, D. Kim, S. Munir and M. Park, JHEP **1507** (2015) 002 [arXiv:1504.05085 [hepph]].
- [60] C. T. Potter, Eur. Phys. J. C **76** (2016) no.1, 44 [arXiv:1505.05554 [hep-ph]].
- [61] R. Enberg, S. Munir, C. Pérez de los Heros and D. Werder, arXiv:1506.05714 [hep-ph].
- [62] J. Cao, L. Shang, P. Wu, J. M. Yang and Y. Zhang, JHEP **1510** (2015) 030 [arXiv:1506.06471 [hep-ph]].
- [63] A. Butter, T. Plehn, M. Rauch, D. Zerwas, S. Henrot-Versillé and R. Lafaye, Phys. Rev. D 93 (2016) 015011 [arXiv:1507.02288 [hep-ph]].
- [64] D. Barducci, G. Bélanger, C. Hugonie and A. Pukhov, JHEP 1601 (2016) 050 [arXiv:1510.00246 [hep-ph]].
- [65] M. Badziak, M. Olechowski and P. Szczerbiak, JHEP 1603 (2016) 179 [arXiv:1512.02472 [hep-ph]].
- [66] Q. F. Xiang, X. J. Bi, P. F. Yin and Z. H. Yu, Phys. Rev. D 94 (2016) no.5, 055031 [arXiv:1606.02149 [hep-ph]].
- [67] J. Cao, Y. He, L. Shang, W. Su and Y. Zhang, JHEP 1608 (2016) 037 [arXiv:1606.04416 [hep-ph]].
- [68] J. Cao, Y. He, L. Shang, W. Su, P. Wu and Y. Zhang, JHEP 1610 (2016) 136 [arXiv:1609.00204 [hep-ph]].
- [69] C. Beskidt, W. de Boer, D. I. Kazakov and S. Wayand, Phys. Lett. B 771 (2017) 611 [arXiv:1703.01255 [hep-ph]].
- [70] Q. Mou and S. Zheng, "Light Neutralino Dark Matter in the NMSSM," arXiv:1703.00343 [hep-ph].

- [71] M. Badziak, M. Olechowski and P. Szczerbiak, JHEP **1707** (2017) 050 [arXiv:1705.00227 [hep-ph]].
- [72] U. Ellwanger, JHEP **02** (2017), 051 [arXiv:1612.06574 [hep-ph]].
- [73] S. Baum, M. Carena, N. R. Shah and C. E. M. Wagner, JHEP 04 (2018), 069 [arXiv:1712.09873 [hep-ph]].
- [74] L. Shang, Y. He, J. Lian and Y. Pan, Eur. Phys. J. C 78 (2018) no.5, 390 [arXiv:1804.08797 [hep-ph]].
- [75] U. Ellwanger and C. Hugonie, Eur. Phys. J. C 78 (2018) no.9, 735 [arXiv:1806.09478 [hep-ph]].
- [76] J. Cao, Y. He, L. Shang, Y. Zhang and P. Zhu, Phys. Rev. D 99 (2019) no.7, 075020 [arXiv:1810.09143 [hep-ph]].
- [77] F. Domingo, J. S. Kim, V. M. Lozano, P. Martin-Ramiro and R. Ruiz de Austri, Phys. Rev. D 101 (2020) no.7, 075010 [arXiv:1812.05186 [hep-ph]].
- [78] W. Abdallah, A. Chatterjee and A. Datta, JHEP 09 (2019), 095 [arXiv:1907.06270 [hepph]].
- [79] J. Cao, L. Meng, Y. Yue, H. Zhou and P. Zhu, Phys. Rev. D 101 (2020) no.7, 075003 [arXiv:1910.14317 [hep-ph]].
- [80] K. Wang and J. Zhu, Chin. Phys. C 44 (2020) no.6, 061001 [arXiv:1911.08319 [hep-ph]].
- [81] K. Wang and J. Zhu, JHEP **06** (2020), 078 [arXiv:2002.05554 [hep-ph]].
- [82] K. Wang and J. Zhu, Phys. Rev. D **101** (2020) no.9, 095028 [arXiv:2003.01662 [hep-ph]].
- [83] M. Guchait and A. Roy, Phys. Rev. D **102** (2020) no.7, 075023 [arXiv:2005.05190 [hep-ph]].
- [84] R. K. Barman, G. Bélanger, B. Bhattacherjee, R. Godbole, D. Sengupta and X. Tata, Phys. Rev. D 103 (2021) no.1, 015029 [arXiv:2006.07854 [hep-ph]].
- [85] K. Wang, J. Zhu and Q. Jie, Chin. Phys. C 45 (2021) no.4, 041003 [arXiv:2011.12848 [hep-ph]].
- [86] H. Zhou, J. Cao, J. Lian and D. Zhang, Phys. Rev. D 104 (2021) no.1, 015017 [arXiv:2102.05309 [hep-ph]].
- [87] J. Cao, D. Li, J. Lian, Y. Yue and H. Zhou, JHEP 06 (2021), 176 [arXiv:2102.05317 [hep-ph]].
- [88] J. Cao, J. Lian, Y. Pan, D. Zhang and P. Zhu, JHEP 09 (2021), 175 [arXiv:2104.03284 [hep-ph]].
- [89] J. Cao, J. Lian, Y. Pan, Y. Yue and D. Zhang, JHEP 03 (2022), 203 [arXiv:2201.11490 [hep-ph]].

- [90] A. Chatterjee, A. Datta and S. Roy, JHEP 06 (2022), 108 [arXiv:2202.12476 [hep-ph]].
- [91] J. Cao, F. Li, J. Lian, Y. Pan and D. Zhang, Sci. China Phys. Mech. Astron. 65 (2022) no.9, 291012 [arXiv:2204.04710 [hep-ph]].
- [92] M. M. Almarashi, F. Alhazmi, R. Abdulhafidh and S. A. Basir, Results Phys. 49 (2023), 106531 [arXiv:2204.06082 [hep-ph]].
- [93] J. Cao, X. Jia, L. Meng, Y. Yue and D. Zhang, JHEP 03 (2023), 198 [arXiv:2210.08769 [hep-ph]].
- [94] L. Wang, J. M. Yang, Y. Zhang, P. Zhu and R. Zhu, Universe 9 (2023) no.4, 178 [arXiv:2302.05719 [hep-ph]].
- [95] Z. Heng, S. Yang, X. Li and L. Shang, Symmetry 15 (2023) no.2, 456 [arXiv:2302.07465 [hep-ph]].
- [96] J. Cao, L. Meng and Y. Yue, Phys. Rev. D 108 (2023) no.3, 035043 [arXiv:2306.06854 [hep-ph]].
- [97] J. Cao, X. Jia and J. Lian, "Unified Interpretation of Muon g-2 anomaly, 95 GeV Diphoton, and bb Excesses in the General Next-to-Minimal Supersymmetric Standard Model," [arXiv:2402.15847 [hep-ph]].
- [98] R. Barate et al. [LEP Working Group for Higgs boson searches, ALEPH, DELPHI, L3 and OPAL], Phys. Lett. B 565 (2003), 61-75 [arXiv:hep-ex/0306033 [hep-ex]].
- [99] CMS Collaboration, "Search for new resonances in the diphoton final state in the mass range between 80 and 115 GeV in pp collisions at $\sqrt{s}=8$ TeV", CMS-PAS-HIG-14-037
- [100] A. M. Sirunyan et al. [CMS], Phys. Lett. B 793 (2019), 320-347 [arXiv:1811.08459 [hepex]].
- [101] CMS Collaboration, "Search for a standard model-like Higgs boson in the mass range between 70 and 110 GeV in the diphoton final state in proton-proton collisions at $\sqrt{s}=13$ TeV", CMS-PAS-HIG-20-002
- [102] ATLAS Collaboration, "Search for resonances in the 65 to 110 GeV diphoton invariant mass range using 80 fb⁻¹ of pp collisions collected at \sqrt{s} =13 TeV with the ATLAS detector", ATLAS-CONF-2018-025
- [103] ATLAS Collaboration, "Search for diphoton resonances in the 66 to 110 GeV mass range using 140 fb⁻¹ of 13 TeV pp collisions collected with the ATLAS detector", ATLAS-CONF-2023-035
- [104] A. Tumasyan *et al.* [CMS], JHEP **07** (2023), 073 [arXiv:2208.02717 [hep-ex]].
- [105] J. Cao, X. Guo, Y. He, P. Wu and Y. Zhang, Phys. Rev. D 95 (2017) no.11, 116001 [arXiv:1612.08522 [hep-ph]].

- [106] T. Biekötter, S. Heinemeyer and G. Weiglein, "The 95.4 GeV di-photon excess at ATLAS and CMS," [arXiv:2306.03889 [hep-ph]].
- [107] G. Belanger, U. Ellwanger, J. F. Gunion, Y. Jiang, S. Kraml and J. H. Schwarz, JHEP 01 (2013), 069 [arXiv:1210.1976 [hep-ph]].
- [108] T. Biekötter, S. Heinemeyer and C. Muñoz, Eur. Phys. J. C 78 (2018) no.6, 504 [arXiv:1712.07475 [hep-ph]].
- [109] F. Domingo, S. Heinemeyer, S. Paßehr and G. Weiglein, Eur. Phys. J. C 78 (2018) no.11, 942 [arXiv:1807.06322 [hep-ph]].
- [110] W. G. Hollik, S. Liebler, G. Moortgat-Pick, S. Paßehr and G. Weiglein, Eur. Phys. J. C 79 (2019) no.1, 75 [arXiv:1809.07371 [hep-ph]].
- [111] K. Wang, F. Wang, J. Zhu and Q. Jie, Chin. Phys. C 42 (2018) no.10, 103109-103109 [arXiv:1811.04435 [hep-ph]].
- [112] K. Choi, S. H. Im, K. S. Jeong and C. B. Park, Eur. Phys. J. C 79 (2019) no.11, 956 [arXiv:1906.03389 [hep-ph]].
- [113] J. Cao, X. Jia, Y. Yue, H. Zhou and P. Zhu, Phys. Rev. D 101 (2020) no.5, 055008 [arXiv:1908.07206 [hep-ph]].
- [114] W. G. Hollik, C. Li, G. Moortgat-Pick and S. Paasch, Eur. Phys. J. C 81 (2021) no.2, 141 [arXiv:2004.14852 [hep-ph]].
- [115] T. Biekötter, A. Grohsjean, S. Heinemeyer, C. Schwanenberger and G. Weiglein, Eur. Phys. J. C 82 (2022) no.2, 178 [arXiv:2109.01128 [hep-ph]].
- [116] A. Adhikary, R. K. Barman, B. Bhattacherjee, A. De, R. M. Godbole and S. Kulkarni, Phys. Rev. D 108 (2023) no.3, 035020 [arXiv:2207.00600 [hep-ph]].
- [117] W. Li, J. Zhu, K. Wang, S. Ma, P. Tian and H. Qiao, "A light Higgs boson in the NMSSM confronted with the CMS di-photon and di-tau excesses," [arXiv:2212.11739 [hep-ph]].
- [118] U. Ellwanger and C. Hugonie, Eur. Phys. J. C 83 (2023) no.12, 1138 [arXiv:2309.07838 [hep-ph]].
- [119] J. Cao, X. Jia, J. Lian and L. Meng, "95 GeV Diphoton and bb Excesses in the General Next-to-Minimal Supersymmetric Standard Model," [arXiv:2310.08436 [hep-ph]].
- [120] W. Li, H. Qiao, K. Wang and J. Zhu, "Light dark matter confronted with the 95 GeV diphoton excess," [arXiv:2312.17599 [hep-ph]].
- [121] S. Roy and C. E. M. Wagner, "Dark Matter searches with photons at the LHC," [arXiv:2401.08917 [hep-ph]].
- [122] U. Ellwanger, J. F. Gunion and C. Hugonie, JHEP 02 (2005), 066 [arXiv:hep-ph/0406215 [hep-ph]].

- [123] U. Ellwanger and C. Hugonie, Comput. Phys. Commun. 175 (2006), 290-303 [arXiv:hep-ph/0508022 [hep-ph]].
- [124] https://www.lupm.in2p3.fr/users/nmssm/index.html
- [125] G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, Comput. Phys. Commun. 185 (2014), 960-985 [arXiv:1305.0237 [hep-ph]].
- [126] G. W. Bennett et al. [Muon g-2], Phys. Rev. D 73 (2006), 072003 [arXiv:hep-ex/0602035 [hep-ex]].
- [127] B. Abi et al. [Muon g-2], Phys. Rev. Lett. 126 (2021) no.14, 141801 [arXiv:2104.03281 [hep-ex]].
- [128] F. Domingo, U. Ellwanger and C. Hugonie, Eur. Phys. J. C 82 (2022) no.11, 1074 [arXiv:2209.03863 [hep-ph]].
- [129] S. Kraml, S. Kulkarni, U. Laa, A. Lessa, W. Magerl, D. Proschofsky-Spindler and W. Waltenberger, Eur. Phys. J. C 74 (2014), 2868 [arXiv:1312.4175 [hep-ph]].
- [130] J. Dutta, S. Kraml, A. Lessa and W. Waltenberger, LHEP 1 (2018) no.1, 5-12 [arXiv:1803.02204 [hep-ph]].
- [131] C. K. Khosa, S. Kraml, A. Lessa, P. Neuhuber and W. Waltenberger, LHEP 2020 (2020), 158 [arXiv:2005.00555 [hep-ph]].
- [132] G. Alguero, J. Heisig, C. K. Khosa, S. Kraml, S. Kulkarni, A. Lessa, H. Reyes-González, W. Waltenberger and A. Wongel, JHEP 08 (2022), 068 [arXiv:2112.00769 [hep-ph]].
- [133] G. Aad *et al.* [ATLAS], "A statistical combination of ATLAS Run 2 searches for charginos and neutralinos at the LHC," [arXiv:2402.08347 [hep-ex]].
- [134] A. Tumasyan *et al.* [CMS], Nature **607** (2022) no.7917, 60-68 [arXiv:2207.00043 [hep-ex]].
- [135] [ATLAS], Nature 607 (2022) no.7917, 52-59 [erratum: Nature 612 (2022) no.7941, E24] [arXiv:2207.00092 [hep-ex]].
- [136] W. Abdallah, A. Datta and S. Roy, JHEP **04** (2021), 122 [arXiv:2012.04026 [hep-ph]].
- [137] A. Datta, M. Guchait, A. Roy and S. Roy, JHEP 11 (2023), 081 [arXiv:2211.05905 [hepph]].
- [138] A. Hayrapetyan *et al.* [CMS], "Search for exotic decays of the Higgs boson to a pair of pseudoscalars in the $\mu\mu$ bb and $\tau\tau$ bb final states," [arXiv:2402.13358 [hep-ex]].