QUANTUM TRANSITION PROBABILITIES

STAN GUDDER

ABSTRACT. Transition probabilities are an important and useful tool in quantum mechanics. However, in their present form, they are limited in scope and only apply to pure quantum states. In this article we extend their applicability to mixed states and to transitions between quantum effects. We also present their dependence on a measured operation or instrument. We begin by defining our concepts on a general quantum effect algebra. These concepts are illustrated using Holevo operations and instruments. We then present transition probabilities in the special case of the Hilbert space formulation of quantum mechanics. We show that for pure states and particular types of operations the transition probabilities reduce to their usual form. We give examples in terms of Lüders operations and instruments.

1. INTRODUCTION

In the usual Hilbert-space formulation of quantum mechanics, an important and useful role is played by transition probabilities [6,10]. For example, suppose a quantum system is described by a Hilbert space H and the system is in a pure state given by a unit vector $\psi \in H$. We now measure the energy of the system which is described by a self-adjoint operator A on H. The energy values are given by eigenvalues λ_i of A with corresponding unit eigenvectors ϕ_i so $A\phi_i = \lambda_i\phi_i$, $i = 1, 2, \dots$ According to the axioms of quantum mechanics, the probability that the energy of the system is λ_i becomes $P(\psi, \phi_i) = |\langle \psi, \phi_i \rangle|^2$ [6,10]. We call $P(\psi, \phi_i)$ the transition proba*bility* of ψ to ϕ_i . This definition is appropriate because $\{\phi_i : i = 1, 2, ...\}$ is an orthonormal basis for H so we have $\sum |\langle \psi, \phi_i \rangle|^2 = 1$. Also, another axiom of quantum mechanics says that if the energy value observed is λ_i then the state ψ updates to the state ϕ_i [6,10]. Thus, $P(\psi, \phi_i)$ is the probability that this update occurs. In general, if $\psi, \phi \in H$ specify two states then calling $P(\psi, \phi) = |\langle \psi, \phi \rangle|^2$ the transition probability of ψ to ϕ does not stress the fact that this transition is caused by a measurement of an observable A. Also, what about mixed states described by density operators on H? Moreover, what about more general measurements such as operations and instruments [1, 2, 8]?

In Section 2, we study probabilistic models that generalize quantum mechanics. We begin by introducing an effect algebra E [1,2]. An effect $a \in E$ is a yes-no (true-false) experiment that either occurs (yes) or does not occur (no) when measured. For $a, b \in E$ there is a partial operation $a \oplus b$ that describes a statistical sum of a, b (when it is defined). A substate on E is function $s: E \to [0,1] \subseteq \mathbb{R}$ that satisfies $s(a \oplus b) = s(a) + s(b)$ when $a \oplus b$ is defined. There is a special effect $1 \in E$ that is always true and a substate s is a state if s(1) = 1. We denote the set of states on E by $\mathcal{S}(E)$ and the set of substates by $\operatorname{Sub}(E)$. If $s \in \mathcal{S}(E)$, then s(a) is the probability that aoccurs when the system is in state s. A convex map $\mathcal{I}: \operatorname{Sub}(E) \to \operatorname{Sub}(E)$ is called an operation [1,2]. An operation corresponds to an apparatus that can be employed to measure an effect and we denote the set of operations by $\mathcal{O}(E)$. If $s \in \mathcal{S}(E), \ \mathcal{I} \in \mathcal{O}(E)$ with $\mathcal{I}(s) \neq 0, a, b \in E$ we define the transition probability of a to b relative to $s, \ \mathcal{I}$ by

$$P_{s,\mathcal{I}}(a,b) = \frac{s(a)}{\mathcal{I}(s)(1)} \mathcal{I}(s)(b)$$

If $\mathcal{I} \in \mathcal{O}(E)$, $s_1, s_2 \in \mathcal{S}(E)$ we define the transition probability of s_1 to s_2 relative to \mathcal{I} by

$$P_{\mathcal{I}}(s_1, s_2) = \mathcal{I}(s_1)(1)\mathcal{I}\left(\mathcal{I}(s_2)(1)\right)$$

Section 2 derives properties of $P_{s,\mathcal{I}}(a,b)$ and $P_{\mathcal{I}}(s_1,s_2)$. We also define repeatable operations and discuss their properties. Moreover, Section 2 discusses generalizations of effects and operations called observables and instruments, respectively. An observable is an effect-valued measure $A_x \in E$ where A_x is the effect that occurs when the observable has the outcome xupon being measure [3,4]. An instrument is an operation-valued measure $\mathcal{I}_x \in \mathcal{O}(E)$ where \mathcal{I}_x is the operation that occurs when the instrument has the outcome x upon being measured [3–5]. We then have the transition probabilities $P_{\mathcal{I}_x}(s_1, s_2)$. If A_x, B_y are observables, we have the transition probabilities $P_{\mathcal{I}_x}(A_x, B_y)$.

Section 2 ends with a discussion of Holevo operations and instruments which are used to illustrate the previously introduced concepts [7]. If $\alpha \in S(E)$, $a \in E$, a pure Holevo operation has the form $\mathcal{H}^{(a,\alpha)}(s) = s(a)\alpha$. We also define mixed Holevo operations and Holevo instruments in a natural way [7]. Transition probabilities relative to these operations and instruments are computed. We also find updated states after these operations and instruments are measured.

Section 3 specializes the theory of effect algebras to the traditional Hilbert space formulation of quantum mechanics. In this formulation, effects are described by operators A satisfying $0 \le A \le I$ and states are described by effects ρ with trace tr (ρ) = 1. We call ρ a *density operator* and the corresponding state satisfies $s(A) = \operatorname{tr}(\rho A)$. In this formalism an operation is a completely positive trace nonincreasing linear map from the set of density operators to the set of positive operators on the Hilbert space H [1, 2, 6, 10]. We point out that every operation has a Kraus decomposition $\mathcal{I}(\rho) = \sum K_i \rho K_i^*$ where K_i are linear operators on H satisfying $\sum K_i^* K_i \leq I$. In particular, we study Lüders operations which have the form $\mathcal{I}(\rho) = A^{1/2} \rho A^{1/2}$ where A is an effect [9]. Lüders instruments are defined in the natural way. We then compute transition probabilities relative to these operations and instruments.

2. Basic Concepts and Definitions

An effect algebra $\mathcal{E} = (E, 0, 1, \oplus)$ [1,3,4,6] is a set of effects E with two special elements $0, 1 \in E$ and a partial binary operation \oplus satisfying the following conditions:

(EA1) if $a \oplus b$ is defined, then $b \oplus a$ is defined and $b \oplus a = a \oplus b$, (EA2) if $a \oplus (b \oplus a)$ is defined then $(a \oplus b) \oplus a$ is defined and

(EA2) if $a \oplus (b \oplus c)$ is defined, then $(a \oplus b) \oplus c$ is defined and

$$(a \oplus b) \oplus c = a \oplus (b \oplus c)$$

(EA3) for every $a \in E$, there exists a unique effect $a' \in E$ such that $a \oplus a' = 1$,

(EA4) if $a \oplus 1$ is defined, then a = 0.

An effect corresponds to a *yes-no* experiment that either occurs or does not occur when it is measured. The effect 0 never occurs and the effect 1 always occurs when measured. The *complement* a' of a occurs if and only if a does not occur. The sum $a \oplus b$ corresponds to a statistical sum of a and b when it is defined. A substate on \mathcal{E} is a function $s \colon E \to [0,1] \subseteq \mathbb{R}$ that satisfies $s(a \oplus b) = s(a) + s(b)$ whenever $a \oplus b$ is defined. A substate s that satisfies s(1) = 1 is called a *state*. We denote the set of substates by $\operatorname{Sub}(\mathcal{E})$ and the set of states by $\mathcal{S}(\mathcal{E})$. A state s describes the initial condition of a physical system and s(a) is the probability that a occurs when the system is in the state s. If $s \in \mathcal{S}(\mathcal{E})$, $\lambda \in [0, 1]$, then λs defined by $(\lambda s)(a) = \lambda s(a)$ is a substate. Conversely, if $s \in \operatorname{Sub}(\mathcal{E})$ and $s \neq 0$, then $\tilde{s}(a) = s(a)/s(1)$ is a state.

A function $J: \operatorname{Sub}(\mathcal{E}) \to \operatorname{Sub}(\mathcal{E})$ that satisfies $J\left(\sum_{i=1}^{n} \lambda_i s_i\right) = \sum_{i=1}^{n} \lambda_i J(s)$ when $\sum_{i=1}^{n} \lambda_i \leq 1$ is called a *convex function* or an *operation* [1, 2, 8]. We denote the set of operations on \mathcal{E} by $\mathcal{O}(\mathcal{E})$ and it follows that if $J \in \mathcal{O}(\mathcal{E})$ then $J(\lambda s) = \lambda J(s)$ for all $\lambda \in [0, 1]$, $s \in \operatorname{Sub}(\mathcal{E})$. If $J \in \mathcal{O}(\mathcal{E})$ satisfies $J(s) \in \mathcal{S}(\mathcal{E})$ for all $s \in \mathcal{S}(\mathcal{E})$, then J is a *channel* [1, 6, 10]. We say that $J \in \mathcal{O}(\mathcal{E})$ measures $a \in E$ if s(a) = J(s)(1) for all $s \in \mathcal{S}(\mathcal{E})$. We say that

STAN GUDDER

 $\mathcal{S}(\mathcal{E})$ is separating if s(a) = s(b) for all $s \in \mathcal{S}(\mathcal{E})$ implies a = b. If $\mathcal{S}(\mathcal{E})$ is separating, then J measures the unique effect \widehat{J} satisfying $s(\widehat{J}) = J(s)(1)$ for all $s \in \mathcal{S}(\mathcal{E})$. We then consider J as an apparatus that measures the effect \widehat{J} . As with effects, an operation is considered to be a *yes-no* measurement. If $s \in \mathcal{S}(\mathcal{E})$, then J(s)(1) is the *probability* J occurs when the system is in the state s. If $J(s) \neq 0$, we call $J(s)^{\sim} = J(s)/J(s)(1) \in \mathcal{S}(\mathcal{E})$ the *updated state* after J is measured where $0 \in \operatorname{Sub}(\mathcal{E})$ means 0(a) = 0 for all $a \in E$.

If $s \in \mathcal{S}(\mathcal{E})$, $J \in \mathcal{O}(\mathcal{E})$ with $J(s) \neq 0$ and $a, b \in E$, the transition probability of a then b relative to s, J is

$$P_{s,J}(a,b) = \frac{s(a)}{J(s)(1)} J(s)(b) = s(a)J(s)^{\sim}(b)$$

Thus, $P_{s,J}(a, b)$ is the probability that a occurs in the state s times the probability that b occurs in the updated state $J(s)^{\sim}$ after J is measured. If $\widehat{\mathcal{I}} = a$, then $P_{s,\mathcal{I}}(a,b) = \mathcal{I}(s)(b)$. Notice that $P_{s,J}(a, \bullet) \in \mathrm{Sub}(\mathcal{E})$ and $P_{s,J}(\bullet,b) \in \mathrm{Sub}(\mathcal{E})$. Moreover, $P_{s,J}(a,1) = s(a)$ for all $a \in E$, $P_{s,J}(1,b) = \mathcal{I}(s)^{\sim}(b)$ for all $b \in E$ and $P_{s,J}(1,1) = 1$.

If $f, g: S \to S$ are functions, their composition $f \circ g: S \to S$ is $f \circ g(t) = f(g(t))$. We say that $J \in \mathcal{O}(\mathcal{E})$ is repeatable if $J \circ J = J$. If $J \in \mathcal{O}(\mathcal{E})$, $s_1, s_2 \in \mathcal{S}(\mathcal{E})$, the transition probability of s_1 to s_2 relative to J is

$$P_J(s_1, s_2) = J(s_1)(1)J \circ J(s_2)(1) = J[J(s_1)(1)J(s_2)](1)$$

Thus, $P_{\mathcal{J}}(s_1, s_2)$ is the probability J occurs when the system is in state s_1 times the probability the composition $J \circ J$ occurs when the system is in state s_2 . We see that if J is repeatable, then

$$P_J(s_1, s_2) = J(s_1)(1)J(s_2)(1)$$

in which case $P_J(s_1, s_2) = P_J(s_2, s_1)$ which does not happen in general. Moreover, if J also measures a, then $P_J(s_1, s_2) = s_1(a)s_2(a)$. The following lemma shows the relationship between these two types of transition probabilities. We denote the product of two real-valued functions f, g by $f \bullet g(x) = f(x)g(x)$.

Lemma 2.1. If $\mathcal{S}(\mathcal{E})$ is separating, then $P_J(s_1, s_2) = P_{s_1,J} \bullet P_{J(s_2),J}(\widehat{J}, 1)$.

Proof. Since

$$s_2 \left[(J \circ J)^{\wedge} \right] = (J \circ J)(s_2)(1) = J \left[J(s_2) \right](1) = P_{J(s_2),J}(\widehat{J}, 1)$$

we obtain

$$P_J(s_1, s_2) = s_1(\widehat{J}) s_2 \left[(J \circ J)^{\wedge} \right] = P_{s_1, J}(\widehat{J}, 1) P_{J(s_2), J}(\widehat{J}, 1)$$

= $P_{s_1, J} \bullet P_{J(s_2), J}(\widehat{J}, 1)$

An observable on \mathcal{E} is a finite set of effects $A = \{A_x : x \in \Omega_A\}, A_x \in E$, such that $\sum_{x \in \Omega_A} A_x = 1$. We call Ω_A the outcome space of A and A_x is the effect that occurs when A has outcome $x \in \Omega_A$. For $\Delta \subseteq \Omega_A$ we define the effect-valued measure $A(\Delta) = \sum_{x \in \Delta} A_x$ [1, 6]. We see that $A(\Omega_A) = 1$ and $A(\Delta_1 \cup \Delta_2) = A(\Delta_1) + A(\Delta_2)$ when $\Delta_1 \cap \Delta_2 = \emptyset$. We denote the set of observables on \mathcal{E} by Ob (\mathcal{E}). If $s \in \mathcal{S}(\mathcal{E}), A \in \text{Ob}(\mathcal{E})$, the distribution of A in the state s is the probability measure $\Phi_s^A(\Delta) = \sum_{x \in \Delta} s(A_x)$ on Ω_A . Let $A, B \in \text{Ob}(\mathcal{E}), s \in \mathcal{S}(\mathcal{E}), J \in \mathcal{O}(\mathcal{E})$ with $J(s) \neq 0$ and consider the transition probability

$$P_{xy} = P_{s,J}(A_x, B_y) = \frac{s(A_x)}{J(s)(1)} J(s)(B_y)$$

For $\Delta \subseteq \Omega_A \times \Omega_B$ we define $P(\Delta) = \sum \{ P_{xy} \colon (x, y) \in \Delta \}.$

Lemma 2.2. *P* is a probability measure on $\Omega_A \times \Delta_B$ with marginals

$$P^{1}(\Delta_{1}) = P(\Delta_{1} \times \Omega_{B}) = s [A(\Delta_{1})] = \Phi_{s}^{A}(\Delta_{1}) \text{ where } \Delta_{1} \subseteq \Omega_{A}$$
$$P^{2}(\Delta_{2}) = P(\Omega_{A} \times \Delta_{2}) = J(s)^{\sim} [B(\Delta_{2})] = \Phi_{J(s)^{\sim}}^{B}(\Delta_{2}) \text{ where } \Delta_{2} \subseteq \Omega_{B}$$

Proof. We have that

$$P(\Omega_A \times \Omega_B) = \sum \{P_{xy} \colon (x, y) \in \Omega_A \times \Omega_B\}$$

= $\frac{1}{J(s)(1)} \sum \{s(A_x)J(s)(B_y) \colon (x, y) \in \Omega_A \times \Omega_B\}$
= $\frac{1}{J(s)(1)} \sum_{x \in \Omega_A} s(A_x) \sum_{y \in \Omega_B} J(s)(B_y) = \frac{1}{J(s)(1)} J(s)(1) = 1$

If $\Delta_1, \Delta_2 \subseteq \Omega_A \times \Omega_B$ with $\Delta_1 \cap \Delta_2 = \emptyset$, then

$$P(\Delta_1 \cup \Delta_2) = \sum \{P_{xy} \colon (x, y) \in \Delta_1 \cup \Delta_2\}$$

=
$$\sum \{P_{xy} \colon (x, y) \in \Delta_1\} + \sum \{P_{xy} \colon (x, y) \in \Delta_2\}$$

=
$$P(\Delta_1) + P(\Delta_2)$$

Hence, P is a probability measure on $\Omega_A \times \Omega_B$. The marginals of P become

$$P^{1}(\Delta_{1}) = P(\Delta_{1} \times \Omega_{B}) = \sum_{(x,y) \in \Delta_{1} \times \Omega_{B}} \frac{s(A_{x})}{J(s)(1)} J(s)(B_{y}) = s \left[A(\Delta_{1})\right] = \Phi_{s}^{A}(\Delta_{1})$$

where $\Delta_1 \subseteq \Omega_A$ and

$$P^{2}(\Delta_{2}) = P(\Omega_{A} \times \Delta_{2}) = \sum_{(x,y)\in\Omega_{A}\times\Delta_{2}} \frac{s(A_{x})}{J(s)(1)} J(s)(B_{y})$$
$$= J(s)^{\sim} [B(\Delta_{2})] = \Phi^{B}_{J(s)^{\sim}}(\Delta_{2})$$

where $\Delta_2 \subseteq \Omega_B$.

An instrument \mathcal{I} on \mathcal{E} is a finite set of operations $\mathcal{I} = \{\mathcal{I}_x : x \in \Omega_{\mathcal{I}}\}, \mathcal{I}_x \in \mathcal{O}(\mathcal{E})$ such that $\sum_{x \in \Omega_{\mathcal{I}}} \mathcal{I}_x$ is a channel [1,2,5,6]. We call $\Omega_{\mathcal{I}}$ the outcome space of \mathcal{I} . If $s \in \mathcal{S}(\mathcal{E})$, then $\mathcal{I}_x(s)(1)$ is the probability that the outcome x occurs when \mathcal{I} is measured and the system is in state s. Moreover, we call $\Phi_s^{\mathcal{I}}(\Delta) = \sum_{x \in \Delta} \mathcal{I}_x(s)(1)$ the distribution of \mathcal{I} in the state s. Since $\overline{\mathcal{I}} = \sum_{x \in \Omega_{\mathcal{I}}} \mathcal{I}_x$ is a channel we have

$$\Phi_s^{\mathcal{I}}(\Omega_{\mathcal{I}}) = \sum_{x \in \Omega_{\mathcal{I}}} \mathcal{I}_x(s)(1) = \left[\sum_{x \in \Omega_{\mathcal{I}}} \mathcal{I}_x(s)\right](1) = 1$$

As in Lemma 2.2, if $\Delta_1, \Delta_2 \subseteq \Omega_{\mathcal{I}}$ and $\Delta_1 \cap \Delta_2 = \emptyset$, then

$$\Phi_s^{\mathcal{I}}(\Delta_1 \cup \Delta_2) = \Phi_s^{\mathcal{I}}(\Delta_1) + \Phi_s^{\mathcal{I}}(\Delta_2)$$

so $\Phi_s^{\mathcal{I}}$ is a probability measure on $\Omega_{\mathcal{I}}$. We denote the set of instruments on \mathcal{E} by In (\mathcal{E}). If $\mathcal{I} \in \text{In}(\mathcal{E})$, we say that \mathcal{I} measures an observable $A \in \text{Ob}(\mathcal{E})$ with $\Omega_A = \Omega_{\mathcal{I}}$ if $\mathcal{I}_x(s)(1) = s(A_x)$ for all $s \in \mathcal{S}(\mathcal{E}) \ x \in \Omega_{\mathcal{I}}$. If \mathcal{E} has a separating set of states, A is unique and we write $A = \widehat{\mathcal{I}}$.

If $\mathcal{I} \in \text{In}(\mathcal{E})$ and $A, B \in \text{Ob}(\mathcal{E})$ we obtain the transition probability

$$P_{s,\mathcal{I}_z}(A_x, B_y) = \frac{s(A_x)}{\mathcal{I}_z(s)(1)} \mathcal{I}_z(s)(B_y)$$

Multiplying by $\mathcal{I}_z(s)(1)$, we have the function on $\Omega_{\mathcal{I}} \times \Omega_A \times \Omega_B$ given by

$$Q_{s,\mathcal{I}_z}(A_x, B_y) = \mathcal{I}(s)(1)P_{s,\mathcal{I}_z}(A_x, B_y) = s(A_x)\mathcal{I}_x(s)(B_y)$$

Since

$$\sum \left\{ Q_{s,\mathcal{I}_z}(A_x B_y) \colon (z, x, y) \in \Omega_\mathcal{I} \times \Omega_A \times \Omega_B \right\} = s(1)\overline{\mathcal{I}}(s)(1) = 1$$

we conclude that Q determines a probability measure on $\Omega_{\mathcal{I}} \times \Omega_A \times \Omega_B$. The marginals of Q become

$$Q^{2}(x) = s(A_{x}) \sum \{ \mathcal{I}_{z}(s)(B_{y}) \colon (z, y) \in \Omega_{\mathcal{I}} \times \Omega_{B} \}$$
$$= s(A_{x})\overline{\mathcal{I}}(s)(1) = s(A_{x})$$

so $Q^2(\Delta) = \Phi_s^A(\Delta), \ \Delta \subseteq \Omega_A,$ $Q^3(y) = \sum \{s(A_x)\mathcal{I}_z(s)(B_y) \colon (z,x) \in \Omega_\mathcal{I} \times \Omega_A\}$ $= \overline{\mathcal{I}}(s)(B_y)$

so $Q^3(\Delta) = \Phi^B_{\overline{\mathcal{I}}(s)}(\Delta), \ \Delta \subseteq \Omega_B,$

$$Q^{1}(z) = \sum \{ s(A_{x})\mathcal{I}_{z}(s)(B_{y}) \colon (x,y) \in \Omega_{A} \times \Omega_{B} \}$$
$$= \mathcal{I}_{z}(s)(1)$$

so $Q^1(\Delta) = \Phi_s^{\mathcal{I}}(\Delta), \ \Delta \subseteq \Omega_{\mathcal{I}}$. We conclude that $Q^1, \ Q^2, \ Q^3$ give the distributions of \mathcal{I} in the state s, A in the state s and B in the state $\overline{\mathcal{I}}(s)$.

If $\mathcal{I} \in \text{In}(\mathcal{E})$, we define the *transition probability* from s_1 to s_2 relative to \mathcal{I} and $(x, y) \in \Omega_{\mathcal{I}} \times \Omega_{\mathcal{I}}$ by

$$(P_{\mathcal{I}})_{(x,y)}(s_1,s_2) = \mathcal{I}_x(s_1)(1)\mathcal{I}_y\left(\overline{\mathcal{I}}(s_2)\right)(1)$$

We see that $P_{\mathcal{I}}$ is a probability measure on $\Omega_{\mathcal{I}} \times \Omega_{\mathcal{I}}$ because

$$\sum_{x,y} (P_{\mathcal{I}})_{(x,y)}(s_1, s_2) = \overline{\mathcal{I}}(s_1)(1)\overline{\mathcal{I}}\left(\overline{\mathcal{I}}(s_2)\right)(1) = 1$$

The marginals of $P_{\mathcal{I}}$ become

$$\sum_{y \in \Omega_{\mathcal{I}}} (P_{\mathcal{I}})_{(x,y)}(s_1, s_2) = \mathcal{I}_x(s_1)(1)$$

which is the probability of outcome x when \mathcal{I} is measured in the state s_1

$$\sum_{x \in \Omega_{\mathcal{I}}} (P_{\mathcal{I}})_{(x,y)}(s_1, s_2) = \mathcal{I}_y \overline{\mathcal{I}}(s_2)(1)$$

which is the probability of outcome y when \mathcal{I} is measured in the state $\overline{\mathcal{I}}(s_2)$.

We now illustrate the previous theory with a particular type of operation and instrument. If $\alpha \in \mathcal{S}(\mathcal{E})$, $a \in E$, we call $\mathcal{H}^{(a,\alpha)} \in \mathcal{O}(E)$ defined by $\mathcal{H}^{(a,\alpha)}(s) = s(a)\alpha$ a pure Holevo operation [7]. Since $[\mathcal{H}^{(a,\alpha)}(s)](1) = s(a)$ for all $s \in \mathcal{S}(\mathcal{E})$ we see that $\mathcal{H}^{(a,\alpha)}$ measures a. If $A \in Ob(\mathcal{E})$ and $\alpha_x \in \mathcal{S}(\mathcal{E})$ for $x \in \Omega_A$ we call

$$\mathcal{H}^{(A,\{\alpha\})}(s) = \sum_{x \in \Omega_A} s(A_x) \alpha_x$$

a mixed Holevo operation [7]. For $s \in \mathcal{S}(\mathcal{E})$ we have

$$\sum_{x \in \Omega_A} s(A_x) = s\left(\sum_{x \in \Omega_A} A_x\right) = s(1) = 1$$

so $\mathcal{H}^{(A,\{\alpha\})}$ is a channel. We also call

$$\mathcal{H}_x^{(A,\{\alpha\})}(s) = s(A_x)\alpha_x$$

a Holevo instrument [7]. Notice that $\mathcal{H}_x^{(A,\{\alpha\})}$ is an instrument because

$$\sum_{x \in \Omega_A} \left[\mathcal{H}_x^{(A,\{\alpha\})}(s) \right] (1) = \sum_{x \in \Omega_A} s(A_x) = 1$$

Also, $\mathcal{H}_x^{(A,\{\alpha\})}$ measures A because $\left[\mathcal{H}_x^{(A,\{\alpha\})}(s)\right](1) = s(A_x)$ for all $s \in \mathcal{S}(\mathcal{E})$ and $x \in \Omega_A$.

For a pure Holevo operation $\mathcal{H}^{(a,\alpha)}$ and $b, c \in E$, if $s(a) \neq 0$ we have

$$P_{s,\mathcal{H}^{(a,\alpha)}}(b,c) = \frac{s(b)}{\left[\mathcal{H}^{(a,\alpha)}(s)\right](1)} \left[\mathcal{H}^{(a,\alpha)}(s)\right](c) = \frac{s(b)}{s(a)}s(a)\alpha(c) = s(b)\alpha(c)$$

We see that $P_{s,\mathcal{H}^{(a,\alpha)}}$ is independent of a if $s(a) \neq 0$. For a mixed Holevo operation we have

$$\left[\mathcal{H}^{(A,\{\alpha\})}(s)\right](c) = \sum_{x \in \Omega_A} s(A_x)\alpha_x(c)$$

so that

$$P_{s,\mathcal{H}^{(A,\{\alpha\})}}(b,c) = s(b) \sum_{x \in \Omega_A} s(A_x) \alpha_x(c)$$

The state transition probability becomes

$$P_{\mathcal{H}^{(a,\alpha)}}(s_1, s_2) = \mathcal{H}^{(a,\alpha)}(s_1)(1)\mathcal{H}^{(a,\alpha)}\left[\mathcal{H}^{(a,\alpha)}(s_2)\right](1)$$
$$= s_1(a)\mathcal{H}^{(a,\alpha)}\left[s_2(a)\alpha\right](1) = s_1(a)s_2(a)\left[\mathcal{H}^{(a,\alpha)}(\alpha)\right](1)$$
$$= s_1(a)s_2(a)\alpha(a)$$

Since a mixed Holevo operation is a channel, we have $P_{\mathcal{H}^{(A,\{\alpha\})}}(s_1, s_2) = 1$ for all $s_1, s_2 \in \mathcal{S}(\mathcal{E})$. For Holevo instruments we have

$$P_{s,\mathcal{H}_{x}^{(A,\{\alpha\})}}(b,c) = \frac{s(b)}{\left[\mathcal{H}_{x}^{(A,\{\alpha\})}(s)\right](1)} \left[\mathcal{H}_{x}^{(A,\{\alpha\})}(s)\right](c) = \frac{s(b)}{s(A_{x})}s(A_{x})\alpha_{x}(c)$$
$$= s(b)\alpha_{x}(c)$$

and

$$P_{\mathcal{H}_{x}^{(A,\{x\})}}(s_{1},s_{2}) = \mathcal{H}_{x}^{(A,\{\alpha\})}(s_{1})(1)\mathcal{H}_{x}^{(A,\{x\})}\left[\mathcal{H}_{x}^{(A,\{\alpha\})}(s_{2})\right](1)$$
$$= s_{1}(A_{x})\mathcal{H}_{x}^{(A,\{\alpha\})}\left[s_{2}(A_{x})\alpha_{x}\right](1) = s_{1}(A_{x})s_{2}(A_{x})\alpha_{x}(A_{x})$$

An operation $\mathcal{I} \in \mathcal{O}(\mathcal{E})$ is repeatable if $\mathcal{I}(\mathcal{I}(s)) = \mathcal{I}(s)$ for all $s \in \mathcal{S}(\mathcal{E})$. Thus, $\mathcal{I} \circ \mathcal{I} = \mathcal{I}$ so we can repeat \mathcal{I} without changing the state. If \mathcal{I} is repeatable, the transition probability satisfies

$$P_{\mathcal{I}}(s_1, s_2) = \mathcal{I}(s_1)(1)\mathcal{I}\left(\mathcal{I}(s_2)\right)(1) = \mathcal{I}(s_1)(1)\mathcal{I}(s_2)(1) = P_{\mathcal{I}}(s_2, s_1)$$

We now show that the converse does not hold. We have seen that

$$P_{\mathcal{H}^{(a,\alpha)}}(s_1,s_2) = P_{\mathcal{H}^{(a,\alpha)}}(s_2,s_1)$$

However, if $s(a) \neq 0, 1$ we obtain

$$\mathcal{H}^{(a,\alpha)}\left(\mathcal{H}^{(a,\alpha)}(s)\right) = \mathcal{H}(a,\alpha)\left(s(a)\alpha\right) = s(a)\mathcal{H}^{(a,\alpha)}(\alpha)$$
$$= s(a)\alpha(a)\alpha \neq s(a)\alpha = \mathcal{H}^{(a,\alpha)}(s)$$

Hence, if $s(a) \neq 0$, then $\mathcal{H}^{(a,\alpha)}$ is repeatable if and only if s(a) = 1.

3. HILBERT SPACE TRANSITION PROBABILITIES

In this section we consider Hilbert space effects, operations, instruments, states and their transition probabilities. These are the basic concepts for the standard quantum probability theory. Let H be a complex Hilbert space which for simplicity, we take to be finite dimensional. Letting $\mathcal{L}_+(H)$ be the set of positive linear operators on H, an *effect* is represented by an operator $a \in \mathcal{L}_+(H)$ satisfying $0 \leq a \leq I$ where 0, I are the zero and identity operators respectively [1,2,6]. The set of effects is denoted by $\mathcal{E}(H)$. A *density operator* $\rho \in \mathcal{L}_+(H)$ satisfies tr $(\rho) = 1$ and we denote the set of density operators on H by $\mathcal{D}(H)$. According to Born's rule, every state $s \in \mathcal{S}(H)$ has the form $s(a) = \operatorname{tr}(\rho a)$ for a unique $\rho \in \mathcal{D}(H)$. This can actually be proved using Gleason's theorem [6] so we will identify $\mathcal{S}(H)$ and $\mathcal{D}(H)$. In the Hilbert space formulation of quantum mechanics, an *operation* is a completely positive linear map $J : \mathcal{D}(H) \to \mathcal{L}_+(H)$ satisfying tr $[J(\rho)] \leq \operatorname{tr}(\rho)$ for all $\rho \in \mathcal{D}(H)$ [1,2,6,10]. We denote the set of operations H by $\mathcal{O}(H)$. If tr $[J(\rho)] = 1$ for all $\rho \in \mathcal{D}(H)$, then J is a *channel*.

Denoting the set of linear operators on H by $\mathcal{L}(H)$, it can be shown that every $J \in \mathcal{O}(H)$ has a Kraus decomposition $J(\rho) = \sum_{i=1}^{n} K_i \rho K_i^*$ where $K_i \in \mathcal{L}(H)$ with $\sum_{i=1}^{n} K_i^* K_i \leq I$ [6,8,10]. Notice that J is a channel if and only if $\sum_{i=1}^{n} K_i^* K_i = I$. The operators K_i are called Kraus operators for J and they are not unique. If J has the form $J(\rho) = a^{1/2} \rho a^{1/2}$ where $a \in \mathcal{E}(H)$ we call J a Lüders operation. If the Kraus operators are projections, it follows that $K_i K_j = 0$ for $i \neq j$ and hence, in this case, J is repeatable. Conversely, if $J(\rho) = a^{1/2} \rho a^{1/2}$ is a repeatable Lüders operation then

$$a^{1/2}\rho a^{1/2} = J(\rho) = J(J(\rho)) = a\rho a$$

for every $\rho \in \mathcal{D}(H)$. Letting $\rho = \frac{1}{n}I$ where $n = \dim H$, we conclude that $a = a^2$ so a is a projection.

In the Hilbert space case, it is well-known that $\mathcal{S}(H)$ is separating. It follows that an operation $J \in \mathcal{O}(H)$ measures a unique effect $\hat{\mathcal{J}} \in \mathcal{E}(H)$ satisfying tr $(\rho \hat{J}) = \text{tr } [J(\rho)]$ for all $\rho \in \mathcal{D}(H)$. Notice that if J is a channel, then $\hat{J} = I$. Although an operation measures a unique effect, as we shall see, every effect is measured by many operations. If J has Kraus decomposition $J(\rho) = \sum K_i \rho K_i^*$, since

$$\operatorname{tr} \left[J(\rho)\right] = \operatorname{tr} \left(\sum K_i \rho K_i^*\right) = \sum \operatorname{tr} \left(K_i \rho K_i^*\right) = \sum \operatorname{tr} \left(\rho K_i^* K_i\right)$$
$$= \operatorname{tr} \left(\rho \sum K_i^* K_i\right)$$

for all $\rho \in \mathcal{D}(H)$, we conclude that $\widehat{J} = \sum K_i^* K_i$. In particular, if $J(\rho) = a^{1/2} \rho a^{1/2}$ is a Lüders operation, then $\widehat{J} = a$.

An observable is given by a finite sequence of effects $A = \{A_x : x \in \Omega_A\}$ where $A_x \in \mathcal{E}(H)$ with $\sum A_x = I$ [1–4, 6]. Then Ω_A is the outcome space of A and A determines an effect-valued measure $A(\Delta) = \sum_{x \in \Delta} A_x, \Delta \subseteq \Omega_A$. The probability that A has outcome x in the state $\rho \in \mathcal{D}(H)$ is tr (ρA_x) and the ρ -distribution of A is $\Phi_{\rho}^A(\Delta) = \sum_{x \in \Delta} \operatorname{tr}(\rho A_x)$. The set of observables on H is denoted by $\operatorname{Ob}(H)$. In a similar way, an *instrument* is given by a finite sequence of operations $\mathcal{I} = \{J_x : x \in \Omega_\mathcal{I}\}, J_x \in \mathcal{O}(H)$ where $\sum J_x$ is a channel [1–3, 5, 6]. Then $\Omega_\mathcal{I}$ is the outcome space of \mathcal{I} and \mathcal{I} defines an operation-valued measure $\mathcal{I}(\Delta) = \sum_{x \in \Delta} J_x, \Delta \in \Omega_\mathcal{I}$. The probability that \mathcal{I} has outcome x in the state $\rho \in \mathcal{D}(H)$ is tr $[J_x(\rho)]$ and the ρ -distribution of \mathcal{I} is $\phi_{\rho}^{\mathcal{I}}(\Delta) = \sum_{x \in \Delta} \operatorname{tr}[J_x(\rho)]$. We denote the set of instruments on H by In (H). An instrument \mathcal{I} measures the unique observable \widehat{J}_x . If $A \in \operatorname{Ob}(H)$, the

corresponding Lüders instrument is
$$\mathcal{L}_x^A(\rho) = A_x^{1/2} \rho A_x^{1/2}, x \in \Omega_A$$
. Since

tr
$$\left[\mathcal{L}_x^A(\rho)\right]$$
 = tr $\left(A_x^{1/2}\rho A_x^{1/2}\right)$ = tr $\left(\rho A_x\right)$

for all $\rho \in \mathcal{D}(H)$, we have $(\mathcal{L}_x^A)^{\wedge} = A_x$ so \mathcal{L}^A measures A.

We now consider transition probabilities in the Hilbert space formulation of quantum mechanics. Let $\mathcal{I} \in \mathcal{O}(H)$, $\rho \in \mathcal{D}(H)$, $A, B \in \mathcal{E}(H)$. If \mathcal{I} has Kraus decomposition $\mathcal{I}(\rho) = \sum K_i \rho K_i^*$ with $\sum K_i^* K_i \leq I$, then

$$\operatorname{tr} \left[\mathcal{I}(\rho) \right] = \operatorname{tr} \left(\sum K_i \rho K_i^* \right) = \sum \operatorname{tr} \left(K_i \rho K_i^* \right) = \sum \operatorname{tr} \left(\rho K_i^* K_i \right)$$
$$= \operatorname{tr} \left(\rho \sum K_i^* K_i \right)$$

Assuming that tr $[\mathcal{I}(\rho)] \neq 0$, we have the updated state

$$\mathcal{I}(\rho)^{\sim} = \frac{\mathcal{I}(\rho)}{\operatorname{tr} (\rho \sum K_i^* K_i)} = \frac{\sum K_i \rho K_i^*}{\operatorname{tr} (\rho \sum K_i^* K)}$$

If $\mathcal{I} = \mathcal{L}^C$, $C \in \mathcal{E}(H)$ is a Lüders operation $\mathcal{L}^C(\rho) = C^{1/2}\rho C^{1/2}$, then

$$\mathcal{L}^C(\rho)^{\sim} = \frac{\mathcal{L}^C(\rho)}{\operatorname{tr}\left[\mathcal{L}^C(\rho)\right]} = \frac{C^{1/2}\rho C^{1/2}}{\operatorname{tr}\left(\rho C\right)}$$

For $A, B \in \mathcal{E}(H)$ and general $\mathcal{I} \in \mathcal{O}(H)$, the transition probability of A to B relative to ρ, \mathcal{I} becomes

$$P_{\rho,\mathcal{I}}(A,B) = \operatorname{tr}(\rho A)\operatorname{tr}\left[\mathcal{I}(\rho)^{\sim}B\right] = \frac{\operatorname{tr}(\rho A)}{\operatorname{tr}\left(\rho \sum K_{i}^{*}K_{i}\right)}\operatorname{tr}\left(\sum K_{i}\rho K_{i}^{*}B\right)$$
$$= \frac{\operatorname{tr}(\rho A)}{\operatorname{tr}\left(\rho \sum K_{i}^{*}K_{i}\right)}\operatorname{tr}\left(\rho \sum K_{i}^{*}BK_{i}\right)$$

If $\mathcal{I} = \mathcal{L}^C$ is a Lïders operation, we obtain

$$P_{\rho,\mathcal{L}^C}(A,B) = \frac{\operatorname{tr}(\rho A)}{\operatorname{tr}(C^{1/2}\rho C^{1/2})}\operatorname{tr}(\rho C^{1/2}BC^{1/2}) = \frac{\operatorname{tr}(\rho A)}{\operatorname{tr}(\rho C)}\operatorname{tr}(C^{1/2}\rho C^{1/2}B)$$

When ψ is a unit vector in H, $\rho = |\psi\rangle\langle\psi|$ is a pure state and we have

$$P_{\rho,\mathcal{L}^C}(A,B) = \frac{\operatorname{tr}(|\psi\rangle\langle\psi|A)}{\operatorname{tr}(|\psi\rangle\langle\psi|C)}\operatorname{tr}\left(|\psi\rangle\langle\psi|C^{1/2}BC^{1/2}\right)$$
$$= \frac{\langle A\psi,\psi\rangle}{\langle C\psi,\psi\rangle}\left\langle C^{1/2}BC^{1/2}\psi,\psi\right\rangle$$

Now let $\rho_1, \rho_2 \in \mathcal{D}(H)$ and $\mathcal{I} \in \mathcal{O}(H)$. We then have the transition probability

$$P_{\mathcal{I}}(\rho_1, \rho_2) = \mathcal{I}(\rho_1)(I)\mathcal{I}\left(\mathcal{I}(\rho_2)\right)(I) = \operatorname{tr}\left[\mathcal{I}(\rho_1)\right]\operatorname{tr}\left[\mathcal{I}\left(\mathcal{I}(\rho_2)\right)\right]$$

If \mathcal{I} has the above Kraus decomposition, then

$$P_{\mathcal{I}}(\rho_1, \rho_2) = \operatorname{tr}\left(\sum_i K_i \rho_1 K_i^*\right) \operatorname{tr}\left(\sum_{i,j} K_j K_i \rho_2 K_i^* K_j^*\right)$$
$$= \operatorname{tr}\left(\rho_1 \sum_i K_i^* K_i\right) \operatorname{tr}\left(\rho_2 \sum_{i,j} K_i^* K_j^* K_j K_i\right)$$

In case of a Lüders operation $\mathcal{I} = \mathcal{L}^A$ we obtain

$$P_{\mathcal{L}^A}(\rho_1, \rho_2) = \operatorname{tr}(\rho_1 A) \operatorname{tr}(\rho_2 A^2)$$

When $\rho_1 = |\psi_i\rangle\langle\psi_1|, \ \rho_2 = |\psi_2\rangle\langle\psi_2|$ are pure states, this becomes

$$P_{\mathcal{L}^A}(\rho_1,\rho_2) = \operatorname{tr} \left(|\psi_1\rangle \langle \psi_1 | A \right) \operatorname{tr} \left(|\psi_2\rangle \langle \psi_2 | A^2 \right) = \langle A\psi_1,\psi_1\rangle \langle A^2\psi_2,\psi_2\rangle$$

If A is a projection, we have

$$P_{\mathcal{L}^A}(\rho_1,\rho_2) = \langle A\psi_1,\psi_1\rangle \langle A\psi_2,\psi_2\rangle$$

and when $A = |\psi\rangle\langle\psi|$ is a one-dimensional projection, this becomes

$$P_{\mathcal{L}^{A}}(\rho_{1},\rho_{2}) = \left| \langle \psi,\psi_{1} \rangle \right|^{2} \left| \langle \psi,\psi_{2} \rangle \right|^{2}$$

Finally, if $\psi = \psi_1$ we obtain $P_{\mathcal{L}^A}(\rho_1, \rho_2) = |\langle \psi_1, \psi_2 \rangle|^2$ which is the usual transition probability for pure states. We conclude that the usual transition probability is a very special case.

STAN GUDDER

References

- P. Busch, M. Grabowski and P. Lahti, *Operational Quantum Physics*, Springer-Verlag, Berlin, 1995.
- [2] E. Davies and J. Lewis, An operational approach to quantum probability, Comm. Math. Phys. 17, 239–260 (1970).
- [3] S. Gudder, Quantum instruments and conditioned observables, arXiv:quant-ph 2005.08117 (2020).
- [4] —, Combinations of quantum observables and instruments, J. Math. Phys. A 54 (36): 364002 (2021).
- [5] —, A theory of quantum instruments, Quanta 12, 27–40 (2023).
- [6] T. Heinosaari and M. Ziman, *The Mathematical Language of Quantum Theory*, Cambridge University Press, Cambridge, 2012.
- [7] A. Holevo, Probabilistic and Statistical Aspects of Quantum Theory, North-Holland, Amsterdam, 1982.
- [8] K. Kraus, States, Effects and Operations, Lecture Notes in Physics, 190, Springer, Berlin, 1983.
- [9] G. Lüders, Über due Zustandsänderung durch den Messprozess, Annalen der Physik 443, 322–328 (1951).
- [10] M. Nielson and I. Chuang, Quantum Computation and Quantum Information, Cambridge University Press, Cambridge, 2000.

Department of Mathematics, University of Denver, Denver, Colorado 80208 Email address: sgudder@du.edu