Achieving quantum advantage in a search for a minimal Goldbach partition with driven atoms in tailored potentials

Oleksandr V. Marchukov¹, Andrea Trombettoni^{2,3}, Giuseppe Mussardo³, and Maxim Olshanii⁴

The famous Goldbach conjecture [1] states that any even natural number Ngreater than 2 can be written as the sum of two prime numbers p and p', with p, p'referred to as a Goldbach pair. In this article we present a quantum analogue protocol for detecting - given a even number N – the existence of a *minimal Goldbach* **partition** N = p + p' with $p \equiv p_{\min}(N)$ being the so-called minimal Goldbach prime, i.e. the least possible value for p among all the Goldbach pairs of N. The proposed protocol is effectively a quantum Grover algorithm with a modified final stage. Assuming that an approximate smooth upper bound $\mathcal{N}(N)$ for the number of primes less than or equal to $p_{\min}(N)$ is known, our protocol will identify if the set of $\mathcal{N}(N)$ lowest primes contains the minimal Goldbach prime in approximately $\sqrt{\mathcal{N}(N)}$ steps, against the corresponding classical value $\mathcal{N}(N)$.

In the larger context of a search for violations of Goldbach's conjecture, the quantum advantage provided by our scheme appears to be potentially convenient. E.g., referring to the current state-of-art numerical search for violations of the Goldbach conjecture among all even numbers up to $N_{\rm max} = 4 \times 10^{18}$ [T. O. e Silva, S. Herzog, and S. Pardi, Mathematics of Computation 83, 2033 (2013)], a quantum realization of the search would deliver a quantum advantage factor of $\sqrt{N(N_{\rm max})} \approx 37$

Oleksandr V. Marchukov: ovmarchukov@gmail.com

Andrea Trombettoni: atrombettoni@units.it Giuseppe Mussardo: mussardo@sissa.it Maxim Olshanii: maxim.olchanyi@umb.edu and it will require a Hilbert space spanning $\mathcal{N}(N_{\text{max}}) \approx 1376$ basis states.

1 Introduction

1.1 Statement of the problem

The Goldbach conjecture [1] is a consequential [2, 3, 4] but yet unproven statement in number theory. It states that for every even natural number N, there exists a pair of prime numbers, (p, p') such that N is a sum of these two numbers,

$$p + p' = N .$$

Of course, there can be more than one Goldbach pair for a given N. We will use the convention that $p' \geq p$ and call (p, p') a Goldbach pair.

The Goldbach conjecture has been successfully verified numerically up to a maximal even number $N_{\text{max}} = 4 \times 10^{18}$ [5]. For a given N, the complexity of the computation is presumed to be

(classical complexity)_{given N} =
$$\mathcal{O}(\mathcal{N}(N))$$
,

where $\mathcal{N}(N)$ is a smooth estimate for the upper bound for the number of primes less then or equal to the minimal prime $p_{\min}(N)$ participating in a Goldbach pair [6] (called, from now on, the *min*imal Goldbach prime). Our goal is to reduce the complexity of the problem to

(quantum complexity)_{given N} =
$$\mathcal{O}(\sqrt{\mathcal{N}(N)})$$
.

It has been suggested [7, 8] that

$$p_{\min}(N) \lesssim p_{\min, \sup}(N)$$
 , (1)

with

$$p_{\min, \sup}(N) = C \ln^2(N) \ln(\ln(N)) . \tag{2}$$

¹Technische Universität Darmstadt, Institut für Angewandte Physik, Hochschulstraße 4a, 64289 Darmstadt, Germany

²Department of Physics, University of Trieste, Strada Costiera 11, I-34151 Trieste, Italy

³SISSA and INFN, Sezione di Trieste, Via Bonomea 265, I-34136 Trieste, Italy

⁴Department of Physics, University of Massachusetts Boston, Boston, Massachusetts 02125, USA

A very simple argument to get (2) is to use the Hardy-Littelwood estimate of Godlbach pairs associated to N [9], which is $\sim N \ln^2 N$. Assuming the Godbach pairs are equispaced, one can get an estimate the minimal Goldbach prime p_{\min} obtaining $p_{\min}(N) \sim \ln^2(N)$.

If one studies the minimal Goldbach primes $p_{\min}(N)$ as a function of N one realizes that the are rather well separated maxima. We call such maxima Goldbach champions. More precisely, the Goldbach champion $p_{\text{champ}}(N)$ the maximum of the minimal Goldbach primes in the interval [4, N]: $p_{\text{champ}}(N) = \max_{i=4,6,\dots,N} p_{\min}(i)$.

Given the available numerical data that covers all even numbers up to to N=24488559567 reported [8] (where the Goldbach champions are also listed,) we estimate the prefactor C as

$$C \approx 1.5$$

. For the number of primes up to that N, we get, accordingly:

$$\mathcal{N}(N) \approx 0.75 \, \ln^2(N) \, . \tag{3}$$

Note that according to [8], all minimal Goldbach primes obey $p_{\min}(N) \leq 1.5 \ln^2(N) \ln(\ln(N))$ with no exceptions.

1.2 The state of the art

Our proposal assumes an ability to build a one-body potential with an arbitrarily tailored spectrum. By using holographic traps [10, 11], such potentials recently became an experimental reality [12] with the implementation of the quantum prime potential for the first 15 prime numbers, i.e. of a potential V(x) such $p^2/2m + V(x)$ has as eigenvalues the first 15 prime numbers (apart from an overall ebergy scale). In [12] the potential v(x) giving the first 10 lucky primes.

The possibility of having a one-body potential opens the possibility of implementing schemes for translating number theory problems in quantum physical settings [13, 14, 15, 16, 17, 18, 19, 20, 21, 22], including applications to few-body quantum prime factorization [14, 16, 17, 18, 19, 21, 22]. Also in [23], we present a quantum proposal for testing the Goldbach conjecture, with no aspiration for a quantum advantage.

2 Protocol and its implementation

2.1 A (lightly) modified Grower protocol

The canonical Grover database search protocol [24] allows, in approximately $\frac{\pi}{4}\sqrt{\mathcal{N}}$ steps, to identify a state $|w\rangle$ hidden in a unitary transformation

$$\hat{U}_w = \sum_{n=1}^{\mathcal{N}} \left\{ \begin{array}{cc} -1 & \text{if} & n=w \\ +1 & \text{otherwise} \end{array} \right\} |n\rangle\langle n| \ .$$

The protocol uses a state

$$|s\rangle \equiv \frac{1}{\sqrt{N}} \sum_{n=1}^{N} |n\rangle ,$$

and the second unitary transformation

$$\hat{U}_s = \sum_{m=1}^{\mathcal{N}} \left\{ \begin{array}{cc} -1 & \text{if} & m = s \\ +1 & \text{otherwise} \end{array} \right\} |m\rangle\langle m| \ ,$$

where $\{|m\rangle\}$ is an orthonormalized basis of which $|s\rangle$ is a member. Often, in literature, \hat{U}_s is presented with an opposite sign.

It can be shown that a two operator sequence $\hat{U}_w\hat{U}_s$, applied to the state $|s\rangle$

$$r(\mathcal{N}) \approx \frac{\pi}{4} \sqrt{\mathcal{N}}$$

times, leads to the state $|w\rangle$ sought after, with a probability close to unity:

$$\left(\hat{U}_s\hat{U}_w\right)^{r(\mathcal{N})}|s\rangle \approx |w\rangle$$
. (4)

The sought key $|w\rangle$ be read out using the state measurement in the $\{|n\rangle\} \otimes |g\rangle$ basis.

As we will see below, the problem of identifying a minimal Goldbach prime is slightly different. In (hypothetical) the case of the Goldbach conjecture being violated for some N, there will not be any minimal Goldbach primes whatsoever. Therefore, in a quantum version of the Goldbach test, we are presented with a unitary

$$\hat{U}_X = \text{either } \hat{U}_w \text{ or } \hat{\mathcal{I}} ,$$

with an unknown $|w\rangle$. Here $\hat{\mathcal{I}}$ is a unity transformation. Our goal is to identify which of the two choices we were given. On the other hand, the actual value of w, even if it existed, will not be discoverable.

Our suggestion is to use the protocol (4), with \hat{U}_w replaced by \hat{U}_X and measure the result in the $\{|m\rangle\} \otimes |g\rangle$ basis instead. In that case, we get

$$(\hat{U}_X \hat{U}_s)^{r(\mathcal{N})} |s\rangle \approx \begin{cases} |s\rangle & \text{if } \hat{U}_X = \hat{\mathcal{I}} \\ |m\rangle \neq |s\rangle & \text{if } \hat{U}_X = \hat{U}_w \end{cases} .$$

$$(5)$$

Finding the state $|s\rangle$ would point to either a rare case of an anomalously large minimal Goldbach prime $p > p_{\min, \sup}(N)$ (to be further investigated classically) or to a true violation of the Goldbach

conjecture, yet undetected.

2.2 Hamiltonians

Our scheme involves a single atom with two internal states, $|g\rangle$ and $|e\rangle$ and two different potentials for each of the internal states and two time-dependent perturbations. In the course of the protocol, we alternate between the two pairs of potentials and two different perturbations. These two Hamitonians read:

$$\hat{H}^{(w)} = U_0 \left\{ \sum_{n=n_{\min}}^{n_{\max}} (-p_n) |n\rangle \langle n| \otimes |g\rangle \langle g| + \sum_{n'=n_{\min}}^{n'_{\max}} (p_{n'} - N) |n'\rangle \langle n'| \otimes |e\rangle \langle e| \right\} + \Theta^{(w)}(t) \left\{ \sum_{n=n_{\min}}^{n_{\max}} \sum_{n'=n'_{\min}}^{n'_{\max}} V_{n',n}^{(w)} |n'\rangle \langle n| |e\rangle \langle e| + \text{h.c.} \right\}.$$
(6)

and

$$\hat{H}^{(s)} = U_0 \left\{ \sum_{m=m_{\min}}^{m_{\max}} (-l_m) |m\rangle \langle m| \otimes |g\rangle \langle g| + \sum_{m'=m'_{\min}}^{m'_{\max}} k_{m'} |m'\rangle \langle m'| \otimes |e\rangle \langle e| \right\}$$

$$+ \Theta^{(s)}(t) \left\{ \sum_{m=m_{\min}}^{m_{\max}} \sum_{m'=m_{\min}}^{m'_{\max}} V_{m',m}^{(s)} |m'\rangle \langle m| |e\rangle \langle e| + \text{h.c.} \right\}$$

$$(7)$$

Here, U_0 is a global energy scale, N is the natural even number to be Goldbach partitioned, p_n is a contiguous sequence of prime numbers greater than 2,

$$p_1 = 3$$
; $p_2 = 5$; $p_3 = 7$; $p_4 = 11$; ...

 l_m is an arbitrary sequence of widely spaced natural numbers, with a typical spacing

$$\Delta l \gg 1$$
,

 $k_{m'}$ is a widely spaced subset of negative integers,

$$\Delta k \gg 1$$
,

with

$$k_{m'_{\min}} = -l_{m_{\max}}$$

and no other coincidences between the $\{k_{m'}\}$ and $\{-l_{m}\}$ sets; functions $\Theta^{(s)}(t)$ and $\Theta^{(w)}(t)$ are allowed to have values that are either 0 or 1, the

matrix elements of both perturbations are assumed to be small,

$$V_{n',n}^{(w)} \ll U_0$$

 $V_{m',m}^{(s)} \ll U_0 \Delta l$ (8)

The bounds for the sequences of indices m and n are indicated at the Fig. 1.

The state $|\bar{n}\rangle \otimes |g\rangle$, with

$$p_{\bar{n}} = p_{\min}(N)$$

—if $p_{\min}(N)$ is an element of the set of primes from p_1 through $p_{\mathcal{N}(N)}$ —plays a role of Grover's state $|w\rangle$.

Two further assumptions are harder to realize, and an additional work will be needed to find a proper empirical setting supporting them.

We furthermore assume that the matrix elements of the $\hat{V}^{(w)}$ perturbation are approximately




Figure 1: The major ingredients of the protocol. Energy is expressed in units of U_0 .

equal to each other, in magnitude:

$$|V_{n',n}^{(w)}| \approx \text{const} \equiv V^{(w)}$$
.

Further, we will associate the ground state of the Hamiltonian \hat{H}_s , $|m_{\rm max}\rangle \otimes |g\rangle$, with Grover's state $|s\rangle$. However, for the Grover procedure to work, we must arrange for the potentials that produce a balanced decomposition of the $|m_{\rm max}\rangle$ onto the members of the $\{|n\rangle\}$ basis:

$$|m_{\text{max}}\rangle = \frac{1}{\sqrt{\mathcal{N}(N)}} \sum_{n=1}^{\mathcal{N}(N)} |n\rangle.$$

2.3 Initial state

The system is initialized in the ground state of the Hamiltonian \hat{H}_s , $|m_{\text{max}}\rangle$: this state is associated with Grover's state $|s\rangle$.

2.4 Realizing \hat{U}_w

To realize the unitary \hat{U}_w , we place the atom to the $\hat{H}^{(w)}$ potential, in the internal ground state $|g\rangle$, and apply the corresponding perturbation for a time

$$\tau^{(w)} = \pi \frac{\hbar}{V^{(w)}}$$

(a 2π pulse, for a transition between the minimal Goldbach prime state $|\bar{n}\rangle \otimes |g\rangle$ and the second member of the pair, $|\bar{n}'\langle \otimes |e\rangle$ (so that $p_{\bar{n}}+p_{\bar{n}'}=N$) and then wait for the nearest integer multiple of $\frac{\hbar}{U_0}$. In the end, the prefactor in front of minimal Goldbach prime state $|\bar{n}\rangle \otimes |g\rangle$, in the decomposition in the $\{|n\rangle \otimes |g\rangle\}$ basis will change sign. For a sufficiently small perturbation strength, the residual off-resonant processes will play no role in the process [25, 23].

2.5 Realizing \hat{U}_s

To realize the unitary \hat{U}_s , we can place our atom in the potential of the Hamiltonian $\hat{H}^{(s)}$, in the ground state $|g\rangle$, and apply the perturbation for a time

$$\tau^{(s)} = \pi \frac{\hbar}{V_{m_{\min},m_{\max}}^{(s)}}$$

(hence realizing another 2π pulse) and then wait for the nearest integer multiple of $\frac{\hbar}{U_0}$. The latter step is designed to annul the effect of the phases accumulated in the free propagation. In the end, the prefactor in front of $|m_{\rm max}\rangle\rangle\otimes|g\rangle$, in the decomposition in the $\{|m\rangle\otimes|g\rangle\}$ basis will change sign.

2.6 Readout

In the end, we measure the atom state in the $\{|m\rangle\rangle\otimes|g\rangle\}$ basis. Finding the system in the state $|m_{\max}\rangle\rangle\otimes|g\rangle$ would indicate that either the lowest Goldbach prime, $p_{\bar{n}}$ does not belong to the set of the first $\mathcal{N}(N)$ primes or the Goldbach conjecture is violated altogether. As in the case of the conventional Grover algorithm [24], this result can be achieved in $\mathcal{O}(\sqrt{\mathcal{N}}(N))$ steps leading to a $\mathcal{O}(\sqrt{\mathcal{N}}(N))$ quantum advantage over the classical listing the set of primes of interest.

As the next step, broader sets of primes should be tested, potentially classically.

3 Conclusion and Outlook

In this article, we suggested a quantum polynomial speed improvement to a query whether a given even number N admits a Goldbach partition N = p + p', with p being taken from the first $\mathcal{N}(N)$ primes. The bound $\mathcal{N}(N)$ was chosen in such a way that for the majority of N's, such partition is known to exist. E.g. for a $\mathcal{N}(N)$ given

by (3), all even numbers N up to 2×10^{10} can be Goldbach partitioned with the lowest prime being within the first $\mathcal{N}(N)$ primes. We were also implicitly assuming that the bound $\mathcal{N}(N)$ was low enough to exclude the appearance of another Goldbach prime within the same set.

Our proposal is an analogue version of Grover's algorithm [24] realized with a single one-dimensional two-level atom in a trap. We altered the last step of Goldbach's procedure, i.e. the detection step of the algorithm: in doing so, we replaced the question "which of the first $\mathcal N$ primes is a Goldbach prime?" by "is there a Goldbach prime among the first $\mathcal N$ primes?"

Two weakest parts of the proposal are (a) the assumption of constancy of the transition matrix elements $V_{n',n}^{(w)}$, and (b) the assumption of constancy of the coefficients of decomposition of the ground state of the unperturbed Hamiltonian $\hat{H}^{(s)}$ in the $|g\rangle$ sector $(|m_{\text{max}}\rangle \otimes |g\rangle = |s\rangle)$ over the $\hat{H}^{(w)}$ (again the $|g\rangle$ sector) unperturbed eigenstates.

Tailoring the matrix elements of experimentally realizable, i.e. coordinate-dependent perturbations is addressed in a separate publication [25].

The role of the shape of the decomposition of $|s\rangle$ over the basis also deserves further research. In our case, the requirements to the decomposition of the $|s\rangle$ are much weaker than in the original Grover problem. The final state no longer needs to be aligned with a particular state (i.e. $|w\rangle$) but instead sufficiently misaligned away from a particular state (i.e. $|s\rangle$). The requirement on the relative difference between the coefficients of decomposition of $|s\rangle$ over the $|g\rangle$ eigenstates of the Hamiltonian $\hat{H}^{(w)}$, $|w'\rangle$ is going to be much more forgiving than in the case of the conventional Grover database search: one can show that for a securely turning the system state away from $|s\rangle$ it is sufficient that the variance of the weights $|\langle w'|s\rangle|^2$ is smaller than its mean:

$$\Delta |\langle w'|s\rangle|^2 \ll |\langle w'|s\rangle|^2$$
 .

We expect an appearance of a generous criterion on the relative difference between the coefficients of decomposition that may allow for a broad variety of potentials to be used.

In the larger context of a search for violations of Goldbach's conjecture, the quantum advantage provided by our scheme appears to be very mild. E.g. the current state-of-art numerical (obviously unsuccessful) search among all the even numbers up to $N_{\rm max} = 4 \times 10^{18}$ [5], a quantum realization of the search would deliver a quantum advantage factor of $\sqrt{\mathcal{N}(N_{\rm max})} \approx 37$ and it will require a Hilbert space spanning $\mathcal{N}(N_{\rm max}) \approx 1376$ basis states. In the future, we would like to find a way to quantum-parallelize the so far classical listing of the even numbers N to be tested.

Acknowledgments

OVM acknowledges the support by the DLR German Aerospace Center with funds provided by the Federal Ministry for Economic Affairs and Energy (BMWi) under Grant No. 50WM1957 and No. 50WM2250E. MO was supported by the NSF Grant No. PHY-1912542. The authors would like to thank the Institut Henri Poincaré (UAR 839 CNRS-Sorbonne Université) and the LabEx CARMIN (ANR-10-LABX-59-01) for their support. - and the Institut Henri Poincaré (UAR 839 CNRS-Sorbonne Université) and LabEx CARMIN (ANR-10-LABX-59-01).

A significant portion of this work was produced during the thematic trimester on "Quantum Many-Body Systems Out-of-Equilibrium", at the Institut Henri Poincaré (Paris): AT, GM, and MO are heartfully grateful to the organizers of the trimester, Rosario Fazio, Thierry Giamarchi, Anna Minguzzi, and Patrizia Vignolo, for the opportunity to be part of the program.

References

- [1] Underwood Dudley. "Elementary number theory". W. H. Freeman and Company. San Francisco (1969).
- [2] Chenglian Liu, Chin-Chen Chang, Zhi-Pan Wu, and Shi-Lin Ye. "A study of relationship between rsa public key cryptosystem and goldbach's conjecture properties". International Journal of Network Security 4, 445 (2015).
- [3] M.K. Viswanath1 and M. Ranjith Kumar. "Goldbach conjecture and cryptography". International Journal of Pure and Applied Mathematics 116, 1311 (2017).
- [4] Subhash Kak. "Goldbach partitions and sequences". Resonance November, 1028 (2014).

- [5] Tomás Oliveira e Silva, Siegfried Herzog, and Silvio Pardi. "Empirical verification of the even goldbach conjecture and computation of prime gaps up to 4 · 10¹⁸". Mathematics of Computation 83, 2033 (2013).
- [6] Tomás Oliveira e Silva. private communication.
- [7] A. Granville, J. Van de Lune, and H.J.J Rielle. "Checking the goldbach conjecture on a vector computer". Number Theory and ApplicationsPage 423 (1989).
- [8] Matti K. Sinisalo. "Checking the goldbach conjecture up to $4 \cdot 10^{11}$ ". Proceedings of the twenty-eighth annual ACM symposium on Theory of computingPage 931 (1993).
- [9] G. H. Hardy and J. E. Littlewood. "Some problems of 'Partitio numerorum'; III: On the expression of a number as a sum of primes". Acta Mathematica 44, 1 70 (1923).
- [10] Matthew Pasienski and Brian DeMarco. "A high-accuracy algorithm for designing arbitrary holographic atom traps". Opt. Express 16, 2176–2190 (2008).
- [11] D. Bowman, T. L. Harte, V. Chardonnet, C. De Groot, S. J. Denny, G. Le Goc, M. Anderson, P. Ireland, D. Cassettari, and G. D. Bruce. "High-fidelity phase and amplitude control of phase-only computer generated holograms using conjugate gradient minimisation". Opt. Express 25, 11692– 11700 (2017).
- [12] Donatella Cassettari, Giuseppe Mussardo, and Andrea Trombettoni. "Holographic realization of the prime number quantum potential". PNAS Nexus 2, pgac279 (2022).
- [13] G. Mussardo. "The quantum mechanical potential for the prime numbers" (1997). arXiv:cond-mat/9712010.
- [14] Christoph Weiss, Steffen Page, and Martin Holthaus. "Factorising numbers with a bose-einstein condensate". Physica A: Statistical Mechanics and its Applications 341, 586-606 (2004).
- [15] R. Mack, J. P. Dahl, H. Moya-Cessa, W. T. Strunz, R. Walser, and W. P. Schleich. "Riemann ζ function from wave-packet dynamics". Phys. Rev. A 82, 032119 (2010).
- [16] Dániel Schumayer and David A. W. Hutchinson. "Colloquium: Physics of the riemann

- hypothesis". Rev. Mod. Phys. **83**, 307–330 (2011).
- [17] F Gleisberg, R Mack, K Vogel, and W P Schleich. "Factorization with a logarithmic energy spectrum". New Journal of Physics 15, 023037 (2013).
- [18] F. Gleisberg, M. Volpp, and W.P. Schleich. "Factorization with a logarithmic energy spectrum of a two-dimensional potential". Physics Letters A **379**, 2556–2560 (2015).
- [19] F. Gleisberg, F. Di. Pumpo, G. Wolff, and W P Schleich. "Prime factorization of arbitrary integers with a logarithmic energy spectrum". J. Phys. A 51, 035009 (2018).
- [20] Giuseppe Mussardo, Andrea Trombettoni, and Zhao Zhang. "Prime suspects in a quantum ladder". Phys. Rev. Lett. 125, 240603 (2020).
- [21] F. Gleisberg and W.P. Schleich. "Factorization with a logarithmic energy spectrum of a central potential". Acta Physica Polonica A ISSN 1898-794X 143, S112 (2023).
- [22] Giuseppe Mussardo and Andrea Trombettoni. "Integer factorization by quantum measurements". arXiv:math/2309.10757 (2023).
- [23] Oleksandr V. Marchukov, Benjamin Carruthers, Harry Kendell, Joanna Ruhl, Donatella Cassettari, Carrie A. Weidner, Andrea Trombettoni, Maxim Olshanii, and Giuseppe Mussardo. "Quantum dynamics of atoms in number-theory-inspired potentials". in preparation.
- [24] Lov K. Grover. "A fast quantum mechanical algorithm for database search". Proceedings of the twenty-eighth annual ACM symposium on Theory of computingPage 212 (1996).
- [25] Oleksandr V. Marchukov and Maxim Olshanii. "Resonance cascades and number theory". arXiv:2402.04361 (2024).