

ON THE INDEX OF POWER COMPOSITIONAL POLYNOMIALS

SUMANDEEP KAUR, SURENDER KUMAR, AND LÁSZLÓ REMETE

ABSTRACT. The index of a monic irreducible polynomial $f(x) \in \mathbb{Z}[x]$ having a root θ is the index $[\mathbb{Z}_K : \mathbb{Z}[\theta]]$, where \mathbb{Z}_K is the ring of algebraic integers of the number field $K = \mathbb{Q}(\theta)$. If $[\mathbb{Z}_K : \mathbb{Z}[\theta]] = 1$, then $f(x)$ is monogenic. In this paper, we give necessary and sufficient conditions for a monic irreducible power compositional polynomial $f(x^k)$ belonging to $\mathbb{Z}[x]$, to be monogenic. As an application of our results, for a polynomial $f(x) = x^d + A \cdot h(x) \in \mathbb{Z}[x]$, with $d > 1$, $\deg h(x) < d$ and $|h(0)| = 1$, we prove that for each positive integer k with $\text{rad}(k) \mid \text{rad}(A)$, the power compositional polynomial $f(x^k)$ is monogenic if and only if $f(x)$ is monogenic, provided that $f(x^k)$ is irreducible. At the end of the paper, we give infinite families of polynomials as examples.

1. INTRODUCTION AND STATEMENTS OF THE RESULTS

Let $f(x) \in \mathbb{Z}[x]$ be a monic irreducible polynomial with a root θ , and set $K = \mathbb{Q}(\theta)$. If \mathbb{Z}_K denotes the ring of algebraic integers of K , then the index of $f(x)$ is the index $[\mathbb{Z}_K : \mathbb{Z}[\theta]]$. It is well known that the index of $f(x)$ is always finite. We say that $f(x)$ is monogenic, if the index of $f(x)$ is equal to 1.

Let k be a positive integer and q be a prime divisor of k . Suppose $f(x) \in \mathbb{Z}[x]$ is a monic polynomial. If $f(x^k)$ is irreducible, then the roots α and β of $f(x^q)$ and $f(x^k)$ respectively, give rise to a tower of fields as shown below.

$$\begin{array}{c} \mathbb{Q}(\beta) \\ \uparrow \\ \mathbb{Q}(\alpha) \\ \uparrow \\ \mathbb{Q} \end{array}$$

Note that $\beta^{\frac{k}{q}} = \alpha$. In this paper, our main motive is to discuss the relation between

2010 *Mathematics Subject Classification.* 11R04, 11R09.

Key words and phrases. discriminant, monogeneity of polynomials, power compositional polynomials.

The second author is grateful to the University Grants Commission, New Delhi for providing financial support in the form of Junior Research Fellowship through Ref No.1129/(CSIR-NET JUNE 2019). The research of the third named author was supported in part by the Eötvös Loránd Research Network (ELKH) and by the University of Debrecen Scientific Research Bridging Fund (DETKA).

the indices $[\mathbb{Z}_L : \mathbb{Z}[\beta]]$ and $[\mathbb{Z}_K : \mathbb{Z}[\alpha]]$, where $L = \mathbb{Q}(\beta)$ and $K = \mathbb{Q}(\alpha)$, i.e., the relation between the index of $f(x^q)$ and the index of $f(x^k)$, where $q \mid k$. Moreover, we give necessary and sufficient conditions for these indices to be equal to 1.

It is well known, that if $f(x)$ is the minimal polynomial of an algebraic integer θ over \mathbb{Q} , then the discriminant $D(f)$ of $f(x)$ and the discriminant $\Delta(K)$ of $K = \mathbb{Q}(\theta)$ are related by the formula

$$D(f) = [\mathbb{Z}_K : \mathbb{Z}[\theta]]^2 \cdot \Delta(K). \quad (1.1)$$

Clearly, if $D(f)$ is squarefree, then $f(x)$ is monogenic. But the converse is not always true. Several classes of monogenic polynomials with non squarefree discriminants are known and their properties have been studied (cf. [3] - [4], [6] - [15], [19]). L. Jones in [13] studied the monogeneity of power compositional Shanks polynomials. In [14], L. Jones constructed infinite collections of monic Eisenstein polynomials $f(x) \in \mathbb{Z}[x]$ such that the power compositional polynomials $f(x^{d^n})$ are monogenic for all integers $n \geq 0$ and any integer $d > 1$, where $f(x)$ is Eisenstein with respect to every prime divisor of d . In this paper, we study the monogeneity of a monic irreducible power compositional polynomial; i.e., a monic irreducible polynomial of the form $f(x^k) \in \mathbb{Z}[x]$.

Throughout the paper, the irreducibility of a monic polynomial belonging to $\mathbb{Z}[x]$ is meant over \mathbb{Q} , unless stated otherwise. Note, that for a positive integer k , the irreducibility of a monic polynomial $f(x^k)$ implies the irreducibility of $f(x)$. For a positive integer z , the radical of z , denoted by $\text{rad}(z)$, is defined as the largest squarefree integer dividing z .

Next, we state our main theorems.

Theorem 1.1. Let $f(x)$ be a monic polynomial with integer coefficients. Let $k \geq 2$ be an integer such that $f(x^k)$ is irreducible. Then $f(x^k)$ is monogenic if and only if

- (1) $f(x)$ is monogenic,
- (2) p does not divide the index of $f(x^p)$ for all primes $p \mid k$ and
- (3) $f(0)$ is squarefree.

Remark 1.2. Note, that for a monic polynomial $f(x) \in \mathbb{Z}[x]$, if the irreducible polynomial $f(x^k)$ is monogenic, then by using Lemma 2.6 it is easy to conclude that $f(x^t)$ is also monogenic for each divisor t of k .

In the proof of Theorem 1.1 we will see that if $f(x)$ is monogenic, then only the prime divisors of k and $f(0)$ can divide the index of $f(x^k)$, so we obtain the following corollary to be used in the proof of Theorem 1.8.

Corollary 1.3. Let $f(x)$ be a monic polynomial having integer coefficients. Let k be a positive integer such that $f(x^k)$ is irreducible. Suppose, that

- (1) $f(x)$ is monogenic and
- (2) p does not divide the index of $f(x^p)$ for any prime $p \mid k$.

Then every prime divisor of the index of $f(x^k)$ will divide $f(0)$.

In view of the above corollary, if $|f(0)| = 1$ or the divisors of $f(0)$ do not divide the index of $f(x^k)$, then (1) and (2) together are equivalent to the monogeneity of $f(x^k)$. This happens, for example, in the case of polynomials that are Eisenstein with respect to any prime divisor of $f(0)$, see Lemma 2.12. These polynomials can be given of the form $f(x) = x^d + A \cdot h(x)$, where $|h(0)| = 1$ and A is squarefree. Adding a further condition, the monogeneity of $f(x)$ will be equivalent to the monogeneity of $f(x^k)$.

Corollary 1.4. Let $f(x) \in \mathbb{Z}[x]$ be a polynomial of the form $f(x) = x^d + A \cdot h(x)$, where $d > 1$, $|h(0)| = 1$ and $\deg h(x) < d$. Let k be a positive integer such that $f(x^k)$ is irreducible. If $\text{rad}(k)$ divides $\text{rad}(A)$, then $f(x)$ is monogenic if and only if $f(x^k)$ is monogenic.

Note, that Theorem 1.1 implies that if $f(x^{p^u})$ is monogenic, then $f(x)$ is monogenic and $f(0)$ is squarefree. In the following theorem we prove, that its converse is true with an additional requirement. More precisely, we give sufficient conditions for the monogeneity of a monic irreducible prime power compositional polynomial.

Theorem 1.5. Let u be a positive integer, p a prime number and $f(x) \in \mathbb{Z}[x]$ a monic polynomial. Suppose that $f(x^{p^u})$ is irreducible. Then the following statements hold:

- (1) If $f(x)$ is monogenic and $f(0)$ is squarefree, then the index of $f(x^{p^u})$ is equal to p^s , with some non-negative integer s .
- (2) p does not divide the index of $f(x^{p^u})$ if and only if $\frac{f(x^p) - f(x)^p}{p}$ is coprime to $f(x)$ modulo p .

Remark 1.6. Note, that if $u = 1$ in the above theorem, then the condition (2) of Theorem 1.1 is equivalent to the condition that $\frac{f(x^p) - f(x)^p}{p}$ is coprime to $f(x)$ modulo p , for every prime $p \mid k$.

Combining the Corollary 1.4 and Theorem 1.5 we get the statement below, which also follows from Lemma 3.1 of L. Jones [14].

Corollary 1.7. Let p be a prime number. Suppose, that $f(x) \in \mathbb{Z}[x]$ is a monic and Eisenstein polynomial with respect to p . If $f(0)$ is squarefree, then $f(x)$ is monogenic if and only if $f(x^{p^u})$ is monogenic for any positive integer u .

Taking Remark 1.6 into account, when $f(x) = x^d + A \cdot h(x) \in \mathbb{Z}[x]$, with $|h(0)| = 1$ and $\deg h(x) < d$, we can provide necessary and sufficient conditions for the monogeneity of the irreducible power compositional polynomial $f(x^k)$. As an application of this theorem, we constructed an infinite family of polynomials for a specific choice of $h(x)$, given in the last section.

Theorem 1.8. Let $f(x) \in \mathbb{Z}[x]$ be a polynomial of the form $f(x) = x^d + A \cdot h(x)$, where $d > 1$, $|h(0)| = 1$ and $\deg h(x) < d$. Let k be a positive integer such that $f(x^k)$ is irreducible. Then $f(x^k)$ is monogenic if and only if

- (1) $f(x)$ is monogenic, and
- (2) $\frac{A \cdot h(x^p) + (-A \cdot h(x))^p}{p}$ is coprime to $f(x)$ modulo p , for any prime $p \mid k$.

2. PRELIMINARY RESULTS

In what follows, for a prime p and a polynomial $g(x) \in \mathbb{Z}[x]$, $\bar{g}(x)$ will denote the polynomial obtained by reducing each coefficient of $g(x)$ modulo p . For a subset $S \subset \mathbb{Z}[x]$, let $\langle S \rangle$ be the ideal generated by S . If $S = \{s_1, s_2, \dots, s_k\}$ is finite, then

$$\langle S \rangle = \{s_1 a_1 + s_2 a_2 + \dots + s_k a_k \mid a_i \in \mathbb{Z}[x], 1 \leq i \leq k\}.$$

To investigate the monogeneity of $f(x)$, we will follow the approach of K. Uchida, so we recall Lemma 2.A and Theorem 2.B proved in [20].

Lemma 2.A. An ideal \mathfrak{M} of $\mathbb{Z}[x]$ containing a monic polynomial is maximal if and only if $\mathfrak{M} = \langle p, g(x) \rangle$ for some prime number p and a monic polynomial $g(x)$ belonging to $\mathbb{Z}[x]$ which is irreducible modulo p .

Theorem 2.B. Let $K = \mathbb{Q}(\theta)$ with θ in \mathbb{Z}_K having minimal polynomial $f(x)$ over \mathbb{Q} , then $\mathbb{Z}_K = \mathbb{Z}[\theta]$ if and only if $f(x)$ does not belong to \mathfrak{M}^2 for any maximal ideal \mathfrak{M} of the polynomial ring $\mathbb{Z}[x]$.

In particular, a prime p divides the index $[\mathbb{Z}_K : \mathbb{Z}[\theta]]$ if and only if $f(x)$ belongs to the square of a maximal ideal of the form $\mathfrak{M} = \langle p, g(x) \rangle$, where $g(x) \in \mathbb{Z}[x]$ is a monic polynomial which is irreducible modulo p . We can split the condition $f(x) \in \langle p, g(x) \rangle^2$ into two parts.

Lemma 2.1. Let p be a prime and $g(x) \in \mathbb{Z}[x]$ be a monic polynomial which is irreducible modulo p . Then $\langle p, g(x) \rangle^2 = \langle p^2, g(x) \rangle \cap \langle p, g^2(x) \rangle$.

Proof. Clearly, $\langle p, g(x) \rangle^2 \subseteq \langle p^2, g(x) \rangle \cap \langle p, g^2(x) \rangle$. To obtain the other direction of the containment, suppose that the polynomial $f(x)$ belongs to both $\langle p^2, g(x) \rangle$ and $\langle p, g^2(x) \rangle$, i.e., there exist $a(x), b(x), c(x), d(x)$ belonging to $\mathbb{Z}[x]$, such that

$$p^2 \cdot a(x) + g(x) \cdot b(x) = f(x) = p \cdot c(x) + g^2(x) \cdot d(x).$$

Since $g(x)$ is irreducible modulo p , it is irreducible in $\mathbb{Z}[x]$. So $\langle g(x) \rangle$ is a prime ideal of $\mathbb{Z}[x]$. By the above equation, $p \cdot (c(x) - p \cdot a(x))$ is a multiple of $g(x)$, i.e., it belongs to the prime ideal $\langle g(x) \rangle$. Obviously, p does not belong to $\langle g(x) \rangle$, thus $c(x) - p \cdot a(x) \in \langle g(x) \rangle$ and therefore $c(x) \in \langle p, g(x) \rangle$. This implies that $p \cdot c(x) \in \langle p^2, p \cdot g(x) \rangle$, and from there $f(x) = p \cdot c(x) + g^2(x) \cdot d(x) \in \langle p^2, p \cdot g(x), g^2(x) \rangle$. Hence $f(x) \in \langle p, g(x) \rangle^2$. \square

If $f(x) \in \langle p, g^2(x) \rangle$, then $\bar{g}(x)$ is a multiple factor of $\bar{f}(x)$. On the other hand, $f(x) \in \langle p^2, g(x) \rangle$ implies that the remainder of $f(x)$ divided by $g(x)$ is a multiple of p^2 . Therefore, the results of Uchida [20] say that p divides the index of $f(x)$ if and only if there exists a monic polynomial $g(x) \in \mathbb{Z}[x]$, which is a multiple irreducible factor of $f(x)$ modulo

p , such that the remainder of $f(x)$ divided by $g(x)$ is a multiple of p^2 . This also follows from the results of Ore [17].

Corollary 2.2. Let $f(x)$ and $g(x)$ belonging to $\mathbb{Z}[x]$ be monic polynomials and let p be a prime number such that $g(x)$ is irreducible modulo p . Suppose that $f(x) \in \langle p, g^2(x) \rangle$, then

$$f(x) \in \langle p, g(x) \rangle^2 \iff f(x) \in \langle p^2, g(x) \rangle.$$

Considering a prime divisor p of $f(0)$, it is easy to see that x is a factor of $f(x)$ modulo p and it is a multiple factor of $f(x^\ell)$ modulo p if $\ell > 1$. So, by using the above corollary, we obtain the following result.

Proposition 2.3. Let $f(x) \in \mathbb{Z}[x]$ be a monic polynomial and p be a prime, such that $p^2 \mid f(0)$. Then p divides the index of $f(x^\ell)$ for any positive integer $\ell > 1$.

Proof. Since p^2 divides $f(0)$, we can see that x is a multiple factor of $f(x^\ell)$ modulo p for any integer $\ell > 1$. Therefore $f(x^\ell) \in \langle p, x^2 \rangle$. Moreover, the remainder of $f(x^\ell)$ divided by x is $f(0)$, which is a multiple of p^2 , thus $f(x^\ell) \in \langle p^2, x \rangle$. By Corollary 2.2 and Theorem 2.B, p divides the index of $f(x^\ell)$. \square

In the following theorem, we show that if $\bar{g}(x)$ is a multiple irreducible factor of $\bar{f}(x)$, then the remainder of $f(x)$ divided by $g(x)$ is a multiple of p^2 if and only if the remainder of $f(x^p)$ divided by $g(x)$ is a multiple of p^2 . This observation is important for the proof of our main results and it is of independent interest as well.

Theorem 2.4. Let $f(x)$ and $g(x)$ belonging to $\mathbb{Z}[x]$ be monic polynomials and let p be a prime number such that $g(x)$ is irreducible modulo p . Suppose that $f(x) \in \langle p, g^2(x) \rangle$. Then

$$f(x) \in \langle p, g(x) \rangle^2 \iff f(x^p) \in \langle p, g(x) \rangle^2.$$

Proof. Write $f(x) = g^2(x) \cdot q(x) + p \cdot r(x)$ for some $q(x), r(x)$ belonging to $\mathbb{Z}[x]$. Let $r(x) = g(x) \cdot s(x) + t(x)$, where $\deg t(x) < \deg g(x)$. Then

$$f(x) = g^2(x) \cdot q(x) + p \cdot g(x) \cdot s(x) + p \cdot t(x). \quad (2.1)$$

Since $g^2(x) \in \langle p, g(x) \rangle^2$ and $p \cdot g(x) \in \langle p, g(x) \rangle^2$, we get that

$$f(x) \in \langle p, g(x) \rangle^2 \iff p \cdot t(x) \in \langle p, g(x) \rangle^2. \quad (2.2)$$

Substituting x^p in place of x in (2.1),

$$f(x^p) = g^2(x^p) \cdot q(x^p) + p \cdot g(x^p) \cdot s(x^p) + p \cdot t(x^p). \quad (2.3)$$

Using the fact, that $p \mid (g(x^p) - g(x)^p)$ we get $g(x^p) \in \langle p, g(x) \rangle$. This implies that both $g^2(x^p)$ and $p \cdot g(x^p)$ belong to the ideal $\langle p, g(x) \rangle^2$. Thus, $f(x^p) \in \langle p, g(x) \rangle^2$ if and only if $p \cdot t(x^p) \in \langle p, g(x) \rangle^2$. Since $p \mid (t(x)^p - t(x^p))$ for any polynomial $t(x) \in \mathbb{Z}[x]$, then

$$p \cdot t(x^p) \in \langle p, g(x) \rangle^2 \iff p \cdot t(x)^p \in \langle p, g(x) \rangle^2. \quad (2.4)$$

Thus, from (2.3) and (2.4),

$$f(x^p) \in \langle p, g(x) \rangle^2 \iff p \cdot t(x)^p \in \langle p, g(x) \rangle^2. \quad (2.5)$$

In view of (2.2) and (2.5), to prove the statement it is enough to show that

$$p \cdot t(x) \in \langle p, g(x) \rangle^2 \text{ if and only if } p \cdot t(x)^p \in \langle p, g(x) \rangle^2.$$

Clearly, $p \cdot t(x) \in \langle p, g(x) \rangle^2$ implies that $p \cdot t(x)^p \in \langle p, g(x) \rangle^2$. Assuming that $p \cdot t(x)^p \in \langle p, g(x) \rangle^2$, we can write

$$p \cdot t(x)^p = p^2 \cdot a(x) + p \cdot g(x) \cdot b(x) + g^2(x) \cdot c(x),$$

for some $a(x), b(x), c(x) \in \mathbb{Z}[x]$. Therefore, $g^2(x) \cdot c(x)$ belongs to the prime ideal $\langle p \rangle$ of $\mathbb{Z}[x]$. Since $g(x)$ is monic and irreducible modulo p , then $c(x) \in \langle p \rangle$ and $t(x)^p \in \langle p, g(x) \rangle$. Equivalently, $t(x) \in \langle p, g(x) \rangle$ and therefore $p \cdot t(x) \in \langle p, g(x) \rangle^2$. This completes the proof. \square

Corollary 2.5. Let u be a positive integer. For a monic polynomial $f(x) \in \mathbb{Z}[x]$, a prime p divides the index of $f(x^p)$ if and only if p divides the index of $f(x^{p^u})$, provided that $f(x^{p^u})$ is an irreducible polynomial.

Proof. Let $g(x) \in \mathbb{Z}[x]$ be a monic polynomial such that $\bar{g}(x)$ is an irreducible factor of $f(x)$ modulo p . As $f(x^p) \equiv f(x)^p \pmod{p}$, $g(x)^p$ is a factor of $f(x^p)$ modulo p , i.e., $f(x^p) \in \langle p, g^2(x) \rangle$. By Theorem 2.4,

$$f(x^p) \in \langle p, g(x) \rangle^2 \iff f(x^{p^2}) \in \langle p, g(x) \rangle^2.$$

Applying induction on u , we get

$$f(x^p) \in \langle p, g(x) \rangle^2 \iff f(x^{p^u}) \in \langle p, g(x) \rangle^2.$$

Keeping Theorem 2.B in mind, p divides the index of $f(x^p)$ if and only if p divides the index of $f(x^{p^u})$. \square

Lemma 2.6. Let $f(x) \in \mathbb{Z}[x]$ be a monic polynomial and ℓ a positive integer such that $f(x^\ell)$ is irreducible. If a prime p divides the index of $f(x)$, then p divides the index of $f(x^\ell)$.

Proof. If p divides the index of $f(x)$, then by Theorem 2.B, there exists a monic polynomial $g(x) \in \mathbb{Z}[x]$, such that $g(x)$ is irreducible modulo p and $f(x) \in \langle p, g(x) \rangle^2$. Therefore, $f(x^\ell) \in \langle p, g(x^\ell) \rangle^2$. Let $h(x)$ be an irreducible factor of $g(x^\ell)$ modulo p . Then $g(x^\ell) \in \langle p, h(x) \rangle$, so $\langle p, g(x^\ell) \rangle \subset \langle p, h(x) \rangle$. This implies, that $f(x^\ell)$ belongs to the square of the maximal ideal $\langle p, h(x) \rangle$. Hence, by Theorem 2.B, p divides the index of $f(x^\ell)$. \square

To investigate all of the possible primes that may divide the index of $f(x^\ell)$, we need to find its discriminant. We shall use the following lemma, which describes the properties of the resultant of two polynomials.

Lemma 2.7. Let $f(x) = a_m x^m + a_{m-1} x^{m-1} + \cdots + a_0$ and $g(x) = b_n x^n + b_{n-1} x^{n-1} + \cdots + b_0$ be two polynomials with integer coefficients of degrees m and n , and let $\lambda_1, \dots, \lambda_m$ and μ_1, \dots, μ_n be the roots of $f(x)$ and $g(x)$, respectively. Then

$$\begin{aligned} (1) \quad R(f, g) &= a_m^n b_n^m \prod_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} (\lambda_i - \mu_j), \\ (2) \quad R(f, g) &= a_m^n \prod_{i=1}^m g(\lambda_i) = (-1)^{mn} b_n^m \prod_{j=1}^n f(\mu_j), \\ (3) \quad R(f, gh) &= R(f, g)R(f, h), \text{ for any } h(x) \in \mathbb{Z}[x], \\ (4) \quad R(f, ag) &= a^m R(f, g), \text{ for any } a \in \mathbb{Z}. \end{aligned}$$

Note, that the discriminant of a monic polynomial $f(x) \in \mathbb{Z}[x]$ of degree n and the resultant $R(f, f')$ are related by the following formula.

$$D(f) = (-1)^{\frac{n(n-1)}{2}} R(f, f'), \quad (2.6)$$

where f' is the derivative of f .

Lemma 2.8. Let $f(x) \in \mathbb{Z}[x]$ be a monic polynomial of degree d , and ℓ a positive integer. Then the discriminant of $f(x^\ell)$ is

$$D(f(x^\ell)) = \pm D(f(x))^\ell \cdot \ell^{d\ell} \cdot f(0)^{\ell-1}.$$

Proof. Using (2.6), we get

$$D(f(x^\ell)) = (-1)^{\frac{d\ell(d\ell-1)}{2}} \cdot R(f(x^\ell), \ell x^{\ell-1} f'(x^\ell)).$$

Applying (3) and (4) of Lemma 2.7,

$$D(f(x^\ell)) = (-1)^{\frac{d\ell(d\ell-1)}{2}} \cdot \ell^{d\ell} \cdot R(f(x^\ell), x^{\ell-1}) \cdot R(f(x^\ell), f'(x^\ell)).$$

Using (2) of Lemma 2.7,

$$\begin{aligned} D(f(x^\ell)) &= (-1)^{\frac{d\ell(d\ell-1)}{2} + d\ell + \ell - 1} \cdot \ell^{d\ell} \cdot \prod_{i=1}^{\ell-1} f(0) \cdot R(f(x^\ell), f'(x^\ell)), \\ &= (-1)^{\frac{d\ell(d\ell+1)}{2} + \ell - 1} \cdot \ell^{d\ell} \cdot f(0)^{\ell-1} \cdot R(f(x^\ell), f'(x^\ell)), \\ &= (-1)^{\frac{d\ell(d\ell+1)}{2} + \ell - 1} \cdot \ell^{d\ell} \cdot f(0)^{\ell-1} \cdot \prod_{i=1}^{d\ell} f'(\lambda_i^\ell), \end{aligned}$$

where $f(\lambda_i^\ell) = 0$, for $1 \leq i \leq d\ell$. Note, that if μ_1, \dots, μ_d are roots of $f(x)$, then we can assume that $\lambda_{i+j-1}^\ell = \mu_j$ for $1 \leq i \leq \ell$ and $1 \leq j \leq d$. Therefore, (1) and (2) of Lemma 2.7 imply that

$$D(f(x^\ell)) = (-1)^{\frac{d\ell(d\ell+1)}{2} + \ell - 1} \cdot \ell^{d\ell} \cdot f(0)^{\ell-1} \cdot \left(\prod_{i=1}^d f'(\mu_i) \right)^\ell.$$

Hence, the result follows from (2) of Lemma 2.7. \square

The following lemma is proved in [16, Proposition 4.15], and it suggests that if $f(x)$ is monogenic, then the discriminant of the field generated by a root of $f(x^\ell)$ cannot be too small.

Lemma 2.9. Let K and L be number fields with $K \subset L$. Then

$$\Delta(K)^{[L:K]} \mid \Delta(L).$$

The next proposition is essential for the proof of our main theorems. It completely describes the possible prime divisors of the index of $f(x^\ell)$.

Proposition 2.10. Let $f(x) \in \mathbb{Z}[x]$ be a monic polynomial of degree d , p a prime and ℓ an integer greater than 1. If $p \nmid \ell$ and $f(x^\ell)$ is irreducible, then p divides the index of $f(x^\ell)$ if and only if either

- (1) p divides the index of $f(x)$, or
- (2) $p^2 \mid f(0)$.

Proof. By Lemma 2.6 and Proposition 2.3, both (1) and (2) imply that p divides the index of $f(x^\ell)$. To prove the converse, we assume that p does not divide the index of $f(x)$ and $p^2 \nmid f(0)$. We split the proof into two cases.

Case 1. $p \nmid f(0)$. Equation (1.1) shows, that

$$[\mathbb{Z}_K : \mathbb{Z}[\theta]]^2 = \frac{D(f(x))}{\Delta(K)},$$

where $\Delta(K)$ is the discriminant of the number field K generated by a root θ of $f(x)$. Let L be a number field generated by a root of $f(x^\ell)$. From Lemma 2.8,

$$D(f(x^\ell)) = \pm D(f(x))^\ell \cdot \ell^{d\ell} \cdot f(0)^{\ell-1}.$$

Using Lemma 2.9, we get

$$\frac{D(f(x^\ell))}{\Delta(L)} \mid \frac{D(f(x))^\ell}{\Delta(K)^\ell} \cdot \ell^{d\ell} \cdot f(0)^{\ell-1},$$

where $\Delta(L)$ is the discriminant of the number field L . Since $p \nmid \frac{D(f(x))}{\Delta(K)}$ and $p \nmid \ell \cdot f(0)$, then p does not divide the index of $f(x^\ell)$.

Case 2. $p \mid f(0)$. In view of Theorem 2.B, we have to show that $f(x^\ell)$ does not belong to $\langle p, h(x) \rangle^2$, where $h(x) \in \mathbb{Z}[x]$ is monic and irreducible modulo p . By Lemma 2.1, it is enough to show that $f(x^\ell)$ does not belong to either $\langle p, h^2(x) \rangle$ or $\langle p^2, h(x) \rangle$. If $h(x) = x$, then $f(x^\ell) - f(0) \in \langle x \rangle \subset \langle p^2, x \rangle$. But $f(0) \notin \langle p^2, x \rangle$, so $f(x^\ell) \notin \langle p^2, x \rangle$. First, we show that $f(x^\ell)$ has a multiple irreducible factor modulo p different from x if and only if $f(x)$ has a multiple irreducible factor modulo p different from x .

Let $g(x)$ be a monic irreducible polynomial modulo p , then $\gcd(g(x), g'(x)) = 1$. Since the derivative of $g(x^\ell)$ is $\ell x^{\ell-1} g'(x^\ell)$ and $p \nmid \ell$, $g(x^\ell)$ is separable if and only if $g(x) \neq x$. Thus, $\gcd(g(x^\ell), g'(x^\ell)) = 1$. We can conclude, that whenever a monic irreducible polynomial $g(x) \neq x$ is a single factor of $f(x)$ modulo p , then all irreducible factors of $g(x^\ell)$ are single factors of $f(x^\ell)$ modulo p . Furthermore, if $g_1(x)$ and $g_2(x)$ are distinct irreducible factors of $f(x)$ modulo p , then $\gcd(g_1(x), g_2(x)) = 1$ and so $\gcd(g_1(x^\ell), g_2(x^\ell)) = 1$. Thus, $h(x) \neq x$ is a multiple irreducible factor of $f(x^\ell)$ modulo p if and only if $f(x)$ has a multiple irreducible factor $g(x)$ modulo p such that $\bar{h}(x) \mid \bar{g}(x^\ell)$.

Let $g(x) \in \mathbb{Z}[x]$ be a monic polynomial which is a multiple irreducible factor of $f(x)$ modulo p . Then $f(x) \in \langle p, g^2(x) \rangle$. By Corollary 2.2 and Theorem 2.B, with the hypothesis that p does not divide the index of $f(x)$, we get $f(x) \notin \langle p^2, g(x) \rangle$. Let us write $f(x) = g(x) \cdot c(x) + r(x)$, where $\deg r(x) < \deg g(x)$ and $c(x), r(x) \in \mathbb{Z}[x]$. Clearly, $r(x) \in \langle p \rangle$ and $\bar{g}(x) \mid \bar{c}(x)$. Set $r(x) = p \cdot s(x)$, for some $s(x) \in \mathbb{Z}[x]$. Then, for certain polynomials $a(x), b(x) \in \mathbb{Z}[x]$, we can write $f(x) = g^2(x) \cdot a(x) + p \cdot g(x) \cdot b(x) + p \cdot s(x)$. Consequently,

$$f(x^\ell) = g^2(x^\ell) \cdot a(x^\ell) + p \cdot g(x^\ell) \cdot b(x^\ell) + p \cdot s(x^\ell). \quad (2.7)$$

As $g(x)$ is irreducible modulo p and $\deg s(x) = \deg r(x) < \deg g(x) = \deg \bar{g}(x)$, then $\gcd(\bar{g}(x), \bar{s}(x)) = 1$. Thus, there exists a polynomial $t(x) \in \mathbb{Z}[x]$ such that $s(x) \cdot t(x) \in 1 + \langle p, g(x) \rangle$ and so $s(x^\ell) \cdot t(x^\ell) \in 1 + \langle p, g(x^\ell) \rangle$. Let $h(x) \in \mathbb{Z}[x]$ be a monic polynomial which is an irreducible factor of $g(x^\ell)$ modulo p , then $g(x^\ell) \in \langle p, h(x) \rangle$, and

$$s(x^\ell) \cdot t(x^\ell) \in 1 + \langle p, h(x) \rangle. \quad (2.8)$$

Since $h(x)$ is a multiple factor of $f(x^\ell)$ modulo p , $f(x^\ell) \in \langle p, h^2(x) \rangle$. We have to show that $f(x^\ell) \notin \langle p^2, h(x) \rangle$. Since $g(x^\ell) \in \langle p, h(x) \rangle$, we can see that

$$g^2(x^\ell) \in \langle p, h(x) \rangle^2 \subset \langle p^2, h(x) \rangle$$

and

$$p \cdot g(x^\ell) \in \langle p^2, p \cdot h(x) \rangle \subset \langle p^2, h(x) \rangle.$$

Using Equation (2.7), we get that $f(x^\ell) \in \langle p^2, h(x) \rangle$ if and only if $p \cdot s(x^\ell) \in \langle p^2, h(x) \rangle$. We can write $s(x^\ell) = h(x) \cdot u(x) + v(x)$, where $\deg v(x) < \deg h(x)$ and $u(x), v(x) \in \mathbb{Z}[x]$. Thus, $p \cdot s(x^\ell) \in \langle p^2, h(x) \rangle$ if and only if $p \cdot v(x) \in \langle p^2, h(x) \rangle$. As $\deg v(x) < \deg h(x)$, $p \cdot v(x) \in \langle p^2, h(x) \rangle$ if and only if $v(x) \in \langle p \rangle$. If $v(x)$ is a multiple of p , then $s(x^\ell) \in \langle p, h(x) \rangle$, which is a contradiction to Equation (2.8). Therefore, $f(x^\ell) \notin \langle p^2, h(x) \rangle$. Hence, Theorem 2.B together with Lemma 2.1 implies that p does not divide the index of $f(x^\ell)$. \square

Remark 2.11. The main idea behind the last part of the proof is that if $g(x)$ is irreducible modulo p and $\deg s(x) < \deg g(x)$, then $\bar{g}(x^\ell)$ and $\bar{s}(x^\ell)$ are coprime. But $f(x^\ell)$ belongs to $\langle p^2, h(x) \rangle$ if and only if $\bar{h}(x)$ is a common divisor of $\bar{g}(x^\ell)$ and $\bar{s}(x^\ell)$.

The following lemma proved in [16, Lemma 2.17] will be used in the proof of the Theorem 1.8.

Lemma 2.12. Let α be an algebraic integer, and let $L = \mathbb{Q}(\alpha)$ be the field generated by it. If the minimal polynomial $f(x)$ of α over \mathbb{Q} is an Eisenstein polynomial with respect to the prime p , i.e., it has the form $x^n + a_{n-1}x^{n-1} + \cdots + a_0$, with a_0, \dots, a_{n-1} divisible by p and $p^2 \nmid a_0$, then the index of $f(x)$ is not divisible by p .

3. PROOF OF THE MAIN RESULTS

Proof of Theorem 1.1. First, assume that $f(x^k)$ is monogenic. Then by Lemma 2.6, $f(x)$ and $f(x^p)$ are also monogenic for all $p \mid k$, so (1) and (2) hold. Furthermore, there is no prime p that divides the index of $f(x^k)$, so by Proposition 2.3, $p^2 \nmid f(0)$ for any prime p , i.e., $f(0)$ is squarefree.

Conversely, assume that (1), (2) and (3) hold. By Proposition 2.10, if $p \nmid k$, then (1) and (3) together imply that p does not divide the index of $f(x^k)$. Let $p \mid k$ and write $k = p^u \ell$, where $p \nmid \ell$. Using Corollary 2.5 and (2), we see that p does not divide the index of $f(x^{p^u})$. Now, using (3) and Proposition 2.10, we get that p does not divide the index of $f(x^k)$. Thus, $f(x^k)$ is monogenic. \square

Proof of Corollary 1.3. Let p be a prime divisor of the index of $f(x^k)$. Suppose, that $p \nmid f(0)$. If $p \nmid k$, then by Proposition 2.10, p divides the index of $f(x)$, which contradicts (1). On the other hand, if $p \mid k$, then we can write $k = p^u \ell$, where $p \nmid \ell$. By Proposition 2.10, p divides the index of $f(x^{p^u})$, hence, by Corollary 2.5, p divides the index of $f(x^p)$. This contradicts (2), so p must be a divisor of $f(0)$. \square

Proof of Corollary 1.4. One can easily see that the monogeneity of an irreducible polynomial of the type $f(x) = x^d + A \cdot h(x) \in \mathbb{Z}[x]$, with $d > 1$, and $\deg h(x) < d$, implies that A is squarefree. Let $p \mid A$. Then $f(x) \in \langle p, x^2 \rangle$, so by Corollary 2.2 and Theorem 2.B, the monogeneity of $f(x)$ implies that A is squarefree. In this situation, $f(x)$ is an Eisenstein polynomial with respect to p and hence by Lemma 2.12, p does not divide the index of $f(x)$. Therefore the proof is complete keeping Theorem 1.1 in mind. \square

Proof of Theorem 1.5. Set $\deg f(x) = d$. Using Lemma 2.8, we get

$$D(f(x^{p^u})) = \pm D(f(x))^{p^u} \cdot p^{udp^u} \cdot f(0)^{p^u-1}.$$

(1) In view of Lemma 2.9, Equation (1.1) and the hypothesis that $f(x)$ is monogenic, we conclude that any prime divisor of the index of $f(x^{p^u})$ is either p or a divisor of $f(0)$. If $|f(0)| = 1$, then the index of $f(x^{p^u})$ is equal to p^s , for some non-negative integer s . Let $q \neq p$ be a prime divisor of $f(0)$. Since $f(x)$ is monogenic and $f(0)$ is squarefree, Proposition 2.10 implies that q does not divide the index of $f(x^{p^u})$. Hence, p is the only possible prime divisor of the index of $f(x^{p^u})$.

(2) Using Corollary 2.5, it is enough to prove that p does not divide the index of $f(x^p)$

if and only if $\frac{f(x^p)-f(x)^p}{p}$ is coprime to $f(x)$ modulo p . Let $\bar{f}(x) = \prod_{i=1}^t \bar{g}_i^{e_i}(x)$ be the factorization of $\bar{f}(x)$ into a product of powers of distinct irreducible polynomials $\bar{g}_i(x)$, where each $g_i(x) \in \mathbb{Z}[x]$ monic. Here, $f(x^p) \in \langle p, g_i^2(x) \rangle$ for all $1 \leq i \leq t$. We can write

$$f(x) = \prod_{i=1}^t g_i^{e_i}(x) + p \cdot a(x),$$

for some $a(x) \in \mathbb{Z}[x]$. Then

$$f(x)^p = \left(\prod_{i=1}^t g_i^{e_i}(x) + p \cdot a(x) \right)^p = \left(\prod_{i=1}^t g_i^{e_i}(x) \right)^p + p^2 \cdot t(x),$$

for some $t(x) \in \mathbb{Z}[x]$. Therefore, $f(x)^p \in \langle p^2, g_i(x) \rangle$ for each $1 \leq i \leq t$. Notice, that $f(x^p) \in \langle p^2, g_i(x) \rangle$ if and only if $f(x^p) - f(x)^p \in \langle p^2, g_i(x) \rangle$. Set $f(x^p) - f(x)^p = p \cdot c(x)$, for some $c(x) \in \mathbb{Z}[x]$. Then, $f(x^p) \in \langle p^2, g_i(x) \rangle$ if and only if $p \cdot c(x) \in \langle p^2, g_i(x) \rangle$. Since $g_i(x) \in \mathbb{Z}[x]$ is monic and irreducible modulo p for each i , it is clear that $p \cdot c(x) \in \langle p^2, g_i(x) \rangle$ if and only if $c(x) = g_i(x) \cdot u(x) + p \cdot v(x)$, for certain $u(x), v(x) \in \mathbb{Z}[x]$. Equivalently, $f(x^p) \in \langle p^2, g_i(x) \rangle$ if and only if $\frac{f(x^p)-f(x)^p}{p}$ is divisible by $g_i(x)$ modulo p . Thus, we get that $f(x^p) \notin \langle p^2, g_i(x) \rangle$ for any $1 \leq i \leq t$ if and only if $\frac{f(x^p)-f(x)^p}{p}$ is coprime to $f(x)$ modulo p . Using Corollary 2.2, $f(x^p) \notin \langle p, g_i(x) \rangle^2$ for any $1 \leq i \leq t$ if and only if $\frac{f(x^p)-f(x)^p}{p}$ is coprime to $f(x)$ modulo p . Hence, our result follows from Theorem 2.B. \square

Proof of Theorem 1.8. Let us start by proving that any prime divisor p of k does not divide the index of the irreducible polynomial $f(x^p)$ if and only if (2) holds. Let p be a prime divisor of k , and write

$$\begin{aligned} f(x^p) &= (x^d + A \cdot h(x) - A \cdot h(x))^p + A \cdot h(x^p) \\ &= (f(x) - A \cdot h(x))^p + A \cdot h(x^p) \\ &= f(x)^p + (-1)^p A^p \cdot h(x)^p + p \cdot f(x) \cdot r(x) + A \cdot h(x^p), \end{aligned}$$

where

$$r(x) = \frac{1}{p} \left(\sum_{i=1}^{p-1} \binom{p}{i} f^{i-1}(x) (-A \cdot h(x))^{p-i} \right).$$

This implies, that

$$\frac{f(x^p) - f(x)^p}{p} = \frac{(-A \cdot h(x))^p + p \cdot f(x) \cdot r(x) + A \cdot h(x^p)}{p}.$$

Therefore, $\frac{f(x^p)-f(x)^p}{p}$ is coprime to $f(x)$ modulo p if and only if $\frac{1}{p}[(-A \cdot h(x))^p + A \cdot h(x^p)]$ is coprime to $f(x)$ modulo p . Thus, using Theorem 1.5, p does not divide the index of $f(x^p)$ if and only if $\frac{1}{p}[(-A \cdot h(x))^p + A \cdot h(x^p)]$ is coprime to $f(x)$ modulo p .

Using Lemma 2.6, the monogeneity of $f(x^k)$ implies that both (1) and (2) hold. Conversely, suppose that both (1) and (2) hold. If $|A| = 1$, then the result follows from Corollary 1.3. Let p be a prime dividing A . If $p^2 \mid A$ then Proposition 2.3 shows that p divides the index of $f(x^p)$, a contradiction to (2). Thus, A must be squarefree. Since $|h(0)| = 1$ and A is squarefree, we can see that $f(x^k)$ is an Eisenstein polynomial with respect to p . Therefore, by Lemma 2.12, p does not divide the index of $f(x^k)$. This shows, that the index of $f(x^k)$ and A are coprime. Hence, Corollary 1.3 implies that $f(x^k)$ is monogenic. \square

4. INFINITE EXAMPLES

In this section, we introduce some well-known infinite parametric families of polynomials for which Theorem 1.1 can be applied efficiently. The first example is the family of pure polynomials obtained by choosing $f(x) = x - A$, where A is an integer. In this case, our theorems imply the following well-known result (see [5, Theorem 1.3]).

Proposition 4.1. Let $k > 1$ be an integer and $f(x) = x - A \in \mathbb{Z}[x]$. If $f(x^k)$ is irreducible, then $f(x^k)$ is monogenic if and only if

- (i) $p^2 \nmid A^p - A$ for any prime $p \mid k$ and
- (ii) A is squarefree.

Proof. The polynomial $f(x) = x - A$ is always monogenic, so (1) of Theorem 1.1 is true for any $A \in \mathbb{Z}$. In view of Theorem 1.1, it is enough to prove that (2) of Theorem 1.1 and (i) are equivalent. Here $f(x^p) \equiv (x - A)^p \pmod{p}$. Theorem 2.B together with Corollary 2.2 imply that p divides the index of $f(x^p)$ if and only if the remainder of $f(x^p)$ divided by $x - A$, which is $A^p - A$, is a multiple of p^2 . This completes the proof. \square

The next example is the family of simplest cubic polynomials of the form

$$f(x) = x^3 - mx^2 - (m + 3)x - 1.$$

In some special cases, we can deal with this family by using Theorem 1.1.

Proposition 4.2. Let $m \in \mathbb{Z}$ and $f(x) = x^3 - mx^2 - (m + 3)x - 1$. Let k be a positive integer such that $f(x)$ is reducible modulo every odd prime $p \mid k$. Then $f(x^k)$ is monogenic if and only if

- (i) $f(x)$ is monogenic, and
- (ii) $m \not\equiv \frac{r^{3p} - 3r^p - 1}{r^p(r^p + 1)} \pmod{p^2}$, for any odd prime $p \mid k$ and $r = 1, \dots, p - 2$.

Proof. We note, that $f(x^k)$ is always irreducible in $\mathbb{Z}[x]$ (see [13, Lemma 3.1]). Referring back to Theorem 1.1, we only have to prove that (2) of Theorem 1.1 is equivalent to (ii). Let p be an odd prime. The Galois group of $f(x)$ is the cyclic group of order three, so the

reducibility of $f(x)$ modulo p implies that it splits into three linear factors. Therefore, the multiple irreducible factors of $f(x^p)$ modulo p are of the form $x - r$. Since $f(0) = -1$ and $f(-1) = 1$, then $r \not\equiv 0, -1 \pmod{p}$. By Theorem 2.B and Corollary 2.2, p divides the index of $f(x^p)$ if and only if $f(x^p) \in \langle p^2, x - r \rangle$ for some $r = 1, \dots, p-2$, i.e., if and only if $f(r^p) \equiv 0 \pmod{p^2}$. Since $r \not\equiv 0, -1 \pmod{p}$, we have $f(r^p) = r^{3p} - mr^{2p} - (m+3)r^p - 1 \equiv 0 \pmod{p^2}$ if and only if

$$m \equiv \frac{r^{3p} - 3r^p - 1}{r^p(r^p + 1)} \pmod{p^2}.$$

Finally, it can be easily checked that for $p = 2$, $f(x)$ is always irreducible modulo 2, and $f(x^2)$ is never contained in $\langle 2, f(x) \rangle^2$, so 2 can never divide the index of $f(x^k)$. \square

Remark 4.3. In Proposition 4.2, the condition $f(x)$ is reducible modulo every odd prime $p \mid k$ can not be dropped. For example, if $k = 13$ and $m \equiv 6 \pmod{13}$, then $f(x)$ is irreducible modulo 13, however, if $m = 71$, then both (i) and (ii) are true, but 13 divides the index of $f(x^k)$. For the sake completeness, see [13, Lemma 3.3], in which the author gives a characterization of the monogeneity of $f(x^p)$, where $f(x) = x^3 - mx^2 - (m+3)x - 1$ is irreducible modulo p .

In the last example, we apply Theorem 1.8 to $f(x) = x^d + A(Bx+1)^m$. The monogeneity of $f(x)$ is investigated in [15].

Proposition 4.4. Let A and B be integers and d and m positive integers such that $d > m$. Assume, that $\gcd(d, mB) = 1$. Let $f(x) = x^d + A(Bx+1)^m$ and let k be a positive integer such that $f(x^k)$ is irreducible. Then $f(x^k)$ is monogenic if and only if the following statements hold:

- (1) Both A and $d^d + (-1)^{d+m} B^d m^m (d-m)^{d-m} A$ are squarefree.
- (2) For any prime p dividing k ,

$$\frac{(-A(Bx+1)^m)^p + A(Bx^p+1)^m}{p}$$

is coprime to $f(x)$ modulo p .

Furthermore, if $\text{rad}(k) \mid \text{rad}(A)$, then $f(x^k)$ is monogenic if and only if both A and $d^d + (-1)^{d+m} B^d m^m (d-m)^{d-m} A$ are squarefree.

Proof. If $A \neq \pm 1$ is a squarefree integer, then $f(x^k)$ is an Eisenstein polynomial with respect to a prime divisor of k , thus $f(x^k)$ is irreducible. This implies, that there exist an infinite number of irreducible polynomials of the form $f(x^k) = x^{dk} + A(Bx^k+1)^m$. By [15, Theorem 1.1], $f(x)$ is monogenic if and only if both A and $d^d + (-1)^{d+m} C^d m^m (d-m)^{d-m} A$ are squarefree. Set $h(x) = (Bx+1)^m$, then $h(0) = 1$ and $\deg h(x) < d$, and our result follows from Theorem 1.8. \square

REFERENCES

- [1] H. Cohen, *A Course in Computational Algebraic Number Theory*, Springer-Verlag, Berlin-Heidelberg, 1993.
- [2] R. Dedekind, Über den Zusammenhang zwischen der Theorie der Ideale und der Theorie der höheren Kongruenzen, *Göttingen Abh.* 23 (1878), 1-23.
- [3] I. Gaál, *Diophantine Equations and Power Integral Bases: Theory and algorithms*, Second edition, Springer-Verlag, Berlin, 2019.
- [4] R. Ibarra, H. Lembeck, M. Ozaslan, H. Smith and K. Stange, Monogenic fields arising from trinomials, arXiv:1908.09793v2.
- [5] B. Jhorar and S.K. Khanduja, When is $R[\theta]$ integrally closed?, *J. Algebra Appl.*, 15:5 (2016), 1650091.
- [6] L. Jones, A brief note on some infinite families of monogenic polynomials, *Bull. Aust. Math. Soc.* 100 (2019), 239-244.
- [7] L. Jones, Generating infinite families of monogenic polynomials using a new discriminant formula, *Albanian J. Math.* 14 (2020), 37-45.
- [8] L. Jones, Monogenic polynomials with non squarefree discriminant, *Proc. of Amer. Math. Soc.* 148 (2020), 1527-1533.
- [9] L. Jones, Monogenic reciprocal trinomials and their Galois groups, *J. Algebra Appl.* 21:2 (2022), 2250026.
- [10] L. Jones and D. White, Monogenic trinomials with non squarefree discriminant, *Int. J. Math.* 32:13 (2021), 2150089.
- [11] L. Jones, Some new infinite families of monogenic polynomials with non-squarefree discriminant, *Acta Arith.* 197 (2021), 213-219.
- [12] L. Jones, On necessary and sufficient conditions for the monogeneity of a certain class of polynomials, *Math. Slovaca*, 72:3 (2022), 591-600.
- [13] L. Jones, On the monogeneity of power-compositional Shanks polynomials, <https://doi.org/10.48550/arXiv.2303.11872>.
- [14] L. Jones, The monogeneity of power-compositional Eisenstein polynomials, *Annal. Math. et Inf.*, 55 (2022), 93-113.
- [15] S. Kaur and S. Kumar, On a conjecture of Lenny Jones about certain monogenic polynomials, *Bull Aust Math Soc.* DOI:10.1017/S0004972723001119.
- [16] W. Narkiewicz, *Elementary and analytic theory of algebraic numbers*, 3rd edn., Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2004.
- [17] Ø. Ore, Newtonsche Polygone in der Theorie der algebraischen Körper, *Math. Ann.*, 99 (1928), 84-117.
- [18] D. Shanks, The simplest cubic fields, *Math. Comput.*, 28 (1974), 1137-1152.
- [19] H. Smith, The monogeneity of Kummer extensions and radical extensions, *Acta Arith.* **198** (2020), 313-327.
- [20] K. Uchida, When is $\mathbb{Z}[\theta]$ the ring of integers?, *Osaka J. Math.* 14 (1977), 155-157.

(Sumandeep Kaur) DEPARTMENT OF MATHEMATICS, PANJAB UNIVERSITY, CHANDIGARH

(Surender Kumar) DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY (IIT), BHILAI

(László Remete¹) INSTITUTE OF MATHEMATICS, UNIVERSITY OF DEBRECEN, HUNGARY AND ELKH-
DE EQUATIONS, FUNCTIONS, CURVES AND THEIR APPLICATIONS RESEARCH GROUP

Email address, Sumandeep Kaur: `sumandhunay@gmail.com`

Email address, Surender Kumar: `surenderk@iitbhillai.ac.in`

Email address, László Remete: `remete.laszlo@science.unideb.hu`

¹Corresponding author