CHARACTERISTIC MODULES OVER A LOCAL RING

MOHSEN GHEIBI AND RYO TAKAHASHI

ABSTRACT. Let R be a commutative noetherian local ring, and let M be a finitely generated R-module. Inspired by works of Vasconcelos and Briggs on characterization of complete intersection local rings through the homological properties of the conormal module, in this paper, we define the characteristic module T_M and the cocharacteristic module T_M and investigate their properties. Our main results include characterizations of Cohen-Macaulay and Gorenstein local rings.

1. Introduction

Let (R, \mathfrak{m}) be a (commutative noetherian) local ring. Let $\pi: Q \twoheadrightarrow \widehat{R}$ be a Cohen presentation of R, that is, a surjective homomorphism from a regular local ring Q to the \mathfrak{m} -adic completion \widehat{R} of R. Let I be the kernel of π . Then the conormal module of R is defined as the R-module C_R such that $\widehat{C_R} \cong I/I^2$. The notion of conormal modules has been studied actively for more than half a century in its algebraic and geometric aspects; see [1, 2, 5, 12, 13] and references therein. The celebrated theorem of Ferrand, Vasconcelos, and Briggs states that if $\operatorname{pd}_R C_R < \infty$, then the ring R is complete intersection. One may think of the conormal module as the first Tor module of \widehat{R} over Q. That is

$$\widehat{\mathcal{C}_R} \cong \operatorname{Tor}_1^Q(\widehat{R}, \widehat{R}).$$

A natural question stems from the aforementioned isomorphism. It is whether the ring R can be characterized regarding certain properties of higher Tor modules of \widehat{R} over Q. In this paper, we focus on the last nonzero Tor module of \widehat{R} over Q and define the *characteristic module* of R to be such an R-module T_R that satisfies

$$\widehat{\mathrm{T}_R} \cong \mathrm{Tor}_s^Q(\widehat{R}, \widehat{R}),$$

where $s = \operatorname{pd}_Q \widehat{R}$. More generally, for each (finitely generated) R-module M we shall define the *characteristic module* T_M of M by the existence of an isomorphism $\widehat{T_M} \cong \operatorname{Tor}_s^Q(\widehat{R}, \widehat{M})$. We also consider the dual notion and define the *cocharacteristic module* of M as an R-module E_M such that $\widehat{E_M} \cong \operatorname{Ext}_O^s(\widehat{R}, \widehat{M})$.

First we discuss the existence and uniqueness of (co)characteristic modules and show that if they exist, they do not depend on the choice of a Cohen presentation. More precisely, we prove the following theorem.

Theorem 1.1. Let R be a local ring and M an R-module.

- (1) If R is a homomorphic image of a Gorenstein local ring, then both T_M and E_M exist.
- (2) If T_M or E_M exists, then it is uniquely determined from M up to isomorphism.

As to the structure of (co)characteristic modules and their completions, we present the theorem below.

Theorem 1.2. Let R be a local ring. Then the following statements hold true.

- (1) Suppose that E_R exists. If R is a Cohen–Macaulay ring, then E_R is a canonical module of R.
- (2) Suppose that E_R exists. For every Cohen presentation $Q \to \widehat{R}$, there are isomorphisms

$$E_{\widehat{R}} \cong \widehat{E_R} \cong \operatorname{Ext}_Q^s(\widehat{R}, \widehat{R}) \cong \operatorname{Ext}_Q^s(\widehat{R}, Q), \qquad s := \operatorname{pd}_Q \widehat{R} = \dim Q - \operatorname{depth} R.$$

(3) If E_M exists, then $E_{\widehat{M}} \cong \widehat{E_M} \cong E_{\widehat{R}} \otimes_{\widehat{R}} \widehat{M}$. If T_M exists, then $T_{\widehat{M}} \cong \widehat{T_M} \cong \operatorname{Hom}_{\widehat{R}}(E_{\widehat{R}}, \widehat{M})$.

Cohen-Macaulay local rings are characterized in terms of the dimensions of (co)characteristic modules.

Theorem 1.3. Suppose that R is a homomorphic image of a Gorenstein local ring. The following conditions are equivalent.

 $^{2020\} Mathematics\ Subject\ Classification.\ 13D05,\ 13D07,\ 13H10.$

Takahashi was partly supported by JSPS Grant-in-Aid for Scientific Research 23K03070.

- (1) The local ring R is Cohen–Macaulay.
- (2) One has dim $E_R = \dim R$.
- (3) One has dim $T_R = \dim R$.
- (4) There exists an R-module M such that dim $E_M = \dim R$.
- (5) There exists an R-module M such that dim $T_M = \dim R$.

As an application of Theorem 1.3, we can also relate a (co)characteristic module to the generic Gorensteinness, and obtain a characterization of the Gorenstein local ring.

Theorem 1.4. Let R be a local ring. The following conditions are equivalent.

- (1) The local ring R is Gorenstein.
- (2) One has that E_R exists and is nonzero and free.
- (3) One has that T_R exists and is nonzero and free.
- (4) There is an R-module M with dim $M = \dim R$ such that E_M exists and $E_M \cong M$.
- (5) There is an R-module M with dim $M = \dim R$ and G-dim $M < \infty$ such that T_M exists and $T_M \cong M$.
- (6) There is an R-module M with $G\operatorname{-dim}_R M < \infty$ such that E_M exists, $\operatorname{pd}_R E_M < \infty$ and $\operatorname{dim} E_M = \operatorname{dim} R$.
- (7) There is an R-module M with $\operatorname{pd}_R M < \infty$ such that T_M exists, $\operatorname{G-dim}_R \operatorname{T}_M < \infty$ and $\operatorname{dim} \operatorname{T}_M = \operatorname{dim} R$.

The organization of this paper is as follows. In Section 2, we collect definitions and properties of basic notions used in this paper. In Section 3, we state the precise definitions of a quasi-canonical module of R, a characteristic module and a cocharacteristic module of a given R-module. We investigate their basic properties in this section. In Section 4, we study characteristic and cocharacteristic modules over a quotient of a regular local ring from functorial approaches. In the final Section 5, we give the proofs of the theorems mentioned above and state some other related results.

2. Preliminaries

This section is devoted to preliminaries for the later section. We begin with stating our convention adopted throughout this paper.

Convention 2.1. Throughout this paper, we assume that all rings are commutative and noetherian and all modules are finitely generated. For a local ring (R, \mathfrak{m}) we denote by $\widehat{(-)}$ the \mathfrak{m} -adic completion. A *Cohen presentation* is by definition a surjective ring homomorphism $Q \to \widehat{R}$ where Q is a regular local ring.

We recall the definition of a dualizing complex of R and some of its fundamental properties.

- **Definition 2.2.** (1) Let D be an R-complex with finitely generated homology (note than that D is homologically bounded). We say that D is *dualizing* if it has finite injective dimension and the homothety morphism $R \to \mathbf{R}\mathrm{Hom}_R(D,D)$ is a quasi-isomorphism.
- (2) We denote by D_R a normalized dualizing complex of R, that is, D_R is a dualizing R-complex with $\inf\{i \mid H^i(D_R) \neq 0\} = 0$.
- **Remark 2.3.** (1) A dualizing complex of R is uniquely determined up to shifts in the derived category of R; see [7, Chapter V, Theorem 3.1] and [4, (A.8.3)]. Hence a normalized dualizing complex D_R of R is uniquely determined up to quasi-isomorphism.
- (2) A local ring R admits a dualizing complex if and only if R is a homomorphic image of a Gorenstein local ring; see [9, Corollary 1.4].
- (3) If R is a homomorphic image of a Gorenstein local ring S, then

$$D_R \cong \mathbf{R} \operatorname{Hom}_S(R, S) [\dim S - \dim R]$$

in the derived category of R. This is a consequence of the derived Hom- \otimes adjointness and the fact that S is a dualizing complex of S.

(4) If R is Cohen–Macaulay, then $H^0(D_R)$ is a canonical module of R by (3) and [3, Theorem 3.3.7(b)].

Next we recall some basic numerical invariants for local rings and modules over them.

Definition 2.4. Let (R, \mathfrak{m}, k) be a local ring.

- (1) We denote by cmd R the Cohen–Macaulay defect of R, that is, cmd $R = \dim R \operatorname{depth} R$.
- (2) We denote by type R the type of R, that is, type $R = \dim_k \operatorname{Ext}_R^{\operatorname{depth} R}(k, R)$.

(3) For an R-module M we denote by $\nu_R(M)$ the minimal number of generators of M, that is, $\nu_R(M) = \dim_k M \otimes_R k$.

The Auslander class and the Bass class of a Cohen–Macaulay local ring with a canonical module, and the Poincaré and the Bass series of a complex play an important role in Section 4.

Definition 2.5. Let R be a Cohen–Macaulay local ring with a canonical module ω . The Auslander class $\mathcal{A}(R)$ is defined as the class of R-modules M such that the natural map $M \to \operatorname{Hom}_R(\omega, \omega \otimes_R M)$ is an isomorphism, $\operatorname{Tor}_{>0}^R(\omega, M) = 0$ and $\operatorname{Ext}_R^{>0}(\omega, \omega \otimes_R M) = 0$. The Bass class $\mathcal{B}(R)$ is defined as the class of R-modules M such that the natural map $\omega \otimes_R \operatorname{Hom}_R(\omega, M) \to M$ is an isomorphism, $\operatorname{Ext}_R^{>0}(\omega, M) = 0$ and $\operatorname{Tor}_{>0}^R(\omega, \operatorname{Hom}_R(\omega, M)) = 0$.

Definition 2.6. Let X be an R-complex with finitely generated homology. Then the formal Laurent series

$$P_X(t) = \sum_{n \in \mathbb{Z}} \beta_n^R(X) t^n, \qquad I^X(t) = \sum_{n \in \mathbb{Z}} \mu_R^n(X) t^n$$

are respectively called the *Poincaré series* and the *Bass series* of X, where $\beta_n^R(X) = \dim_k \operatorname{Tor}_n^R(k, X)$ and $\mu_R^n(X) = \dim_k \operatorname{Ext}_R^n(X, k)$ are the *n*th Betti and Bass numbers of X.

3. Basic properties of characteristic modules

In this section, we state basic properties of characteristic modules and cocharacteristic modules. First of all, we define a quasi-canonical module of a local ring.

Definition 3.1. We define a quasi-canonical module of R to be an R-module \mathcal{E}_R such that

$$\widehat{\mathcal{E}_R} \cong \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, Q),$$

where $Q \to \widehat{R}$ is a Cohen presentation.

The following proposition guarantees the existence and uniqueness of a quasi-canonical module of a local ring, and explains that it is a generalization of a canonical module of a Cohen–Macaulay local ring.

Proposition 3.2. (1) Suppose that R is a homomorphic image of a Gorenstein local ring. Then a quasi-canonical module of R exists. Indeed, $H^{\operatorname{cmd} R}(D_R)$ is a quasi-canonical module of R.

- (2) If a quasi-canonical module of R exists, then it is uniquely determined up to isomorphism.
- (3) If R is Cohen-Macaulay, then a quasi-canonical module of R is a canonical module of R.

Proof. (1) As R and \widehat{R} are homomorphic images of Gorenstein local rings, there exist normalized dualizing complex D_R and $D_{\widehat{R}}$, and we easily see that $\widehat{D_R} \cong D_{\widehat{R}}$. Set $H = H^{\operatorname{cmd} R}(D_R)$. Let $Q \twoheadrightarrow \widehat{R}$ be a Cohen presentation. We have

$$\widehat{H} \cong \operatorname{H}^{\operatorname{cmd} R}(\widehat{\mathcal{D}_R}) \cong \operatorname{H}^{\operatorname{cmd} R}(\mathcal{D}_{\widehat{R}}) \cong \operatorname{Ext}_O^{\operatorname{cmd} R + (\dim Q - \dim \widehat{R})}(\widehat{R}, Q) = \operatorname{Ext}_O^{\dim Q - \operatorname{depth} R}(\widehat{R}, Q).$$

Therefore, H is a quasi-canonical module of R.

(2) Let X and Y be quasi-canonical modules of R. Then there exist Cohen presentations $Q \to \widehat{R}$ and $Q' \to \widehat{R}$ such that $\widehat{X} \cong \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, Q)$ and $\widehat{Y} \cong \operatorname{Ext}_{Q'}^{\dim Q' - \operatorname{depth} R}(\widehat{R}, Q')$. Since \widehat{R} is a homomorphic image of the Gorenstein local rings Q and Q', a normalized dualizing complex $D_{\widehat{R}}$ of \widehat{R} exists, and we have

$$\mathbf{R}\mathrm{Hom}_Q(\widehat{R},Q)[\dim Q - \dim \widehat{R}] \cong \mathrm{D}_{\widehat{R}} \cong \mathbf{R}\mathrm{Hom}_{Q'}(\widehat{R},Q')[\dim Q' - \dim \widehat{R}]$$

in the derived category of \widehat{R} . It follows that

$$\begin{split} \widehat{X} &\cong \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, Q) = \operatorname{Ext}_Q^{\operatorname{cmd} \widehat{R} + (\dim Q - \dim \widehat{R})}(\widehat{R}, Q) \cong \operatorname{H}^{\operatorname{cmd} \widehat{R}}(\operatorname{D}_{\widehat{R}}), \\ \widehat{Y} &\cong \operatorname{Ext}_{Q'}^{\dim Q' - \operatorname{depth} R}(\widehat{R}, Q') = \operatorname{Ext}_{Q'}^{\operatorname{cmd} \widehat{R} + (\dim Q' - \dim \widehat{R})}(\widehat{R}, Q') \cong \operatorname{H}^{\operatorname{cmd} \widehat{R}}(\operatorname{D}_{\widehat{R}}). \end{split}$$

We thus obtain $\widehat{X} \cong \widehat{Y}$, which implies $X \cong Y$.

(3) This is an immediate consequence of [3, Theorem 3.3.7(b)].

A quasi-canonical module is compatible with completion.

Corollary 3.3. If \mathcal{E}_R exists, then there is an isomorphism $\widehat{\mathcal{E}_R} \cong \mathcal{E}_{\widehat{R}}$.

Proof. Let $Q \twoheadrightarrow \widehat{R}$ be a Cohen presentation. Then $D_{\widehat{R}} = \mathbf{R} \operatorname{Hom}_Q(\widehat{R}, Q)[\dim Q - \dim \widehat{R}]$, and $\mathcal{E}_{\widehat{R}} = \operatorname{H}^{\operatorname{cmd} \widehat{R}}(D_{\widehat{R}}) = \operatorname{Ext}_Q^{\operatorname{cmd} \widehat{R} + \dim Q - \dim \widehat{R}}(\widehat{R}, Q) = \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, Q) \cong \widehat{\mathcal{E}_R}$ by Proposition 3.2(1).

Now we define a characteristic module and a cocharacteristic module of a given R-module.

Definition 3.4. Let M be an R-module. We define a *characteristic module* of M to be an R-module T_M such that

$$\widehat{\mathcal{T}_M} \cong \operatorname{Tor}_{\dim Q - \operatorname{depth} R}^Q(\widehat{R}, \widehat{M})$$

where $Q woheadrightarrow \widehat{R}$ is a Cohen presentation of R. Dually, we define a cocharacteristic module of M to be an R-module \mathcal{E}_M such that

$$\widehat{\mathcal{E}_M} \cong \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, \widehat{M}).$$

The definition of characteristic modules is motivated by the notion of the conormal module of a local ring.

Remark 3.5. Let R be a local ring. The *conormal module* of R is by definition an R-module C_R such that $\widehat{C_R} \cong I/I^2$, where I is the kernel of a minimal Cohen presentation $Q \twoheadrightarrow \widehat{R}$. Hence

$$\widehat{\mathcal{C}_R} \cong \operatorname{Tor}_1^Q(\widehat{R}, \widehat{R}).$$

One has $\operatorname{pd}_Q \widehat{R} = n - t$, where $n = \dim Q$ and $t = \operatorname{depth} R$. The minimal free resolution of \widehat{R} over Q is of the form $F = (0 \to F_{n-t} \to F_{n-t-1} \to \cdots \to F_0 \to 0)$. There is a complex

$$F \otimes_Q \widehat{R} = (0 \to F_{n-t} \otimes_Q \widehat{R} \to F_{n-t-1} \otimes_Q \widehat{R} \to \cdots \to F_0 \otimes_Q \widehat{R} \to 0).$$

The completion of C_R is the first homology of this complex, while the completion of T_R is the left-end homology of this complex.

Spectral sequence arguments give rise to equalities of types and minimal numbers of generators.

Proposition 3.6. Let (R, \mathfrak{m}, k) be a local ring of depth t, and let M be a finitely generated R-module. Assume T_M and E_M exist. Let $Q \twoheadrightarrow \widehat{R}$ be a Cohen presentation, where Q is a regular local ring of dimension n.

- (1) One has $\nu_R(\mathbf{E}_M) = \operatorname{type}(R)\nu_R(M)$.
- (2) Assume T_M is nonzero and $\operatorname{depth}_{\widehat{R}}\operatorname{Tor}_{n-t-i}^Q(\widehat{R},\widehat{M})\geqslant t-i+1$ for all $1\leqslant i\leqslant n-t$. Then one has $\operatorname{depth}_R T_M = \operatorname{depth} R$ and

$$\operatorname{type}_{R}(T_{M}) = \operatorname{type}(R) \operatorname{type}_{R}(M).$$

Proof. Passing to the completion, we may assume $R = \widehat{R}$. Let F be a free resolution of M over Q, U a free resolution of K over K, K an injective resolution of K over K, and K an injective resolution of K over itself.

(1) The double complex $U \otimes_R \operatorname{Hom}_Q(R, J')$ induces an spectral sequence

$$\mathrm{E}^2_{p,q}=\mathrm{Tor}^R_p(k,\mathrm{Ext}^q_Q(R,M))\Longrightarrow \mathrm{H}_{p-q}(U\otimes_R\mathrm{Hom}_Q(R,J')).$$

Since Q is regular, there exists an isomorphism of complexes $U \otimes_R \operatorname{Hom}_Q(R, J') \cong \operatorname{Hom}_Q(\operatorname{Hom}_R(U, R), J')$ from which we get the following isomorphisms

$$\begin{array}{ll} \operatorname{H}_m(U \otimes_R \operatorname{Hom}_Q(R,J')) & \cong \operatorname{H}_m(\operatorname{Hom}_Q(\operatorname{Hom}_R(U,R),J')) \\ & \cong \operatorname{H}_m(\operatorname{Hom}_Q(\operatorname{Hom}_R(k,J),J')) \\ & \cong \operatorname{H}_m(\operatorname{Hom}_R(k,J) \otimes_k \operatorname{Hom}_Q(k,J')) \\ & \cong \oplus_{i-j=m} \operatorname{Ext}_R^i(k,R) \otimes_k \operatorname{Ext}_Q^j(k,M). \end{array}$$

Since $\operatorname{pd}_Q R = n - t$ and the maps $d_{p,q}^r$ on E^r page are bidegree (-r, -r + 1), we have $\operatorname{E}_{0,n-t}^2 \cong \operatorname{E}_{0,n-t}^\infty$. Therefore $k \otimes_R \operatorname{E}_M \cong \operatorname{Ext}_R^t(k,R) \otimes_k \operatorname{Ext}_Q^n(k,M)$. Since $\operatorname{Ext}_Q^n(k,M) \cong k \otimes_Q M$, we are done.

(2) The double complex $\operatorname{Hom}_R(U, R \otimes_Q F)$ induces an spectral sequence

$$\mathrm{E}_{p,q}^2 \cong \mathrm{Ext}_R^p(k,\mathrm{Tor}_q^Q(R,M)) \Longrightarrow \mathrm{H}^{p-q}(\mathrm{Hom}_R(U,R\otimes_Q F)).$$

Since F is a perfect complex over Q, there is a natural isomorphism of complexes $\operatorname{Hom}_R(U,R)\otimes_Q F\cong \operatorname{Hom}_R(U,R\otimes_Q F)$. Therefore one has

$$\begin{array}{ll} \mathrm{H}^m(\mathrm{Hom}_R(U,R\otimes_Q F)) & \cong \mathrm{H}^m(\mathrm{Hom}_R(U,R)\otimes_Q F) \\ & \cong \mathrm{H}^m(\mathrm{Hom}_R(k,J)\otimes_Q F) \\ & \cong \mathrm{H}^m(\mathrm{Hom}_R(k,J)\otimes_k (k\otimes_Q F)) \\ & \cong \oplus_{i-j=m} \operatorname{Ext}^i_R(k,R)\otimes_k \operatorname{Tor}^Q_i(k,M). \end{array}$$

Since the maps $d_{p,q}^r$ on \mathbf{E}^r page are bidegree (r,r-1), the spectral sequence and the assumptions show that $\operatorname{Ext}_R^j(k,\mathbf{T}_M)=0$ for all $0\leqslant j\leqslant t-1$, and $\operatorname{E}_{t,n-t}^2\cong\operatorname{Ext}_{t,n-t}^{\infty}\cong\operatorname{Ext}_{t,n-t}^{\infty}\cong\operatorname{Ext}_{t,n-t}^{\infty}(k,R)\otimes_k\operatorname{Tor}_{n-t}^{\mathbb{Q}}(k,M)$. Therefore $\operatorname{depth}_R\mathbf{T}_M=t$ and since \mathbb{Q} is regular, we have $\operatorname{Tor}_{n-t}^{\mathbb{Q}}(k,M)\cong\operatorname{Ext}_{\mathbb{Q}}^t(k,M)$. This finishes the proof. \square

The following example shows that the equality of Proposition 3.6(2) may fail if the depth assumption on Tor modules is removed. We used Macaulay2 for computations.

Example 3.7. Let R be the completion of the third Veronese subring of the polynomial ring k[s,t] over a field k, that is, $R = k[s^3, s^2t, st^2, t^3]$. Then R is a Cohen–Macaulay complete local ring of dimension 2. We have $R \cong Q/I$ where Q = k[w, x, y, z] and $I = (x^2 - yw, y^2 - xz, xy - wz)$. One has type(R) = 2 (see [3, Exercise 3.6.21]) but type $_R(T_R) = 3 \neq \text{type}(R)^2 = 4$. Moreover, the module $T_R = \text{Tor}_2^Q(R, R)$ is maximal Cohen–Macaulay and depth $_R$ Tor $_1^Q(R, R) = 1$. Since dim Tor $_1^Q(R, R) = \text{dim } I/I^2 = 2$, the conormal module of R is not Cohen–Macaulay.

One obtains a sufficient condition for an artinian local ring R to be Gorenstein.

Corollary 3.8. Let R be an artinian local ring. If there exists a nonzero R-module M such that $\operatorname{type}_R(M) \geqslant \operatorname{type}_R(T_M)$, then R is Gorenstein. In particular, if $M \cong T_M$, then R is Gorenstein.

The following example shows that Corollary 3.8 is not true if R is not Cohen–Macaulay.

Example 3.9. Let k be a field, Q = k[[x, y]], $I = (x^2, xy)$, and R = Q/I. Then R is a one-dimensional complete local ring and depth R = 0. One has $\operatorname{type}(R) = 1$ and $\operatorname{T}_k \cong k$ but R is not Gorenstein. Also for M = R/(x), one has $\operatorname{pd}_Q M = 1$ and therefore $\operatorname{T}_M = \operatorname{Tor}_2^Q(R, M) = 0$. This shows the assumption $\operatorname{T}_M \neq 0$ is necessary in Proposition 3.6.

4. Functorial approaches to characteristic modules and applications

In this section, we restrict to the case of a quotient of a regular local ring and discuss functorial aspects of characteristic modules and cocharacteristic modules over it. Throughout this section, let Q be an n-dimensional regular local ring, I an ideal of Q, R = Q/I and $t = \operatorname{depth} R$. Then $\operatorname{pd}_Q R = n - t$. Let $F = (0 \to F_{n-t} \xrightarrow{\partial_{n-t}} \cdots \xrightarrow{\partial_1} F_0 \to 0)$ be a minimal free resolution of R over Q. Then $D_R = \mathbf{R}\operatorname{Hom}_Q(R,Q)[n - \dim R]$ and $E := \mathcal{E}_R = \operatorname{H}^{\operatorname{cmd} R}(D_R) = \operatorname{Ext}_Q^{n-t}(R,Q)$ by Proposition 3.2(1).

For an R-module M we define the R-modules $\mathbb{T}(M)$ and $\mathbb{E}(M)$, which play a main role in this section.

Definition 4.1. For an R-module M we define the modules $\mathbb{T}(M)$ and $\mathbb{E}(M)$ as follows.

$$\mathbb{T}(M) = \operatorname{Tor}_{n-t}^Q(R,M), \qquad \mathbb{E}(M) = \operatorname{Ext}_Q^{n-t}(R,M).$$

The modules $\mathbb{T}(M)$ and $\mathbb{E}(M)$ can be described by using the quasi-canonical module E.

Proposition 4.2. There are isomorphisms of covariant functors from mod R to mod R:

$$\mathbb{T}(-) \cong \operatorname{Hom}_R(E, -), \qquad \mathbb{E}(-) \cong E \otimes_R -.$$

In particular, one has an adjoint pair

$$\operatorname{mod} R \xrightarrow{\mathbb{E}} \operatorname{mod} R$$

and an R-module isomorphism $\mathbb{E}(R) \cong E$.

Proof. Let M be an R-module. By definition, the module $\mathbb{T}(M)$ is the kernel of the map $\partial_{n-t} \otimes_Q M$, while the modules E and $\mathbb{E}(M)$ is the cokernels of the maps $\operatorname{Hom}_Q(\partial_{n-t}, Q)$ and $\operatorname{Hom}_Q(\partial_{n-t}, M)$, respectively. There is a commutative diagram

of R-modules, where the vertical maps are natural isomorphisms. We get an isomorphism $\operatorname{Hom}_Q(E,M) \to \mathbb{T}(M)$. As E and M are R-modules, we have $\operatorname{Hom}_Q(E,M) = \operatorname{Hom}_R(E,M)$. Hence $\mathbb{T}(M) \cong \operatorname{Hom}_R(E,M)$. There is also a commutative diagram

$$\operatorname{Hom}_{Q}(F_{n-t-1}, M) \xrightarrow{\operatorname{Hom}_{Q}(\partial_{n-t}, M)} \operatorname{Hom}_{Q}(F_{n-t}, M) \xrightarrow{} \mathbb{E}(M) \xrightarrow{} 0$$

$$\cong \bigwedge \qquad \qquad \cong \bigcap \qquad \cong \bigcap$$

of R-modules, where the vertical maps are natural isomorphisms. We get an isomorphism $E \otimes_Q M \to \mathbb{E}(M)$. As E and M are R-modules, we have $E \otimes_Q M \cong E \otimes_Q R \otimes_R M \cong E \otimes_R M$. Hence $\mathbb{E}(M) \cong E \otimes_R M$.

Next we define the canonical homomorphisms α_M and β_M for each R-module M.

Definition 4.3. Let M be an R-module. Define the maps

$$\alpha_M: M \to \operatorname{Hom}_R(E, E \otimes_R M), \qquad \beta_M: E \otimes_R \operatorname{Hom}_R(E, M) \to M$$

by $\alpha_M(m)(e) = e \otimes m$ and $\beta_M(e \otimes f) = f(e)$ for $e \in E$, $m \in M$ and $f \in \text{Hom}_R(E, M)$. Proposition 4.2 says that we may think

$$\alpha_M: M \to \mathbb{TE}(M), \qquad \beta_M: \mathbb{ET}(M) \to M.$$

The homomorphisms $\alpha_{\mathbb{T}(M)}$ and $\beta_{\mathbb{E}(M)}$ split.

Proposition 4.4. (1) There is an equality $\mathbb{T}(\beta_M) \circ \alpha_{\mathbb{T}(M)} = 1$. Hence $\alpha_{\mathbb{T}(M)} : \mathbb{T}(M) \to \mathbb{TET}(M)$ is a split monomorphism.

(2) There is an equality $\beta_{\mathbb{E}(M)} \circ \mathbb{E}(\alpha_M) = 1$. Hence $\beta_{\mathbb{E}(M)} : \mathbb{ETE}(M) \to \mathbb{E}(M)$ is a split epimorphism.

Proof. Using Proposition 4.2, we have

$$\begin{split} \mathbb{T}(\beta_M) &= \operatorname{Hom}_R(E,\beta_M) : \operatorname{Hom}_R(E,E \otimes_R \operatorname{Hom}_R(E,M)) \to \operatorname{Hom}_R(E,M), \\ \alpha_{\mathbb{T}(M)} &= \alpha_{\operatorname{Hom}_R(E,M)} : \operatorname{Hom}_R(E,M) \to \operatorname{Hom}_R(E,E \otimes_R \operatorname{Hom}_R(E,M)), \\ \beta_{\mathbb{E}(M)} &= \beta_{E \otimes_R M} : E \otimes_R \operatorname{Hom}_R(E,E \otimes_R M) \to E \otimes_R M, \\ \mathbb{E}(\alpha_M) &= E \otimes_R \alpha_M : E \otimes_R M \to E \otimes_R \operatorname{Hom}_R(E,E \otimes_R M). \end{split}$$

Now we easily verify that the assertions hold.

Here we remark when α_M and β_M are isomorphisms.

Remark 4.5. Assume that R is Cohen–Macaulay. In view of [4, (3.4.11)], one has that if $\operatorname{pd}_R M < \infty$, then α_M is an isomorphism, and if $\operatorname{id}_R M < \infty$, then β_M is an isomorphism.

The following corollary provides criteria for detecting infinite injective dimension over a local ring by using minimal numbers of generators of modules.

Corollary 4.6. For a finitely generated R-module M, if $\nu_R(\mathbb{T}(M)) > \nu_R(M)$, then $\mathrm{id}_R M = \infty$.

Proof. Assume that M is nonzero and $\mathrm{id}_R M < \infty$. Then by using Proposition 3.6(1) and Remark 4.5, we have $\nu_R(M) = \mathrm{type}(R)\nu_R(\mathbb{T}(M))$. Therefore $\nu_R(\mathbb{T}(M)) \leqslant \nu_R(M)$.

The proposition below says about the compatibility of a quasi-canonical module with localization.

Proposition 4.7. Let $\mathfrak{p} \in \operatorname{Supp}_R(\mathcal{E}_R)$, and write $\mathfrak{p} = \mathfrak{q}/I$ where \mathfrak{q} is a prime ideal of Q containing I. One then has $\operatorname{ht} \mathfrak{q} - \operatorname{depth} R_{\mathfrak{p}} = n - t$ and there is an isomorphism

$$(\mathcal{E}_R)_{\mathfrak{p}} \cong \mathcal{E}_{R_{\mathfrak{p}}}$$
.

Proof. We have $0 \neq (\mathcal{E}_R)_{\mathfrak{p}} = \operatorname{Ext}_Q^{n-t}(R,Q)_{\mathfrak{p}} \cong \operatorname{Ext}_{Q_{\mathfrak{q}}}^{n-t}(R_{\mathfrak{p}},Q_{\mathfrak{q}})$. Since $\operatorname{pd}_{Q_{\mathfrak{q}}}R_{\mathfrak{p}} \leqslant \operatorname{pd}_QR = n-t$, we have $\operatorname{Ext}_{Q_{\mathfrak{q}}}^{>(n-t)}(R_{\mathfrak{p}},Q_{\mathfrak{q}}) = 0$. As $Q_{\mathfrak{q}}$ is a Gorenstein local ring, it is seen that $\dim Q_{\mathfrak{q}} - \operatorname{depth} R_{\mathfrak{p}} = n-t = \dim Q - \operatorname{depth} R$. Thus $\mathcal{E}_{R_{\mathfrak{p}}} \cong \operatorname{Ext}_{Q_{\mathfrak{q}}}^{n-t}(R_{\mathfrak{p}},Q_{\mathfrak{q}}) \cong (\mathcal{E}_R)_{\mathfrak{p}}$.

Now we state and prove the main result of this section.

Theorem 4.8. The following conditions are equivalent.

- (1) There exists an R-module M such that $\dim \mathbb{T}(M) = \dim R$.
- (2) There exists an R-module M such that $\dim \mathbb{E}(M) = \dim R$.
- (3) One has dim $\mathbb{T}(R) = \dim R$.
- (4) One has dim $\mathbb{E}(R) = \dim R$.
- (5) The local ring R is Cohen–Macaulay.
- (6) The map α_R is an isomorphism.
- (7) The map β_E is an isomorphism.

Proof. First of all, note by Proposition 4.2 that $\mathbb{E}(R) \cong E = \mathcal{E}_R$. The implications $(4) \Rightarrow (2)$ and $(3) \Rightarrow (1)$ are obvious. If R is Cohen–Macaulay, then E is a canonical module of R by Proposition 3.2(3). Hence the implication $(5) \Rightarrow (4)$ holds. Note that the supports of $\operatorname{Hom}_R(E, M)$ and $E \otimes_R M$ are contained in the support of E for each R-module M. Using Proposition 4.2, we see that the implications $(1) \Rightarrow (4) \Leftarrow (2)$ hold.

Assume (4). The equality dim $E = \dim R$ implies that Supp $E \cap \operatorname{Assh} R$ is nonempty. We have

$$\emptyset \neq \operatorname{Supp} E \cap \operatorname{Assh} R \subseteq \operatorname{Supp} E \cap \operatorname{Ass} R = \operatorname{Ass} \operatorname{Hom}_R(E, R) = \operatorname{Ass} \mathbb{T}(R),$$

where the last equality follows from Proposition 4.2. We observe that $\dim \mathbb{T}(R) = \dim R$. Thus (3) follows. There exists $\mathfrak{p} \in \operatorname{Supp} E \cap \operatorname{Assh} R$. Take a prime ideal \mathfrak{q} of Q that contains I and satisfies $\mathfrak{p} = \mathfrak{q}/I$. Proposition 4.7 implies $\operatorname{ht} \mathfrak{q} - \operatorname{depth} R_{\mathfrak{p}} = n - t$. As $\mathfrak{p} \in \operatorname{Assh} R$, we have $\operatorname{depth} R_{\mathfrak{p}} = 0$ and $\dim R/\mathfrak{p} = \dim R$. Hence $\operatorname{depth} R = t = n - \operatorname{ht} \mathfrak{q} = \dim Q/\mathfrak{q} = \dim R/\mathfrak{p} = \dim R$. Therefore, R is Cohen–Macaulay. Thus (5) follows. We have got the implications (3) \Leftarrow (4) \Rightarrow (5), and now the conditions (1)–(5) are equivalent.

The implications $(7) \Leftarrow (5) \Rightarrow (6)$ follow from the fact that if R is Cohen–Macaulay, then E is a canonical module of R. If (6) holds, then $R \cong \mathbb{TE}(R)$ and therefore $\dim_R \mathbb{TE}(R) = \dim R$. By setting $M = \mathbb{E}(R)$, we have condition (2) holds. In a similar way, we can show that (7) implies (1).

Remark 4.9. The implication $(4) \Rightarrow (5)$ in the above theorem can also be deduced by using [3, Theorem 8.1.1(b)]. Indeed, we have dim $E = \dim R / \operatorname{Ann}_R \operatorname{Ext}_Q^{n-t}(R,Q) \leqslant t = \operatorname{depth} R$. Hence, if dim $E = \dim R$, then R is Cohen–Macaulay.

Applying the above theorem, we get the following two corollaries, which give criteria for Gorensteinness in terms of characteristic and cocharacteristic modules.

Corollary 4.10. The following conditions are equivalent.

- (1) There exists an R-module M such that dim $M = \dim R$, G-dim $M < \infty$ and $M \cong \mathbb{T}(M)$.
- (2) There exists an R-module M such that dim $M = \dim R$ and $M \cong \mathbb{E}(M)$.
- (3) The local ring R is Gorenstein.

Proof. (2) \Rightarrow (3): Theorem 4.8 implies that R is Cohen–Macaulay, while type R=1 by Proposition 3.6(1).

- $(3) \Rightarrow (1)$ and (2): As R is Gorenstein, Proposition 3.2(3) implies $E \cong R$. We obtain $\mathbb{E}(R) \cong E \cong R$ and $\mathbb{T}(R) \cong \operatorname{Hom}_R(R,R) \cong R$ by Proposition 4.2. Thus, letting M = R, we are done.
- $(1)\Rightarrow (3)$: Theorem 4.8 and Proposition 3.2(3) imply that R is Cohen–Macaulay and E is a canonical module of R. By Proposition 4.2 and [4, (3.1.11)] we have $\operatorname{Hom}_R(E, M) \cong \mathbb{T}(M) \cong M \in \mathcal{A}(R)$. We get $M \in \mathcal{B}(R)$ by [11, Theorem 2.8(a)], which implies that $\operatorname{Ext}_R^{>0}(E, M) = 0$. It follows that $M \cong \mathbf{R}\operatorname{Hom}_R(E, M)$ in the derived category of R. Using [4, (A.7.7)], we get $\operatorname{I}^M(t) = \operatorname{I}^{\mathbf{R}\operatorname{Hom}_R(E, M)}(t) = \operatorname{P}_E(t)\operatorname{I}^M(t)$. Comparing the coefficients of $t^{\operatorname{depth} M}$ in $\operatorname{I}^M(t)$ and $\operatorname{P}_E(t)\operatorname{I}^M(t)$, we observe that E is cyclic. Thus the ring R is Gorenstein. \square

Corollary 4.11. The following conditions are equivalent.

- (1) There exists an R-module M such that $\operatorname{pd}_R M < \infty$, $\operatorname{G-dim}_R \mathbb{T}(M) < \infty$ and $\operatorname{dim} \mathbb{T}(M) = \operatorname{dim} R$.
- (2) There exists an R-module M with G-dim_R $M < \infty$ such that $\operatorname{pd}_R \mathbb{E}(M) < \infty$ and $\dim \mathbb{E}(M) = \dim R$.
- (3) The R-module $\mathbb{T}(R)$ is nonzero and free.
- (4) The R-module $\mathbb{E}(R)$ is nonzero and free.
- (5) The ring R is Gorenstein.

Proof. It follows from Theorem 4.8 and Propositions 3.2(3), 4.2 that under any of those five conditions which are given in the corollary, R is a Cohen–Macaulay local ring and $E \cong \mathbb{E}(R)$ is a canonical module of R.

- $(5) \Rightarrow (3)$ and (4): By the argument in the proof of the implication $(3) \Rightarrow (1)$ and (2) in Corollary 4.10, we have $\mathbb{T}(R) \cong \mathbb{E}(R) \cong R$. Hence the R-modules $\mathbb{T}(R)$ and $\mathbb{E}(R)$ are nonzero and free.
 - $(3) \Rightarrow (1)$ and $(4) \Rightarrow (2)$: Letting M = R, we are done.

- $(1) \Rightarrow (5)$: As $\mathbb{T}(M)$ has finite G-dimension, it is in $\mathcal{A}(R)$ by [4, (3.1.11)]. Since $\mathbb{T}(M) \cong \operatorname{Hom}_R(E, M)$ by Proposition 4.2, we have $M \in \mathcal{B}(R)$ by [11, Theorem 2.8(a)]. Since M has finite projective dimension, it follows from [4, (3.4.12)] that R is Gorenstein.
- (2) \Rightarrow (5): As M has finite G-dimension, it is in $\mathcal{A}(R)$ by [4, (3.1.11)]. Hence $\mathbb{E}(M) \cong E \otimes M \in \mathcal{B}(R)$ by Proposition 4.2 and [11, Theorem 2.8(b)]. Since $\mathbb{E}(M)$ has finite projective dimension, it follows from [4, (3.4.12)] that R is Gorenstein.

To state our next result, we recall a basic fact on faithful modules.

Remark 4.12. An R-module X is faithful if and only if the homothety map $R \to \operatorname{Hom}_R(X,X)$ is injective. Indeed, the kernel of this map coincides with the annihilator of X.

The following lemma is necessary to get our next corollary.

Lemma 4.13. Let M be an R-module such that $M \cong \mathbb{T}(M)$.

- (1) If M is faithful, then so is E.
- (2) If E is faithful, then $E_{\mathfrak{p}} \cong R_{\mathfrak{p}}$ for every $\mathfrak{p} \in \operatorname{Ass}_R M$.
- *Proof.* (1) Proposition 4.2 yields $\mathbb{T}(M) \cong \operatorname{Hom}_R(E, M)$. We get $\operatorname{Ann}_R E \subseteq \operatorname{Ann}_R \mathbb{T}(M) \subseteq \operatorname{Ann}_R M$, where the latter inclusion follows from the assumption that $M \cong \mathbb{T}(M)$. Thus, the assertion holds.
- (2) Since $M \cong \mathbb{T}(M) \cong \operatorname{Hom}_R(E, M)$, we have $\mathfrak{p} \in \operatorname{Ass} M \subseteq \operatorname{Supp} M \subseteq \operatorname{Supp} E$. In view of Propositions 4.2 and 4.7, there are isomorphisms

$$M_{\mathfrak{p}} \cong \mathbb{T}_{R}(M)_{\mathfrak{p}} \cong \operatorname{Hom}_{R}(E, M)_{\mathfrak{p}} \cong \operatorname{Hom}_{R_{\mathfrak{p}}}(E_{\mathfrak{p}}, M_{\mathfrak{p}}) \cong \operatorname{Hom}_{R_{\mathfrak{p}}}(\mathcal{E}_{R_{\mathfrak{p}}}, M_{\mathfrak{p}}) \cong \mathbb{T}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}),$$

 $\operatorname{Hom}_{R_{\mathfrak{p}}}(\kappa(\mathfrak{p}), M_{\mathfrak{p}}) \cong \operatorname{Hom}_{R_{\mathfrak{p}}}(\kappa(\mathfrak{p}), \operatorname{Hom}_{R_{\mathfrak{p}}}(E_{\mathfrak{p}}, M_{\mathfrak{p}})) \cong \operatorname{Hom}_{R_{\mathfrak{p}}}(E_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} \kappa(\mathfrak{p}), M_{\mathfrak{p}}).$

Since \mathfrak{p} is in Ass M, the $\kappa(\mathfrak{p})$ -vector space $\operatorname{Hom}_{R_{\mathfrak{p}}}(\kappa(\mathfrak{p}), M_{\mathfrak{p}})$ is nonzero. Substituting $\kappa(\mathfrak{p})^{\oplus c}$ for $E_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} \kappa(\mathfrak{p})$, we see that c = 1, so that $E_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} \kappa(\mathfrak{p}) \cong \kappa(\mathfrak{p})$. Nakayama's lemma implies that $E_{\mathfrak{p}}$ is cyclic over $R_{\mathfrak{p}}$. Faithfulness localizes by Remark 4.12, and $E_{\mathfrak{p}}$ is a faithful $R_{\mathfrak{p}}$ -module. It follows that $E_{\mathfrak{p}} \cong R_{\mathfrak{p}}$.

Now we can prove another corollary of Theorem 4.8.

Corollary 4.14. Let M be an R-module such that $M \cong \mathbb{T}(M)$. Consider the following two conditions.

- (i) The R-module M is faithful.
- (ii) The R-module E is faithful and dim $M = \dim R$.

Then the following statements hold true.

- (1) If (i) holds, then (ii) holds as well.
- (2) If (ii) holds, then R is a Cohen-Macaulay local ring and E is a canonical module of R.
- (3) If (ii) holds, then the ring R is locally Gorenstein on Ass M.
- (4) If (i) holds, then the ring R is generically Gorenstein.

Proof. The equality Ann M=0 particularly says dim $M=\dim R$. Lemma 4.13(1) implies that E is faithful as well.

- (2) Since dim $M = \dim R$, there exists $\mathfrak{p} \in \operatorname{Assh} R \cap \operatorname{Supp} M$. Then $\mathfrak{p} \in \operatorname{Min} M \subseteq \operatorname{Ass} M$. Lemma 4.13(2) implies $E_{\mathfrak{p}} \cong R_{\mathfrak{p}}$, and hence dim $E = \dim R$. As $E \cong \mathbb{E}(R)$ by Proposition 4.2, it follows from Theorem 4.8 and Proposition 3.2(3) that R is Cohen–Macaulay and E is a canonical module of R.
- (3) Let $\mathfrak{p} \in \operatorname{Ass} M$. By (2), the local ring $R_{\mathfrak{p}}$ is Cohen–Macaulay, and $E_{\mathfrak{p}}$ is a canonical module of $R_{\mathfrak{p}}$. It follows from Lemma 4.13(2) that $E_{\mathfrak{p}}$ is isomorphic to $R_{\mathfrak{p}}$, which means that $R_{\mathfrak{p}}$ is Gorenstein.
- (4) Since the R-module M is faithful, R is embedded in a finite direct sum of copies of M. This implies that Ass R is contained in Ass M. The assertion is now a direct consequence of (3) and (1).

5. Proofs of our main results

This section is devoted to giving proofs of the results stated in the Introduction and stating related results.

Proof of Theorem 1.1. (1) A quasi-canonical module \mathcal{E}_R of R exists by Proposition 3.2(1). Set $X = \operatorname{Hom}_R(\mathcal{E}_R, M)$ and $Y = \mathcal{E}_R \otimes_R M$. Let $Q \twoheadrightarrow \widehat{R}$ be a Cohen presentation. Applying Proposition 4.2 and Corollary 3.3, we have

$$\widehat{X} \cong \operatorname{Hom}_{\widehat{R}}(\widehat{\mathcal{E}_R}, \widehat{M}) \cong \operatorname{Hom}_{\widehat{R}}(\mathcal{E}_{\widehat{R}}, \widehat{M}) \cong \mathbb{T}_{\widehat{R}}(\widehat{M}) = \operatorname{Tor}_{\dim Q - \operatorname{depth} R}^Q(\widehat{R}, \widehat{M}),$$

and similarly $\widehat{Y} \cong \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, \widehat{M})$. Therefore, X is a characteristic module of M, and Y is a cocharacteristic module of M.

(2) Suppose that T_M exists. Then there exists a Cohen presentation $Q \to \widehat{R}$ such that $\widehat{T_M} \cong \operatorname{Tor}_{\dim Q-\operatorname{depth} R}^Q(\widehat{R},\widehat{M})$. Using Proposition 4.2, we have $\widehat{T_M} \cong \mathbb{T}_{\widehat{R}}(\widehat{M}) \cong \operatorname{Hom}_{\widehat{R}}(\mathcal{E}_{\widehat{R}},\widehat{M})$. Hence T_M is uniquely determined from M up to isomorphism. Similarly, E_M is uniquely determined from M up to isomorphism, if it exists.

Proof of Theorem 1.2. Using Propositions 3.2, 4.2 and Theorem 1.1, we see that $\mathcal{E}_{\widehat{R}}$ and $\mathcal{E}_{\widehat{R}}$ exist and $\mathcal{E}_{\widehat{R}} \cong \mathcal{E}_{\widehat{R}}$.

We prove (1) and (2) together. Suppose that E_R exists. By definition, there exists a Cohen presentation $P \to \widehat{R}$ such that $\widehat{E_R} \cong \operatorname{Ext}_P^{\dim P - \operatorname{depth} R}(\widehat{R}, \widehat{R})$. Using Propositions 3.2(2), 4.2 and Theorem 1.1(2), we get

$$\operatorname{Ext}_P^{\dim P-\operatorname{depth} R}(\widehat{R},P) \cong \mathcal{E}_{\widehat{R}} \cong \mathbb{E}_{\widehat{R}}(\widehat{R}) \cong \operatorname{Ext}_P^{\dim P-\operatorname{depth} R}(\widehat{R},\widehat{R}) \cong \operatorname{E}_{\widehat{R}}.$$

Hence E_R is a quasi-canonical module of R, and we have $\mathcal{E}_R \cong E_R$ by Proposition 3.2(2). If R is Cohen–Macaulay, then E_R is a canonical module of R by Proposition 3.2(3).

Let $Q \twoheadrightarrow \widehat{R}$ be a Cohen presentation. By Propositions 3.2(2) and 4.2, we get $\mathcal{E}_{\widehat{R}} \cong \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, Q) \cong \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, \widehat{R})$. Hence $\operatorname{E}_{\widehat{R}} \cong \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, \widehat{R}) \cong \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, Q)$.

(3) Suppose that T_M exists. Then $\widehat{T_M} \cong \operatorname{Tor}_{\dim Q-\operatorname{depth} R}^Q(\widehat{R}, \widehat{M})$ for some Cohen presentation $Q \twoheadrightarrow \widehat{R}$. Theorem 1.1 implies that $T_{\widehat{M}}$ exists and is isomorphic to $\operatorname{Tor}_{\dim Q-\operatorname{depth} R}^Q(\widehat{R}, \widehat{M})$. Hence $\widehat{T_M} \cong T_{\widehat{M}}$. Proposition 4.2 shows that $T_{\widehat{M}} \cong \operatorname{Hom}_{\widehat{R}}(\mathcal{E}_{\widehat{R}}, \widehat{M}) \cong \operatorname{Hom}_{\widehat{R}}(E_{\widehat{R}}, \widehat{M})$. In a similar way, we see that if E_M exists, then $E_{\widehat{M}} \cong \widehat{E_M} \cong E_{\widehat{R}} \otimes_{\widehat{R}} \widehat{M}$.

Proof of Theorem 1.3. Passing to the completion, we may assume that R is complete. Then the assertion follows from Theorem 4.8.

Proof of Theorem 1.4. By completion, we may assume that R is complete, and then we are done by Corollaries 4.10, 4.11.

From now on, we shall state several other results than the ones given in the Introduction. We begin with showing that the cocharacteristic module of a local ring R and the quasi-canonical module of R are the same.

Corollary 5.1. Let R be a local ring. Suppose that either \mathcal{E}_R or E_R exists. Then the other exists as well, and both are isomorphic: one has $\mathcal{E}_R \cong E_R$.

Proof. We have already seen in the proof of Theorem 1.2 that if E_R exists, then \mathcal{E}_R also exists and $E_R \cong \mathcal{E}_R$. Suppose that \mathcal{E}_R exists. Then $\widehat{\mathcal{E}_R} \cong \mathcal{E}_{\widehat{R}} \cong \mathbb{E}_{\widehat{R}}(\widehat{R}) = \operatorname{Ext}_Q^{\dim Q - \operatorname{depth} R}(\widehat{R}, \widehat{R})$ for some Cohen presentation $Q \twoheadrightarrow \widehat{R}$, where the two isomorphisms follow from Corollary 3.3 and Proposition 4.2, respectively. By the definition of a cocharacteristic module, we see that \mathcal{E}_R is a cocharacteristic module of R. It follows from Theorem 1.1(2) that \mathcal{E}_R is isomorphic to E_R .

Finally, we remove the assumption from Corollary 4.14(3)(4) that R is a homomorphic image of a regular local ring. We do this after stating a remark.

Remark 5.2. Let R be a local ring. For an R-module M it holds that

$$\operatorname{Ass}_R M = \{ P \cap R \mid P \in \operatorname{Ass}_{\widehat{R}} \widehat{M} \}.$$

Indeed, let $\mathfrak{p} \in \operatorname{Ass}_R M$. Then $\operatorname{Ass}_{\widehat{R}} \widehat{M} = \bigcup_{\mathfrak{q} \in \operatorname{Ass}_R M} \operatorname{Ass}_{\widehat{R}} \widehat{R}/\mathfrak{q}\widehat{R}$ by [10, Theorem 12(ii)], which contains $\operatorname{Ass}_{\widehat{R}} \widehat{R}/\mathfrak{p}\widehat{R}$. Since $\widehat{R}/\mathfrak{p}\widehat{R}$ is nonzero, there exists $P \in \operatorname{Ass}_{\widehat{R}} \widehat{R}/\mathfrak{p}\widehat{R} \subseteq \operatorname{Ass}_{\widehat{R}} \widehat{M}$. Then $P \cap R \in \{P' \cap R \mid P' \in \operatorname{Ass}_{\widehat{R}} \widehat{R}/\mathfrak{p}\widehat{R}\} = \{\mathfrak{p}\}$ by [10, Theorem 12(i)], and hence $P \cap R = \mathfrak{p}$.

Conversely, let $P \in \operatorname{Ass}_{\widehat{R}} \widehat{M}$ and put $\mathfrak{p} = P \cap R$. Then P is in $\bigcup_{\mathfrak{q} \in \operatorname{Ass}_R M} \operatorname{Ass}_{\widehat{R}} \widehat{R}/\mathfrak{q}\widehat{R}$, so that $P \in \operatorname{Ass}_{\widehat{R}} \widehat{R}/\mathfrak{q}\widehat{R}$ for some $\mathfrak{q} \in \operatorname{Ass}_R M$. It follows that $\mathfrak{p} = P \cap R \in \{P' \cap R \mid P' \in \operatorname{Ass}_{\widehat{R}} \widehat{R}/\mathfrak{q}\widehat{R}\} = \{\mathfrak{q}\}$, which implies that $\mathfrak{p} = \mathfrak{q} \in \operatorname{Ass}_R M$.

Corollary 5.3. Let R be a local ring and M an R-module. Suppose that E_R and T_M exist and that $T_M \cong M$.

- (1) If E_R is faithful and dim $M = \dim R$, then R is locally Gorenstein on $\operatorname{Ass}_R M$.
- (2) If M is faithful, then R is generically Gorenstein.

Proof. There are isomorphisms $\widehat{M} \cong \widehat{\mathrm{T}_M} \cong \mathbb{T}_{\widehat{R}}(\widehat{M})$.

- (1) By Remark 4.12, the \widehat{R} -module $\mathbb{E}_{\widehat{R}}(\widehat{R}) = \widehat{E}_R$ is faithful. Also, $\dim \widehat{M} = \dim \widehat{R}$. If \widehat{R} is locally Gorenstein on $\operatorname{Ass}_{\widehat{R}} \widehat{M}$, then R is locally Gorenstein on $\operatorname{Ass}_R M$ by Remark 5.2. Thus the assertion follows from Corollary 4.14(3).
- (2) By Remark 4.12, the \widehat{R} -module \widehat{M} is faithful. If \widehat{R} is generically Gorenstein, then so is R by Remark 5.2. Hence the assertion follows from Corollary 4.14(4).

We close the section by posing a natural question, asking whether one can remove from Theorem 1.4(5) the assumption that M has finite G-dimension.

Question 5.4. Let R be a local ring. Suppose that there is an R-module M with dim $M = \dim R$ such that T_M exists and $T_M \cong M$. Is then R Gorenstein?

Corollary 3.8 guarantees that this question is affirmative in the case when the local ring R is artinian.

Acknowledgments. Part of this work was done during Takahashi's visit to the Florida A&M University in February, 2024. He is grateful to the members of the Department of Mathematics for their hospitality.

References

- L. L. AVRAMOV; J. HERZOG, Jacobian criteria for complete intersections, The graded case, *Invent. Math.* 117 (1994), no. 1, 75–88.
- [2] B. Briggs, Vasconcelos' conjecture on the conormal module, Invent. Math. 227 (2022), no. 1, 415-428.
- [3] W. Bruns; J. Herzog, Cohen-Macaulay rings, revised edition, Cambridge Studies in Advanced Mathematics 39, Cambridge University Press, Cambridge, 1998.
- [4] L. W. Christensen, Gorenstein dimensions, Lecture Notes in Math. 1747, Springer-Verlag, Berlin, 2000.
- [5] D. FERRAND, Suite régulière et intersection complète, C. R. Acad. Sci. Paris Sér. A-B 264 (1967), A427-A428.
- [6] A. GROTHENDIECK, Éléments de géométrie algébrique IV, Étude locale des schémas et des morphismes de schémas IV, *Inst. Hautes Études Sci. Publ. Math.* (1967), no. 32, 361 pp.
- [7] R. Hartshorne, Residues and duality, Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64, With an appendix by P. Deligne, Lecture Notes in Math. 20, Springer-Verlag, Berlin-New York, 1966.
- [8] C. Huneke; R. Wiegand, Tensor products of modules and the rigidity of Tor, Math. Ann. 299 (1994), no. 3, 449–476.
- [9] T. KAWASAKI, On arithmetic Macaulayfication of Noetherian rings, Trans. Amer. Math. Soc. 354 (2002), no. 1, 123-149.
- [10] H. MATSUMURA, Commutative algebra, Math. Lecture Note Ser. 56, Benjamin/Cummings Publishing Co., Inc., Reading, MA, 1980.
- [11] R. TAKAHASHI; D. WHITE, Homological aspects of semidualizing modules, Math. Scand. 106 (2010), no. 1, 5–22.
- [12] W. V. VASCONCELOS, Ideals generated by R-sequences, J. Algebra 6 (1967), 309-316.
- [13] W. V. VASCONCELOS, On the homology of I/I^2 , Comm. Algebra 6 (1978), 1801–1809.

(Mohsen Gheibi) DEPARTMENT OF MATHEMATICS, FLORIDA A&M UNIVERSITY, TALLAHASSEE FL, USA $Email\ address$: mohsen.gheibi@famu.edu

(Ryo Takahashi) Graduate School of Mathematics, Nagoya University, Furocho, Chikusaku, Nagoya 464-8602, Japan

 $Email\ address: \ {\tt takahashi@math.nagoya-u.ac.jp} \\ URL: \ {\tt https://www.math.nagoya-u.ac.jp/~takahashi/}$