# Generation of kink-antikink pairs through the excitation of an oscillon in the $\phi^4$ model

Fabiano C. Simas<sup>1,3</sup> and E. da Hora<sup>2,3\*</sup>

<sup>1</sup> Centro de Ciências de Chapadinha (CCCh), Universidade Federal do Maranhão (UFMA), 65000-000 Chapadinha, MA, Brazil

<sup>2</sup> Coordenação do Curso de Bacharelado Interdisciplinar em Ciência e Tecnologia,

Universidade Federal do Maranhão, 65080-805, São Luís, Maranhão, Brazil. <sup>3</sup> Programa de Pós-Graduação em Física, Universidade Federal do Maranhão (UFMA),

Campus Universitário do Bacanga, 65085-580, São Luís, Maranhão, Brazil

# Abstract

We examine the oscillon decay in a model with a single scalar field and two asymmetric fundamental states. Specifically, we verify how the evolution of an initially excited oscillon leads to the formation of antikink-kink pairs in the presence of a centered long-lived pulse. In such a context, the magnitude of the original perturbation stands for the most important factor influencing the process. For intermediary values of this magnitude, we find a transitory behavior which suggests that the energy exchange between translational and vibrational modes begins to play an important role in the overall evolution. As the magnitude increases, the centered pulse intermediates an interaction between the inner structures which compose the pairs whose behavior mimics that of a standard antikink-kink collision and therefore reinforces the role played by the energy transfer mechanism. We depict the resulting resonant structure and note that it mimics that of an usual kink-antikink scattering. We also reproduce some of our results now via an initial lumplike perturbation proposed as a kink-antikink pair. For different values of width, *an almost pure* antikink-kink pair escapes to the infinity after multiple bounces.

<sup>\*</sup>Electronic address: fc.simas@ufma.br, carlos.hora@ufma.br

## I. INTRODUCTION

In the context of classical field models, topological defects are usually described as the solutions to highly nonlinear equations of motion. In this case, the nonlinearity is introduced via the potential which describes the self-interaction of the scalar matter and, as a consequence, also determines the vacuum structure of the corresponding theory [1].

In such a scenario, localized configurations connect two adjacent minima of the potential, with both the solution and its anti-version having the very same total energy, but topological charges with opposite signs.

Under special circumstances, these profiles can be also obtained via a set of first-order differential equations which emerges from the minimization of the energy of the model through the implementation of the so-called BPS technique [2, 3]. In this case, it is possible to verify that these first-order equations lead to the standard second-order equations of motion, from which one concludes that the BPS configurations are legitimate solutions of the effective theory indeed.

In this overall context, localized structures have been studied in connection to a wide variety of physical systems throughout the last few years, ranging from the subatomic to the cosmological scale [4, 5]. Furthermore, topological defects are particularly common in high-energy issues [6], optical communication [7] and DNA [8].

The topological structures which emerge in a (1 + 1)-dimensional scalar field theory are known as kinks. In such a relativistic ambient, these originally time-independent BPS solutions acquire dynamics via the implementation of a Lorentz boost, from which the evolution of the resulting travelling profiles must be studied via the numerical treatment of the effective equations of motion.

The evolution of travelling kinks have been widely investigated. Among some of the fields of research, we can highlight the study of kink scattering, which revealed a surprisingly complex structure of bounces and annihilation. The final result of the scattering process is intimately influenced by the value of the initial velocity of the colliding structures, therefore giving rise to resonance windows [9–11]. Additional discussions in the context of most diverse models can also be found in [12–31].

At the same time, the study of the so-called oscillons and spharelons has been the subject of recent investigations. The sphaleron stands for a static, finite and unstable solution which appears in Skyrme models and electroweak theories [32]. It was demonstrated in the Ref. [33] that the spharelon serves as the source for the internal mode which gives rise to the fractal structures in a kink-antikink collision. Oscillons, on the other hand, are spatially localized oscillatory solutions which present a long lifetime. Their stability can not assured because due to the lack of a well-defined topological charge [34, 35]. These solutions naturally occur from the collapse of bubbles [36] and can be also formed in the context of a domain wall network [37].

Interestingly, connections between spharelons and oscillons have been discovered recently. In this sense, the Ref. [38] has examined a spharelon which, when slightly perturbed, decays into a large amplitude oscillon. In addition, the Ref. [39] has also investigated the spharelonoscillon relation, with the authors showing that the presence of the internal mode of the spharelon is not a necessary ingredient in the dynamics of the oscillon.

A recent research has revealed the creation of kink-antikink pairs as a result of collisions between two identical wave trains. In that work, the authors have considered the  $\phi^4$  model, from which they have identified that the resonant excitation plays an important role in the formation of the new pair, with the process being determined by the amplitude [40]. Most recently, a scenario with only one symmetric vacuum was proposed. In such a case, the formation of a fractal structure from the decay of excited oscillons [41] was verified. Here, the most intriguing aspect is that the effective model does not support any static solutions. However, the resonant window structure, which is based on the energy exchange between the translational and vibrational modes, is realized. Examine also the Refs. [42–45], which address pair production.

The aim of the present manuscript is to contribute with such a research. In this sense, we study the production of antikink-kink pairs as a result of the decay of an oscillon initially excited around a vacuum state in the  $\phi^4$  theory. In order to present our developments, this work is organized as follows: in the next Section II, we introduce the particular model and comment on its general aspects. In the sequence, we define the initial condition (a Gaussianlike oscillatory perturbation) which depends on two parameters, i.e., its amplitude and width. We are particularly interested in the effects brought about by a perturbation of different intensities. Still in the Sec. II, we observe that the corresponding solution for the field at the center of mass mimics the profile inherent to a multiple-bounce window with nontrivial order, which therefore suggests that the exchange of energy between the translational and vibrational modes via the energy transfer mechanism begins to play a relevant role. We then study the corresponding resonant structure and show that it reveals a fractal pattern which surprisingly mimics that of a standard kink-antikink scattering. Motivated by such a coincidence, we dedicate the Section III to the reproduction of the effects caused by a Gaussian perturbation now via a lumplike profile proposed as a kink-antikink pair, from which we argue that such an approach holds well in the limit of small amplitudes. Finally, the Sec. IV brings a brief summary and our perspectives regarding future investigations.

### II. THE MODEL AND ITS RESULTS

In this manuscript, we consider a (1+1)-dimensional theory which contains a single real scalar field  $\phi(x,t)$  only, its Lagrangian density being given by

$$\mathcal{L} = \frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi - V(\phi), \qquad (1)$$

where both the field, coordinates and coupling constants were made dimensionless via the implementation of an appropriate rescaling defined in terms of the mass scale of the model. As usual, the Greek index  $\mu$  runs from 0 to 1 and therefore counts the spacetime coordinates. Moreover, we have chosen to work with  $\eta^{\mu\nu} = (+-)$  as the metric signature of the Minkowski spacetime.

In such a context, the potential  $V(\phi)$  determines the self-interaction inherent to the real field via the specification of the vacuum structure of the corresponding theory. In the present case, we consider the well-known  $\phi^4$  model, whose potential reads

$$V(\phi) = \frac{1}{2}(1 - \phi^2)^2,$$
(2)

which has many applications in various physics scenarios. For instance, in condensed matter physics, a one-dimensional Ginzburg-Landau theory has been proposed to describe the phase transition in shape memory alloys [46]. Moreover, in Ref. [47], the authors found that temperature has a significant influence on the formation of kink-antikink pairs.

The resulting nonintegrable theory is known to support energetically stable timeindependent solutions which connect the two adjacent vacuum predicted by the  $\phi^4$  potential above. Specifically, while the resulting kink is given by  $\phi_K(x) = \tanh(x)$  and interpolates between the states  $\phi_K(-\infty) = -1$  and  $\phi_K(\infty) = 1$ , the corresponding antikink can be promptly obtained based on the  $Z_2$  symmetry inherent to the original model and therefore connects the adjacent vacuum in an opposite way, i.e.  $\phi_{\bar{K}}(-\infty) = 1$  and  $\phi_{\bar{K}}(\infty) = -1$ . These two profiles can be verified to have the very same energy, but topological charges with opposite signs. Furthermore, the stability analysis reveals the existence of two distinct modes - translational and vibrational ones - inherent to the kink profile. In such a case, the evolution of a kink-antikink collision is directly affected by the existence of the vibrational mode (related to  $\omega_1 = \sqrt{3}$ ) via the so-called *energy transfer mechanism*. As a result, two-bounce windows appear and a typical fractal structure can be observed.

In this manuscript, instead of colliding kinks, we focus our attention on the generation of oscillating pulses and kinks as a result of oscillons decay. With such a purpose in mind, we consider as the initial condition

$$\phi(x,0) = 1 + A_0 e^{-x^2/\sigma},\tag{3}$$

where the parameter  $A_0$  represents the amplitude and  $\sigma$  stands for the width of a perturbation (with a Gaussian profile) around the positive vacuum  $\phi_0 = +1$  of the  $\phi^4$  potential. As we demonstrate below, depending on the values of these parameters, we find more or less excited oscillons, which eventually give rise to different final states of the field  $\phi(x,t)$ . Here, it is worthwhile to point out that such an initial data is comparable to that very recently used in the Ref. [41], where the authors have successfully studied oscillons decay in a model with a symmetric vacuum only and whose potential can be approximated by the sine-Gordon one.

In what follows, we investigate the evolution of the field configuration in the full space for different values of  $A_0$  and with an initially fixed  $\sigma$ . At this first moment, we are interested in small perturbations around the vacuum state, from which we consider very small values of the amplitude.

The Figure 1 (top) shows the evolution of  $\phi(x,t)$  for  $\sigma = 10$  and  $A_0 = 0.05$  (left) and 0.15 (right). In such a context, it is important to note how the original (small) perturbation produces an excited oscillon around the vacuum which decays and annihilates. In this sense, the field profile at the center of mass as a function of t is displayed in the Fig. 1 (bottom), which reveals how the amplitude of the oscillations around  $\phi_0 = +1$  decreases. It is therefore reasonable to conclude that, when submitted to a relatively small Gaussian-like perturbation around its vacuum value, the scalar sector tends to return to its fundamental state. In other

words, the corresponding perturbation are not intense enough to remove the field from its vacuum value and then generate a nontrivial configuration. Remarkably, unlike what is seen in the Ref. [41], the original perturbation in the present  $\phi^4$  scenario does not decompose into two (or more) constituent oscillons.



FIG. 1: Evolution of the scalar sector  $\phi(x, t)$  in the full space (top) and the field profile at the center of mass (bottom) for  $A_0 = 0.05$  (left) and 0.15 (right). Here,  $\sigma = 10$ . The oscillon annihilates and the field returns to its fundamental state.

In the sequence, in order to explore the impact of parameter values on the oscillon development process, we consider also the case of not so small perturbations, from which we solve the numerical problem for additional values of  $A_0$ . The corresponding solutions appear in the figures below.

The Figure 2 brings the results for intermediary  $A_0$ , from which it is possible to infer the existence of a range of values for which the solutions present a transitory behavior. In particular, for  $A_0 = 0.30$  and 0.40, the perturbation produces a long-lived oscillon which, in view of the nonintegrable nature of the theory, annihilates after an extremely long time and induces the real field to finally return to its vacuum state. On the other hand, the solution for  $A_0 = 0.50$  reveals the occurrence of a *perturbed pulse (i.e. with anharmonic oscillations)* whose existence might be connected to the exchange of energy between the translational and vibrational modes through the energy transfer mechanism.



FIG. 2: Evolution of the scalar sector  $\phi(x,t)$  in the full space (top) and the field profile at the center of mass (bottom) for  $A_0 = 0.30$  (left), 0.40 (center) and 0.50 (right). Here,  $\sigma = 10$ . The solutions present a transitory behavior, with the anharmonic pulse possibly indicating the exchange of energy between the translational and vibrational modes.

The suggestion above is corroborated by the solutions which appear in the Figure 3. Here, for  $A_0 = 0.60$ , the Gaussian perturbation oscillates and, through the emission of radiation, rapidly decays into a antikink-kink pair with a central long-lived small oscillon around the negative vacuum  $\phi_0 = -1$ . In addition, the evolution for  $A_0 = 0.70$  shows the scattering of two *almost pure* antikink-kink pairs, from which we conclude that an increasing amplitude generates an even more excited oscillon whose decay leads to the creation of such pairs followed by the escape of the radiation. Similar phenomenon was studied in the Ref. [48], where the scattering of two lumps was verified to produce kink-antikink pairs, while multiple antikink-kink pairs were obtained in a double sine-Gordon model [49]. An antikink-kink pair can also be obtained via the collision of two identical wave trains, see the Ref. [40]. Still regarding our solution for  $A_0 = 0.70$ , we highlight that, once the time evolution begins, a centered oscillon emerges and then intermediates a "collision" between the inner structures whose corresponding pattern mimics the one typically related to a four-bounce window, see the resulting profile for  $\phi(0, t)$ . In particular, between the first two bounces, the scalar field performs two complete oscillations in a clear manifestation of the energy exchange between the translational and vibrational modes, such a behavior reinforcing the role played by the energy transfer mechanism. Moreover, when  $A_0 = 0.75$ , the perturbation produces a pair of bions and a central oscillon around  $\phi_0 = +1$ .



FIG. 3: Evolution of the scalar sector  $\phi(x,t)$  in the full space (top) and the field profile at the center of mass (bottom) for  $A_0 = 0.60$  (left), 0.70 (center) and 0.75 (right). Here,  $\sigma = 10$ . The Gaussian perturbation rapidly decays into kink-antikink pairs.

The final configurations obtained for high amplitudes are shown in the Figure 4. In general, the initial condition oscillates and then generates two antikink-kink pairs which propagate without considerable radiation escape, while the central region is distinguished by the existence of oscillons which travel around while emitting a certain quantity of radiation. Interestingly, there are regions where the initial excitation produces two oscillons which escape to the infinity, again irradiating. In particular, two antikink-kink pairs appear for  $A_0 = 0.80$  and 0.90. However, there are now two basic differences in comparison to the previous occurrence: long-lived oscillations around  $\phi_0 = +1$  take place at the center of mass, and the arising antikink-kink pairs do not bounce. Note that three non-bouncing pairs emerge for  $A_0 = 0.95$ .



FIG. 4: Evolution of the scalar sector  $\phi(x, t)$  in the full space (top) and the field profile at the center of mass (bottom) for  $A_0 = 0.80$  (left), 0.90 (center) and 0.95 (right). Here,  $\sigma = 10$ . The initial condition generates antikink-kink pairs which propagate without considerable radiation escape. In particular, three non-bouncing pairs emerge for  $A_0 = 0.95$ .

It sounds relevant to study the resonant structure related to the evolution of the scalar field in view of the Gaussian perturbation. In this sense, the Figure 5 summarizes the main behaviors involving the oscillon decay, as well as the appearance of antikink-kink pairs. Specifically, this plot shows the temporal evolution of the field value at the center of mass for fixed  $\sigma = 10$  and different values of  $A_0$ , from which we identify well-defined segments which alternate between blue and red colors. Here, in view of the previous figures which bring the evolution of  $\phi(x, t)$  in the full space, it is possible to understand that the blue regions represent a field oscillation around  $\phi_0 = +1$ , whilst the red portions means the emergence of an antikink-kink pair at the origin. For instance, small values of  $A_0$  lead to either an annihilation of the original perturbation or the generation of a long-lived oscillon. On the other hand, when  $0.5 < A_0 < 0.7$ , there are stripes in red which represent a single antikink-kink pair.

Notably, we also note a fractal structure which resembles the outcome of kink-antikink collisions. In this case, in order to recognize the details of the resonant structure, we have included a zoom, see the plots in the center and in the left of the Figure 5. It is important to make clear that in kink-antikink scatterings, the resonant structure reflects the number of interactions between the pair. In our case, however, the graphic depicts the alternation between the emergence of an oscillation and an antikink-kink pair.



FIG. 5: The resonant structure related to the evolution of the scalar field in view of a Gaussian perturbation. Blue regions represent a field oscillation around  $\phi_0 = +1$ , whilst the red portions means the emergence of an antikink-kink pair at the origin. The resulting fractal structure resembles that of a kink-antikink collision. Here, we fixed  $\sigma = 10$ .

Next, a new blue band appears as the value of the amplitude increases. This area comprises either the formation of oscillatory pulses in the central region or the emergence of two antikink-kink pairs which escape to infinity. For those values of  $A_0$  greater than 0.90, an intriguing aspect is revealed, i.e. this region presents multiple red peaks, as can be seen in the Fig. 5 (left), which suggests the formation of two pairs which scatter symmetrically and an additional pair located at x = 0.

We end this section by exploring the effects related to an initial oscillation with a different width. The numerical results are depicted in the Figure 6 for  $\sigma = 20$  and different values of  $A_0$ . In this case, a single antikink-kink pair is generated by the original perturbation whose amplitude now determines the number of bounces performed by the pair. For instance, when  $A_0 = 0.55$ , the initial configuration oscillates and briefly separates. However, it lacks sufficient kinetic energy and hence the pair collides. Only after this first collision, the process of complete separation of the pair begins. Interestingly, we see that the pair formation occurs only after two collisions (which corresponds to a first-order two-bounce window) for  $A_0 = 0.53$ . In the last case, the solution formed when  $A_0 = 0.5346$  remains trapped for a long time, i.e., it performs three complete collisions (with a second-order three-bounce window near to a seventh-order two-bounce one) and only then the antikink-kink pair escapes to the infinity. Again, these results emphasize the role played by the energy transfer mechanism during the generation of antikink-kink pairs due to an original Gaussian perturbation.



FIG. 6: Evolution of the scalar sector  $\phi(x,t)$  in the full space (top) and the field profile at the center of mass (bottom) for  $A_0 = 0.55$  (left), 0.53 (center) and 0.5346 (right). Here,  $\sigma = 20$ . The amplitude determines the number of bounces performed by the single pair, with two (three) collisions occurring for  $A_0 = 0.53$  ( $A_0 = 0.5346$ ).

# III. THE PERTURBATION ITSELF AS A KINK-ANTIKINK PAIR

As we have pointed out previously, the  $\phi^4$  model is known to support the existence of time-independent kinks with finite energy and whose scattering process has been intensively studied, see the Refs. [9, 10]. Interestingly, the temporal evolution of the real field  $\phi(x, t)$ due to a Gaussian perturbation applied to its fundamental state not only reveals a fractal pattern which is similar to that of a standard kink-antikink collision, but it is also suggested to occur according to the energy exchange between the translational and vibrational modes related to the aforementioned kink profile. In view of these facts, it sounds reasonable to conjecture that the evolution of  $\phi(x,t)$  when submitted to a Gaussian perturbation can eventually be mimicked by a lumplike profile.

The present section is devoted to the study of such a possibility. In this sense, we first highlight that the original Gaussian perturbation can be approximated by a static lump constructed as a kink-antikink pair. Therefore, instead of  $\phi(x,0) = 1 + A_0 e^{-x^2/\sigma}$ , we now adopt the initial condition as

$$\phi(x,0) = 1 + \mathcal{A}_0 \left[ \tanh(x + |x_0|) - \tanh(x - |x_0|) \right], \tag{4}$$

where  $\mathcal{A}_0$  represents the magnitude of the lumplike perturbation, while  $2x_0$  stands for the distance between the kink and the antikink which compose the pair (therefore representing the "width" of the corresponding lump, whose center is assumed to be positioned at x = 0). As we argue below, this general form can be used to map the results which arise from the Gaussian perturbation in terms of the properties inherent to a canonical kink-antikink interaction.

It is useful to write down the Taylor series expansions for both the Gaussian and the lumplike perturbations. In this case, whether we compare the relevant terms in both series, one gets that  $\mathcal{A}_0$  and  $x_0$  can be expressed in terms of  $\mathcal{A}_0$  and  $\sigma$  as

$$\mathcal{A}_0 = \frac{A_0}{2\sqrt{1 - \frac{1}{\sigma}}},\tag{5}$$

and

$$|x_0| = \operatorname{arctanh}\left[\sqrt{1 - \frac{1}{\sigma}}\right],\tag{6}$$

via which we promptly rewrite the initial condition in the form

$$\phi(x,0) = 1 + \frac{A_0}{2\sqrt{1-\frac{1}{\sigma}}} \left[ \tanh\left(x + \operatorname{arcsech}\left(\sigma^{-\frac{1}{2}}\right)\right) - \tanh\left(x - \operatorname{arcsech}\left(\sigma^{-\frac{1}{2}}\right)\right) \right], \quad (7)$$

where we have assumed  $\sigma > 1$  in order to satisfy  $|x_0| > 0$ .

The expression above can be used as a good approximation to describe the results obtained in the limit of small  $A_0$ . For instance, the Figure 7 brings a comparison between the profiles for  $\phi(0, t)$  obtained via the Gaussian (blue line) and lumplike (red line) perturbations. Here, we have fixed  $\sigma = 10$  and again solved the problem for  $A_0 = 0.05$ , 0.15 and 0.25. It is possible to see that, despite some differences (of order  $10^{-2}$  and which decrease with time) in the corresponding amplitudes, the solutions behave in the very same qualitative way. In particular, they suggest that the central long-lived pulse which emerges for small values of the original amplitude can be eventually treated as a kink-antikink pair which perform several collisions without getting enough kinetic energy to escape from each other. Due to the nonintegrable nature of the overall scenario, the colliding pair continuously emits radiation and finally disappears, forcing the field to return to its fundamental state, therefore mimicking the very same [qualitative] picture which arises from a Gaussian perturbation.



FIG. 7: Field profile at the center of mass for  $A_0 = 0.05$  (left),  $A_0 = 0.15$  (center) and  $A_0 = 0.25$  (right). Here, we have fixed  $\sigma = 10$ . The results obtained via the Gaussian (lumplike) perturbation appears as the blue (red) line. For small  $A_0$ , the solutions behave in the very same qualitative way.

As the value of  $A_0$  increases, the resulting solutions assume different profiles (both in magnitude and frequency). As a consequence, the aforementioned correspondence loses its accuracy and can not be applied anymore, see the profiles for  $A_0 = 0.25$ , for instance.

It is always possible to improve the approximation above by fitting the Gaussian and the lumplike profiles via a rescaling applied on x in the expression for the last one. As the interested reader can verify, such a procedure gives rise to even better results (in comparison to the ones displayed in the previous Figure 7). However, the corresponding accuracy is still limited by the values of  $A_0$ .

Here, in order to improve the accuracy of the Taylor series based approach, we adopt an alternative way, i.e. we assume small values of the width  $\sigma$ . In this case, the Gaussian profile is even more localized around x = 0, from which the series expansion approximation consequently applies in a even better way. The results for  $\sigma = 2$  and  $A_0 = 0.30$ , 0.60 and 0.90 appear in the Figure 8, from which it is possible to see how accurate [both qualitatively and quantitatively] the results provided by the lumplike perturbation are. It is however important to reinforce that the corresponding final configurations dramatically depend on the values of  $\sigma$  itself, see the results displayed in the Figure 6 for  $\sigma = 20$ , for instance. In this sense, we clarify that, when  $\sigma = 2$ , the final structure solution is a short-lived (longlived) centered pulse for  $A_0 = 0.30$  ( $A_0 = 0.60$ ), while it represents the scattering of a single antikink-kink pair (with a central pulse around the negative vacuum) for  $A_0 = 0.90$  (i.e. multiple antikink-kink pairs were not observed).



FIG. 8: Field profile at the center of mass for  $A_0 = 0.30$  (left),  $A_0 = 0.50$  (center) and  $A_0 = 0.90$  (right). Here, we have fixed  $\sigma = 2$ . The Gaussian profile is even more localized around the origin, from which the series expansion provides better results.

In all these cases, the original Gaussian perturbation was approached via a lumplike profile which was proposed as a static kink-antikink pair. The results obtained via the Gaussian configuration were then mimicked via those related to a kink-antikink interaction. We believe that such an approach contributes to the comprehension on the role played by the energy transfer mechanism in the formation of the final structures. In addition, the lumplike perturbation approach is expected to provide some insight about the occurrence of a fractal pattern which resembles that of a kink-antikink collision.

## IV. SUMMARY AND PERSPECTIVES

In this manuscript, we have studied some of the results caused by a perturbation applied to a fundamental state of a classical field.

With this aim in mind, we have considered a model with a single real scalar field  $\phi(x, t)$ only whose self-interaction potential  $V(\phi)$  has the usual  $\phi^4$  profile. The resulting nonintegrable scenario is known to support energetically stable time-independent solutions which connect the two adjacent minima of the potential. In addition, these solutions can be verified to have the very same energy, but topological charges with opposite signs. Moreover, the stability analysis reveals that such a kink profile admits two well-defined internal modes - translational and vibrational - which can be applied to explain the temporal evolution of an scattering in such a classical ambient via the so-called *energy transfer mechanism*.

Motivated by the prominent developments recently presented in the Ref. [41], we have here implemented a Gaussian-like perturbation around the vacuum state  $\phi_0 = +1$ . The perturbation is controlled by two distinct parameters, namely,  $A_0$  and  $\sigma$ , which rotulate its amplitude and width, respectively.

We have then focused our attention on how different values of  $A_0$  give rise to distinct configurations of the real sector  $\phi(x, t)$ , from which we have considered some  $\sigma$  values in our numerical investigations.

Initially, we have investigated the impact of small amplitudes. These solutions have revealed that the perturbation produces an excited oscillon which decays and annihilates. We have also explored the scenario of intermediate values of  $A_0$ , where the dynamics has indicated the creation of long-lived oscillons. As the amplitude increases, we have observed the formation of antikink-kink pairs followed by the eventual emission of radiation. In particular, we have identified the appearance of a centered short-lived oscillon which intermediates the interaction between the inner structures which compose the emergent pairs. Curiously, we have noted that the corresponding solution for  $\phi(0, t)$  mimics the pattern inherent to a four-bounce window, with the scalar field performing two complete oscillations between the first two bounces, from which we have argued that such a behavior reinforces the role played by the energy transfer mechanism. For large amplitudes, the initial condition oscillates and then generates two pairs which propagate in opposite directions, followed by an oscillon at the center of mass. Additionally, we have found that a new antikink-kink pair was formed.

Motivated by these results, we have obtained the resonant structure inherent to the evolution of the scalar field in view of an initial Gaussian perturbation, see the Fig. 5. We have identified field oscillations around  $\phi_0 = +1$  (blue regions) and the emergence of an antikink-kink pair positioned at x = 0 (red portions). Notably, we have noted that the resulting fractal structure mimics that of a kink-antikink scattering. Moreover, we have clarified that, while the resonant structure related to a standard kink-antikink collision reflects the number of interactions between the pair, the structure which we have obtained in the present manuscript depicts the alternation between the emergence of an oscillatory pulse and an antikink-kink pair.

We have also investigated the impact of an initial oscillation with a different width, from which we have fixed  $\sigma = 20$ . In these circumstances, we have observed that a single antikinkkink pair is generated by the perturbation, with the amplitude now determining the number of bounces performed by the pair before its escape to the infinity. We have pointed out that these multiple bounces can be rotulated as usual, i.e., based on the order of the window they belong. As a result, we have highlighted that the existence of these windows with nontrivial order can be understood as a clear consequence of the energy exchange between the translational and vibrational modes.

The coincidence between the fractal patterns has inspired us to study the reproduction of the effects caused by the Gaussian perturbation via a lumplike profile constructed as a kink-antikink pair. We have connected these two scenarios via the comparison between the respective Taylor series expansions, from which we have written the parameters which characterize the lumplike perturbation in terms of the  $A_0$  and  $\sigma$ . We have then depicted the corresponding results, which have revealed that such a Taylor series based approach holds well for small values of  $A_0$ , its accuracy decreasing as the values of  $A_0$  increase. In particular, when  $\sigma$  is small, the results practically coincide with each other, from what we have pointed out that such a successful mapping may contribute to the understanding not only of the coincidence between the fractal pattern, but also of the fundamental role played by the energy transfer mechanism in the definition of the final configurations due to a Gaussian perturbation.

Perspectives regarding future investigations include the study of the results related to an initial perturbation around the fundamental states inherent to both the  $\phi^6$  and the sine-Gordon potentials. In this context, the last one sounds particularly interesting due to the integrable nature of the corresponding ambient, from which emission of radiation is not expected to occur. Also, we plan to depict the resonant structures of the resulting configurations, from which we intend to verify whether the corresponding fractal patterns can be eventually compared to the well-established ones or, on the other hand, they reveal a novel profile to be explored. These issues are currently under investigation, and positive results will be eventually reported in a future contribution.

## Acknowledgements

The authors thank João Campos and Azadeh Mohammadi for their useful comments on the present manuscript. EH also thanks the partial financial support received from the Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq (Brazillian agency) via the Grant N° 309604/2020-6. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

- N. Manton, P. Sutcliffe, *Topological Solitons*, Cambridge University Press, Cambridge U. K. (2004).
- [2] M. K. Prasad, Charles M. Sommerfield, Exact Classical Solution for the 't Hooft Monopole and the Julia-Zee Dyon, Phy. Rev. Lett. 35, 12 (1975).
- [3] E. B. Bogomolny, Stability of Classical Solutions, Sov. J. Nucl. Phys. 24, 449 (1976).
- [4] T. Dauxois, *Physics of Solitons*, Cambridge University Press, Cambridge (2010).
- [5] T. Vachaspati, Kinks and Domain Walls: An Introduction to Classical and Quantum Solitons, Cambridge University Press, Cambridge U. K. (2006).
- [6] John T. Giblin Jr, Lam Hui, E. A. Lim, I-Sheng Yang, How to run through walls: dynamics of bubble and soliton collisions, Phys. Rev. D 82, 045019 (2010).

- [7] L. F. Mollenauer, J. P. Gordon, Solitons in Optical Fibers Fundamentals and Applications, Academic Press, Burlington, MA, U.S.A. (2006).
- [8] L.V. Yakushevich, Nonlinear Physics of DNA, Wiley-VCH, Weinheim, Germany, (2004).
- [9] D. K. Campbell, J. F. Schonfeld, C. A. Wingate, Resonance structure in kink-antikink interactions in φ<sup>4</sup> theory, Physica D 9, 1 (1983).
- [10] P. Anninos, S. Oliveira, R. A. Matzner, Fractal structure in the scalar  $\lambda(\phi^2 1)^2$  theory, Phys. Rev. D 44, 1147 (1991).
- [11] P. Dorey, K. Mersh, T. Romańczukiewicz, Ya. Shnir, Kink-antikink collisions in the φ<sup>6</sup> model,, Phys. Rev. Lett. **107**, 091602 (2011).
- [12] I. Takyi, S. Gyampoh, B. Barnes, J. Ackora-Prah, G. A. Okyere, Kink collision in the noncanonical φ<sup>6</sup> model: A model with localized inner structures, Results in Physics 44, 106197 (2023).
- F. C. E. Lima, C. A. S. Almeida, Aspects of Kink-Like Structures in 2D Dilaton Gravity, Fortschr. Phys. 71, 2300051 (2023).
- [14] T. Mendonça, H. de Oliveira, The collision of two-kinks defects, J. High Energ. Phys. 2015, 120 (2015).
- [15] H. Weigel, Kink-Antikink Scattering in  $\phi^4$  and  $\phi^6$  Models, J. Phys. Conf. Ser. 482, 012045 (2014).
- [16] V. A. Gani, A. Moradi Marjaneh, A. Askari, E. Belendryasova, D. Saadatmand, Scattering of the double sine-Gordon kinks, Eur. Phys. J. C 78, 345 (2018).
- [17] Dionisio Bazeia, João G. F. Campos, Azadeh Mohammadi, Kink-antikink collisions in the  $\phi^8$ model: short-range to long-range journey, J. High Energ. Phys. 05, 116 (2023).
- [18] F. C. Simas, Adalto R. Gomes, K. Z. Nobrega, J. C. R. E. Oliveira, Suppression of two-bounce windows in kink-antikink collisions, J. High Energ. Phys. 2016, 104 (2016).
- [19] D. Bazeia, E. Belendryasova, V. A. Gani, Scattering of kinks of the sinh-deformed φ<sup>4</sup> model, Eur. Phys. J. C 78, 340 (2018).
- [20] A. Moradi Marjaneh, F. C. Simas, D. Bazeia, Collisions of kinks in deformed  $\phi^4$  and  $\phi^6$ models, Chaos, Solitons and Fractals **164**, 112723 (2022).
- [21] Dionisio Bazeia, João G. F. Campos, Azadeh Mohammadi, Resonance mediated by fermions in kink-antikink collisions, J. High Energ. Phys. 2022, 85 (2022).
- [22] Alonso-Izquierdo Alberto, Non-topological kink scattering in a two-component scalar field the-

ory model, Commun. Nonlin. Sci. Numer. Simul. 85, 105251 (2020).

- [23] A. Halavanau, T. Romanczukiewicz, Y. Shnir, Resonance structures in coupled two-component φ<sup>4</sup> model, Phys. Rev. D 86, 085027 (2012).
- [24] F. C. Lima, F. C. Simas, K. Z. Nobrega, A. R. Gomes, Boundary scattering in the φ<sup>6</sup> model,
  J. High Ener. Phys. 10, 147 (2019).
- [25] Ivan C. Christov, Robert J. Decker, A. Demirkaya, Vakhid A. Gani, P. G. Kevrekidis, Avinash Khare, Avadh Saxena, *Kink-kink and kink-antikink interactions with long-range tails*, Phys. Rev. Lett. **122**, 171601 (2019).
- [26] A. Alonso Izquierdo, J. Queiroga-Nunes, L. M. Nieto, Scattering between wobbling kinks, Phys. Rev. D 103, 045003 (2021).
- [27] C. Adam, K. Oles, T. Romanczukiewicz, A. Wereszczynski, Spectral Walls in Soliton Collisions, Phys. Rev. Lett. 122, 241601 (2019).
- [28] C. Adam, K. Oles, T. Romanczukiewicz, A. Wereszczynski, W. J. Zakrzewski, Spectral walls in multifield kink dynamics, J. High Ener. Phys. 08, 147 (2021).
- [29] C. Adam, K. Oles, T. Romanczukiewicz, A. Wereszczynski, Spectral Walls in Soliton Collisions, Phys. Rev. Lett. 122, 241601 (2019).
- [30] J. G. F. Campos, A. Mohammadi, J. M. Queiruga, A. Wereszczynski, W. J. Zakrzewski, *Fermionic spectral walls in kink collisions*, J. High Ener. Phys. **01**, 071 (2023).
- [31] João G. F. Campos, Mohammadi Azadeh, Fermions on wobbling kinks: normal versus quasinormal modes, J. High Ener. Phys. 2021, 103 (2021).
- [32] N. S. Manton, The inevitability of sphalerons in field theory, Philos. Trans. R. Soc. Lond. A 377 (2161), 20180327 (2019).
- [33] C. Adam, D. Ciurla, K. Oleś, T. Romańczukiewicz, A. Wereszczyński, Sphalerons and resonance phenomenon in kink-antikink collisions, Phys. Rev. D 104, 105022 (2021).
- [34] Marcelo Gleiser, *Pseudostable bubbles*, Phys. Rev. D 49, 2978 (1994).
- [35] Marcelo Gleiser, David Sicilia, General theory of oscillon dynamics, Phys. Rev. D 80, 125037 (2009).
- [36] E. J. Copeland, M. Gleiser, H. -R. Müller, Oscillons: Resonant configurations during bubble collapse, Phys. Rev. D 52, 1920 (1995).
- [37] Mark Hindmarsh, Petja Salmi, Oscillons and domain walls, Phys. Rev. D 77, 105025 (2008).
- [38] N. S. Manton, T. Romańczukiewicz, Simplest oscillon and its sphaleron, Phys. Rev. D 107,

085012(2023).

- [39] K. Oles, J. Queiruga, T. Romanczukiewicz, A. Wereszczynski, Sphaleron without shape mode and its oscillon, Phys. Lett. B 847, 138300 (2023).
- [40] Y. Shnir, T. Romanczukiewicz, Oscillon resonances and creation of kinks in particle collisions, Phys. Rev. Lett. 105, 081601 (2010).
- [41] F. Blaschke, T. Romańczukiewicz, K. Slawińska, A. Wereszczyński, Amplitude modulations and resonant decay of excited oscillons, arXiv:2403.00443 2403, 00443 (2024).
- [42] Omer Albayrak, Tanmay Vachaspati, Creating kinks with quantum mediation, Phys. Rev. D 109, 036001 (2024).
- [43] S. V. Demidov and D. G. Levkov, Soliton-Antisoliton Pair Production in Particle Collisions, Phys. Rev. Lett. 107, 071601 (2011).
- [44] S. V.Demidov, D. G. Levkov, Soliton pair creation in classical wave scattering, J. High Energ. Phys. 2011, 16 (2011).
- [45] C. S. Carvalho, L. Perivolaropoulos, Kink-antikink formation from an oscillation mode by sudden distortion of the evolution potential, Phys. Rev. D 79, 065032 (2009).
- [46] F. Falk, Ginzburg-Landau theory and solitary waves in shape-memory alloys, Z. Phys. B 54, 159 (1984).
- [47] D. Yu. Grigoriev, V. A. Rubakov, Soliton pair creation at finite temperatures: Numerical study in (1,1) dimensions, Nuclear Phys. B 299, 67 (1988).
- [48] F. C. Lima, F. C. Simas, K. Z. Nobrega, A. R. Gomes, Scattering of metastable lumps in a model with a false vacuum, Phys. Lett. B 822, 136707 (2021).
- [49] F. C. Simas, F. C. Lima, K. Z. Nobrega, A. R. Gomes, Solitary oscillations and multiple antikink-kink pairs in the double sine-Gordon model, J. High Energ. Phys. 2020, 143 (2020).