

# Affine Yangians and some cosets of non-rectangular $W$ -algebras

Mamoru Ueda\*

Department of Mathematical and Statistical Sciences, University of Alberta,  
11324 89 Ave NW, Edmonton, AB T6G 2J5, Canada

## Abstract

We construct a homomorphism from the affine Yangian associated with  $\widehat{\mathfrak{sl}}(q_u - q_{u+1})$  to the universal enveloping algebra of a  $W$ -algebra associated with  $\mathfrak{gl}(\sum_{s=1}^l q_s)$  and a nilpotent element of type  $(1^{q_1-q_2}, 2^{q_2-q_3}, \dots, (l-1)^{q_{l-1}-q_l}, l^{q_l})$  by using the coproduct and evaluation map for the affine Yangian and the Miura map for non-rectangular  $W$ -algebras. As an application, we give a homomorphism from the affine Yangian to some cosets of non-rectangular  $W$ -algebras.

## 1 Introduction

The AGT conjecture suggests the existence of a representation of the principal  $W$ -algebra of type  $A$  on the equivariant homology space of the moduli space of  $U(r)$ -instantons. Schiffmann and Vasserot [23] gave this representation by using an action of the Yangian associated with  $\widehat{\mathfrak{gl}}(1)$  on this equivariant homology space. More generally, Crutzig-Diaconescu-Ma [4] conjectured that an action of an iterated  $W$ -algebra of type  $A$  on the equivariant homology space of the affine Laumon space will be given through an action of a shifted affine Yangian constructed in [7].

In order to resolve the Crutzig-Diaconescu-Ma's conjecture, we need to construct a surjective homomorphism from the shifted affine Yangian to the universal enveloping algebra of the iterated  $W$ -algebra of type  $A$ . The iterated  $W$ -algebra of type  $A$  is expected to be the tensor of a  $W$ -algebra of type  $A$  and  $\beta\gamma$ -systems. Thus, the Crutzig-Diaconescu-Ma's conjecture can be considered as the affine analogue of Brundan-Kleshchev [3], which gives a surjective homomorphism from the shifted Yangian to a finite  $W$ -algebra of type  $A$ . However, the shifted affine Yangian is so complicated that it is difficult to construct the homomorphism of the Crutzig-Diaconescu-Ma's conjecture directly. In finite setting, there exist homomorphisms from finite Yangians to finite  $W$ -algebras of type  $A$ , which are restirictions of the one in [3]. One of these homomorphisms was given by De Sole-Kac-Valeri [6] by using the Lax operator. In [30] and [25], the author constructed the affine analogue of the homomorphism in De Sole-Kac-Valeri [6]. Let us take a positive number  $N$  and its partition:

$$N = q_1 + q_2 + \dots + q_l, \quad q_1 \geq q_2 \geq \dots \geq q_l$$

and set  $f$  as a nilpotent element of type  $(1^{q_1-q_2}, 2^{q_2-q_3}, \dots, (l-1)^{q_{l-1}-q_l}, l^{q_l})$ . In [25], we constructed a homomorphism from the affine Yangian associated with  $\widehat{\mathfrak{sl}}(q_l)$  to the universal enveloping algebra of a  $W$ -algebra  $\mathcal{W}^k(\mathfrak{gl}(N), f)$ , which is a  $W$ -algebra associated with  $\mathfrak{gl}(N)$  and  $f$  by using the the coproduct and evaluation map for the affine Yangian and the Miura map for non-rectangular  $W$ -algebras. Moreover, in [25], we show that the coproduct for the affine Yangian is compatible with the parabolic induction for a non-rectangular  $W$ -algebra via this homomorphism.

---

\*mueda@ualberta.ca

In [27], we gave the generalization of [30]. We constructed a homomorphism from the affine Yangian associated with  $\widehat{\mathfrak{sl}}(q_u - q_{u+1})$  to the universal enveloping algebra of  $\mathcal{W}^k(\mathfrak{gl}(N), f)$ . In this article, we gave another proof to [27]. Let us take positive integers  $\{n_a\}_{a=1}^{s+1}$  and  $\{b_a\}_{a=1}^s$  by

$$n_a > n_{a+1}, \quad n_{a-1} = q_{b_{a-1}} > q_{b_{a-1}+1} = \cdots = q_{b_a} = n_a > q_{b_a+1} = n_{a+1}, \quad n_{s+1} = 0.$$

By using the coproduct for the affine Yangian, we can construct a homomorphism

$$\Delta_l: \bigotimes_{1 \leq a \leq s} Y_{\hbar, \varepsilon - (n_a - q_l) \hbar} (\widehat{\mathfrak{sl}}(n_a - n_{a+1})) \rightarrow \widehat{\bigotimes}_{1 \leq v \leq l} Y_{\hbar, \varepsilon - (q_v - q_l) \hbar} (\widehat{\mathfrak{sl}}(q_v)),$$

where  $\widehat{\bigotimes}_{1 \leq v \leq l} Y_{\hbar, \varepsilon - (q_v - q_l) \hbar} (\widehat{\mathfrak{sl}}(q_v))$  is the standard degreewise completion of the tensor of affine Yangians. The Miura map ([15]) induces a homomorphism

$$\tilde{\mu}: \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f)) \hookrightarrow \widehat{\bigotimes}_{1 \leq i \leq l} U(\widehat{\mathfrak{gl}}(q_i)),$$

where  $\widehat{\bigotimes}_{1 \leq i \leq l} U(\widehat{\mathfrak{gl}}(q_i))$  is the standard degreewise completion of  $\bigotimes_{1 \leq i \leq l} U(\widehat{\mathfrak{gl}}(q_i))$ . We can connect  $\Delta_l$  to the Miura map  $\tilde{\mu}$  by using the evalution map for the Yangian:

$$\text{ev}^n: Y_{\hbar, \varepsilon - (n_a - q_l) \hbar} (\widehat{\mathfrak{sl}}(n)) \rightarrow \mathcal{U}(\widehat{\mathfrak{gl}}(n)),$$

where  $\mathcal{U}(\widehat{\mathfrak{gl}}(n))$  is the standard degreewise completion of the universal enveloping algebra of  $\widehat{\mathfrak{gl}}(n)$ .

**Theorem 1.1.** *There exists a homomorphism*

$$\Phi: \bigotimes_{1 \leq a \leq s} Y_{\hbar, \varepsilon - (n_a - q_l) \hbar} (\widehat{\mathfrak{sl}}(n_a - n_{a+1})) \rightarrow \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f))$$

satisfying the relation

$$\tilde{\mu} \circ \Phi = \bigotimes_{1 \leq i \leq l} \text{ev}^{q_i} \circ \Delta_l.$$

The homomorphism induces a homomorphism

$$\Phi_a: Y_{\hbar, \varepsilon - (n_a - q_l) \hbar} (\widehat{\mathfrak{sl}}(n_a - n_{a+1})) \rightarrow \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f))$$

for  $1 \leq a \leq s$ , which are homomorphisms given in [30] and [27]. Theorem 1.1 will be connected to the Gaiotto-Rapcak's triality.

Gaiotto and Rapcak [9] introduced a vertex algebra called the  $Y$ -algebra. The  $Y$ -algebra can be interpreted as a truncation of  $\mathcal{W}_{1+\infty}$ -algebra ([8]) whose universal enveloping algebra is isomorphic to the affine Yangian of  $\widehat{\mathfrak{gl}}(1)$  up to suitable completions (see [2], [24] and [21]). Gaiotto-Rapcak [9] conjectured a triality of the isomorphism of  $Y$ -algebras. Let us take a nilpotent element  $f \in \mathfrak{sl}(m+n)$  of type  $(n^1, 1^m)$ . It is known that some kinds of  $Y$ -algebras can be realized as a coset of the pair of  $\mathcal{W}^k(\mathfrak{sl}(m+n), f_{n,m})$  and the universal affine vertex algebra associated with  $\mathfrak{gl}(m)$  up to Heisenberg algebras. In this case, Creutzig-Linshaw [5] have proved the triality conjecture. This result is the generalization of the Feigin-Frenkel duality and the coset realization of principal  $W$ -algebra.

By Theorem 1.1, the images of  $\Phi_a$  are commutative with each other. For non-critical level, the completion of image of  $\Phi_1$  coincides with the subalgebra isomorphic to the universal enveloping algebra of a subalgebra which is expected to be isomorphic to the rectangular  $W$ -algebra associated with  $\mathfrak{gl}((n_1 - n_2)b_1)$  and a nilpotent element of type  $(n_1 - n_2)^{b_1}$ . Especially, in the case that  $b_1 = 1, 2$ , this subalgebra coincides with the rectangular  $W$ -algebra. By Theorem 1.1, for a non-critical level, we find that  $\Phi_a$  becomes a homomorphism from the affine Yangian to the universal enveloping algebra of the coset of the pair of  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  and this subalgebra. We expect that this result will be connected to obtain the generalization of the Gaiotto-Rapcak's triality.

## 2 $W$ -algebras of type $A$

We fix some notations for vertex algebras. For a vertex algebra  $V$ , we denote the generating field associated with  $v \in V$  by  $v(z) = \sum_{n \in \mathbb{Z}} v_{(n)} z^{-n-1}$ . We also denote the OPE of  $V$  by

$$u(z)v(w) \sim \sum_{s \geq 0} \frac{(u_{(s)}v)(w)}{(z-w)^{s+1}}$$

for all  $u, v \in V$ . We denote the vacuum vector (resp. the translation operator) by  $|0\rangle$  (resp.  $\partial$ ). We denote the universal affine vertex algebra associated with a finite dimensional Lie algebra  $\mathfrak{g}$  and its inner product  $\kappa$  by  $V^\kappa(\mathfrak{g})$ . By the PBW theorem, we can identify  $V^\kappa(\mathfrak{g})$  with  $U(t^{-1}\mathfrak{g}[t^{-1}])$ . In order to simplify the notation, here after, we denote the generating field  $(ut^{-1})(z)$  as  $u(z)$  for  $u \in \mathfrak{g}$ . By the definition of  $V^\kappa(\mathfrak{g})$ , the generating fields  $u(z)$  and  $v(z)$  satisfy the OPE

$$u(z)v(w) \sim \frac{[u, v](w)}{z-w} + \frac{\kappa(u, v)}{(z-w)^2} \quad (2.1)$$

for all  $u, v \in \mathfrak{g}$ .

We take a positive integer and its partition:

$$N = \sum_{i=1}^l q_i, \quad q_1 \geq q_2 \geq \cdots \geq q_l. \quad (2.2)$$

For  $\{q_i\}_{i=1}^l$  in (2.2), we set  $\{n_s\}_{s=1}^{v+1}$  and  $\{b_s\}_{s=1}^v$  as

$$q_{b_{s-1}} = n_{s-1} < q_{b_{s-1}+1} = q_{b_{s-1}+2} = \cdots = q_{b_s} = n_s < q_{b_s+1} = n_{s+1}, n_{v+1} = 0.$$

We also fix an inner product of  $\mathfrak{gl}(N)$  determined by

$$(E_{i,j}|E_{p,q}) = k\delta_{i,q}\delta_{p,j} + \delta_{i,j}\delta_{p,q}.$$

For  $1 \leq i \leq N$ , we set  $1 \leq \text{col}(i) \leq l$  and  $q_1 - q_{\text{col}(i)} < \text{row}(i) \leq q_1$  satisfying

$$\text{col}(i) = s \text{ if } \sum_{j=1}^{s-1} q_j < i \leq \sum_{j=1}^s q_j, \quad \text{row}(i) = i - \sum_{j=1}^{\text{col}(i)-1} q_j + q_1 - q_{\text{col}(i)}.$$

We take a nilpotent element  $f$  as

$$f = \sum_{1 \leq j \leq N} e_{\hat{j},j},$$

where the integer  $1 \leq \hat{j} \leq N$  is determined by

$$\text{col}(\hat{j}) = \text{col}(j) + 1, \quad \text{row}(\hat{j}) = \text{row}(j).$$

The  $W$ -algebra is a vertex algebra associated with the reductive Lie algebra  $\mathfrak{g}$  and a nilpotent element  $f$ . We denote the  $W$ -algebra associated with  $\mathfrak{gl}(N)$  and  $f$  by  $\mathcal{W}^k(\mathfrak{gl}(N), f)$ .

We set the inner product on  $\mathfrak{gl}(q_s)$  by

$$\kappa_s(E_{i,j}, E_{p,q}) = \delta_{j,p}\delta_{i,q}\alpha_s + \delta_{i,j}\delta_{p,q},$$

where  $\alpha_s = k + N - q_s$ . Then, by Corollary 5.2 in [10], the  $W$ -algebra  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  can be embedded into  $\bigotimes_{1 \leq s \leq l} V^{\kappa_s}(\mathfrak{gl}(q_s))$ . This embedding is called the Miura map.

**Theorem 2.3** (Theorem 3.6 in [27]). *We set  $\gamma_a = \sum_{u=a+1}^l \alpha_u$ . For positive integers  $q_1 - q_v < p, q \leq q_1 - q_{v+1}$ , the following elements of  $\bigotimes_{1 \leq s \leq l} V^{\kappa_s}(\mathfrak{gl}(q_s))$  are contained in  $\mathcal{W}^k(\mathfrak{gl}(N), f)$ :*

$$\begin{aligned} W_{p,q}^{(1)} &= \sum_{1 \leq r \leq l} e_{p,q}^{(r)}[-1], \\ W_{p,q}^{(2)} &= - \sum_{1 \leq r \leq l} \gamma_r e_{p,q}^{(r)}[-2] + \sum_{\substack{r_1 < r_2 \\ u > q_1 - q_v}} e_{u,q}^{(r_1)}[-1] e_{p,u}^{(r_2)}[-1] \\ &\quad - \sum_{\substack{r_1 \geq r_2 \\ q_1 - q_{r_2} < u \leq q_1 - q_v}} e_{u,q}^{(r_1)}[-1] e_{p,u}^{(r_2)}[-1], \end{aligned}$$

where we denote  $E_{i,j}t^{-u} \in U(t^{-1}\mathfrak{gl}(q_s)[t^{-1}]) = V^\kappa(\mathfrak{gl}(q_s))$  by  $e_{i+(q_1-q_s), j+(q_1-q_s)}^{(s)}[-u]$ .

Let us compute the OPEs of  $W_{i,j}^{(2)}$  and  $W_{p,q}^{(1)}$  for  $q_1 - q_v < i, j, p, q \leq q_{v+1}$ .

**Theorem 2.4.** *The following relations hold:*

$$\begin{aligned} (W_{p,q}^{(1)})_{(0)} W_{i,j}^{(2)} &= \delta_{i,q} W_{p,j}^{(2)} - \delta_{p,j} W_{i,q}^{(2)}, \\ (W_{p,q}^{(1)})_{(1)} W_{i,j}^{(2)} &= \delta_{p,j} \gamma_1 W_{i,q}^{(1)} - \delta_{p,q} q_v (b_v - 1) W_{i,j}^{(1)} + \sum_{1 \leq u \leq q_1 - q_v} \delta_{p,j} \delta_{i,q} W_{u,u}^{(1)}, \\ (W_{p,q}^{(1)})_{(2)} W_{i,j}^{(2)} &= (-2 \sum_{1 \leq r \leq l} \gamma_r \alpha_r \delta_{p,j} \delta_{i,q} - 2 \sum_{1 \leq r \leq l} \gamma_r \delta_{p,q} \delta_{i,j} + \sum_{1 \leq r \leq l} (q_v - q_r) \alpha_r) |0\rangle. \end{aligned}$$

We denote by  $\mathcal{W}^k(\mathfrak{gl}(ln), (l^n))$  the  $W$ -algebra  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  in the case that  $q_1 = q_2 = \dots = q_l = n$  and call a rectangular  $W$ -algebra of type  $A$ . Arakawa-Molev [1] gave strong generators of  $\mathcal{W}^k(\mathfrak{gl}(ln), (l^n))$  explicitly. In the appendix, we will give more elements of  $\mathcal{W}^k(\mathfrak{gl}(N), f)$ :

$$\{\widetilde{W}_{i,j}^{(r)} \mid 1 \leq i, j \leq n_1 - n_2, 1 \leq r \leq b_1\},$$

which corresponds to strong generators of Arakawa-Molev in the case that  $q_1 = q_2 = \dots = q_l = n$ . In the case that  $b_1 = 2$ , we have known the relationship between the elements  $\widetilde{W}_{i,j}^{(r)}$  and  $W_{i,j}^{(r)}$ .

**Theorem 2.5.** *There exists an embedding*

$$\iota: \mathcal{W}^{k+n_2+\sum_{s=3}^l q_s}(\mathfrak{gl}(2(n_1 - n_2)), (2^{n_1 - n_2})) \rightarrow \mathcal{W}^k(\mathfrak{gl}(N), f)$$

defined by

$$\widetilde{W}_{i,j}^{(1)} \mapsto W_{i,j}^{(1)}, \quad \widetilde{W}_{i,j}^{(2)} + \gamma_1 \partial W_{i,j}^{(1)}.$$

*Proof.* By the form of  $W_{i,j}^{(r)}$  and  $W_{i,j}^{(2)}$ , we find that the OPEs of  $\{W_{i,j}^{(r)} \mid r = 1, 2, 1 \leq i, j \leq n_1 - n_2\}$  in  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  is the same as those of

$$\{\widetilde{W}_{i,j}^{(1)}, \widetilde{W}_{i,j}^{(2)} - \gamma_1 \partial \widetilde{W}_{i,j}^{(1)} \mid r = 1, 2, 1 \leq i, j \leq n_1 - n_2\}$$

in  $\mathcal{W}^{k+\sum_{s=3}^l q_s}(\mathfrak{gl}(2n_1), (2^{n_1}))$ . In [22], the OPEs of rectangular  $W$ -algebras have been computed (see also [26]). Then, we find an embedding

$$\iota_1: \mathcal{W}^{k+n_2+\sum_{s=3}^l q_s}(\mathfrak{gl}(2(n_1 - n_2)), (2^{n_1 - n_2})) \rightarrow \mathcal{W}^{k+\sum_{s=3}^l q_s}(\mathfrak{gl}(2n_1), (2^{n_1}))$$

given by  $\widetilde{W}_{i,j}^{(r)} \mapsto \widetilde{W}_{i,j}^{(r)}$  for  $1 \leq i, j \leq n_1 - n_2$ . □

Next, let us recall the universal enveloping algebra of a vertex algebra. For any vertex algebra  $V$ , let  $L(V)$  be the Borcherds Lie algebra, that is,

$$L(V) = V \otimes \mathbb{C}[t, t^{-1}] / \text{Im}(\partial \otimes \text{id} + \text{id} \otimes \frac{d}{dt}), \quad (2.6)$$

where the commutation relation is given by

$$[ut^a, vt^b] = \sum_{r \geq 0} \binom{a}{r} (u_{(r)} v) t^{a+b-r}$$

for all  $u, v \in V$  and  $a, b \in \mathbb{Z}$ .

**Definition 2.7** (Section 6 in [20]). We set  $\mathcal{U}(V)$  as the quotient algebra of the standard degreewise completion of the universal enveloping algebra of  $L(V)$  by the completion of the two-sided ideal generated by

$$(u_{(a)} v) t^b - \sum_{i \geq 0} \binom{a}{i} (-1)^i (ut^{a-i} vt^{b+i} - (-1)^a vt^{a+b-i} ut^i), \quad (2.8)$$

$$|0\rangle t^{-1} - 1. \quad (2.9)$$

We call  $\mathcal{U}(V)$  the universal enveloping algebra of  $V$ .

Induced by the Miura map  $\mu$ , we obtain the embedding

$$\tilde{\mu}: \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f)) \rightarrow \widehat{\bigoplus}_{1 \leq s \leq l} U(\widehat{\mathfrak{gl}}(q_s)),$$

where  $\widehat{\bigoplus}_{1 \leq s \leq l} U(\widehat{\mathfrak{gl}}(q_s))$  is the standard degreewise completion of  $\bigotimes_{1 \leq s \leq l} U(\widehat{\mathfrak{gl}}(q_s))$ .

### 3 The affine Yangian and its coproduct

The affine Yangian of type  $A$  was first introduced by Guay ([11] and [12]). The affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  is a 2-parameter Yangian and is the deformation of the universal enveloping algebra of the central extension of  $\mathfrak{sl}(n)[u^{\pm 1}, v]$ . The affine Yangian has several presentations. In this section, we define the affine Yangian of type  $A$  by using the minimalistic presentation given in Guay-Nakajima-Wendlandt [13].

Hereafter, we sometimes identify  $\{0, 1, 2, \dots, n-1\}$  with  $\mathbb{Z}/n\mathbb{Z}$ . We also set  $\{X, Y\} = XY + YX$  and

$$a_{i,j} = \begin{cases} 2 & \text{if } i = j, \\ -1 & \text{if } j = i \pm 1, \\ 0 & \text{otherwise} \end{cases}$$

for  $i \in \mathbb{Z}/n\mathbb{Z}$ .

**Definition 3.1.** Suppose that  $n \geq 3$ . The affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  is the associative algebra generated by  $X_{i,r}^+, X_{i,r}^-, H_{i,r}$  ( $i \in \{0, 1, \dots, n-1\}$ ,  $r = 0, 1$ ) subject to the following defining relations:

$$[H_{i,r}, H_{j,s}] = 0, \quad (3.2)$$

$$[X_{i,0}^+, X_{j,0}^-] = \delta_{i,j} H_{i,0}, \quad (3.3)$$

$$[X_{i,1}^+, X_{j,0}^-] = \delta_{i,j} H_{i,1} = [X_{i,0}^+, X_{j,1}^-], \quad (3.4)$$

$$[H_{i,0}, X_{j,r}^\pm] = \pm a_{i,j} X_{j,r}^\pm, \quad (3.5)$$

$$[\tilde{H}_{i,1}, X_{j,0}^\pm] = \pm a_{i,j} (X_{j,1}^\pm), \text{ if } (i, j) \neq (0, n-1), (n-1, 0), \quad (3.6)$$

$$[\tilde{H}_{0,1}, X_{n-1,0}^\pm] = \mp \left( X_{n-1,1}^\pm + (\varepsilon + \frac{n}{2}\hbar)X_{n-1,0}^\pm \right), \quad (3.7)$$

$$[\tilde{H}_{n-1,1}, X_{0,0}^\pm] = \mp \left( X_{0,1}^\pm - (\varepsilon + \frac{n}{2}\hbar)X_{0,0}^\pm \right), \quad (3.8)$$

$$[X_{i,1}^\pm, X_{j,0}^\pm] - [X_{i,0}^\pm, X_{j,1}^\pm] = \pm a_{ij} \frac{\hbar}{2} \{X_{i,0}^\pm, X_{j,0}^\pm\} \text{ if } (i,j) \neq (0,n-1), (n-1,0), \quad (3.9)$$

$$[X_{0,1}^\pm, X_{n-1,0}^\pm] - [X_{0,0}^\pm, X_{n-1,1}^\pm] = \mp \frac{\hbar}{2} \{X_{0,0}^\pm, X_{n-1,0}^\pm\} + (\varepsilon + \frac{n}{2}\hbar)[X_{0,0}^\pm, X_{n-1,0}^\pm], \quad (3.10)$$

$$(\text{ad } X_{i,0}^\pm)^{1+|a_{i,j}|}(X_{j,0}^\pm) = 0 \text{ if } i \neq j, \quad (3.11)$$

where we set  $\tilde{H}_{i,1} = H_{i,1} - \frac{\hbar}{2}H_{i,0}^2$ .

Let us denote Chevalley generators of  $\widehat{\mathfrak{sl}}(n)$  by  $\{h_i, x_i^\pm \mid 0 \leq i \leq n-1\}$ . By the defining relations (3.2)-(3.11), there exists a homomorphism from the universal enveloping algebra of  $\widehat{\mathfrak{sl}}(n)$  to the affine Yangian  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  given by  $h_i \mapsto H_{i,0}$  and  $x_i^\pm \mapsto X_{i,0}^\pm$ . We denote the image of  $x \in U(\widehat{\mathfrak{sl}}(n))$  via this homomorphism by  $x \in Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ .

Let us set the degree of  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  by

$$\deg(H_{i,r}) = 0, \quad \deg(X_{i,r}^\pm) = \begin{cases} \pm 1 & \text{if } i = 0, \\ 0 & \text{if } i \neq 0. \end{cases}$$

This degree is compatible with the natural degree on  $\widehat{\mathfrak{sl}}(n)$ . We denote the standard degreewise completion of  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  by  $\widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ . By using the minimalistic presentation of the Yangian, Guay-Nakajima-Wendlandt [13] gave a coproduct for the Yangian associated with a Kac-Moody Lie algebra of the affine type.

**Theorem 3.12** (Theorem 5.2 in [13]). *There exists an algebra homomorphism*

$$\Delta^n: Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \widehat{\otimes} Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$$

determined by

$$\begin{aligned} \Delta^n(X_{j,0}^\pm) &= X_{j,0}^\pm \otimes 1 + 1 \otimes X_{j,0}^\pm \text{ for } 0 \leq j \leq n-1, \\ \Delta^n(\tilde{H}_{i,1}) &= \tilde{H}_{i,1} \otimes 1 + 1 \otimes \tilde{H}_{i,1} + A_i \text{ for } 1 \leq i \leq n-1, \end{aligned}$$

where  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \widehat{\otimes} Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  is the standard degreewise completion of  $\otimes^2 Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  and

$$\begin{aligned} A_i &= -\hbar(E_{i,i} \otimes E_{i+1,i+1} + E_{i+1,i+1} \otimes E_{i,i}) \\ &\quad + \hbar \sum_{s \geq 0}^i \sum_{u=1}^i (-E_{u,i} t^{-s-1} \otimes E_{i,u} t^{s+1} + E_{i,u} t^{-s} \otimes E_{u,i} t^s) \\ &\quad + \hbar \sum_{s \geq 0}^n \sum_{u=i+1}^n (-E_{u,i} t^{-s} \otimes E_{i,u} t^s + E_{i,u} t^{-s-1} \otimes E_{u,i} t^{s+1}) \\ &\quad - \hbar \sum_{s \geq 0}^i \sum_{u=1}^i (-E_{u,i+1} t^{-s-1} \otimes E_{i+1,u} t^{s+1} + E_{i+1,u} t^{-s} \otimes E_{u,i+1} t^s) \\ &\quad - \hbar \sum_{s \geq 0}^n \sum_{u=i+1}^n (-E_{u,i+1} t^{-s} \otimes E_{i+1,u} t^s + E_{i+1,u} t^{-s-1} \otimes E_{u,i+1} t^{s+1}). \end{aligned}$$

The homomorphism  $\Delta^n$  is said to be the coproduct for the affine Yangian since  $\Delta^n$  satisfies the coassociativity.

*Remark 3.13.* By (3.3), (3.4), (3.6) and (3.8), we find that the affine Yangian  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  is generated by  $X_{i,0}^\pm$  for  $0 \leq i \leq n-1$  and  $H_{j,1}$  for  $1 \leq j \leq n-1$ . Thus, the homomorphism is defined if the image of these elements are determined. In this article, we will define homomorphisms by giving the image of these elements.

## 4 Evaluation map for the affine Yangian

Guay [12] gave the evaluation map for the affine Yangian and Kodera [17] showed the surjectivity of the evaluation map. The evaluation map for the affine Yangian is a non-trivial homomorphism from the affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  to the completion of the universal enveloping algebra of the affinization of  $\mathfrak{gl}(n)$ . We set a Lie algebra

$$\widehat{\mathfrak{gl}}(n) = \mathfrak{gl}(n) \otimes \mathbb{C}[z^{\pm 1}] \oplus \mathbb{C}\tilde{c} \oplus \mathbb{C}z$$

whose commutator relations are given by

$$z \text{ and } \tilde{c} \text{ are central elements of } \widehat{\mathfrak{gl}}(n),$$

$$[E_{i,j} \otimes t^u, E_{p,q} \otimes t^v] = (\delta_{j,p}E_{i,q} - \delta_{i,q}E_{p,j}) \otimes t^{u+v} + \delta_{u+v,0}u(\delta_{i,q}\delta_{j,p}\tilde{c} + \delta_{i,j}\delta_{p,q}z).$$

We take the grading of  $U(\widehat{\mathfrak{gl}}(n))/U(\widehat{\mathfrak{gl}}(n))(z-1)$  as  $\deg(Xt^s) = s$  and  $\deg(\tilde{c}) = 0$ . We denote the degreewise completion of  $U(\widehat{\mathfrak{gl}}(n))/U(\widehat{\mathfrak{gl}}(n))(z-1)$  by  $\mathcal{U}(\widehat{\mathfrak{gl}}(n))$ .

**Theorem 4.1** (Theorem 3.8 in [18] and Theorem 4.18 in [17]). 1). Suppose that  $\tilde{c} = \frac{\varepsilon}{\hbar}$ . For  $a \in \mathbb{C}$ , there exists an algebra homomorphism

$$\text{ev}_{\hbar, \varepsilon}^{n,a}: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \mathcal{U}(\widehat{\mathfrak{gl}}(n))$$

uniquely determined by

$$\text{ev}_{\hbar, \varepsilon}^{n,a}(X_{i,0}^+) = \begin{cases} E_{n,1}t & \text{if } i = 0, \\ E_{i,i+1} & \text{if } 1 \leq i \leq n-1, \end{cases} \quad \text{ev}_{\hbar, \varepsilon}^{n,a}(X_{i,0}^-) = \begin{cases} E_{1,n}t^{-1} & \text{if } i = 0, \\ E_{i+1,i} & \text{if } 1 \leq i \leq n-1, \end{cases}$$

and

$$\begin{aligned} \text{ev}_{\hbar, \varepsilon}^{n,a}(H_{i,1}) &= (a - \frac{i}{2}\hbar) \text{ev}_{\hbar, \varepsilon}^{n,a}(H_{i,0}) - \hbar E_{i,i}E_{i+1,i+1} \\ &\quad + \hbar \sum_{s \geq 0} \sum_{k=1}^i E_{i,k}t^{-s}E_{k,i}t^s + \hbar \sum_{s \geq 0} \sum_{k=i+1}^n E_{i,k}t^{-s-1}E_{k,i}t^{s+1} \\ &\quad - \hbar \sum_{s \geq 0} \sum_{k=1}^i E_{i+1,k}t^{-s}E_{k,i+1}t^s - \hbar \sum_{s \geq 0} \sum_{k=i+1}^n E_{i+1,k}t^{-s-1}E_{k,i+1}t^{s+1} \end{aligned}$$

for  $i \neq 0$ .

2). In the case that  $\varepsilon \neq 0$ , the image of  $\text{ev}_{\hbar, \varepsilon}^{n,a}$  is dense in the target.

## 5 Tensor algebras of the affine Yangians

In this section, we will construct a homomorphism from tensor of affine Yangians to another tensor of affine Yangians.

**Theorem 5.1** (Theorem 3.1 in [26] and Theorem 3.1 in [28]). 1). For  $m \geq 1$  and  $n \geq 3$ , there exists a homomorphism

$$\Psi_1^{n,m+n}: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \widetilde{Y}_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(m+n))$$

given by

$$\Psi_1^{n,m+n}(X_{i,0}^+) = \begin{cases} E_{n,1}t & \text{if } i = 0, \\ E_{i,i+1} & \text{if } i \neq 0, \end{cases} \quad \Psi_1^{n,m+n}(X_{i,0}^-) = \begin{cases} E_{1,n}t^{-1} & \text{if } i = 0, \\ E_{i+1,i} & \text{if } i \neq 0, \end{cases}$$

and

$$\Psi_1^{n,m+n}(H_{i,1}) = H_{i,1} - \hbar \sum_{s \geq 0} \sum_{k=n+1}^{m+n} E_{i,k} t^{-s-1} E_{k,i} t^{s+1} + \hbar \sum_{s \geq 0} \sum_{k=n+1}^{m+n} E_{i+1,k} t^{-s-1} E_{k,i+1} t^{s+1}$$

for  $i \neq 0$ .

2). For  $m \geq 3$  and  $n \geq 1$ , there exists a homomorphism

$$\Psi_2^{m,m+n}: Y_{\hbar,\varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow \widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(m+n))$$

determined by

$$\Psi_2^{m,m+n}(X_{i,0}^+) = \begin{cases} E_{m+n,n+1}t & \text{if } i=0, \\ E_{n+i,n+i+1} & \text{if } i \neq 0, \end{cases} \quad \Psi_2^{m,m+n}(X_{i,0}^-) = \begin{cases} E_{n+1,m+n}t^{-1} & \text{if } i=0, \\ E_{n+i+1,n+i} & \text{if } i \neq 0, \end{cases}$$

and

$$\begin{aligned} \Psi_2^{m,m+n}(H_{i,1}) &= H_{i+n,1} + \hbar \sum_{s \geq 0} \sum_{k=1}^n E_{k,n+i} t^{-s-1} E_{n+i,k} t^{s+1} \\ &\quad - \hbar \sum_{s \geq 0} \sum_{k=1}^n E_{k,n+i+1} t^{-s-1} E_{n+i+1,k} t^{s+1} \end{aligned}$$

for  $i \neq 0$ .

3). For  $m, n \geq 3$ , two homomorphisms  $\Psi_1$  and  $\Psi_2$  induce the homomorphism

$$\Psi_1^{n,m+n} \otimes \Psi_2^{m,m+n}: Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \otimes Y_{\hbar,\varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow \widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(m+n)).$$

Suppose that  $q_s - q_{s+1} = 0$  or  $\geq 3$ . Let us define the following algebra

$$Y_{\hbar,\varepsilon}^{n_1,n_2,\dots,n_v} = \bigotimes_{1 \leq s \leq v} Y_{\hbar,\varepsilon_s}(\widehat{\mathfrak{sl}}(n_s - n_{s+1})),$$

where we set  $\varepsilon_s = \varepsilon - (n_s - n_v)\hbar$ . We denote  $x \in Y_{\hbar,\varepsilon_s}(\widehat{\mathfrak{sl}}(n_s - n_{s+1})) \subset Y^{n_1,n_2,\dots,n_v}$  by  $x^{(s)}$ . Then, we can construct two homomorphisms  $\Delta_A^{n_1,\dots,n_v}$  and  $\Delta_B^{n_1,\dots,n_v}$ :

$$\Delta_A^{n_1,\dots,n_v}: Y^{n_1,n_2,\dots,n_v} \rightarrow Y^{n_1,n_2,\dots,n_v} \widehat{\otimes} Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n_v))$$

determined by

$$\Delta_A^{n_1,\dots,n_v}(x^{(r)}) = \begin{cases} x^{(r)} \otimes 1 & \text{if } r \neq v, \\ \Delta(x)^{(v)} & \text{if } r = v \end{cases}$$

and

$$\Delta_B^{n_1,\dots,n_v}: Y^{n_1,n_2,\dots,n_l} \rightarrow Y^{n_1,n_2,\dots,n_{v-2},n_{v-1}} \widehat{\otimes} Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n_v))$$

determined by

$$\Delta_B^{n_1,\dots,n_v}(x^{(r)}) = \begin{cases} x^{(r)} \otimes 1 & \text{if } r \neq v-2, \\ (\Psi_1^{n_{v-1}-n_v,n_{v-1}}(x))^{(v-1)} \otimes 1 & \text{if } r = v-1, \\ 1^{\otimes(v-2)} \otimes ((\Psi_2^{n_v,n_{v-1}} \otimes 1) \circ \Delta)(x) & \text{if } r = v, \end{cases}$$

where  $S \widehat{\otimes} U$  means the standard degreewise completion of the tensor product  $S \otimes U$ .

We set a homomorphism

$$\Delta_l = \prod_{s=1}^{l-1} (\Delta_s \otimes \text{id}^{l-s-1}) : Y_{\hbar, \varepsilon}^{n_1, n_2, \dots, n_v} \rightarrow \widehat{\bigotimes}_{1 \leq s \leq l} Y_{\hbar, \varepsilon - (n_s - n_v)\hbar}(\widehat{\mathfrak{sl}}(q_s)),$$

where  $\widehat{\bigotimes}_{1 \leq s \leq l} Y_{\hbar, \varepsilon - (n_s - n_v)\hbar}(\widehat{\mathfrak{sl}}(q_s))$  is the standard degreewise completion of the tensor of affine Yangians and

$$\Delta_s = \begin{cases} \Delta_B^{n_1, \dots, n_a} & \text{if } n_{a-1} = q_{s-1} > q_s = n_a, \\ \Delta_A^{n_1, \dots, n_a} & \text{if } q_{s-1} = q_s = n_a. \end{cases}$$

By the definition of  $\Psi^{n,m+n}$  and  $\Delta^n$ , we obtain

$$\Delta_l(H_{i,1})^{(a)} = \sum_{1 \leq s \leq b_a} H_{i+q_s-n_a}^{(s)} + B_i + C_i + D_i,$$

where we set  $H_{i+q_s-n_a}^{(s)} = 1^{\otimes s-1} \otimes H_{i+q_s-n_a} \otimes 1^{\otimes l-s}$ ,  $E_{i,j}^{(s)} t^s = 1^{\otimes s-1} \otimes E_{i,j} \otimes 1^{\otimes l-s}$  and

$$\begin{aligned} B_i = & -\hbar \sum_{1 \leq r_1 < r_2 \leq b_a} E_{q_{r_1}-n_a+i, q_{r_1}-n_a+i}^{(r_1)} E_{q_{r_2}-n_a+i+1, q_{r_2}-n_a+i+1}^{(r_2)} \\ & - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} E_{q_{r_1}-n_a+i+1, q_{r_1}-n_a+i+1}^{(r_1)} E_{q_{r_2}-n_a+i, q_{r_2}-n_a+i}^{(r_2)} \\ & - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-n_a+i} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-n_a+i}^{(r_1)} t^{-s-1} E_{q_{r_2}-n_a+i, u}^{(r_2)} t^{s+1} \\ & + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-n_a+i} E_{q_{r_1}-n_a+i, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s} E_{u, q_{r_2}-n_a+i}^{(r_2)} t^s \\ & - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+i+1}^{q_{r_2}} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-n_a+i}^{(r_1)} t^{-s} E_{q_{r_2}-n_a+i, u}^{(r_2)} t^s \\ & + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+i+1}^{q_{r_2}} E_{q_{r_1}-n_a+i, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s-1} E_{u, q_{r_2}-n_a+i}^{(r_2)} t^{s+1} \\ & + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-n_a+i} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-n_a+i+1}^{(r_1)} t^{-s-1} E_{q_{r_2}-n_a+i+1, u}^{(r_2)} t^{s+1} \\ & - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-n_a+i} E_{q_{r_1}-n_a+i+1, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s} E_{u, q_{r_2}-n_a+i+1}^{(r_2)} t^s \\ & + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+i+1}^{q_{r_2}} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-q_{l+i+1}}^{(r_1)} t^{-s} E_{q_{r_2}-n_a+i+1, u}^{(r_2)} t^s \\ & - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+i+1}^{q_{r_2}} E_{q_{r_1}-n_a+i+1, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s-1} E_{u, q_{r_2}-n_a+i+1}^{(r_2)} t^{s+1}, \\ C_i = & \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=1}^{q_r-n_a} E_{u, q_r-n_a+i}^{(r)} t^{-s-1} E_{q_r-n_a+i, u}^{(r)} t^{s+1} \\ & - \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=1}^{q_r-n_a} E_{u, q_r-n_a+i+1}^{(r)} t^{-s-1} E_{q_r-n_a+i+1, u}^{(r)} t^{s+1} \end{aligned}$$

$$\begin{aligned}
& + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-n_a} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-n_a+i}^{(r_1)} t^{-s-1} E_{q_{r_2}-n_a+i, u}^{(r_2)} t^{s+1} \\
& - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-n_a} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-n_a+i+1}^{(r_1)} t^{-s-1} E_{q_{r_2}-n_a+i+1, u}^{(r_2)} t^{s+1} \\
& + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-n_a} E_{u, q_{r_2}-n_a+i}^{(r_2)} t^{-s-1} E_{q_{r_1}-n_a+i, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{s+1} \\
& - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-n_a} E_{u, q_{r_2}-n_a+i+1}^{(r_2)} t^{-s-1} E_{q_{r_1}-n_a+i+1, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{s+1}, \\
D_i = & -\hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_r-n_{a+1}+1}^{q_r} E_{q_r-n_a+i, u}^{(r)} t^{-s-1} E_{u, q_r-n_a+i}^{(r)} t^{s+1} \\
& + \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_r-n_{a+1}+1}^{q_r} E_{q_r-n_a+i+1, u}^{(r)} t^{-s-1} E_{u, q_r-n_a+i+1}^{(r)} t^{s+1} \\
& - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_{a+1}+1}^{q_{r_2}} E_{q_{r_1}-n_a+i, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s-1} E_{u, q_{r_2}-n_a+i}^{(r_2)} t^{s+1} \\
& + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_{a+1}+1}^{q_{r_2}} E_{q_{r_1}-n_a+i+1, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s-1} E_{u, q_{r_2}-n_a+i+1}^{(r_2)} t^{s+1} \\
& - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_{a+1}+1}^{q_{r_2}} E_{q_{r_2}-n_a+i, u}^{(r_2)} t^{-s-1} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-n_a+i}^{(r_1)} t^{s+1} \\
& + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_{a+1}+1}^{q_{r_2}} E_{q_{r_2}-n_a+i+1, u}^{(r_2)} t^{-s-1} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-n_a+i+1}^{(r_1)} t^{s+1}.
\end{aligned}$$

We note that  $B_i$ ,  $C_i$  and  $D_i$  come from the coproduct for the affine Yangian, the homomorphism  $\Psi_2^{m,m+n}$  and the homomorphism  $\Psi_1^{n,m+n}$  respectively.

## 6 The tensor algebra of affine Yangians and non-rectangular $W$ -algebras

In this section, we construct a homomorphism from the tensor algebra of affine Yangians to the universal enveloping algebra of the non-rectangular  $W$ -algebra.

**Theorem 6.1.** *Assume that  $\varepsilon = \hbar(k + N - q_l)$ . Then, there exists an algebra homomorphism*

$$\Phi: Y_{\hbar, \varepsilon}^{n_1, \dots, n_v} \rightarrow \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f))$$

determined by

$$\begin{aligned}
\Phi((X_{i,0}^+)^{(a)}) &= \begin{cases} W_{n_1-n_{a+1}, n_1-n_a+1}^{(1)} t & \text{if } i = 0, \\ W_{n_1-n_a+i, n_1-n_a+i+1}^{(1)} & \text{if } 1 \leq i \leq n_a - n_{a+1} - 1, \end{cases} \\
\Phi((X_{i,0}^-)^{(a)}) &= \begin{cases} W_{n_1-n_a+1, n_1-n_a+1}^{(1)} t^{-1} & \text{if } i = 0, \\ W_{n_1-n_a+i+1, n_1-n_a+i}^{(1)} & \text{if } 1 \leq i \leq n_a - n_{a+1} - 1, \end{cases}
\end{aligned}$$

and

$$\Phi((H_{i,1})^{(a)}) = -\hbar(W_{n_1-n_a+i, n_1-n_a+i}^{(2)} t - W_{n_1-n_a+i+1, n_1-n_a+i+1}^{(2)} t)$$

$$\begin{aligned}
& - \frac{i}{2} \hbar (W_{n_1-n_a+i, n_1-n_a+i}^{(1)} - W_{n_1-n_a+i+1, n_1-n_a+i+1}^{(1)}) \\
& + \hbar W_{n_1-n_a+i, n_1-n_a+i}^{(1)} W_{n_1-n_a+i+1, n_1-n_a+i+1}^{(1)} \\
& + \hbar \sum_{s \geq 0} \sum_{u=1}^i W_{n_1-n_a+i, n_1-n_a+u}^{(1)} t^{-s} W_{n_1-n_a+u, n_1-n_a+i}^{(1)} t^s \\
& + \hbar \sum_{s \geq 0} \sum_{u=i+1}^n W_{n_1-n_a+i, n_1-n_a+u}^{(1)} t^{-s-1} W_{n_1-n_a+u, n_1-n_a+i}^{(1)} t^{s+1} \\
& - \hbar \sum_{s \geq 0} \sum_{u=1}^i W_{n_1-n_a+i+1, n_1-n_a+u}^{(1)} t^{-s} W_{n_1-n_a+u, n_1-n_a+i+1}^{(1)} t^s \\
& - \hbar \sum_{s \geq 0} \sum_{u=i+1}^n W_{n_1-n_a+i+1, n_1-n_a+u}^{(1)} t^{-s-1} W_{n_1-n_a+u, n_1-n_a+i+1}^{(1)} t^{s+1}
\end{aligned}$$

for  $1 \leq i \leq n_a - n_{a+1} - 1$ .

Similarly to the main theorem in [19] and [25], it is enough to show the following theorem.

**Theorem 6.2.** *We obtain the following relation:*

$$\bigotimes_{1 \leq s \leq l} \text{ev}_{\hbar, \varepsilon - (q_s - q_l) \hbar}^{q_s, \gamma_s \hbar} \circ \Delta_l = \tilde{\mu} \circ \Phi.$$

*Proof.* It is enough to show that the relation holds for  $X_{j,0}^\pm$  ( $0 \leq j \leq n_a - n_{a+1} - 1$ ) and  $H_{i,1}$  ( $1 \leq i \leq n_a - n_{a+1} - 1$ ). It is trivial that the relation holds for  $X_{j,0}^\pm$ . Hence, we only show the latter one, that is,

$$\bigotimes_{1 \leq s \leq l} \text{ev}_{\hbar, \varepsilon - (q_s - q_l) \hbar}^{q_s, \gamma_s \hbar} \circ \Delta_l (H_{i,1}) = \tilde{\mu} \circ \Phi (H_{i,1}) \quad (6.3)$$

We denote  $E_{i,j}^{(r)} t^a \in \widehat{\bigotimes}_{1 \leq s \leq l} U(\widehat{\mathfrak{gl}}(q_s))$  by  $e_{q_1-q_s+i, q_1-q_s+j}^{(r)} t^a$ .

By the definition of the evaluation map and the last of Section 5, we find that the left hand side of (6.3) is equal to the sum of the following four terms:

$$\begin{aligned}
& \bigotimes_{1 \leq s \leq b_a} \text{ev}_{\hbar, \varepsilon - (q_s - q_l) \hbar}^{q_s, \gamma_s} (\bigotimes_{1 \leq s \leq l} H_{i+q_s-n_a}^{(s)}) \\
& = \sum_{r=1}^{b_a} \left( \gamma_r - \frac{i}{2} \hbar \right) (e_{q_1-n_a+i, q_1-n_a+i}^{(r)} - e_{q_1-n_a+i+1, q_1-n_a+i+1}^{(r)}) \\
& - \sum_{r=1}^{b_a} \hbar e_{q_1-n_a+i, q_1-n_a+i}^{(r)} e_{q_1-n_a+i+1, q_1-n_a+i+1}^{(r)} \\
& + \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_1-q_r+1}^{q_1-n_a+i} e_{q_1-n_a+i, u}^{(r)} t^{-s} e_{u, q_1-n_a+i}^{(r)} t^s \\
& + \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_1-n_a+i+1}^{q_1-n_a+i} e_{q_1-n_a+i, u}^{(r)} t^{-s-1} e_{u, q_1-n_a+i+1}^{(r)} t^{s+1} \\
& - \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_1-q_r+1}^{q_1-n_a+i} e_{q_1-n_a+i+1, u}^{(r)} t^{-s} e_{u, q_1-n_a+i+1}^{(r)} t^s \\
& - \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_1-n_a+i+1}^{q_1-n_a+i} e_{q_1-n_a+i+1, u}^{(r)} t^{-s-1} e_{u, q_1-n_a+i+1}^{(r)} t^{s+1},
\end{aligned} \quad (6.4)$$

$$\begin{aligned}
& \bigotimes_{1 \leq s \leq b_a} \text{ev}_{\hbar, \varepsilon - (q_s - q_l) \hbar}^{q_s, \gamma_s}(B_i) \\
&= -\hbar \sum_{1 \leq r_1 < r_2 \leq b_a} e_{q_1 - n_a + i, q_1 - n_a + i}^{(r_1)} e_{q_1 - n_a + i + 1, q_1 - n_a + i + 1}^{(r_2)} \\
&\quad - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} e_{q_1 - n_a + i + 1, q_1 - n_a + i + 1}^{(r_1)} e_{q_1 - n_a + i, q_1 - n_a + i}^{(r_2)} \\
&\quad - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2} - n_a + i} e_{q_1 - q_{r_2} + u, q_1 - n_a + i}^{(r_1)} t^{-s-1} E_{q_1 - q_{r_2} + i, q_1 - q_{r_2} + u}^{(r_2)} t^{s+1} \\
&\quad + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2} - n_a + i} e_{q_1 - n_a + i, q_1 - q_{r_2} + u}^{(r_1)} t^{-s} e_{q_1 - q_{r_2} + u, q_1 - n_a + i}^{(r_2)} t^s \\
&\quad - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2} - n_a + i + 1}^{q_{r_2}} e_{q_1 - q_{r_2} + u, q_1 - n_a + i}^{(r_1)} t^{-s} e_{q_1 - n_a + i, q_1 - q_{r_2} + u}^{(r_2)} t^s \\
&\quad + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2} - n_a + i + 1}^{q_{r_2}} e_{q_1 - n_a + i, q_1 - q_{r_2} + u}^{(r_1)} t^{-s-1} e_{q_1 - q_{r_2} + u, q_1 - n_a + i}^{(r_2)} t^{s+1} \\
&\quad + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2} - n_a + i} e_{q_1 - q_{r_2} + u, q_1 - n_a + i + 1}^{(r_1)} t^{-s-1} E_{q_1 - n_a + i + 1, q_1 - q_{r_2} + u}^{(r_2)} t^{s+1} \\
&\quad - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2} - n_a + i} e_{q_1 - n_a + i + 1, q_1 - q_{r_2} + u}^{(r_1)} t^{-s} e_{q_1 - q_{r_2} + u, q_1 - n_a + i + 1}^{(r_2)} t^s \\
&\quad + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2} - n_a + i + 1}^{q_{r_2}} e_{q_1 - q_{r_2} + u, q_1 - n_a + i + 1}^{(r_1)} t^{-s} e_{q_1 - n_a + i + 1, q_1 - q_{r_2} + u}^{(r_2)} t^s \\
&\quad - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2} - n_a + i + 1}^{q_{r_2}} e_{q_1 - n_a + i + 1, q_1 - q_{r_2} + u}^{(r_1)} t^{-s-1} e_{q_1 - q_{r_2} + u, q_1 - n_a + i + 1}^{(r_2)} t^{s+1}, \quad (6.5) \\
& \bigotimes_{1 \leq s \leq b_a} \text{ev}_{\hbar, \varepsilon - (q_s - q_l) \hbar}^{q_s, \gamma_s}(C_i) \\
&= \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=1}^{q_r - n_a} e_{q_1 - q_r + u, q_1 - n_a + i}^{(r)} t^{-s-1} e_{q_1 - q_r + i, q_1 - n_a + u}^{(r)} t^{s+1} \\
&\quad - \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=1}^{q_r - n_a} e_{q_1 - q_r + u, q_1 - q_r + i + 1}^{(r)} t^{-s-1} e_{q_1 - q_r + i + 1, q_1 - q_r + u}^{(r)} t^{s+1} \\
&\quad + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2} - n_a} e_{q_1 - q_{r_2} + u, q_1 - n_a + i}^{(r_1)} t^{-s-1} e_{q_1 - n_a + i, q_1 - q_{r_2} + u}^{(r_2)} t^{s+1} \\
&\quad - \hbar \sum_{s \geq 0} \sum_{u=1}^{q_{r_2} - n_a} e_{q_1 - q_{r_2} + u, q_1 - n_a + i + 1}^{(r_1)} t^{-s-1} e_{q_1 - n_a + i + 1, q_1 - q_{r_2} + u}^{(r_2)} t^{s+1} \\
&\quad + \hbar \sum_{s \geq 0} \sum_{u=1}^{q_{r_2} - n_a} e_{q_1 - q_{r_2} + u, q_1 - n_a + i}^{(r_2)} t^{-s-1} e_{q_1 - n_a + i, q_1 - q_{r_2} + u}^{(r_1)} t^{s+1} \\
&\quad - \hbar \sum_{s \geq 0} \sum_{u=1}^{q_{r_2} - n_a} e_{q_1 - q_{r_2} + u, q_1 - n_a + i + 1}^{(r_2)} t^{-s-1} e_{q_1 - n_a + i + 1, q_1 - q_{r_2} + u}^{(r_1)} t^{s+1}, \quad (6.6)
\end{aligned}$$

$$\begin{aligned}
& \bigotimes_{1 \leq s \leq l} \text{ev}_{\hbar, \varepsilon - (q_s - q_l) \hbar}^{q_s, \gamma_s}(D_i) \\
&= -\hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_r - n_a + 1 + 1}^{q_r} e_{q_1 - n_a + i, q_1 - q_r + u}^{(r)} t^{-s-1} e_{q_1 - q_r + u, q_1 - n_a + i}^{(r)} t^{s+1} \\
&\quad + \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_r - n_a + 1 + 1}^{q_r} e_{q_1 - n_a + i + 1, q_1 - q_r + u}^{(r)} t^{-s-1} e_{q_1 - q_r + u, q_1 - n_a + i + 1}^{(r)} t^{s+1} \\
&\quad - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2} - n_a + 1 + 1}^{q_{r_2}} e_{q_1 - n_a + i, q_1 - q_{r_2} + u}^{(r_1)} t^{-s-1} e_{q_1 - q_{r_2} + u, q_1 - n_a + i}^{(r_2)} t^{s+1} \\
&\quad + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2} - n_a + 1 + 1}^{q_{r_2}} e_{q_1 - n_a + i + 1, q_1 - q_{r_2} + u}^{(r_1)} t^{-s-1} e_{q_1 - q_{r_2} + u, q_1 - n_a + i + 1}^{(r_2)} t^{s+1} \\
&\quad - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2} - n_a + 1 + 1}^{q_{r_2}} e_{q_1 - n_a + i, q_1 - q_{r_2} + u}^{(r_2)} t^{-s-1} e_{q_1 - q_{r_2} + u, q_1 - n_a + i}^{(r_1)} t^{s+1} \\
&\quad + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2} - n_a + 1 + 1}^{q_{r_2}} e_{q_1 - n_a + i + 1, q_1 - q_{r_2} + u}^{(r_2)} t^{-s-1} e_{q_1 - q_{r_2} + u, q_1 - n_a + i + 1}^{(r_1)} t^{s+1}. \tag{6.7}
\end{aligned}$$

By the definition of  $\tilde{\mu}$ , we have

$$\begin{aligned}
& \hbar \tilde{\mu}(W_{q_1 - n_a + i, q_1 - n_a + i}^{(2)} t) - \hbar \tilde{\mu}(W_{q_1 - n_a + i + 1, q_1 - n_a + i + 1}^{(2)} t) \\
&= \hbar \sum_{r=1}^{b_a} \gamma_r e_{q_1 - n_a + i, q_1 - n_a + i}^{(r)} + \sum_{s \in \mathbb{Z}} \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{u > q_1 - n_a} e_{u, q_1 - n_a + i}^{(r_1)} t^{-s} e_{q_1 - n_a + i, u}^{(r_2)} t^s \\
&\quad - \hbar \sum_{s \geq 0} \sum_{r=1}^{b_a} \sum_{1 \leq u \leq q_r - n_a} e_{q_1 - q_r + u, q_1 - n_a + i}^{(r)} t^{-s-1} e_{q_1 - n_a + i, q_1 - q_r + u}^{(r)} t^{s+1} \\
&\quad - \hbar \sum_{s \geq 0} \sum_{r=1}^{b_a} \sum_{1 \leq u \leq q_r - n_a} e_{q_1 - n_a + i, q_1 - q_r + u}^{(r)} t^{-s} e_{q_1 - q_r + u, q_1 - n_a + i}^{(r)} t^s \\
&\quad - \hbar \sum_{s \in \mathbb{Z}} \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{1 \leq u \leq q_{r_2} - n_a} e_{q_1 - n_a + i, q_1 - q_{r_2} + u}^{(r_1)} t^{s+1} e_{q_1 - q_{r_2} + u, q_1 - n_a + i}^{(r_2)} t^{-s-1} \\
&\quad - \hbar \sum_{r=1}^{b_a} \gamma_r e_{q_1 - n_a + i + 1, q_1 - n_a + i + 1}^{(r)} - \sum_{s \in \mathbb{Z}} \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{u > q_1 - n_a} e_{u, q_1 - n_a + i + 1}^{(r_1)} t^{-s} e_{q_1 - n_a + i + 1, u}^{(r_2)} t^s \\
&\quad + \hbar \sum_{s \geq 0} \sum_{r=1}^{b_a} \sum_{1 \leq u \leq q_r - n_a} e_{q_1 - q_r + u, q_1 - n_a + i + 1}^{(r)} t^{-s-1} e_{q_1 - n_a + i + 1, q_1 - q_r + u}^{(r)} t^{s+1} \\
&\quad + \hbar \sum_{s \geq 0} \sum_{r=1}^{b_a} \sum_{1 \leq u \leq q_r - n_a} e_{q_1 - n_a + i + 1, q_1 - q_r + u}^{(r)} t^{-s} e_{q_1 - q_r + u, q_1 - n_a + i + 1}^{(r)} t^s \\
&\quad + \hbar \sum_{s \in \mathbb{Z}} \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{1 \leq u \leq q_{r_2} - n_a} e_{q_1 - n_a + i + 1, q_1 - q_{r_2} + u}^{(r_1)} t^{s+1} e_{q_1 - q_{r_2} + u, q_1 - n_a + i + 1}^{(r_2)} t^{-s-1}. \tag{6.8}
\end{aligned}$$

Here after, in order to simplify the notation, we denote the right hand side of  $i$ -th term of equation labeled  $( )$  by  $( )_i$ . The sum of (6.4)-(6.8) is equal to the sum of the left hand side of (6.3) and  $\hbar \tilde{\mu}(W_{q_1 - n_a + i, q_1 - n_a + i}^{(2)} t) - \hbar \tilde{\mu}(W_{q_1 - n_a + i + 1, q_1 - n_a + i + 1}^{(2)} t)$ . We divide this sum into 8 pieces:

$$(6.4)_1 + (6.8)_1 + (6.8)_6$$

$$= -\frac{i}{2}\hbar\tilde{\mu}(W_{q_1-n_a+i,q_1-n_a+i}^{(1)} - W_{q_1-n_a+i+1,q_1-n_a+i+1}^{(1)}), \quad (6.9)$$

$$(6.4)_2 + (6.5)_1 + (6.5)_2$$

$$= -\hbar\tilde{\mu}(W_{q_1-n_a+i,q_1-n_a+i}^{(1)} W_{q_1-n_a+i+1,q_1-n_a+i+1}^{(1)}), \quad (6.10)$$

$$(6.4)_3 + (6.4)_4 + (6.6)_1 + (6.7)_1 + (6.8)_3 + (6.8)_4$$

$$\begin{aligned} &= \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_1-n_a+1}^{q_1-n_a+i} e_{q_1-n_a+i,u}^{(r)} t^{-s} e_{u,q_1-n_a+i}^{(r)} t^s \\ &\quad + \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_1-n_a+i+1}^{q_1-n_a+i+1} e_{q_1-n_a+i,u}^{(r)} t^{-s-1} e_{u,q_1-n_a+i}^{(r)} t^{s+1}, \end{aligned} \quad (6.11)$$

$$(6.4)_5 + (6.4)_6 + (6.6)_1 + (6.7)_2 + (6.8)_8 + (6.8)_9$$

$$\begin{aligned} &= -\hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_1-n_a+1}^{q_1-n_a+i} e_{q_1-n_a+i+1,u}^{(r)} t^{-s} e_{u,q_1-n_a+i+1}^{(r)} t^s \\ &\quad - \hbar \sum_{r=1}^{b_a} \sum_{s \geq 0} \sum_{u=q_1-n_a+i+1}^{q_1-n_a+i+1} e_{q_1-n_a+i+1,u}^{(r)} t^{-s-1} e_{u,q_1-n_a+i+1}^{(r)} t^{s+1} \end{aligned} \quad (6.12)$$

$$(6.5)_3 + (6.5)_5 + (6.6)_3 + (6.7)_5 + (6.8)_2$$

$$\begin{aligned} &= \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+1}^{q_{r_2}-n_a+i} e_{q_1-q_{r_2}+u,q_1-n_a+i}^{(r_1)} t^s E_{q_1-q_{r_2}+i,q_1-q_{r_2}+u}^{(r_2)} t^{-s} \\ &\quad + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+i+1}^{q_{r_2}-n_a+i+1} e_{q_1-q_{r_2}+u,q_1-n_a+i}^{(r_1)} t^{s+1} e_{q_1-n_a+i,q_1-q_{r_2}+u}^{(r_2)} t^{-s-1}, \end{aligned} \quad (6.13)$$

$$(6.5)_4 + (6.5)_6 + (6.6)_5 + (6.7)_3 + (6.8)_5$$

$$\begin{aligned} &= \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+1}^{q_{r_2}-n_a+i} e_{q_1-n_a+i,q_1-q_{r_2}+u}^{(r_1)} t^{-s} e_{q_1-q_{r_2}+u,q_1-n_a+i}^{(r_2)} t^s \\ &\quad + \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+i+1}^{q_{r_2}-n_a+i+1} e_{q_1-n_a+i,q_1-q_{r_2}+u}^{(r_1)} t^{-s-1} e_{q_1-q_{r_2}+u,q_1-n_a+i}^{(r_2)} t^{s+1}, \end{aligned} \quad (6.14)$$

$$(6.5)_7 + (6.5)_9 + (6.6)_4 + (6.7)_6 + (6.8)_7$$

$$\begin{aligned} &= -\hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+1}^{q_{r_2}-n_a+i} e_{q_1-q_{r_2}+u,q_1-n_a+i+1}^{(r_1)} t^s E_{q_1-n_a+i+1,q_1-q_{r_2}+u}^{(r_2)} t^{-s} \\ &\quad - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+i+1}^{q_{r_2}-n_a+i+1} e_{q_1-q_{r_2}+u,q_1-n_a+i+1}^{(r_1)} t^{s+1} e_{q_1-n_a+i+1,q_1-q_{r_2}+u}^{(r_2)} t^{-s-1}, \end{aligned} \quad (6.15)$$

$$(6.5)_8 + (6.5)_{10} + (6.6)_6 + (6.7)_4 + (6.8)_{10}$$

$$\begin{aligned} &= -\hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+1}^{q_{r_2}-n_a+i} e_{q_1-n_a+i+1,q_1-q_{r_2}+u}^{(r_1)} t^{-s} e_{q_1-q_{r_2}+u,q_1-n_a+i+1}^{(r_2)} t^s \\ &\quad - \hbar \sum_{1 \leq r_1 < r_2 \leq b_a} \sum_{s \geq 0} \sum_{u=q_{r_2}-n_a+i+1}^{q_{r_2}-n_a+i+1} e_{q_1-n_a+i+1,q_1-q_{r_2}+u}^{(r_1)} t^{-s-1} e_{q_1-q_{r_2}+u,q_1-n_a+i+1}^{(r_2)} t^{s+1}. \end{aligned} \quad (6.16)$$

By a direct computation, we find that the sum of (6.13)-(6.16) is equal to

$$\begin{aligned} & \hbar \sum_{u=1}^i W_{q_1-n_a+i, q_1-n_a+u}^{(1)} t^{-s} W_{q_1-n_a+u, q_1-n_a+i}^{(1)} t^s \\ & + \hbar \sum_{u=i+1}^{n_a-n_a+1} W_{q_1-n_a+i, q_1-n_a+u}^{(1)} t^{-s-1} W_{q_1-n_a+u, q_1-n_a+i}^{(1)} t^{s+1} \\ & - \hbar \sum_{u=1}^i W_{q_1-n_a+i+1, q_1-n_a+u}^{(1)} t^{-s} W_{q_1-n_a+u, q_1-n_a+i+1}^{(1)} t^s \\ & - \hbar \sum_{u=i+1}^{n_a-n_a+1} W_{q_1-n_a+i+1, q_1-n_a+u}^{(1)} t^{-s-1} W_{q_1-n_a+u, q_1-n_a+i+1}^{(1)} t^{s+1}. \end{aligned}$$

Thus, we have obtained the relation (6.3).  $\square$

## 7 Affine Yangians and some cosets of non-rectangular $W$ -algebras

Restricting the homomorphism  $\Phi$  to  $Y_{\hbar, \varepsilon_s}(\widehat{\mathfrak{sl}}(n_s - n_{s+1}))$ , we obtain the homomorphism

$$\Phi_s: Y_{\hbar, \varepsilon_s}(\widehat{\mathfrak{sl}}(n_s - n_{s+1})) \rightarrow \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f)).$$

These are homomorphisms given in Theorem 4.7 of [27]. By Theorem 6.1, we obtain the following theorem.

**Theorem 7.1.** *The images of  $\Phi_s$  are commutative with each other.*

In this section, we consider these homomorphisms from the perspective of the coset of a vertex algebra. For a vertex algebra  $A$  and its vertex subalgebra  $B$ , we set a coset vertex algebra of the pair  $(A, B)$  as follows:

$$C(A, B) = \{v \in A \mid w_{(r)}v = 0 \text{ for } w \in B \text{ and } r \geq 0\}.$$

### 7.1 The case that $b_1 = 1$

In the case that  $b_1 = 1$ , we have an embedding from  $\mathcal{U}(\widehat{\mathfrak{gl}}(n_1 - n_2))$  to  $\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f))$  given by  $E_{i,j}[-1]t^s$  to  $W_{i,j}^{(1)}t^s$ . Through this embedding,  $\Phi_1$  becomes an evaluation map for the affine Yangian.

We note that  $\mathcal{U}(\widehat{\mathfrak{gl}}(n_1 - n_2))$  can be considered as the universal enveloping algebra of the universal affine vertex algebra associated with  $\mathfrak{gl}(n_1 - n_2)$  and the inner product defined by

$$\kappa(E_{i,j}, E_{p,q}) = (k + N - q_1)\delta_{i,q}\delta_{p,j} + \delta_{i,j}\delta_{p,q}.$$

By Theorem 7.1 and Theorem 4.1, we obtain the following theorem by the same way as Theorem 5.9 in [28]:

**Theorem 7.2.** *In the case that  $k + N - q_1 \neq 0$ , the images of  $\Phi_a$  are contained in the universal enveloping algebra of  $C(\mathcal{W}^k(\mathfrak{gl}(N), f), V^\kappa(\mathfrak{sl}(n_1 - n_2)))$ .*

**Remark 7.3.** In the case that  $l = 2$  and  $q_1 = m > q_2 = n$ , we show in Theorem 9.3 of [25] that the image of  $\Phi_2$  is contained in the universal enveloping algebra of  $C(\mathcal{W}^k(\mathfrak{gl}(m+n), f), V^\kappa(\mathfrak{sl}(m-n)))$  for any  $k \in \mathbb{C}$ . We expect that Theorem 7.2 holds for any  $k \in \mathbb{C}$ .

## 7.2 The case that $b_1 = 2$

First, let us recall the main results of [29], which gave  $\Phi_1$  for a rectangular  $W$ -algebra.

**Theorem 7.4** (Theorem 3.17 and Theorem 5.1 in [29]). 1). In the case that  $k + (l - 1)n \neq 0$ , the rectangular  $W$ -algebra  $\mathcal{W}^k(\mathfrak{gl}(ln), (l^n))$  is generated by  $\{W_{i,j}^{(r)} \mid 1 \leq i, j \leq n, r = 1, 2\}$ .

2). The image of the homomorphism

$$\Phi_1: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(ln), (l^n)))$$

is dense in the target if  $k + (l - 1)n \neq 0$ .

Let us denote by  $\widetilde{\mathcal{W}}^k(\mathfrak{gl}(b_1(n_1 - n_2)), (b_1^{n_1 - n_2}))$  the subalgebra of  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  generated by  $\{W_{i,j}^{(r)} \mid r = 1, 2, 1 \leq i, j \leq n_1 - n_2\}$ . The following theorem follows from Theorem 2.5 and Theorem 7.4.

**Lemma 7.5.** 1). The subalgebra  $\widetilde{\mathcal{W}}^k(\mathfrak{gl}(2(n_1 - n_2)), (2^{n_1 - n_2}))$  is isomorphic to the rectangular  $W$ -algebra  $\mathcal{W}^{k+n_2+\sum_{s=3}^l q_s}(\mathfrak{gl}(2(n_1 - n_2)), (2^{n_1 - n_2}))$ .

2). In the case that  $b_1 = 2$ , the completion of the image of  $\Phi_1$  coincides with the universal enveloping algebra of  $\widetilde{\mathcal{W}}^k(\mathfrak{gl}(2(n_1 - n_2)), (2^{n_1 - n_2}))$ .

By Lemma 7.5, Theorem 7.1 and Theorem 6.1, we obtain the following theorem.

**Theorem 7.6.** The homomorphism  $\Phi_s$  the homomorphism  $\Phi_s$  induces a homomorphism

$$\tilde{\Phi}_s: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n_s - n_{s+1})) \rightarrow C(\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f)), \mathcal{U}(\mathcal{W}^{k+n_2+\sum_{s=3}^l q_s}(\mathfrak{gl}(2(n_1 - n_2)), (2^{n_1 - n_2})))),$$

where  $C(\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f)), \mathcal{U}(\mathcal{W}^{k+n_2+\sum_{s=3}^l q_s}(\mathfrak{gl}(2(n_1 - n_2)), (2^{n_1 - n_2}))))$  is the centralizer algebra of  $\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f))$  and  $\mathcal{U}(\mathcal{W}^{k+n_2+\sum_{s=3}^l q_s}(\mathfrak{gl}(2(n_1 - n_2)), (2^{n_1 - n_2})))$ .

By the same way as Theorem 6.5 in [28], we obtain the following theorem from Lemma 2.4.

**Theorem 7.7.** The homomorphism  $\Phi_s$  the homomorphism  $\Phi_s$  induces a homomorphism

$$\tilde{\Phi}_s: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n_s - n_{s+1})) \rightarrow \mathcal{U}(C(\mathcal{W}^k(\mathfrak{gl}(N), f), \mathcal{W}^{k+n_2+\sum_{s=3}^l q_s}(\mathfrak{sl}(2(n_1 - n_2)), (2^{n_1 - n_2})))),$$

In the appendix, we will consider the case that  $b_1 > 2$ .

## A Elements of $\mathcal{W}^k(\mathfrak{gl}(N), f)$

In this section, we construct elements of  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  which are correpponding to Arkawa-Molev's strong generators of rectangular  $W$ -algebras.

We consider two universal affine vertex algebras. The first one is associated with a Lie algebra

$$\mathfrak{b} = \bigoplus_{\substack{1 \leq i, j \leq N \\ \text{col}(i) \geq \text{col}(j)}} \mathbb{C}E_{i,j} \subset \mathfrak{gl}(N)$$

and its inner product

$$\kappa_{\mathfrak{b}}(E_{i,j}, E_{p,q}) = \alpha_{\text{col}(i)}\delta_{i,q}\delta_{p,j} + \delta_{i,j}\delta_{p,q}.$$

The second one is the universal affine vertex algebra associated with a Lie superalgebra  $\mathfrak{a} = \mathfrak{b} \oplus \bigoplus_{\substack{1 \leq i, j \leq N \\ \text{col}(i) > \text{col}(j)}} \mathbb{C}\psi_{i,j}$  with the following commutator relations;

$$[E_{i,j}, \psi_{p,q}] = \delta_{j,p}\psi_{i,q} - \delta_{i,q}\psi_{p,j}, \quad [\psi_{i,j}, \psi_{p,q}] = 0,$$

where  $e_{i,j}$  is an even element and  $\psi_{i,j}$  is an odd element. We set the inner product on  $\mathfrak{a}$  such that

$$\tilde{\kappa}_{\mathfrak{b}}(E_{i,j}, E_{p,q}) = \kappa_{\mathfrak{b}}(E_{i,j}, E_{p,q}), \quad \tilde{\kappa}_{\mathfrak{b}}(E_{i,j}, \psi_{p,q}) = \tilde{\kappa}_{\mathfrak{b}}(\psi_{i,j}, \psi_{p,q}) = 0.$$

By the definition of  $V^{\tilde{\kappa}}(\mathfrak{a})$  and  $V^{\kappa}(\mathfrak{b})$ ,  $V^{\tilde{\kappa}}(\mathfrak{a})$  contains  $V^{\kappa}(\mathfrak{b})$ .

In this section, we regard  $V^{\tilde{\kappa}}(\mathfrak{a})$  (resp.  $V^{\kappa}(\mathfrak{b})$  and  $V^{\tilde{\kappa}}(\mathfrak{a}) \otimes \mathbb{C}[\tau]$ ) as a non-associative superalgebra whose product  $\cdot$  is defined by

$$u[-w] \cdot v[-s] = (u[-w])_{(-1)} v[-s], \quad [\tau, u[-s]] = su[-s],$$

where  $\tau$  is an even element. We sometimes omit  $\cdot$  and in order to simplify the notation. By [15] and [16], a  $W$ -algebra  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  can be realized as a subalgebra of  $V^{\kappa}(\mathfrak{b})$ .

Let us define an odd differential  $d_0: V^{\kappa}(\mathfrak{b}) \rightarrow V^{\tilde{\kappa}}(\mathfrak{a})$  determined by

$$d_0 1 = 0, \tag{A.1}$$

$$[d_0, \partial] = 0, \tag{A.2}$$

$$\begin{aligned} [d_0, e_{i,j}[-1]] &= \sum_{\text{col}(i) > \text{col}(r) \geq \text{col}(j)} e_{r,j}[-1] \psi_{i,r}[-1] - \sum_{\text{col}(j) < \text{col}(r) \leq \text{col}(i)} \psi_{r,j}[-1] e_{i,r}[-1] \\ &\quad + \delta(\text{col}(i) > \text{col}(j)) \alpha_{\text{col}(i)} \psi_{i,j}[-2] + \psi_{\tilde{i},j}[-1] - \psi_{i,\tilde{j}}[-1]. \end{aligned} \tag{A.3}$$

By using Theorem 2.4 in [14], we can define the  $W$ -algebra  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  as follows.

**Definition A.4.** The  $W$ -algebra  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  is the vertex subalgebra of  $V^{\kappa}(\mathfrak{b})$  defined by

$$\mathcal{W}^k(\mathfrak{gl}(N), f) = \{y \in V^{\kappa}(\mathfrak{b}) \subset V^{\tilde{\kappa}}(\mathfrak{a}) \mid d_0(y) = 0\}.$$

*Remark A.5.* The projection from  $\mathfrak{b}$  to  $\bigotimes_{1 \leq i \leq l} \mathfrak{gl}(q_i)$  induces the Miura map  $\mu$ , that is, an embedding from  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  to  $\bigotimes_{1 \leq i \leq l} V^{\kappa_i}(\mathfrak{gl}(q_i))$ .

By using Definition A.4, we construct some elements of  $\mathcal{W}^k(\mathfrak{gl}(N), f)$ .

We denote by  $T(C)$  a non-associative free algebra associated with a vector space  $C$  and by  $\mathfrak{gl}(l)_{\leq 0}$  the Lie algebra  $\bigoplus_{1 \leq j \leq i \leq l} \mathbb{C} E_{i,j}$ . Let us set an  $q_1 \times q_1$  matrix  $B = (b_{i,j})_{1 \leq i,j \leq q_1}$  as

$$\left[ \begin{array}{ccccc} \alpha_1 \tau + E_{1,1}[-1] & -1 & 0 & \dots & 0 \\ E_{2,1}[-1] & \alpha_1 \tau + E_{2,2}[-1] & -1 & \dots & 0 \\ \vdots & \vdots & \ddots & & \vdots \\ E_{l-1,1}[-1] & E_{l-1,2}[-1] & \dots & \alpha_1 \tau + E_{l-1,l-1}[-1] & -1 \\ E_{l,1}[-1] & E_{l,2}[-1] & \dots & E_{l,l-1}[-1] & \pi \tau + E_{l,l}[-1] \end{array} \right] \tag{A.6}$$

whose entries are elements of  $T(\mathfrak{gl}(l)_{\leq 0}[t^{-1}]t^{-1}) \otimes \mathbb{C}[\tau]$ . For any matrix  $A = (a_{i,j})_{1 \leq i,j \leq s}$ , we define  $\text{cdet}(A)$  as

$$\sum_{\sigma \in \mathfrak{S}_s} \text{sgn}(\sigma) a_{\sigma(1),1} (a_{\sigma(2),2} (a_{\sigma(3),3} \cdots a_{\sigma(s-1),s-1}) a_{\sigma(s),s}) \in T(\mathfrak{gl}(l)_{\leq 0}[t^{-1}]t^{-1}) \otimes \mathbb{C}[\tau].$$

We regard  $\mathfrak{gl}(q_1)$  as an associative superalgebra whose product  $\cdot$  is determined by  $E_{i,j} \cdot E_{s,u} = \delta_{j,s} E_{i,u}$ . Then, we obtain a non-associative superalgebra  $\mathfrak{gl}(q_1) \otimes V^{\kappa}(\mathfrak{b}) \otimes \mathbb{C}[\tau]$ . We construct a homomorphism

$$T: T(\mathfrak{gl}(l)_{\leq 0}[t^{-1}]t^{-1}) \otimes \mathbb{C}[\tau] \rightarrow \mathfrak{gl}(m|n) \otimes V^{\kappa}(\mathfrak{b}) \otimes \mathbb{C}[\tau]$$

determined by

$$T_{i,j}(E_{u,v}[-s]) = E_{(u-1)q_1+i, (v-1)q_1+j}[-s] \in \mathfrak{b}[t^{-1}]t^{-1}, \quad T(\tau) = \tau,$$

where  $T_{i,j}(x)$  is defined as  $E_{j,i} \otimes T_{i,j}(x) = T(x)$ . Since  $T$  is a homomorphism, we obtain

$$T_{i,j}(xy) = \sum_{r=1}^{q_1} T_{r,i}(x) T_{j,r}(y).$$

By the commutator relation of  $V^\kappa(\mathfrak{b})$  and  $\mathbb{C}[\tau]$ , we define  $\widetilde{W}_{i,j}^{(r)} \in V^{\kappa_b}(\mathfrak{b})$  is defined by

$$T_{j,i}(\text{cdet}(B)) = \sum_{r=0}^l \widetilde{W}_{j,i}^{(r)} (\alpha_1 \tau)^{l-r}. \quad (\text{A.7})$$

**Theorem A.8.** *The  $W$ -algebra  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  contains  $\{\widetilde{W}_{i,j}^{(r)} \mid 1 \leq r \leq l, 1 \leq i, j \leq n_1 - n_2\}$ .*

The proof of Theorem A.8 is as the one of Arakawa-Molev [1] (see also Theorem 3.11 and Remark 3.9) since we take  $1 \leq i, j \leq n_1 - n_2$ .

In particular, by (A.7), we have

$$\mu(\widetilde{W}_{i,j}^{(1)}) = \sum_{1 \leq s \leq b_1} e_{(s-1)q_1+i, (s-1)q_1+j}[-1], \quad (\text{A.9})$$

$$\begin{aligned} \mu(\widetilde{W}_{i,j}^{(2)}) &= \alpha_1 \sum_{1 \leq s \leq b_1} (s-1) e_{(s-1)q_1+i, (s-1)q_1+j}[-2] \\ &\quad + \sum_{\substack{r_1 < r_2 \\ 1 \leq t \leq m+n}} (-1)^{p(t)+p(e_{i,t})p(e_{j,t})} e_{t,i}^{(r_1)}[-1] e_{j,t}^{(r_2)}[-1]. \end{aligned} \quad (\text{A.10})$$

By the form of  $W_{i,j}^{(1)}$  and  $W_{i,j}^{(2)}$ , we find that the OPEs of  $\{W_{i,j}^{(r)} \mid r = 1, 2, 1 \leq i, j \leq n_1 - n_2\}$  in  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  is the same as those of

$$\{\widetilde{W}_{i,j}^{(1)}, W_{i,j}^{(2)} - \gamma_1 \partial W_{i,j}^{(1)} \mid r = 1, 2, 1 \leq i, j \leq n_1 - n_2\} \quad (\text{A.11})$$

in  $\mathcal{W}^{k+\sum_{s=3}^l q_s}(\mathfrak{gl}(q_1 b_1), (b_1^{q_1}))$ . Similarly to Theorem 7.7, we find the following theorem.

**Theorem A.12.** *The homomorphism  $\Phi_s$  the homomorphism  $\Phi_s$  induces a homomorphism*

$$\Phi: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n_s - n_{s+1})) \rightarrow C(\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f)), \mathcal{U}(\widetilde{\mathcal{W}}^k(\mathfrak{gl}(b_1(n_1 - n_2)), (b_1^{n_1 - n_2})))),$$

where  $C(\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f)), \mathcal{U}(\widetilde{\mathcal{W}}^k(\mathfrak{gl}(b_1(n_1 - n_2)), (b_1^{n_1 - n_2}))))$  is the centralizer of  $\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f))$  and  $\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(b_1(n_1 - n_2)), (b_1^{n_1 - n_2})))$ .

The OPEs of the elements (A.11) in the rectangular  $W$ -algebra  $\mathcal{W}^{k+\sum_{s=b_1+1}^l q_s}(\mathfrak{gl}(q_1 b_1), (b_1^{q_1}))$  is the same as those in  $\mathcal{W}^{k+n_2(b_1-1)+\sum_{s=b_1+1}^l q_s}(\mathfrak{gl}((n_1 - n_2)b_1), (b_1^{n_1 - n_2}))$ .

**Conjecture A.13.** The subalgebra  $\widetilde{\mathcal{W}}^k(\mathfrak{gl}(b_1(n_1 - n_2)), (b_1^{n_1 - n_2}))$  is isomorphic to the rectangular  $W$ -algebra  $\mathcal{W}^{k+n_2(b_1-1)+\sum_{s=b_1+1}^l q_s}(\mathfrak{gl}((n_1 - n_2)b_1), (b_1^{n_1 - n_2}))$ .

If we can prove this conjecture, we can obtain the homomorphism

$$\widetilde{\Phi}_s: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n_s - n_{s+1})) \rightarrow \mathcal{U}(C(\mathcal{W}^k(\mathfrak{gl}(N), f), \mathcal{W}^{k+(b_1-1)n_2+\sum_{s=3}^l q_s}(\mathfrak{sl}(b_1(n_1 - n_2)), (b_1^{n_1 - n_2}))))$$

by the same way as Theorem 6.5 in [28].

## References

- [1] T. Arakawa and A. Molev. Explicit generators in rectangular affine  $W$ -algebras of type  $A$ . *Lett. Math. Phys.*, 107(1):47–59, 2017, <https://doi.org/10.1007/s11005-016-0890-2>.
- [2] N. Arbesfeld and O. Schiffmann. A presentation of the deformed  $W_{1+\infty}$  algebra. In *Symmetries, integrable systems and representations*, volume 40 of *Springer Proc. Math. Stat.*, pages 1–13. Springer, Heidelberg, 2013.
- [3] J. Brundan and A. Kleshchev. Shifted Yangians and finite  $W$ -algebras. *Adv. Math.*, 200(1):136–195, 2006, <https://doi.org/10.1016/j.aim.2004.11.004>.
- [4] T. Creutzig, D. E. Diaconescu, and M. Ma. Affine Laumon spaces and iterated  $\mathcal{W}$ -algebras. *Comm. Math. Phys.*, 402(3):2133–2168, 2023.
- [5] T. Creutzig and A. R. Linshaw. Trialities of  $\mathcal{W}$ -algebras. *Camb. J. Math.*, 10(1):69–194, 2022.
- [6] A. De Sole, V. G. Kac, and D. Valeri. A Lax type operator for quantum finite  $W$ -algebras. *Selecta Math. (N.S.)*, 24(5):4617–4657, 2018.
- [7] M. Finkelberg and A. Tsymbaliuk. Multiplicative slices, relativistic Toda and shifted quantum affine algebras. In *Representations and nilpotent orbits of Lie algebraic systems*, volume 330 of *Progr. Math.*, pages 133–304. Birkhäuser/Springer, Cham, 2019.
- [8] M. R. Gaberdiel and R. Gopakumar. Triality in minimal model holography. *J. High Energy Phys.*, (7):127, front matter+26, 2012.
- [9] D. Gaiotto and M. Rapčák. Vertex algebras at the corner. *J. High Energy Phys.*, (1):160, front matter+85, 2019.
- [10] N. Genra. Screening operators for  $\mathcal{W}$ -algebras. *Selecta Math. (N.S.)*, 23(3):2157–2202, 2017, <https://doi.org/10.1007/s00029-017-0315-9>.
- [11] N. Guay. Cherednik algebras and Yangians. *Int. Math. Res. Not.*, (57):3551–3593, 2005, <https://doi.org/10.1155/IMRN.2005.3551>.
- [12] N. Guay. Affine Yangians and deformed double current algebras in type A. *Adv. Math.*, 211(2):436–484, <https://doi.org/10.1016/j.aim.2006.08.007>, 2007.
- [13] N. Guay, H. Nakajima, and C. Wendlandt. Coproduct for Yangians of affine Kac-Moody algebras. *Adv. Math.*, 338:865–911, 2018, <https://doi.org/10.1016/j.aim.2018.09.013>.
- [14] V. Kac, S. S. Roan, and M. Wakimoto. Quantum reduction for affine superalgebras. *Comm. Math. Phys.*, 241(2-3):307–342, 2003, <https://doi.org/10.1007/s00220-003-0926-1>.
- [15] V. G. Kac and M. Wakimoto. Quantum reduction and representation theory of superconformal algebras. *Adv. Math.*, 185(2):400–458, 2004, <https://doi.org/10.1016/j.aim.2003.12.005>.
- [16] V. G. Kac and M. Wakimoto. Corrigendum to: “Quantum reduction and representation theory of superconformal algebras” [Adv. Math. **185** (2004), no. 2, 400–458; mr2060475]. *Adv. Math.*, 193(2):453–455, 2005, <https://doi.org/10.1016/j.aim.2005.01.001>.
- [17] R. Kodera. Braid group action on affine Yangian. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 15:Paper No. 020, 28, 2019, <https://doi.org/10.3842/SIGMA.2019.020>.
- [18] R. Kodera. On Guay’s evaluation map for affine Yangians. *Algebr. Represent. Theory*, 24(1):253–267, 2021, <https://doi.org/10.1007/s10468-019-09945-w>.
- [19] R. Kodera and M. Ueda. Coproduct for affine Yangians and parabolic induction for rectangular  $W$ -algebras. *Lett. Math. Phys.*, 112(1):Paper No. 3, 37, 2022.

- [20] A. Matsuo, K. Nagatomo, and A. Tsuchiya. Quasi-finite algebras graded by Hamiltonian and vertex operator algebras. *London Math. Soc. Lecture Note Ser.*, 372:282–329, 2010, <https://doi.org/10.1017/CBO9780511730054.015>.
- [21] D. Maulik and A. Okounkov. Quantum groups and quantum cohomology. *Astérisque*, (408):ix+209, 2019.
- [22] M. Rapčák. On extensions of  $\widehat{\mathfrak{gl}(m|n)}$  Kac-Moody algebras and Calabi-Yau singularities. *J. High Energy Phys.*, (1):042, 34, 2020, [https://doi.org/10.1007/jhep01\(2020\)042](https://doi.org/10.1007/jhep01(2020)042).
- [23] O. Schiffmann and E. Vasserot. Cherednik algebras, W-algebras and the equivariant cohomology of the moduli space of instantons on  $\mathbf{A}^2$ . *Publ. Math. Inst. Hautes Études Sci.*, 118:213–342, 2013, <https://doi.org/10.1007/s10240-013-0052-3>.
- [24] A. Tsymbaliuk. The affine Yangian of  $\mathfrak{gl}_1$  revisited. *Adv. Math.*, 304:583–645, 2017.
- [25] M. Ueda. The coproduct for the affine Yangian and the parabolic induction for non-rectangular  $W$ -algebras. arXiv:2404.14096.
- [26] M. Ueda. A homomorphism from the affine Yangian  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  to the affine Yangian  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n+1))$ . arXiv:2312.09933.
- [27] M. Ueda. Notes on a homomorphism from the affine Yangian associated with  $\widehat{\mathfrak{sl}}(n)$  to the affine Yangian associated with  $\widehat{\mathfrak{sl}}(n)$ . arXiv:2402.01870.
- [28] M. Ueda. Two homomorphisms from the affine Yangian associated with  $\widehat{\mathfrak{sl}}(n)$  to the affine Yangian associated with  $\widehat{\mathfrak{sl}}(n)$ . arXiv:2404.10923.
- [29] M. Ueda. Affine super Yangians and rectangular  $W$ -superalgebras. *J. Math. Phys.*, 63(5):Paper No. 051701, 34, 2022.
- [30] M. Ueda. Guay’s affine Yangians and non-rectangular  $W$ -algebras. *Adv. Math.*, 438:Paper No. 109468, 44, 2024.