

SPECTRAL SHIFT FUNCTIONS OF ALL ORDERS

ARUP CHATTOPADHYAY, TEUN D.H. VAN NULAND, AND CHANDAN PRADHAN

ABSTRACT. Let $n \in \mathbb{N}$ and let H_0, V be self-adjoint operators such that V is bounded and $V(H_0 - i)^{-p} \in \mathcal{S}^{n/p}$ for $p = 1, \dots, n$. We prove the existence, uniqueness up to polynomial summands, and regularity properties of all higher order spectral shift functions associated to the perturbation theory of $t \mapsto H_0 + tV$ ($t \in [0, 1]$). Such perturbations arise in situations in noncommutative geometry and mathematical physics with dimension $n - 1$. Our proof is simpler than that of [13, 14] and works at all orders regardless of n .

1. INTRODUCTION

The spectral shift function is a useful notion for the spectral analysis of quantum systems, and the question of its existence has sparked enormous progress in mathematical perturbation theory. It appears when analysing the movement of the spectrum of $H_0 + tV$ ($t \in [0, 1]$) as a bounded self-adjoint perturbation V of a self-adjoint operator H_0 is gradually switched on. First defined by Krein [8], and generalized to higher order by Koplienko [7], the spectral shift function of order k is the function $\eta_k = \eta_{H_0, V, k}$ such that for all sufficiently regular $f : \mathbb{R} \rightarrow \mathbb{C}$ we have

$$\mathrm{Tr} \left(f(H_0 + V) - \sum_{m=0}^{k-1} \frac{d^m}{dt^m} f(H_0 + tV) \Big|_{t=0} \right) = \int_{\mathbb{R}} \eta_k(x) f^{(k)}(x) dx.$$

In [15] the spectral shift function η_k of order $k \geq n$ was shown to exist whenever $V \in \mathcal{S}^n$. Because physical situations – for example when H_0 is a differential operator and V a multiplication operator – require V to have continuous spectrum, the paper [13] used the much weaker assumption $V(H_0 - i)^{-1} \in \mathcal{S}^n$ ($n \in \mathbb{N}$) to obtain the spectral shift functions of order $k \geq n$. Under a similar assumption, [14] added existence of η_{n-1} when n is even. Throughout the literature one finds many such assumptions which generalise the case of $(H_0 - i)^{-1} \in \mathcal{S}^n$ to a ‘locally compact’ setting. Often, existence of the spectral shift function depends on n , which for a differential operator H_0 depends on its order and the dimension of the underlying space. The notable exception is that η_1 was shown to exist independently of n in [22]. However, for general $n \in \mathbb{N}$, existence of $\eta_2, \dots, \eta_{n-1}$ remained open.

In this paper we prove existence, uniqueness and regularity properties of η_k for all orders $k \in \mathbb{N}$ under the assumptions that $V = V^* \in \mathcal{B}(\mathcal{H})$ and

$$V(H_0 - i)^{-p} \in \mathcal{S}^{n/p} \quad (p = 1, \dots, n), \quad (1.1)$$

2010 *Mathematics Subject Classification.* 47A55.

Key words and phrases. Spectral action functional, multiple operator integral, weighted relative perturbation.

for an arbitrary $n \in \mathbb{N}$. In addition, our proof is much simpler than that of [13, 14]. The assumption (1.1) unifies the ‘locally compact’ assumption $V(H_0 - i)^{-n} \in \mathcal{S}^1$ with the ‘relative Schatten’ assumption $V(H_0 - i)^{-1} \in \mathcal{S}^n$ and is not new – cf. [17, 19, 21]. We exemplify its applicability in Section 5.

Our simple yet novel contribution is that, under the right assumptions, existence of the spectral shift function η_k can be deduced from the existence of η_{k-1} . Our proof is thus split into induction basis – existence of η_1 – in Section 3, and induction step in Section 4. Under different assumptions, existence of η_1 was already known, and the reader only interested in the cases already covered by [22] needs only to read Section 4.

Both parts of the proof make use of the theory of multiple operator integration, which is well-equipped to deal with these sorts of questions.

As we will see, assumption (1.1) is especially convenient to obtain Schatten class membership and bounds of the multiple operator integral, and thus in particular, of higher order derivatives of functions of operators, as well as differences of functions of operators. Combining this with the ease with which multiple operator integrals can be manipulated algebraically, we are encouraged that the use of multiple operator integrals under the standing assumption (1.1) can be useful for similar investigations. We are particularly referring to the development of a perturbation theoretical framework that can handle quantum theory and geometry at the same time, through the use of operator algebra [3, 4, 6, 17, 21].

This paper is structured as follows. We introduce the relevant function spaces and theory of multiple operator integration in the preliminary section 2. The induction basis is proved in Section 3 and the induction step in Section 4. Examples are discussed in Section 5.

1.1. Acknowledgements. We gratefully acknowledge Anna Skripka and Fedor Sukochev for useful discussions. A. Chattopadhyay is supported by the Core Research Grant (CRG), File No: CRG/2023/004826, by SERB, DST, Govt. of India. C. Pradhan acknowledges support from the IoE post-doctoral fellowship at IISc Bangalore. T.D.H. van Nuland was supported in part by ARC (Australian Research Council) grant FL17010005, and in part by the NWO project ‘Noncommutative multi-linear harmonic analysis and higher order spectral shift’, OCENW.M.22.070.

2. PRELIMINARIES

We denote positive constants by letters c, d, C, D with subscripts indicating dependence on their parameters. For instance, the symbol c_α denotes a constant depending only on the parameter α .

2.1. Function spaces. Let $n \in \mathbb{N} = \{1, 2, \dots\}$ and let X be an interval in \mathbb{R} possibly coinciding with \mathbb{R} . Let $C(X)$ the space of all continuous functions on X , $C_0(\mathbb{R})$ the space of continuous functions on \mathbb{R} decaying to 0 at infinity, $C^n(X)$ the space of n -times continuously differentiable functions on X . Let $C_b^n(X)$ denote the subset of $C^n(X)$ of such f for which $f^{(n)}$ is bounded. Let $C_c^n(\mathbb{R})$ denote the subspace of $C^n(\mathbb{R})$ consisting of compactly supported functions. We also use the notation $C^0(\mathbb{R}) = C(\mathbb{R})$. Let $a, b \in \mathbb{R}$. Let $C_c^n((a, b))$ denote the

subspace of $C_c^n(\mathbb{R})$ consisting of the functions whose closed support is contained in (a, b) . We will need the following elementary lemma.

Lemma 2.1. *Let $a, b \in \mathbb{R}$, $a < b$, $k \in \mathbb{N}$, and $f \in C_c^k((a, b))$. Then,*

$$\|f^{(j)}\|_\infty \leq (b-a)^{k-j} \|f^{(k)}\|_\infty, \quad j = 0, \dots, k.$$

Let $L^p(\mathbb{R})$ denote the space of measurable f for which $|f|^p$ is Lebesgue integrable on \mathbb{R} equipped with the standard norm $\|f\|_p = \|f\|_{L^p(\mathbb{R})} := (\int_{\mathbb{R}} |f(x)|^p dx)^{1/p}$, $1 \leq p < \infty$, and let $L^\infty(\mathbb{R})$ denote the space of essentially bounded functions on \mathbb{R} equipped with the essential supremum norm $\|\cdot\|_\infty$. Let $L_{\text{loc}}^1(\mathbb{R})$ denote the space of functions locally integrable on \mathbb{R} equipped with the seminorms $f \mapsto \int_{-a}^a |f(x)| dx$, $a > 0$. For an $L^1(\mathbb{R})$ -function f , \hat{f} denotes the Fourier transform of f . Whenever we write $\hat{f} \in L^1(\mathbb{R})$, it is implicitly assumed that $f \in C_0(\mathbb{R})$.

Let us introduce the function class that features in our main theorem. Although it looks technical at first, the class neatly shows the influences of the summability parameter n , and the order k of the spectral shift function, on the required decay and differentiability of the function f . Let $u(\lambda) = (\lambda - i)$, $\lambda \in \mathbb{R}$. For $a, b \in \mathbb{R}$, let $a \vee b = \max\{a, b\}$.

Definition 2.2. *Let $n, k \in \mathbb{N}$. Let $\mathfrak{Q}_n^k(\mathbb{R})$ denote the space of all functions $f \in C^k(\mathbb{R})$ such that*

- (i) $fu^{2n} \in C_b(\mathbb{R})$,
- (ii) $f^{(l)}u^{n+l+1} \in C_0(\mathbb{R})$, $1 \leq l \leq k$, and
- (iii) $\widehat{f^{(k)}u^{k \vee n}} \in L^1(\mathbb{R})$.

We note the following properties of $f \in \mathfrak{Q}_n^k(\mathbb{R})$.

Proposition 2.3. *Let $n, k \in \mathbb{N}$. Then for each $f \in \mathfrak{Q}_n^k(\mathbb{R})$,*

- (i) $\widehat{f^{(l)}u^s} \in L^1(\mathbb{R})$ for $s, l \in \mathbb{Z}_{\geq 0}$ such that $s \leq n \vee l$ and $l \leq k$,
- (ii) $((fu^s)^{(l)})^\wedge \in L^1(\mathbb{R})$ for $s, l \in \mathbb{Z}_{\geq 0}$ such that $s \leq n \vee l$ and $l \leq k$.

Proof. Item (i) follows from Definition 2.2, the convolution theorem of the Fourier transform, and the well-known fact that $\hat{g} \in L^1(\mathbb{R})$ for all $g \in C^1(\mathbb{R})$ with $g, g' \in L^2(\mathbb{R})$. Item (ii) follows from (i) and the Leibniz rule. \square

Let $\mathfrak{R}_n := \{(\cdot - z)^{-l} : \text{Im}(z) \neq 0, l \in \mathbb{N}, l \geq 2n+1\}$, and let $\mathcal{S}(\mathbb{R})$ denote the class of Schwartz functions on \mathbb{R} . Then it follows from the definition of $\mathfrak{Q}_n^k(\mathbb{R})$ that

$$C_c^{k+1}(\mathbb{R}) \subset \mathfrak{Q}_n^k(\mathbb{R}) \subset C_0(\mathbb{R}), \quad \mathcal{S}(\mathbb{R}) \subset \mathfrak{Q}_n^k(\mathbb{R}), \quad \text{and} \quad \mathfrak{R}_n \subset \mathfrak{Q}_n^k(\mathbb{R}).$$

As one straightforwardly checks that $\mathfrak{Q}_n^k(\mathbb{R})$ is an algebra, arbitrary products of functions in $C_c^{n+1}(\mathbb{R}) \cup \mathcal{S}(\mathbb{R}) \cup \mathfrak{R}_n$ are in $\mathfrak{Q}_n^k(\mathbb{R})$ as well.

Remark 2.4. *For all $n, k_1, k_2 \in \mathbb{N}$, Proposition 2.3(i) implies that*

$$\mathfrak{Q}_n^{k_1}(\mathbb{R}) \subseteq \mathfrak{Q}_n^{k_2}(\mathbb{R}) \quad \text{whenever} \quad k_2 \leq k_1.$$

For proof of the following Lemma, we refer to [14, Lemma 2.4].

Lemma 2.5. *Let $n, m \in \mathbb{N} \cup \{0\}$, and let μ be a finite complex Radon measure on \mathbb{R} . For every $\epsilon \in (0, 1]$ and every $k \in \mathbb{N} \cup \{0\}$ there exists a complex Radon measure $\tilde{\mu}_{k,\epsilon}$ with*

$$\|\tilde{\mu}_{k,\epsilon}\| \leq \|u^{-1-\epsilon}\|_1 \|\mu\|$$

and

$$\int_{\mathbb{R}} g^{(n)} u^m d\mu = \int_{\mathbb{R}} g^{(n+k)} u^{m+k+\epsilon} d\tilde{\mu}_{k,\epsilon} \quad (2.1)$$

for all $g \in C^{n+k}(\mathbb{R})$ satisfying $g^{(n+l)} u^{m+l} \in C_0(\mathbb{R})$, $l = 0, \dots, k-1$ and $g^{(n+k)} u^{m+k-1} \in L^1(\mathbb{R})$.

2.2. Multiple Operator Integrals (MOI). Let $f \in C^n(\mathbb{R})$. Recall that the divided difference of order n is defined recursively as follows:

$$\begin{aligned} f^{[0]}(\lambda) &= f(\lambda), \\ f^{[n]}(\lambda_0, \lambda_1, \dots, \lambda_n) &= \begin{cases} \frac{f^{[n-1]}(\lambda_0, \lambda_1, \dots, \lambda_{n-2}, \lambda_n) - f^{[n-1]}(\lambda_0, \lambda_1, \dots, \lambda_{n-2}, \lambda_{n-1})}{\lambda_n - \lambda_{n-1}} & \text{if } \lambda_n \neq \lambda_{n-1}, \\ \frac{\partial}{\partial \lambda} f^{[n-1]}(\lambda_0, \lambda_1, \dots, \lambda_{n-2}, \lambda) \Big|_{\lambda=\lambda_{n-1}} & \text{if } \lambda_n = \lambda_{n-1}. \end{cases} \end{aligned}$$

Throughout this article we fix a complex separable Hilbert space \mathcal{H} . Let $\mathcal{B}(\mathcal{H})$ denote the algebra of bounded linear operators on \mathcal{H} . The Schatten p -norm of an operator $A \in \mathcal{B}(\mathcal{H})$ is given by

$$\|A\|_p := \left(\sum_{n \in \mathbb{N}} s_n(A)^p \right)^{1/p},$$

where the numbers $s_n(A)$ are the singular values of A . For $p \geq 1$ we denote by $\mathcal{S}^p := \mathcal{S}^p(\mathcal{H})$ the space of all Schatten p -class operators on \mathcal{H} , defined as follows:

$$\mathcal{S}^p = \{A \in \mathcal{B}(\mathcal{H}) : \|A\|_p < \infty\}.$$

We use the notation $\mathcal{S}^\infty := \mathcal{S}^\infty(\mathcal{H})$ to represent the set of all compact operators on \mathcal{H} . For any $A \in \mathcal{S}^\infty$, the norm $\|A\|_\infty$ is defined as $\max_{n \in \mathbb{N}} \{s_n(A)\} = \|A\|$. It is well-known that for $1 \leq p \leq \infty$, \mathcal{S}^p form a Banach space with respect to the Schatten norm $\|\cdot\|_p$.

Our proof employs the elegant framework of multiple operator integration. The following is a simple, yet widely applicable definition of the multiple operator integral based on [2]. Many variations of this definition are available [20].

Definition 2.6. *Let $n \in \mathbb{Z}_{\geq 0}$, let H_0, \dots, H_n be self-adjoint operators in \mathcal{H} and let $\phi : \mathbb{R}^{n+1} \rightarrow \mathbb{C}$ be such that there exists a finite measure space (Ω, μ) and measurable bounded functions $a_0, \dots, a_n : \mathbb{R} \times \Omega \rightarrow \mathbb{C}$ such that, for all $\lambda_0, \dots, \lambda_n \in \mathbb{R}$,*

$$\phi(\lambda_0, \dots, \lambda_n) = \int_{\Omega} a_0(\lambda_0, x) \cdots a_n(\lambda_n, x) d\mu(x). \quad (2.2)$$

Then, the multiple operator integral is the multilinear map $T_\phi^{H_0, \dots, H_n} : \mathcal{B}(\mathcal{H})^{\times n} \rightarrow \mathcal{B}(\mathcal{H})$ defined on $V_1, \dots, V_n \in \mathcal{B}(\mathcal{H})$ by

$$T_\phi^{H_0, \dots, H_n}(V_1, \dots, V_n)\psi := \int_{\Omega} a_0(H_0, x) V_1 a_1(H_1, x) \cdots V_n a_n(H_n, x) \psi d\mu(x),$$

for all $\psi \in \mathcal{H}$. By the results of [2], the above Bochner integral exists and, as suggested by the notation, $T_\phi^{H_0, \dots, H_n}$ depends on ϕ , but not on the measure space (Ω, μ) and the functions a_0, \dots, a_n chosen to represent ϕ through (2.2).

As symbols ϕ we almost exclusively use divided differences $\phi = f^{[n]}$ of sufficiently regular functions $f : \mathbb{R} \rightarrow \mathbb{C}$. In this case the multiple operator integral is explicit and admits a useful Schatten bound.

Theorem 2.7. *Let $n \in \mathbb{N}$, H_0, \dots, H_n be self-adjoint operators in \mathcal{H} and let $f \in C^n(\mathbb{R})$ be such that $\widehat{f^{(n)}} \in L^1(\mathbb{R})$.*

(1) *For all $V_1, \dots, V_n \in \mathcal{B}(\mathcal{H})$ and all $\psi \in \mathcal{H}$ we have*

$$T_{f^{[n]}}^{H_0, \dots, H_n}(V_1, \dots, V_n)\psi := \int_{\mathbb{R}} \int_{\Delta_n} \widehat{f^{(n)}}(t) e^{its_0 H_0} V_1 e^{its_1 H_1} \cdots V_n e^{its_n H_n} \psi ds dt,$$

where the simplex $\Delta_n = \{s = (s_0, \dots, s_n) \in \mathbb{R}_{\geq 0}^{n+1} : s_0 + \cdots + s_n = 1\}$ is endowed with the flat measure with total measure $1/n!$.

(2) *For all $\alpha, \alpha_j \in [1, \infty]$ with $\frac{1}{\alpha_1} + \cdots + \frac{1}{\alpha_n} = \frac{1}{\alpha}$,*

$$\|T_{f^{[n]}}^{H_0, \dots, H_n}\|_{S^{\alpha_1} \times \cdots \times S^{\alpha_n} \rightarrow S^\alpha} \leq \frac{1}{n!} \|\widehat{f^{(n)}}\|_1. \quad (2.3)$$

Proof. See [11, Lemma 1] and [11, Lemma 4]. \square

The following algebraic property of the multiple operator integral is proven in [13, Thm 3.10(i)].

Lemma 2.8. *Let $n \in \mathbb{N}$, let H_0, \dots, H_n be self-adjoint operators in \mathcal{H} , and let $V_1, \dots, V_n \in \mathcal{B}(\mathcal{H})$. For every $f \in C^n$ satisfying $\widehat{f^{(n)}}, \widehat{(fu)^{(n)}}, \widehat{f^{(n-1)}} \in L^1(\mathbb{R})$, we have*

$$\begin{aligned} T_{f^{[n]}}^{H_0, \dots, H_n}(V_1, \dots, V_n) &= T_{(fu)^{[n]}}^{H_0, \dots, H_n}(V_1, V_2, \dots, V_n (H_n - i)^{-1}) \\ &\quad - T_{f^{[n-1]}}^{H_0, \dots, H_{n-1}}(V_1, \dots, V_{n-1}) V_n (H_n - i)^{-1}. \end{aligned} \quad (2.4)$$

The following estimate is a consequence of [20, Theorem 4.4.7 and Remark 4.4.2], first proven for $\tilde{H} = H$ in [15, Theorem 5.3 and Remark 5.4].

Theorem 2.9. *Let $k \in \mathbb{N}$ and let $\alpha, \alpha_1, \dots, \alpha_k \in (1, \infty)$ satisfy $\frac{1}{\alpha_1} + \cdots + \frac{1}{\alpha_k} = \frac{1}{\alpha}$. Let H, \tilde{H} be self-adjoint operators in \mathcal{H} . Assume that $V_\ell \in \mathcal{S}^{\alpha_\ell}$, $1 \leq \ell \leq k$. Then there exists $c_{\alpha, k} > 0$ such that*

$$\|T_{f^{[k]}}^{\tilde{H}, H, \dots, H}(V_1, V_2, \dots, V_k)\|_\alpha \leq c_{\alpha, k} \|f^{(k)}\|_\infty \prod_{1 \leq \ell \leq k} \|V_\ell\|_{\alpha_\ell} \quad (2.5)$$

for every $f \in C_b^k(\mathbb{R})$ such that $\widehat{f^{(k)}} \in L^1(\mathbb{R})$.

The following result is a consequence of [20, Theorems 4.4.6 and 5.3.5].

Theorem 2.10. *Let $n \in \mathbb{N}$ and let $f \in C^n(\mathbb{R})$ be such that $\widehat{f^{(k)}} \in L^1(\mathbb{R})$, $k = 0, 1, \dots, n$. Let H be a self-adjoint operator in \mathcal{H} and let V be a self-adjoint operator in $\mathcal{B}(\mathcal{H})$. Then, the Gâteaux derivative $\frac{d^k}{dt^k} f(H + tV)|_{t=0}$ exists in the uniform operator topology and admits the multiple operator integral representation*

$$\frac{1}{k!} \frac{d^k}{ds^k} f(H + sV)|_{s=t} = T_{f^{[k]}}^{H+tV, \dots, H+tV}(V, \dots, V), \quad 1 \leq k \leq n. \quad (2.6)$$

The map $t \mapsto \frac{1}{k!} \frac{d^k}{ds^k} f(H + sV)|_{s=t}$ is continuous in the strong operator topology and, when $V \in \mathcal{S}^k$, in the trace norm of \mathcal{S}^1 .

Throughout this paper, we shall refer to the following hypothesis.

Hypothesis 2.11. *Let $n \in \mathbb{N}$, and let $n \geq 2$. Let H_0 be a self-adjoint (unbounded) operator in \mathcal{H} , and let V be a bounded self-adjoint operator on \mathcal{H} such that*

$$V(H_0 - i)^{-p} \in \mathcal{S}^{n/p}, \quad p \in \{1, \dots, n\}.$$

We remark that Hypothesis 2.11 for (n, H_0, V) implies Hypothesis 2.11 for $(n+1, H_0, V)$ *et cetera*. We may therefore restrict to $n \geq 2$, which will help us avoid some ugly case distinctions. For improved function classes for the case $n = 1$, see [13, Theorem 4.1].

3. KREIN TRACE FORMULA

In this section, we will obtain the Krein trace formula (first-order spectral shift formula) and the associated spectral shift function under Hypothesis 2.11. The following two lemmas are essential to prove our main results in this section.

Lemma 3.1. *Assume Hypothesis 2.11. Then there exists a sequence $\{V_k\}_{k \in \mathbb{N}}$ of finite rank self-adjoint operators such that*

(i) *$SOT - \lim_{k \rightarrow \infty} V_k = V$, where SOT means strong operator topology,*

(ii) *$\|\cdot\|_{n/p} - \lim_{k \rightarrow \infty} V_k(H_0 - i)^{-p} = V(H_0 - i)^{-p}$ for $1 \leq p \leq n$,*

(iii) *$\|V_k\| \leq \|V\|$, and*

(iv) *$\|V_k(H_0 - i)^{-p}\|_{n/p} \leq \|V(H_0 - i)^{-p}\|_{n/p}$ for $1 \leq p \leq n$.*

Proof. Let $E_{H_0}(\cdot)$ be the spectral measure of H_0 . Let $k \in \mathbb{N}$, and let $P_k = E_{H_0}((-k, k))$. Then the following statements are true.

- $SOT - \lim_{k \rightarrow \infty} P_k = I$, and
- $SOT - \lim_{k \rightarrow \infty} P_k V P_k = V$.

Note that

$$P_k V P_k ((H_0 - i)^{-n} P_k + P_k^\perp) = P_k V P_k (H_0 - i)^{-n} P_k = P_k V (H_0 - i)^{-n} P_k \in \mathcal{S}^1. \quad (3.1)$$

By functional calculus it follows that $((H_0 - i)^{-n}P_k + P_k^\perp)$ is boundedly invertible. Hence, from (3.1), we have

$$P_k V P_k = P_k V (H_0 - i)^{-n} P_k ((H_0 - i)^{-n} P_k + P_k^\perp)^{-1} \in \mathcal{S}^1. \quad (3.2)$$

For each fixed $k \in \mathbb{N}$, by the spectral theorem there exists a sequence $\{E_{P_k V P_k}^l\}_{l \in \mathbb{N}}$ of finite rank projections such that

$$E_{P_k V P_k}^l P_k V P_k = P_k V P_k E_{P_k V P_k}^l, \text{ and } \|\cdot\|_1 - \lim_{l \rightarrow \infty} E_{P_k V P_k}^l P_k V P_k = P_k V P_k.$$

Therefore, there exists a strictly increasing sequence of natural numbers $\{l_k\}_{k \in \mathbb{N}}$ such that

$$\left\| E_{P_k V P_k}^{l_k} P_k V P_k - P_k V P_k \right\|_1 < \frac{1}{k}.$$

Let $V_k = E_{P_k V P_k}^{l_k} P_k V P_k$. Then

- (i) $SOT - \lim_{k \rightarrow \infty} V_k = V$,
- (ii)

$$\begin{aligned} & \|V_k(H_0 - i)^{-p} - V(H_0 - i)^{-p}\|_{n/p} \\ & \leq \|V_k(H_0 - i)^{-p} - P_k V P_k(H_0 - i)^{-p}\|_{n/p} + \|P_k V(H_0 - i)^{-p} P_k - V(H_0 - i)^{-p}\|_{n/p} \\ & \leq \frac{1}{k} + \|P_k V(H_0 - i)^{-p} P_k - V(H_0 - i)^{-p}\|_{n/p} \longrightarrow 0 \text{ as } k \rightarrow \infty, \end{aligned}$$

- (iii) $\|V_k\| \leq \|V\|$,

- (iv) $\|V_k(H_0 - i)^{-p}\|_{n/p} = \|E_{P_k V P_k}^{l_k} P_k V(H_0 - i)^{-p} P_k\|_{n/p} \leq \|V(H_0 - i)^{-p}\|_{n/p}.$

This completes the proof. \square

Lemma 3.2. *Let n, H_0, V satisfy Hypothesis 2.11. Let $(R, W) \in \{(0, V), (0, V_k), (V_k, V - V_k), (V_k, V_l - V_k)\}$, where $\{V_k\}_{k \in \mathbb{N}}$ is given by Lemma 3.1. For $t \in [0, 1]$, we denote*

$$H^t := H_0 + R + tW.$$

For $1 \leq p \leq n$, define

$$\alpha_{n, V, H_0} := \max_{1 \leq p \leq n} \|V(H_0 - i)^{-p}\|_{n/p}. \quad (3.3)$$

Then

- (a) $(H^t - i)^{-p} W, W(H^t - i)^{-p} \in \mathcal{S}^{n/p}$, and
- (b) $\|(H^t - i)^{-p} W\|_{n/p} = \|W(H^t - i)^{-p}\|_{n/p} \leq 2^p(1 + \|V\|) \alpha_{n, W, H_0} \max_{\ell=0, p-1} \{\alpha_{n, V, H_0}^\ell\}.$

Proof. Note that the equality in (b) follows directly from $(H^t + i)^{-l} W = (W(H^t - i)^{-(m+1)})^*$, as this operator equals $(H^t - i)^{-l} W$ up to a unitary.

We prove the remaining results using mathematical induction on p . Let $p = 1$. Let $V_t = H^t - H_0$. Note that

$$W[(H^t - i)^{-1} - (H_0 - i)^{-1}] = -W(H_0 - i)^{-1} V_t (H^t - i)^{-1}.$$

By noting that $\|V_t\| \leq \|V\|$, we obtain from the above identity that

$$\|W(H^t - i)^{-1}\|_n \leq (1 + \|V\|) \|W(H_0 - i)^{-1}\|_n. \quad (3.4)$$

Thus (a) and (b) are true for $p = 1$.

Suppose (a) and (b) are true for $p = 1, 2, \dots, m$ for some $m \in \{1, \dots, n-1\}$. We will prove that (a) and (b) also hold for $p = m+1$. Indeed, note that

$$W[(H^t - i)^{-(m+1)} - (H_0 - i)^{-(m+1)}] = - \sum_{l=0}^m W(H^t - i)^{-(l+1)} V_t (H_0 - i)^{-(m+1-l)}.$$

Therefore, using Hölder's inequality from the above identity we conclude

$$\begin{aligned} & \|W(H^t - i)^{-(m+1)}\|_{n/(m+1)} \\ & \leq \|W(H_0 - i)^{-(m+1)}\|_{n/(m+1)} + \|W(H^t - i)^{-1}\|_n \|V(H_0 - i)^{-m}\|_{n/m} \\ & \quad + \sum_{l=1}^m \|W(H^t - i)^{-l}\|_{n/l} \|V(H_0 - i)^{-(m+1-l)}\|_{n/(m+1-l)}. \end{aligned} \quad (3.5)$$

Let us write $C_p = (1 + \|V\|) \alpha_{n,W,H_0} \max_{\ell=0, p-1} \{\alpha_{n,V,H_0}^\ell\}$, and consider the three terms on the right-hand side of (3.5). The first term is bounded by C_{m+1} , and thanks to (3.4) the second term is also bounded by C_{m+1} . By induction hypothesis, the third term is bounded by $\sum_{l=1}^m 2^l C_l \|V(H_0 - i)^{-(m+1-l)}\|_{n/(m+1-l)} \leq \sum_{l=1}^m 2^l C_{m+1}$. In total, we obtain

$$\|W(H^t - i)^{-(m+1)}\|_{n/(m+1)} \leq 2^{m+1} C_{m+1},$$

proving the induction step. The lemma follows. \square

The following theorem gives a key estimate to establish Krein's trace formula under Hypothesis 2.11.

Theorem 3.3. *Let n, H_0, V satisfy Hypothesis 2.11. Let $(R, W) \in \{(0, V), (V_l, V - V_l), (V_l, V_k - V_l)\}$, where $\{V_k\}_{k \in \mathbb{N}}$ is given by Lemma 3.1. Let $t \in [0, 1]$, and denote $H^t = H_0 + R + tW$. Then for each $f \in \mathfrak{Q}_n^1(\mathbb{R})$, we have*

$$T_{f^{[1]}}^{H^t, H^t}(W) = T_{(fu^n)^{[1]}}^{H^t, H^t}(W(H^t - i)^{-n}) - \sum_{k=0}^{n-1} (fu^k)(H^t)W(H^t - i)^{-(k+1)} \in \mathcal{S}^1, \quad (3.6)$$

and each of the above $n+1$ summands is \mathcal{S}^1 -continuous in t . Moreover,

$$\left| \text{Tr} \left(T_{(fu^n)^{[1]}}^{H^t, H^t}(W(H^t - i)^{-n}) \right) \right| \leq \| (fu^n)^{(1)} \|_\infty \|W(H^t - i)^{-n}\|_1, \quad (3.7)$$

$$\left| \text{Tr} \left(\sum_{k=0}^{n-1} (fu^k)(H^t)W(H^t - i)^{-(k+1)} \right) \right| \leq n \|fu^{n-1}\|_\infty \|W(H^t - i)^{-n}\|_1. \quad (3.8)$$

Proof. Let $f \in \mathfrak{Q}_n^1(\mathbb{R})$. Then by repeated applications of Proposition 2.3 and Lemma 2.8, we get

$$\begin{aligned} T_{f^{[1]}}^{H^t, H^t}(W) &= T_{(fu)^{[1]}}^{H^t, H^t}(W(H^t - i)^{-1}) - f(H^t)W(H^t - i)^{-1} \\ &= T_{(fu^2)^{[1]}}^{H^t, H^t}(W(H^t - i)^{-2}) - (fu)(H^t)W(H^t - i)^{-2} - f(H^t)W(H^t - i)^{-1} \end{aligned}$$

$$= T_{(fu^n)^{[1]}}^{H^t, H^t}(W(H^t - i)^{-n}) - \sum_{k=0}^{n-1} (fu^k)(H^t)W(H^t - i)^{-(k+1)}. \quad (3.9)$$

As Proposition 2.3 implies $\widehat{(fu^n)^{(1)}} \in L^1(\mathbb{R})$, an application of Theorem 2.7 and Lemma 3.2 yields

$$T_{(fu^n)^{[1]}}^{H^t, H^t}(W(H^t - i)^{-n}) \in \mathcal{S}^1. \quad (3.10)$$

Since $t \mapsto (H^t - i)^{-1}$ is strongly continuous, it follows (cf. [16, Theorem VIII.20]) that $t \mapsto e^{isH^t}$ is strongly continuous. Hence, Theorem 2.7 and Lemma 3.2 also yield \mathcal{S}^1 -continuity of the above operator in t .

Since $fu^{2n} \in C_b(\mathbb{R})$ by Definition 2.2(i), for each $0 \leq k \leq n-1$, we have

$$(fu^k)(H^t)W = (fu^{k+n})(H^t)(H^t - i)^{-n}W \in \mathcal{S}^1. \quad (3.11)$$

Thus, the identity (3.9) and the properties (3.10) and (3.11) together establish (3.6). The \mathcal{S}^1 -continuity of (3.11) in t follows from Lemma 3.2 and strong continuity of $t \mapsto (fu^{k+n})(H^t)$, where the latter fact follows (cf. [16, Theorem VIII.20]) from the inclusion $fu^{k+n} \in C_b(\mathbb{R})$ and the strong continuity of $t \mapsto (H^t - i)^{-1}$.

By a standard property of the double operator integral (which can be derived from Theorem 2.7), we have

$$\mathrm{Tr} \left(T_{(fu^n)^{[1]}}^{H^t, H^t}(W(H^t - i)^{-n}) \right) = \mathrm{Tr} \left((fu^n)^{(1)}(H^t) (W(H^t - i)^{-n}) \right).$$

Hence, by utilizing Hölder's inequality, we derive the estimate (3.7).

By using (3.11) and the cyclicity of the trace we conclude

$$\begin{aligned} \mathrm{Tr} \left((fu^k)(H^t)W(H^t - i)^{-(k+1)} \right) &= \mathrm{Tr} \left((H^t - i)^{-(k+1)}(fu^k)(H^t)W \right) \\ &= \mathrm{Tr} \left((fu^{n-1})(H^t)(H^t - i)^{-n}W \right). \end{aligned} \quad (3.12)$$

Applying Hölder's inequality to the right-hand side of (3.12), combined with the invariance of $\|\cdot\|_1$ under adjoints, results in the estimate (3.8). This concludes the proof. \square

We briefly recall the classical result that the first-order spectral shift function exists for perturbations of trace class. This result is due to M. G. Krein [8], see also [23, Theorem 8.3.3].

Theorem 3.4 (Krein). *Let H_0 be a self-adjoint operator in \mathcal{H} , and let $V = V^* \in \mathcal{S}^1$. Then there exists an integrable function ξ such that for all $f \in C^1(\mathbb{R})$ satisfying $\widehat{f'} \in L^1(\mathbb{R})$ (slightly more generally, $\widehat{f'}$ is only assumed to be a finite complex measure) we have*

$$\mathrm{Tr}(f(H_0 + V) - f(H_0)) = \int_{-\infty}^{\infty} \xi(\lambda) f'(\lambda) d\lambda.$$

The following is the main result of this section.

Theorem 3.5. *Let n, H_0, V satisfy Hypothesis 2.11. Then there exists a locally integrable function $\eta_{1,H_0,V} \in L^1\left(\mathbb{R}, \frac{d\lambda}{(1+|\lambda|)^{n+\epsilon}}\right)$ such that*

$$\int_{\mathbb{R}} \frac{|\eta_{1,H_0,V}(\lambda)|}{(1+|\lambda|)^{n+\epsilon}} d\lambda \leq n 2^n (2n+3+2(n+1)\epsilon^{-1})(1+\|V\|) \max_{\ell=1,n} \{\alpha_{n,V,H_0}^\ell\} \quad (3.13)$$

for all $\epsilon \in (0, 1]$, where α_{n,V,H_0} is given by (3.3), and, moreover,

$$\mathrm{Tr}(f(H_0 + V) - f(H_0)) = \int_{\mathbb{R}} f'(\lambda) \eta_{1,H_0,V}(\lambda) d\lambda \quad (3.14)$$

for every $f \in \mathfrak{Q}_n^1(\mathbb{R})$. The locally integrable function $\eta_{1,H_0,V}$ is determined by (3.14) uniquely up to an additive constant.

Proof. Let $f \in \mathfrak{Q}_n^1(\mathbb{R})$. Let $H_t = H_0 + tV, t \in [0, 1]$. By the fundamental theorem of calculus and using Theorem 2.10 we get the (a priori pointwise) integral

$$f(H_0 + V) - f(H_0) = \int_0^1 T_{f^{[1]}}^{H_t, H_t}(V) dt. \quad (3.15)$$

We now apply Theorem 3.3 to the above. By (3.7), (3.8), Lemma 3.2, and \mathcal{S}^1 -continuity of the trace we may take the trace of the integral of (3.6) and find

$$\mathrm{Tr}(f(H_0 + V) - f(H_0)) \quad (3.16)$$

$$= \mathrm{Tr} \left(\int_0^1 T_{(fu^n)^{[1]}}^{H_t, H_t}(V(H_t - i)^{-n} dt \right) - \mathrm{Tr} \left(\int_0^1 \sum_{k=0}^{n-1} (fu^k)(H_t) V(H_t - i)^{-(k+1)} dt \right) \quad (3.17)$$

$$= \alpha_1((fu^n)^{(1)}) + \alpha_2(fu^{n-1}) \quad (3.18)$$

for certain linear functionals α_1, α_2 defined on suitable subspaces of $C_0(\mathbb{R})$, and all $f \in \mathfrak{Q}_n^1(\mathbb{R})$. The above integrands are \mathcal{S}^1 -continuous by Theorem 3.3, which shows that the above expressions are well defined and we may interchange trace with integral everywhere.

From (3.7) and (3.8) it follows that α_1 and α_2 are continuous in the supremum norm, and hence we may extend α_1 and α_2 to continuous functionals on $C_0(\mathbb{R})$. By the Riesz–Markov representation theorem, there exist two complex Borel measures μ_1, μ_2 such that

$$\begin{aligned} \mathrm{Tr}(f(H_0 + V) - f(H_0)) &= \int_{\mathbb{R}} (fu^n)^{(1)}(\lambda) d\mu_1(\lambda) + \int_{\mathbb{R}} (fu^{n-1})(\lambda) d\mu_2(\lambda) \\ &= \int_{\mathbb{R}} \left(f^{(1)}(\lambda) u^n(\lambda) + n f(\lambda) u^{n-1}(\lambda) \right) d\mu_1(\lambda) + \int_{\mathbb{R}} (fu^{n-1})(\lambda) d\mu_2(\lambda), \end{aligned} \quad (3.19)$$

and with total variation bounded by (3.7), (3.8) and Lemma 3.2(b) as

$$\begin{aligned} \|\mu_1\|, \|\mu_2\| &\leq n \|V(H_t - i)^{-n}\|_1 \\ &\leq n 2^n (1 + \|V\|) \max_{\ell=1,n} \{\alpha_{n,V,H_0}^\ell\}. \end{aligned} \quad (3.20)$$

For $\epsilon \in (0, 1]$, applying Lemma 2.5 to (3.19) ensures the existence of a measure ν_ϵ such that

$$\mathrm{Tr}(f(H_0 + V) - f(H_0)) = \int_{\mathbb{R}} f^{(1)}(\lambda) d\nu_\epsilon(\lambda) \quad (f \in \mathfrak{Q}_n^1(\mathbb{R})) \quad (3.21)$$

with

$$\int_{\mathbb{R}} |u^{-n-\epsilon}(\lambda)| d|\nu_\epsilon|(\lambda) \leq \|\mu_1\| + \|u^{-1-\epsilon}\|_1 (n\|\mu_1\| + \|\mu_2\|).$$

As $\frac{1}{1+|\lambda|} \leq |u(\lambda)^{-1}|$, this further implies that

$$\begin{aligned} \int_{\mathbb{R}} \frac{d|\nu_\epsilon|(\lambda)}{(1+|\lambda|)^{n+\epsilon}} &\leq \|\mu_1\| + \|u^{-1-\epsilon}\|_1 (n\|\mu_1\| + \|\mu_2\|) \\ &\leq n 2^n (2n + 3 + 2(n+1)\epsilon^{-1})(1 + \|V\|) \max_{\ell=1, n} \{\alpha_{n, V, H_0}^\ell\}, \end{aligned} \quad (3.22)$$

where we have estimated $\|u^{-1-\epsilon}\|_1 = 2 \int_0^1 |u^{-1-\epsilon}| + 2 \int_1^\infty |u^{-1-\epsilon}| \leq 2 + 2\epsilon^{-1}$.

Next, we will demonstrate that the measure ν_ϵ in (3.21) is absolutely continuous with respect to the Lebesgue measure. Let $\{V_k\}_{k \in \mathbb{N}}$ be the sequence given in Lemma 3.1. Therefore, by Theorem 3.4, there exists a sequence of integrable functions $\{\eta_{k,1}\}_{k \in \mathbb{N}}$ on \mathbb{R} such that, for all $f \in C_c^\infty(\mathbb{R})$,

$$\mathrm{Tr}(f(H_0 + V_k) - f(H_0)) = \int_{\mathbb{R}} f^{(1)}(\lambda) \eta_{k,1}(\lambda) d\lambda. \quad (3.23)$$

If $V_{k,l} := V_k - V_l$, then from (3.23), subsequently applying a basic property of the multiple operator integral, and Theorem 3.3, we gather

$$\begin{aligned} &\left| \int_{\mathbb{R}} f^{(1)}(\lambda) (\eta_{k,1} - \eta_{l,1})(\lambda) d\lambda \right| \\ &= |\mathrm{Tr}(f(H_0 + V_k) - f(H_0 + V_l))| \\ &= \left| \int_0^1 \mathrm{Tr} \left(T_{f^{[1]}}^{H_0 + V_l + tV_{k,l}, H_0 + V_l + tV_{k,l}}(V_{k,l}) \right) dt \right| \\ &\leq (\|(fu^n)^{(1)}\|_\infty + n\|fu^{n-1}\|_\infty) \int_0^1 dt \|V_{k,l}(H_0 + V_l + tV_{k,l} - i)^{-n}\|_1. \end{aligned} \quad (3.24)$$

Thanks to Lemma 3.2, from (3.24) we obtain

$$\begin{aligned} &\left| \int_{\mathbb{R}} f^{(1)}(\lambda) (\eta_{k,1} - \eta_{l,1})(\lambda) d\lambda \right| \\ &\leq (\|(fu^n)^{(1)}\|_\infty + n\|fu^{n-1}\|_\infty) 2^n (1 + \|V\|) \alpha_{n, V_{k,l}, H_0} \max_{p=0, n-1} \{\alpha_{n, V, H_0}^p\}. \end{aligned} \quad (3.25)$$

Let $f \in C_c^\infty((-a, a))$ for some fixed $a > 0$. Then using Lemma 2.1, from (3.25) we get

$$\begin{aligned} &\left| \int_{\mathbb{R}} f^{(1)}(\lambda) (\eta_{k,1} - \eta_{l,1})(\lambda) d\lambda \right| \\ &\leq (1 + a + 4na) (1 + a)^{n-1} \|f^{(1)}\|_\infty 2^n (1 + \|V\|) \alpha_{n, V_{k,l}, H_0} \max_{p=0, n-1} \{\alpha_{n, V, H_0}^p\}. \end{aligned} \quad (3.26)$$

Therefore, subsequently using (3.26) and Lemma 3.1(ii) implies that

$$\begin{aligned} \|\eta_{k,1} - \eta_{l,1}\|_{L^1((-a,a))} &= \sup_{\|f^{(1)}\|_\infty \leq 1} \left| \int_{\mathbb{R}} f^{(1)}(\lambda) (\eta_{k,1} - \eta_{l,1})(\lambda) d\lambda \right| \\ &\leq (1 + a + 4na) (1 + a)^{n-1} 2^n (1 + \|V\|) \alpha_{n,V_k,l,H_0} \max_{p=0,n-1} \{\alpha_{n,V,H_0}^p\} \\ &\longrightarrow 0 \text{ as } k, l \rightarrow \infty. \end{aligned}$$

Thus $\{\eta_{k,1}\}_{k \in \mathbb{N}}$ is a Cauchy sequence in $L_{loc}^1(\mathbb{R})$; we let η_1 be its $L_{loc}^1(\mathbb{R})$ -limit. By a calculation completely analogous to (3.24)-(3.25), we conclude for $f \in C_c^\infty(\mathbb{R})$ that

$$\begin{aligned} \text{Tr}(f(H_0 + V) - f(H_0)) &= \lim_{k \rightarrow \infty} \text{Tr}(f(H_0 + V_k) - f(H_0)) \\ &= \lim_{k \rightarrow \infty} \int_{\mathbb{R}} f^{(1)}(\lambda) \eta_{k,1}(\lambda) d\lambda \\ &= \int_{\mathbb{R}} f^{(1)}(\lambda) \eta_1(\lambda) d\lambda. \end{aligned} \tag{3.27}$$

Now combining (3.21) and (3.27) we have the following equality

$$\int_{\mathbb{R}} f^{(1)}(\lambda) d\nu_\epsilon(\lambda) = \int_{\mathbb{R}} f^{(1)}(\lambda) \eta_1(\lambda) d\lambda \quad \text{for all } f \in C_c^\infty(\mathbb{R}). \tag{3.28}$$

Let $d\mu(\lambda) = \eta_1(\lambda) - d\nu_\epsilon(\lambda)$. Then, it follows from (3.28) that

$$\int_a^b f^{(1)}(\lambda) d\mu(\lambda) = 0 \quad \text{for all } f \in C_c^\infty(\mathbb{R}). \tag{3.29}$$

Now consider the distribution T_μ defined by

$$T_\mu(\phi) := \int_a^b \phi d\mu(\lambda)$$

for every $\phi \in C_c^\infty(\mathbb{R})$. By (3.29) and the definition of the derivative of a distribution, $T_\mu^{(1)} = 0$. Hence by [1, Theorem 3.10 and Example 2.21], $d\mu(\lambda) = c d\lambda$ for some constant c . Consequently, $\eta_1 \in L_{loc}^1(\mathbb{R})$ satisfying (3.27) is unique up to an additive constant. In particular, we can assume $d\nu_\epsilon(\lambda) = \eta_1(\lambda) d\lambda$, and obtain

$$\int_{\mathbb{R}} \frac{|\eta_1(\lambda)|}{(1 + |\lambda|)^{n+\epsilon}} d\lambda = \int_{\mathbb{R}} \frac{d|\nu_\epsilon|(\lambda)}{(1 + |\lambda|)^{n+\epsilon}}.$$

By setting $\eta_{1,H_0,V} = \eta_1$, the above equality together with (3.21)-(3.22) completes the proof. \square

4. HIGHER ORDER TRACE FORMULA

The aim of this section is to provide all higher order spectral shift functions corresponding to a pair of self adjoint operators (H_0, V) satisfying the Hypothesis 2.11.

The following lemma is essential to reach our goal.

Lemma 4.1. *Let n, H_0, V satisfy Hypothesis 2.11. Then for $1 \leq k \leq n-1$, and $f \in \mathfrak{Q}_n^k(\mathbb{R})$,*

$$T_{f^{[k]}}^{H_0, \dots, H_0}(V, \dots, V) \in \mathcal{S}^1,$$

and

$$\begin{aligned} & T_{f^{[k]}}^{H_0, \dots, H_0}(V, \dots, V) \\ &= \sum_{l=0}^k \sum_{\substack{j_1 + \dots + j_{l+1} = k \\ j_1, \dots, j_l \geq 1, j_{l+1} \geq 0}} \sum_{r=0}^{\min(n-k, l)} (-1)^{k-l+r} \sum_{\substack{p_0 \geq 0, p_1, \dots, p_r \geq 1 \\ p_0 + \dots + p_r = n-k}} T_{(fu^{n-k+l-r})^{[l-r]}}^{H_0, \dots, H_0}(\tilde{V}^{j_1}, \dots, \tilde{V}^{j_{l-r}}) \\ & \quad \times (H_0 - i)^{-p_0} \tilde{V}^{j_{l-r+1}} \dots (H_0 - i)^{-p_r} \tilde{V}^{j_{l+1}}. \end{aligned} \quad (4.1)$$

We recall that by standard convention, a product $B_1 \dots B_m$ for $m < 1$ is 1, a list (B_1, \dots, B_m) for $m < 1$ is the empty list $()$, and, by definition, $T_{g^{[0]}}^{H_0}() = g(H_0)$.

Proof. It was obtained in [13, Equation 39] that for any $k \geq 1$, and $f \in \mathfrak{Q}_n^k(\mathbb{R})$,

$$T_{f^{[k]}}^{H_0, \dots, H_0}(V, \dots, V) = \sum_{l=0}^k \sum_{\substack{j_1 + \dots + j_{l+1} = k \\ j_1, \dots, j_l \geq 1, j_{l+1} \geq 0}} (-1)^{k-l} T_{(fu^l)^{[l]}}^{H_0, \dots, H_0}(\tilde{V}^{j_1}, \dots, \tilde{V}^{j_l}) \tilde{V}^{j_{l+1}}. \quad (4.2)$$

We may now apply [11, Lemma 9] to each of the terms above, for $s = n - k$. Indeed, for all $l \in \{0, \dots, k\}$, from Proposition 2.3 it follows that fu^l is in the desired function class $\mathcal{W}^{n-k, l}$, in the notation of [11, Section 3]. The lemma follows immediately. \square

The following theorem provides the existence of measures which are necessary to obtain the spectral shift functions of any order in the main theorem of this section.

Theorem 4.2. *Let n, H_0, V satisfy Hypothesis 2.11. Let $\epsilon \in (0, 1]$. Then for each natural number $k \geq 2$, there exists a complex Borel measure $\mu_{k, \epsilon}$ such that*

$$\mathrm{Tr} \left(T_{f^{[k]}}^{H_0, \dots, H_0}(V, \dots, V) \right) = \int_{\mathbb{R}} f^{(k)}(\lambda) u^{k \vee n + \epsilon}(\lambda) d\mu_{k, \epsilon}(\lambda) \quad (4.3)$$

for every $f \in \mathfrak{Q}_n^k(\mathbb{R})$, with

$$\|\mu_{k, \epsilon}\| \leq c_{n, k, \epsilon} \alpha_{n, V, H_0}^k, \quad (4.4)$$

where $c_{n, k, \epsilon}$ is some positive constant, and α_{n, V, H_0} is given by (3.3).

Proof. For $2 \leq k < n$: By Lemma 4.1, we have the identity (4.1). Next we show that each term in the right hand side of (4.1) is a trace class operator and hence we can estimate the trace of those terms. Assuming the notations involved in (4.1), we note the following.

- (i) Recalling that $T_{g^{[0]}}^{H_0}() = g(H_0)$, we notice that the summands corresponding to $r = l$ are of the form of (a minus sign times)

$$(fu^{n-k})(H_0)(H_0 - i)^{-p_0} \tilde{V}^{j_1} \dots (H_0 - i)^{-p_l} \tilde{V}^{j_{l+1}}.$$

Employing a similar argument as given for (3.11), $fu^{2n} \in C_b(\mathbb{R})$ implies that

$$(fu^{n-k})(H_0)\tilde{V}^{j_1} \dots (H_0 - i)^{-p_l}\tilde{V}^{j_{l+1}} \in \mathcal{S}^1,$$

and using the cyclicity property of the trace along with Hölder's inequality for Schatten norms, this yields

$$\begin{aligned} & \left| \text{Tr} \left((fu^{n-k})(H_0)(H_0 - i)^{-p_0}\tilde{V}^{j_1}(H_0 - i)^{-p_1} \dots \tilde{V}^{j_l}(H_0 - i)^{-p_l}\tilde{V}^{j_{l+1}} \right) \right| \\ &= \left| \text{Tr} \left((fu^{n-k})(H_0)\tilde{V}^{j_1}(H_0 - i)^{-p_1} \dots \tilde{V}^{j_l}(H_0 - i)^{-p_l}\tilde{V}^{j_{l+1}}(H_0 - i)^{-p_0} \right) \right| \\ &\leq \|fu^{n-k}\|_\infty \alpha_{n,V,H_0}^k, \end{aligned} \quad (4.5)$$

where α_{n,V,H_0} is given by (3.3).

- (ii) For the rest of the terms in the summation of right hand side of (4.1): Now Theorem 2.7 and Hölder's inequality for Schatten norms together implies

$$T_{(fu^{n-k+l-r})^{[l-r]}}^{H_0, \dots, H_0}(\tilde{V}^{j_1}, \dots, \tilde{V}^{j_{l-r}})(H_0 - i)^{-p_1}\tilde{V}^{j_{l-r+1}} \dots (H_0 - i)^{-p_{r+1}}\tilde{V}^{j_{l+1}} \in \mathcal{S}^1.$$

If the product on the right of the multiple operator integral is nontrivial, then by applying Hölder's inequality for Schatten norms and using Theorem 2.9 we obtain

$$\begin{aligned} & \left| \text{Tr} \left(T_{(fu^{n-k+l-r})^{[l-r]}}^{H_0, \dots, H_0}(\tilde{V}^{j_1}, \dots, \tilde{V}^{j_{l-r}})(H_0 - i)^{-p_1}\tilde{V}^{j_{l-r+1}} \dots (H_0 - i)^{-p_{r+1}}\tilde{V}^{j_{l+1}} \right) \right| \\ &\leq d_{n,k} \|fu^{n-k+l-r}\|_\infty \alpha_{n,V,H_0}^k, \end{aligned} \quad (4.6)$$

where $d_{n,k}$ is a constant depending only on n and k .

The case where $r = 0$ and $j_{l+1} = 0$ takes a bit more care, because in the corresponding term $\text{Tr}(T_{(fu^{n-k+l})^{[l]}}^{H_0, \dots, H_0}(\tilde{V}^{j_1}, \dots, \tilde{V}^{j_l})(H_0 - i)^{-(n-k)})$ the resolvent is not necessarily summable, and Theorem 2.9 is not applicable for $\alpha = 1$. Luckily, this case is similarly covered by [14, Lemma 2.9(i)] after commuting the resolvent inside the multiple operator integral.

Thus, (4.1) together with (i) and (ii) implies

$$T_{f^{[k]}}^{H_0, \dots, H_0}(V, \dots, V) \in \mathcal{S}^1.$$

Having noted the remarks above, the estimates (4.5) and (4.6), the Hahn-Banach theorem, and the Riesz–Markov representation theorem together imply that there exist complex Borel measures $\mu_{n,k,l,r}$, $0 \leq l \leq k$, $0 \leq r \leq \min\{n-k, l\}$ such that

$$\|\mu_{n,k,l,r}\| \leq \tilde{d}_{n,k} \alpha_{n,V,H_0}^k,$$

where $\tilde{d}_{n,k}$ is a constant depending only on n and k , and

$$\begin{aligned} & \text{Tr} \left(T_{f^{[k]}}^{H_0, \dots, H_0}(V, \dots, V) \right) \\ &= \sum_{l=0}^k \sum_{r=0}^{\min\{n-k, l\}} \int_{\mathbb{R}} (fu^{n-k-r})^{(l-r)}(\lambda) d\mu_{n,k,l,r}(\lambda) \end{aligned}$$

$$\begin{aligned}
&= \sum_{l=0}^k \sum_{r=0}^{\min\{n-k, l\}} \sum_{s=0}^{l-r} \binom{l-r}{s} \int_{\mathbb{R}} f^{(l-r-s)}(\lambda) (u^{n-k-r})^{(s)}(\lambda) d\mu_{n,k,l,r}(\lambda) \\
&= \sum_{\substack{0 \leq l \leq k \\ 0 \leq r \leq \min\{n-k, l\} \\ 0 \leq s \leq l-r}} \binom{l-r}{s} \frac{(n-k-r)!}{(n-k-r-s)!} \int_{\mathbb{R}} f^{(l-r-s)}(\lambda) u^{n-k-r-s}(\lambda) d\mu_{n,k,l,r}(\lambda).
\end{aligned} \tag{4.7}$$

Thanks to Lemma 2.5 and Equation (4.7), there exists a measure $\mu_{k,\epsilon}$ that satisfies (4.3) and (4.4).

For $k \geq n$: Thanks to Equation (4.2), Lemma 2.9, and Hölder's inequality, we can argue similarly to the case $k < n$ and conclude the existence of a measure $\mu_{k,\epsilon}$ satisfying (4.3) and (4.4). This completes the proof. \square

Now, we proceed to present and demonstrate the central theorem of this section. In the theorem and its proof, the Taylor remainder is denoted

$$\mathcal{R}_{k,f,H_0}(V) := f(H_0 + V) - \sum_{m=0}^{k-1} \frac{d^m}{dt^m} \Big|_{t=0} f(H_0 + tV).$$

Theorem 4.3. *Let n, H_0, V satisfy Hypothesis 2.11. Then for each fixed $k \in \mathbb{N}$, there exists a locally integrable function $\eta_k := \eta_{k,H_0,V}$ such that*

$$\int_{\mathbb{R}} \frac{|\eta_k(\lambda)|}{(1+|\lambda|)^{n+k+\epsilon}} d\lambda \leq C_{n,k,\epsilon} (1 + \|V\|) \max_{\ell=1, k \vee n} \{\alpha_{n,V,H_0}^\ell\} \tag{4.8}$$

for all $\epsilon \in (0, 1]$, where $C_{n,k,\epsilon}$ are some positive constants depending only on n, k, ϵ , and α_{n,V,H_0} is given by (3.3), and, moreover,

$$\text{Tr}(\mathcal{R}_{k,f,H_0}(V)) = \int_{\mathbb{R}} f^{(k)}(\lambda) \eta_k(\lambda) d\lambda \tag{4.9}$$

for each $f \in \mathfrak{Q}_n^k(\mathbb{R})$. The locally integrable function η_k is determined by (4.9) uniquely up to a polynomial summand of degree at most $k-1$.

Proof. We prove (4.8) and (4.9) using the principle of mathematical induction on k . The base case, $k=1$, of the induction follows from Theorem 3.5. Now we assume (4.8) and (4.9) are true for $k \in \{1, 2, \dots, m\}$ for some $m \in \mathbb{N}$. Let $\epsilon \in (0, 1]$. Let $f \in \mathfrak{Q}_n^{m+1}(\mathbb{R})$. Thanks to Theorem 2.10 and Theorem 4.2, there exists a finite complex Borel measure $\mu_{m,\epsilon}$ such that

$$\begin{aligned}
\text{Tr}(\mathcal{R}_{m+1,f,H_0}(V)) &= \text{Tr}(\mathcal{R}_{m,f,H_0}(V)) - \text{Tr}\left(T_{f^{[m]}}^{H_0, \dots, H_0}(V, \dots, V)\right) \\
&= \int_{\mathbb{R}} f^{(m)}(\lambda) \eta_m(\lambda) d\lambda - \int_{\mathbb{R}} f^{(m)}(\lambda) u^{m \vee n + \epsilon}(\lambda) d\mu_{m,\epsilon}(\lambda).
\end{aligned} \tag{4.10}$$

Next, by performing integration by parts on the right hand side of (4.10), we obtain

$$\text{Tr}(\mathcal{R}_{m+1,f,H_0}(V)) = \int_{\mathbb{R}} f^{(m)}(\lambda) \eta_m(\lambda) d\lambda - \int_{\mathbb{R}} f^{(m)}(\lambda) u^{m \vee n + \epsilon}(\lambda) d\mu_{m,\epsilon}(\lambda) d\lambda$$

$$\begin{aligned}
&= \int_{\mathbb{R}} f^{(m+1)}(\lambda) \left(- \int_0^\lambda \eta_m(t) dt \right) d\lambda + \int_{\mathbb{R}} f^{(m+1)}(\lambda) u^{m \vee n + \epsilon}(\lambda) \mu_{m, \epsilon}((-\infty, \lambda]) d\lambda \\
&\quad - (m \vee n + \epsilon) \int_{\mathbb{R}} f^{(m+1)}(\lambda) \left(\int_0^\lambda u^{m \vee n + \epsilon - 1}(t) \mu_{m, \epsilon}((-\infty, t]) dt \right) d\lambda \\
&= \int_{\mathbb{R}} f^{(m+1)}(\lambda) \eta_{m+1}(\lambda) d\lambda,
\end{aligned} \tag{4.11}$$

where

$$\eta_{m+1}(\lambda) = u^{m \vee n + \epsilon}(\lambda) \mu_{m, \epsilon}((-\infty, \lambda]) - \int_0^\lambda \eta_m(t) dt - (m \vee n + \epsilon) \int_0^\lambda u^{m \vee n + \epsilon - 1}(t) \mu_{m, \epsilon}((-\infty, t]) dt. \tag{4.12}$$

Now it follows from (4.4), (4.12), Lemma 2.5, and the induction hypothesis, that

$$\int_{\mathbb{R}} \frac{|\eta_{m+1}(\lambda)|}{(1 + |\lambda|)^{n+m+1+\epsilon}} d\lambda \leq \tilde{C}_{n, m+1, \epsilon} (1 + \|V\|) \max_{\ell=1, k \vee n} \{\alpha_{n, V, H_0}^\ell\}, \tag{4.13}$$

for every $\epsilon \in (0, 1]$, where $\tilde{C}_{n, m+1, \epsilon}$ is some positive constant. This completes the induction on k . The uniqueness of η_k can be established in a similar manner as we did in Theorem 3.5. Indeed, suppose η_k and ξ_k satisfy (4.9), and let $\gamma_k = \eta_k - \xi_k$. Then from (4.9), we conclude

$$\int_{\mathbb{R}} f^{(k)}(\lambda) \gamma_k(\lambda) d\lambda = 0 \quad \text{for all } f \in C_c^\infty(\mathbb{R}). \tag{4.14}$$

Now consider the distribution T_{γ_k} defined by

$$T_{\gamma_k}(\phi) = \int_{\mathbb{R}} \phi(\lambda) \gamma_k(\lambda) d\lambda$$

for every $\phi \in C_c^\infty(\mathbb{R})$. By (4.14) and the definition of the derivative of a distribution, $T_{\gamma_k}^{(k)} = 0$. Hence by [1, Theorem 3.10 and Example 2.21], $\gamma_k(\lambda) d\lambda = f_k(\lambda) d\lambda$ for some polynomial f_k of degree at most $k-1$. Consequently, $\eta_k \in L_{loc}^1(\mathbb{R})$ satisfying (4.9) is unique up to a polynomial summand of degree at most $k-1$. Hence, we can assert that the estimates (4.9) hold true for all $\epsilon \in (0, 1]$. This completes the proof. \square

Remark 4.4. By following our proof, yet replacing the application of Theorem 3.4 with an application of [22], under slightly stronger conditions on H_0 and V one obtains a more moderate growth of the spectral shift function, namely

$$\int_{\mathbb{R}} \frac{|\eta_k(\lambda)|}{(1 + |\lambda|)^{k-1+\epsilon}} d\lambda < \infty \tag{4.15}$$

for all $k \in \mathbb{N}$ and all $\epsilon > 0$. It is however unclear if one can obtain bounds similar to (4.8) of our main theorem.

5. EXAMPLES

In this section we collect some known classes of examples of self-adjoint operators H_0 and V satisfying our assumptions, namely $V \in \mathcal{B}(\mathcal{H})$ and

$$V(H_0 - i)^{-p} \in \mathcal{S}^{n/p} \quad (p = 1, \dots, n). \quad (5.1)$$

We merely scratch the surface of the collection of examples; more can be found in [13, 17, 21, 18], and references therein.

A result about Schatten class membership typically comes along with a bound on the Schatten norm, and by our main result these bounds imply estimates on the weighted integral norm of the spectral shift functions, for each specific situation. For brevity we omit these bounds, as they can be found in the respective cited references. Such bounds are called Cwikel estimates and Birman–Solomyak estimates, and form an active area of research [9, 10].

5.1. Dirac and Laplace operators on Euclidean spaces. If, for suitable functions f, g , we have $V = f(x)$ and $(H_0 - i)^{-p} = g(-i\nabla)$, where x is the canonical vector of multiplication operators on $L^2(\mathbb{R}^d)$ and ∇ is the vector whose components are the operators $\{\partial_i\}_{i=1}^d$, then conditions for (5.1) to hold can be found in [5] and [18, Chapter 4]. As shown in the latter reference, such results typically require different proofs for $p \leq n/2$ and $p > n/2$. Moreover, the spaces

$$\ell^p(L^q)(\mathbb{R}^d) := \left\{ f : \mathbb{R}^d \rightarrow \mathbb{C} \text{ measurable} : \sum_{k \in \mathbb{Z}^d} (\|f \upharpoonright_{(0,1)^d+k}\|_{L^q((0,1)^d+k)})^p < \infty \right\},$$

for $p \in [1, \infty)$ and $q \in [1, \infty]$, are frequently useful.

Two especially important examples of H_0 are the the Laplacian $H_0 = \Delta = \sum_{k=1}^d \frac{\partial^2}{\partial x_k^2}$ in $L^2(\mathbb{R}^d)$ and the Dirac operator $H_0 = D$, defined as

$$D := -i \sum_{k=1}^d e_k \otimes \frac{\partial}{\partial x_k}$$

in $\mathbb{C}^N \otimes L^2(\mathbb{R}^d)$, where $e_1, \dots, e_d \in M_N(\mathbb{C})$ are self-adjoint matrices satisfying $e_k e_l + e_l e_k = 2\delta_{k,l}$ for all $k, l \in \{1, \dots, d\}$. Here $N = 2^{d/2}$ if d is even and $N = 2^{(d+1)/2}$ if d is odd. The following result is proven in [19, Theorem III.4].

Theorem 5.1 ([19]). *Let $m, d \in \mathbb{N}$ and $f \in \ell^1(L^2)(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ be real-valued, with associated multiplication operator M_f .*

(1) *If $n > d$, then*

$$(I \otimes M_f)(D - i)^{-p} \in \mathcal{S}^{n/p}, \quad p = 1, \dots, n.$$

(2) *If $n > \frac{d}{2}$, then*

$$M_f(\Delta - i)^{-p} \in \mathcal{S}^{n/p}, \quad p = 1, \dots, n.$$

5.2. The Hodge–Dirac operator on a Riemannian space. Part of the above result generalizes to suitable Riemannian manifolds, in the sense of the following theorem. For its proof we refer to [21, Equation (3.11)], which in that paper is proven in order to obtain [21, Theorem 3.4.1] by specializing to $p = d$.

Theorem 5.2 ([21]). *Let (X, g) be a second countable d -dimensional complete smooth Riemannian manifold. Let $\Omega_c^k(X)$ be the space of smooth compactly supported differential k -forms, and \mathcal{H}_k its completion with respect to the natural Hilbert space structure defined by g . Defined in the Hilbert space $\mathcal{H} := \bigoplus_{k=0}^d \mathcal{H}_k$, let D_g be the associated Hodge–Dirac operator. For $f \in C_c^\infty(X)$, let M_f denote the associated multiplication operator on \mathcal{H} . Then, for all $p \geq 1$ we have*

$$M_f(D_g - i)^{-p} \in \mathcal{S}^{d/p, \infty}.$$

Our main result thus becomes applicable by setting $n = d + 1$, and, independently of dimension, implies existence of spectral shift functions of all orders for Dirac operators on manifolds as in the above theorem.

In light of the above, it is reasonable to expect a generalization of Theorem 5.1 to exist for suitable Riemannian manifolds, but this is well beyond the scope of this text. A similar remark applies to matrix-valued differential operators and pseudodifferential operators.

5.3. Noncommutative Geometry. Noncommutative geometry is a prime motivator for this paper, as it too is involved with proving results about geometrical objects – such as differential operators – without reference to their underlying space. The typical benefit for the noncommutative geometer is that the thus obtained results hold for more general objects which may not be geometrical, but may still be useful to describe physics.

A first application of the spectral shift function is to the theory of spectral flow [2], as used in index theory. Typically, one takes

$$H_0 = D, \quad V = u[D, u^*], \quad \text{so that } H_0 + V = uDu^*,$$

for a unitary u . More generally, perturbations in gauge theory take the form $V = \sum_{j=1}^n a_j[D, b_j]$, when a_j and b_j are elements of a ‘smooth’ algebra $\mathcal{A} \subseteq \mathcal{B}(\mathcal{H})$. Higher order spectral shift functions also provide analytical information on the Taylor remainders of the spectral action, which may be useful for the description of classical and quantum field theories [6, 12].

The following result shows that our Hypothesis 2.11 holds for a wide class of nonunital spectral triples. It is a special case of [17, Proposition 10].

Theorem 5.3 ([17]). *Let $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ be a smooth local (d, ∞) -summable spectral triple in the sense of [17, Definition 7]. Then for all $a \in \mathcal{B}(\mathcal{H})$ satisfying $a\phi = \phi a = a$ for some $\phi \in \mathcal{A}_c$ we have*

$$a(\mathcal{D} - i)^{-p} \in \mathcal{S}^{(d+1)/p}, \quad p = 1, \dots, d + 1.$$

In particular cases one may expect strengthenings of the above result. This is indeed the case for the Moyal plane \mathbb{R}_θ^d , which is a prominent motivating example of a nonunital spectral triple. The result below follows from [9], as is written out explicitly below, for convenience of the reader.

Theorem 5.4 ([9]). *Let $d \geq 2$ be even, $V \in W^{d,1}(\mathbb{R}_\theta^d)$, $h \in \mathbb{N}$, and set $n := \lfloor \frac{d}{h} \rfloor + 1$. For all $p \in \{1, \dots, n\}$, if we assume that $g \in \ell^{d/(hp),\infty}(L^\infty)(\mathbb{R}^d)$, then we have that $Vg(-i\nabla_\theta) \in \mathcal{S}^{n/p}$. In particular,*

$$V(D_\theta - i)^{-p} \in \mathcal{S}^{n/p} \quad (p \in \{1, \dots, n\}, \quad n = d + 1),$$

and

$$V(\Delta_\theta - i)^{-p} \in \mathcal{S}^{n/p} \quad (p \in \{1, \dots, n\}, \quad n = d/2 + 1),$$

where D_θ and Δ_θ are the noncommutative Dirac and Laplace operators (cf. [9, p. 29]).

Proof. A combination of [9, Proposition 6.15(iii)], [9, Definition 6.14], and [9, Definition 6.7] yields $V \in W^{d,q}(\mathbb{R}_\theta^d)$ and thus $V \in L^q(\mathbb{R}_\theta^d)$ for all $q \geq 1$.

We define $n := \lfloor \frac{d}{h} \rfloor + 1$ and let $p \in \{1, \dots, n\}$, $h \in \mathbb{N}$, and $g \in \ell^{d/(hp),\infty}(L^\infty)(\mathbb{R}^d) \subseteq L^{d/(hp),\infty}(\mathbb{R}^d)$.

First suppose that $p \leq n/2$. Setting $q := d/(hp)$, a case distinction shows that $q \geq 1$. We find $g \in L^{q,\infty}(\mathbb{R}^d)$ and $V \in L^q(\mathbb{R}_\theta^d)$, hence [9, Lemma 2.3] implies that $V \otimes g \in \mathcal{L}_{q,\infty}(\mathcal{N})$, the noncommutative $L^{q,\infty}$ -space of the von Neumann algebra $\mathcal{N} := L^\infty(\mathbb{R}_\theta^d) \otimes L^\infty(\mathbb{R}^d)$. In particular, since $n/p > d/(hp)$, we have $V \otimes g \in (\mathcal{L}_{n/p} \cap \mathcal{L}_\infty)(\mathcal{N})$. We note that $\frac{n}{p} \geq 2$ by assumption, so $L^{n/p} \cap L^\infty$ is an interpolation space for (L^2, L^∞) . Therefore, [9, Theorem 7.2] gives

$$Vg(-i\nabla_\theta) \in \mathcal{L}_{n/p}(\mathcal{B}(L^2(\mathbb{R}^d))) = \mathcal{S}^{n/p}.$$

Now suppose that $p > n/2$. Set $q := \frac{n}{p} \in [1, 2)$ and note that $g \in \ell^q(L^\infty)(\mathbb{R}^d)$. We have $V \in W^{d,q}(\mathbb{R}_\theta^d)$ by Sobolev embedding, hence [9, Theorem 7.7] implies that

$$Vg(-i\nabla_\theta) \in \mathcal{L}_q(\mathcal{B}(L^2(\mathbb{R}^d))) = \mathcal{S}^{n/p},$$

completing the proof of the first statement.

To prove the second statement, one employs the functions g on \mathbb{R}^d defined by

$$g(x) := (\gamma \cdot x - i)^{-p},$$

$$g(x) := (\|x\|^2 - i)^{-p},$$

where γ is the vector of Clifford matrices (cf. [9, Definition 6.17]). We indeed have $g \in \ell^{d/(hp),\infty}(L^\infty)(\mathbb{R}^d)$ for $h = 1$, $h = 2$, respectively. In the former case g is matrix-valued and this inclusion can be interpreted component-wise, the finite component $\mathbb{C}^{2^{\lfloor d/2 \rfloor}}$ of the Hilbert space factors through, and one indeed obtains $V(D_\theta - i)^{-p} \in \mathcal{S}^{n/p}(\mathbb{C}^{2^{\lfloor d/2 \rfloor}} \otimes L^2(\mathbb{R}^d))$ as well as $V(\Delta_\theta - i)^{-p} \in \mathcal{S}^{n/p}(L^2(\mathbb{R}^d))$. \square

REFERENCES

- [1] M. A. Al-Gwaiz, *Theory of distributions*. Monographs and Textbooks in Pure and Applied Mathematics, 159. Marcel Dekker, Inc., New York, 1992, XII+257 pp.
- [2] N. A. Azamov, A. L. Carey, P. G. Dodds, F. Sukochev, *Operator integrals, spectral shift function, and spectral flow*, Canad. J. Math. 61 (2009), no. 2, 241–263.
- [3] A. Connes, *Noncommutative geometry*. 1994.
- [4] A. Connes, H. Moscovici, *The local index formula in noncommutative geometry*. Geom. Funct. Anal. 5 (1995), 174–243.

- [5] M. Cwikel, *Weak type estimates for singular values and the number of bound states of Schrödinger operators*, Ann. Math. 106 (1977), 93–100.
- [6] B. Iochum, C. Levy, D. Vassilevich, *Spectral action beyond the weak-field approximation*, Commun. Math. Phys. 316 (2012), 595–613.
- [7] L. S. Koplienko, *The trace formula for perturbations of nonnuclear type*. Sibirsk. Mat. Zh. 25 (1984), no. 5, 62–71; English translation, Siberian Math. J. 25 (1984), 735–743.
- [8] M. G. Krein, *On the trace formula in perturbation theory*, Mat. Sbornik N.S. 33 (1953), no. 75, 597–626.
- [9] G. Levitina, F. Sukochev, D. Zanin, *Cwikel estimates revisited*, P. Lond. Math. Soc. 120 (2020), no. 2, 265–304.
- [10] E. McDonald, R. Ponge, *Cwikel estimates and negative eigenvalues of Schrödinger operators on noncommutative tori*, J. Math. Phys. 63 (2022) no. 4.
- [11] T. D. H. van Nuland, W. D. van Suijlekom, *Cyclic cocycles in the spectral action*, J. Noncommut. Geom. 16 (2022), no. 3, 1103–1135.
- [12] T. D. H. van Nuland, W. D. van Suijlekom, *One-loop corrections to the spectral action*, J. High Energy Phys. (2022), no. 5, Paper No. 078, 14 pp.
- [13] T. D. H. van Nuland, A. Skripka, *Spectral shift for relative Schatten class perturbations*, J. Spectr. Theory 12 (2022), no. 4, 1347–1382.
- [14] T. D. H. van Nuland, A. Skripka *Higher-order spectral shift function for resolvent comparable perturbations*, to appear in J. Operator Theory, <https://arxiv.org/abs/2211.03330>.
- [15] D. Potapov, A. Skripka, F. Sukochev, *Spectral shift function of higher order*, Invent. Math. 193 (2013), no. 3, 501–538.
- [16] M. Reed, B. Simon, *Methods of modern mathematical physics. I. Functional analysis*, Academic Press, New York-London, 1972.
- [17] A. C. Rennie, *Summability for nonunital spectral triples*, K-Theory 31 (2004), no. 1, 71–100.
- [18] B. Simon, *Trace ideals and their applications*. Mathematical surveys and monographs, vol. 120, American mathematical society, Providence, RI, 2005.
- [19] A. Skripka, *Lipschitz estimates for functions of Dirac and Schrödinger operators*, J. Math. Phys. 62 (2021), no. 1, Paper No. 013506, 28 pp.
- [20] A. Skripka, A. Tomsikova, *Multilinear Operator Integrals: Theory and Applications*, Lecture Notes in Math. 2250, Springer International Publishing, 2019, XI+192 pp.
- [21] F. Sukochev, D. Zanin, *The Connes character formula for locally compact spectral triples*, Astérisque 445 (2023), 150 pp.
- [22] D. R. Yafaev, *A trace formula for the Dirac operator*, Bull. London Math. Soc. 37 (2005), no. 6, 908–918.
- [23] D. R. Yafaev, *Mathematical scattering theory. General theory*. Translations of Mathematical Monographs, 105. American Mathematical Society, Providence, RI, 1992.

A. C., DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY GUWAHATI, GUWAHATI, 781039, INDIA

Email address: arupchatt@iitg.ac.in, 2003arupchattopadhyay@gmail.com

T. v. N., TU DELFT, EWI/DIAM, 2600 GA DELFT, THE NETHERLANDS

Email address: t.d.h.vannuland@tudelft.nl, teunvn@gmail.com

C. P., DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE, 560012, INDIA

Email address: chandan.pradhan2108@gmail.com