ON POSSIBLE EMBEDDINGS OF THE STANDARD MODELS OF PARTICLE PHYSICS AND GRAVITY IN E_8

ROBERT ARNOTT WILSON

ABSTRACT. I investigate the structure of E_8 under the action of the subalgebra/subgroup $A_1 + G_2 + C_3$, as a potential route to unification of the fundamental forces of nature into a single algebraic structure. The particular real form $E_{8(-24)}$ supports a decomposition into compact G_2 plus split $A_1 + C_3$, which allows a restriction from G_2 to SU(3) for the strong force, together with split $SL_2(\mathbb{R})$ to break the symmetry of the weak interaction and give mass to the intermediate vector bosons. The factor C_3 contains various copies of the Lorentz group $SL_2(\mathbb{C})$ and extends the 'spacetime' symmetries to the full group of symplectic symmetries of real 3 + 3-dimensional phase space.

Restricting G_2 to the Standard Model SU(3) extends C_3 to A_5 , in the real form SU(3,3), acting on a complex phase space that includes both momentum and current. There is then a natural restriction from SU(3,3) to SO(3,3), describing the action of $SL_4(\mathbb{R})$ on phase space. The resulting action of $SL_4(\mathbb{R})$ on E_8 includes tensors that are equivalent to the stress-energy tensor, the Ricci tensor and the Riemann tensor, and therefore permits the formalism of general relativity to be developed inside $E_{8(-24)}$. The model then suggests unexpected and perhaps subtle ways in which general relativity and particle physics may be forced to modify each other, in order to produce a unified theory.

1. INTRODUCTION

A number of E_8 models of fundamental physics have been proposed in recent years [1, 2, 3, 4, 5], but none of them has been sufficiently compelling to persuade large numbers of people that they are useful. The key issue is how to split up the symmetries of E_8 to get something that looks like the Standard Model, and in particular, how to do this in a reasonably 'natural' way. In addition there are a number of technical issues which cause a lot of trouble, particularly to do with complex structures and chirality, and with implementing three generations of fermions when there appears on the face of it to be only enough room for two [6].

The approach to 'naturality' taken in [1] is to take the Freudenthal–Tits 'magic square' [7, 8, 9] as a guide. The Lie structure of the magic square is

and most emphasis has been put on the fourth row and the fourth column, where the exceptional Lie groups are found. However, the other entries in the table are also interesting. For example, the route taken in [1] from top left to bottom right goes via A_2 , A_5 and E_6 in the second row (see also [10, 11]).

Date: First draft: 20th April 2024; this version; 6th May 2024.

The corresponding 'magic square' for 2×2 matrices is

where I use the notation B and D to emphasise that these are all spin groups, with $B_n = Spin(2n + 1)$ and $D_n = Spin(2n)$. However, I would like to draw particular attention to the isomorphisms between spin groups (types B and D) and unitary (type A) and symplectic (type C) groups:

(3)

$$B_1 = A_1,$$

 $D_2 = A_1 + A_1,$
 $B_2 = C_2,$
 $D_3 = A_3.$

All of the exceptional isomorphisms listed here arise from the triality automorphism of D_4 , which links together all the groups in the table that are not in the fourth row or fourth column.

The isomorphism $B_1 = A_1$ gives the basic fact that $Spin(3) \cong SU(2)$, which is the foundation of quantum mechanics (QM), while $D_2 = A_1 + A_1$ extends this to $Spin(3,1) \cong SL(2,\mathbb{C})$ for relativistic QM. The isomorphism $D_3 = A_3$ is needed to explain the complex structure of the Dirac algebra, as the complex Clifford algebra of Minkowski spacetime, since the smallest real Lie algebra that contains all the Dirac matrices, including γ_5 , is of this type. This isomorphism also plays a significant role in various Grand Unified Theories (GUTs) from 1974 onwards [12, 13, 14], in various real forms including $Spin(6) \cong SU(4)$, $Spin(4, 2) \cong SU(2, 2)$ and $Spin(3, 3) \cong SL_4(\mathbb{R})$. The isomorphism $B_2 = C_2$ similarly lies at the heart of the AdS/CFT correspondence [15], and describes the Lie algebra of the even part of the Dirac algebra.

Extending then to the 3×3 case, we extend C_2 to C_3 , which looks a lot more 'natural' than extending B_2 to C_3 . In this note, therefore, I concentrate on the third row of the magic square [16, 17], and especially the first group, $Sp_6(\mathbb{R})$, that has a classical interpretation as the symmetry group of phase space [18]. The second group extends this to a complex phase space, so that both momentum and current can be included, as well as the position of both mass and charge. This extends the symplectic group to $Sp_6(\mathbb{C}) \cong SU(3,3)$, whose centralizer in E_8 is $SU(3) \times SL_2(\mathbb{R})$. The latter is a real form of the gauge group $SU(3) \times SU(2)$ of the strong and weak nuclear forces, and I shall argue in this paper that the group SU(3,3) describes both classical and quantum forms of both electromagnetism and gravity.

2. Embedding C_2 in D_8

In the semi-split version of the magic square, both the compact and split real forms of C_2 occur, but because I want to use C_2 to implement the Dirac algebra, I want the split form, that is the group $Spin(3,2) \cong Sp_4(\mathbb{R})$. Embedding into Spin(12,4) we see the centralizer Spin(9,2), which we can split into three pieces, Spin(6), Spin(3) and Spin(2), if we want to get the Standard Model gauge groups SU(3), SU(2) and U(1). However, the reason for this splitting will not become clear until we consider the embedding of C_3 in E_8 . In the notation of [1, 19], we need to choose a copy of the split quaternions \mathbb{H}' in the split octonions, say the copy with basis U, K, L, KL. Then the corresponding copy of Spin(3,2) acts on the indices u := 1, U, K, L, KL, leaving IL, JL for Spin(2), and i, j, k, il, jl, kl for Spin(6) and/or SU(3), so that Spin(3) acts on l, I, J. In particular we see some symmetry-breaking for Spin(3), generated by X_{lI} , X_{lJ} and $D_{I,J}$, already in the notation. Now for the Dirac part of the algebra, the labels u, U, K, L, KL correspond to the matrices $\gamma_1, \gamma_2, \gamma_3, \gamma_0$ and γ_5 respectively, and the products of pairs of these gamma matrices generate the algebra $\mathfrak{so}(3, 2)$:

(4)
$$X_1, X_K, D_K, X_L, D_L, D_{K,L}, X_{KL}, D_{KL}, D_{K,KL}, D_{L,KL}, X_{KL}, D_{K,KL}, D_{L,KL}, X_{KL}, X_{KL}$$

in which the first row contains the rotations in $SL_2(\mathbb{C})$, and the second row contains the boosts, while the third row contains a Lorentzian 4-vector.

Since this algebra is quaternionic rather than octonionic, its elements can be written as ordinary 2×2 anti-Hermitian quaternion matrices, which make it easier to understand the structure. The Xs are off-diagonal, the single-index Ds are diagonal traceless, and the double-index Ds add the imaginary traces:

(5)
$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & K \\ K & 0 \end{pmatrix}, \begin{pmatrix} K & 0 \\ 0 & -K \end{pmatrix}, \\ \begin{pmatrix} 0 & L \\ L & 0 \end{pmatrix}, \begin{pmatrix} L & 0 \\ 0 & -L \end{pmatrix}, \begin{pmatrix} KL & 0 \\ 0 & KL \end{pmatrix}, \\ \begin{pmatrix} 0 & KL \\ KL & 0 \end{pmatrix}, \begin{pmatrix} KL & 0 \\ 0 & -KL \end{pmatrix}, \begin{pmatrix} L & 0 \\ 0 & L \end{pmatrix}, \begin{pmatrix} K & 0 \\ 0 & K \end{pmatrix}$$

These matrices can be regarded as a split quaternionic version of the Pauli matrices, which occur in the first row, in the mathematicians' anti-Hermitian convention rather than the physicists' Hermitian convention. The last matrix in the third row is the scalar matrix that is the product of the three Pauli matrices (Hermitian convention).

3. Extending to C_3

To extend from C_2 to C_3 we have to generalise the X terms to Y and Z, and the D terms to E:

(6)
$$Y_1, Y_K, Y_L, Y_{KL}, Z_1, Z_K, Z_L, Z_{KL}, E_K, E_L, E_{KL}.$$

Equivalently, we extend from 2×2 matrices to 3×3 , to obtain a split quaternionic version of the Gell-Mann matrices [20]. It turns out that the 'scalar' 2×2 matrices extend to traceless 3×3 matrices, in the same way that the Gell-Mann matrices are traceless. In the notation of [19], this condition is enforced by the identity:

$$(7) D_p + E_p + F_p = 0$$

for all single index p, here K, L and KL. I make no claims as to how these Gell-Mann matrices should be interpreted, or whether they have anything to do with the Gell-Mann matrices used in QCD [21].

In fact, the compact part of this group $Sp_6(\mathbb{R})$ is a copy of U(3) generated by all the elements that have an even number of copies of L in their labels:

(8)
$$X_1, X_K, Y_1, Y_K, Z_1, Z_K, D_K, E_K, D_{L,KL}$$

where again the double-index Ds add an imaginary trace to the matrices. As matrices, these are just the anti-Hermitian matrices over the complex subalgebra \mathbb{C} of \mathbb{H}' . There are also two copies of $GL_3(\mathbb{R})$ obtained by replacing K by L or KL (or indeed any linear combination of the two):

(9)
$$X_1, X_L, Y_1, Y_L, Z_1, Z_L, D_L, E_L, D_{K,KL}; X_1, X_{KL}, Y_1, Y_{KL}, Z_1, Z_{KL}, D_{KL}, E_{KL}, D_{K,L}.$$

These are anti-Hermitian matrices over the respective copies of the split complex numbers \mathbb{C}' in \mathbb{H}' . Leaving off the double-index Ds restricts to $SL_3(\mathbb{R})$, and the subtle relationships between these two copies of $SL_3(\mathbb{R})$ and SU(3) will play an important role in this paper.

4. The centralizer of C_3

The centralizer of $Sp_6(\mathbb{R})$ in E_8 is a group of type $A_1 + G_2$, in which the copy of G_2 is compact, acting on the indices i, j, k, l, il, jl, kl, and the copy of A_1 is split, acting on the indices I, J, IL, JL in one of its chiral spinor (weak isospin?) representations. This copy of A_1 is obviously not the same as the copy, acting on l, I, J, that we suggested in Section 2 simply by looking in D_8 . However, we effectively chose a copy of $\mathfrak{su}(2) + \mathfrak{u}(1)$ acting on l, I, J, IL, JL, so that by 'mixing' $D_{I,J}$ with $D_{IL,JL}$ we obtain one of the elements of the centralizer. To get the rest we need to replace X_{lI} and X_{lJ} by one combination of $D_{I,IL}$ and $D_{J,JL}$ and another combination of $D_{I,JL}$.

In other words, the conversion between these two copies of A_1 , one of which is compact and the other split, is very reminiscent of the 'symmetry-breaking' of the weak SU(2) in the Standard Model, that converts from a 'primordial' massless SU(2) to $SL_2(\mathbb{R})$ via the complexification $SL_2(\mathbb{C})$, in order to give the intermediate vector bosons non-zero masses. Thus the E_8 model provides a fundamental mathematical reason for this symmetry-breaking, namely enforcement of the condition that the gauge group must commute not only with D_2 , as the Coleman–Mandula theorem [22] requires, or with C_2 , as the embedding in D_8 requires, but with the whole of C_3 .

To see the details of how these two copies of A_1 are related to each other, we can work in the group they generate, which is another copy of Spin(3, 2), acting on the labels l, I, J, IL, JL, and commuting with the first copy. The ten dimensions of this group are represented by

(10)
$$X_{lI}, X_{lJ}, D_{I,J}, D_{IL,JL}, X_{IIL}, X_{IJL}, X_{IJL}, D_{I,IL}, D_{J,IL}, D_{J,IL}, D_{J,JL}$$

Since we are genuinely using the octonions at this point, it is not possible to write these elements of the algebra as ordinary matrices. However, the notation of [19] may help to visualise what is going on. The generators of the two copies of A_1 are related as follows (where the signs are determined by the chirality, or equivalently by the embedding in G_2):

(11)

$$D_{I,J} \rightarrow D_{I,J} - D_{IL,JL}$$

$$X_{lI} \rightarrow D_{I,IL} - D_{J,JL}$$

$$X_{lJ} \rightarrow D_{J,IL} + D_{I,JL}.$$

Turning now to the remaining factor, which is presumably related to the strong force, we compare the group Spin(6) that appears in the centralizer of Spin(3,2), with the group G_2 that appears in the centralizer of $Sp_6(\mathbb{R})$. After replacing the original (unbroken symmetry) copy of A_1 by the chiral copy on I, J, IL, JL, we no longer require the label l for the weak interaction, which can be added to the strong force, to extend the gauge group SU(3) acting on 3 + 3 colours and anti-colours to G_2 acting on 7 'colours'. This extension is reminiscent of the Pati–Salam model, which uses SU(4) for four colours and four anti-colours, but is group-theoretically completely different.

Notice also that we have a chiral pair of left-handed and right-handed copies of $SL_2(\mathbb{R})$, so that there is a close parallel between the two models:

(12)
$$SU(4) \times SU(2)_L \times SU(2)_R$$
$$G_2 \times SL_2(\mathbb{R})_L \times SL_2(\mathbb{R})_R.$$

Generators for $SL_2(\mathbb{R})_R$ in this case are obtained by changing the signs in (11). But this group $SL_2(\mathbb{R})_R$ is not contained in $Sp_6(\mathbb{R})$, so we do not find it useful, and shall not use it. In Section 5 we consider a different type of 'right-handed' $SL_2(\mathbb{R})$, whose generators are given in (22), which is contained in $Sp_6(\mathbb{R})$, and has more useful symmetry properties.

Here we have a total of 20 degrees of freedom, compared to 21 in the Pati–Salam model. Therefore our full model is of type $C_3 + A_1 + G_2$, based on the group

(13)
$$Sp_6(\mathbb{R}) \times SL_2(\mathbb{R}) \times G_2$$

with the first factor generalising the Lorentz group, the second factor representing a real form of weak SU(2), and the third factor generalizing strong SU(3).

5. Extending to A_5

The occurrence of G_2 in the decomposition $A_1+G_2+C_3$, rather than A_2 , that we would expect for the strong force, suggests that we should move to the second group in the third row of the magic square, of type A_5 , and the associated decomposition $A_1 + A_2 + A_5$ of E_8 . The particular real forms that arise are

(14)
$$SL_2(\mathbb{R}) \times SU(3) \times SU(3,3),$$

which is analogous to the decomposition

(15)
$$SU(2) \times SL_3(\mathbb{R}) \times SL_3(\mathbb{H})$$

studied in [1, Section II.C]. I propose these different real forms as a potentially closer match to the Standard Model, since the compact group SU(3) is more suitable for massless gluons, while the split group $SL_2(\mathbb{R})$ has two boosts that are suitable for masses of the intermediate vector bosons. This is because mass is usually introduced by complexifying the compact gauge group, precisely in order to generate boosts. To extend $Sp_6(\mathbb{R})$ to SU(3,3) we add 14 dimensions, all involving the complex structure *l*. The analogous 2×2 extension is from Spin(3,2) acting on u, U, K, L, KL to Spin(4,2) acting on l, u, U, K, L, KL, so that the five new elements are

Since we are not really using the octonions here, all these elements can be written as 2×2 anti-Hermitian matrices over

(17)
$$\mathbb{C} \otimes \mathbb{H}' = \langle u, l \rangle \otimes \langle U, K, L, KL \rangle$$

so that the new matrices are

(18)
$$\begin{pmatrix} 0 & l \\ l & 0 \end{pmatrix}, \begin{pmatrix} l & 0 \\ 0 & -l \end{pmatrix}, \begin{pmatrix} 0 & lK \\ -lK & 0 \end{pmatrix}, \begin{pmatrix} 0 & lL \\ -lL & 0 \end{pmatrix}, \begin{pmatrix} 0 & lKL \\ -lKL & 0 \end{pmatrix}.$$

To get the whole of SU(3,3), therefore, we need to extend from 2×2 to 3×3 matrices, which means adding in the corresponding Ys and Zs, and one E:

(19)
$$E_l, Y_l, Y_{lK}, Y_{lL}, Y_{lKL}, Z_l, Z_{lK}, Z_{lL}, Z_{lKL}.$$

The 'diagonal' part of this group is $U(1) \times U(1) \times SL_2(\mathbb{R}) \times SL_2(\mathbb{R}) \times SL_2(\mathbb{R})$, an 11-dimensional group generated by

$$(20) D_l, E_l, D_K, E_K, D_{L,KL}, D_L, E_L, D_{K,KL}, D_{KL}, E_{KL}, D_{K,L}, D_{K,L}, E_{KL}, D_{K,L}, D_$$

and the off-diagonal part consists of 8 dimensions each of elements of type X(bosonic), Y and Z (fermionic). Adding any one of these three types gives a group $U(1) \times SL_2(\mathbb{R}) \times SU(2,2)$ of dimension 19. The subgroup $Sp_6(\mathbb{R})$ loses the first two diagonal elements of type D, and half of the off-diagonal elements of types X, Y, Z. In this case the Ds and Xs generate a group $SL_2(\mathbb{R}) \otimes Sp_4(\mathbb{R})$ of dimension 13. This last copy of $SL_2(\mathbb{R})$ is generated by

(21)
$$E_K - F_K, E_L - F_L, E_{KL} - F_{KL},$$

or equivalently by

$$D_{I,J} + D_{IL,JL} + D_{KL,L},$$

$$D_{I,IL} + D_{J,JL} + D_{K,KL},$$

$$D_{J,IL} + D_{JL,I} + D_{K,L}.$$
(22)

and is a type of 'right-handed' $SL_2(\mathbb{R})$, as described in Section 4.

The particular real form SU(3,3) suggested here as a (huge) generalisation of the Lorentz group, from 6 dimensions to 35, is closely related to Penrose twistors, since the corresponding entry in the magic square of 2×2 matrix groups is SU(2,2). In other words, SU(3,3) combines the group $Sp_6(\mathbb{R})$ of symmetries of phase space with the group SU(2,2) of symmetries of twistors, into a single symmetry group. If this mathematical unification can lead to a physical unification, then it could have far-reaching consequences for the fundamental theory.

It should be noted that SU(2, 2) embeds in Spin(12, 4) in two different ways, as Spin(2, 4) and as Spin(4, 2), centralizing Spin(10) and Spin(8, 2) respectively. The former was used in [5] and extended to $SU(2, 3) \times SU(5)$ in an attempt to understand how twistors relate to E_8 models. Here we use the latter instead, so that the centralizer splits as $Spin(6) \otimes Spin(2, 2)$ to give a different real form of the Georgi Spin(10) GUT, and a different embedding of the twistors into E_8 . Comparing with [1], we see that the latter uses $Spin(3, 3) \otimes Spin(4)$ inside Spin(7, 3), as yet another real form of Spin(10). It is worthwhile considering which real form is most appropriate. In the Standard Model, the strong force SU(3) is definitely compact, and the mediators are correspondingly massless, but the weak force SU(2) is definitely not compact, because the complexification is used to allow the mediators to be massive. So of the three choices Spin(10), Spin(7,3) and Spin(8,2), only the last has a reasonable chance of agreeing with the Standard Model.

The group $Sp_6(\mathbb{R})$ is studied in detail in [25], embedded in a different real form of SU(6), namely $SL_3(\mathbb{H}')$. It is straightforward to translate that work into SU(3,3), simply by multiplying the Hermitian matrices by l to make them anti-Hermitian over $\mathbb{C} \otimes \mathbb{H}'$. However, any interpretation in [25] that is based on the particular real form is suspect. The study of $SL_3(\mathbb{H})$ in [1] is also not difficult to translate into, or out of, these other two real forms. But again, the interpretation offered in [1] is quite different from the interpretation I offer here. The great advantage of using SU(3,3) rather than $SL_3(\mathbb{H}')$ is that it removes the contradiction with general relativity that was apparent in [25] (see Section 9 below).

6. Representations

Let us first look at the restriction to $A_1 + A_2 + A_5$ of the adjoint representation of E_8 . The real constituents for the real form $SL_2(\mathbb{R}) \times SU(3) \times SU(3,3)$ are as follows:

$$3 = 3 \otimes 1 \otimes 1$$
$$8 = 1 \otimes 8 \otimes 1$$
$$35 = 1 \otimes 1 \otimes 35$$
$$40 = 2 \otimes 1 \otimes 20$$
$$90 = 1 \otimes 3_{\mathbb{C}} \otimes_{\mathbb{C}} 15_{\mathbb{C}}$$
$$(23) \qquad 72 = 2 \otimes 3_{\mathbb{C}} \otimes_{\mathbb{C}} 6_{\mathbb{C}}$$

The first three constituents are the adjoint representations of the three factors, and the last three involve the real weak doublet representation $\mathbf{2}$ and the complex colour triplet representation $\mathbf{3}_{\mathbb{C}}$. The representations of SU(3,3) are the natural 6-dimensional complex representation $\mathbf{6}_{\mathbb{C}}$, its anti-symmetric square $\mathbf{15}_{\mathbb{C}}$ and its anti-symmetric cube (real $\mathbf{20}$).

Restricting to SU(2,2) to separate bosonic and fermionic representations we have

(24)

$$6_{\mathbb{C}} \rightarrow (1+1) + 4$$

 $20 \rightarrow (6+6) + (4+4)$
 $15_{\mathbb{C}} \rightarrow (1+6) + (4+4)$

which gives us a total of 16 + 48 + 48 = 112 dimensions of spinors, of which 48 are right-handed and 64 are left-handed. The left-handed spinors split 16 + 48 into leptons and quarks, while the right-handed spinors here represent only quarks. The remaining 16 dimensions of right-handed spinors lie inside the group SU(3,3). In order to allocate some of these spinors to right-handed electrons, therefore, we need to break the symmetry back down to $Sp_6(\mathbb{R})$. This gives us 8 dimensions of right-handed leptons, which is the ratio that we expect.

One way to study the splitting of E_8 into representations of $A_1 + C_3 + G_2$ is to embed it first in $F_4 + G_2$, where we have a decomposition

(25)
$$248 = 52 + 14 + 26 \otimes 7.$$

Then we restrict from F_4 to $A_1 + C_3$ to get

$$\mathbf{52} = \mathbf{3} + \mathbf{21} + \mathbf{2} \otimes \mathbf{14}b$$

$$(26) 26 = 2 \otimes 6 + 14a$$

where **6** is the natural representation of $Sp_6(\mathbb{R})$, and the other representations are defined by

(27)
$$\begin{aligned} \Lambda^{2}(\mathbf{6}) &= \mathbf{1} + \mathbf{14}a, \\ S^{2}(\mathbf{6}) &= \mathbf{21}, \\ \Lambda^{3}(\mathbf{6}) &= \mathbf{6} + \mathbf{14}b. \end{aligned}$$

Here, $\Lambda^2(\mathbf{6})$ has a natural structure as a Jordan algebra, while $S^2(\mathbf{6})$ has a natural structure as a Lie algebra. An alternative way to see these splittings is by restriction from SU(3,3):

$$egin{aligned} & \mathbf{6}_{\mathbb{C}}
ightarrow \mathbf{6} \ & \mathbf{15}_{\mathbb{C}}
ightarrow \mathbf{1} + \mathbf{14}a \ & \mathbf{20}
ightarrow \mathbf{6} + \mathbf{14}b \ \end{aligned}$$

As representations of $A_1 + G_2 + C_3$ we have the following irreducible constituents of E_8 :

$$3 = 3 \otimes 1 \otimes 1$$

$$14 = 1 \otimes 14 \otimes 1$$

$$21 = 1 \otimes 1 \otimes 21$$

$$28 = 2 \otimes 1 \otimes 14b$$

$$98 = 1 \otimes 7 \otimes 14a$$

$$(29)$$

$$84 = 2 \otimes 7 \otimes 6$$

All the right-handed spinors (including the electrons) are now inside the **98**, while the left-handed lepton spinors have been split 8 + 8 between the **28** and the **84**, as a result of the breaking of **20** into 14b + 6. This curious phenomenon will no doubt repay closer scrutiny. It looks at first sight like a distinction between (massless) neutrinos and (massive) electrons, but that is not consistent with the identification of the A_1 factor as acting on weak doublets. Hence one or other of these suggested interpretations has to change.

This analysis gives us a total of five different types of spinors:

- 8 dof in 28, left-handed leptons;
- 8 + 48 dof in 98, right-handed leptons and quarks;
- 8 + 48 dof in 84, left-handed leptons and quarks.

In total, then, there are 24 dof for leptons, or 6 Weyl spinors, compared to the 9 that are usually expected for three generations. Similarly, there are 96 dof for quarks, or 24 Weyl spinors, compared to the 36 that are usually expected. Thus we need a mechanism similar to that proposed in [1] for reducing the number of

independent spinors required by one-third. This is not surprising, of course, as it is well-known that the standard interpretation requires 180 dof for spinors [6]. However, a *discrete* symmetry of order 3 can be implemented in a 2-dimensional real space using the symmetries of an equilateral triangle, so there is no theoretical reason why a 3-space is needed for this symmetry.

An alternative interpretation is that this model describes only the first-generation fermions, since we have broken the generation symmetry by choosing K. In that case, we have not too few spinors, but too many. In particular, half of the spinors that we assumed were leptons are actually associated with quarks under the action of G_2 . Hence it may be more reasonable to assume that they are (first generation) baryons, i.e. proton and neutron. This assumption allows us to use the representation **20** to describe all 'normal' matter, made out of electrons, protons and neutrons, while leaving more exotic particles to be described by other representations.

7. Restricting to $A_2 + A_2$

An alternative strategy for producing spinors for right-handed electrons is to restrict from A_5 to $A_2 \times A_2$ instead of C_3 . This extends the centralizer from $A_1 + A_2$ to $A_2 + A_2$, and gives rise to the following subgroup of $E_{8(-24)}$:

$$(30) SL_3(\mathbb{C}) \times SU(3) \times SL_3(\mathbb{R}).$$

This group provides an obvious embedding of the Lorentz group $SL_2(\mathbb{C})$ in $SL_3(\mathbb{C})$, defining the splitting into fermions and bosons, and offers various possibilities for $SL_2(\mathbb{R})$ or SO(3) for the weak force. However, $SL_3(\mathbb{R})$ acts identically on vectors and both types of spinors, so does not provide any obvious way to implement the chirality of the weak force. We considered this possibility in the work that led to [1], but did not find a way to make it work. Of course, that does not necessarily mean that it cannot be done.

The group $SL_3(\mathbb{C})$ is generated by the 16 elements

$$(31) D_l, D_L, X_1, X_l, X_L, X_{lL}, E_l, E_L, Y_1, Y_l, Y_L, Y_{lL}, Z_1, Z_l, Z_L, Z_{lL}$$

in which the top row is the Lorentz group $SL_2(\mathbb{C})$, centralized by a complex scalar generated by

(32)
$$E_L - F_L = D_L + 2E_L, \\ E_l - F_l = D_l + 2E_l.$$

The group SU(3) acts on the labels i, j, k, il, jl, kl, identically on X, Y and Z. The group $SL_3(\mathbb{R})$ acts similarly on the labels I, J, K, IL, JL, KL. The 120 spinors therefore split as 24 + 24 + 72, in which the 72 have both an i, j, k and an I, J, K in the label, and the 24s have one or the other.

We now have to break the symmetry of I, J, K in order to separate left-handed and right-handed spinors, so let us separate I, J from K to break 72 = 24 + 48 and one of the 24 = 8 + 16. This gives us a splitting of quarks in 24 + 24 + 48, such that 24 + 24 are right-handed, and 48 are left-handed. Similarly, the leptons split as 8 right-handed and 16 left-handed, again in agreement with the Standard Model. To be more explicit, we give the labels in the form of a table, with the labels for the C_3 version for comparison:

$$(33) \qquad \begin{array}{c} \operatorname{RH} 1 \\ \operatorname{LH} 1 \\ \operatorname{RH} q \\ \operatorname{LH} q \\ \operatorname{LH} q \\ \operatorname{LH} i, j, k, il, jl, kl \\ \operatorname{LH} i, j, k, il, jl, kl \\ \operatorname{LH} jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, J, IL, JL \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, j, kl \\ \operatorname{I}, j, k, il, jl, kl \\ \operatorname{I}, j, kl$$

The actual signs for the right-handed leptons in the C_3 case are different in the Y and Z spinors, but only one sign occurs in each case. The allocation of individual particles in the C_3 and $A_2 + A_2$ cases is not necessarily the same, but the overall picture is very similar. But only the C_3 case has the projection with U - KL that corresponds to $1 - \gamma_5$ in the Standard Model. This appears to be a decisive vote in favour of the C_3 model over the $A_2 + A_2$ model, at least for the Dirac algebra.

On the other hand, the Dirac equation is probably better modelled with $A_2 + A_2$, where there are 12 Lorentzian 4-vectors available for the gamma matrices. Nine of these extend Spin(3,1) to Spin(4,1), and therefore have an obvious place to put a mass that is defined by the Einstein mass equation:

$$\begin{array}{cccc} & D_{I}, X_{I}, X_{lI}, D_{I,L}, & X_{i}, D_{i}, D_{i,l}, X_{iL} & X_{il}, D_{il}, D_{il,l}, X_{ilL} \\ (34) & D_{J}, X_{J}, X_{lJ}, D_{J,L}, & X_{j}, D_{j}, D_{j,l}, X_{jL}, & X_{jl}, D_{jl}, D_{jl,l}, X_{jlL}, \\ & D_{K}, X_{K}, X_{lK}, D_{K,L} & X_{k}, D_{k}, D_{k,l}, X_{kL} & X_{kl}, D_{kl}, D_{kl,l}, X_{klL} \end{array}$$

The first column look like electrons, and the others look like quarks, although it is also possible to take arbitrary linear combinations. The other three extend Spin(3, 1) to Spin(3, 2), and therefore do not have an Einstein/Dirac mass term:

(35)
$$D_{IL}, X_{IL}, X_{UL}, D_{IL,L}, \\ D_{JL}, X_{JL}, X_{IJL}, D_{JL,L}, \\ D_{KL}, X_{KL}, X_{IKL}, D_{KL,L}$$

These are presumably neutrinos.

8. Symmetry-breaking

Nevertheless, there are many questions remaining about the differences between C_3 and $A_2 + A_2$, particularly concerning the physical interpretations, and the reason for the symmetry-breaking from $SL_3(\mathbb{R})$ to $SL_2(\mathbb{R})$. In order to bring the questions into focus, it is useful to consider the square of groups A_2 , $A_2 + A_2$, C_3 and A_5 , together with the corresponding block of the 2×2 magic square, and the centralizers.

	$SL_3(\mathbb{R}) \\ Sp_6(\mathbb{R})$	$SL_3(\mathbb{C})$ SU(3,3)	Spin Spin	$(2,1) \\ (3,2)$	$\begin{array}{c} Spin(3,1)\\ Spin(4,2) \end{array}$
(36)		$G_2 ightarrow G_2 ightarrow G_2 ightarrow$	$\langle SL_3(\mathbb{R}) \\ \langle SL_2(\mathbb{R}) \rangle$	$\frac{SU(3)}{SU(3)}$	$\begin{array}{l} \times SL_3(\mathbb{R}) \\ \times SL_2(\mathbb{R}) \end{array}$

It may also be helpful to display the generators for the four groups in a table

	X_1, X_L, D_L	X_l, X_{lL}, D_l
(37)	Y_1, Y_L, Z_1, Z_L, E_L	$Y_l, Y_{lL}, Z_l, Z_{lL}, E_l$
(37)	$X_K, X_{KL}, D_K, D_{KL}, D_{L,KL}, D_{K,L}, D_{K,KL}$	X_{lK}, X_{lKL}
	$Y_K, Y_{KL}, Z_K, Z_{KL}, E_K, E_{KL}$	$Y_{lK}, Y_{lKL}, Z_{lK}, Z_{lKL}$

where each box contains the extra generators needed in each case, to add to the generators of the group(s) above and/or to the left of the box.

We would expect to use Spin(3, 1) in the top right corner for the Lorentz group, acting on the four labels u, U, l, L for spacetime and/or 4-momentum. However, in the Dirac algebra including γ_5 we need Spin(3, 2), which appears in the bottomleft, acting on the five labels u, U, K, L, KL. From this labelling we see that we have lost one of the three dimensions of momentum, labelled l, and gained two dimensions of something else, labelled K, KL. In particular, we have broken the symmetry of spacetime, to include a preferred direction in space, and we have broken the symmetry of I, J, K, to obtain a weak force with a broken symmetry group $SL_2(\mathbb{R})$. The breaking of the I, J, K symmetry seems most likely to be a breaking of the generation symmetry of fundamental fermions. We can, of course, avoid the breaking of the spacetime-symmetry by extending to Spin(4, 2) and the complex Dirac algebra, as is effectively done in the Standard Model.

But in the E_8 model it seems more likely that the symmetry-breaking from complex quaternions and Spin(4, 2) down to real split quaternions and Spin(3, 2)is important for the description of those parts of physics that use (mathematically real) momentum but not (mathematically imaginary) current. In other words we may need this symmetry-breaking for the description of mass and gravity. The particular space direction chosen must be something physically important such as the direction of acceleration relative to an inertial frame, or the direction of the ambient gravitational field, or the direction of the ambient angular momentum.

This suggests that the top right corner, in the form

$$(38) SL_3(\mathbb{C}) \times SU(3) \times SL_3(\mathbb{R}),$$

represents the Standard Model of elementary particles, with three colours of quarks and three generations of fermions (but only for half of the right-handed quarks!). The restriction from $SL_3(\mathbb{C})$ to $SL_2(\mathbb{C})$ splits fermions from bosons, and allows every observer to choose their own preferred copy of the Lorentz group $SL_2(\mathbb{C})$. The ten remaining dimensions of $SL_3(\mathbb{C})$ consist of a complex scalar and a Dirac spinor. Hence there is an 8-parameter family of copies of $SL_2(\mathbb{C})$ available for different observers. This compares to a 9-parameter family of copies of SO(3, 1) inside $SL_4(\mathbb{R})$ that describes the analogous phenomenon in General Relativity. Clearly, therefore, this model does not resolve the basic problem of incompatibility of GR with QM. It does, however, provide an explanation of sorts for the so-called 'righthanded neutrinos': these are not interpreted as particles, but as transformations between different coordinate systems preferred by different observers.

The interpretation of the bottom left corner, in the form

$$(39) Sp_6(\mathbb{R}) \times G_2 \times SL_2(\mathbb{R}),$$

is now freed from the necessity to include the Lorentz group in $Sp_6(\mathbb{R})$, and hence freed from the necessity to extend Minkowski spacetime, with symmetry group SO(3,1), to anti-de Sitter spacetime, with symmetry group SO(3,2). Indeed, the latter group, or rather its double cover $Sp_4(\mathbb{R})$, now only has to act on two of the three spatial coordinates, and can therefore plausibly be identified with the group of symmetries of phase space for 2-dimensional dynamics. The choice of which two dimensions these are is the same as the choice of restriction from SO(3,1) to SO(2,1) above, and therefore has the same relationship to acceleration, rotation and/or the gravitational field. In practice, most experiments are horizontal, so that the most likely direction in most cases will be the direction of the gravitational field. However, other directions may enter into the model at various points.

ROBERT ARNOTT WILSON

Finally, the extension from $Sp_4(\mathbb{R})$ to $Sp_6(\mathbb{R})$ allows individual observers to choose whichever 2-dimensional part of dynamics they wish to model with the Standard Model, and hence to modify the Standard Model for different ambient conditions of acceleration, rotation and gravity. If we want a model that is truly relativistic (independent of the observer), then we can treat $Sp_6(\mathbb{R})$ as the symmetry group of phase space for 3-dimensional dynamics, which entails abandoning the concept of 'spacetime', since it is now observer-dependent, and therefore no longer useful for a fundamental theory. The embedding of $Sp_4(\mathbb{R})$ via $Sp_4(\mathbb{R}) \times Sp_2(\mathbb{R})$ into $Sp_6(\mathbb{R})$ shows that there is again an 8-parameter family of copies of $Sp_4(\mathbb{R})$ available, for an 8-parameter family of observers. This extends to an 11-parameter family of copies of $SL_2(\mathbb{C})$, in case this is useful.

A more radical proposal is made in [25], in which it is noted that both SU(3)and $SL_3(\mathbb{R})$ are subgroups of $Sp_6(\mathbb{R})$, and it is therefore proposed to identify the colour symmetry group SU(3) and the generation symmetry group $SL_3(\mathbb{R})$ with the corresponding subgroups of $Sp_6(\mathbb{R})$, so that the latter group now contains all of the required symmetries of fundamental physics. It may be that this proposal is too radical, but it is certainly necessary to have *some* mechanism for linking the generation symmetry group $SL_3(\mathbb{R})$ to *some* concept of mass.

9. GRAVITY

It must be stressed again that with the standard interpretation of the Lorentz group $SL_2(\mathbb{C})$ acting on spacetime, labelled by u, U, l, L, there is no conceivable way of implementing General Relativity inside $E_{8(-24)}$. The fact that $SL_4(\mathbb{R})$ has no double cover that acts on spinors categorically rules this out. However, there are bits of E_8 that appear not to be used in the Standard Model, that could in principle be used for a quantum theory of gravity, and that might approximate to GR in appropriate circumstances. Or it may be that a suitable tweak to the interpretation might allow GR in to the model.

For example, $SL_4(\mathbb{R})$ acting on spacetime translates to SO(3,3) acting on phase space, and there is an obvious copy of SO(3,3) in SU(3,3), that extends $SL_3(\mathbb{R})$ in $Sp_6(\mathbb{R})$. This suggests that complexifying phase space, in order to implement momentum and current separately, may be key to including quantum gravity in the model. In other words, A_5 is the battleground in which the QM and GR models must resolve their differences. These differences mainly lie in the fundamental properties of spacetime, so the hope is that by shifting the emphasis from spacetime to phase space, a compromise might be reached in which spacetime is never defined or used at all. If QM uses $Sp_6(\mathbb{R})$ to act on phase space, as it should in a Hamiltonian theory, and GR uses SO(3,3), then both can be embedded in SU(3,3), and each can be allowed to add corrections to the other.

The key to this process is to re-interpret the various spin groups as acting on 2+2dimensional (real, complex or quaternionic) phase space, and not on an abstract space of 'spinors' that have no concrete physical interpretation. We have already done this [25] with the group $Spin(3,2) \cong Sp_4(\mathbb{R})$ in C_3 , interpreted as a subalgebra of the Dirac algebra, so we must now do the same with $Spin(4,2) \cong SU(2,2)$ in A_5 , and interpret this both as the full complex Dirac algebra, and as a complex version of the subgroup SO(2,2) of SO(3,3), corresponding to a subgroup $SL_2(\mathbb{R}) \times SL_2(\mathbb{R})$ of $SL_4(\mathbb{R})$ in GR. Within the 2×2 magic square, we therefore have a copy of

(40)
$$SO(2,2) = Spin(2,1) \otimes Spin(2,1)$$

inside Spin(4,2), from which we can write down generators for SO(3,3) as follows:

(41)
$$D_L, D_{K,KL}, X_1, X_L, X_{lK}, X_{lKL}, E_L, Y_1, Y_L, Y_{lK}, Y_{lKL}, Z_1, Z_L, Z_{lK}, Z_{lKL}$$

These are the elements in the top left and bottom right of (37), together with $D_{K,KL}$. The first row of (41) consists of generators for SO(2,2). The elements whose labels do not include l generate $GL_3(\mathbb{R})$ inside $Sp_6(\mathbb{R})$.

This choice of SO(2,2) amounts to splitting the 6 labels into u, U, L for one copy of Spin(2,1) and l, K, KL for the other. In matrix terms we have

(42)
$$\begin{pmatrix} L \\ 0 \\ -L \end{pmatrix}, \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ L \\ 0 \end{pmatrix}, \begin{pmatrix} L \\ 0 \\ 0 \\ L \end{pmatrix}, \begin{pmatrix} 0 \\ -lK \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ -lKL \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ -lKL \\ 0 \end{pmatrix},$$

where the first row is SO(2,1) acting on 2 + 1-dimensional spacetime.

This splitting is similar to Woit's splitting [23, 24] into 'right-handed' spacetime and 'left-handed' gauge groups, respectively, except that he complexifies everything and interprets these groups as compact SU(2) rather than split $SL_2(\mathbb{R})$. Moreover, he interprets SU(2) as acting on a Euclidean rather than Lorentzian spacetime, or a spinor. In our interpretation, the 'right-handed' copy acts on 2 + 1-spacetime, which we must presumably interpret as being perpendicular to the local direction of the gravitational field, and the 'left-handed' copy acts on 2 + 1 fermions in a single generation.

At this point we may notice that we have acquired two copies of $SL_2(\mathbb{R})$, both labelled 'left-handed', one acting on labels l, K, KL, the other acting on labels I, J, IL, JL. Some mixing of the two may be required in order to define masses for the intermediate vector bosons. We also have a second 'right-handed' copy acting on the six labels I, J, K, IL, JL, KL, but not acting on l. Again it is not clear whether there is any relationship between this right-handed copy, analogous to the one defined in [1], and the one acting on u, U, L (with or without l), analogous to the one defined in [23, 24]. What is clear, at least, is that the terms 'righthanded' and 'left-handed' have multiple different meanings in the literature, and are consequently best avoided.

10. General relativity

The tensors used in GR are irreducible representations of SO(3,3) of dimensions 6 (field strength tensor), 10 (Ricci tensor, stress-energy tensor) and 20 (Riemann curvature tensor). The representations of SU(3,3) that are available in E_8 are complex $\mathbf{6}_{\mathbb{C}}$ and $\mathbf{15}_{\mathbb{C}}$, and real **20** and **35**, which restrict to SO(3,3) as follows:

$$\begin{array}{c} {\bf 6}_{\mathbb C} \to {\bf 6} \\ {\bf 15}_{\mathbb C} \to {\bf 15} \\ {\bf 20} \to {\bf 10}a + {\bf 10}b \\ {\bf 35} \to {\bf 15} + {\bf 20} \end{array}$$

It follows that the Einstein field equations must be written in terms of the representation $\mathbf{2} \otimes \mathbf{20}$ on left-handed leptons, that relates matter to neutrinos, and therefore expresses the gravitational field in terms of neutrinos. On the other hand, the Riemann Curvature Tensor (RCT) lies in the adjoint representation of SU(3,3), and consists of the 20 dimensions outside SO(3,3). Thus the RCT becomes part of the symmetry group of the model, and consists of the elements in the top right and bottom left of (37), excluding the 'scalar' $D_{K,KL}$. Finally, the field strength tensor is coupled to the quark colour/charge representation $\mathbf{2} \otimes \mathbf{3}_{\mathbb{C}}$, and hence to baryonic matter.

From this analysis, we see that the only essential thing that is missing from GR is the distinction between 10a and 10b, which are usually regarded as being self-dual and therefore equivalent, but are in fact dual to each other. In the usual formalism in terms of the Lorentz group SO(3, 1), they both restrict to 1 + 9, so that the distinction between them is less obvious. The introduction of a second scalar (the cosmological constant) only extends from 10 to 11 variables, when 20 are required for the full theory. A consequence of this generalisation is that electrons are far more important for gravity than is usually supposed, since all three generations of neutrinos and antineutrinos, and therefore all three generations of electrons, participate in an essential way.

Indeed, it is possible to include the whole of GR and the weak force inside E_6 by adding the representation $\mathbf{2} \otimes \mathbf{20}$ to the adjoint representation of SU(3,3), so that on restriction to $A_1 + A_3$ the adjoint representation of E_6 has the structure

(44)
$$\mathbf{3} \otimes \mathbf{1} + \mathbf{1} \otimes (\mathbf{15} + \mathbf{20}) + \mathbf{2} \otimes (\mathbf{10}a + \mathbf{10}b).$$

The representation $\mathbf{3} \otimes \mathbf{1}$ is the adjoint representation of $SL_2(\mathbb{R})$, generated by

(45)
$$D_{I,J} - D_{IL,JL}, \quad D_{I,IL} - D_{J,JL}, \quad D_{J,IL} + D_{I,JL}.$$

and generators for the adjoint representation $\mathbf{1} \otimes \mathbf{15}$ of SO(3,3) are listed in (41). A translation to 6×6 real matrices is obtained via the map

(46)
$$lKL \mapsto \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad L \mapsto \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad lK \mapsto \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

but it is noteworthy that this translation does *not* map **15** to antisymmetric matrices, as one might naively expect. This is because of the difference between the split and compact signatures. For example, we have

$$(47) X_{1} \mapsto \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, X_{L} \mapsto \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \\ \begin{pmatrix} 47 \end{pmatrix} X_{lK} \mapsto \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, X_{lKL} \mapsto \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

The Riemann Curvature Tensor **20** is generated by

$$(48) \begin{array}{c} D_{l}, E_{l}, D_{K}, E_{K}, D_{KL}, E_{KL}, D_{L,KL}, D_{K,L}, \\ X_{l}, X_{lL}, X_{K}, X_{KL}, \\ Y_{l}, Y_{lL}, Y_{K}, Y_{KL}, \\ Z_{l}, Z_{lL}, Z_{K}, Z_{KL}. \end{array}$$

We can take the five elements with label l to correspond to the diagonal elements of the tensor as 6×6 matrices.

The four 10-dimensional tensors consist of all the elements containing I or J in their labels, and they each split into Ds, Xs, Ys and Zs as 4 + 2 + 2 + 2. Bases for the four representations are given below. The top two can be taken as the two copies of $\mathbf{10}a$, mixed by $SL_2(\mathbb{R})$, in which case the bottom two are $\mathbf{10}b$.

	$X_{lI(1+L)}, Y_{lI(1+L)}, Z_{lI(1+L)},$		$X_{lI(1-L)}, Y_{lI(1-L)}, Z_{lI(1-L)},$
	$X_{J(1+L)}, Y_{J(1+L)}, Z_{J(1+L)},$		$X_{J(1-L)}, Y_{J(1-L)}, Z_{J(1-L)},$
	$D_{K,I(1+L)}, E_{K,I(1+L)},$		$D_{K,I(1-L)}, E_{K,I(1-L)},$
	$D_{KL,I(1+L)}, E_{KL,I(1+L)}.$		$D_{KL,I(1-L)}, E_{KL,I(1-L)}.$
	$X_{I(1+L)}, Y_{I(1+L)}, Z_{I(1+L)},$]	$X_{I(1-L)}, Y_{I(1-L)}, Z_{I(1-L)},$
(49)	$X_{lJ(1+L)}, Y_{lJ(1+L)}, Z_{lJ(1+L)},$		$X_{lJ(1-L)}, Y_{lJ(1-L)}, Z_{lJ(1-L)},$
	$D_{K,J(1+L)}, E_{K,J(1+L)},$		$D_{K,J(1-L)}, E_{K,J(1-L)},$
	$D_{KL,J(1+L)}, E_{KL,J(1+L)}.$		$D_{KL,J(1-L)}, E_{KL,J(1-L)}.$

11. MOND

The proposed extension to General Relativity involves extending the standard 10-dimensional symmetric rank 2 tensors on spacetime to 20-dimensional antisymmetric rank 3 tensors on phase space. Restricting to the Lorentz group $SO_3(\mathbb{C})$, the latter breaks up as 1 + 9 + 9 + 1, containing two real scalars, with spin (0, 0), and two real representations with spin (1, 1). With respect to a basis of phase space consisting of 3 directions of momentum and 3 directions of position, the antisymmetric cube breaks up into pieces with 0, 1, 2 and 3 position coordinates, of dimensions 1, 9, 9 and 1 respectively.

Newton-Einstein gravity consists mainly of the inverse-square law, with two position vectors, and a relativistic 'mass' (or energy) that involves the momentum vector. This occupies one of the 9-dimensional pieces, mixed with a scalar 'rest mass', that involves all three position coordinates. In other words, the rest mass is defined by an inverse cube law that relates an object to its surroundings. The other scalar has no position coordinates at all, and therefore plays the role of a cosmological constant, or a 'dark energy' field that in principle may not be constant throughout the universe [26]. Indeed, this 'dark energy' depends on the anti-symmetric cube of momentum, and therefore relates an object to the rest of the universe via Mach's Principle.

This leaves one 9-dimensional piece of the representation unaccounted for in Newton–Einstein gravity. This has a single position vector, and therefore corresponds to an inverse-linear term in the universal law of gravity. Such a term was first proposed by Milgrom [27] in 1983, in order to reconcile the theory of gravity with observations that appeared to be inconsistent with the existing theory. This proposed modification to Newtonian dynamics (MOND) has since been confirmed in a wide variety of studies of astronomical objects on varying scales [28, 29, 30].

ROBERT ARNOTT WILSON

The model proposed here suggests a dependence of this inverse-linear term on a square of momentum, which may mean the momentum of each body relative to the 'centre-of-mass' frame, assuming that such a frame can be adequately defined. However, it may be necessary to embed the system in a larger system (or external field), in which case we have two independent momenta relative to the external system. In particular, for a star on the outskirts of a spiral galaxy, with a reasonably 'standard' stellar mass, and where the speed of rotation with respect to the centre of the galaxy is observationally close to a constant value (in contrast to the expected Keplerian decline), this momentum-squared factor is close to being a constant. But the point at which the inverse-linear term becomes dominant is not yet clear from this picture, and I have not found a convincing explanation for the empirical 'critical acceleration' a_0 . What appears to be critical is a ratio of momentum to distance, which translates in Newtonian physics to a ratio of mass to duration.

A number of models [31, 32, 33] have attempted to reconcile GR with MOND by adding extra scalar and/or vector fields to the Einsteinian tensors. However, the analysis of symmetries carried out in this paper suggests that it is not sufficient to add scalars or vectors, but that it is necessary to add a second tensor, dual to the first. On the other hand it has been proposed by Yahalom [34] that the inverselinear term is merely an artefact of the fact that the gravitational field propagates at a finite and not infinite speed. In other words, it is not enough to know where an object is, it is also necessary to know how fast its mass is travelling. It may or may not be necessary to separate its momentum into mass and velocity, but it is certainly necessary to know the momentum. Thus it may be possible to implement this 'retarded gravity' approach within the phase space model that I propose.

12. CONCLUSION

The problem of unification of particle physics and gravity goes back almost a century, and occupied Einstein for at least a quarter of that century. Yet the problem seems no nearer to a solution today than it did fifty years ago, and even further away than it seemed forty or thirty years ago. This indicates that there must be something subtly wrong in the basic assumptions somewhere. My analysis locates this problem in the concept of spacetime itself. The way that spacetime is treated in relativity, using the Lorentz group in the form SO(3, 1), is mathematically (never mind physically) inconsistent with the way that spacetime is treated in quantum mechanics, using the Lorentz group in the form $SL_2(\mathbb{C})$. It isn't a question of one of them being 'right' and the other one 'wrong', it is a question of there being no consistent definition of spacetime at all, and no possible way to measure spacetime in the absence of objects embedded in spacetime.

Therefore I have considered the possibility of describing physics without using spacetime, but instead using only phase space, as Hamilton taught us to do. This involves re-interpreting the Dirac algebra, the Einstein field equations and the Riemann curvature tensor in terms of a complex phase space, in order to include both momentum and current. The symmetry group of the complex phase space in three dimensions is the complex symplectic group $Sp_6(\mathbb{C}) \cong SU(3,3)$. Taking my cue from the variety of models based on E_8 and the Freudenthal–Tits magic square, from $E_8 \times E_8$ heterotic string theory down, I focus on the group SU(3,3) embedded in $E_{8(-24)}$ via $E_{6(2)}$, and find within these groups all the mathematical structures that a unified model appears to require.

This is not, of course, in itself a unified theory of fundamental physics. But it is a unified mathematical model, in which all the ingredients of all the fundamental theories of physics can be found. This includes the complex Dirac algebra, the gauge groups of the weak and strong nuclear forces, including symmetry-breaking of the weak force, a classification of elementary fermions in which there are no righthanded neutrinos, general covariance, and all the tensors used in GR. Moreover, these ingredients fit together in ways that are broadly consistent with experiment. I therefore suggest that this is a promising foundation on which to try to build a unified theory of fundamental physics.

References

- C. A. Manogue, T. Dray and R. A. Wilson (2022), Octions: an E₈ description of the standard model, J. Math. Phys. 63, 081703. arXiv:2204.05310
- [2] A. G. Lisi (2007), An exceptionally simple theory of everything, arXiv:0711.0770
- [3] A. G. Lisi (2010), An explicit embedding of gravity and the standard model in E8, arXiv:1006.4908
- [4] D. Chester, A. Marrani and M. Rios (2020), Beyond the standard model with six-dimensional spacetime, arXiv:2002.02391.
- [5] R. A. Wilson (2022), Chirality in an E_8 model of elementary particles, arXiv:2210.06029.
- [6] J. Distler and S. Garibaldi (2010), There is no E₈ theory of everything, Communications in Math. Phys. 298 (2), 419–436.
- [7] H. Freudenthal, Beziehungen der E₇ und E₈ zur Oktavenebene. X-XI. Indag. Math. 25 (1963), 457–487.
- [8] J. Tits, Algèbres alternatives, algèbres de Jordan et algèbres de Lie exceptionelles, Indag. Math. 28 (1966), 223–237.
- [9] C. H. Barton and A. Sudbery, Magic squares and matrix models of Lie algebras, Advances in Math. 180 (2003), 596–647.
- [10] C. A. Manogue and T. Dray (2010), Octonions, E6, and particle physics, J. Phys. Conf. Ser. 254, 012005.
- [11] I. Todorov and M. Dubois-Violette (2018), Deducing the symmetry of the standard model from the automorphism and structure groups of the exceptional Jordan algebra, Int. J. Mod. Phys. A 33, 1850118
- [12] J. C. Pati and A. Salam (1974), Lepton number as the fourth 'color', Phys. Rev. D 10 (1), 275–289.
- [13] R. Penrose (1967), Twistor algebra, J. Math. Phys. 8 (2), 345-366.
- [14] T. Adamo (2017), Lectures on twistor theory. arXiv:1712.02196.
- [15] J. Maldecena (1998), The large N limit of superconformal field theories and supergravity, Adv. Theor. Math. Phys. 2, 231–252.
- [16] T. Dray. C. A. Manogue and R. A. Wilson (2014), A symplectic representation of E₇, Comment. Math. Univ. Carolin. 55, 387.
- [17] T. Dray, C. A. Manogue and R. A. Wilson (2024), A new division algebra representation of E₇, arXiv:2401.105
- [18] M. de Gosson and B. Hiley (2011), Imprints of the quantum world in classical mechanics, Found. Phys. 41, 1415–1436.
- [19] R. A. Wilson, T. Dray and C. A. Manogue (2022), An octonionic construction of E_8 and the Lie algebra magic square, arXiv:2204.04996
- [20] M. Gell-Mann (1961), The eightfold way: a theory of strong interaction symmetry, Synchrotron Lab. Report CTSL-20, Cal. Tech.
- [21] W. Greiner, S. Schramm and E. Stein (2007), Quantum Chromodynamics, Springer.
- [22] S. Coleman and J. Mandula (1967), All possible symmetries of the S matrix, *Physical Review* 159 (5), 1251.
- [23] P. Woit (2021), Euclidean spinors and twistor unification, arXiv:2104.05099
- [24] P. Woit (2023), Spacetime is right-handed, arXiv:2311.00608.
- [25] R. A. Wilson (2024), A Clifford algebra model in phase space, arXiv:2404.04278.
- [26] J. Colin, R. Mohayaee, M. Rameez and S. Sarkar (2019), Evidence for anisotropy of cosmic acceleration, Astronomy and Astrophysics 631, L13. arXiv:1808.04597.

ROBERT ARNOTT WILSON

- [27] M. Milgrom (1983), A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis, Astrophysical J. 270, 365–370.
- [28] B. Famaey and S. McGaugh (2012), Modified Newtonian dynamics (MOND): observational phenomenology and relativistic extensions, *Living reviews in relativity* 15, 10. arXiv:1112.3960.
- [29] K.-H. Chae (2023), Breakdown of the Newton–Einstein standard gravity at low accelerations in internal dynamics of wide binary stars, *The Astrophysical Journal* **952**, 128.
- [30] X. Hernandez (2023), MNRAS 525, 1401.
- [31] R. H. Sanders (1997), A stratified framework for scalar-tensor theories of Modified Dynamics, Astrophys. J. 480, 492–502.
- [32] J. D. Bekenstein (2004), Relativistic gravitation theory for the modified Newtonian dynamics paradigm, *Phys. Rev. D* 70, 083509.
- [33] C. Skordis and T. Złośnik (2021), A new relativistic theory for Modified Newtonian Dynamics, *Phys. Rev. Lett.* **127**, 161302.
- [34] A. Yahalom (2019), The effect of retardation on galactic rotation curves, J. Phys.: Conf. Ser. 1239, 012006.

QUEEN MARY UNIVERSITY OF LONDON Email address: r.a.wilson@qmul.ac.uk