

Environmental cosmic acceleration from a phase transition in the dark sector

Ø. Christiansen,¹ F. Hassani,¹ and D. F. Mota¹

¹ *Institute for Theoretical Astrophysics, University of Oslo, Sem Sælands vei 13, Oslo, 0371, Norway*

(Dated: May 2, 2024)

A new degravitation mechanism within the framework of scalar tensor gravity is proposed. The mechanism eliminates all constant contributions from the potential to the Friedmann equation, leaving only the kinematic and the dynamic terms of the potential to drive cosmic acceleration. We explore a scenario involving a density-triggered phase transition in the late-time universe, and argue that the resulting effective energy density and equation of state parameter can explain late-time cosmology when extrapolated to a region of the parameter space.

Cosmological data indicate that the expansion of the universe is currently accelerating [1]. The simplest way to explain this is a fine-tuned cosmological constant, which is unexplained by the Standard Model of particle physics [2]. Moreover, tensions in recent observational evidence, if not due to systematics, seem to require a more phenomenologically complex ingredient than a constant [3]. Observations also suggest tensions for the dark sector at smaller scales [4], suggesting the need for more detailed modelling of the entire dark sector. This has motivated studies of extensions to the Standard Model of cosmology and general relativity.

Here we present results from a novel explanation of the late-time accelerating expansion caused by a modified version of the symmetron [5, 6]. The symmetron is a real scalar field with an environment-dependent mass term using the Anderson-Higgs mechanism; in low-density environments, its vacuum expectation value (VEV) drifts away from the origin, causing a phase of out-of-equilibrium transition for the scalar field. The VEV also degenerates and takes on a non-trivial topology, leading to the appearance of domain walls at cosmological scales. The resulting model has a wide range of phenomenological consequences, which individually may seem to be able to account for the cosmological observations and alleviate tensions [6–12].

Whether the symmetron, or some variant of it, can simultaneously account for the cosmological tensions remains an open question. The classical symmetron contains a cosmological constant that is included to drive the late-time accelerating expansion [6], which in turn is fine-tuned. In the following, we define a model that continuously removes constant contributions to the vacuum energy; instead we try to account for the cosmic acceleration through the dynamic behaviour of the scalar field. Recent advances in cosmological simulations of the symmetron [11, 13] allow the comprehensive treatment of the scalar field's energy contribution for some parameter ranges, that we present the results for here. By extrapolating our results, we indicate a region of the model parameter space that is interesting with regards to producing the late-time cosmic expansion.

The field equations of the symmetron $\phi \in \mathbb{R}$ on a flat

cosmological background is [6]

$$\ddot{\phi} - \frac{1}{a^2} \nabla^2 \phi + 3H\dot{\phi} = V_{\text{eff},\phi} = V_{,\phi} - \frac{A_{,\phi}(\phi)}{A(\phi)} T, \quad (1)$$

where a is the scale factor, H is the Hubble function, T is the trace of the stress-energy tensor of the matter sector, dotted quantities are differentiated with respect to cosmic time, V_{eff} is the effective scalar field potential and V is the potential

$$V = -\frac{1}{2}\mu^2\phi^2 + \frac{\lambda}{4}\phi^4, \quad (2)$$

with mass μ and self-coupling strength λ . In the non-relativistic limit, $T \sim -\rho_m$ is the energy density of matter. $A(\phi) \equiv 1 + \frac{1}{2} \left(\frac{\phi}{M} \right)^2$ is the conformal factor, with the conformal coupling strength M . This gives a term for the effective potential which provides a stable minimum at the origin as long as $\rho_m > \rho_* = \mu^2 M^2$. As ρ_m becomes smaller, the effective potential becomes the Mexican hat potential, and develops two stable minima \tilde{v} (VEVs) at non-zero field values. The VEVs are found locally (at time t and position x) by minimising V_{eff} and are given by

$$\tilde{v}_{\pm}(t, x) = \pm \frac{\mu}{\sqrt{\lambda}} \sqrt{1 + \frac{T(t, x)}{\rho_*}}, \quad (3)$$

We define the true VEV $v_0 \equiv \tilde{v}_+(T=0)$. The potential V , equation (2), in the true vacuum can be related to phenomenological quantities [13]

$$V_0 = \frac{\lambda}{4} v_0^4 = \frac{9\beta_*^2 L_C^2}{2a_*^6} H_0^4 \Omega_{m,0}^2, \quad (4)$$

where $a_* = (1 + z_*)^{-1}$ is the scale factor of the phase transition in a homogeneous universe, β_* is the relative strength of the gravitational force in the true VEV v_0 , and L_C is the Compton wavelength related to the mass μ as $L_C = \frac{1}{\sqrt{2}\mu}$. H_0 is the current time Hubble parameter and $\Omega_{m,0}$ is the current time energy fraction of matter.

For nonlinear fields such as the symmetron, that may develop large spatial gradients due to screening and topological defects, the gradients may have a significant effect

on the background evolution. It has been shown in [11] that inhomogeneities in the density field cause the field to undergo phase transition much earlier than it would on the background. To include this effect into the vacuum, we define the generalised potential V_{gen} whose derivative $V_{\text{gen},\phi} \equiv V_{\text{eff},\phi} + \frac{1}{a^2} \nabla^2 \phi$ includes the Laplacian term from equation (1). The generalised VEV v is found by minimising V_{gen} , i.e. solving

$$\nabla^2 v^2 = -a^2 \partial_v V_{\text{eff}}(v), \quad (5)$$

and is equivalent to finding the symmetron's quasistatic limit (equation (1) with $\dot{\phi}, \ddot{\phi} \rightarrow 0$), which is also referred to as the adiabatic limit in the literature.

In the usual approach, such as for quintessence dark energy, the potential contains a constant term, $\langle V(\phi = 0) \rangle$, which contributes to the Friedmann equation, and is the driver of cosmic acceleration. To get a constant term that can account for cosmic acceleration requires a high degree of fine-tuning (this is the smallness problem of the cosmological constant). We therefore propose a model where the potential contribution to the Friedmann equation is¹

$$\Omega_V(\phi, v) = \frac{\lambda}{4\rho_c} \langle (\phi^2 - v^2)^2 \rangle, \quad (6)$$

which automatically removes all vacuum contributions, and eliminates the need to fine-tune the cosmological constant. The full contribution to the Friedmann equation includes the kinetic and gradient terms that are expanded around the VEV,

$$\Omega_\phi = \Omega_V + \frac{1}{2\rho_c} \langle (\dot{\phi} - \dot{v})^2 + \partial_i(\phi - v)\partial^i(\phi - v) \rangle, \quad (7)$$

and we will refer to the last two terms as Ω_{kin} and Ω_{grad} respectively. $\rho_c(a)$ is the critical density of the universe and the brackets mean that we take the spatial average of their argument, because the Friedmann equation is evaluated on the background.

Since v is the quasistatic solution of the field equation, Ω_V has contributions from the dynamic parts of the field. In the absence of large inhomogeneities in the scalar field (when $\nabla^2 \phi \rightarrow 0$, and $V_{\text{gen},\phi} = V_{\text{eff},\phi}$), then $v \rightarrow \tilde{v}_\pm$, and we recover the classical symmetron energy contribution, with an additional term proportional to v_\pm^4 .

We define the dynamical and topological contributions as $\Omega_V = \Omega_{\text{dyn}} + \Omega_{\text{top}}$. If the dynamic field, ϕ , develops topological defects through a phase transition, then there is a question of how to solve the VEV v in equation (5). If we solve it using the same vacuum manifold as is obtained in the dynamical solution, by which we mean that we are

solving the VEV around the same topological defects, then we remove the contribution of topological defects in equation (6). Alternatively, we can solve the VEV using a trivial vacuum manifold (i.e. all fields with the same positive sign minimum and no defects). In this case, the topological defects that develop in the field ϕ will introduce large gradients for ϕ that are absent in the VEV v and will therefore give large contributions in equation (6). Considering both solutions for the VEV allows us to separate the effect of topology Ω_{top} .

In figure 1 we plot the volume average, equation (6), for some different parameter choices. We see that 1) In all cases, there is an initial spike in the energy density at the start of the phase transition; 2) for this choice of parameters, the energy scale is of the order of $\Omega_V(z=0) \sim 10^{-9}$. To obtain the energy scale $\mathcal{O}(1)$ required for the accelerating expansion, we have to extrapolate in the model parameters; 3) the effect of changing β_* is very well approximated² by the change in amplitude given by the equation (4), which means that $\langle V \rangle \propto \beta_*^2$. Considering all of the parameter scalings, we find $V \propto \frac{\beta_*^2 L_C^5}{a_*^9}$; and 4) in all cases where the domain wall contribution (topology) is included in the energy density, there is a rapidly increasing potential energy in the later stages of the phase transition.

We define the effective equation of state through the energy contribution of the scalar field [13], ρ , as

$$\omega = \frac{1}{3} \frac{d \log \rho}{d \log a} - 1. \quad (8)$$

Figure 2 shows the effective equation of state parameter carrying the energy of the potential in equation (6). We are neglecting the kinematic and gradient contributions $\Omega_{\text{kin}}, \Omega_{\text{grad}}$ for now. In all cases in figure 2 where we remove the effect of the domain walls (topology), the effective equation of state parameter is centred around -1 . There is a slight drift from a phantom regime ($\omega < -1$) to larger values and some oscillation around this drift. An earlier phase transition seems to reduce the slope of the drift. In the case where the effect of domain walls (topology) is included, the topological defects seem to push the equation of state parameter towards the phantom regime. Usually in literature, having a ‘phantom crossing’ is considered problematic[15], but their argument does not apply in our case due to the degravitation mechanism and matter coupling.

Finally, there are the kinetic and gradient terms, Ω_{kin} and Ω_{grad} , which we also expect to contribute to the Friedmann equation. Our results show that they both peak early in the phase transition, after which the gradi-

¹ Using v_\pm instead, without a prescription for smoothing of T , we would have contributions from arbitrarily small underdense regions.

² We have generally found a very small effect of varying β on the scalar field configuration [11].




FIG. 1: Energy density contribution from the scalar field potential, normalised to the critical density of the universe ($\Omega = \rho/\rho_c$), as a function of redshift z . The left panel shows the case where the domain wall contribution is removed. Right panel includes the contribution of the domain walls.




FIG. 2: Effective equation of state parameters as a function of cosmological redshift z . The kinematic and gradient contributions (7) are neglected here. The left panel shows the case where the domain wall contribution is removed. The right panel includes the domain wall contribution. Error bars show 95 % confidence intervals due to the intrinsic scatter in the data. The band labelled DESI+ demonstrates the 68% preferred region for the flat $\omega_0\omega_a$ CDM model, presented in [14].

ent contribution decays rapidly and Ω_{kin} dominates between the two. We also find that the vacuum v generally evolves slowly $|\dot{v}| \ll |\dot{\phi}|$ and so Ω_{kin} is the contribution of the scalar field kinetic energy $\dot{\phi}^2/2$ coming from the oscillations of the field. We find that the kinetic term is larger than the potential contribution, equation (6), for all of the models. The kinetic energy decays as $\rho_{\text{kin}} \sim a^{-2}$.

Comparing different simulations, we see no effect of parameter variation on the relative amplitude of the kinetic and potential components. However, we do see a smaller injection of kinetic energy at the start of the phase transition in the case of stronger screening (or larger a_*). The height of the peak compares to the plateau as $\Omega_{\text{kin}}^{(\text{peak})}/\Omega_{\text{kin}} \propto \sqrt{L_C/a_*}$. This factor is $\Omega_{\text{kin}}^{(\text{peak})}/\Omega_{\text{kin}} \approx 2.7$ for the fiducial simulation ($z_*, L_C, \beta_* = 0.1, 1, 8$). While for the same parameters, today ($z = 0$), $\Omega_{\text{kin}}/\Omega_V \sim 4$.

From the above, we can see that in order to dynamically generate an energy density that mimics the cosmological constant by the mechanism proposed here, the following problems must be solved by moving to an appropriate region of the parameter space

- A The initial spike of energy density occurring at the start of the phase transition.
- B The late rise in energy owing to a growing contribution of the domain walls to the energy budget.
- C The large amplitude of the kinetic energy relative to the potential $\Omega_{\text{kin}}/\Omega_V$.
- D The small amplitude of the energy density Ω_V .

Therefore, we note the following part of the symmetron parameter space as interesting with regards to issues A-D: small critical energy densities ρ_* (equivalently large $a_* > 1$) and small interaction length scales L_C . Finally

the force strength β_* is chosen to obtain the correct energy density amplitude.

By having small critical energy densities ρ_* we ensure that only the least dense regions, which are mostly disjunct up to the present time undergo the phase transition. Since the domains are mostly disjunct, we expect a suppression of the final increase in energy density (B) that is caused by the formation of domain walls. $a_* > 1$ increasingly depends on nonlinearities in the density field to undergo collapse before the current time, so the scale L_C is chosen to make the field interact with these small-scale nonlinearities. In this regime, we expect an almost quasistatic transition, since the timescale of the scalar field is smaller and the spatial scale resolves the filaments. We therefore expect a suppression of the initial spike (A) that we understand to be caused by an initial period of out-of-equilibrium dynamics for the scalar.

We do not see a clear effect of varying the parameters on the relative amplitude between the kinetic and potential energies, but we have found nonlinearities in the dependence on z_* and L_C , that make the extrapolation difficult. As the initial spike (A) is the first injection of kinetic energy into the field, we expect a suppression of the relative amplitude of the kinetic energy (C). The problem (C), may therefore be resolved in the interesting limit of small L_C, z_* .

To be explicit, we make the following estimates: To remove the initial spike (A), we assume that it is sufficient that the spatial scale of the symmetron is of order $L_C \lesssim \mathcal{O}(\text{kpc})$. For such a choice, considering the inferred scaling of $a_{\text{SSB}} \propto L_C^{0.22} a_*^{0.59}$, keeping $z_{\text{SSB}} \sim 0.6$, we set $a_* \sim 12$. This is motivated by cosmological data, where the cosmic acceleration happens around $0.4 \lesssim z \lesssim 1$ [1]. Finally, to obtain the correct energy amplitude (D), given the reported scaling of V , we set $\beta_* = 10^{18}$. Although this may seem unreasonably large compared to the values of $\beta_* \sim \mathcal{O}(1)$ typically considered in the literature, we point out that the screening is also set very large, so that the force operates only inside of the least dense voids, where the clustering constraints are at their worst and the effect of a fifth force is minimal. Moreover, this force strength is still 10^7 times weaker than the weak force and 10^{18} times weaker than electromagnetism.

Despite the field being strongly screened, we expect observables such as a modification of the density profile within large cosmic voids, which is expected to give clear signatures in observables such as the Integrated Sachs-Wolfe (ISW) effect [11, 16–18]. We expect upcoming constraints on the equation of state parameter from among other the continuation of the DESI survey [14]. And the oscillating energy density and topological defects can leave observable imprints in gravitational waves [12, 19].

The appeal of this mechanism is manifold:

1. The nearly quasistatic evolution of the field addresses the smallness problem of the cosmological constant by suppressing the energy density contribution to the Friedmann equation relative to that of the original potential (2)
2. Since the phase transition is triggered by nonlinearities in the overdensity field, the mechanism partially solves the coincidence problem by making ‘now’ (i.e. $z \lesssim 1$) a generic time for the accelerated expansion
3. The addition of a fifth force at a strength scale midway between electromagnetism and gravity improves the hierarchy problem of why gravity is uniquely weak, and provides a more continuous range of force scales among the five forces
4. In terms of minimality, the symmetron has been considered as a dark matter candidate for similar parameter choices for the interaction scale $L_C \sim \text{kpc}$ [7, 8, 10], in which case it behaves as a fuzzy dark matter with an extended mass spectrum that depends on the environment
5. The relatively large energy scale of the degravitated potential V_0 and the phantom behaviour caused by the formation of topological defects may allow the connection of the low energy, late-time acceleration with the higher energy scale expansion during inflation [20] in a cyclic universe scenario. In this scenario, the symmetron has the additional advantage of being symmetry restored during reheating. Looking at the degravitated potential V_0 from the equation (4), we see that it has a scale $V_0 \sim 10^{16} \cdot \rho_{c,0} \sim 10^{19} \text{ eV/cm}^3$, which can be made available in the topological defects or upon symmetry restoration.

In conclusion, the large discrepancy between the cosmological constant expected from quantum field theory and that required to explain the cosmological acceleration motivates: 1) the introduction of a degravitation mechanism of constant term contributions to the Friedmann equation and 2) dynamical drivers of the cosmological acceleration. In this work we have introduced a model where there is a late-time, environment-dependent phase transition in the dark sector whose dynamic energy component drives the late-time accelerated expansion. We have studied the evolution and parameter scaling of the energy components within the parameter space computationally accessible to our simulations. By extrapolation, we point to a specific part of the symmetron parameter space, $(L_C, a_*, \beta_*) \sim (\text{kpc}, 12, 10^{18})$, which can account for the late-time cosmic acceleration.

Acknowledgements: $\emptyset.C.$ thanks Martin Kunz for discussions. We thank the Research Council of Norway for their support and the resources provided by UNINETT Sigma2 – the National Infrastructure for High Performance Computing and Data Storage in Norway, as well as the Swiss National Supercomputing Centre for their support under project IDs 301367 and s1051 respectively.

-
- [1] Planck, *Astronomy & Astrophysics* **641**, A6 (2020), ISSN 0004-6361, 1432-0746, arXiv: 1807.06209, URL <http://arxiv.org/abs/1807.06209>.
- [2] S. Weinberg, *Reviews of Modern Physics* **61**, 1 (1989), publisher: American Physical Society, URL <https://link.aps.org/doi/10.1103/RevModPhys.61.1>.
- [3] E. Abdalla, G. F. Abellán, A. Aboubrahim, A. Agnello, O. Akarsu, Y. Akrami, G. Alestas, D. Aloni, L. Amendola, L. A. Anchordoqui, et al., *Journal of High Energy Astrophysics* **34**, 49 (2022), ISSN 22144048, arXiv:2203.06142 [astro-ph, physics:hep-ph], URL <http://arxiv.org/abs/2203.06142>.
- [4] J. S. Bullock and M. Boylan-Kolchin, *Annual Review of Astronomy and Astrophysics* **55**, 343 (2017), ISSN 0066-4146, 1545-4282, arXiv: 1707.04256, URL <http://arxiv.org/abs/1707.04256>.
- [5] K. Hinterbichler and J. Khoury, *Physical Review Letters* **104**, 231301 (2010), ISSN 0031-9007, 1079-7114, arXiv: 1001.4525, URL <http://arxiv.org/abs/1001.4525>.
- [6] K. Hinterbichler, J. Khoury, A. Levy, and A. Matas, *Physical Review D* **84**, 103521 (2011), ISSN 1550-7998, 1550-2368, URL <https://link.aps.org/doi/10.1103/PhysRevD.84.103521>.
- [7] C. Burrage, E. J. Copeland, and P. Millington, *Physical Review D* **95**, 064050 (2017), ISSN 2470-0010, 2470-0029, arXiv:1610.07529 [astro-ph, physics:gr-qc, physics:hep-ph], URL <http://arxiv.org/abs/1610.07529>.
- [8] C. Burrage, E. J. Copeland, C. Kading, and P. Millington, *Physical Review D* **99**, 043539 (2019), ISSN 1550-7998/0556-2821, aDS Bibcode: 2019PhRvD..99d3539B, URL <https://ui.adsabs.harvard.edu/abs/2019PhRvD..99d3539B>.
- [9] A. P. Naik and C. Burrage, *Journal of Cosmology and Astroparticle Physics* **2022**, 020 (2022), ISSN 1475-7516, arXiv:2205.00712 [astro-ph], URL <http://arxiv.org/abs/2205.00712>.
- [10] C. Kading, *Astronomy* **2**, 128 (2023), eprint: 2304.05875.
- [11] Ø. Christiansen, F. Hassani, and D. F. Mota, *asimulation: Domain formation and impact on observables in resolved cosmological simulations of the (a)symmetron* (2024), arXiv:2401.02410 [astro-ph, physics:gr-qc, physics:hep-th], URL <http://arxiv.org/abs/2401.02410>.
- [12] Ø. Christiansen, J. Adamek, F. Hassani, and D. F. Mota, *Gravitational waves from dark domain walls* (2024), iSSN: 2693-5015, URL <https://www.researchsquare.com/article/rs-3857229/v1>.
- [13] Ø. Christiansen, F. Hassani, M. Jalilvand, and D. F. Mota, *Journal of Cosmology and Astroparticle Physics* **2023**, 009 (2023), ISSN 1475-7516, publisher: IOP Publishing, URL <https://dx.doi.org/10.1088/1475-7516/2023/05/009>.
- [14] D. Collaboration, A. G. Adame, J. Aguilar, S. Ahlen, S. Alam, D. M. Alexander, M. Alvarez, O. Alves, A. Anand, U. Andrade, et al., *DESI 2024 VI: Cosmological Constraints from the Measurements of Baryon Acoustic Oscillations* (2024), arXiv:2404.03002 [astro-ph], URL <http://arxiv.org/abs/2404.03002>.
- [15] A. Vikman, *Physical Review D* **71**, 023515 (2005), ISSN 1550-7998, 1550-2368, arXiv:astro-ph/0407107, URL <http://arxiv.org/abs/astro-ph/0407107>.
- [16] A. Kovács, C. Sánchez, J. García-Bellido, S. Nadathur, R. Crittenden, D. Gruen, D. Huterer, D. Bacon, J. DeRose, S. Dodelson, et al., *Monthly Notices of the Royal Astronomical Society* **465**, 4166 (2017), ISSN 0035-8711, 1365-2966, arXiv:1610.00637 [astro-ph], URL <http://arxiv.org/abs/1610.00637>.
- [17] A. Kovács, *Monthly Notices of the Royal Astronomical Society* **475**, 1777 (2018), ISSN 0035-8711, 1365-2966, arXiv:1701.08583 [astro-ph], URL <http://arxiv.org/abs/1701.08583>.
- [18] A. Kovács, C. Sánchez, J. García-Bellido, J. Elvin-Poole, N. Hamaus, V. Miranda, S. Nadathur, T. Abbott, F. B. Abdalla, J. Annis, et al., *Monthly Notices of the Royal Astronomical Society* **484**, 5267 (2019), ISSN 0035-8711, 1365-2966, arXiv:1811.07812 [astro-ph], URL <http://arxiv.org/abs/1811.07812>.
- [19] A. Afzal, G. Agazie, A. Anumarlapudi, A. M. Archibald, Z. Arzoumanian, P. T. Baker, B. Bécsy, J. J. Blanco-Pillado, L. Blecha, K. K. Boddy, et al., *The Astrophysical Journal Letters* **951**, L11 (2023), ISSN 2041-8205, 2041-8213, arXiv:2306.16219 [astro-ph, physics:gr-qc, physics:hep-ph], URL <http://arxiv.org/abs/2306.16219>.
- [20] A. Vilenkin, *Physical Review Letters* **72**, 3137 (1994), ISSN 0031-9007, arXiv:hep-th/9402085, URL <http://arxiv.org/abs/hep-th/9402085>.