

Gravitational algebras and the generalized second law

Thomas Faulkner^{*1} and Antony J. Speranza^{†1}

¹*Department of Physics, University of Illinois, Urbana-Champaign, Urbana IL 61801, USA*

May 1, 2024

Abstract

We derive the generalized second law (GSL) for arbitrary cuts of Killing horizons from the perspective of crossed-product gravitational algebras, making use of a recent proposal by one of us for the construction of local gravitational algebras. This construction relies on the existence of a state whose modular flow is geometric on the horizon. In both free and interacting quantum field theories, such states are guaranteed to exist by the properties of half-sided translations on the horizon. Using geometric identities derived from the canonical analysis of general relativity on null surfaces, we show that the crossed product entropy agrees with the generalized entropy of the horizon cut in a semiclassical limit, and further reproduce Wall's result relating the GSL to monotonicity of relative entropy of the quantum field algebras. We also give a novel generalization of the GSL for interacting theories in asymptotically flat spacetimes involving the concept of an algebra at infinity for a half-sided translation, which accounts for triviality of the algebra of fields smeared only on the horizon. Going beyond the semiclassical limit, we compute subleading corrections to the crossed product entropy, but are unable to determine if the GSL continues to hold after accounting for these. We speculate that an improved GSL could follow from a hidden subalgebra structure of the crossed products, assuming the existence of an operator-valued weight between horizon cut algebras.

^{*}tomf@illinois.edu

[†]asperanz@gmail.com

Contents

1	Introduction	3
1.1	Summary of results	5
2	Crossed product and modular theory	8
3	Semiclassical generalized second law	11
3.1	Half-sided modular inclusions for horizon cuts	12
3.2	Geometric horizon identities	17
3.3	Generalized second law for semiclassical states	23
4	Algebra at infinity	28
4.1	Semiclassical GSL	29
4.2	Gravitational algebra	36
4.3	Rotating black holes and Unruh states	43
5	Perturbative corrections to crossed product entropy	45
6	Monotonicity from crossed product subalgebras	52
7	Discussion	58
A	Tomita operators for crossed product algebra	63
B	Null surfaces	67
B.1	Null surface geometry	67
B.2	Gaussian null coordinates	68
B.3	Quasi-Killing vectors	69
C	Canonical formalism and gravitational charges	70

1 Introduction

Generalized entropy is a fundamental quantity in semiclassical quantum gravity providing a link between dynamical geometry, thermodynamics, and entanglement. It originally appeared in the context of black hole thermodynamics, where Bekenstein recognized that one could violate the second law of thermodynamics by depositing entropy into a black hole unless the black hole possessed an intrinsic entropy proportional to its horizon area [1, 2]. The resulting generalized entropy consists of a sum

$$S_{\text{gen}} = \frac{A}{4\hbar G_N} + S_{\text{out}}, \quad (1.1)$$

with the Bekenstein-Hawking area term representing the intrinsic black hole entropy, and S_{out} denoting the matter entropy outside of the black hole. The generalized second law is then the statement that S_{gen} cannot decrease under time evolution, despite the fact that the matter entropy S_{out} may decrease as matter falls across the horizon.

The connection to entanglement comes from interpreting S_{out} as the fine-grained entanglement entropy of quantum fields restricted to the black hole exterior [3–6]. Not only does this give a concrete definition of S_{out} , it also leads to a UV-finite notion of entropy in quantum gravity, owing to an argument of Susskind and Uglum [7] that the divergences in the quantum field entanglement entropy cancel against the renormalization of G_N in S_{gen} . This cancellation of divergences has been verified in a numerous examples [8–12], and ultimately motivates interpreting S_{gen} as a true count of the microscopic quantum gravitational degrees of freedom of the black hole.

Since entanglement entropy can be associated with generic causally complete subregions in quantum field theory, it is natural to consider generalized entropy for arbitrary subregions in semiclassical quantum gravity [12–14]. In this more general context, the area of the codimension-2 boundary of a Cauchy surface for the subregion takes the place of the horizon area in the Bekenstein-Hawking term in S_{gen} . By viewing the generalized entropy as the quantum analog of the area in semiclassical gravity, one may upgrade various classical theorems in general relativity to quantum versions. Bekenstein’s generalized second law is one such example, being the quantum version of Hawking’s classical area theorem [15, 16]. The quantum focusing conjecture as a semiclassical version of the classical focusing theorem is another prominent example [12], which has far-reaching applications including quantum entropy bounds (see [17] for a review) and field theoretic inequalities such as the quantum null energy condition [18–20]. Generalized entropy also features in holographic entanglement entropy, where the Ryu-Takayanagi and quantum extremal surface formulas relate the generalized entropy of entanglement wedges in the bulk to entanglement entropies of subregions in the boundary CFT dual [21–26].

One obstruction to obtaining rigorous proofs of semiclassical entropy relations such as the quantum focusing conjecture is the occurrence of UV divergences in entanglement entropies in quantum field theory. These can be eliminated by employing regulators and constructing renormalized entropies, but this procedure can often obscure important quantum information theoretic relations such as strong subadditivity. On the other hand, the expected UV-

finiteness of the generalized entropy suggests that coupling the theory to gravity results in a universal renormalized notion of entropy in semiclassical quantum gravity. If this is the case, there ought to be a continuum description of this renormalized entropy that is manifestly regulator-independent.

Recent progress in this direction comes from considerations of von Neumann algebras in quantum gravity. These began with the work of Leutheusser and Liu [27, 28], who considered the strict large N limit of a holographic CFT above the Hawking-Page transition, and found an emergent type III₁ von Neumann algebra, which they took to be a signature of the emergence of a black hole horizon in the bulk dual. Subsequently, Witten argued that $\frac{1}{N}$ corrections implement a crossed product on the algebra, resulting in a type II_∞ von Neumann algebra, which notably possesses a well-defined notion of renormalized entropy [29]. This entropy agrees with the generalized entropy of the horizon in semiclassical states of the algebra up to an overall constant, which can be interpreted as a universal large background value of the entropy [30]. The agreement between algebraic entropy and generalized entropy is closely tied to the implementation of the gravitational constraints, and therefore holds directly in the semiclassical bulk description. This allows for the construction of type II algebras and associated entropies in configurations with no currently known holographic dual description, such as the static patch of de Sitter, or general black holes in asymptotically flat space [31, 32].

The bulk semiclassical gravity picture further suggests that generic subregions give rise to type II gravitational algebras with a well-defined renormalized entropy [33]. These algebras each contain a type III₁ subalgebra describing quantum fields within the subregion in a fixed background geometry, as one expects from the $G_N \rightarrow 0$ limit, which suppresses gravitational backreaction. The emergence of such type III₁ subregion algebras in the large N limit of holography has been explored in a number of recent works [34–37]. The local gravitational algebras also contain additional operators representing the asymptotic ADM energy or the energy of a localized observer degree of freedom, which serve as anchors to dress the local operators in the gravitational descriptions. Imposing the gravitational constraints on both sets of operators results in a type II crossed product algebra that describes the fields and semiclassical geometry of the subregion. As in the case of Killing horizons, the renormalized entropy computed for semiclassical states in this algebra agrees with the generalized entropy up to a state-independent constant. Hence, crossed product algebras are a viable framework for a continuum, regulator-independent description of semiclassical geometry and entropies.

Since crossed-product gravitational algebras have manifestly UV-finite renormalized entropies, they potentially could provide a new set of tools for analyzing semiclassical entropy relations such as the quantum focusing conjecture [12]. The goal of the present work is to initiate such investigations in a more restricted context by revisiting the generalized second law (GSL) for Killing horizons. In addition to their global boost symmetry, Killing horizons possess an enhanced near-horizon symmetry that enables a detailed analysis of the vacuum entanglement properties of quantum fields in the spacetime containing the horizon. Due to this symmetry, one can obtain an explicit expression for the vacuum modular Hamiltonian of any subregion whose entangling surface lives on an arbitrary cut of the horizon [38–40].

These subregions therefore provide nontrivial examples of the *geometric modular flow conjecture* that is necessary in the construction of gravitational algebras for arbitrary subregions proposed by Jensen, Sorce, and Speranza (JSS) [33]. Hence, a second motivation for investigating the GSL is to verify the general construction proposed by JSS in situations where the modular Hamiltonian is not determined by global spacetime symmetries.

Wall gave a proof of the GSL for rapidly evolving perturbed Killing horizons by relating the generalized entropy of cuts of the horizon to the relative entropy of quantum fields in the region defined by the cut [38]. After this, the GSL becomes equivalent to the monotonicity of relative entropy under algebra inclusions. The first result of the present paper is to reproduce this proof using crossed product gravitational algebras. The entropy of certain semiclassical states for the gravitational algebra manifestly takes the form of a relative entropy for the quantum fields, up to a correction involving the asymptotic gravitational charge. Hence, by performing the crossed product separately at two different cuts of a horizon, we immediately conclude in section 3.3 a monotonicity result for the entropies of these states, thereby showing that crossed product algebras can reproduce Wall’s proof of the generalized second law.

This raises the question of whether crossed product entropies can lead to stronger monotonicity results beyond the strict semiclassical limit for the quantum state. The entropy formula for type II gravitational algebras receives corrections to the semiclassical expression due to entanglement between the asymptotic gravitational charges and quantum field degrees of freedom. One can ask whether the crossed product entropy continues to satisfy monotonicity after these corrections are included. We explore this question by computing the corrections to the semiclassical entropy formula in section 5. We demonstrate a number of properties of these corrections including explicit bounds on their size, but are unable to determine if the crossed product entropy satisfies a monotonicity result after accounting for these corrections. We also explore the possibility in section 6 that the crossed product algebras could exhibit a subalgebra structure that is obscured in their standard presentation. This would follow from the existence of an operator-valued weight between horizon cut algebras, and assuming its existence, we derive a modified monotonicity result for crossed product entropies.

The overall picture that emerges from these investigations is that crossed products for subregions indeed capture a useful notion of gravitational entropy, and they provide powerful tools for investigating entropy in semiclassical quantum gravity. They raise the possibility of finding strengthened data-processing inequalities associated with the change in relative entropy between horizon cuts, or otherwise characterizing temporary violations of the second law by gravitational effects. Ultimately, they could lead to a better understanding entropy inequalities such as the quantum focusing conjecture, consequences of causality in quantum field theory, and low energy constraints on quantum theories of gravity.

1.1 Summary of results

This paper contains a number of new results concerning crossed product algebras and semiclassical generalized entropy which we briefly summarize here.

Section 2 reviews the definition of the modular crossed product algebra, and goes over the modular theory for the classical-quantum states that are used in the remainder of the paper. We present novel expressions for the Tomita operators for these states, the details of which are given in appendix A, which lead to a simplified derivation of the density matrices for these states originally obtained in [33]. This also produces expressions for the modular conjugations for these states which have not appeared previously.

We then turn to the discussion of gravitational algebras and the GSL for Killing horizons in section 3. We begin in section 3.1 reviewing the argument for the form of the vacuum modular Hamiltonian for arbitrary cuts of a Killing horizon. Versions of this argument have appeared in various forms across several works [38–40], and this section serves as a summary that presents the derivation in a unified manner. We emphasize the connection between the form of this modular Hamiltonian and properties of half-sided translations and half-sided modular inclusions, introduced respectively by Borchers and Wiesbrock [41–43]. Focusing on half-sided translations leads to a derivation of the modular Hamiltonian that holds even in interacting field theories, which specifically does not rely on the existence of bounded operators localized strictly on the black hole horizon.

Following this, we present in section 3.2 various geometric identities associated with the gravitational constraint equations on the black hole horizon. These identities are derived using the canonical analysis of general relativity on null surfaces, the details of which are given in appendix C. This leads to the identification of the global gravitational constraint (3.35) that relates the asymptotic gravitational charges to the bulk generator of modular flow, and which is responsible for producing the crossed product when imposed on the algebra of quantum fields and asymptotic charges. We also note that a semilocal gravitational constraint leads to a geometric identity (3.32) which relates the area of a horizon cut to the one-sided modular Hamiltonian and the late-time area. This identity was originally derived by Wall by directly integrating the Raychaudhuri equation [38]; our derivation instead shows how it arises from the canonical analysis, which directly ties the identity to the underlying diffeomorphism invariance of the gravitational theory. This has the added advantage of transparently identifying the gravitational contribution to the average null energy on the horizon, which in the $G_N \rightarrow 0$ limit reduces to the square of the perturbative horizon shear, equation (3.41). This identity also corresponds to the local Smarr relation discussed by JSS [33], specialized to the present case of a subregion bounded by a cut of a Killing horizon.

We then turn to the construction of gravitational algebras and analysis of the GSL in section 3.3. We show that the gravitational algebra resulting from imposing the global constraint on the horizon cut algebras takes the form of a modular crossed product, using the form of the vacuum modular Hamiltonian derived in section 3.1. We can then directly compute the entropy of classical-quantum states on these algebras, and show using the semi-classical expansion of this entropy (2.14) that the GSL reduces to monotonicity of relative entropy for the underlying quantum field theory algebra inclusion. This resulting GSL relies on choosing an appropriate normalization for the traces defined on the gravitational algebras for different cuts, which in general is not canonically determined. We resolve this by requiring that the trace agree on functions of the asymptotic charge \hat{q} , which is contained in all horizon cut algebras. This implies that we can use the same formula for the trace, (2.9),

for any horizon cut, and it is important that the same quantum field vacuum state $|\Omega\rangle$ can be used in this formula for different horizon cuts.

To simplify the discussion, the analysis of section 3 was restricted to AdS black holes, for which the horizon provides a complete Cauchy surface. Section 4 generalizes the arguments to spacetimes with other asymptotics, such as asymptotically flat or de Sitter black holes. For free fields, this generalization simply consists of tensoring in an additional factor to the algebra associated with fields that pass through \mathcal{I}^+ or the cosmological horizon. However, this description fails for interacting theories, since there is no algebra consisting of fields strictly localized to codimension-1 black hole horizon, and hence naively there is no algebra with which to take the tensor product. In section (4.1), we show that one can handle the case of interacting theories using the concept of an algebra at infinity for the half-sided translation. This leads to a natural definition of a subalgebra associated with fields that never enter the black hole, but with a trivial relative commutant, corresponding to triviality of the horizon algebra. This captures the notion that any operator with finite fluctuations must be smeared in a region near the horizon, and cannot be strictly localized to the horizon. Using the existence of a conditional expectation to this algebra at infinity, we show that one again recovers the semiclassical GSL, generalizing the argument of Wall that used the existence of tensor-factorized states [38].

Having established the structure of this algebra at infinity, we describe in section 4.2 the construction of the gravitational algebra. Adapting the arguments of Kudler-Flam, Leutheusser, and Satishchandran (KFLS) [32], we show that the gravitational constraints can separately be imposed on the algebra at infinity and on the horizon. Once this is done, there is a final matching constraint associated with the junction where these null surfaces meet on the Penrose diagram. This condition takes the form of a constraint, equation (4.12), matching the two time variables arising in the separate crossed products for each horizon. Along with the matching conditions for the asymptotic charges, this eliminates the asymptotic horizon area in favor of the asymptotic ADM mass as the extra gravitational charge in the algebra. We discuss conditions for which the final algebra remains semifinite and possesses a trace, and show that these are satisfied in the example of the Schwarzschild black hole in the Hartle-Hawking vacuum. Owing to the existence of the conditional expectation to the algebra at infinity, we are able to conclude that the crossed product algebras satisfy a global second law (4.35) which generalizes a similar result obtained in [30], and holds for all states, without any semiclassical assumption. We also note that the matching constraint can be used to eliminate the observer in the discussion of the algebra for the Schwarzschild-de Sitter black hole considered by KFLS [32]. In section 4.3, we point out some subtleties that may arise in obtaining a semifinite algebra at infinity when working with asymptotically flat rotating black holes or black holes formed from collapse.

The versions of the GSL obtained in sections 3.3 and 4.2 rely on the semiclassical approximation of the quantum states. This naturally leads to the question of whether crossed product algebras satisfy a GSL beyond the semiclassical limit. To begin addressing this question, we compute in section 5 the subleading corrections to the semiclassical entropy formula (2.14). Noting that the classical-quantum state $|\Phi, f\rangle$ can be viewed as the image under a Petz-recovery channel from the quantum field theory state $|\Phi\rangle$ to a state on the crossed

product, we show that monotonicity of relative entropy for this channel implies that the corrections to the semiclassical crossed product entropy are strictly nonnegative. We then obtain an explicit expression for this correction, given by equation (5.23). We can directly evaluate the first nontrivial contribution to this correction in the semiclassical expansion, which we find is second order in the expansion parameter, and we verify that this correction is positive. Unfortunately, we are unable to determine if the crossed product entropy, including these corrections, satisfies a monotonicity result, but note that if one existed, it would imply an improved data-processing inequality (5.32).

Finally, in section 6, we explore the possibility that the crossed-product GSL could follow from a hidden subalgebra structure for the gravitational algebras. We show that if there is an operator-valued weight [44, 45] between the horizon cut quantum field theory algebras, there is a canonical way to realize the later horizon-cut gravitational algebra as a subalgebra of the earlier gravitational algebra. Although we do not at present have a proof for the existence of this operator-valued weight, we conjecture one should exist by analogy with the case of a split inclusions, for which the operator-valued weight is guaranteed. Assuming its existence, the result crossed product subalgebra structure should imply a version of the GSL stemming from monotonicity of relative entropy for this algebra inclusion. In order to leverage this, we consider a free energy quantity (6.15) defined as a relative entropy between the state of interest and the dual weight that naturally arises in the crossed product construction. This leads to the monotonicity results (6.17) and (6.22), which are nontrivial, but not quite the GSL we were seeking. However, we note that in the semiclassical limit, these inequalities reduce to the GSL, and we leave open the possibility that they may lead to the desired result with a more careful analysis.

We conclude in section 7 by outlining some questions raised by the present work, and describing ideas for future applications to problems in semiclassical quantum gravity.

Note added: During the final stages of writing this paper, [46] appeared which contains some overlap with the results presented here.

2 Crossed product and modular theory

We begin with a brief overview of the crossed product construction in semiclassical gravity, before describing the application of this construction to black hole horizons in section 3.

Several recent investigations into semiclassical quantum gravity have noted the relevance of crossed products in the algebraic formulation of the theory [29–33]. The $G_N \rightarrow 0$ limit of quantum gravity admits a description in terms of quantum field theory in curved space-time, in which local field algebras are assigned to arbitrary causally complete subregions. These algebras are generically von Neumann algebra factors of type III₁, and this structure is expected to persist to all orders in the $\sqrt{G_N}$ expansion [27–29]. However, there exist certain nonlocal observables such as the global ADM Hamiltonian that interact nontrivially with these local algebras when going beyond the strict $G_N \rightarrow 0$ limit. In particular, the

gauge constraints arising from diffeomorphism invariance yield relations between the nonlocal observables and local algebras which affect the structure of the algebras even at $G_N = 0$. Crossed product algebras are the result of imposing the constraints on the combined system of local algebras and global observables.

The construction of gravitational crossed product algebras begins with the algebra \mathcal{A}_{QFT} of quantum fields restricted to a causally complete subregion R in spacetime. The subregions of interest in the present work are the exterior of a bifurcation surface of an eternal black hole, or more generally the exterior of a cut of the future horizon. \mathcal{A}_{QFT} acts on a Hilbert space \mathcal{H}_{QFT} , and we take it to be a factor of type III₁ in accord with arguments for the universal type of subregion algebras in quantum field theory [47–49]. The additional nonlocal degree of freedom is an asymptotic gravitational charge, which will generally be proportional to the perturbation of late-time area of a black hole horizon when working with Killing horizons, and can be related to other asymptotic charges such as the ADM Hamiltonian of the spacetime. We represent this asymptotic charge as minus the position operator acting on the Hilbert space $\mathcal{H}_{\text{asy}} = L^2(\mathbb{R})$, which reflects the semiclassical assumption that the spectrum of \hat{q} is assumed to be continuous and unbounded above and below. The crossed product algebra $\widehat{\mathcal{A}}$ is the subalgebra of $\mathcal{A}_{\text{QFT}} \otimes \mathcal{B}(\mathcal{H}_{\text{asy}})$ that commutes with the complementary asymptotic charge associated, for example, with the left asymptotic area of a two-sided black hole. On the Hilbert space $\widehat{\mathcal{H}} = \mathcal{H}_{\text{QFT}} \otimes \mathcal{H}_{\text{asy}}$, this left asymptotic charge is represented by the operator

$$\hat{q}_L = \hat{q} + h_\Omega, \quad (2.1)$$

where $h_\Omega = -\log \Delta_\Omega$ is the modular Hamiltonian of a specific vacuum state $|\Omega\rangle \in \mathcal{H}_{\text{QFT}}$. The interpretation of \hat{q}_L as a left asymptotic charge follows from a diffeomorphism constraint in semiclassical quantum gravity, and additionally relies on the modular flow generated by h_Ω agreeing with the diffeomorphism flow along a vector ξ^a on a Cauchy surface for the subregion R . As we review in section 3.1, vacuum states of the horizon average null energy operators satisfy this property for algebras associated to cuts of the Killing horizon. We will refer to $|\Omega\rangle$ as the vacuum state for the QFT algebra since it is a vacuum for the average null energy operators, although it may not coincide with a global minimal energy state.

$\widehat{\mathcal{A}}$ can be expressed as

$$\widehat{\mathcal{A}} = \langle e^{i\hat{p}h_\Omega} \mathbf{a} e^{-i\hat{p}h_\Omega}, \hat{q} \rangle, \quad \mathbf{a} \in \mathcal{A}_{\text{QFT}}, \quad (2.2)$$

where the notation $\langle \cdot \rangle$ means the von Neumann algebra generated by the displayed operators by taking a double commutant. The commutant algebra $\widehat{\mathcal{A}}'$ naturally describes operators in the causal complement R' dressed to the left asymptotic charge \hat{q}_L . It consists of the operators

$$\widehat{\mathcal{A}}' = \langle \mathbf{a}', \hat{q} + h_\Omega \rangle, \quad \mathbf{a}' \in \mathcal{A}'_{\text{QFT}}. \quad (2.3)$$

We are interested in characterizing entropies of states on the algebra $\widehat{\mathcal{A}}$ and relating them to generalized entropies for the Killing horizon. The specific states we will focus on are the classical-quantum states $|\widehat{\Phi}\rangle = |\Phi, f\rangle := |\Phi\rangle_{\text{QFT}} \otimes f(q)$. The density matrix for this state can be computed by first determining the Tomita operator $S_{\widehat{\Phi}}$, defined as the unbounded,

antilinear operator which acts on states of the form $\widehat{\mathbf{a}}|\widehat{\Phi}\rangle$, $\widehat{\mathbf{a}} \in \widehat{\mathcal{A}}$ as

$$S_{\widehat{\Phi}}\widehat{\mathbf{a}}|\widehat{\Phi}\rangle = \widehat{\mathbf{a}}^\dagger|\widehat{\Phi}\rangle \quad (2.4)$$

By solving this relation for a spanning set of operators of the form $\widehat{\mathbf{a}} = e^{i\hat{p}h_\Omega}\mathbf{a}e^{-i\hat{p}h_\Omega}e^{iu\hat{q}}$ (see appendix A), one finds that

$$S_{\widehat{\Phi}} = J_{\Phi|\Omega}e^{-i\hat{p}h_\Omega}\widehat{\mathbf{s}}_{\widehat{\Phi}}\widehat{\mathbf{s}}'_{\widehat{\Phi}} \quad (2.5)$$

$$\widehat{\mathbf{s}}_{\widehat{\Phi}} = e^{i\hat{p}h_\Omega}\Delta_{\Phi|\Omega}^{\frac{1}{2}}f^*(\hat{q} - h_\Omega)e^{\frac{\hat{q}}{2}}e^{-i\hat{p}h_\Omega} \quad (2.6)$$

$$\widehat{\mathbf{s}}'_{\widehat{\Phi}} = \frac{1}{e^{\frac{\hat{q}}{2}}f(\hat{q} + h_\Omega)}J_\Omega J_{\Omega|\Phi}\Delta_{\Omega|\Phi}^{\frac{1}{2}} \quad (2.7)$$

where $\widehat{\mathbf{s}}_{\widehat{\Phi}}$ is a linear operator affiliated with $\widehat{\mathcal{A}}$ and $\widehat{\mathbf{s}}'_{\widehat{\Phi}}$ with $\widehat{\mathcal{A}}'$.

In the special case of a vacuum classical-quantum state $|\widehat{\Omega}\rangle = |\Omega, f\rangle$ with real wavefunction $f^*(\hat{q}) = f(\hat{q})$, the expression simplifies dramatically such that the polar decomposition $S_{\widehat{\Omega}} = J_{\widehat{\Omega}}\Delta_{\widehat{\Omega}}^{\frac{1}{2}}$ becomes manifest, with

$$J_{\widehat{\Omega}} = J_\Omega e^{-i\hat{p}h_\Omega}, \quad \Delta_{\widehat{\Omega}} = [|f(\hat{q})|^2 e^{\hat{q}}] \cdot \left[\frac{e^{-\hat{q}}\Delta_\Omega}{|f(\hat{q} + h_\Omega)|^2} \right] = \rho_{\widehat{\Omega}} \cdot (\rho'_{\widehat{\Omega}})^{-1}. \quad (2.8)$$

The density matrix $\rho_{\widehat{\Omega}}$ is an operator affiliated with $\widehat{\mathcal{A}}$, and can be used to define a trace on the algebra according to [29]

$$\text{Tr}(\widehat{\mathbf{a}}) = \langle \Omega, f | \rho_{\widehat{\Omega}}^{-1} \widehat{\mathbf{a}} | \Omega, f \rangle = 2\pi \langle \Omega, 0_p | e^{-\hat{q}} \widehat{\mathbf{a}} | \Omega, 0_p \rangle, \quad (2.9)$$

where $|0_p\rangle$ is the δ -function normalized zero \hat{p} eigenstate. Note that the density matrix used to define this trace depends only on the operator \hat{q} , and not on the modular operator Δ_Ω . This will be important in section 3.3 when comparing traces for crossed product algebras associated with different subregions. These algebras do not obviously form subalgebras for nested subregions since the modular Hamiltonian h_Ω depends on the choice of subregion. However, all the algebras share the operator \hat{q} , and the density matrix $\rho_{\widehat{\Omega}}$ remains the same in each algebra, leading to a standard choice for its normalization and the normalization of the trace in different crossed product algebras.

Returning to the more general classical-quantum state $|\widehat{\Phi}\rangle$, we obtain the density matrix for $\widehat{\mathcal{A}}$ from the relation $\rho_{\widehat{\Phi}} = \widehat{\mathbf{s}}_{\widehat{\Phi}}^\dagger \widehat{\mathbf{s}}_{\widehat{\Phi}}$. This reproduces the expression derived in [33],

$$\rho_{\widehat{\Phi}} = e^{i\hat{p}h_\Omega} f(\hat{q} - h_\Omega) e^{\hat{q}} \Delta_{\Phi|\Omega} f^*(\hat{q} - h_\Omega) e^{-i\hat{p}h_\Omega}. \quad (2.10)$$

For computations of the entropy, it is convenient to rewrite this density matrix as

$$\begin{aligned} \rho_{\widehat{\Phi}} &= f(\hat{q}) e^{\frac{\hat{q}}{2}} e^{i\hat{p}h_\Omega} \Delta_\Omega^{-\frac{1}{2}} \Delta_{\Phi|\Omega} \Delta_\Omega^{-\frac{1}{2}} e^{-i\hat{p}h_\Omega} e^{\frac{\hat{q}}{2}} f^*(\hat{q}) \\ &= f(\hat{q}) e^{\frac{\hat{q}}{2}} e^{i\hat{p}h_{\Omega|\Phi}} \Delta_{\Omega|\Phi}^{-\frac{1}{2}} \Delta_\Phi \Delta_{\Omega|\Phi}^{-\frac{1}{2}} e^{-i\hat{p}h_{\Omega|\Phi}} e^{\frac{\hat{q}}{2}} f^*(\hat{q}) \\ &= e^{i\hat{p}h_{\Omega|\Phi}} f(\hat{q} - h_{\Omega|\Phi}) e^{\hat{q}} \Delta_\Phi f^*(\hat{q} - h_{\Omega|\Phi}) e^{-i\hat{p}h_{\Omega|\Phi}} \end{aligned} \quad (2.11)$$

where in the second line we have used the cocycle relations $\Delta_\Omega^{-\frac{1}{2}} \Delta_{\Phi|\Omega}^{\frac{1}{2}} = \Delta_{\Omega|\Phi}^{-\frac{1}{2}} \Delta_\Phi^{\frac{1}{2}}$ and the fact that these operators are affiliated with \mathcal{A}_{QFT} , as well as the fact that h_Ω and $h_{\Omega|\Phi}$ generate the same action on operators in \mathcal{A}_{QFT} (see e.g. [33, Appendix C]). The logarithm is then given by

$$-\log \rho_{\hat{\Phi}} = -h_{\Omega|\Phi} - \hat{q} - \log \left(f(\hat{q}) e^{i\hat{p}h_{\Omega|\Phi}} \Delta_\Phi e^{-i\hat{p}h_{\Omega|\Phi}} f^*(\hat{q}) \right). \quad (2.12)$$

Because the remaining operators inside the logarithm do not commute, the exact expression for the entropy, described in section 5, is somewhat complicated. The reason for the complications is that despite appearances, the state $|\Phi, f\rangle$ describes a state with entanglement between the quantum field degrees of freedom and the asymptotic charge \hat{q} , coming from the dressing factors $e^{i\hat{p}h_\Omega}(\cdot)e^{-i\hat{p}h_\Omega}$ for quantum field theory operators in the crossed product algebra.

However, when comparing to standard treatments of the generalized entropy, there is a useful further restriction on the class of states. This involves taking the wavefunction $f(q)$ to be slowly varying, or, alternatively, taking its Fourier transform $\tilde{f}(p)$ to be sharply peaked at low momentum. Since \hat{p} has the interpretation of the time shift generated by the asymptotic charge, these states describe semiclassical geometries in which this global time variable is well-localized. Away from this limit, the quantum field operators experience significant smearing in time due to the factors of $e^{i\hat{p}h_\Omega}$ acting on the wavefunction $f(q)$. In keeping with the notation of previous works [30, 31, 33], we will refer to states with $f(q)$ slowly varying as *semiclassical states*.

For semiclassical states, it is valid to treat the operator \hat{p} as a small parameter, and expand the logarithm in (2.12) in powers of \hat{p} . Since we will be taking an expectation value in the state $|f\rangle$, each factor of \hat{p} in the expansion will eventually act on a wavefunction, resulting in a suppression by derivatives of f (see section 5 for a precise derivation of this expansion). The strict semiclassical entropy comes from keeping only the leading $\mathcal{O}(\hat{p}^0)$ term in the expansion, for which the term inside the logarithm in (2.12) becomes $|f(\hat{q})|^2 \Delta_\Phi$. The entropy formula in the semiclassical limit then becomes [30, 31, 33]

$$S(\rho_{\hat{\Phi}}) = \langle \hat{\Phi} | -\log \rho_{\hat{\Phi}} | \hat{\Phi} \rangle \approx \langle \hat{\Phi} | -h_{\Omega|\Phi} + h_\Phi - \hat{q} - \log |f(\hat{q})|^2 | \hat{\Phi} \rangle \quad (2.13)$$

$$= -S_{\text{rel}}(\Phi||\Omega) - \langle \hat{q} \rangle_{\hat{\Phi}} + S_f^q \quad (2.14)$$

This entropy formula will be key for arriving at a second law for gravitational crossed product algebras in section 3.3.

3 Semiclassical generalized second law

Having introduced crossed product algebras and derived the entropy formula for semiclassical states, the next task is to relate this formula to the generalized second law for Killing horizons. This involves comparing the entropies computed for a state on two different crossed product algebras associated with different cuts of the future Killing horizon, and demonstrating that this entropy monotonically increases. Doing so invokes two key insights from Wall's original proof of the GSL [38]. The first is the observation that for vacuum states of the horizon




Figure 1: The two-sided AdS black hole is shown in the Penrose diagram above. The bifurcation surface is labeled as the point b , and \mathcal{H}_b^+ denotes the half of the event horizon to the future of b . \mathcal{H}_b^+ serves as a Cauchy surface for the right exterior region R_b , shown in blue. The causal complement R'_b involving the left exterior of the black hole is shown in green. The red dot labeled c_λ denotes a cut of the future horizon, and the portion of the horizon to the future of this cut, \mathcal{H}_λ^+ , is a Cauchy surface for the subregion R_λ in the right exterior, depicted by the red lines. The causal complement of this region is R'_λ , labeled by the orange lines, and consists of the entire left exterior as well as a portion of the black hole interior.

algebra, the modular Hamiltonian for an arbitrary horizon cut generates a geometric flow along the horizon. The algebraic explanation for this property is that nested cuts of the horizon define half-sided modular inclusions [42, 43], which follows, as we review below, from the average null energy condition. The second key insight is the relation between the relative entropy of states of the quantum fields outside a horizon cut and the generalized entropy of the same region. We give a novel derivation of the geometric formulas leading to this identity in section 3.2, utilizing recent analyses of the phase space of general relativity on null Cauchy surfaces. This derivation directly relates the identity to similar relations employed by JSS in the construction of gravitational algebras for general subregions [33]. Applied to semiclassical states for the crossed product algebras, these two insights reduce the GSL to the monotonicity of relative entropy under algebra inclusions, thereby reproducing Wall’s proof.

3.1 Half-sided modular inclusions for horizon cuts

The first step in formulating the GSL in terms of crossed product algebras is to demonstrate that quantum field theory algebras associated with nested horizon cuts define half-sided modular inclusions. This fact was originally established for free and superrenormalizable theories by Wall [38] (although not using the terminology of half-sided modular inclusions), and arguments extending this result to arbitrary interacting quantum field theories have been given by Casini, Teste, and Torroba [39], and Witten [40]. This section serves as a

review of these results, presented in a way that is directly amenable to application to the generalized second law for black holes.

For the initial discussion, we consider the case of a two-sided black hole in asymptotically anti-de Sitter space, shown in figure 1; generalizations including asymptotically flat and de Sitter black holes will be discussed in section 4. AdS asymptotics have the advantage that boundary conditions prevent radiation from escaping through infinity, which implies that the black hole event horizon \mathcal{H} for the right asymptotic region defines a complete Cauchy surface for the spacetime. \mathcal{H} forms one component of a bifurcate Killing horizon, and we denote the associated Killing vector as ξ^a and the bifurcation surface at which ξ^a vanishes as b . The bifurcation surface divides \mathcal{H} into two pieces, \mathcal{H}_b^+ to the future and \mathcal{H}_b^- to the past. \mathcal{H}_b^+ is a Cauchy surface for the right exterior region of the black hole, denoted R_b , and we define $\mathcal{B} = \mathcal{A}(R_b)$ to be the type III₁ von Neumann algebra of quantum fields restricted to this region.

To describe the horizon cut algebra, it is convenient to employ Gaussian null coordinates (u, v, y^A) near \mathcal{H} (see appendix B.2). In these coordinates, \mathcal{H} is located at $u = 0$, and v is the affine parameter for the null horizon generator l^a , which on \mathcal{H} is related to the Killing vector according to $l^a = \frac{1}{\kappa v} \xi^a$, where κ is the horizon surface gravity. The bifurcation surface lies at $u = v = 0$, and all surfaces of constant v are null hypersurfaces associated with ingoing light rays that intersect \mathcal{H} . The coordinate u is an affine parameter for the null tangent vector n^a for these light rays, which is normalized at \mathcal{H} to satisfy $n \cdot l = -1$. A horizon cut c_λ is specified by a function $\lambda(y^A)$ as the surface $v = \lambda(y^A)$ on \mathcal{H} . The cut divides the horizon into a future component \mathcal{H}_λ^+ and a past component \mathcal{H}_λ^- , with \mathcal{H}_λ^+ serving as a Cauchy surface for the region R_λ in the right exterior that is spacelike separated from the cut. The quantum field theory algebra for this region is denoted $\mathcal{C}_\lambda = \mathcal{A}(R_\lambda)$, and the nesting of the regions $R_\lambda \subset R_b$ implies the inclusion of the associated algebras $\mathcal{C}_\lambda \subset \mathcal{B}$.

The crucial object in demonstrating that this inclusion is half-sided modular is the average null energy operator, defined in terms of an integral of the quantum field stress tensor T_{ab} over given null geodesic γ_y on the horizon at fixed y^A as

$$P(y^A) = \int_{\gamma_y} dv T_{ab} l^a l^b = \int_{-\infty}^{\infty} dv T_{vv}(v, y^A). \quad (3.1)$$

We will make use of smeared versions of this operator with respect to a function $\rho(y^A)$, defined as the stress-energy flux associated with the vector ρl^a ,

$$P_\rho = - \int_{\mathcal{H}} T^a_b \rho^b \epsilon_{a\dots} = \int_{-\infty}^{\infty} dv \int dy^A \sqrt{q} T_{vv}(v, y^A) \rho(y^A). \quad (3.2)$$

Here, the orientation conventions for the volume forms are $\epsilon = -l \wedge \eta$, where ϵ is the spacetime volume form, and $\eta = \sqrt{q} dv \wedge dy^1 \wedge \dots \wedge dy^{d-2}$ is the induced volume form on the horizon relative the affinely parameterized null normal, i.e. $\eta = i_l \epsilon$, and \sqrt{q} determinant of the spatial metric on the horizon. P_ρ generates a geometric action on fields localized on the horizon, translating a local operator $\mathcal{O}(v, y^A)$ along the null coordinate to an operator $e^{isP_\rho} \mathcal{O}(v, y^A) e^{-isP_\rho} = \tilde{\mathcal{O}}(v + s\rho(y^A), y^A)$ at the shifted affine coordinate $v + s\rho(y^A)$. The precise form of the translated operator $\tilde{\mathcal{O}}$ depends on its spin and scale dimension, but the

important point is that P_ρ sends local operators on the horizon to local operators at the translated coordinate.

Away from \mathcal{H} , P_ρ generates a nonlocal action on individual operators since the vector field ρ^a does not extend to a Killing vector on the full spacetime. However, its geometric action on horizon operators implies an overall geometric action on the algebras determined by horizon cuts. Schematically, because \mathcal{H}_λ^+ serves as a Cauchy surface for the region R_λ exterior to the cut, the operators in R_λ are dynamically determined by the operators on \mathcal{H}_λ^+ . Hence, because e^{isP_ρ} translates operators located on \mathcal{H}_λ^+ to operators located on the translated future horizon $\mathcal{H}_{\lambda+s\rho}^+$, we should expect it to map the full algebra \mathcal{C}_λ associated with R_λ into the algebra $\mathcal{C}_{\lambda+s\rho}$ associated with the exterior region $R_{\lambda+s\rho}$ for the translated cut. This argument for the geometric action of P_ρ on the algebras \mathcal{C}_λ was first made by Wall in the case of free field theories by taking advantage of the fact that one can obtain a nontrivial horizon algebra by smearing certain local operators with functions supported purely on \mathcal{H} [38]. For interacting field theories, the argument is more subtle because local operators must be smeared over open regions in spacetime in order to produce operators with finite fluctuations [50]. Here, we will describe an argument due to Witten [40] for the validity of this geometric action for interacting field theories; closely related arguments were originally given by Casini, Teste, and Torroba [39].

Although there are no well-defined operators that are smeared solely on the horizon in interacting theories, the unsmeared local operators $\mathcal{O}(x)$ nevertheless have finite matrix elements $\langle \chi | \mathcal{O}(x) | \phi \rangle$ for a sufficiently large number of states $|\phi\rangle, |\chi\rangle$ in the QFT Hilbert space \mathcal{H}_{QFT} . For example, these states could be obtained by acting on the vacuum with operators smeared in a region spacelike separated from $\mathcal{O}(x)$. Such an object is known as a *sesquilinear form*, defined as a map $\mathfrak{s}(|\chi\rangle, |\phi\rangle)$ from vectors in a domain $\mathcal{D}(\mathfrak{s}) \subset \mathcal{H}_{\text{QFT}}$ to \mathbb{C} that is conjugate linear in the first argument and linear in the second [51]. Sesquilinear forms generalize the notion of an unbounded operator to objects such as $\mathcal{O}(x)$ which make sense inside of certain matrix elements but tend to produce unnormalizable states when acting on a vector. The domain associated to a given local operator $\mathcal{O}(x)$, viewed as a sesquilinear form, is expected to be dense in \mathcal{H}_{QFT} , since, for example, the object $\frac{1}{(1+H)^n} \mathcal{O}(x) \frac{1}{(1+H)^n}$ will be a bounded operator for sufficiently high powers of n , where H is the positive Hamiltonian associated with global future-directed time evolution [52, 53]; the domain of $\mathcal{O}(x)$ is then contained in the domain of H^n . Hence, the future horizon of a cut \mathcal{H}_λ^+ is associated with a set \mathcal{S}_λ of densely defined sesquilinear forms constructed from local operators on \mathcal{H}_λ^+ , even though none of these define true operators with finite fluctuations on \mathcal{H}_{QFT} .

The full algebra \mathcal{C}_λ can be constructed from \mathcal{S}_λ by taking a double commutant. A bounded operator \mathbf{b}' is said to commute with a sesquilinear form \mathfrak{s} if \mathbf{b}' preserves the domain $\mathcal{D}(\mathfrak{s})$ and $\mathfrak{s}((\mathbf{b}')^\dagger |\chi\rangle, |\phi\rangle) = \mathfrak{s}(|\chi\rangle, \mathbf{b}' |\phi\rangle)$ for all $|\chi\rangle, |\phi\rangle \in \mathcal{D}(\mathfrak{s})$. For the forms defined by local operators, this condition simply states that $\langle \chi | \mathbf{b}' \mathcal{O}(x) | \phi \rangle = \langle \chi | \mathcal{O}(x) \mathbf{b}' | \phi \rangle$ in states for which these expressions are finite. The commutant \mathcal{S}'_λ is then an algebra of bounded operators localized in the region R'_λ to the left of the cut, shown in figure 1. Taking a second commutant results in the algebra of operators to the right of the cut, $\mathcal{C}_\lambda = \mathcal{S}''_\lambda$, which is a von Neumann algebra since it is defined as a commutant of a collection of bounded operators \mathcal{S}'_λ . Note that \mathcal{S}'_λ itself need not be a von Neumann algebra, since \mathcal{S}_λ is not an algebra of

bounded operators, and hence is not contained in \mathcal{C}_λ . The full algebra for the region R'_λ can be defined as $\mathcal{A}(R'_\lambda) = \mathcal{C}'_\lambda = \mathcal{S}'''_\lambda$. The geometric action of P_ρ on local operators on the horizon implies that $e^{isP_\rho}\mathcal{S}_\lambda e^{-isP_\rho} = \mathcal{S}_{\lambda+s\rho}$. It is also clear then that e^{isP_ρ} will conjugate elements of \mathcal{S}'_λ into operators that commute with $\mathcal{S}_{\lambda+s\rho}$, and hence $e^{isP_\rho}\mathcal{S}'_\lambda e^{-isP_\rho} = \mathcal{S}'_{\lambda+s\rho}$. The same argument applies to $\mathcal{C}_\lambda = \mathcal{S}''_\lambda$, and hence we conclude that $e^{isP_\rho}\mathcal{C}_\lambda e^{-isP_\rho} = \mathcal{C}_{\lambda+s\rho}$, showing that the geometric action of P_ρ extends to the horizon cut algebras as a whole.

We now apply this result on the geometric action of the null translations to the algebra $\mathcal{B} = \mathcal{C}_0$ associated with the exterior of the horizon bifurcation surface. Fixing a nonnegative profile for the smearing function $\lambda(y^A)$, we define $U_\lambda(s) = e^{isP_\lambda}$ to be the unitary implementing a finite null translation. By the above discussion, we find that for $s \geq 0$,

$$U_\lambda(s)\mathcal{B}U_\lambda(s)^\dagger = \mathcal{C}_{s\lambda} \subseteq \mathcal{B}. \quad (3.3)$$

This is one necessary condition in order to conclude that $U_\lambda(s)$ generates a *half-sided translation* for \mathcal{B} , a concept introduced by Borchers [41, 42]. The other conditions are that $U_\lambda(s)$ have a positive generator, and that it preserve a vacuum state $|\Omega\rangle$ that is cyclic and separating for \mathcal{B} . Positivity of the generator P_λ is implied by the average null energy condition (ANEC) [54–56], since P_λ is an integral of the average null energy of each horizon generator, weighted by a positive function λ . Rigorous proofs of the ANEC exist for interacting field theories in Minkowski space [57–59]. In curved spacetimes, it is generally expected to hold for achronal null geodesics for backgrounds satisfying the semiclassical Einstein equation [60–64]. In the present context, since the horizon generators are achronal, it is reasonable to assume that the ANEC holds for them, implying that the P_λ are positive operators. The vacuum state $|\Omega\rangle$ is then simply a ground state for the smeared average null energy operator, satisfying $P_\lambda|\Omega\rangle = 0$. Note that since average null energy operators on different horizon generators commute, any other choice of smearing ρ yields an operator satisfying $[P_\rho, P_\lambda] = 0$, and hence $|\Omega\rangle$ can be defined as a simultaneous ground state of all smeared horizon average null energy operators [38].

Together, the conditions (i) $P_\lambda \geq 0$, (ii) $U_\lambda(s)\mathcal{B}U_\lambda(s)^\dagger \subseteq \mathcal{B}$ for $s \geq 0$, and (iii) $P_\lambda|\Omega\rangle = 0$ with $|\Omega\rangle$ cyclic and separating for \mathcal{B} imply that $U_\lambda(s)$ defines a half-sided translation for \mathcal{B} . Borchers’s theorem for half-sided translations then states that the modular flow Δ_Ω^{-is} for the vector $|\Omega\rangle$ combines with translations to form a unitary representation of the one-dimensional affine group, with the relation

$$\Delta_\Omega^{-is}U_\lambda(t)\Delta_\Omega^{is} = U_\lambda(e^{2\pi st}). \quad (3.4)$$

This relation holds for any choice of smearing λ since $|\Omega\rangle$ is the simultaneous ground state of all smeared translation generators P_λ .

Specializing for the moment to the constant smearing $\lambda = 1$, we can demonstrate that modular flow acts geometrically on local operators on the horizon. We first express a given horizon operator $\mathcal{O}(v)$ (leaving the y^A dependence implicit) as the translation of an operator at $v = 0$, $\mathcal{O}(v) = U_1(v)\mathcal{O}_1(0)U_1(v)^\dagger$. Because $\mathcal{O}_1(0)$ lies on the bifurcation surface, it should commute with operators in both \mathcal{B} and \mathcal{B}' , and hence modular flow should map it to another

local operator at $v = 0$, $\Delta_{\Omega}^{-is} \mathcal{O}_1(0) \Delta_{\Omega}^{is} = \mathcal{O}_2(0)$. Then we find that

$$\begin{aligned} \Delta_{\Omega}^{-is} \mathcal{O}(v) \Delta_{\Omega}^{is} &= \Delta_{\Omega}^{-is} U_1(v) \Delta_{\Omega}^{is} \mathcal{O}_2(0) \Delta_{\Omega}^{-is} U_1(v)^{\dagger} \Delta_{\Omega}^{is} = U_1(e^{2\pi s} v) \mathcal{O}_2(0) U_1(e^{2\pi s} v) \\ &= \mathcal{O}_3(e^{2\pi s} v), \end{aligned} \quad (3.5)$$

for some local operator \mathcal{O}_3 . This is precisely the form of a geometric flow along the vector field $2\pi v \left(\frac{\partial}{\partial v}\right)^a = \frac{2\pi}{\kappa} \xi^a$, whose generator can be expressed as an integral over the horizon of the stress tensor weighted by ξ^a . From this argument, we conclude that the vacuum modular Hamiltonian $h_{\Omega}^{\mathcal{B}}$ for \mathcal{B} is given by

$$h_{\Omega}^{\mathcal{B}} = -\log \Delta_{\Omega} = -\frac{2\pi}{\kappa} \int_{\mathcal{H}} T^a_b \xi^b \epsilon_{a\dots} = 2\pi \int_{-\infty}^{\infty} dv \int dy^A \sqrt{q} T_{vv} v. \quad (3.6)$$

Because we did not specify the form of the operator \mathcal{O}_3 in equation (3.5), in principle we only determine $h_{\Omega}^{\mathcal{B}}$ up to transformations that fix its location such as local Lorentz or internal symmetry transformations. However, the fact that equation (3.4) implies the commutation relation

$$[h_{\Omega}^{\mathcal{B}}, P_1] = -i2\pi P_1 \quad (3.7)$$

fixes the form of $h_{\Omega}^{\mathcal{B}}$ in terms of the stress tensor, up to terms that commute with P_1 . Such terms are related to the algebra at infinity, and are discussed in section 4. Equation (3.6) is the standard expression for the modular Hamiltonian of the Hartle-Hawking state (in situations where it exists) in terms of the generator of the Killing symmetry. However, for more general Killing horizons such as Kerr black holes and Schwarzschild-de Sitter, equation (3.6) remains valid even though there is no global Hartle-Hawking state.

Finally, we obtain the form of the modular Hamiltonian for an arbitrary horizon cut \mathcal{C}_{λ} . First we note that $\mathcal{C}_{\lambda} = U_{\lambda}(1) \mathcal{B} U_{\lambda}(1)^{\dagger}$, which, along with the relation (3.4) demonstrates that for $s \geq 0$,

$$\Delta_{\Omega, \mathcal{B}}^{-is} \mathcal{C}_{\lambda} \Delta_{\Omega, \mathcal{B}}^{is} = U_{\lambda}(e^{2\pi s} - 1) \mathcal{C}_{\lambda} U_{\lambda}(e^{2\pi s} - 1)^{\dagger} \subseteq \mathcal{C}_{\lambda}. \quad (3.8)$$

Assuming $|\Omega\rangle$ is cyclic for \mathcal{C}_{λ} , this property implies that $\mathcal{C}_{\lambda} \subset \mathcal{B}$ is a *half-sided modular inclusion*, as defined by Wiesbrock [43]. Since $U_{\lambda}(1)|\Omega\rangle = |\Omega\rangle$, the modular operator for \mathcal{C}_{λ} is simply obtained from $\Delta_{\mathcal{B}}$ by conjugation: $\Delta_{\mathcal{C}_{\lambda}} = U_{\lambda}(1) \Delta_{\mathcal{B}} U_{\lambda}(-1)$ (leaving the dependence on $|\Omega\rangle$ implicit). Again applying (3.4), this results in

$$\Delta_{\mathcal{B}}^{-is} \Delta_{\mathcal{C}_{\lambda}}^{is} = U_{\lambda}(e^{2\pi s} - 1), \quad (3.9)$$

which leads to the relation for the modular Hamiltonians

$$h_{\mathcal{B}} - h_{\mathcal{C}_{\lambda}} = 2\pi P_{\lambda}. \quad (3.10)$$

Since both $h_{\mathcal{B}}$ and P_{λ} have been expressed in terms of integrals of the stress tensor in equations (3.6) and (3.2), we see that $h_{\mathcal{C}_{\lambda}}$ may also be expressed as a stress tensor integral over the horizon according to

$$h_{\mathcal{C}_{\lambda}} = -\frac{2\pi}{\kappa} \int_{\mathcal{H}} T^a_b \zeta_{\lambda}^b \epsilon_{a\dots} = 2\pi \int_{-\infty}^{\infty} dv \int dy^A \sqrt{q} T_{vv} \cdot (v - \lambda(y^A)), \quad (3.11)$$

where we have defined $\zeta_\lambda^a = \xi^a - \kappa\lambda^a$. This is Wall's expression for the horizon-cut modular Hamiltonian [38].

The argument described above for deriving this equation for interacting theories using half-sided translations and modular inclusions was first given for Rindler horizons by Casini, Teste, and Torroba [39]. The idea to define the exterior algebras \mathcal{C}_λ as a double commutant of a set \mathcal{S}_λ of sesquilinear forms consisting of local operators on the horizon appears in a lecture by Witten [40], and immediately allows one to see that the horizon average null energy operators generate half-sided translations. The idea to relate the geometric action of modular Hamiltonians on Killing horizons to half-sided modular inclusions appears in work by Summers and Verch [65], and is closely related to older arguments by Sewell for the Hawking temperature of Killing horizons in the Wightman formulation of axiomatic quantum field theory [66]. These latter two works attempt to make sense of a net of operators strictly localized to the Killing horizon, which appears to only be valid in free field theory. In interacting theories, there are arguments that no bounded operators can be strictly localized to a finite affine parameter interval of a codimension-1 null surface [50], and in this case one must use the more sophisticated arguments developed by Casini, Teste, and Torroba, and Witten to conclude that horizon cuts define half-sided modular inclusions.

3.2 Geometric horizon identities

Connecting the results in section 3.1 on the form of the horizon cut modular Hamiltonian to the generalized entropy requires a discussion of how the quantum fields couple to gravity in the semiclassical limit $G_N \rightarrow 0$. This limit suppresses backreaction, allowing for a description in terms of quantum fields on the fixed black hole geometry. However, we must also account for the gravitational constraints when constructing a theory that consistently embeds into the interacting theory away from $G_N = 0$. These constraints lead to relations between the stress-energy on the horizon and the values of asymptotic gravitational charges evaluated at the future and past boundaries of the horizon $c_{\pm\infty}$ located at $v \rightarrow \pm\infty$. In addition, semilocal constraints associated with one side of a horizon cut lead to formal relations between the area of the cut and the asymptotic charges, which are necessary for establishing the equivalence of the entropy of the gravitational algebra and the horizon generalized entropy. In this section, we review how these constraints arise and present a novel derivation of the quasilocal identity relating the horizon area and one-sided boost energy. This identity is originally due to Wall [38]; see also [32] for a related recent treatment.

The structure of the constraints and asymptotic charges can be derived from the classical theory at finite G_N ; from these one can obtain perturbative expressions in the $\sqrt{G_N}$ expansion that then apply in the semiclassical quantum theory. Due to the choice of AdS asymptotics, the black hole horizon \mathcal{H} serves as a null Cauchy surface for the spacetime. The canonical formulation of general relativity on null surfaces has been the subject of several recent investigations [67–73] and can immediately be applied in the present context to derive the form of the Hamiltonians and constraints for the geometric flows considered in section 3.1. We will simply quote some of the main results needed from the canonical analysis in this section; additional details can be found in appendix C and in the cited references.

We denote by ϕ the dynamical fields in the theory, which consist of the metric g_{ab} and any matter fields ψ . On the null Cauchy surface, one can define a symplectic potential current $\mathcal{E}[\phi; \delta\phi]$ which, assuming minimal coupling between matter and gravity, is expressed as a sum of a gravitational term and a matter term,

$$\mathcal{E} = \mathcal{E}^g + \mathcal{E}^\psi. \quad (3.12)$$

The gravitational piece, after imposing the gauge conditions given in (C.7), takes the form

$$\mathcal{E}^g[g_{ab}; \delta g_{ab}] = \frac{1}{16\pi G_N} \eta \left(\sigma^{ab} \delta q_{ab} - \frac{d-3}{d-2} \Theta q^{ab} \delta q_{ab} \right), \quad (3.13)$$

where η is the volume form on \mathcal{H} , q_{ab} is the degenerate induced metric, $\Theta = \nabla_a l^a$ is the expansion, and $\sigma_{ab} = \frac{1}{2} \mathcal{L}_l q_{ab} - \frac{1}{d-2} \Theta q_{ab}$ is the shear, which vanishes in the background but is nonvanishing once we consider perturbative fluctuations of the geometry. The matter symplectic potential \mathcal{E}^ψ is theory dependent, but for definiteness one can have in mind a scalar field φ , for which

$$\mathcal{E}^\varphi = \eta(l^a \nabla_a \varphi) \delta\varphi. \quad (3.14)$$

The symplectic form is obtained from \mathcal{E} by integrating its antisymmetric variation over the Cauchy surface \mathcal{H} ,

$$\Omega[\delta_1\phi, \delta_2\phi] = \int_{\mathcal{H}} \left(\delta_1 \mathcal{E}[\delta_2\phi] - \delta_2 \mathcal{E}[\delta_1\phi] \right). \quad (3.15)$$

For a vector field ζ^a tangent to \mathcal{H} , the associated Hamiltonian H_ζ can be defined as (see equation (C.18))

$$H_\zeta = \int_{\mathcal{H}} \mathcal{E}[\phi; \mathcal{L}_\zeta \phi] = \frac{1}{16\pi G_N} \int_{\mathcal{H}} \eta \left(\sigma^{ab} \mathcal{L}_\zeta q_{ab} - \frac{d-3}{d-2} \Theta q^{ab} \mathcal{L}_\zeta q_{ab} \right) - \int_{\mathcal{H}} (T_{(\psi)})^a_b \zeta^b \epsilon_{a\dots}. \quad (3.16)$$

which displays the contributions from both the gravitational field and the matter stress tensor $(T_{(\psi)})^a_b$. This object generates the flow of the diffeomorphism ζ^a on the gravitational phase space since it satisfies Hamilton's equation

$$\delta H_\zeta = \Omega[\delta\phi, \mathcal{L}_\zeta \phi], \quad (3.17)$$

which holds off-shell as long as ζ^a preserves the asymptotic boundary conditions imposed at $v \rightarrow \pm\infty$.

Due to diffeomorphism invariance, the stress-energy density $\mathcal{E}[\phi; \mathcal{L}_\zeta \phi]$ can be written as a total derivative dM_ζ , with M_ζ defined in equation (C.13), up to a term involving the constraint equation C_ζ ,

$$\mathcal{E}[\phi; \mathcal{L}_\zeta \phi] = dM_\zeta + C_\zeta, \quad (3.18)$$

Integrating this relation over \mathcal{H} yields a formula expressing the integrated constraint $\mathfrak{C}[\zeta] = \int_{\mathcal{H}} C_\zeta$ in terms of the generator of the diffeomorphism H_ζ and the asymptotic charges \mathcal{Q}_ζ^\pm ,

$$\mathfrak{C}[\zeta] = H_\zeta - \mathcal{Q}_\zeta^+ + \mathcal{Q}_\zeta^-, \quad (3.19)$$

where

$$\mathcal{Q}_\zeta^\pm = \int_{c_{\pm\infty}} M_\zeta. \quad (3.20)$$

We now specialize to the vector fields that are relevant to the discussion of half-sided modular inclusions considered in section 3.1, all of which are parallel to the null generator l^a on \mathcal{H} . They have the property of being *quasi-Killing vectors*,¹ meaning that they satisfy Killing's equation $\mathcal{L}_\zeta g_{ab} = 0$ only on \mathcal{H} , i.e. they are symmetries of the horizon which may not extend to symmetries of the full spacetime. A generic quasi-Killing vector parallel to l^a on \mathcal{H} can be written in terms of a function $\tau(v, y^A)$ as (see appendix B.3)

$$\zeta^a = \tau l^a + u \nabla^a \tau. \quad (3.21)$$

The choice $\tau = \kappa v$ yields the global Killing vector $\xi^a = \kappa v l^a - \kappa u n^a$, while $\tau = \lambda(y^A)$ results in the smeared null translation that appears in the average null energy operator (3.2).

The boost vector ζ_λ^a occurring in the modular Hamiltonian for a horizon cut is also quasi-Killing, arising from the smearing $\tau(v, y^A) = \kappa(v - \lambda(y^A))$ which leads to the expression

$$\zeta_\lambda^a = \kappa \left[(v - \lambda) l^a - u \lambda n^a - u \nabla^a \lambda \right]. \quad (3.22)$$

This vector vanishes at the horizon cut c_λ at which $v = \lambda$, and its covariant derivative there satisfies

$$\nabla_a (\zeta_\lambda)_b \stackrel{c_\lambda}{=} \kappa n_{ab}^\lambda, \quad (3.23)$$

where

$$n_{ab}^\lambda = 2l_{[a}(n_{b]} + \nabla_{b]}\lambda) \quad (3.24)$$

is the unit binormal to c_λ . To see this, note that the cut is defined by the relations $u = 0$ and $v - \lambda(y^A) = 0$, so (3.24) is proportional to the unit binormal since $n_a = -\nabla_a v$. To verify it is unit-normalized, we compute

$$n_{ab}^\lambda n_\lambda^{ab} = 2(l \cdot l) |n + d\lambda|^2 - 2 \left(l \cdot (n + d\lambda) \right)^2 \stackrel{\mathcal{H}}{=} -2. \quad (3.25)$$

It is notable that κ in equation (3.23) is constant, since it means that the modular flow for the horizon cut corresponds to a boost with constant surface gravity. This is a nontrivial verification of the geometric modular flow conjecture from [33], which states that a modular flow should look like a constant-surface-gravity boost near the entangling surface. Constancy of κ can be viewed as a *zeroth law of modular flow*, since it is closely related to the KMS condition for the modular flow on the horizon cut algebra. It is notable that κ is constant for a generic cut of the horizon, and does not depend on quantities such as the extrinsic curvature of the codimension-2 cut, which is nonzero in the u direction away from the bifurcation surface.

¹This terminology was introduced by Bob Wald in a talk related to his work with Hollands and Zhang on dynamical black hole entropy [74].

For the translation quasi-Killing vector $\lambda l^a + u \nabla^a \lambda$, the gravitational contribution to the energy density obtained from (3.13) is

$$\mathcal{E}^g[\mathcal{L}_{\lambda l} g_{ab}] = \frac{1}{8\pi G_N} \eta \lambda \left(\sigma^{ab} \sigma_{ab} - \frac{d-3}{d-2} \Theta^2 \right), \quad (3.26)$$

which motivates the definition of the gravitational null energy density as²

$$t_{vv}^{(g)} = \frac{1}{8\pi G_N} \left(\sigma^{ab} \sigma_{ab} - \frac{d-3}{d-2} \Theta^2 \right). \quad (3.27)$$

Note that this definition of gravitational null energy makes sense in principle on generic null surfaces, not only for perturbations of a Killing horizon. On the other hand, the potential $M_{\lambda l}$ evaluates to (see equation (C.25))

$$M_{\lambda l} = -\frac{1}{8\pi G_N} \mu \lambda \Theta, \quad (3.28)$$

with μ the induced $(d-2)$ -form that pulls back to the area form on cuts of \mathcal{H} . From this, we can then integrate equation (3.18) between two horizon cuts $v = \tau_1(y^A)$ and $v = \tau_2(y^A)$ to derive a relation between the expansions at the two cuts,

$$\frac{1}{8\pi G_N} \left(\int_{\tau_2} \mu \lambda \Theta - \int_{\tau_1} \mu \lambda \Theta \right) = - \int_{\mathcal{H}_1^2} \eta \lambda \left(t_{vv}^{(g)} + T_{ab}^{(\psi)} l^a l^b \right) + \mathfrak{C}_1^2[\lambda l] \quad (3.29)$$

where $\mathfrak{C}_1^2[\lambda l]$ is the integral of the constraint $C_{\lambda l}$ over \mathcal{H}_1^2 .

Equation (3.29) can be viewed as an integrated version of the Raychaudhuri equation,³ which highlights the interpretation of the terms in the equation as contributions from the matter and gravitational null energy. As the second cut τ_2 approaches future infinity on the horizon, the expansion must go to zero as a boundary condition specifying that \mathcal{H} is an event horizon [38, 75]. This limit results in an expression for the expansion on a cut c_τ in terms of the semi-infinite null energy (once the constraints $C_{\lambda l} = 0$ are imposed)

$$\frac{1}{8\pi G_N} \int_{c_\tau} \mu \lambda \Theta = P_\tau^+ = \int_{\mathcal{H}_\tau^+} \eta \lambda \left(t_{vv}^{(g)} + T_{ab}^{(\psi)} l^a l^b \right). \quad (3.30)$$

Another useful relation comes from examining the charges for the boost vector ζ_λ^a . From equation (C.25), the potential M_{ζ_λ} evaluates to

$$M_{\zeta_\lambda} = \frac{\kappa}{8\pi G_N} \mu \left(1 - (v - \lambda) \Theta \right), \quad (3.31)$$

²Closely related expressions for the gravitational energy associated with generic null surfaces were recently explored in [70].

³An alternative way to obtain this formula is to write $d(\Theta \mu) = (\dot{\Theta} + \Theta^2) \eta$ with $\dot{\Theta} = l^a \nabla_a \Theta$. One then uses the Raychaudhuri equation to obtain $\dot{\Theta} + \Theta^2 = \frac{d-3}{d-2} \Theta^2 - \sigma^{ab} \sigma_{ab} - T_{ab}^{(\psi)} l^a l^b + C_l$, with $C_l = 0$ a component of the Einstein equation.

and after integrating the flux-balance equation (3.18) between the cut c_λ at which ζ_λ^a vanishes and infinity, we obtain

$$\frac{\kappa}{8\pi G_N}(A_\infty - A_\lambda) = H_{\zeta_\lambda}^+ = \kappa \int_{\mathcal{H}_\lambda^+} \eta(v - \lambda) \left(t_{vv}^{(g)} + T_{ab}^{(\psi)} l^a l^b \right), \quad (3.32)$$

again using the future boundary condition $\Theta_{+\infty} = 0$ and imposing the constraint $C_{\zeta_\lambda} = 0$. This is the nonlinear version of Wall's relation [38] between the difference of horizon areas ($A_\infty - A_\lambda$) and the one-sided boost energy $H_{\zeta_\lambda}^+$. It also is the identity corresponding to the local Smarr relation discussed by JSS [33], specialized to the case of a cut of a Killing horizon. Note that the boost energy contains both a matter contribution from $T_{ab}^{(\psi)}$ and a gravitational contribution $t_{vv}^{(g)}$ involving the shear and expansion according to (3.27). This latter contribution will become the graviton null energy density when quantizing around the Killing horizon background.

The crossed product algebra is the result of imposing the global constraint (3.19) for ζ_λ^a , which is the quasi-Killing vector that preserves the horizon cut. We already found in equation (3.32) that the future asymptotic charge is given by the late time area of the event horizon,

$$\mathcal{Q}_{\zeta_\lambda}^+ = \frac{\kappa A_\infty}{8\pi G_N}. \quad (3.33)$$

The asymptotic gravitational charge at past infinity is given by

$$\mathcal{Q}_{\zeta_\lambda}^- = \lim_{\tau \rightarrow -\infty} \frac{\kappa}{8\pi G_N} \left(A_\tau - \int_{c_\tau} \mu(v - \lambda) \Theta \right). \quad (3.34)$$

Unlike the future horizon boundary, we cannot impose $\Theta \xrightarrow{\tau \rightarrow -\infty} 0$ since the asymptotic value of Θ is determined via (3.30) by the total average null energy on the horizon. This raises the possibility that $\mathcal{Q}_{\zeta_\lambda}^-$ could diverge in the limit. However, at least when working perturbatively about a Killing horizon, the expansion Θ remains small up to late time scales of order $v \sim \mathcal{O}(\frac{1}{G_N})$. In this perturbative regime, the terms growing linearly with v in the area A_τ cancel against the linear term in v in the integral involving the expansion in (3.34), resulting in a finite expression if we take $\tau \rightarrow -\infty$ and $G_N \rightarrow 0$ with $|\tau| \ll \frac{1}{G_N}$.

Since we will mostly be concerned with the algebra to the future of a cut, the precise expression for the left asymptotic charge will not enter into the entropy calculations. It suffices to know that such a left asymptotic charge can be defined, after which the constraint (3.19) can be written

$$\mathfrak{C}[\zeta_\lambda] = H_{\zeta_\lambda} - \frac{\kappa}{8\pi G_N} A_\infty + \mathcal{Q}_{\zeta_\lambda}^- \quad (3.35)$$

It is, however, interesting to note that the expression $A_\tau - \int_{c_\tau} \mu v \Theta$ has recently appeared in the dynamical entropy proposal of Hollands, Wald, and Zhang [74], who showed that this combination can be interpreted as the area of an apparent horizon inside the black hole event horizon (see also [76]). As $\tau \rightarrow -\infty$, this apparent horizon approaches the white hole event horizon for the left asymptotic region. In general, when there is nonzero null energy flux through \mathcal{H} , the Einstein-Rosen bridge connecting the left and right regions becomes longer,

causing the left white hole event horizon to differ from the right black hole event horizon. Hence, we can use the above result to rewrite the left asymptotic charge in terms of the left white hole event horizon asymptotic area $A_{-\infty}^{\text{wh}}$ and the asymptotic expansion $\Theta_{-\infty}$ as

$$\mathcal{Q}_{\zeta_\lambda}^- = \frac{\kappa}{8\pi G_N} \left(A_{-\infty}^{\text{wh}} + \int_{c_{-\infty}} \mu \lambda \Theta_{-\infty} \right). \quad (3.36)$$

We now turn to the description of the various relations above in the small G_N limit, which allows for perturbative quantization of the matter fields and gravitons about the Killing horizon background. The metric can be expanded about its background value g_{ab}^0 as

$$g_{ab} = g_{ab}^0 + \varkappa h_{ab} \quad (3.37)$$

where $\varkappa = \sqrt{32\pi G_N}$; this choice ensures that h_{ab} has a canonical kinetic term in the quadratic Lagrangian. Similarly, the degenerate metric on \mathcal{H} is expanded as

$$q_{ab} = q_{ab}^0 + \varkappa \rho_{ab} + \frac{\varkappa}{d-2} \rho q_{ab}^0, \quad q_0^{ab} \rho_{ab} = 0, \quad (3.38)$$

and because the shear and expansion vanish in the background, to leading order in \varkappa they are given by

$$\sigma_{ab} = \frac{\varkappa}{2} \mathcal{L}_l \rho_{ab} \equiv \frac{\varkappa}{2} \dot{\rho}_{ab} \quad (3.39)$$

$$\Theta = \frac{\varkappa}{2} \mathcal{L}_l \rho \equiv \frac{\varkappa}{2} \dot{\rho}. \quad (3.40)$$

This then implies that the graviton null energy density, given by (3.27), is finite as $G_N \rightarrow 0$. Additionally, the expansion is constrained to satisfy (3.30) for all choices of smearing λ . Since the right hand side is $\mathcal{O}(\varkappa^0)$, it must be that the expansion can be nonzero only at $\mathcal{O}(\varkappa^2)$, implying that in the linear theory, $\dot{\rho} = 0$, and the graviton null energy density simplifies to

$$t_{vv}^{(g)} = \dot{\rho}^{ab} \dot{\rho}_{ab}. \quad (3.41)$$

Note that this energy density differs from the Hollands-Wald canonical energy considered in [75] by a total derivative which integrates to zero over the horizon.

The vanishing of the expansion at $\mathcal{O}(\varkappa)$ implies the horizon area is constant at linear order. Its value is then set by $A_\infty^{(1)}$, the $\mathcal{O}(\varkappa)$ term in the expansion of the late time horizon area, which we choose to set to zero as a boundary condition; fluctuations in the area at this order have the interpretation of describing different background spacetimes.⁴ On the other hand, the $\mathcal{O}(\varkappa^2)$ term in the late time area, $A_\infty^{(2)}$, does interact nontrivially with the matter fields, and hence we include it as a global gravitational charge. Together with the second

⁴Fluctuations at this order are expected in the canonical ensemble for the black hole, but allowing for such large fluctuations leads to complications in the construction of the gravitational algebras, and require working in a formal power series in \varkappa , as in [29]. Restricting fluctuations in the area to be $\mathcal{O}(\varkappa^2)$ corresponds to a microcanonical ensemble, as discussed in [30].

order contribution to the left gravitational charge $(\mathcal{Q}_{\zeta\lambda}^-)^{(2)}$, the global constraint (3.35) for the linearized theory becomes

$$\mathfrak{C}^{(2)}[\zeta\lambda] = \kappa \int_{\mathcal{H}} \eta \left(t_{vv}^{(g)} + T_{ab}^{(\psi)} l^a l^b \right) (v - \lambda) - \frac{\kappa}{2\pi} \frac{A_\infty^{(2)}}{4G_N} + (\mathcal{Q}_{\zeta\lambda}^-)^{(2)} \quad (3.42)$$

$$= \frac{\kappa}{2\pi} \left(h_{\mathcal{C}_\lambda} - \frac{A_\infty^{(2)}}{4G_N} \right) + (\mathcal{Q}_{\zeta\lambda}^-)^{(2)} \quad (3.43)$$

where in the second line we have applied the formula (3.11) to write the constraint in terms of the vacuum modular Hamiltonian of the horizon cut, which, including the graviton contribution, takes the form

$$h_{\mathcal{C}_\lambda} = 2\pi \int_{\mathcal{H}} \eta \left(t_{vv}^{(g)} + T_{ab}^{(\psi)} l^a l^b \right) (v - \lambda) = 2\pi \int dy^A \int_{-\infty}^{\infty} dv \sqrt{q} \left(t_{vv}^{(g)} + T_{vv}^{(\psi)} \right) (v - \lambda(y^A)). \quad (3.44)$$

3.3 Generalized second law for semiclassical states

We now have all the components needed to construct gravitational algebras associated with horizon cuts and use them to prove a generalized second law. The starting point is the algebra \mathcal{C}_λ associated with the region R_λ spacelike separated from the horizon cut defined by the function $\lambda(y^A)$. This algebra describes both matter and graviton degrees of freedom. The matter will generically involve an interacting quantum field theory; in particular, we will not make use of any free field relations in the matter sector. In this case, the classical description of the matter in terms of a Lagrangian and symplectic form may be a poor approximation, but the general arguments leading to the form of the matter modular Hamiltonian, equation (3.11) only rely on the existence of a stress tensor as the generator of diffeomorphisms and positivity of the average null energy. Since all matter couples universally to gravity via its stress tensor, the matter contribution to the gravitational constraint in equation (3.43) remains valid even for interacting theories.

On the other hand, the gravitons are quantized as a decoupled theory of free, massless, spin-2 particles, obtained by promoting the metric perturbation h_{ab} in (3.37) to a quantum operator and performing canonical quantization with appropriate gauge-fixing conditions for linearized diffeomorphisms. This is the correct description in the $G_N \rightarrow 0$ limit, which suppresses self-interactions between gravitons as well as direct matter-graviton couplings. The ability to treat gravitons as free fields is important at a technical level, since gravitational interactions are nonrenormalizable, causing the theory to become strongly coupled at the Planck length. A description in this regime would require a fully nonperturbative theory of quantum gravity, and it is widely expected that the language of quantum field theory is not applicable at such energy scales. However, assuming the low energy gravitational theory admits an effective field theory description [77, 78], the free graviton approximation should give an accurate description at leading order in G_N . In this case, the classical phase space analysis described in section 3.2 is directly relevant, and the graviton contribution to the constraint (3.42) is given by $t_{vv}^{(g)}$ defined in (3.41), which is quadratic in the graviton field.

The gravitational algebra additionally includes degrees of freedom associated with the asymptotic charges \mathcal{Q}_λ^\pm . As discussed in section 3.2, these charges are constructed from the $\mathcal{O}(\varkappa^2)$ contribution to the metric perturbation, and hence are not present in the linearized theory involving only gravitons [32]. In the interacting theory away from $G_N = 0$, these asymptotic charges are present and are related to the matter and graviton stress energy, which is nontrivial even in the linearized theory. Hence, in order to obtain a theory at $G_N = 0$ that consistently lifts to the interacting theory, we must include the asymptotic charges explicitly and couple them to the matter fields via the constraint (3.42).⁵

The $\mathcal{O}(\varkappa^2)$ contribution to the asymptotic horizon area comprises the future asymptotic charge \mathcal{Q}_λ^+ that must be included. We represent this operator on an independent Hilbert space $\mathcal{H}_{\text{asy}} = L^2(\mathbb{R})$ as the position operator,

$$\hat{q} = -\frac{A_\infty^{(2)}}{4G_N}. \quad (3.45)$$

This choice reflects two semiclassical assumptions about the spectrum of the asymptotic area operator, namely that it is continuous and unbounded above and below. The continuous spectrum of this operator implies that we are working in a description that does not resolve individual microstates of the black hole, which ultimately manifests in the constant shift ambiguity in the crossed product entropy. The unboundedness of the spectrum corresponds to the fact that we are considering perturbations of the area around a fixed background, and because the perturbation only describes changes in the geometry at order \varkappa^2 , it can take on large positive or negative values in the $\varkappa \rightarrow 0$ limit without producing a significant change in the background value of the area.

The algebra of gravitationally dressed operators is denoted $\widehat{\mathcal{C}}_\lambda$, and is constructed by imposing the constraint (3.43) to obtain a gauge-invariant Hilbert space and operators, following the procedure described in [31, 33]. This results in the crossed product algebra discussed in section 2. It is represented on the Hilbert space $\widehat{\mathcal{H}} = \mathcal{H}_{\text{QFT}} \otimes \mathcal{H}_{\text{asy}}$, and is given by

$$\widehat{\mathcal{C}}_\lambda = \langle e^{i\hat{p}h_\lambda} \mathbf{a} e^{-i\hat{p}h_\lambda}, \hat{q} \rangle, \quad \mathbf{a} \in \mathcal{C}_\lambda, \quad (3.46)$$

where $\hat{p} = -i\frac{d}{dq}$, and $h_\lambda \equiv h_{\mathcal{C}_\lambda}$ is the modular Hamiltonian for the horizon cut in the average null energy vacuum $|\Omega\rangle$, given in terms of the matter and graviton stress tensors by equation (3.44).

To examine the generalized second law, we would like to compute entropies of states of the form $|\widehat{\Phi}\rangle = |\Phi\rangle \otimes |f\rangle$, where $|\Phi\rangle$ is a state in \mathcal{H}_{QFT} and f is a wavefunction on $L^2(\mathbb{R})$ that is slowly varying in the position basis, and hence whose derivatives are suppressed by

⁵In a complete description, one should also add charges associated with other asymptotic symmetries such as rotations [29–31]. Including these charges results in additional crossed products with respect to the symmetries that they generate. They do not change the overall type of the algebra, so they will not affect the results involving finiteness of the entropy and should not affect the arguments for the semiclassical generalized second law, so we do not consider them in detail in this work. However, rotational asymptotic charges are important when obtaining the correct matching conditions for rotating black hole backgrounds [32], so it would be worth considering them in more detail in the future.

a small parameter ε . Using the formula for the density matrix for this state given in (2.12), we can apply the semiclassical expression for the entropy (2.14) to arrive at

$$S_\lambda(\rho_{\widehat{\Phi}}) = -S_{\text{rel}}^{\mathcal{C}_\lambda}(\Phi||\Omega) + \left\langle \frac{A_\infty^{(2)}}{4G_N} \right\rangle_{\widehat{\Phi}} + S_f^{A_\infty} + \mathcal{O}(\varepsilon^2). \quad (3.47)$$

The $\mathcal{O}(\varepsilon^2)$ corrections to this semiclassical entropy formula are discussed in section 5.

We can convert this expression into something involving the generalized entropy by employing the horizon-cut identity (3.32). The term on the right hand side is proportional to the one-sided vacuum modular Hamiltonian K_λ , defined by restricting the range of integration of v in (3.44) to the future of the horizon cut, $v > \lambda(y^A)$. This defines the formal split of the modular Hamiltonian $h_\lambda = K_\lambda - K'_\lambda$ into an object K_λ affiliated with \mathcal{C}_λ and K'_λ affiliated with the commutant \mathcal{C}'_λ . The one-sided modular Hamiltonians K_λ, K'_λ make sense as densely-defined sesquilinear forms (but not as unbounded operators), similar to the local operators discussed in section 3.1. K_λ therefore has well-defined expectation values for a dense set of states. When coupling to gravity, the constraint (3.32) then implies that the area of the horizon cut, expanded to second order in \varkappa , must also define a sesquilinear form affiliated with the crossed product algebra,

$$\frac{A_\lambda^{(2)}}{4G_N} = \frac{A_\infty^{(2)}}{4G_N} - K_\lambda = -\hat{q} - K_\lambda. \quad (3.48)$$

The relative entropy in (3.47) also has an expression in terms of one-sided quantities. The starting point is the standard expression for the relative entropy between two density matrices $S_{\text{rel}}(\rho_\Phi||\rho_\Omega) = \text{Tr}(\rho_\Phi \log \rho_\Phi - \rho_\Phi \log \rho_\Omega) = -S(\rho_\Phi) + \langle \Phi | -\log \rho_\Omega | \Phi \rangle$. Assuming a regulator to make the one-sided QFT entanglement entropies finite, we can apply this formula to the relative entropy appearing in (3.47). The density matrix for the horizon vacuum state $|\Omega\rangle$ is thermal with respect to the one-sided modular Hamiltonian, and hence is formally expressed as

$$\rho_\Omega = \frac{1}{Z} e^{-K_\lambda}, \quad (3.49)$$

with the normalization set by the requirement that K_λ have zero expectation value in the vacuum, $\langle \Omega | K_\lambda | \Omega \rangle = 0$. The vacuum entanglement entropy is then simply given by $\log Z$:

$$S_\Omega^{\text{QFT}} = \langle \Omega | -\log \rho_\Omega | \Omega \rangle = \log Z. \quad (3.50)$$

From these relations, the relative entropy takes the form

$$-S_{\text{rel}}^{\mathcal{C}_\lambda}(\Phi||\Omega) = S_{\Phi,\lambda}^{\text{QFT}} - \log Z - \langle K_\lambda \rangle_\Phi. \quad (3.51)$$

Applying this to (3.47) and employing the identity (3.48), we arrive at the semiclassical expression for the crossed product entropy

$$S_\lambda(\rho_{\widehat{\Phi}}) = \left\langle \frac{A_\lambda^{(2)}}{4G_N} \right\rangle_{\widehat{\Phi}} + S_{\Phi,\lambda}^{\text{QFT}} + S_f^{A_\infty} - \log Z + \mathcal{O}(\varepsilon^2). \quad (3.52)$$

This agrees with the horizon generalized entropy up to a constant, which can be formally determined by writing the area perturbation in terms of the background value $A_\lambda^{(0)}$ as $A_\lambda^{(2)} = A_\lambda - A_\lambda^{(0)}$. The resulting entropy formula is then

$$S_\lambda(\rho_{\widehat{\Phi}}) = \left\langle \frac{A_\lambda}{4G_N} \right\rangle_{\widehat{\Phi}} + S_{\Phi, \lambda}^{\text{QFT}} + S_f^{A_\infty} - \left(\frac{A_\lambda^{(0)}}{4G_N} + \log Z \right) + \mathcal{O}(\varepsilon^2) \quad (3.53)$$

The term $\log Z$ should be interpreted as quantum field vacuum entanglement entropy, and hence the above constant in parentheses can be viewed as the generalized entropy in the vacuum state $|\Omega\rangle$ (assuming no contribution from the entropy associated with the asymptotic area). Crucially, this constant is independent of the horizon cut λ , since the vacuum $|\Omega\rangle$ is a stationary state. This constant will therefore drop out when considering differences in entropies between horizon cuts.

We now examine the algebra associated with a later cut defined by $\tilde{\lambda}(y^A) \geq \lambda(y^A)$. The quantum field theory algebra $\mathcal{C}_{\tilde{\lambda}}$ acts on the same Hilbert space as before \mathcal{H}_{QFT} , and we once again adjoin the same asymptotic charge $\hat{q} = -\frac{A_\infty^{(2)}}{4G_N}$ acting on $\mathcal{H}_{\text{asy}} = L^2(\mathbb{R})$ when constructing the gravitational algebra. It is significant at this point that the asymptotic charge does not depend on the horizon cut λ , which follows from the form of the constraint (3.43) for a generic boost quasi-Killing vector ζ_λ^α . The gravitational algebra again takes the form of a crossed product,

$$\widehat{\mathcal{C}}_{\tilde{\lambda}} = \langle e^{i\hat{p}h_{\tilde{\lambda}}} \mathbf{a} e^{-i\hat{p}h_{\tilde{\lambda}}}, \hat{q} \rangle, \quad \mathbf{a} \in \mathcal{C}_{\tilde{\lambda}}. \quad (3.54)$$

Even though $\mathcal{C}_{\tilde{\lambda}} \subset \mathcal{C}_\lambda$, the crossed product algebra $\widehat{\mathcal{C}}_{\tilde{\lambda}}$ is not a subalgebra of $\widehat{\mathcal{C}}_\lambda$ in this representation, since the modular Hamiltonians $h_{\tilde{\lambda}}$ and h_λ associated with different cuts are not the same.

An issue that arises when comparing entropies between two type II algebras is that there may not be an obvious choice for the relative normalizations of the traces. This results in a potential ambiguity when comparing entropies between the two algebras. In the present context, we can take advantage of the fact that the algebras share a common operator, the asymptotic charge \hat{q} , to find a canonical relative normalization for the traces. For both algebras, the state $|\widehat{\Omega}\rangle = |\Omega\rangle \otimes |f\rangle$ has a density matrix determined by equation (2.8), and is given by

$$\rho_{\widehat{\Omega}} = |f(\hat{q})|^2 e^{\hat{q}}, \quad (3.55)$$

which is solely a function of \hat{q} . Normalizing the density matrices for both algebras to agree precisely with (3.55) then determines the relative normalization for the traces on the algebras, and in both cases is given by the formula (2.9). This choice ensures that operators in the subalgebra generated by \hat{q} have the same trace in both $\widehat{\mathcal{C}}_{\tilde{\lambda}}$ and $\widehat{\mathcal{C}}_\lambda$.

Considering the same state $|\widehat{\Phi}\rangle = |\Phi\rangle \otimes |f\rangle$ as before, the computation of the entropy for the algebra $\widehat{\mathcal{C}}_{\tilde{\lambda}}$ now proceeds analogously, giving

$$S_{\tilde{\lambda}}(\rho_{\widehat{\Phi}}) = -S_{\text{rel}}^{\mathcal{C}_{\tilde{\lambda}}}(\Phi||\Omega) + \left\langle \frac{A_\infty^{(2)}}{4G_N} \right\rangle_{\widehat{\Phi}} + S_f^{A_\infty} + \mathcal{O}(\varepsilon^2) \quad (3.56)$$

Since $\langle \frac{A_\infty^{(2)}}{4G_N} \rangle_{\widehat{\Phi}}$ and $S_f^{A_\infty}$ depend only on the wavefunction f and the operator \hat{q} , these quantities agree when computed in either algebra. They will therefore drop out when considering the difference in entropies $S_{\widehat{\lambda}} - S_\lambda$. Hence, in the semiclassical limit in which we can ignore the $\mathcal{O}(\varepsilon^2)$ contributions to the entropy, we have that

$$S_{\widehat{\lambda}} - S_\lambda = -S_{\text{rel}}^{\mathcal{C}_{\widehat{\lambda}}}(\Phi||\Omega) + S_{\text{rel}}^{\mathcal{C}_\lambda}(\Phi||\Omega) \geq 0 \quad (3.57)$$

where the last inequality follows from the monotonicity of relative entropy under the inclusion $\mathcal{C}_{\widehat{\lambda}} \subset \mathcal{C}_\lambda$ [79–81]. Hence, we have succeeded in showing that in the semiclassical limit $\varepsilon \ll 1$, the crossed product entropies increase upon restricting to horizon cuts that are further to the future. This is the statement of the second law for the crossed product algebras.

An attractive feature of the second law formulated in terms of crossed product algebras is that it directly relates the increase in entropy to the monotonicity of relative entropy, without appealing to relations involving regulated quantum field theory entanglement entropies. Of course, we argued above that the crossed product entropy agrees semiclassically with the generalized entropy by invoking the local gravitational constraint (3.32), so this monotonicity result can instead be stated directly in terms of generalized entropy. The formula (3.53) is applicable for the $\widehat{\mathcal{C}}_\lambda$ algebra as well, and since the constant in parentheses is independent of the horizon cut, it drops out of the entropy difference. Hence, equation (3.57) also implies the standard statement of the generalized second law in terms of generalized entropies,

$$\left\langle \frac{A_{\widehat{\lambda}}}{4G_N} \right\rangle_{\widehat{\Phi}} + S_{\widehat{\lambda}}^{\text{out}} - \left\langle \frac{A_\lambda}{4G_N} \right\rangle_{\widehat{\Phi}} - S_\lambda^{\text{out}} \geq 0, \quad (3.58)$$

where $S_{\widehat{\lambda}}^{\text{out}} := S_{\widehat{\Phi}, \widehat{\lambda}}^{\text{QFT}} + S_f^{A_\infty}$, and similarly for S_λ^{out} . This shows that the crossed product entropies reproduce Wall's proof of the generalized second law, but they have the added benefit of giving an algebraic interpretation to the generalized entropies appearing in it.

One important aspect of this derivation is that it is necessary to work in the representation where the crossed product algebras take the form (3.46) and (3.54). These representations have the property that the algebraic intersection consists of operators in the asymptotic algebra, i.e. $\widehat{\mathcal{C}}_\lambda \wedge \widehat{\mathcal{C}}_{\widehat{\lambda}} = \langle \hat{q} \rangle$. In some recent investigations of gravitational algebras [29–31], a unitarily equivalent representation was employed where the algebras take the form

$$\widetilde{\mathcal{C}}_\lambda = e^{-i\hat{p}h_\lambda} \widehat{\mathcal{C}}_\lambda e^{i\hat{p}h_\lambda} = \langle \mathbf{a}, \hat{q} - h_\lambda \rangle, \quad \mathbf{a} \in \mathcal{C}_\lambda, \quad (3.59)$$

$$\widetilde{\mathcal{C}}_{\widehat{\lambda}} = e^{-i\hat{p}h_{\widehat{\lambda}}} \widehat{\mathcal{C}}_{\widehat{\lambda}} e^{i\hat{p}h_{\widehat{\lambda}}} = \langle \mathbf{b}, \hat{q} - h_{\widehat{\lambda}} \rangle, \quad \mathbf{b} \in \mathcal{C}_{\widehat{\lambda}}. \quad (3.60)$$

The procedure described above of considering the same global state $|\widehat{\Phi}\rangle = |\Phi\rangle \otimes |f\rangle$ for the two algebras and then comparing the entropies leads to the conclusion that the entropy *decreases* in the semiclassical limit.⁶ A possibly related pathology with this parameterization

⁶This entropy is equal to the entropy of the twirled state $|\widetilde{\Phi}\rangle = e^{i\hat{p}h}|\Phi, f\rangle$ on the original crossed product algebra $\widehat{\mathcal{C}}_\lambda$. The density matrix for this state is given by equation (A.27), and assuming a slowly varying wavefunction, the entropy can be approximated by

$$S(\rho_{\widehat{\Phi}}) \approx \langle \Phi | h_{\Phi|\Omega} | \Phi \rangle - \langle f | \hat{q} + \log g(\hat{q}) | f \rangle = -S_{\text{rel}}^{\mathcal{C}'_\lambda}(\Phi||\Omega) - \langle f | \hat{q} + \log g(\hat{q}) | f \rangle, \quad (3.61)$$

is that the asymptotic charge is now given by $\frac{A_\infty^{(2)}}{4G_N} = -\hat{q} + h_\lambda$, which now depends on the horizon cut λ . The expectation value of this asymptotic charge then evolves as we change the cut, despite fixing the global state $|\widehat{\Phi}\rangle$. These issues indicate there we should not use above parameterization while also fixing the state $|\widehat{\Phi}\rangle$.

As the formulas (3.47) and (3.56) indicate, there are corrections to the entropy in the gravitational algebras starting at second order in the semiclassical expansion parameter ε . This immediately raises the question of whether the generalized second law continues to hold even after including these corrections. This question is nontrivial because there is not an obvious way in which the algebra $\widehat{\mathcal{C}}_\lambda$ embeds as a subalgebra of $\widehat{\mathcal{C}}_\lambda$, making it difficult to appeal to monotonicity results of entropies under algebra inclusions. In order to begin addressing these questions, in section 5 we compute the corrections to the semiclassical entropy expression, and discuss various bounds on these corrections. Unfortunately, these corrections do not ostensibly obey any monotonicity properties, but we leave open the possibility that a more careful analysis would produce such a result. We also explore in section 6 another approach in which we identify a subalgebra of $\widehat{\mathcal{C}}_\lambda$ unitarily equivalent to $\widehat{\mathcal{C}}_\lambda$, assuming the existence of an operator-valued weight between the underlying quantum field theory algebra. This may ultimately lead to improved monotonicity results for the subleading corrections to the crossed product entropies.

4 Algebra at infinity

The discussion of the GSL to this point has focused for simplicity on asymptotically AdS black holes. The Killing horizon is a Cauchy surface for these spacetimes since the Dirichlet boundary conditions imposed at the AdS boundary cause all outgoing modes to be reflected back through the black hole horizon. In this case, the quantum field algebra associated with the horizon is complete, and the gravitational algebra enlarges this only by a single additional asymptotic charge, the late time area perturbation $\frac{A_\infty^{(2)}}{4G_N}$. However, for black holes with alternative asymptotics, there are outgoing quantum field modes that never enter the black hole horizon. For asymptotically flat black holes, these include modes that propagate out to future timelike infinity i^+ or null infinity \mathcal{I}^+ , while for asymptotically de Sitter black holes, these include degrees of freedom beyond the cosmological horizon (see figure 2). The analysis must therefore be generalized to allow for the possibility of these nontrivial algebras, which we refer to as the algebra at infinity.

In this section, we describe the treatment of the generalized second law with nontrivial algebras at infinity. In section 4.1, we describe how this algebra at infinity is naturally incorporated into algebraic structures associated with half-sided translations, which allow a generalization of Wall's discussion of the GSL with an algebra at infinity that applies to interacting field theories. Following this, section 4.2 describes the general construction of the

which now involves the relative entropy for the commutant quantum field theory algebra \mathcal{C}'_λ . Since the inclusion for the commutant algebras is reversed, $\mathcal{C}'_\lambda \subset \mathcal{C}'_\lambda$, monotonicity of relative entropy now causes the crossed product entropy to decrease between cuts.




Figure 2: The extended Schwarzschild-de Sitter spacetime provides an example where one finds an algebra at infinity for the horizon translation. This spacetime is described as an infinite chain of black hole and cosmological horizons, indicated by the ellipses “...” in the Penrose diagram above. The algebra for a cut c_λ is defined as all quantum fields supported in the region spacelike to the right of the cut, shown by the red lines above. The algebra at infinity is associated with the region R_∞ shown in purple, consisting of fields localized beyond the cosmological horizon.

gravitational algebra in these contexts. Finally, section 4.3 discusses some subtleties that can arise when analyzing rotating black holes and backgrounds involving the Unruh vacuum.

4.1 Semiclassical GSL

Intuitively, the existence of additional degrees of freedom localized far from the black hole should not affect the derivation of the GSL, since any entropy associated with these degrees of freedom would appear in the algebra associated with any horizon cut; it should therefore cancel out in when considering entropy differences between two cuts. Wall gave a precise version of this argument assuming the existence of a factorization between the algebra associated with the horizon and the algebra associated with \mathcal{I}^+ [38]. Such a factorization is expected to hold for free field theories [82], and was recently utilized for the construction of gravitational algebras associated with arbitrary Killing horizons [32].

However, there are reasons to believe that this factorization property fails for interacting theories. Correlation functions of local operators in interacting field theories have stronger divergences than in free theories, which necessitates smearing the fields in timelike directions in order to obtain operators with finite fluctuations [38, 83, 84]. Smearing only on the horizon is not sufficient, implying that there is no nontrivial algebra of bounded operators localized solely on a horizon cut \mathcal{H}_λ^+ [50]. This prevents a decomposition of the horizon cut algebra \mathcal{C}_λ into a tensor product $\mathcal{A}(\mathcal{H}_\lambda^+) \otimes \mathcal{A}(\infty)$, since $\mathcal{A}(\mathcal{H}_\lambda^+)$ is trivial. Fortunately, the existence of a tensor factorization is not strictly necessary in order to derive the GSL. Instead, we will make use of a weaker condition on the structure of the horizon cut algebra \mathcal{C}_λ that follows

from the properties of half-sided translations.

The argument from section 3.1 that the smeared average null energy operator P_λ generates a half-sided translation continues to hold for spacetimes with non-AdS asymptotics, provided that the horizon cut algebra \mathcal{C}_λ is defined as the double commutant of local operators on a Cauchy slice consisting of the future horizon cut \mathcal{H}_λ^+ and a component Σ^{asy} spacelike separated from the horizon. For example, in asymptotically flat spacetimes, Σ^{asy} consists of \mathcal{I}^+ and a contribution from i^+ if massive fields are present. Since P_λ is an integral of the stress tensor only over the horizon \mathcal{H} , it acts trivially on operators localized to Σ^{asy} , which are associated purely with outgoing modes that do not interact with the horizon \mathcal{H} . This allows us to characterize the algebra \mathcal{B}_∞ associated with Σ^{asy} as the collection of operators in \mathcal{B} (the algebra associated with the exterior of the bifurcation surface) that are invariant under the translations,

$$\mathcal{B}_\infty := \{a \in \mathcal{B} | e^{itP_\lambda} a e^{-itP_\lambda} = a\} \quad (4.1)$$

The properties of such a fixed point algebra under the action of a half-sided translation were studied by Borchers [85], who showed that the algebra can equivalently be defined as the operators contained in the translated algebras $\mathcal{C}_{t\lambda} = e^{itP_\lambda} \mathcal{B} e^{-itP_\lambda}$ for all $t \geq 0$,

$$\mathcal{B}_\infty = \bigwedge_{t \geq 0} \mathcal{C}_{t\lambda}. \quad (4.2)$$

Since $\mathcal{C}_{t\lambda}$ defines an algebra associated with a cut further to the future for larger values of t , the limiting algebra \mathcal{B}_∞ consists of modes that never fall across the horizon, and hence escape to the region associated with Σ^{asy} . The fact that the definitions (4.1) and (4.2) agree justifies the statement above that the operators localized to the asymptotic regions are fixed by the action of P_λ .

One can further show that the algebra \mathcal{B}_∞ is preserved by the modular flow of \mathcal{B} in a vacuum state $|\Omega\rangle$ for P_λ , $\Delta_{\mathcal{B}}^{-is} \mathcal{B}_\infty \Delta_{\mathcal{B}}^{is} = \mathcal{B}_\infty$. By Takesaki's theorem [86], this implies the existence of a conditional expectation from \mathcal{B} to \mathcal{B}_∞ . Recall that a conditional expectation is a completely positive linear map $E : \mathcal{B} \rightarrow \mathcal{B}_\infty$ such that $E(b) = b$ for all $b \in \mathcal{B}_\infty$ and satisfies the bimodule property $E(b_1 a b_2) = b_1 E(a) b_2$ for $b_1, b_2 \in \mathcal{B}_\infty$, $a \in \mathcal{B}$ [81, 87]. Takesaki's theorem further guarantees that the conditional expectation preserves the state $\omega = \langle \Omega | \cdot | \Omega \rangle$ on \mathcal{B} , in the sense that $\omega \circ E(b) \equiv \omega(E(b)) = \omega(b)$ for $b \in \mathcal{B}$. It is straightforward to see that \mathcal{B}_∞ is also preserved by the modular flow for any horizon-cut algebra \mathcal{C}_λ , and therefore each of these has an associated conditional expectation $E_\lambda : \mathcal{C}_\lambda \rightarrow \mathcal{B}_\infty$.

Assuming \mathcal{B}_∞ is a factor, the existence of the conditional expectation E_λ implies that \mathcal{B}_∞ and its relative commutant $\mathcal{B}_{\infty,\lambda}^c = \mathcal{B}'_\infty \wedge \mathcal{C}_\lambda$ embed into \mathcal{C}_λ as a tensor product, i.e. $\mathcal{B}_\infty \otimes \mathcal{B}_{\infty,\lambda}^c \cong \mathcal{B}_\infty \vee \mathcal{B}_{\infty,\lambda}^c \subset \mathcal{C}_\lambda$ [86]. If in fact we have equality $\mathcal{B}_\infty \vee \mathcal{B}_{\infty,\lambda}^c = \mathcal{C}_\lambda$, the inclusion $\mathcal{B}_\infty \subset \mathcal{C}_\lambda$ is called *conormal* [88], and along with the existence of the conditional expectation, this implies that \mathcal{C}_λ tensor factorizes $\mathcal{B} = \mathcal{B}_\infty \otimes \mathcal{B}_{\infty,\lambda}^c$ into an algebra at infinity \mathcal{B}_∞ and its relative commutant $\mathcal{B}_{\infty,\lambda}^c$. In this case the relative commutant is naturally associated with a nontrivial horizon algebra $\mathcal{B}_{\infty,\lambda}^c = \mathcal{A}(\mathcal{H}_\lambda^+)$, which is the setup expected for free quantum fields. For interacting quantum fields, we do not expect a nontrivial horizon algebra, and hence may conjecture that the relative commutant is trivial, $\mathcal{B}_{\infty,\lambda}^c = \mathbb{C}\mathbb{1}$, in which case the




Figure 3: The Penrose diagram of the conformally extended Schwarzschild solution is shown above. The standard maximally extended Schwarzschild solution is the region shown in yellow, and the triangular regions in the corners are the conformal extension of the spacetime beyond \mathcal{I}^+ and \mathcal{I}^- . The algebra of massless or conformal fields in this spacetime naturally allows for smearing of the fields in the extended regions. This motivates defining the algebra at infinity for the Schwarzschild black hole to be associated with the region R_∞ shown in purple, and this region captures the notion of outgoing quantum field modes that never enter the black hole horizon.

inclusion $\mathcal{B}_\infty \subset \mathcal{C}_\lambda$ is called *singular* [88]. Since our discussion of the GSL generically involves interacting matter fields but free gravitons, we will be in an intermediate case where $\mathcal{B}_{\infty,\lambda}^c$ is nontrivial and describes the graviton horizon algebra, but $\mathcal{B}_\infty \vee \mathcal{B}_{\infty,\lambda}^c \neq \mathcal{C}_\lambda$ since the matter fields do not have a horizon algebra.

This discussion indicates that the existence of the conditional expectation E_λ is a generalization of the notion of a tensor factorization of the algebra \mathcal{C}_λ . When \mathcal{C}_λ does factorize, there is a class of tensor product states $\omega = \omega_\infty \otimes \omega_{\mathcal{H}_\lambda^+}$ on $\mathcal{C}_\lambda = \mathcal{B}_\infty \otimes \mathcal{B}_{\infty,\lambda}^c$ that contain no correlations between the two subfactors, i.e. $\omega(\mathbf{bc}) = \omega(\mathbf{b})\omega(\mathbf{c})$ for $\mathbf{b} \in \mathcal{B}_\infty$ and $\mathbf{c} \in \mathcal{B}_{\infty,\lambda}^c$. The states that are preserved by the conditional expectation, $\omega = \omega \circ E_\lambda$, are product states which satisfy $E_\lambda(\mathbf{c}) = \omega_{\mathcal{H}_\lambda^+}(\mathbf{c})\mathbb{1}$ for all $\mathbf{c} \in \mathcal{B}_{\infty,\lambda}^c$. This shows that the conditional expectation E_λ encodes the information of a state $\omega_{\mathcal{H}_\lambda^+}$ on $\mathcal{B}_{\infty,\lambda}^c$, and different choices of this state result in different conditional expectations from \mathcal{C}_λ to \mathcal{B}_∞ . When the inclusion $\mathcal{B}_\infty \subset \mathcal{C}_\lambda$ is singular, the conditional expectation E_λ is unique [87, Section IX.4]. In this case, there is no algebra on which to define a horizon state $\omega_{\mathcal{H}_\lambda^+}$, but the analog of tensor factorized states are states fixed by the conditional expectation. Such states can be thought of as vacuum states for fields localized very close to the horizon.

A question that arises is why the algebra associated with the horizon is trivial in interacting theories, while the algebra at infinity can remain nontrivial. For example, in the asymptotically flat Schwarzschild black hole, both algebras appear to be associated with operators smeared on null surfaces (\mathcal{H}_λ^+ or \mathcal{I}^+), and so it is not immediately clear what distinguishes the two. The resolution is that the algebra at infinity is nontrivial in interacting theories whenever it is associated with an open region of spacetime. This is easiest to see in the Schwarzschild-de Sitter spacetime, where the algebra at infinity consists of all operators smeared in the region beyond the cosmological horizon, which is an open spacetime region (see figure 2).

For the asymptotically flat Schwarzschild spacetime, we can give a similar argument by considering a conformal extension of the usual spacetime to include regions beyond \mathcal{I}^+ , \mathcal{I}^- into which modes propagating out to infinity enter. These are the triangular regions depicted in figure 3 added on to the standard Penrose diagram of the Schwarzschild spacetime. This conformal extension of Schwarzschild can be constructed by analytic continuation, in which case the regions beyond \mathcal{I}^\pm are conformal to the negative mass (or equivalently, negative r) Schwarzschild solution [89]. This extended spacetime is similar to the conformal embedding of Minkowski spacetime into the Einstein static universe [90], see figure 4. When dealing with conformal matter fields, the net of algebras associated with the Schwarzschild spacetime should naturally define a net on the extended spacetime. In this case, we see that the algebra at infinity can be characterized by all fields smeared in the open region R_∞ added beyond \mathcal{I}^+ in the conformal extension of the spacetime.⁷

Figure 4 also highlights the difference between black hole spacetimes and a Rindler horizon. When embedded into the Einstein static universe, the Rindler patch covers a causal diamond whose boundary intersects \mathcal{I}^+ only along a single generator. This corresponds to the fact that only light rays propagating in the same direction parallel to the Rindler horizon will remain in the Rindler patch when they intersect \mathcal{I}^+ . In this case, the algebra at infinity is trivial, since the region at infinity only consists of this single null generator along \mathcal{I}^+ . Hence, asymptotically flat black holes provide an interesting arena to study half-sided translations with nontrivial algebras at infinity, which is not available when restricting to quantum field theory in Minkowski space.

As mentioned above, any algebraic state arising from an average null energy vacuum $|\Omega\rangle$ is fixed by the conditional expectation E_λ . When the algebra \mathcal{B}_∞ is nontrivial, the vacuum $|\Omega\rangle$ is no longer unique; any other state of the form $\mathbf{b}|\Omega\rangle$ with $\mathbf{b} \in \mathcal{B}_\infty$ is also a vacuum state, since $[P_\lambda, \mathbf{b}] = 0$. For any such vacuum, its modular flow maps \mathcal{B}_∞ into itself. This suggests that the modular Hamiltonian will decompose into a component associated with \mathcal{B}_∞ and a component associated with the horizon. To describe this decomposition, we first note that P_λ also generates an inverse half-sided translation for the commutant algebras \mathcal{C}'_λ , and these similarly must contain a nontrivial translation-invariant algebra $\mathcal{B}_{-\infty}'$ that is naturally

⁷There may be pathologies associated with the existence of naked singularities in this conformal extension, since the new region has the geometry of the negative mass Schwarzschild spacetime. It seems likely that these can be handled with appropriate boundary conditions, the details of which do not matter for the present discussion. It is only relevant that there is an open region beyond \mathcal{I}^+ in which we can smear fields in order to obtain a nontrivial algebra at infinity.




Figure 4: Shown above are the conformal extensions of the Schwarzschild and Minkowski spacetimes which are asymptotically conformal to the Einstein static universe near spatial infinity i^0 , which shrinks to a point in this conformal compactification. The algebra at infinity for the Schwarzschild solution lives in the purple region R_∞ . Note that the spacetime can have pathologies such as naked singularities in the region R_∞ and singular behavior near i^+ , but the figure still motivates defining a nontrivial algebra for these regions. In the conformal extension of Minkowski space, the maroon region indicates the Rindler wedge, and we see that the region at infinity for this patch is the single generator of \mathcal{S}^+ shown in purple. Because this is not an open region in the Einstein static universe, we do not expect an algebra at infinity for half-sided translations associated with Rindler horizons in higher than two spacetime dimensions.

associated with the algebra at minus infinity on the left half of the black hole spacetime, containing \mathcal{S}_L^- and i_L^- . This algebra is equivalently characterized as $\mathcal{B}_{-\infty'} = J_\Omega \mathcal{B}_\infty J_\Omega$, where J_Ω is the modular conjugation for a vacuum state $|\Omega\rangle$. Modular flow in the commutant algebras again maps $\mathcal{B}_{-\infty'}$ into itself, and there is a collection of conditional expectations $E'_\lambda : \mathcal{C}'_\lambda \rightarrow \mathcal{B}_{-\infty'}$ which fix the algebraic states on \mathcal{C}'_λ defined by the vacuum vector $|\Omega\rangle$.

Because states fixed by a conditional expectation are the analogs of tensor factorized states, they represent configurations in which there is no correlation between operators in \mathcal{B}_∞ and the ingoing modes on the horizon. However, $|\Omega\rangle$ is a separating state for \mathcal{B}_∞ , meaning that this algebra is entangled with something. The purifying degrees of freedom must be the complementary algebra at minus infinity, $\mathcal{B}_{-\infty'}$. Defining the two-sided algebra at infinity as the algebraic union of these two,

$$\mathcal{A}_\infty = \mathcal{B}_\infty \vee \mathcal{B}_{-\infty'}, \quad (4.3)$$

this discussion suggests that $|\Omega\rangle$ defines a pure state on \mathcal{A}_∞ , which, in particular, implies that \mathcal{A}_∞ is type I. Its commutant $\mathcal{A}_{\mathcal{H}} = \mathcal{A}'_\infty$ contains operators localized to the complete Killing horizon. Note that $\mathcal{A}_{\mathcal{H}}$ is nontrivial even in the interacting case where there is no nontrivial horizon algebra localized to the future of a cut. At minimum, $\mathcal{A}_{\mathcal{H}}$ contains the spectral projections of the translation generator P_λ .

Since the modular operator h_Ω^λ for the horizon cut algebra \mathcal{C}_λ maps both \mathcal{B}_∞ and $\mathcal{B}_{-\infty'}$ into themselves, it also maps \mathcal{A}_∞ into itself and similarly for the horizon algebra $\mathcal{A}_{\mathcal{H}}$. As both \mathcal{A}_∞ and $\mathcal{A}_{\mathcal{H}}$ are type I, this suggests that h_Ω^λ generates inner automorphisms of these algebras, meaning that it would be possible to find operators h_∞ affiliated with \mathcal{A}_∞ and $h_{\mathcal{H}}^\lambda$ affiliated with $\mathcal{A}_{\mathcal{H}}$ such that

$$h_\Omega^\lambda = h_\infty + h_{\mathcal{H}}^\lambda. \quad (4.4)$$

At present we do not have a proof of this factorization of the modular Hamiltonian, in part due to the fact that the algebras \mathcal{A}_∞ and $\mathcal{A}_{\mathcal{H}}$ are not factors when the inclusion $\mathcal{B}_\infty \subset \mathcal{C}_\lambda$ has trivial relative commutant. However, we can motivate the existence of $h_{\mathcal{H}}^\lambda$ as a well-defined operator by appealing to the free field case. In that case, the horizon cut algebra factorizes as $\mathcal{C}_\lambda = \mathcal{C}_\lambda^{\mathcal{H}} \otimes \mathcal{B}_\infty$, and since there is now a quantum field algebra localized to the horizon to the future of the cut, we expect that the full horizon algebra $\mathcal{A}_{\mathcal{H}}$ becomes a factor. In this case the modular operator factorizes between a horizon component and component at infinity, $\Delta_{\Omega,\lambda} = \Delta_{\mathcal{H},\lambda} \Delta_\infty$, implying that the modular Hamiltonian exhibits the decomposition (4.4). On the other hand, $h_{\mathcal{H}}^\lambda$ has an expression (3.11) in terms of the integral of the stress tensor over the horizon, and its independent definition in terms of modular theory means that this operator has finite fluctuations in a dense class of states, despite only being smeared on a codimension-1 null surface. Since the two-point function of the stress tensor in an interacting CFT takes the same form as in a free theory, we expect that $h_{\mathcal{H}}^\lambda$ will have finite fluctuations for interacting theories as well. Hence, it is plausible that the factorization (4.4) holds even in interacting theories where the inclusion $\mathcal{B}_\infty \subset \mathcal{C}_\lambda$ is singular.

An important aspect of this splitting is that any vacuum state $|\Omega\rangle$ gives rise to the same horizon component of the modular Hamiltonian $h_{\mathcal{H}}^\lambda$. This follows from the fact that any two such vacua $|\Omega_{1,2}\rangle$ induce states on \mathcal{C}_λ that are fixed by the conditional expectation E_λ , and similarly the states $\omega'_{1,2}$ induced on \mathcal{C}'_λ are fixed by the complementary conditional expectation E'_λ . Hence the Connes cocycle $u_{2|1}(s) = \Delta_{\Omega_2}^{is} \Delta_{\Omega_1|\Omega_2}^{-is}$ for these two states is an element of \mathcal{B}_∞ , and the commutant cocycle $u'_{2|1}(s) = \Delta_{\Omega_1|\Omega_2}^{-is} \Delta_{\Omega_1}^{is}$ is an element of $\mathcal{B}_{-\infty'}$ [87, Chapter IX, corollary 4.22(ii)], which then implies that $\mathbf{b} := h_{\Omega_2}^\lambda - h_{\Omega_1|\Omega_2}^\lambda$ is affiliated with \mathcal{B}_∞ and $\mathbf{b}' := h_{\Omega_1|\Omega_2}^\lambda - h_{\Omega_1}^\lambda$ is affiliated with $\mathcal{B}_{-\infty'}$. We therefore have that

$$h_{\Omega_2}^\lambda - h_{\Omega_1}^\lambda = \mathbf{b} + \mathbf{b}' \quad \implies \quad h_{\mathcal{H},2}^\lambda - h_{\mathcal{H},1}^\lambda = \mathbf{b} + \mathbf{b}' - h_{\infty,2} + h_{\infty,1} \text{ affiliated with } \mathcal{A}_\infty. \quad (4.5)$$

Thus, the difference $h_{\mathcal{H},2}^\lambda - h_{\mathcal{H},1}^\lambda$ must be affiliated with the center of $\mathcal{A}_{\mathcal{H}}$. Since the decomposition (4.4) is ambiguous up to compensating shifts of $h_{\mathcal{H}}^\lambda$ and h_∞ by elements of the center, we can consistently choose the decomposition such that $h_{\mathcal{H}}^\lambda$ is the same for all vacuum states $|\Omega\rangle$. This point connects with the discussion of section 3.1 that the commutation relation with P_λ uniquely fixes the modular Hamiltonian of a vacuum state up to terms that commute with P_λ . Since h_∞ is affiliated with $\mathcal{A}'_{\mathcal{H}}$, it commutes with P_λ , and hence $h_{\mathcal{H}}^\lambda$ alone satisfies the commutation relation (3.7) associated with the half-sided translation.

We now apply this reasoning to the generalized second law. Given a vacuum state $|\Omega\rangle$, we formally split the modular Hamiltonian into one-sided components $h_\Omega^\lambda = K_\lambda - K'_\lambda$, defined as usual as sesquilinear forms respectively affiliated with \mathcal{C}_λ and \mathcal{C}'_λ . This decomposition can be refined by performing a similar split on each of h_∞ and $h_{\mathcal{H}}^\lambda$ appearing in (4.4), resulting in $K_\lambda = K_{\mathcal{H}}^\lambda + K_\infty$, with K_∞ affiliated with \mathcal{B}_∞ and $K_{\mathcal{H}}^\lambda$ affiliated with \mathcal{B}'_∞ . The validity of this decomposition only relies on the existence of the conditional expectation $E_\lambda : \mathcal{C}_\lambda \rightarrow \mathcal{B}_\infty$, and holds even in interacting theories where we cannot assume a tensor product structure between \mathcal{B}_∞ and a horizon algebra. This decomposition of K_λ is all that is needed to apply Wall's argument for the GSL in setups where the algebra at infinity is nontrivial. The relative entropy between an arbitrary state $|\Phi\rangle$ and a translation vacuum $|\Omega\rangle$ is give by

$$\begin{aligned} S_{\text{rel}}^{\mathcal{C}_\lambda}(\Phi||\Omega) &= -S_{\Phi,\lambda}^{\text{QFT}} + \langle K_{\mathcal{H}}^\lambda + K_\infty \rangle_\Phi + S_{\Omega,\lambda}^{\text{QFT}} \\ &= -S_{\Phi,\lambda}^{\text{QFT}} - \left\langle \frac{A_\lambda}{4G_N} \right\rangle_\Phi + \left\langle \frac{A_\infty}{4G_N} \right\rangle_\Phi + \langle K_\infty \rangle_\Phi + S_{\Omega,\lambda}^{\text{QFT}} \end{aligned} \quad (4.6)$$

applying the identity (3.32) and the fact that $K_{\mathcal{H}}^\lambda$ continues to be given by the stress tensor integral $\frac{2\pi}{\kappa} H_{\zeta_\lambda}^+$ appearing in that relation. When considering relative entropy differences between two horizon cuts, the last three terms in (4.6) will cancel since the vacuum entanglement entropy is independent of the horizon cut and because K_∞ and $\frac{A_\infty}{4G_N}$ are forms affiliated with the algebra at infinity whose expectation values are the same in the state $|\Phi\rangle$ for any value of λ . Monotonicity of relative entropy again reduces to the GSL for this setup.

One interesting aspect of this discussion is that we can employ any vacuum state $|\Omega\rangle$ when applying monotonicity of relative entropy in (4.6). This independence of the choice of vacuum can be made more explicit by invoking an identity satisfied by the relative entropy in the presence of a conditional expectation [81, 91–93]. Using the notation $\omega = \langle \Omega | \cdot | \Omega \rangle$, $\varphi = \langle \Phi | \cdot | \Phi \rangle$ as states on \mathcal{C}_λ , and recalling $\omega = \omega \circ E_\lambda$, the identity reads

$$S_{\text{rel}}^{\mathcal{C}_\lambda}(\varphi||\omega) = S_{\text{rel}}^{\mathcal{B}_\infty}(\varphi|_{\mathcal{B}_\infty} || \omega|_{\mathcal{B}_\infty}) + S_{\text{rel}}^{\mathcal{C}_\lambda}(\varphi||\varphi \circ E_\lambda). \quad (4.7)$$

The first term on the right involves the relative entropy of the two states restricted to \mathcal{B}_∞ , and hence is the same when computed for two different horizon cuts. Monotonicity of the relative entropy on the left between horizon cuts therefore implies that $S_{\text{rel}}^{\mathcal{C}_\lambda}(\varphi||\varphi \circ E_\lambda)$ is also monotonic. Since this relative entropy only refers to the state φ and the conditional expectation E_λ , it is manifestly independent of the choice of vacuum state ω . In terms of entropies, it can be expressed as

$$\begin{aligned} S_{\text{rel}}^{\mathcal{C}_\lambda}(\varphi||\varphi \circ E_\lambda) &= -S_{\Phi,\lambda}^{\text{QFT}} + \langle K_{\mathcal{H}}^\lambda + K_\infty^\Phi \rangle_\Phi + S_{\varphi \circ E_\lambda}^{\text{QFT}} \\ &= -S_{\Phi,\lambda}^{\text{QFT}} - \left\langle \frac{A_\lambda}{4G_N} \right\rangle_\Phi + \left\langle \frac{A_\infty}{4G_N} \right\rangle_\Phi + S_{\varphi \circ E_\lambda}^{\text{QFT}} \end{aligned} \quad (4.8)$$

using that $\langle K_\infty^\Phi \rangle_\Phi = 0$ and applying (3.32). Since $\varphi \circ E_\lambda$ is a vacuum state, the entropy $S_{\varphi \circ E_\lambda}^{\text{QFT}}$ is independent of the cut λ , and since the expectation value of A_∞ is also independent of the cut, monotonicity of $S_{\text{rel}}^{\mathcal{C}_\lambda}(\varphi||\varphi \circ E_\lambda)$ again implies the GSL.

4.2 Gravitational algebra

The discussion up to this point has been at the level of relative entropies in quantum field theory. We now describe how to incorporate the gravitational constraints to arrive at a semi-classical second law for a gravitational algebra that incorporates asymptotic charges. We will specialize to asymptotically flat spacetimes with massless matter fields in this section, but it should be clear that similar arguments apply to other asymptotics such as cosmological black holes. As emphasized by KFLS [32], because the null Cauchy surface for our subregion naturally splits into a component on the black hole horizon \mathcal{H}_λ^+ and a component associated with \mathcal{I}^+ , the gravitational constraints similarly decompose into independent constraints on each component of the Cauchy surface. This allows for a construction of the gravitational algebra in steps by introducing individual asymptotic charges associated with each constraint and constructing the crossed product algebras. Following this, there is an additional constraint arising from matching conditions where the two Cauchy surfaces meet (i.e. across i^+ in the asymptotically flat case). The matching conditions determine some of the gravitational charges in terms of other quantities, so that the final algebra contains only asymptotic charges at spatial infinity. We describe the details of this procedure below, and then demonstrate it explicitly for an asymptotically flat Schwarzschild black hole.

In order to keep the discussion as general as possible, it is convenient to assume that the constraints have already been imposed on the algebra at infinity, so that \mathcal{B}_∞ already incorporates asymptotic charges associated with \mathcal{I}^+ such as the ADM mass or late-time Bondi mass. We will assume that this results in a semifinite algebra, meaning that \mathcal{B}_∞ is type I or type II and possesses a normal, faithful, semifinite trace, which we denote Tr_∞ . We also expect \mathcal{B}_∞ to be properly infinite (i.e. type I_∞ or type II_∞), meaning that $\text{Tr}_\infty \mathbb{1} = \infty$, so Tr_∞ does not define a normalizable state. Instead, Tr_∞ is a normal, faithful, semifinite weight, which still has a well-defined modular theory associated with it [87, Chapter VIII]. We continue to assume that \mathcal{B}_∞ is the translation-invariant algebra of the horizon cut algebra \mathcal{C}_λ (which remains type III_1), and hence we can define a weight on \mathcal{C}_λ by composing Tr_∞ with the conditional expectation E_λ ,

$$\omega_{\text{Tr}} := \text{Tr}_\infty \circ E_\lambda. \tag{4.9}$$

The modular flow for ω_{Tr} reproduces the modular flow associated with Tr_∞ on operators in \mathcal{B}_∞ , which is trivial. Hence, \mathcal{B}_∞ is the centralizer of this modular flow, meaning that these operators commute with the modular Hamiltonian h_{Tr}^λ for this weight.⁸ Using the decomposition (4.4), valid for modular Hamiltonians of weights fixed by E_λ , we find that $h_{\text{Tr},\infty} = 0$ and hence $h_{\text{Tr}}^\lambda = h_{\mathcal{H}}^\lambda$, which is uniquely determined by the conditional expectation E_λ . Hence when \mathcal{B}_∞ is semifinite, the geometric flow generated by $h_{\mathcal{H}}^\lambda$ which fixes points on \mathcal{I}^+ and acts like a boost around the horizon cut is the modular flow of a weight defined on \mathcal{C}_λ .

⁸To define h_{Tr}^λ as an operator on the full Hilbert space, we must also specify a weight for the commutant algebra \mathcal{C}'_λ , which we can define using the trace $\text{Tr}_{-\infty}$ on $\mathcal{B}_{-\infty}$ and the complementary conditional expectation $E'_\lambda : \mathcal{C}_\lambda \rightarrow \mathcal{B}_{-\infty}$. Given such a pair of weights, we can always find a Hamiltonian h_{Tr}^λ which generates modular flow on \mathcal{C}_λ and backwards modular flow in \mathcal{C}'_λ [45, Lemma 5.11].

The procedure for imposing the gravitational constraints is now exactly the same as in section 3.3. We introduce the additional asymptotic charge

$$\hat{y} = -\frac{A_\infty^{(2)}}{4G_N} \quad (4.10)$$

acting on a Hilbert space $\mathcal{H}_{\text{asy}} = L^2(\mathbb{R}_y)$, and determine the operators that commute with the gravitational constraint associated with the boost diffeomorphism which $h_{\mathcal{H}}^\lambda$ generates. When represented on $\tilde{\mathcal{H}} = \hat{\mathcal{H}} \otimes \mathcal{H}_{\text{asy}}$, where $\hat{\mathcal{H}}$ includes the QFT Hilbert space as well as the Hilbert space for asymptotic charges in \mathcal{B}_∞ , these operators form a crossed product algebra

$$\tilde{\mathcal{C}}_\lambda = \left\langle e^{i\hat{k}h_{\mathcal{H}}^\lambda} \mathbf{a} e^{-i\hat{k}h_{\mathcal{H}}^\lambda}, \hat{y} \right\rangle, \quad \mathbf{a} \in \mathcal{C}_\lambda, \quad \hat{k} = -i \frac{d}{dy}. \quad (4.11)$$

Because $h_{\mathcal{H}}^\lambda$ is the modular Hamiltonian of the weight ω_{Tr} , $\tilde{\mathcal{C}}_\lambda$ is the crossed product of a type III₁ quantum field theory algebra by a modular automorphism group, and hence is type II_∞ and possesses a trace.

$\tilde{\mathcal{C}}_\lambda$ is not the final gravitational algebra associated with the exterior of the cut since we still need to impose the constraint coming from the matching condition at i^+ . This condition arises because the Hilbert space $\tilde{\mathcal{H}}$ contains two operators that measure the time shift at i^+ relative to i_L^- in the complementary region. One of these is the operator \hat{k} appearing in the crossed product, which is conjugate to the asymptotic area operator \hat{y} which generates forward time evolution along the horizon. This gives \hat{k} the interpretation of a renormalized Killing time difference between i_L^- and the cut of \mathcal{H}_λ^+ in the distant future where one matches the horizon to a spacelike hypersurface connecting it to \mathcal{I}^+ . The second time operator is associated with time translations along \mathcal{I}^+ . We suppose that \mathcal{B}_∞ contains an energy operator \hat{x} generating forward time evolution along \mathcal{I}^+ , which, due to the semiclassical assumptions has a continuous spectrum and is unbounded above and below. It therefore possesses a conjugate time operator which we denote \hat{p} which measures the time shift between i_L^- and the cut of \mathcal{I}^+ where one matches the Cauchy surface connecting it to the black hole horizon. Since these two notions of time shift must agree, we arrive at the final matching constraint

$$\mathfrak{C}_{\text{match}} = \hat{k} - \hat{p} = 0. \quad (4.12)$$

To impose this constraint on the algebra, we first look for operators in $\tilde{\mathcal{C}}_\lambda$ that commute with $\hat{k} - \hat{p}$. This subalgebra

$$\tilde{\mathcal{C}}_\lambda^0 = \tilde{\mathcal{C}}_\lambda \wedge \left\langle \hat{k} - \hat{p} \right\rangle' \quad (4.13)$$

of $\tilde{\mathcal{C}}_\lambda$ is isomorphic to the desired gravitational algebra, but represented on a Hilbert space that contains other operators that do not preserve the constraint (4.12). We can remedy this by passing to a Hilbert space of coinvariants, following the procedure outlined by CLPW [31]. This Hilbert space is a quotient of $\tilde{\mathcal{H}}$ obtained by identifying vectors that differ by a state in the image of $\hat{k} - \hat{p}$, so that $|\Omega\rangle \sim |\Omega\rangle + (\hat{k} - \hat{p})|\chi\rangle$. Passing to this space of coinvariants can be used to eliminate the factor of \mathcal{H}_{asy} from the Hilbert space, so that the physical Hilbert space is identified with $\hat{\mathcal{H}}$. This identification is achieved by defining the operator

$$V := \langle 0_k | e^{-i\hat{y}\hat{p}}, \quad (4.14)$$

which maps $\tilde{\mathcal{H}}$ to $\hat{\mathcal{H}}$, and annihilates states of the form $(\hat{k} - \hat{p})|\chi\rangle$. This allows us to map any operator in $\mathfrak{c}_0 \in \tilde{\mathcal{C}}_\lambda^0$ to an operator $\hat{\mathfrak{c}}$ acting on $\hat{\mathcal{H}}$ by the relation

$$V\mathfrak{c}_0 = \hat{\mathfrak{c}}V. \quad (4.15)$$

The algebra in $\mathcal{B}(\hat{\mathcal{H}})$ generated by all such $\hat{\mathfrak{c}}$ is the final result for the gravitational algebra $\hat{\mathcal{C}}_\lambda$.⁹

A subtlety arises at this point regarding the existence of a trace on $\hat{\mathcal{C}}_\lambda$. The issue is that the trace defined on $\tilde{\mathcal{C}}_\lambda$ is a semifinite weight, meaning it assigns infinite values to some operators in $\tilde{\mathcal{C}}_\lambda$, although it is finite on sufficiently many operators that they generate the full algebra. There is no guarantee that the trace defines a semifinite weight when restricted to a subalgebra, since the subalgebra may consist mostly of operators with infinite trace. Hence, we need to separately analyze whether $\hat{\mathcal{C}}_\lambda$ admits a trace. We will describe here a set of sufficient conditions that ensure this is the case.

Since \hat{p} generates an automorphism of \mathcal{B}_∞ , we can assume that this algebra takes the form of a crossed product $\mathcal{B}_\infty = \overline{\mathcal{B}_\infty^0} \rtimes \mathbb{R}$, where $\overline{\mathcal{B}_\infty^0}$ is the subalgebra of operators commuting with \hat{p} , $\overline{\mathcal{B}_\infty^0} = \mathcal{B}_\infty \wedge \langle \hat{p} \rangle'$. Denoting H_∞ as the generator of the automorphism on $\overline{\mathcal{B}_\infty^0}$ for this crossed product, the algebra \mathcal{B}_∞ takes the form

$$\mathcal{B}_\infty = \langle e^{i\hat{p}H_\infty} \mathfrak{b} e^{-i\hat{p}H_\infty}, \hat{x} \rangle, \quad \mathfrak{b} \in \mathcal{B}_\infty^0 \cong \overline{\mathcal{B}_\infty^0}, \quad [\hat{x}, \hat{p}] = i. \quad (4.16)$$

The algebra \mathcal{B}_∞^0 simply consists of quantum field theory operators without the dressing factors $e^{i\hat{p}H_\infty}(\cdot)e^{-i\hat{p}H_\infty}$. The full horizon cut algebra also takes the form of a crossed product: defining $\overline{\mathcal{C}_\lambda^0} = \mathcal{C}_\lambda \wedge \langle \hat{p} \rangle'$, we find that

$$\mathcal{C}_\lambda = \langle e^{i\hat{p}H_\infty} \mathfrak{c} e^{-i\hat{p}H_\infty}, \hat{x} \rangle, \quad \mathfrak{c} \in \mathcal{C}_\lambda^0 \cong \overline{\mathcal{C}_\lambda^0}. \quad (4.17)$$

The unconstrained gravitational algebra $\tilde{\mathcal{C}}_\lambda$ then is a crossed product of \mathcal{C}_λ^0 by \mathbb{R}^2 ,

$$\tilde{\mathcal{C}}_\lambda = \langle e^{i\hat{k}h_{\mathcal{H}}^\lambda} e^{i\hat{p}H_\infty} \mathfrak{c} e^{-i\hat{p}H_\infty} e^{-i\hat{k}h_{\mathcal{H}}^\lambda}, \hat{x}, \hat{y} \rangle, \quad \mathfrak{c} \in \mathcal{C}_\lambda^0, \quad (4.18)$$

and when imposing the matching constraint (4.12), we find that only the combination $\hat{x} + \hat{y}$ remains in the final gravitational algebra. The relation (4.15) then maps $\hat{x} + \hat{y}$ to \hat{x} when representing this algebra on $\hat{\mathcal{H}}$, so that the final gravitational algebra is just a single crossed product by the combined automorphism generated by $h_{\mathcal{H}}^\lambda + H_\infty$,

$$\hat{\mathcal{C}}_\lambda = \langle e^{i\hat{p}(h_{\mathcal{H}}^\lambda + H_\infty)} \mathfrak{c} e^{-i\hat{p}(h_{\mathcal{H}}^\lambda + H_\infty)}, \hat{x} \rangle, \quad \mathfrak{c} \in \mathcal{C}_\lambda^0. \quad (4.19)$$

We can now characterize the status of the trace on $\hat{\mathcal{C}}_\lambda$ in terms of properties of the automorphism that H_∞ generates on \mathcal{B}_∞^0 . The first case to consider is when \hat{p} generates an

⁹ V is not bounded when viewed as an operator acting on $\tilde{\mathcal{H}}$, and is singular even as an unbounded operator since $VV^\dagger = \langle 0_k | e^{-i\hat{y}\hat{p}} e^{i\hat{y}\hat{p}} | 0_k \rangle = \infty$. One interpretation of V is to view it as determining an operator-valued weight T from operators in $\tilde{\mathcal{C}}_\lambda$ to operators in $\hat{\mathcal{C}}_\lambda$ via $T(\tilde{\mathfrak{c}}) = V\tilde{\mathfrak{c}}V^\dagger$ (see section 6 for the definition of operator-valued weights). Equation (4.15) then corresponds to the bimodule property of the operator-valued weight. It is interesting that the BRST procedure described by CLPW [31] has an interpretation in terms of operator-valued weights.

automorphism that preserves the trace on \mathcal{B}_∞ , and the dual automorphism generated by H_∞ on \mathcal{B}_∞^0 is inner. An example of this case would be the algebra defined at \mathcal{I}^- when working with the Unruh state for a black hole, as considered in [32], where the analog of \mathcal{B}_∞^0 is type I_∞ . In this case, \mathcal{B}_∞^0 is itself semifinite and possesses a trace, so modular flow on this algebra is inner. Since H_∞ generates an inner automorphism of \mathcal{B}_∞^0 , by the converse of the cocycle derivative theorem [87, Theorem VIII.3.8], it must be the modular Hamiltonian h_∞ of some weight defined on \mathcal{B}_∞^0 . Pulling back this weight by the conditional expectation E_λ , this leads to a weight on \mathcal{C}_λ^0 whose modular Hamiltonian is $h_{\mathcal{H}}^\lambda + h_\infty$, and therefore $\widehat{\mathcal{C}}_\lambda$ is a modular crossed product of the type III_1 algebra \mathcal{C}_λ^0 , and hence possesses a trace.

The second case is when the automorphism generated by \hat{p} rescales the trace on \mathcal{B}_∞ . An example of this case, considered below, is the Schwarzschild black hole in the Hartle-Hawking state. In this case, we can apply the general structure theorem for type III von Neumann algebras [87, Theorem XII.1.1] to conclude, assuming both \mathcal{B}_∞^0 and \mathcal{B}_∞ are factors, that \mathcal{B}_∞^0 is type III_1 and H_∞ generates modular flow of some weight on \mathcal{B}_∞^0 , whose modular Hamiltonian we denote h_∞ . So once again by composing this weight with the conditional expectation E_λ , we obtain a weight on \mathcal{C}_λ^0 whose modular Hamiltonian is $h_{\mathcal{H}}^\lambda + h_\infty$. Again, the gravitational algebra $\widehat{\mathcal{C}}_\lambda$ is a modular crossed product which possesses a trace.

Finally, if \hat{p} preserves the trace on \mathcal{B}_∞ , but H_∞ generates an outer automorphism on \mathcal{B}_∞^0 , then it is possible to show that \mathcal{B}_∞^0 is already semifinite. However, the outerness of the automorphism generated by H_∞ shows that it is not a modular Hamiltonian for \mathcal{B}_∞^0 , in which case neither is $h_{\mathcal{H}}^\lambda + H_\infty$ for \mathcal{C}_λ^0 . This is enough to conclude that $\widehat{\mathcal{C}}_\lambda$ is not semifinite. This case might arise if H_∞ were a different symmetry generator of \mathcal{B}_∞^0 , such as rotation. In such situations, there are additional gravitational charges that should be added such as the ADM angular momentum, and by incorporating the associated constraints one may be able to argue that the final gravitational algebra is semifinite. An example where these additional constraints are important is for rotating black holes, such as the Kerr example considered in [32].

We conclude this section by applying the general construction to the example of a Schwarzschild black hole. In this case, there is a natural vacuum state $|\Omega\rangle$ provided by the Hartle-Hawking state [94]. The horizon cut algebra \mathcal{C}_λ^0 consists of all quantum fields to the exterior of the cut $v = \lambda(y^A)$, and possesses a subalgebra \mathcal{B}_∞^0 of quantum fields that propagate out through \mathcal{I}^+ . Since $|\Omega\rangle$ is a vacuum state for the horizon average null energy P_λ which generates a half-sided translation and \mathcal{B}_∞^0 is the translation-invariant algebra, the modular Hamiltonian for $|\Omega\rangle$ splits into a sum of two pieces, $h_\Omega^\lambda = h_{\mathcal{H}}^\lambda + h_\infty$, where $h_{\mathcal{H}}^\lambda$ is given, as usual, as an integral of the stress tensor over the horizon (3.11). The component h_∞ affiliated with the algebra at infinity is independent of the choice of horizon cut, and therefore can be identified by taking the cut to lie on the bifurcation surface, $\lambda = 0$. In this case, the modular Hamiltonian is the generator of the global time-translation symmetry associated with the Killing vector ξ^a . In general, this generator will be given by an integral over a complete Cauchy surface Σ in the global spacetime,

$$h_\Omega = -\frac{2\pi}{\kappa} \int_\Sigma T^a_b \xi^b \epsilon_{a\dots}, \quad (4.20)$$

and it is independent of the choice of Cauchy surface since it is the generator of a symmetry. Choosing the Cauchy surface to lie on the horizon and asymptotic infinity $\Sigma = \mathcal{H} \cup \mathcal{I}^+ \cup \mathcal{I}_L^-$ (and assuming massless fields so that contributions from i^+, i_L^- can be neglected), we see that h_∞ is a local integral of the stress tensor over just \mathcal{I}^+ and \mathcal{I}_L^- ,

$$h_\infty = -\frac{2\pi}{\kappa} \int_{\mathcal{I}^+ \cup \mathcal{I}_L^-} T^a_b \xi^b \epsilon_{a\dots} \quad (4.21)$$

This reflects the fact that the state of the fields at \mathcal{I}^+ in the Hartle-Hawking state is thermal at the Hawking temperature $T_H = \frac{\kappa}{2\pi}$. This modular Hamiltonian includes a contribution from gravitons, which is quadratic in the Bondi news tensor at $\mathcal{I}^+, \mathcal{I}_L^-$ [95–97].

To carry out the first step in the construction of the gravitational algebra, we must add the appropriate asymptotic charges to \mathcal{B}_∞^0 and impose constraints to end up with a semifinite algebra. The asymptotic charge is given by the difference between the ADM mass M_{ADM} defined at i^0 and the late time Bondi mass M_+ defined at i^+ . The constraint that relates these operators to the quantum fields is the Bondi mass loss formula [95–101],

$$\mathfrak{C}_{\mathcal{I}^+} = K_\infty + \frac{2\pi}{\kappa} (M_+ - M_{\text{ADM}}) = 0, \quad (4.22)$$

where K_∞ involves only the component of h_∞ affiliated with \mathcal{I}^+ ,

$$K_\infty = -\frac{2\pi}{\kappa} \int_{\mathcal{I}^+} T^a_b \xi^b \epsilon_{a\dots} \quad (4.23)$$

There is a similar constraint involving operators in the complementary algebra at infinity at \mathcal{I}_L^- , the left ADM mass M_{ADM}^L , and the left early time Bondi mass M_-^L at i_L^- , which reads

$$\mathfrak{C}_{\mathcal{I}_L^-} = -K_{-\infty'} - \frac{2\pi}{\kappa} (M_-^L - M_{\text{ADM}}^L) = 0, \quad K_{-\infty'} = \frac{2\pi}{\kappa} \int_{\mathcal{I}_L^-} T^a_b \xi^b \epsilon_{a\dots} \quad (4.24)$$

When imposing these constraints on the algebra, it is necessary to consider the sum of these two constraints $\mathfrak{C}_\infty = \mathfrak{C}_{\mathcal{I}^+} + \mathfrak{C}_{\mathcal{I}_L^-}$, due to the appearance of K_∞ and $K_{-\infty'}$ in the individual constraints (4.22), (4.24). Since \mathcal{B}_∞^0 is an algebra of quantum fields in a thermal state, the one-sided Hamiltonians $K_\infty, K_{-\infty'}$ are only defined as sesquilinear forms due to infinite thermal fluctuations. However, the combination appearing \mathfrak{C}_∞ is $K_\infty - K_{-\infty'} = h_\infty$, which is a well-defined operator generating a strongly continuous flow and therefore amenable to constructing a crossed product algebra. Carrying out the general procedure outlined in sections 2 and 3.3 and references [31, 33] for imposing the constraint, the resulting semifinite algebra \mathcal{B}_∞ is the crossed product

$$\mathcal{B}_\infty = \langle e^{i\hat{p}h_\infty} \mathbf{b} e^{-i\hat{p}h_\infty}, \hat{x} \rangle, \quad \mathbf{b} \in \mathcal{B}_\infty^0, \quad \hat{x} = \frac{2\pi}{\kappa} (M_+ - M_{\text{ADM}}). \quad (4.25)$$

Since h_∞ generates modular flow on \mathcal{B}_∞^0 , this algebra is a modular crossed product and therefore possesses a trace. The algebra \mathcal{C}_λ that includes \mathcal{B}_∞ as a subalgebra is defined exactly as in equation (4.17), which is a type III₁ algebra admitting a half-sided translation with \mathcal{B}_∞ as the translation-invariant subalgebra.

We next impose the horizon constraint by including the asymptotic area operator

$$\hat{y} = -\frac{A_\infty}{4G_N}, \quad (4.26)$$

and forming the crossed product of \mathcal{C}_λ by the horizon modular flow generator $h_{\mathcal{H}}^\lambda$, just as in section 3.3. This leads to the double crossed product algebra $\tilde{\mathcal{C}}_\lambda$ defined in equation (4.18). We then impose the matching constraint (4.12) by determining the subalgebra that commutes with $\hat{k} - \hat{p}$. The only asymptotic charge that survives this constraint is the sum

$$\hat{x} + \hat{y} = -\frac{2\pi}{\kappa} M_{\text{ADM}} + \frac{2\pi}{\kappa} M_+ - \frac{A_\infty}{4G_N}. \quad (4.27)$$

Furthermore, since we have assumed only massless fields, there is no stress energy flux through i^+ , which implies a further condition on the gravitational charges [32],

$$\frac{A_\infty}{4G_N} = \frac{2\pi}{\kappa} M_+, \quad (4.28)$$

so that these terms cancel in (4.27). Therefore, the only asymptotic charge in the final gravitational algebra is the ADM mass. After representing this algebra on a Hilbert space of coinvariants, which we can take to be $\mathcal{H}_{\text{QFT}} \otimes L^2(\mathbb{R})$ as described above, the resulting algebra is

$$\widehat{\mathcal{C}}_\lambda = \left\langle e^{i\hat{p}(h_{\mathcal{H}}^\lambda + h_\infty)} \mathbf{c} e^{-i\hat{p}(h_{\mathcal{H}}^\lambda + h_\infty)}, \hat{x} \right\rangle, \quad \mathbf{c} \in \mathcal{C}_\lambda^0, \quad \hat{x} = -\frac{2\pi}{\kappa} M_{\text{ADM}}. \quad (4.29)$$

One confirms that this final algebra possesses a trace because it is simply the modular crossed product of \mathcal{C}_λ^0 , since $h_{\mathcal{H}}^\lambda + h_\infty$ is the horizon-cut modular Hamiltonian of the Hartle-Hawking state.

The generalized second law for these horizon cut algebras now follows in a similar manner as in section 3.3. We consider semiclassical states $|\widehat{\Phi}\rangle = |\Phi, f\rangle$ with the wavefunction for \hat{x} slowly varying, $\frac{f'}{f} \sim \mathcal{O}(\varepsilon)$. Applying formula (2.14), the semiclassical expansion for the entropy of the algebra $\widehat{\mathcal{C}}_\lambda$ in this state is given by

$$S(\rho_{\widehat{\Phi}}^\lambda) = -S_{\text{rel}}^{\mathcal{C}_\lambda^0}(\Phi||\Omega) - \langle \hat{x} \rangle_{\widehat{\Phi}} + S_f^{M_{\text{ADM}}} + \mathcal{O}(\varepsilon^2). \quad (4.30)$$

This can again be converted into an expression involving the generalized entropy of the horizon cut by writing the relative entropy as the modular free energy $S_{\text{rel}}^{\mathcal{C}_\lambda^0}(\Phi||\Omega) = \langle K_\infty + K_{\mathcal{H}}^\lambda \rangle_\Phi - S_{\Phi, \lambda}^{\text{QFT}}$ and applying the constraints (4.22), (4.28), and (3.32). Since the same formula applies for the entropy of the algebra for a later cut $\tilde{\lambda}$ in the same state $|\widehat{\Phi}\rangle$, and since $\langle \hat{x} \rangle_{\widehat{\Phi}}$ and $S_f^{M_{\text{ADM}}}$ remain unchanged, the entropy difference between the two cuts reduces to a difference of relative entropies,

$$S(\rho_{\tilde{\lambda}}) - S(\rho_\lambda) = -S_{\text{rel}}^{\mathcal{C}_{\tilde{\lambda}}^0}(\Phi||\Omega) + S_{\text{rel}}^{\mathcal{C}_\lambda^0}(\Phi||\Omega) \geq 0, \quad (4.31)$$

up to $\mathcal{O}(\varepsilon^2)$ corrections.

Interestingly, a stronger statement of the generalized second law can be proven for overall differences in entropy between the horizon cut λ and the entropy at very late times. As discussed above, $\widehat{\mathcal{C}}_\lambda$ is a crossed product of a quantum field algebra \mathcal{C}_λ which admits a half-sided translation. The state in which the crossed product is defined is fixed by the conditional expectation E_λ that sends \mathcal{C}_λ to the translation-invariant algebra \mathcal{B}_∞^0 . Modular flow in this state then maps \mathcal{B}_∞^0 to itself, and hence there is a subalgebra of $\widehat{\mathcal{C}}_\lambda$ isomorphic to the crossed product of \mathcal{B}_∞^0 ,

$$\mathcal{B}_\infty = \langle e^{i\hat{p}h_\infty} \mathbf{b} e^{-i\hat{p}h_\infty}, \hat{x} \rangle \subset \widehat{\mathcal{C}}_\lambda. \quad (4.32)$$

Because modular flow commutes with the conditional expectation E_λ , the gravitational algebra $\widehat{\mathcal{C}}_\lambda$ admits a conditional expectation to \mathcal{B}_∞ , denoted \widehat{E}_λ , which acts on operators as

$$\widehat{E}_\lambda \left(e^{i\hat{p}(h_{\mathcal{H}}^\lambda + h_\infty)} \mathbf{c} e^{-i\hat{p}(h_{\mathcal{H}}^\lambda + h_\infty)} e^{iu\hat{x}} \right) = e^{i\hat{p}h_\infty} E_\lambda(\mathbf{c}) e^{-i\hat{p}h_\infty} e^{iu\hat{x}}. \quad (4.33)$$

The formula for the trace on either algebra $\widehat{\mathcal{C}}_\lambda$ or \mathcal{B}_∞ is the same, given by

$$\text{Tr}(\cdot) = 2\pi \langle \Omega, 0_p | e^{-\hat{x}}(\cdot) | \Omega, 0_p \rangle, \quad (4.34)$$

and hence $\text{Tr}_\infty = \text{Tr}_{\lambda|\infty}$. Because the inclusion $\mathcal{B}_\infty \subset \widehat{\mathcal{C}}_\lambda$ is trace-preserving, we are in a position to apply the results of [102], which imply that the entropy on \mathcal{B}_∞ is larger than the entropy on \mathcal{C}_λ for *any* state $|\widehat{\Phi}\rangle$, with no semiclassical assumption on the wavefunction. Hence we arrive at the global second law,

$$S(\rho_{\widehat{\Phi}}^\infty) \geq S(\rho_{\widehat{\Phi}}^\lambda). \quad (4.35)$$

This is similar to the global second law discussed in [30], except that it now involves a large algebra at infinity \mathcal{B}_∞ that consists of more than just the asymptotic charges. We also note that the conditional expectation \widehat{E}_λ preserves the trace $\text{Tr}_\lambda = \text{Tr}_\infty \circ \widehat{E}_\lambda$, but we expect that the index of this inclusion is infinite. Hence, the upper bound on the entropy increase derived in [102] is simply infinite, and does not yield an interesting constraint.

As a final comment, we mention that a double crossed product algebra similar to $\widetilde{\mathcal{C}}_\lambda$ defined in (4.18) appears in the construction of the algebra for Schwarzschild-de Sitter by KFLS [32]. In this construction, there is an additional degree of freedom included in the algebra that describes an observer eternally restricted to the regions between the black hole and cosmological horizons. The observer's energy therefore contributes to the flux through i^- so that the analog of the matching condition (4.28) is modified by the observer's Hamiltonian, which is then related to the sum of the two early time horizon area fluctuations. The presence of such an observer also allows the matching condition (4.12) for the two horizon times to not be imposed, so that the final double crossed product algebra is the correct description in this case. However, it is also natural to ask whether the observer degree of freedom can be eliminated in this setup. This is done precisely by setting the observer Hamiltonian to zero, and imposing the matching constraint (4.12). The resulting algebra is a single crossed product, and semifiniteness follows from the arguments discussed above. In this description, the early time black hole horizon area replaces the observer Hamiltonian as the asymptotic gravitational charge, so that the black hole itself plays the role of the observer in de Sitter space. This setup is quite analogous to the model of an observer in holography as a small black hole, considered in [103, 104].

4.3 Rotating black holes and Unruh states

The discussion of section 4.2 aims to be sufficiently general so as to provide a construction of semifinite gravitational algebras for generic black holes with a variety of asymptotics and angular velocities. However, a number of subtleties arise when carrying out the details of the construction for more complicated cases than the Schwarzschild black hole. In this section, we describe two such subtleties in applying the procedure in the physically relevant contexts of rotating black holes and black holes formed from collapse. We conjecture how these subtleties should be resolved, but do not perform an in-depth analysis, which we leave as an interesting topic for future investigation.

For rotating asymptotically flat black holes such as the Kerr solution or Myers-Perry black holes in $d \geq 5$ [105], the main issue is that the horizon-generating Killing vector ξ^a is not timelike throughout the black hole exterior. In general, the Killing vector has the decomposition [106]

$$\xi^a = t^a + \sum_i \Omega_{\mathcal{H}}^i \phi_i^a, \quad (4.36)$$

where t^a is the Killing vector associated with asymptotic time translations, ϕ_i^a are a collection of rotational Killing vectors, and $\Omega_{\mathcal{H}}^i$ are the horizon rotational velocities. Since the norms of ϕ_i^a grow at large radius while the norm of t^a remains bounded, the vector ξ^a eventually becomes spacelike as long as there is at least one nonzero $\Omega_{\mathcal{H}}^i$. There are therefore operators localized in the region that ξ^a is spacelike that carry negative Killing energy. The existence of negative energy states created by these operators is closely tied to the phenomenon of superradiance for rotating black holes [107, 108].

Negative Killing energy states pose a problem for constructing a thermal state, since a thermal density matrix $\rho \propto e^{-\beta H_\xi}$ is not normalizable when the spectrum of H_ξ is not bounded below. This appears to preclude the existence of a global vacuum state $|\Omega\rangle$ that satisfies a KMS condition with respect to the Killing flow, meaning there is no analog of the Hartle-Hawking state for the Kerr black hole whose modular Hamiltonian globally generates a geometric flow [109]. This does not pose an immediate issue to the semiclassical generalized second law discussed in section 4.1. There, we only required that there is a vacuum state $|\Omega\rangle$ for the horizon average null energy operators P_λ , since any such state has a modular Hamiltonian whose horizon component $h_{\mathcal{H}}^\lambda$ is given by an integral of the stress tensor (3.11). The behavior of the modular flow of $|\Omega\rangle$ on the algebra at infinity \mathcal{B}_∞ does not affect the derivation of the generalized second law in this context.

The actual problem arises when attempting to construct a semifinite gravitational algebra following the procedure of section 4.2. In this case, we would like to argue that imposing the gravitational constraints on the algebra at infinity results in an algebra with a trace; however, this is tantamount to assuming that there is a state ω_∞ on \mathcal{B}_∞ whose modular flow generates time translations along the vector t^a at \mathcal{I}^+ . One possibility is that such a state does exist, and one would further need to argue that it is thermal at the Hawking temperature. Assuming this is possible, this state could be extended to a state on any of the horizon cut algebras by composing it with the conditional expectation E_λ , and then $\omega_\lambda = \omega_\infty \circ E_\lambda$ would be a horizon vacuum state. For rotating black holes, the modular

flow for this state would be geometric only for operators on the horizon and on \mathcal{I}^+ , since it flows along different Killing vectors on these two surfaces (respectively, ξ^a and t^a). Assuming that one can argue that this state is Hadamard throughout the two-sided Kerr geometry, this would provide the necessary state to construct a semifinite gravitational algebra. One would also need to consider modified matching conditions for the gravitational charges and incorporate additional crossed products for rotations, along the lines of KFLS [32], due to the fact that the constraints are associated with different Killing vectors and therefore different asymptotic charges.

Another possibility is to try to work directly with the analog of the Hartle-Hawking state, which in the case of Kerr is given by the Frolov-Thorne vacuum, which has the appropriate thermal behavior for operators near the horizon where ξ^a is timelike [110]. However, this vacuum cannot lead to a well-defined state everywhere in the Kerr exterior, due to issues related to superradiance discussed above. This manifests in a failure of the Hadamard property for correlation functions in the region where ξ^a is spacelike. If one were to treat the Frolov-Thorne vacuum as a normalizable state, it would produce a unitarily inequivalent GNS representation of the local field algebra relative to a state that was Hadamard everywhere in the Kerr exterior. Such a representation would be irrelevant for investigations of the generalized second law for the Kerr black hole. However, it is possible that by reinterpreting the Frolov-Thorne vacuum as a nonnormalizable weight as opposed to a state, it could provide the appropriate modular operator to construct a trace on the algebra at infinity.

Recall that a weight on a von Neumann algebra is a positive linear function from positive operators in the algebra to the extended positive reals $\widehat{\mathbb{R}} = \mathbb{R}^+ \cup \{+\infty\}$ [87, Chapter VII]. The weight is said to be semifinite if the weak closure of the algebra generated by the operators with finite expectation value in the weight gives back the full von Neumann algebra. Since the Frolov-Thorne vacuum assigns finite expectation values to operators in the region where ξ^a is timelike, we can interpret it as a weight on the algebra. Furthermore, the causal completion of this region in spacetime yields back the entire exterior wedge of the Kerr black hole, so it is plausible that the subalgebra of operators with finite expectation value in the Frolov-Thorne vacuum is weakly dense in the full von Neumann algebra of the black hole exterior. If this is the case, the Frolov-Thorne vacuum would define a semifinite weight on this algebra. The modular operator for this weight would presumably generate the Killing flow of ξ^a , and hence by implementing the crossed product of this flow on the algebra at infinity, one would obtain an algebra with a trace. While an intriguing proposal, there are many details to check in order to establish the proposed properties of the Frolov-Thorne vacuum as a semifinite weight. A few of these issues are discussed in section 7.

A second set of questions arises when applying the arguments for the generalized second law to black holes formed from collapse, whose natural vacuum state is the Unruh state [111, 112] as opposed to the Hartle-Hawking state. The Unruh vacuum is defined by initial data on the past black hole horizon and \mathcal{I}^- , and imposes that there is no incoming radiation from past infinity. By contrast, the Hartle-Hawking state has incoming thermal radiation from \mathcal{I}^- in addition to outgoing Hawking radiation at \mathcal{I}^+ , which is an unnatural idealization for astrophysical black holes; such a boundary condition is more appropriate for an eternal black hole. The question therefore arises to what extent the Unruh state describing an evaporating

black hole is covered by the analysis in the present paper which relied on properties of the Hartle-Hawking vacuum.

At first glance it appears that the Unruh state can just be considered a specific excited state for the algebra of quantum fields exterior to a cut of the horizon. It would then describe a state with some amount of entanglement between modes that enter the horizon and modes in \mathcal{B}_∞ that escape through \mathcal{I}^+ , and therefore satisfies a generalized second law according to the discussion of section 4.1. The main issue with this argument is that the Unruh vacuum differs significantly from the Hartle-Hawking vacuum in the infrared, so that in standard treatments these states result in unitarily inequivalent representations of the field algebras in the black hole exterior. This results in a failure in the Hadamard property at the past black hole horizon in the maximally extended spacetime, leading to a divergence in the average null energy operators P_λ coming from contributions near the bifurcation surface.

As in the discussion of the Frolov-Thorne vacuum, the resolution here may involve reinterpreting the states under consideration instead as weights on a single representation of the field algebra. If we take the perspective that the Unruh vacuum is the appropriate state for black holes formed from collapse, this would then require interpreting the Hartle-Hawking vacuum as a weight on the von Neumann algebra constructed from the GNS representation of the field algebra in the Unruh state. Fortunately, the structure theorems of half-sided translations and modular inclusions generalize to the case of normal semifinite weights [113], so assuming the Hartle-Hawking vacuum still defines a weight fixed by the translation, these theorems would continue to allow the derivation of the modular Hamiltonian as a stress tensor integral as in sections 3.1 and 4.1. One possible subtlety in applying the general discussion of section 4.1 is that when the translation-invariant weight is not normalizable, there may no longer be a conditional expectation to the translation-invariant algebra at infinity. A natural conjecture generalizing the results of [85] is that instead, one would find an operator-valued weight to the algebra at infinity. This would likely still allow the discussion of section 4.1 to go through, and would still allow the construction of a gravitational algebra following the procedure of section 4.2. Verifying the details of these conjectures is an important direction for future work.

5 Perturbative corrections to crossed product entropy

In section 3.3, we found that the crossed product entropies satisfy a generalized second law to leading order in the semiclassical expansion for states of the form $|\widehat{\Phi}\rangle = |\Phi, f\rangle$. This raises the question of whether these entropies continue to be monotonic after accounting for corrections in the semiclassical expansion. In this section we begin to address this question. We derive the exact expression for the correction to the semiclassical entropy in the crossed product algebra, and work out some of its main properties. One of these properties is that the correction is strictly nonnegative, which follows from the interpretation of the classical-quantum state as a Petz recovery of the quantum field state $|\Phi\rangle$. We can also obtain an upper bound on the entropy correction using the explicit expression for the entropy correction. Although we are not able to use these bounds to conclude anything about the corrected

generalized second law, which relies on differences in entropy corrections, we note that these bounds were recently employed in [114] to obtain rigorous proofs of the Bekenstein bound and quantum null energy condition in terms of vacuum-subtracted entropies. These bounds indicate that it is difficult to violate the generalized second law with semiclassical corrections, although it remains possible that such a violation could occur over short time scales in regimes where the quantum field relative entropy is nearly constant. We leave open the question of whether the GSL holds at all times for the crossed product entropies, but note that if it did hold, it would represent a novel data-processing inequality bounding the decrease of relative entropy for quantum fields. It would therefore be interesting to connect this proposed bound to the improved data-processing inequalities employed in universal recovery channels [115–117].

We begin by explaining the interpretation of the state $|\Phi, f\rangle$ as a Petz-recovered state, which leads to the lower bound on the entropy correction. Given a state φ for a von Neumann algebra \mathcal{A} , we denote by $|\varphi^{\frac{1}{2}}\rangle$ its canonical purification in the natural cone defined by a vacuum state $|\Omega\rangle$ in a standard representation of \mathcal{A} . We consider a unital, completely positive (UCP) map $\alpha : \mathcal{B} \rightarrow \mathcal{A}$ between von Neumann algebras, and look to construct the associated Petz recovery map associated with a cyclic-separating state φ on \mathcal{A} . We first define the φ -dual map α' , which is a UCP map on the commutants $\alpha'_\varphi : \mathcal{A}' \rightarrow \mathcal{B}'$ defined by the relation

$$\left\langle (\varphi \circ \alpha)^{\frac{1}{2}} \left| \alpha'_\varphi(\mathbf{a}') \mathbf{b} \right| (\varphi \circ \alpha)^{\frac{1}{2}} \right\rangle = \left\langle \varphi^{\frac{1}{2}} \left| \mathbf{a}' \alpha(\mathbf{b}) \right| \varphi^{\frac{1}{2}} \right\rangle, \quad \mathbf{a}' \in \mathcal{A}', \mathbf{b} \in \mathcal{B}. \quad (5.1)$$

By combining α' with the modular conjugations $J_{\mathcal{A}}, J_{\mathcal{B}}$, we obtain a new UCP map $\alpha_\varphi^P : \mathcal{A} \rightarrow \mathcal{B}$ called the *Petz map* [81, 118, 119], defined by

$$\alpha_\varphi^P(\mathbf{a}) = J_{\mathcal{B}} \alpha'_\varphi(J_{\mathcal{A}} \mathbf{a} J_{\mathcal{A}}) J_{\mathcal{B}}. \quad (5.2)$$

We now apply this to the crossed product algebra

$$\widehat{\mathcal{A}} = \langle e^{i\hat{p}h_\Omega} \mathbf{a} e^{-i\hat{p}h_\Omega}, \hat{q} \rangle, \quad (5.3)$$

where $\mathbf{a} \in \mathcal{A}$ are operators in a QFT algebra of interest, and $|\Omega\rangle$ is a cyclic-separating vacuum state whose modular Hamiltonian h_Ω defines the crossed product. The inclusion map

$$\iota(\mathbf{a}) = e^{i\hat{p}h_\Omega} \mathbf{a} e^{-i\hat{p}h_\Omega} \quad (5.4)$$

defines a UCP map from the QFT subalgebra \mathcal{A} into the crossed product $\widehat{\mathcal{A}}$, and hence we would like to construct its Petz map. We consider the vacuum state $|\widehat{\Omega}\rangle = |\Omega, f\rangle$ with wavefunction f chosen to be real and positive. This ensures that $|\widehat{\Omega}\rangle$ is cyclic-separating for $\widehat{\mathcal{A}}$, and reality of f implies that $|\widehat{\Omega}\rangle$ is in the canonical cone of the tracial weight $\text{Tr} = 2\pi \langle \Omega, 0_p | e^{-\hat{q}} \cdot | \Omega, 0_p \rangle$, and hence has the modular conjugation (see equation (2.8))

$$J_{\widehat{\mathcal{A}}} = J_\Omega e^{-i\hat{p}h_\Omega}. \quad (5.5)$$

States of this form pull back to the vacuum state $|\Omega\rangle$ under the action of the dual channel ι^* , meaning that $(\omega_{\widehat{\Omega}} \circ \iota)(\mathbf{a}) = \langle \widehat{\Omega} | \iota(\mathbf{a}) | \widehat{\Omega} \rangle = \langle \Omega | \mathbf{a} | \Omega \rangle$.

We denote $\omega_{\widehat{\Omega}}$ -dual map by $\iota'_f : \widehat{\mathcal{A}}' \rightarrow \mathcal{A}'$, which acts as

$$\langle \Omega | \iota'_f(\widehat{\mathbf{a}}') \mathbf{a} | \Omega \rangle = \langle \Omega, f | \widehat{\mathbf{a}}' \iota(\mathbf{a}) | \Omega, f \rangle = \langle \Omega, f | \widehat{\mathbf{a}}' e^{i\hat{p}h\Omega} \mathbf{a} e^{-i\hat{p}h\Omega} | \Omega, f \rangle = \langle \Omega | \langle f | e^{-i\hat{p}h\Omega} \widehat{\mathbf{a}}' e^{i\hat{p}h\Omega} | f \rangle \mathbf{a} | \Omega \rangle. \quad (5.6)$$

This determines the expression for ι'_f to be

$$\iota'_f(\widehat{\mathbf{a}}') = \langle f | e^{-i\hat{p}h\Omega} \widehat{\mathbf{a}}' e^{i\hat{p}h\Omega} | f \rangle. \quad (5.7)$$

Combining this with the expression (5.5) for the modular conjugation, we find that the Petz map $\iota_f^P : \widehat{\mathcal{A}} \rightarrow \mathcal{A}$ is given by

$$\iota_f^P(\widehat{\mathbf{a}}) = J_\Omega \langle f | e^{-i\hat{p}h\Omega} J_\Omega e^{-i\hat{p}h\Omega} \widehat{\mathbf{a}} e^{i\hat{p}h\Omega} J_\Omega e^{i\hat{p}h\Omega} | f \rangle J_\Omega = \langle f | \widehat{\mathbf{a}} | f \rangle, \quad (5.8)$$

where in the last equality we used that f is real valued to pass the J_Ω past $\langle f |$ and $| f \rangle$. Hence, we find that taking the partial expectation value with respect to the wavefunction f defines a UCP map from $\widehat{\mathcal{A}}$ to \mathcal{A} that serves as a Petz map dual to the inclusion map (5.4). The quantum channel $(\iota_f^P)^*$ maps a state φ on \mathcal{A} to a state $(\iota_f^P)^* \varphi = \varphi \circ \iota_f^P$ on the crossed product $\widehat{\mathcal{A}}$. If φ arises from the vector state $|\Phi\rangle$, it follows immediately from (5.8) that $|\widehat{\Phi}\rangle = |\Phi, f\rangle$ induces the state $\varphi \circ \iota_f^P$ on $\widehat{\mathcal{A}}$. Hence, we find that the classical-quantum states which were the focus of our investigations into the GSL have the interpretation of a Petz recovery of a state $|\Phi\rangle$ on the quantum field theory subalgebra \mathcal{A} .

Because $|\widehat{\Phi}\rangle$ and $|\widehat{\Omega}\rangle$ are images of the states $|\Phi\rangle$ and $|\Omega\rangle$ under the Petz recovery channel $(\iota_f^P)^*$, the relative entropy computed in $\widehat{\mathcal{A}}$ of these states must be less than the corresponding relative entropy computed in \mathcal{A} , which follows from Uhlmann's general monotonicity theorem for relative entropy under the action of a quantum channel [80, 81]. This monotonicity result places a lower bound on the corrections to the semiclassical expression of the entropy for the state $|\widehat{\Phi}\rangle$. To derive the bound, recall from equation (2.8) that the density matrix for the state $|\widehat{\Omega}\rangle$ is given by

$$\rho_{\widehat{\Omega}} = g(\hat{q}) e^{\hat{q}}, \quad (5.9)$$

where $g(\hat{q}) = |f(\hat{q})|^2$. We can then write the relative entropy in $\widehat{\mathcal{A}}$ as

$$\begin{aligned} S_{\text{rel}}^{\widehat{\mathcal{A}}}(\widehat{\Phi} || \widehat{\Omega}) &= \langle \widehat{\Phi} | \log \rho_{\widehat{\Phi}} - \log \rho_{\widehat{\Omega}} | \widehat{\Phi} \rangle = -S(\rho_{\widehat{\Phi}}) - \langle f | \log g(\hat{q}) + \hat{q} | f \rangle \\ &= S_{\text{rel}}^{\mathcal{A}}(\Phi || \Omega) - \Delta S(\rho_{\widehat{\Phi}}), \end{aligned} \quad (5.10)$$

where in the last equality we have used the semiclassical expansion of the entropy $S(\rho_{\widehat{\Phi}}) = -S_{\text{rel}}^{\mathcal{A}}(\Phi || \Omega) - \langle f | \log g(\hat{q}) + \hat{q} | f \rangle + \Delta S(\rho_{\widehat{\Phi}})$ which follows from equation (2.14), with ΔS denoting the subleading corrections. Monotonicity of relative entropy under the channel $(\iota_f^P)^*$ requires that $S_{\text{rel}}^{\mathcal{A}}(\Phi || \Omega) \geq S_{\text{rel}}^{\widehat{\mathcal{A}}}(\widehat{\Phi} || \widehat{\Omega})$, which, together with (5.10), implies

$$\Delta S(\rho_{\widehat{\Phi}}) \geq 0. \quad (5.11)$$

This is our first key result: the correction to the semiclassical expression for the crossed product entropy for the state $|\widehat{\Phi}\rangle$ is strictly nonnegative.

We now proceed to derive an explicit expression for the entropy correction $\Delta S(\rho_{\hat{\Phi}})$. The starting point is the expression (2.12) for the logarithm of the density matrix $\rho_{\hat{\Phi}}$ for the classical-quantum state $|\hat{\Phi}\rangle = |\Phi, f\rangle$. This expression can be written

$$-\log \rho_{\hat{\Phi}} = -h - \hat{q} - \log X; \quad X := f(\hat{q})e^{i\hat{p}h}\Delta e^{-i\hat{p}h}f^*(\hat{q}), \quad (5.12)$$

where we have defined

$$h = h_{\Omega|\Phi}, \quad \Delta = \Delta_{\Phi}, \quad (5.13)$$

and now allow the wavefunction f to be complex.¹⁰ The entropy is obtained by taking the expectation value of $-\log \rho_{\hat{\Phi}}$ in the state $|\Phi, f\rangle$. The leading semiclassical contribution to the entropy comes from

$$S_0(\rho_{\hat{\Phi}}) = \langle \Phi, f | -h - \log \Delta - \hat{q} - \log g(\hat{q}) | \Phi, f \rangle, \quad (5.14)$$

which reproduces the expression (2.14), noting that $\langle \Phi | \log \Delta | \Phi \rangle = 0$. The correction to the entropy is therefore given by

$$\Delta S(\rho_{\hat{\Phi}}) = \langle \Phi, f | -\log X + \log(g(\hat{q})\Delta) | \Phi, f \rangle \quad (5.15)$$

To evaluate this, we use the integral representation of the operator logarithm

$$-\log A = \int_0^\infty d\lambda \left(\frac{1}{\lambda + A} - \frac{1}{\lambda + 1} \right) \quad (5.16)$$

to write

$$\begin{aligned} -\log X + \log(g(\hat{q})\Delta) &= \int_0^\infty d\lambda \left(\frac{1}{\lambda + X} - \frac{1}{\lambda + g(\hat{q})\Delta} \right) \\ &= \int_0^\infty d\lambda \frac{1}{f^*(\hat{q})} \left(e^{i\hat{p}h} \frac{1}{\frac{\lambda}{g(\hat{q}-h)} + \Delta} e^{-i\hat{p}h} - \frac{1}{\frac{\lambda}{g(\hat{q})} + \Delta} \right) \frac{1}{f(\hat{q})} \end{aligned} \quad (5.17)$$

Since we will eventually take the expectation value of this expression in the state $|\Phi, f\rangle$, it is convenient to pass this operator through the UCP map $\langle f | \cdot | f \rangle$, which results in an operator acting on the QFT Hilbert space whose expectation value in the state $|\Phi\rangle$ yields the entropy correction. This simplifies the expression due to the identity

$$\frac{1}{f(\hat{q})} | f \rangle = \int dz | z \rangle = \int ds dz \frac{e^{-isz}}{\sqrt{2\pi}} | s \rangle = \sqrt{2\pi} | 0_p \rangle, \quad (5.18)$$

using the normalization of position and momentum eigenstates $\langle s | z \rangle = \frac{e^{-isz}}{\sqrt{2\pi}}$. This then gives

$$\begin{aligned} \langle f | -\log X + \log(g(\hat{q})\Delta) | f \rangle &= 2\pi \langle 0_p | \int_0^\infty d\lambda \left(\frac{1}{\frac{\lambda}{g(\hat{q}-h)} + \Delta} - \frac{1}{\frac{\lambda}{g(\hat{q})} + \Delta} \right) | 0_p \rangle \\ &= \int_0^\infty d\lambda \int dz \left(\frac{1}{\frac{\lambda}{g(z-h)} + \Delta} - \frac{1}{\frac{\lambda}{g(z)} + \Delta} \right). \end{aligned} \quad (5.19)$$

¹⁰In fact, the phase of the wavefunction has no effect on the entropy, since we can write a generic classical-quantum state as $|\Phi, f\rangle = e^{i\phi(\hat{q})} |\Phi, f_r\rangle$, where $f_r(q)$ is real. Since $e^{i\phi(\hat{q})}$ is a unitary operator in $\hat{\mathcal{A}}$, the states $|\Phi, f\rangle$ and $|\Phi, f_r\rangle$ have the same entropy, since the entropy is invariant under the action of unitaries from the algebra.

In order to simplify this further, we would like to switch the order of integration and change variables to $\lambda = g(z)\mu$. This can only be done if the λ integral converges at fixed z , which does not hold in the present form. We therefore manipulate it into a form that does converge at fixed z , making use of the ability to add terms whose z -integral is zero. In particular, we define $\delta(z, h) := g(z - h) - g(z)$, which satisfies $\int dz \delta(z, h) = 0$, and then express (5.19) as

$$\begin{aligned} \langle f | \cdots | f \rangle &= \int_0^\infty d\lambda \int dz \frac{1}{\lambda + \Delta g(z)} \delta(z, h) \frac{\lambda}{\lambda + \Delta g(z - h)} \\ &= \int_0^\infty d\lambda \int dz \left[-\frac{1}{\lambda + \Delta g(z)} \delta(z, h) \Delta g(z - h) \frac{1}{\lambda + \Delta g(z - h)} \right. \\ &\quad \left. + \left(\frac{1}{\lambda + g(z)\Delta} - \frac{1}{\lambda + 1} \right) \delta(z, h) \right]. \end{aligned} \quad (5.20)$$

Here we have made use of the operator identity $A^{-1} + B^{-1} = A^{-1}(A + B)B^{-1}$, and have added the term $-\frac{1}{\lambda+1}\delta(z, h)$ in the last line which z -integrates to zero. It is now clear that the λ integral of each of the two terms in the final expression in (5.20) converges at fixed z , allowing the order of integration to be reversed.

The integral in the second line can be performed explicitly, resulting in

$$\begin{aligned} \int dz \int_0^\infty d\lambda \left(\frac{1}{\lambda + g(z)\Delta} - \frac{1}{\lambda + 1} \right) \delta(z, h) &= - \int dz \log(g(z)\Delta) \delta(z, h) \\ &= - \int dz \log g(z) [g(z - h) - g(z)] \\ &= - \int dz g(z) \log \frac{g(z + h)}{g(z)} =: G(h), \end{aligned} \quad (5.21)$$

where in the first line we have used that $\log(\Delta)\delta(z, h)$ integrates to zero.

We define the first line of (5.20) to be $-R(\Delta, h)$, and simplify its expression by switching the order of integration and substituting $\lambda = g(z)\mu$. After subtracting from the integrand a term $\frac{1}{\mu+\Delta}\delta(z, h)\frac{\Delta}{\mu+\Delta}$ which integrates to zero, this leads to

$$\begin{aligned} R(\Delta, h) &= \int dz \int_0^\infty d\mu \frac{1}{\mu + \Delta} \delta(z, h) \Delta \left(\frac{1}{\frac{\mu g(z)}{g(z-h)} + \Delta} - \frac{1}{\mu + \Delta} \right) \\ &= \int dz g(z) \int_0^\infty d\mu \frac{1}{\mu + \Delta} \frac{\delta(z, h)}{g(z)} \frac{\mu}{\frac{\mu}{\Delta} + \frac{g(z-h)}{g(z)}} \frac{\delta(z, h)}{g(z)} \frac{1}{\mu + \Delta}. \end{aligned} \quad (5.22)$$

Hence, the final expression for the entropy correction is given by

$$\Delta S(\rho_{\widehat{\Phi}}) = \langle \Phi | -R(\Delta, h) + G(h) | \Phi \rangle, \quad (5.23)$$

with $G(h)$ defined by (5.21) and $R(\Delta, h)$ by (5.22).

The first observation to make about this expression is that $R(\Delta, h)$ is a positive operator. This follows from the fact that the term in the middle of the integrand $\frac{1}{\frac{\mu}{\Delta} + \frac{g(z-h)}{g(z)}}$ takes the

form of a parallel sum $\left(\frac{\Delta}{\mu} : \frac{g(z)}{g(z-h)}\right)$. The parallel sum of two positive, invertable operators is defined by $(A : B) = \frac{1}{A^{-1} + B^{-1}}$, and the resulting operator is also positive [120, Chapter 36]. Since both $\frac{\Delta}{\mu}$ and $\frac{g(z)}{g(z-h)}$ are positive for all values of μ and z in the range of integration, the middle operator in (5.22) is manifestly positive. The full integrand then takes the form $\mu g(z) C^\dagger \left(\frac{\Delta}{\mu} : \frac{g(z)}{g(z-h)}\right) C$, where the operator C depends on μ, z, Δ , and h , and hence $R(\Delta, h)$ is the integral of a collection of positive operators with respect to a positive measure. This demonstrates that $R(\Delta, h)$ is positive.

Positivity of $R(\Delta, h)$ then leads to an upper bound on $\Delta S(\rho_{\hat{\Phi}})$ in terms of the expectation value of $G(h)$, as is clear from (5.23). Combining with the bound (5.11), this leads to

$$0 \leq \Delta S(\rho_{\hat{\Phi}}) \leq \langle \Phi | G(h) | \Phi \rangle. \quad (5.24)$$

Hence, the entropy variation is upper bounded by a function of the relative modular operator $h = h_{\Omega|\Phi}$, whose precise form depends on the wavefunction $f(\hat{q})$. These bounds were recently employed in a nongravitational setting in [114] to obtain rigorous statements about quantum field theory entropies regulated by the crossed product.

The next point about the entropy correction is that it has no first order contribution in the semiclassical expansion, so the correction is generically $\mathcal{O}(\varepsilon^2)$ with $\varepsilon \sim \frac{g'}{g}$. The expression (5.22) for $R(\Delta, h)$ is manifestly $\mathcal{O}(\varepsilon^2)$ since $\delta(z, h) = -g'(z)h + \mathcal{O}(\varepsilon^2)$, and two factors of $\delta(z, h)$ appear in $R(\Delta, h)$. The contribution from $G(h)$ is also $\mathcal{O}(\varepsilon^2)$, since the first order contribution integrates to zero,

$$\begin{aligned} G(h) &= - \int dz g(z) \left(h \frac{g'(z)}{g(z)} + \frac{1}{2} h^2 \frac{g''(z)}{g(z)} - \frac{1}{2} h^2 \left(\frac{g'(z)}{g(z)} \right)^2 \right) + \mathcal{O}(\varepsilon^3) \\ &= \frac{1}{2} \int dz \frac{g'(z)^2}{g(z)} h^2 + \mathcal{O}(\varepsilon^3). \end{aligned} \quad (5.25)$$

This demonstrates that the semiclassical expression (2.14) for the crossed product entropy is valid up to ε^2 corrections. In particular, this implies that it is difficult to violate the second law in perturbation theory, since generically the decrease in relative entropy will more than compensate the change in the $\mathcal{O}(\varepsilon^2)$ terms in the entropy.

We can simplify the contribution of $R(\Delta, h)$ to the entropy correction by evaluating this operator in the state $|\Phi\rangle$. Because $\Delta|\Phi\rangle = |\Phi\rangle$, we find that

$$\begin{aligned} \langle \Phi | R(\Delta, h) | \Phi \rangle &= \int dz g(z) \langle \Phi | \frac{\delta(z, h)}{g(z)} \int_0^\infty d\mu \frac{\mu}{(\mu+1)^2} \frac{1}{\frac{\mu}{\Delta} + \frac{g(z-h)}{g(z)}} \frac{\delta(z, h)}{g(z)} | \Phi \rangle \\ &= \int dz g(z) \langle \Phi | \frac{\delta(z, h)}{g(z)} \Delta^{\frac{1}{2}} \int_0^\infty d\mu \frac{\mu}{(\mu+1)^2} \frac{1}{\mu + \Delta^{\frac{1}{2}} \frac{g(z-h)}{g(z)} \Delta^{\frac{1}{2}}} \Delta^{\frac{1}{2}} \frac{\delta(z, h)}{g(z)} | \Phi \rangle \\ &= \int dz g(z) \langle \Phi | \frac{\delta(z, h)}{g(z)} \Delta^{\frac{1}{2}} \left[R_2 \left(\Delta^{\frac{1}{2}} \frac{g(z-h)}{g(z)} \Delta^{\frac{1}{2}} \right) \right] \Delta^{\frac{1}{2}} \frac{\delta(z, h)}{g(z)} | \Phi \rangle, \end{aligned} \quad (5.26)$$

where we have defined the function

$$R_2(x) := \int_0^\infty d\mu \frac{\mu}{(\mu+1)^2(\mu+x)} = \frac{x \log x + 1 - x}{(1-x)^2}. \quad (5.27)$$

The $\mathcal{O}(\varepsilon^2)$ contribution to this expression can be obtained directly by replacing $\delta(z, h)$ with $-g'(z)h$ and setting $\frac{g(z-h)}{g(z)}$ to 1 inside the argument of R_2 . Combining the resulting expression with (5.25) according to (5.23), we arrive at the expression for the $\mathcal{O}(\varepsilon^2)$ contribution to the entropy

$$S^{(2)}(\rho_{\widehat{\Phi}}) = \int dz \frac{g'(z)^2}{g(z)} \langle \Phi | h \left(\frac{1}{2} - \Delta R_2(\Delta) \right) h | \Phi \rangle. \quad (5.28)$$

The final expression simplifies such that the only dependence on the wavefunction $g(z)$ is through the prefactor

$$k_g := \int dz \frac{g'(z)^2}{g(z)}, \quad (5.29)$$

and the rest of the expression involves an operator constructed purely from the quantum field theory state, involving the relative modular Hamiltonian $h_{\Omega|\Phi}$ and the modular operator Δ_Φ .

This expression allows us to verify the bounds (5.24) at $\mathcal{O}(\varepsilon^2)$. First, because $\Delta R_2(\Delta)$ is a positive operator, we immediately find that $S^{(2)}(\rho_{\widehat{\Phi}}) \leq \frac{1}{2} k_g \langle \Phi | h^2 | \Phi \rangle$. To verify the lower bound, we first note that the operator $(1 - \Delta)$ has vanishing expectation value in the state $h|\Phi\rangle$. This follows from the fact that because $h_\Phi|\Phi\rangle = 0$, we can instead write this state as $(h_{\Omega|\Phi} - h_\Phi)|\Phi\rangle$. The operator $h_{\Omega|\Phi} - h_\Phi$ is affiliated with \mathcal{A} since it is the derivative of the Connes cocycle $\Delta_{\Omega|\Phi}^{is} \Delta_\Phi^{-is}$ at $s = 0$. Because of this, $\langle \Phi | (h_{\Omega|\Phi} - h_\Phi) \Delta_\Phi (h_{\Omega|\Phi} - h_\Phi) | \Phi \rangle = \langle \Phi | (h_{\Omega|\Phi} - h_\Phi)^2 | \Phi \rangle$, which then implies $\langle \Phi | h(1 - \Delta)h | \Phi \rangle = 0$. We can therefore add any multiple of $(1 - \Delta)$ to $\frac{1}{2} - \Delta R_2(\Delta)$ without changing its expectation value in the state $h|\Phi\rangle$. It is straightforward to verify from (5.27) that $\frac{1}{2} - \Delta R_2(\Delta) - \frac{1}{3}(1 - \Delta) \geq 0$, and so

$$\langle \Phi | h \left(\frac{1}{2} - \Delta R_2(\Delta) \right) h | \Phi \rangle = \langle \Phi | h \left(\frac{1}{2} - \Delta R_2(\Delta) - \frac{1}{3}(1 - \Delta) \right) h | \Phi \rangle \geq 0. \quad (5.30)$$

This coincides with the lower bound in (5.24). Thus, the second order contribution to the entropy satisfies the bounds

$$0 \leq S^{(2)}(\rho_{\widehat{\Phi}}) \leq \frac{1}{2} k_g \langle \Phi | h^2 | \Phi \rangle. \quad (5.31)$$

The motivation for computing the subleading corrections to the crossed-product entropy was to try to verify a GSL beyond the leading order in the semiclassical expansion. The correction terms simplify into a fairly compact form, and the contributions from $R(\Delta, h)$ can be expressed in terms of a Kubo-Ando operator mean of Δ_Φ and $\frac{g(z)}{g(z-h)}$ [120–122], defined by the function R_2 appearing in (5.26). Although such operator means are useful in constructing functionals that satisfy data-processing inequalities, in the present context we were unable to prove monotonicity. The issue is that the operator mean appears sandwiched between the factors of $\delta(z, h)$, so that it is unclear whether the full entropy correction $\Delta S(\rho_{\widehat{\Phi}})$ satisfies a monotonicity property when evaluated for a crossed product associated with a later cut of the horizon. It is not even clear if we should expect $\Delta S(\rho_{\widehat{\Phi}})$ to be separately monotonic; the GSL, if it holds for crossed product algebras, would allow $\Delta S(\rho_{\widehat{\Phi}})$ to decrease as long as it is compensated by a larger increase in leading order entropy, determined by the decrease

in relative entropy $S_{\text{rel}}(\Phi||\Omega)$. If such a monotonicity result held, it would lead to a novel data-processing inequality for the relative entropy

$$S_{\text{rel}}^{\mathcal{C}_{\hat{\lambda}}}(\Phi||\Omega) - S_{\text{rel}}^{\mathcal{C}_{\lambda}}(\Phi||\Omega) \leq \Delta S(\rho_{\hat{\Phi}}^{\hat{\lambda}}) - \Delta S(\rho_{\hat{\Phi}}^{\lambda}) \quad (5.32)$$

with $\Delta S(\rho_{\hat{\Phi}})$ determined by (5.23) and an arbitrary positive probability distribution $g(z)$. In any situation where $\Delta S(\rho_{\hat{\Phi}})$ decreases, so that the right side of (5.32) is negative, this inequality would lead to an improvement of the standard data-processing inequality that only requires that the relative entropy decreases.

At this point, the only definitive statement we have is that it is difficult to violate monotonicity of the crossed product entropy perturbatively in the semiclassical expansion, because the corrections are $\mathcal{O}(\varepsilon^2)$. Thus in order to violate the second law, we would need the relative entropy for the quantum fields to be close to constant when moving between two cuts, so that its change is able to compete with the $\mathcal{O}(\varepsilon^2)$ terms. We also know that because the entropy satisfies a global second law, as discussed in section 4.2 and in reference [30], any violation of the local second law for crossed product entropies must necessarily be temporary. We leave the question of whether the second law can be violated at short times scales or whether it holds at all times to future work.

6 Monotonicity from crossed product subalgebras

The perturbative expressions for the crossed product entropies obtained in section 5 obeyed a number of interesting relations, but it was ultimately inconclusive whether these entropies satisfy a second law. Part of the issue is that the crossed product obscures the subalgebra structure of the underlying quantum field theory algebras, as is evidenced by the factors $e^{i\hat{p}h\Omega}$, $e^{-i\hat{p}h\Omega}$ that dress the quantum field operators appearing in the crossed product (5.3). Since the modular Hamiltonians h_{Ω}^{λ} for subalgebras \mathcal{C}_{λ} associated with different horizon cuts are different, generically these dressing factors will mix a subalgebra $\mathcal{C}_{\hat{\lambda}} \subset \mathcal{C}_{\lambda}$ throughout the entire algebra. Once we include the asymptotic charge \hat{q} in the algebra, this mixing will prevent the crossed product of the subalgebra, $\hat{\mathcal{C}}_{\hat{\lambda}} = \langle e^{i\hat{p}h_{\Omega}^{\hat{\lambda}}}\mathbf{c}e^{-i\hat{p}h_{\Omega}^{\hat{\lambda}}}, \hat{q} \rangle$ with $\mathbf{c} \in \mathcal{C}_{\hat{\lambda}}$, from obviously forming a subalgebra of the crossed product of the larger algebra $\hat{\mathcal{C}}_{\lambda} = \langle e^{i\hat{p}h_{\Omega}^{\lambda}}\mathbf{b}e^{-i\hat{p}h_{\Omega}^{\lambda}}, \hat{q} \rangle$ with $\mathbf{b} \in \mathcal{C}_{\lambda}$. Since a second law should have an underlying quantum information theoretic justification such as the existence of a quantum channel (which naturally arises when working with subalgebras), the failure of the crossed product algebras to form subalgebras poses a possible obstruction to the existence of a generalized second law beyond the leading semiclassical approximation.

This suggests that one possibility for arriving at a second law is to look for a subalgebra of $\hat{\mathcal{C}}_{\lambda}$ that is naturally isomorphic to the crossed product of the quantum field subalgebra $\hat{\mathcal{C}}_{\hat{\lambda}}$. The condition for such subalgebras to occur is precisely the existence of an *operator-valued weight* from \mathcal{C}_{λ} to $\mathcal{C}_{\hat{\lambda}}$ [45]. An operator-valued weight can be viewed as a sort of unnormalized conditional expectation between von Neumann algebras that may be finite on only a subset of operators; therefore, it generalizes the notion of a conditional expectation in the same

way that weights generalize the notion of states on the von Neumann algebra. Formally, an operator-valued weight for an algebra inclusion $\mathcal{B} \subset \mathcal{A}$ is a linear map T from \mathcal{A}_+ , the positive operators in \mathcal{A} , into the extended positive cone $\tilde{\mathcal{B}}_+$ of \mathcal{B} , which roughly consists of positive operators in \mathcal{B} along with the addition of certain unbounded positive operators affiliated with \mathcal{B} [87, Section IX.4] [44, 45]. Like conditional expectations, operator-valued weights satisfy the bimodule property $T(\mathfrak{b}^\dagger \mathfrak{a} \mathfrak{b}) = \mathfrak{b}^\dagger T(\mathfrak{a}) \mathfrak{b}$ for all $\mathfrak{b} \in \mathcal{B}$, $\mathfrak{a} \in \mathcal{A}_+$.

If T is an operator-valued weight satisfying the additional technical requirements of faithfulness, normality, and semifiniteness, then we can take a weight φ on \mathcal{B} and compose it with T to obtain a faithful, semifinite, normal weight $\omega = \varphi \circ T$ on the larger algebra \mathcal{A} . This weight has the property that the modular flow with respect to ω preserves the subalgebra \mathcal{B} , and agrees with the modular flow of the weight φ on \mathcal{B} . This result is a direct generalization of Takesaki's theorem for conditional expectations to the case of operator-valued weights, and in fact the existence of the pair of weights ω, φ on \mathcal{A}, \mathcal{B} with this property is equivalent to the existence of the operator-valued weight T [45]. The main difference from a conditional expectation is that T may assign infinite values to the entire subalgebra \mathcal{B} while still being finite on a dense subalgebra of \mathcal{A} , whereas a conditional expectation assigns a finite value to all operators and is normalized to map the identity $\mathbb{1}$ to itself.

Before discussing the implications of the operator-valued weight on the structure of crossed products, we first consider whether one should expect an operator-valued weight to exist between two horizon-cut algebras $\mathcal{C}_{\tilde{\lambda}} \subset \mathcal{C}_\lambda$. One reason to expect that the operator-valued weight exists is due to the analogy with split inclusions. A split inclusion $\mathcal{B} \subset \mathcal{A}$ is one for which there exists a type I factor \mathcal{N} that interpolates \mathcal{B} and \mathcal{A} , in the sense that $\mathcal{B} \subset \mathcal{N} \subset \mathcal{A}$ [123]. Such inclusions often occur for quantum field theory algebras when \mathcal{B} and \mathcal{A} are algebras associated with causally complete nested subregions $R_{\mathcal{A}}$ and $R_{\mathcal{B}}$ such that there is a finite corridor between $R_{\mathcal{B}}$ and $R_{\mathcal{A}}$; the technical requirement on the regions is that the closure of $R_{\mathcal{B}}$ is contained in the interior of $R_{\mathcal{A}}$, and the field theory is said to satisfy the *split property* if in such a situation $\mathcal{B} \subset \mathcal{A}$ is always a split inclusion [49, Section V.5].

When both \mathcal{A} and \mathcal{B} are type III₁ algebras associated with local subregions in quantum field theory, the existence of the type I interpolating algebra \mathcal{N} implies that there cannot be a (normal) conditional expectation E from \mathcal{A} to \mathcal{B} . The reason is that such a conditional expectation would restrict to a conditional expectation from \mathcal{N} to \mathcal{B} , which is not possible since conditional expectations cannot exist to a subalgebra with a higher type, i.e. there are no conditional expectations from a type I algebra to a type III subalgebra. On the other hand, \mathcal{N} at the same time guarantees the existence of a normal, faithful, semifinite operator-valued weight T from \mathcal{A} to \mathcal{B} . This is because there always exists such an operator-valued weight when either the algebra or the subalgebra is type I, so there are operator-valued weights $T_{\mathcal{N}, \mathcal{A}}$ from \mathcal{A} to \mathcal{N} and $T_{\mathcal{B}, \mathcal{N}}$ from \mathcal{N} to \mathcal{B} [44, 45]. Composing these results in an operator-valued weight $T = T_{\mathcal{B}, \mathcal{N}} \circ T_{\mathcal{N}, \mathcal{A}}$ from \mathcal{A} to \mathcal{B} , which is normal, faithful, and semifinite if both $T_{\mathcal{N}, \mathcal{A}}$ and $T_{\mathcal{B}, \mathcal{N}}$ are [87, Proposition IX.4.16].

Unfortunately, the existence of an operator-valued weight in the case of a split inclusion does not give any immediate information about the horizon-cut inclusion $\mathcal{C}_{\tilde{\lambda}} \subset \mathcal{C}_\lambda$, which is instead a half-sided modular inclusion. Half-sided modular inclusions are never split [43],

so the above arguments do not apply for algebras associated with nested cuts of a Killing horizon. In this case, it still seems unlikely that there is a conditional expectation from \mathcal{C}_λ to $\mathcal{C}_{\tilde{\lambda}}$; for example, in the free field theory case where the relative commutant $\mathcal{C}'_{\tilde{\lambda}} \wedge \mathcal{C}_\lambda$ forms a nontrivial factor consisting of fields smeared between the cuts, the conditional expectation would imply a factorization of the algebras across the horizon cut $\tilde{\lambda}$, which is not a property expected in quantum field theory.¹¹ However, this leaves open the possibility that there exists an operator-valued weight from \mathcal{C}_λ to $\mathcal{C}_{\tilde{\lambda}}$. One might expect that you could argue for such an operator-valued weight by considering a sequence of algebras $\mathcal{C}_{\tilde{\lambda},n} \subset \mathcal{C}_{\tilde{\lambda}}$ of regions whose entangling surfaces are located off of the horizon and spacelike separated from the entangling surface of \mathcal{C}_λ , which approach $\mathcal{C}_{\tilde{\lambda}}$ in the limit $n \rightarrow \infty$. Since each inclusion $\mathcal{C}_{\tilde{\lambda},n} \subset \mathcal{C}_\lambda$ is then split, there is a sequence of operator-valued weights T_n to each of these subalgebras. One might hope that these limit to an operator-valued weight T from \mathcal{C}_λ to $\mathcal{C}_{\tilde{\lambda}}$ in the limit $n \rightarrow \infty$; unfortunately, it is difficult to make a rigorous argument to this effect. Nevertheless, there is no argument we know of that obviously precludes such an operator-valued weight, and so we will conjecture that one indeed exists, and explore the implications for the crossed product algebras.

The existence of an operator-valued weight between horizon-cut algebras implies a subalgebra structure for the crossed product algebras. To simplify notation for the rest of this section, we denote the earlier horizon cut algebra as \mathcal{B} and the later algebra as $\mathcal{C} \subset \mathcal{B}$, with T denoting the operator-valued weight from \mathcal{B} to \mathcal{C} . To see how the crossed product $\widehat{\mathcal{C}}$ embeds as a subalgebra of $\widehat{\mathcal{B}}$, we begin by defining a weight χ on \mathcal{B} by composing the vacuum state on \mathcal{C} , $\omega_{\mathcal{C}} = \langle \Omega | \cdot | \Omega \rangle$, with the operator-valued weight,

$$\chi = \omega_{\mathcal{C}} \circ T. \quad (6.1)$$

We denote the unitary operator implementing modular flow for this weight on \mathcal{B} as $\Delta_{\chi|\omega}^{-is} = e^{ish_{\chi|\omega}}$, where ω refers to the vacuum weight on the commutant algebra \mathcal{B}' , and is needed to define how $h_{\chi|\omega}$ acts in the global QFT Hilbert space, including its action on operators in \mathcal{B}' . Since χ is obtained as a pullback via T , Haagerup's theorem for operator-valued weights [45] guarantees that modular flow with respect to $\Delta_{\chi|\omega}^{-is}$ on the subalgebra \mathcal{C} agrees with modular flow for the state $\omega_{\mathcal{C}}$, generated by $\Delta_{\omega_{\mathcal{C}}}^{-is} = e^{ish_{\omega_{\mathcal{C}}}}$,

$$\Delta_{\chi|\omega}^{-is} \mathbf{c} \Delta_{\chi|\omega}^{is} = \Delta_{\omega_{\mathcal{C}}}^{-is} \mathbf{c} \Delta_{\omega_{\mathcal{C}}}^{is}, \quad \forall \mathbf{c} \in \mathcal{C}. \quad (6.2)$$

This relation immediately demonstrates that the crossed product $\widetilde{\mathcal{B}}$ constructed using modular flow for the weight χ contains the crossed product $\widehat{\mathcal{C}}$ as a subalgebra, since $\widehat{\mathcal{C}} = \langle \Delta_{\omega_{\mathcal{C}}}^{-i\hat{p}} \mathbf{c} \Delta_{\omega_{\mathcal{C}}}^{i\hat{p}} e^{it\hat{q}} \rangle = \langle \Delta_{\chi|\omega}^{-i\hat{p}} \mathbf{c} \Delta_{\chi|\omega}^{i\hat{p}} e^{it\hat{q}} \rangle \subset \langle \Delta_{\chi|\omega}^{-i\hat{p}} \mathbf{b} \Delta_{\chi|\omega}^{i\hat{p}} e^{it\hat{q}} \rangle = \widetilde{\mathcal{B}}$. We can then apply Connes's result [125] [87, Section VIII.3] that modular flows with respect to different weights are related to each other by a unitary in the algebra,

$$\Delta_{\chi|\omega}^{-is} \mathbf{b} \Delta_{\chi|\omega}^{is} = u_{\chi|\omega}(-s) \Delta_{\omega}^{-is} \mathbf{b} \Delta_{\omega}^{is} u_{\chi|\omega}(-s)^\dagger, \quad (6.3)$$

¹¹In this case, one can argue that the inclusion extends to a Mobius-covariant net associated with a family of translated horizon cuts, which allows one to prove that no conditional expectation from \mathcal{C}_λ to $\mathcal{C}_{\tilde{\lambda}}$ exists [124].

where the Connes cocycle $u_{\chi|\omega}(-s)$ is defined by

$$u_{\chi|\omega}(-s) = \Delta_{\chi|\omega}^{-is} \Delta_{\omega}^{is} \in \mathcal{B}. \quad (6.4)$$

Following the arguments from [29], we can conjugate the crossed product algebra $\tilde{\mathcal{B}}$ by a unitary constructed from the cocycle

$$U \equiv \mathbf{u}(-\hat{p}) := u_{\chi|\omega}(-\hat{p}) = \Delta_{\chi|\omega}^{-i\hat{p}} \Delta_{\omega}^{i\hat{p}} \quad (6.5)$$

in order to map it into the original crossed product algebra $\hat{\mathcal{B}} = \langle \Delta_{\omega}^{-i\hat{p}} \mathbf{b} \Delta_{\omega}^{i\hat{p}} e^{it\hat{q}} \rangle$. To show this, we need the identity

$$U^\dagger e^{it\hat{q}} U = \Delta_{\omega}^{-i\hat{p}} e^{it(\hat{q} - h_{\chi|\omega})} \Delta_{\omega}^{i\hat{p}} = \Delta_{\omega}^{-i\hat{p}} e^{-ith_{\chi|\omega}} \Delta_{\omega}^{i\hat{p}} e^{ith_{\omega}} e^{it\hat{q}} = \Delta_{\omega}^{-i\hat{p}} u_{\chi|\omega}(t) \Delta_{\omega}^{i\hat{p}} e^{it\hat{q}} \quad (6.6)$$

which is an element of $\hat{\mathcal{B}}$ since $u_{\chi|\omega}(t) \in \mathcal{B}$. Hence, a generic element $\tilde{\mathbf{b}} = \Delta_{\chi|\omega}^{-i\hat{p}} \mathbf{b} \Delta_{\chi|\omega}^{i\hat{p}} e^{it\hat{q}}$ in $\tilde{\mathcal{B}}$ maps to

$$U^\dagger \tilde{\mathbf{b}} U = \Delta_{\omega}^{-i\hat{p}} \mathbf{b} u_{\chi|\omega}(t) \Delta_{\omega}^{i\hat{p}} e^{it\hat{q}} \in \hat{\mathcal{B}}. \quad (6.7)$$

This shows that $U^\dagger \tilde{\mathcal{B}} U = \hat{\mathcal{B}}$.

Now because $\hat{\mathcal{C}} \subset \tilde{\mathcal{B}}$, the conjugated algebra $\bar{\mathcal{C}} = U^\dagger \hat{\mathcal{C}} U$ is a subalgebra of $\hat{\mathcal{B}}$. By conjugating a generic element $\Delta_{\chi|\omega}^{-i\hat{p}} \mathbf{c} \Delta_{\chi|\omega}^{i\hat{p}} e^{it\hat{q}}$ in $\hat{\mathcal{C}}$ by $U^\dagger(\cdot)U$, we find a generating set of operators for $\bar{\mathcal{C}}$:

$$\bar{\mathcal{C}} = \langle \Delta_{\omega}^{-i\hat{p}} \mathbf{c} u_{\chi|\omega}(t) \Delta_{\omega}^{i\hat{p}} e^{it\hat{q}} \rangle \quad (6.8)$$

Thus, we have constructed a subalgebra of $\hat{\mathcal{B}}$ unitarily equivalent to $\hat{\mathcal{C}}$. We can therefore define an inclusion map $\iota_{\hat{\mathcal{B}}, \hat{\mathcal{C}}}$ from $\hat{\mathcal{C}}$ into $\hat{\mathcal{B}}$ by

$$\iota_{\hat{\mathcal{B}}, \hat{\mathcal{C}}}(\hat{\mathbf{c}}) = U^\dagger \hat{\mathbf{c}} U. \quad (6.9)$$

Since this is a UCP map, its dual is a quantum channel from states on $\hat{\mathcal{B}}$ to states on $\hat{\mathcal{C}}$. We would therefore like to leverage this channel and the monotonicity of relative entropy to arrive at some version of a second law for the crossed product algebras.

The most straightforward way to do this is to examine the relative entropy of a generic state $|\hat{\Phi}\rangle$ with respect to a specific vacuum weight on the algebra. In general, it is possible to define the relative entropy $S_{\text{rel}}(\varphi||\psi)$ where φ is a state while ψ is only a weight which may not be normalized, and could even be unbounded in the sense that $\psi(\mathbb{1}) = \infty$ [81, Chapter 7]. In fact, the type II_∞ entropies that we have computed throughout this paper are naturally interpreted as minus the relative entropy with respect to the tracial weight on the algebra [81, 102, 126]. For algebras possessing a trace, one can always find a density matrix ρ_ψ affiliated with the algebra that reproduces expectation values for the weight via

$$\psi(\mathbf{b}) = \text{Tr}(\rho_\psi \mathbf{b}). \quad (6.10)$$

Because the weight ψ need not be normalized, this density matrix no longer necessarily satisfies $\text{Tr} \rho_\psi = 1$.

For the crossed product, choosing the trace as the reference weight is problematic because the trace on $\widehat{\mathcal{B}}$ does not restrict to a semifinite weight on the subalgebra $\widehat{\mathcal{C}}$. Instead, there is an operator-valued weight \widehat{T} from $\widehat{\mathcal{B}}$ to $\widehat{\mathcal{C}}$ constructed from T such that $\text{Tr}_{\widehat{\mathcal{B}}} = \text{Tr}_{\widehat{\mathcal{C}}} \circ \widehat{T}$, where $\text{Tr}_{\widehat{\mathcal{C}}}$ is the semifinite trace on $\widehat{\mathcal{C}}$. However, unless \widehat{T} is bounded so that it can be normalized to be a conditional expectation (which we do not expect for nested horizon cuts), \widehat{T} will blow up on the subalgebra $\widehat{\mathcal{C}}$, showing that $\text{Tr}_{\widehat{\mathcal{B}}}$ is not semifinite on this algebra. A more convenient reference is the dual weight $\widehat{\omega}$ arising from the vacuum state $|\Omega\rangle$ in the crossed product construction. This dual weight can be formally represented as arising from an infinite norm vector $|\widehat{\Omega}\rangle = \sqrt{2\pi}|\Omega, 0_p\rangle$, where $|0_p\rangle$ is the zero momentum eigenstate used in the expression for the trace on the crossed product. This vector also defines the dual weight for the $\widehat{\mathcal{C}}$ crossed product algebra, and furthermore has the property that it is invariant under the unitary $U = u(-\hat{p})$ in equation (6.5) defining the map from $\widehat{\mathcal{C}}$ to $\widehat{\mathcal{C}}$. Defining the weight $\widehat{\omega}_{\mathcal{B}} = 2\pi\langle\Omega, 0_p|(\cdot)|\Omega, 0_p\rangle$ and similarly for $\widehat{\omega}_{\mathcal{C}}$, we find that $\widehat{\omega}_{\mathcal{B}}$ restricts to a semifinite weight on $\widehat{\mathcal{C}}$, and further more pulls back to the weight $\widehat{\omega}_{\mathcal{C}}$ under the inclusion map (6.9),

$$(\widehat{\omega}_{\mathcal{B}} \circ \iota_{\widehat{\mathcal{B}}, \widehat{\mathcal{C}}})(\widehat{\mathcal{C}}) = 2\pi\langle\Omega, 0_p|U^\dagger \widehat{\mathcal{C}} U|\Omega, 0_p\rangle = \widehat{\omega}_{\mathcal{C}}(\widehat{\mathcal{C}}). \quad (6.11)$$

We now consider another state $\widehat{\varphi}$ arising from a vector $|\widehat{\Phi}\rangle$. Monotonicity of relative entropy for the UCP map $\iota_{\widehat{\mathcal{B}}, \widehat{\mathcal{C}}}$ then states that

$$S_{\text{rel}}^{\widehat{\mathcal{B}}}(\widehat{\varphi}||\widehat{\omega}_{\mathcal{B}}) \geq S_{\text{rel}}^{\widehat{\mathcal{C}}}(\widehat{\varphi} \circ \iota_{\widehat{\mathcal{B}}, \widehat{\mathcal{C}}}||\widehat{\omega}_{\mathcal{B}} \circ \iota_{\widehat{\mathcal{B}}, \widehat{\mathcal{C}}}) = S_{\text{rel}}^{\widehat{\mathcal{C}}}(\widehat{\varphi} \circ \iota_{\widehat{\mathcal{B}}, \widehat{\mathcal{C}}}||\widehat{\omega}_{\mathcal{C}}) \quad (6.12)$$

where the second equality applies (6.11). The state $\widehat{\varphi} \circ \iota_{\widehat{\mathcal{B}}, \widehat{\mathcal{C}}}$ on $\widehat{\mathcal{C}}$ is the expectation value in the vector state $U|\widehat{\Phi}\rangle$. To convert this into a statement about entropies, we note that the density matrices for the weights $\widehat{\omega}_{\mathcal{B}}$ and $\widehat{\omega}_{\mathcal{C}}$ are given in both cases by

$$\rho_{\widehat{\omega}} = e^{\hat{q}}, \quad (6.13)$$

which follows immediately from the definition (6.10) of the density matrix and the expression (2.9) for the trace on the two algebras.

In terms of the density matrices, the relative entropy is given by the usual expression

$$S_{\text{rel}}(\widehat{\varphi}||\widehat{\omega}) = \text{Tr}(\rho_{\widehat{\varphi}}(\log \rho_{\widehat{\varphi}} - \log \rho_{\widehat{\omega}})) \quad (6.14)$$

and hence

$$S_{\text{rel}}^{\widehat{\mathcal{B}}}(\widehat{\varphi}||\widehat{\omega}_{\mathcal{B}}) = -S(\rho_{\widehat{\Phi}}^{\widehat{\mathcal{B}}}) - \langle\widehat{\Phi}|\hat{q}|\widehat{\Phi}\rangle, \quad (6.15)$$

showing that this relative entropy can be interpreted as a free energy with respect to the asymptotic charge $-\hat{q}$, which is given by either the asymptotic area or the ADM mass in the crossed product constructions discussed in sections 3.3 and 4.2 (see equations (3.45) and (4.29)). The right hand side of (6.12) evaluates to

$$S_{\text{rel}}^{\widehat{\mathcal{C}}}(\widehat{\varphi} \circ \iota_{\widehat{\mathcal{B}}, \widehat{\mathcal{C}}}||\widehat{\omega}_{\mathcal{C}}) = -S(\rho_{U\widehat{\Phi}}^{\widehat{\mathcal{C}}}) - \langle\widehat{\Phi}|U^\dagger \hat{q} U|\widehat{\Phi}\rangle. \quad (6.16)$$

Hence, the monotonicity result (6.12) implies the entropy inequality

$$S(\rho_{U\widehat{\Phi}}^{\widehat{\mathcal{C}}}) + \langle\widehat{\Phi}|U^\dagger \hat{q} U - \hat{q}|\widehat{\Phi}\rangle \geq S(\rho_{\widehat{\Phi}}^{\widehat{\mathcal{B}}}). \quad (6.17)$$

This is not quite the statement of the second law we were seeking. First, the entropy on the left hand side is computed for a different state $U|\widehat{\Phi}\rangle$ for the algebra $\widehat{\mathcal{C}}$ instead of the original state $|\widehat{\Phi}\rangle$. Second, there is an additional term involving the difference in expectation value of \hat{q} between the state $U|\widehat{\Phi}\rangle$ and $|\widehat{\Phi}\rangle$. As it turns out, at leading order in the semiclassical expansion, these two differences cancel out, and the inequality (6.17) reduces to the semiclassical second law discussed in sections 3.3 and 4.2. To see this, we set $|\widehat{\Phi}\rangle = |\Phi, f\rangle$ with the wavefunction f slowly varying, $\frac{f'}{f} \sim \mathcal{O}(\varepsilon)$. As discussed in [30], because the unitary U is a cocycle flow with parameter \hat{p} , it leaves the state $|\Phi, f\rangle$ invariant up to corrections of order ε . This means that at this order, we can approximate the density matrix $\rho_{U\widehat{\Phi}}$ by $\rho_{\widehat{\Phi}}$, and hence

$$-\log \rho_{U\widehat{\Phi}} \approx -\log \rho_{\widehat{\Phi}} \approx -\hat{q} - h_{\Omega|\Phi} + h_{\Phi} - \log |f(\hat{q})|^2 + \mathcal{O}(\varepsilon). \quad (6.18)$$

The entropy $S(\rho_{U\widehat{\Phi}})$ is obtained as the expectation value of $-\log \rho_{U\widehat{\Phi}}$ in the state $U|\widehat{\Phi}\rangle$. Again because U leaves the state $|\widehat{\Phi}\rangle$ nearly invariant, most terms in this expectation value are the same as in the state $|\widehat{\Phi}\rangle$ at leading order in the semiclassical expansion. The exception is the operator \hat{q} , which shifts when conjugated by the cocycle,

$$U^\dagger \hat{q} U = \hat{q} + h_\omega - e^{i\hat{p}h_\omega} h_{\chi|\omega} e^{-i\hat{p}h_\omega}. \quad (6.19)$$

However, we see that this shift is compensated by the additional terms appearing in the inequality (6.17): the entropy of $\rho_{U\widehat{\Phi}}$ admits the approximation,

$$\begin{aligned} S(\rho_{U\widehat{\Phi}}^{\widehat{\mathcal{C}}}) &\approx \langle \widehat{\Phi} | U^\dagger (-\hat{q} - h_{\Omega|\Phi} + h_{\Phi} - \log |f(\hat{q})|^2) U | \widehat{\Phi} \rangle \\ &\approx \langle \widehat{\Phi} | -\hat{q} - h_{\Omega|\Phi} + h_{\Phi} - \log |f(\hat{q})|^2 | \widehat{\Phi} \rangle + \langle \widehat{\Phi} | \hat{q} - U^\dagger \hat{q} U | \widehat{\Phi} \rangle \\ &\approx S(\rho_{\widehat{\Phi}}^{\widehat{\mathcal{C}}}) + \langle \widehat{\Phi} | \hat{q} - U^\dagger \hat{q} U | \widehat{\Phi} \rangle \end{aligned} \quad (6.20)$$

and plugging this into the inequality (6.17) reproduces the second law for crossed product algebras at leading order in the semiclassical expansion,

$$S(\rho_{\widehat{\Phi}}^{\widehat{\mathcal{C}}}) + \mathcal{O}(\varepsilon) \geq S(\rho_{\widehat{\Phi}}^{\widehat{\mathcal{B}}}). \quad (6.21)$$

Hence, assuming the existence of the operator-valued weight between horizon cuts, we are able to establish a subalgebra explanation for the semiclassical generalized second law derived in sections 3.3 and 4.2.

Finally, it is interesting to note that the terms in (6.17) that could spoil the proposed second law can be written as a relative entropy difference. Noting that $S_{\text{rel}}^{\widehat{\mathcal{C}}}(\widehat{\Phi} || \widehat{\omega}_C) = -S(\rho_{\widehat{\Phi}}^{\widehat{\mathcal{C}}}) - \langle \hat{q} \rangle_{\widehat{\Phi}}$, we can rewrite the inequality (6.17) as

$$S(\rho_{\widehat{\Phi}}^{\widehat{\mathcal{C}}}) + S_{\text{rel}}^{\widehat{\mathcal{C}}}(\widehat{\Phi} || \widehat{\omega}_C) - S_{\text{rel}}^{\widehat{\mathcal{C}}}(U\widehat{\Phi} || \widehat{\omega}_C) \geq S(\rho_{\widehat{\Phi}}^{\widehat{\mathcal{B}}}). \quad (6.22)$$

Hence, depending on the sign of the relative entropy difference $S_{\text{rel}}^{\widehat{\mathcal{C}}}(\widehat{\Phi} || \widehat{\omega}_C) - S_{\text{rel}}^{\widehat{\mathcal{C}}}(U\widehat{\Phi} || \widehat{\omega}_C)$ between the states $|\widehat{\Phi}\rangle$ and $U|\widehat{\Phi}\rangle$ on $\widehat{\mathcal{C}}$, we can either guarantee the second law holds (when the difference is negative or zero), or derive a necessary condition on the state $|\widehat{\Phi}\rangle$ for a violation

to occur (the difference is positive). At present, there does not seem to be any constraint on the sign of this relative entropy difference, but examining it more closely may result in further insights into the crossed product second law. One statement that seems to hold is that this difference should be $\mathcal{O}(\varepsilon^2)$ in the semiclassical expansion, since the perturbative calculations of section 5 demonstrated that the second law for semiclassical states holds up to linear order in ε .

7 Discussion

The overarching goal of this paper has been to explore properties of subsystem gravitational algebras for regions defined by arbitrary cuts of a Killing horizon. These regions provide nontrivial examples where the modular Hamiltonian of a vacuum state $|\Omega\rangle$ is known and generates a geometric flow along the horizon, which facilitates the construction of type II_∞ gravitationally dressed crossed-product algebras following the general procedure of JSS [33]. As reviewed in section 3.1, the geometric flow of this state follows from properties of half-sided translations and the ANEC, and holds even for interacting quantum field theories. We found that entropies of semiclassical states in the crossed product algebra reproduce the generalized entropy of the horizon cut, and for these states we were able to reduce the second law for crossed product algebras to Wall’s derivation of the generalized second law [38]. The crossed-product second law improves on Wall’s result by identifying a gravitational von Neumann algebra whose entropy is computed by the generalized entropy, with the further advantage that crossed-product entropies are manifestly UV finite, so that one does not need to worry about subtleties involving the renormalization of the quantum field entanglement entropy.

After discussing the case of AdS black holes, we presented in section 4.1 a novel approach to treating asymptotically flat and de Sitter black holes using the concept of an algebra at infinity for the half-sided translation. This algebra naturally characterizes the degrees of freedom that never enter the black hole. Using a theorem of Borchers which guarantees the existence of a conditional expectation to this algebra at infinity [85], we discussed how to handle the generalized second law in this context, even in the case of an interacting theory where one expects the algebra of fields strictly localized to the horizon to the future of a cut to be trivial. We also described the construction of gravitational algebras for these alternative asymptotics, which lead to nontrivial matching conditions at asymptotic infinity.

Finally, we explored the question of whether the crossed product algebras satisfy a generalized second law beyond the leading semiclassical approximation of the states. We derived in section 5 formulas for the corrections to the crossed-product entropy, and showed that the first correction appears at second order in the semiclassical expansion. Although we obtained explicit expressions for the entropy corrections, we were unable to determine whether these corrections can be used to violate the second law at short time scales. However, we noted that because the gravitational algebra at infinity defines a trace-preserving inclusion inside the dressed horizon-cut algebras, the entropy at infinity must be larger than that of the horizon cut algebra. This gives a global second law which generalizes a similar result

appearing in [30], and necessarily constrains any possible violation of the second law between horizon cuts to be temporary. After this, we explored in section 6 the possibility that nested horizon-cut gravitational algebras could form a subalgebra structure using the conjectured existence of an operator-valued weight between horizon cuts. This leads to a monotonicity constraint on entropies between two horizon cuts, but involving different states at different cuts. We showed that this monotonicity result still reduces to the second law at leading order in the semiclassical expansion, and left open the question of whether it can be leveraged to produce a stronger result.

The investigations carried out in this work have led to a number of intriguing questions and directions for future work, so we conclude by describing some of these.

Singular vacua as semifinite weights. In the discussion of rotating black holes in section 4.3, we confronted the issue that there does not appear to be a natural vacuum state whose modular flow is geometric on the algebra at infinity. The closest analog is the Frolov-Thorne vacuum, which is well known to become singular in the region far from the black hole where the horizon-generating Killing vector is spacelike. Ordinarily, quantization about such a singular state results in a unitarily inequivalent representation of the quantum field algebra, which is likely not the correct description of quantum fields on a smooth background geometry. However, we raised the possibility that one could reinterpret this vacuum as a nonnormalizable weight on the physical von Neumann algebra of fields in the black hole exterior. Having a weight, as opposed to a state, whose modular flow is geometric is sufficient to arrive at a gravitational algebra with a trace following the JSS construction. Similarly, the Unruh vacuum, which provides the more accurate description of an astrophysical black hole formed from collapse, yields a unitarily inequivalent representation of the quantum field algebra from the Hartle-Hawking vacuum. However, the latter vacuum provides the state whose modular flow is geometric on the horizon, and hence one would like a description where both states can be treated in a single representation of the field algebra. We suggested the possibility that when quantizing around the Unruh vacuum, the Hartle-Hawking vacuum could be treated as a semifinite weight.

These points motivate the broader question of determining when a singular state of the quantum field algebra can be reinterpreted as a semifinite weight. One important aspect of such a procedure is identifying the correct definition subalgebra for the weight (see e.g. [87, Section VII.1]). Given a weight φ on a von Neumann algebra \mathcal{A} , we can identify a subcone of the set of positive operators \mathcal{A}_+ with finite weight,

$$\mathfrak{p}_\varphi = \{\mathfrak{a} \in \mathcal{A}_+ | \varphi(\mathfrak{a}) < \infty\}. \quad (7.1)$$

The linear span of \mathfrak{p}_φ forms the *definition subalgebra* $\mathfrak{m}_\varphi \subset \mathcal{A}$ of the weight φ . In general, \mathfrak{m}_φ is not a C^* -algebra since it need not contain uniform limits of operators, and therefore is not a von Neumann algebra either. The definition subalgebra is also not unital for nonnormalizable weights, since if $\varphi(\mathbb{1}) < \infty$, one could rescale φ to make it a state. If the weight φ is semifinite, the weak closure of \mathfrak{m}_φ yields the full von Neumann algebra \mathcal{A} . Because of this last fact, one can still perform an analog of the GNS representation for this algebra with respect to the weight φ called the *semicyclic representation*, and such a representation with respect to

a faithful, semifinite, normal weight is unitarily equivalent to the GNS representation with respect to a faithful normal state. This fact allows many of the results of Tomita-Takesaki theory to apply to the case of semifinite weights.

The re-interpretation of singular states in quantum field theory as semifinite weights thus requires a careful analysis of the associated definition subalgebra \mathfrak{m}_φ . Since in the case of the Frolov-Thorne vacuum, the region in the black hole exterior where the state is regular has a causal completion consisting of the entire exterior, it is possible this region can be used to identify an appropriate weakly dense definition subalgebra for the Frolov-Thorne vacuum, viewed as a weight. Similar comments apply to the Hartle-Hawking vacuum for the algebra constructed from the Unruh state. We believe that working out the details of this weight interpretation of singular vacua may provide important insights into the relevance of these states and further inform the construction of gravitational algebras.

GSL violation or improved monotonicity. Upon computing the corrections to the crossed product entropy for states of the form $|\Phi, f\rangle$ in section 5, we were ultimately unable to determine whether these corrections can be used to violate the leading order second law found in sections 3.3 and 4.2. The leading order statement follows from monotonicity of relative entropy for the quantum field subalgebra of the crossed product, and so if the entropy still satisfies a second law upon including the subleading corrections, it would require a modified bound on the amount the quantum field relative entropy decreases between horizon cuts. In some cases, this bound would improve the standard data-processing inequality that relative entropy must decrease between horizon cuts. It would therefore be interesting to relate the proposed bound to other recent works on improving the data-processing inequality and its connection to universal recovery channels [115–117].

One possible avenue to arrive at a nontrivial bound is the conjectured subalgebra structure for crossed products discussed in section 6. This structure relies on the existence of an operator-valued weight between algebras for nested cuts of the Killing horizon, whose existence has not been proven. This leads to the following purely mathematical questions in the theory of operator algebras: given a half-sided modular inclusion $\mathcal{N} \subset \mathcal{M}$, can there exist an operator-valued weight T from \mathcal{M} to \mathcal{N} ? Does this always occur, and if one exists, what are its properties? In the context of half-sided modular inclusions arising from horizon cuts, if such an operator-valued weight existed, it should possess a natural construction using tools from quantum field theory. In the interacting case where we expect the relative commutant $\mathcal{N}' \wedge \mathcal{M}$ to be trivial, the operator-valued weight would have to be unique, since if another operator-valued weight existed, the cocycle between the two weights would define a nontrivial element of the relative commutant. Even in the free case where the relative commutant is nontrivial, there should be a preferred operator-valued weight that restricts to the horizon vacuum state on the relative commutant. Hence, a tangible goal for the future is to explicitly construct the operator-valued weight, or prove that one does not exist.

Assuming the existence of such an operator-valued weight, we found a version of the second law for crossed products, equation (6.17), that does not rely on any semiclassical approximation. Unfortunately, we do not at present have a simple interpretation of the

derived inequality, since it involves the entropy of a different state at a later cut of the horizon, as well as the change in the expectation value of the asymptotic charge between the two states. It would be interesting to better understand this inequality, since at the very least it constrains any violation of the crossed product second law to be bounded by a relative entropy difference of two states, as evidenced by equation (6.22).

Connections to dynamical horizon entropy. One output of the construction of gravitational algebras for horizon cuts is that it yields an algebra whose entropy directly coincides with the generalized entropy of the horizon cut. This supports the idea that in the limit of weak gravitational couple, this entropy can be interpreted as an entanglement entropy, subject to the caveat that entropies in type II algebras are most naturally thought of as vacuum-subtracted entropies [29, 102]. Furthermore, it reinforces the interpretation of the horizon area as capturing the classical component of the black hole entropy. Recently, an alternative proposal for the entropy of dynamical black holes has appeared that modifies this classical component by a term involving the perturbative expansion of the horizon [74, 76, 127]. This raises the question of whether this entropy has a quantum generalization, and whether one can account for it in the framework of semiclassical gravitational algebras.

We note that one aspect of the construction of the gravitational algebras in the present work is the use of the event horizon boundary condition in section 3.2 in order to interpret the asymptotic gravitational charge as the late time area of the horizon. Interestingly, the new dynamical entropy proposal can be interpreted as the area of an apparent horizon inside the black hole, as opposed to the area of the event horizon [74, 76]. This is strongly reminiscent of the Engelhardt-Wall proposal for the coarse-grained outer entropy in the context of AdS/CFT [128, 129]. In our setup, we could arrive at a null surface that intersects the apparent horizon by relaxing the event horizon boundary condition that the asymptotic expansion Θ goes to zero. This also suggests that we should include an asymptotic gravitational charge associated with this freedom to choose this boundary condition, since the late time expansion naturally arises as an asymptotic charge associated to the null translation along the horizon. The asymptotic past and future expansions are related to the average null energy along the horizon via a gravitational constraint (3.29), so it raises the possibility that one should include this asymptotic charge and gravitational dressing for this constraint as well. It may be that including this translational gravitational charge would lead to a natural connection to the new dynamical entropy, and may also help in a more complete formulation of the generalized second law, without the possibility of temporary entropy decrease beyond the leading semiclassical approximation.

Applications to quantum focusing. A natural generalization of the present work is to investigate the crossed product construction for more general subregions. Ultimately, we would like to be able to formulate the quantum focusing conjecture (QFC) [12] as a statement about crossed product entropies for subregions bounded by a causal horizon. One benefit of such a description is that it would handle the problem of UV divergences in the entanglement entropy, which are one of the main technical obstructions to obtaining rigorous statements about quantum focusing. On the other hand, there is a question of

what more we could learn from a crossed-product formulation of the QFC. The gravitational algebras considered here appear in the $G_N \rightarrow 0$ limit of quantum gravity, and in this limit the quantum focusing conjecture in most cases reduces to the classical focusing theorem of general relativity. In order to arrive at a nontrivial quantum result, one needs to consider background geometries that classically saturate the focusing inequality, so that perturbative corrections to the geometry can compete with contributions coming from the change in entanglement entropy as one evolves along a light sheet. In this limit, quantum focusing generally reduces to the quantum null energy condition (QNEC), which is a statement purely about quantum field theory in a fixed background geometry, and there now exist proofs of the QNEC independent of any assumption about quantum focusing [18–20]. However, we found even when considering the generalized second law using crossed products, proving that the entropy increases beyond the leading semiclassical approximation is nontrivial, and could result in improved data-processing inequalities in quantum field theory. Therefore, it would be interesting to determine if a similar improvement could arise from the crossed product formulation of the QFC.

One of the main advantages of working with a Killing horizon in the present work was the ability to leverage the ANEC to prove the existence of a state with a geometric modular flow along the horizon. This suggests that a strategy for finding the necessary states conjectured in [33] to carry out the crossed product for generic subregions is to look for related energy conditions adapted to the subregion under consideration. One could look for positive operators involving a smeared stress tensor over the subregion horizon to try to establish the existence of a half-sided translation, which then would determine a vacuum state and modular operator whose flow is boost-like near the subregion entangling surface. It is possible that the novel energy conditions recently investigated in [130–132] could be helpful in establishing these positivity results. This may provide a crucial clue for how to rigorously establish that local gravitational algebras can be realized as a crossed product, and ultimately help in formulating the QFC in this language.

Acknowledgments

We thank Daine Danielson, Ben Freivogel, Stefan Hollands, Ted Jacobson, Nima Lashkari, Roberto Longo, Pranav Pulakkat, Gautam Satishchandran, Erik Verlinde, Bob Wald, Aron Wall, and Victor Zhang for helpful discussions. A.J.S. thanks the Isaac Newton Institute for Mathematical Sciences for support and hospitality during the programme “Black holes: bridges between number theory and holographic quantum information” where work on this paper was undertaken. This work was supported by EPSRC grant EP/R014604/1. This research is supported in part by the Air Force Office of Scientific Research under award number FA9550-19-1-036 and by the DOE award number DE-SC0015655.

A Tomita operators for crossed product algebra

The Tomita operators considered in section 2 for states of the form $|\widehat{\Phi}\rangle = |\Phi, f\rangle$ were crucial for obtaining density matrices and computing the perturbative expansion of entropies for the crossed product algebra in section 5. Here, we derive the formula for these Tomita operators, and additionally obtain expressions for the Tomita operators for a more general class of states obtained from arbitrary superpositions of the $|\Phi, f\rangle$ states. This simplifies the derivation of the density matrices for these states from [33], and additionally yields new results in the explicit expressions for the modular conjugations $J_{\widehat{\Phi}}$. This appendix assumes familiarity with Tomita-Takesaki theory; see [133], [33, Appendix C] for an introduction.

The crossed product of a QFT algebra \mathcal{A} by the modular automorphism group associated with the state $|\Omega\rangle \in \mathcal{H}_{\text{QFT}}$ is generated by the operators

$$\widehat{\mathcal{A}} = \langle e^{i\hat{p}h} \mathbf{a} e^{-i\hat{p}h}, \hat{q} \rangle \quad (\text{A.1})$$

acting on the Hilbert space $\widehat{\mathcal{H}} = \mathcal{H}_{\text{QFT}} \otimes L^2(\mathbb{R})$, where $h = h_{\Omega}$ is the modular Hamiltonian for the state $|\Omega\rangle$. For a given vector state $|\widehat{\Phi}\rangle \in \widehat{\mathcal{H}}$ for the crossed product algebra, the Tomita operator is defined to act as

$$S_{\widehat{\Phi}} \widehat{\mathbf{a}} |\widehat{\Phi}\rangle = \widehat{\mathbf{a}}^\dagger |\widehat{\Phi}\rangle. \quad (\text{A.2})$$

Hence, we can solve for $S_{\widehat{\Phi}}$ by determining the action of a linearly spanning set of operators $\widehat{\mathbf{a}} \in \widehat{\mathcal{A}}$ along with their conjugates $\widehat{\mathbf{a}}^\dagger$ on the state $|\widehat{\Phi}\rangle$. Considering the additive basis of operators

$$\widehat{\mathbf{a}} = e^{i\hat{p}h} \mathbf{a} e^{-i\hat{p}h} e^{iu\hat{q}}, \quad (\text{A.3})$$

we can compute their action on the classical-quantum states $|\widehat{\Phi}\rangle = |\Phi, f\rangle$ which were used in the investigations of the GSL in this paper.

Before doing this, it is helpful to note how the QFT Tomita operators act in the larger Hilbert space $\mathcal{H}_{\text{QFT}} \otimes L^2(\mathbb{R})$. This is somewhat subtle because the Tomita operators are antilinear, so there is a choice to make when defining its action on the larger Hilbert space $\widehat{\mathcal{H}}$. We will define any QFT Tomita operator S to act trivially on the position basis $|y\rangle$ of $L^2(\mathbb{R})$, i.e.

$$S|\Phi, y\rangle = |S\Phi, y\rangle \quad (\text{A.4})$$

This implies that S commutes with \hat{q} :

$$S\hat{q}|\Phi, y\rangle = Sy|\Phi, y\rangle = yS|\Phi, y\rangle = y|S\Phi, y\rangle = \hat{q}|S\Phi, y\rangle = \hat{q}S|\Phi, y\rangle. \quad (\text{A.5})$$

On the other hand, S must anticommute with \hat{p} :

$$\begin{aligned} S\hat{p}|\Phi, y\rangle &= S \int dy' \langle y' | \hat{p} | y \rangle |\Phi, y'\rangle = \int dy' \langle y | \hat{p} | y' \rangle S|\Phi, y'\rangle = - \int dy' \langle y' | \hat{p} | y \rangle |S\Phi, y'\rangle \\ &= -\hat{p}|S\Phi, y\rangle = -\hat{p}S|\Phi, y\rangle \end{aligned} \quad (\text{A.6})$$

where we have used that $\langle y | \hat{p} | y' \rangle = -\langle y' | \hat{p} | y \rangle$. We can remember these commutation relations by noting that \hat{q} acts as multiplication by a real number in the $|y\rangle$ basis, while \hat{p} is a derivative

times an imaginary number, $\hat{p} = -i \frac{d}{dq}$. These relations imply that S acts as a standard time-reversal transformation on the $L^2(\mathbb{R})$ Hilbert space, and hence sends a momentum eigenstate to its negative counterpart, $S|\Phi, s_p\rangle = |S\Phi, -s_p\rangle$. Finally, we note the action on the state $|\Phi, f\rangle$,

$$S|\Phi, f\rangle = \int dy S f(y) |\Phi, y\rangle = \int dy f^*(y) |S\Phi, y\rangle = |S\Phi, f^*\rangle \quad (\text{A.7})$$

These relations also hold for the antiunitary modular conjugations J .

We now compute¹²

$$\begin{aligned} \hat{\mathbf{a}}^\dagger|\widehat{\Phi}\rangle &= e^{-iu\hat{q}} e^{i\hat{p}h} \mathbf{a}^\dagger e^{-i\hat{p}h} |\Phi, f\rangle \\ &= e^{-iu\hat{q}} e^{i\hat{p}h} \mathbf{a}^\dagger e^{-i\hat{p}h} S_{\Phi|\Omega} S_\Omega |\Omega, f\rangle \\ &= S_{\Phi|\Omega} S_\Omega e^{-iu\hat{q}} e^{i\hat{p}h} \mathbf{a}^\dagger |\Omega, f\rangle \\ &= S_{\Phi|\Omega} S_\Omega e^{-iu\hat{q}} e^{i\hat{p}h} S_\Omega \mathbf{a} |\Omega, f^*\rangle \\ &= S_{\Phi|\Omega} e^{iu\hat{q}} e^{-i\hat{p}h} \mathbf{a} |\Omega, f^*\rangle \\ &= S_{\Phi|\Omega} \int ds \widetilde{f}^*(s) e^{-ish} \mathbf{a} e^{iu\hat{q}} |\Omega, s\rangle \\ &= S_{\Phi|\Omega} \int ds \widetilde{f}^*(s) e^{is(\hat{q}-h)} \mathbf{a} |\Omega, u\rangle \\ &= \sqrt{2\pi} S_{\Phi|\Omega} f^*(\hat{q}-h) \mathbf{a} |\Omega, u\rangle \end{aligned} \quad (\text{A.8})$$

In the third line we have used that $S_{\Phi|\Omega} S_\Omega$ is affiliated with $\mathcal{A}'_{\text{QFT}}$ to commute it to the left, and in the fifth line we used that $S_\Omega^2 = \mathbb{1}$ and $S_\Omega e^{i\hat{p}h} S_\Omega = e^{-i\hat{p}h}$. On the other hand, we have that

$$\begin{aligned} \hat{\mathbf{a}}|\widehat{\Phi}\rangle &= e^{i\hat{p}h} \mathbf{a} e^{-i\hat{p}h} e^{iu\hat{q}} |\Phi, f\rangle \\ &= e^{i\hat{p}h} \mathbf{a} e^{-i\hat{p}h} e^{iu\hat{q}} S_{\Phi|\Omega} S_\Omega |\Omega, f\rangle \\ &= S_{\Phi|\Omega} S_\Omega \int ds \tilde{f}(s) e^{i\hat{p}h} \mathbf{a} e^{-i\hat{p}h} e^{iu\hat{q}} |\Omega, s\rangle \\ &= S_{\Phi|\Omega} S_\Omega \int ds \tilde{f}(s) e^{i\hat{p}h} \mathbf{a} e^{is\hat{q}} |\Omega, u\rangle \\ &= \sqrt{2\pi} S_{\Phi|\Omega} S_\Omega e^{i\hat{p}h} f(\hat{q}) \mathbf{a} |\Omega, u\rangle. \end{aligned} \quad (\text{A.9})$$

These produce an equation for the Tomita operator via (A.2),

$$S_{\widehat{\Phi}} S_{\Phi|\Omega} S_\Omega e^{i\hat{p}h} f(\hat{q}) = S_{\Phi|\Omega} f^*(\hat{q}-h) \quad (\text{A.10})$$

whose solution is

$$S_{\widehat{\Phi}} = S_{\Phi|\Omega} f^*(\hat{q}-h) \frac{1}{f(\hat{q})} e^{-i\hat{p}h} S_\Omega S_{\Omega|\Phi}, \quad (\text{A.11})$$

¹²We follow the conventions of [33, Appendix E] for Fourier transforms. In particular, we have $|f\rangle = \int dy f(y) |y\rangle = \int ds \tilde{f}(s) |s\rangle$ and $\langle y|s\rangle = \frac{e^{isy}}{\sqrt{2\pi}}$.

noting that $S_\Omega^{-1} = S_\Omega$, $S_{\Phi|\Omega}^{-1} = S_{\Omega|\Phi}$. We can simplify this to the form

$$S_{\widehat{\Phi}} = J_{\Phi|\Omega} \Delta_{\Phi|\Omega}^{\frac{1}{2}} f^*(\hat{q} - h) e^{\frac{\hat{q}}{2}} e^{-i\hat{p}h} \frac{1}{e^{\frac{\hat{q}}{2}} f(\hat{q} + h)} \Delta_{\Omega}^{\frac{1}{2}} S_\Omega S_{\Omega|\Phi} \quad (\text{A.12})$$

$$= J_{\Phi|\Omega} e^{-i\hat{p}h} \cdot \left[e^{i\hat{p}h} \Delta_{\Phi|\Omega}^{\frac{1}{2}} f^*(\hat{q} - h) e^{\frac{\hat{q}}{2}} e^{-i\hat{p}h} \right] \cdot \left[\frac{1}{e^{\frac{\hat{q}}{2}} f(\hat{q} + h)} J_\Omega J_{\Omega|\Phi} \Delta_{\Omega|\Phi}^{\frac{1}{2}} \right] \quad (\text{A.13})$$

where the first term in brackets is affiliated with $\widehat{\mathcal{A}}$, and the second term is affiliated with $\widehat{\mathcal{A}}'$. Note by computing $S_{\widehat{\Phi}}^\dagger S_{\widehat{\Phi}}$, we immediately arrive at the expression for the density matrices derived in [33],

$$\rho_{\widehat{\Phi}} = e^{i\hat{p}h} f(\hat{q} - h) e^{\frac{\hat{q}}{2}} \Delta_{\Phi|\Omega} f^*(\hat{q} - h) e^{-i\hat{p}h} \quad (\text{A.14})$$

$$\rho'_{\widehat{\Phi}} = \Delta_{\Omega|\Phi}^{-\frac{1}{2}} J_{\Phi|\Omega} J_\Omega e^{\hat{q}} |f(\hat{q} + h)|^2 J_\Omega J_{\Omega|\Phi} \Delta_{\Omega|\Phi}^{-\frac{1}{2}}. \quad (\text{A.15})$$

It is clear from this construction that we can obtain expressions for the Tomita operator of arbitrary superpositions $|\widehat{\Phi}\rangle = \int d\lambda |\Phi_\lambda, f_\lambda\rangle$ as well, where $d\lambda$ is some measure for the wavefunctions indexed by λ , which could in principle include discrete δ -function components. In this case we have

$$\widehat{\mathfrak{a}}^\dagger \int d\lambda |\Phi_\lambda, f_\lambda\rangle = \sqrt{2\pi} \int d\lambda S_{\Phi_\lambda|\Omega} f_\lambda^*(\hat{q} - h) \mathfrak{a}|\Omega, u\rangle \quad (\text{A.16})$$

$$\widehat{\mathfrak{a}} \int d\lambda |\Phi_\lambda, f_\lambda\rangle = \sqrt{2\pi} \int d\lambda S_{\Phi_\lambda|\Omega} S_\Omega e^{i\hat{p}h} f_\lambda(\hat{q}) \mathfrak{a}|\Omega, u\rangle \quad (\text{A.17})$$

and hence

$$\begin{aligned} S_{\widehat{\Phi}} &= \left[\int d\lambda S_{\Phi_\lambda|\Omega} f_\lambda^*(\hat{q} - h) \right] \left[\int d\mu S_{\Phi_\mu|\Omega} S_\Omega e^{i\hat{p}h} f_\mu(\hat{q}) \right]^{-1} \\ &= J_\Omega e^{-i\hat{p}h} \left[\int d\lambda e^{i\hat{p}h} J_\Omega J_{\Phi_\lambda|\Omega} \Delta_{\Phi_\lambda|\Omega}^{\frac{1}{2}} f_\lambda^*(\hat{q} - h) e^{-i\hat{p}h} \right] \left[\int d\mu \Delta_{\Omega|\Phi_\mu}^{-\frac{1}{2}} J_{\Phi_\mu|\Omega} J_\Omega \Delta_{\Omega}^{\frac{1}{2}} f(\hat{q} + h) \right]^{-1} \\ &= J_\Omega e^{-i\hat{p}h} \left[\int d\lambda e^{i\hat{p}h} J_\Omega J_{\Phi_\lambda|\Omega} \Delta_{\Phi_\lambda|\Omega}^{\frac{1}{2}} f_\lambda^*(\hat{q} - h) e^{\frac{\hat{q}}{2}} e^{-i\hat{p}h} \right] \left[\int d\mu \Delta_{\Omega|\Phi_\mu}^{-\frac{1}{2}} J_{\Phi_\mu|\Omega} J_\Omega f(\hat{q} + h) e^{\frac{\hat{q}}{2}} \right]^{-1} \end{aligned} \quad (\text{A.18})$$

Writing it in this way makes it clear that the Tomita operator can always be expressed in the form

$$S_{\widehat{\Phi}} = J_\Omega e^{-i\hat{p}h} \widehat{\mathfrak{s}} \widehat{\mathfrak{s}}', \quad \widehat{\mathfrak{s}} \in \widehat{\mathcal{A}}, \widehat{\mathfrak{s}}' \in \widehat{\mathcal{A}}'. \quad (\text{A.19})$$

Using the polar decomposition $\widehat{\mathfrak{s}} = \widehat{\mathfrak{u}} \rho_{\widehat{\Phi}}^{\frac{1}{2}}$, with $\widehat{\mathfrak{u}}$ a unitary element of $\widehat{\mathcal{A}}$ and $\rho_{\widehat{\Phi}}$ the density matrix for $\widehat{\mathcal{A}}$, and similarly writing $\widehat{\mathfrak{s}}' = \widehat{\mathfrak{u}}' (\rho'_{\widehat{\Phi}})^{-\frac{1}{2}}$, we arrive at the expression for the modular operator and modular conjugation:

$$J_{\widehat{\Phi}} = J_\Omega e^{-i\hat{p}h} \widehat{\mathfrak{u}} \widehat{\mathfrak{u}}', \quad \Delta_{\widehat{\Phi}} = \rho_{\widehat{\Phi}} (\rho'_{\widehat{\Phi}})^{-1} \quad (\text{A.20})$$

It is useful to consider this expression for the vacuum classical-quantum states $|\widehat{\Omega}\rangle = |\Omega, f\rangle$. In that case, the Tomita operator simplifies to

$$S_{\widehat{\Omega}} = J_{\Omega} e^{-i\hat{p}h} \frac{f^*(\hat{q})}{f(\hat{q}+h)} \Delta_{\widehat{\Omega}}^{\frac{1}{2}} = J_{\Omega} e^{-i\hat{p}h} \left(\frac{f^*(\hat{q})f^*(\hat{q}+h)}{f(\hat{q})f(\hat{q}+h)} \right)^{\frac{1}{2}} \frac{|f(\hat{q})|}{|f(\hat{q}+h)|} \Delta_{\widehat{\Omega}}^{\frac{1}{2}} \quad (\text{A.21})$$

which is now in the form of a polar decomposition, allowing the modular conjugation to be read off

$$J_{\widehat{\Omega}} = J_{\Omega} e^{-i\hat{p}h} \left(\frac{f^*(\hat{q})f^*(\hat{q}+h)}{f(\hat{q})f(\hat{q}+h)} \right)^{\frac{1}{2}}, \quad \Delta_{\widehat{\Omega}} = \frac{g(\hat{q})}{g(\hat{q}+h)} \Delta_{\Omega}. \quad (\text{A.22})$$

This is particularly simple when the wavefunction is real, in which case $J_{\widehat{\Omega}} = J_{\Omega} e^{-i\hat{p}h}$.

It is also helpful to have the Tomita operator for the twirled state $|\tilde{\Phi}\rangle = e^{i\hat{p}h} |\Phi, f\rangle$. We proceed as before, computing

$$\begin{aligned} \widehat{\mathbf{a}}|\tilde{\Phi}\rangle &= e^{i\hat{p}h} \mathbf{a} e^{-i\hat{p}h} e^{iu\hat{q}} e^{i\hat{p}h} |\Phi, f\rangle \\ &= \int ds \tilde{f}(s) e^{i\hat{p}h} \mathbf{a} e^{-i\hat{p}h} e^{iu\hat{q}} e^{ish} |\Phi, s\rangle \\ &= \int ds \tilde{f}(s) e^{i\hat{p}h} \mathbf{a} e^{-i(\hat{p}-s)h} e^{is\hat{q}} |\Phi, u\rangle \\ &= \int ds \tilde{f}(s) e^{is(\hat{q}+h)} e^{i\hat{p}h} \mathbf{a} e^{-i\hat{p}h} S_{\Phi|\Omega} S_{\Omega} |\Omega, u\rangle \\ &= \sqrt{2\pi} f(\hat{q}+h) S_{\Phi|\Omega} S_{\Omega} e^{i\hat{p}h} \mathbf{a} |\Omega, u\rangle \end{aligned} \quad (\text{A.23})$$

and

$$\begin{aligned} \widehat{\mathbf{a}}^{\dagger}|\tilde{\Phi}\rangle &= e^{-iu\hat{q}} e^{i\hat{p}h} \mathbf{a}^{\dagger} e^{-i\hat{p}h} e^{i\hat{p}h} |\Phi, f\rangle \\ &= \int ds \tilde{f}(s) e^{ish} \mathbf{a}^{\dagger} e^{-iu\hat{q}} |\Phi, s\rangle \\ &= \int ds \tilde{f}(s) e^{ish} \mathbf{a}^{\dagger} e^{is\hat{q}} |\Phi, -u\rangle \\ &= \sqrt{2\pi} f(\hat{q}+h) \mathbf{a}^{\dagger} |\Phi, -u\rangle \\ &= \sqrt{2\pi} f(\hat{q}+h) S_{\Phi|\Omega} \mathbf{a} |\Omega, u\rangle. \end{aligned} \quad (\text{A.24})$$

Then using that $S_{\tilde{\Phi}} \widehat{\mathbf{a}}|\tilde{\Phi}\rangle = \widehat{\mathbf{a}}^{\dagger}|\tilde{\Phi}\rangle$, we find that

$$S_{\tilde{\Phi}} = f(\hat{q}+h) S_{\Phi|\Omega} e^{-i\hat{p}h} S_{\Omega} S_{\Omega|\Phi} \frac{1}{f(\hat{q}+h)} \quad (\text{A.25})$$

We can then simplify this into a factorized form,

$$\begin{aligned}
S_{\tilde{\Phi}} &= J_{\Omega} f^*(\hat{q} - h) J_{\Omega} J_{\Phi|\Omega} \Delta_{\Phi|\Omega}^{\frac{1}{2}} e^{-i\hat{p}h} \Delta_{\Omega}^{-\frac{1}{2}} J_{\Omega} J_{\Omega|\Phi} \Delta_{\Omega|\Phi}^{\frac{1}{2}} \frac{1}{f(\hat{q} + h)} \\
&= J_{\Omega} e^{-i\hat{p}h} \left[f^*(\hat{q}) e^{-i\hat{p}h} J_{\Omega} J_{\Phi|\Omega} \Delta_{\Phi|\Omega}^{\frac{1}{2}} \Delta_{\Omega}^{-\frac{1}{2}} e^{-i\hat{p}h} e^{\frac{\hat{q}}{2}} \right] J_{\Omega} J_{\Omega|\Phi} \Delta_{\Omega|\Phi}^{\frac{1}{2}} \frac{e^{-\frac{\hat{q}}{2}}}{f(\hat{q} + h)} \\
&= J_{\Omega} e^{-i\hat{p}h} J_{\Omega} J_{\Omega|\Phi} \left[f^*(\hat{q}) e^{-i\hat{p}h} J_{\Omega} J_{\Phi|\Omega} \Delta_{\Phi|\Omega}^{\frac{1}{2}} e^{\frac{\hat{q}}{2}} e^{-i\hat{p}h} \right] \Delta_{\Omega|\Phi}^{\frac{1}{2}} \frac{e^{-\frac{\hat{q}}{2}}}{f(\hat{q} + h)} \\
&= e^{i\hat{p}h} J_{\Omega|\Phi} \left[f^*(\hat{q}) e^{-i\hat{p}h} J_{\Omega} J_{\Phi|\Omega} \Delta_{\Phi|\Omega}^{\frac{1}{2}} e^{\frac{\hat{q}}{2}} e^{-i\hat{p}h} \right] \cdot \left[\Delta_{\Omega|\Phi}^{\frac{1}{2}} \frac{e^{-\frac{\hat{q}}{2}}}{f(\hat{q} + h)} \right] \tag{A.26}
\end{aligned}$$

so that in particular the density matrices are given by,

$$\rho_{\tilde{\Phi}} = e^{i\hat{p}h} \Delta_{\Phi|\Omega}^{\frac{1}{2}} J_{\Omega|\Phi} J_{\Omega} e^{\hat{q}} g(\hat{q} - h) J_{\Omega} J_{\Phi|\Omega} \Delta_{\Phi|\Omega}^{\frac{1}{2}} e^{-i\hat{p}h} \tag{A.27}$$

$$\rho'_{\tilde{\Phi}} = f(\hat{q} + h) e^{\frac{\hat{q}}{2}} \Delta_{\Omega|\Phi}^{-1} e^{\frac{\hat{q}}{2}} f^*(\hat{q} + h) \tag{A.28}$$

again reproducing the expressions from [33].

B Null surfaces

The identities considered in section 3.2 that are crucial in obtaining the crossed product gravitational algebras and relating the crossed product entropy to generalized entropy make extensive use of geometric relations for null surfaces. This appendix summarizes the necessary properties of null surfaces needed to arrive at these identities by way of the canonical analysis on null surfaces, discussed in appendix C.

B.1 Null surface geometry

We begin by briefly reviewing the geometry of null surfaces, see [68, 134, 135] for additional details. Let \mathcal{N} be a null hypersurface with a null normal l_a whose norm vanishes on \mathcal{N} , $l \cdot l \stackrel{\mathcal{N}}{=} 0$. We can define a null rigging vector n^a that satisfies $n \cdot l = -1$ and $n \cdot n = 0$, and with it define a projector onto the null surface

$$\Pi^a_b = \delta^a_b + n^a l_b. \tag{B.1}$$

The metric at \mathcal{N} then naturally decomposes as

$$g_{ab} = q_{ab} + n_a l_b + l_a n_b, \tag{B.2}$$

where q_{ab} is the degenerate induced metric on \mathcal{N} . Additionally, the spacetime volume form ϵ decomposes as

$$\epsilon = -l \wedge \eta \tag{B.3}$$

where

$$\eta = -n \wedge \mu \quad (\text{B.4})$$

is the volume form on \mathcal{N} , and $\mu = i_l \eta$ is the degenerate area form that induces a volume form on horizon cuts. Note that the choice of rigging vector n^a also allows for an inverse degenerate metric q^{ab} to be defined, which satisfies $q^{ab}n_b = q^{abl}_b = 0$, $q^{ab}q_{ac}q_{bd} = q_{cd}$.

The shape operator W^a_b for the null surface is defined as the projection of $\nabla_a l^b$ onto the null surface, and decomposes as

$$W^a_b = \Pi^c_b \nabla_c l^a = -kn_b l^a + \varpi_b l^a + q^{ac}(\sigma_{cb} + \frac{1}{d-2}\Theta q_{cb}) \quad (\text{B.5})$$

where k is the inaffinity, ϖ_b is the Hájiček one-form, σ_{cb} is the shear, and Θ is the expansion, defined by

$$k = -n_b l^a \nabla_a l^b \quad (\text{B.6})$$

$$\varpi_b = -q^c_b n_a \nabla_c l^a \quad (\text{B.7})$$

$$\Theta = \nabla_a l^a - k \quad (\text{B.8})$$

$$\sigma_{cb} = \frac{1}{2} \mathcal{L}_l q_{cb} - \frac{1}{d-2} \Theta q_{cb} \quad (\text{B.9})$$

B.2 Gaussian null coordinates

Gaussian null coordinates (GNuC) [75, 106, 136] provide a convenient coordinate system for describing the metric near a null hypersurface \mathcal{N} . On such a surface, we take l^a to be an affinely parameterized null generator satisfying $l^a \nabla_a l^b \stackrel{\mathcal{N}}{=} 0$, and let v denote the affine parameter so that $l^a \nabla_a v = 1$. Surfaces of constant v then define a foliation of the null hypersurface, and we can define a future-directed transverse null rigging vector n^a orthogonal to these surfaces and normalized such that $n \cdot l = -1$. Extending n^a off of \mathcal{N} as an affinely parameterized null geodesic generator, it can be chosen to satisfy $n \cdot n = 0$, $n^a \nabla_a n^b = 0$, and $\nabla_{[a} n_{b]} = 0$ throughout the open coordinate patch near \mathcal{N} . We can furthermore extend l^a off of \mathcal{N} by requiring $[n, l]^a = 0$. By taking u to be an affine parameter for n^a with \mathcal{N} at $u = 0$, coordinates (u, v, y^A) can be chosen such that

$$l^a = \left(\frac{\partial}{\partial v} \right)^a, \quad n^a = \left(\frac{\partial}{\partial u} \right)^a. \quad (\text{B.10})$$

The metric in this coordinate system takes the form

$$ds^2 = -2dudv - u^2 Adv^2 + 2uB_A dv dy^A + q_{AB} dy^A dy^B \quad (\text{B.11})$$

where A , B_A and q_{AB} are functions of the spacetime coordinates. Note in this coordinate system the relations

$$n_a = -\nabla_a v, \quad l_a = -\nabla_a u + uB_a - u^2 A \nabla_a v, \quad l \cdot l = -u^2 A \quad (\text{B.12})$$

hold, where $B_a = B_A \nabla_a y^A$.

When \mathcal{N} is a Killing horizon with bifurcation surface located at $u = v = 0$, the Killing vector in these coordinates takes the form [75]

$$\xi^a = \kappa v l^a - \kappa u n^a. \quad (\text{B.13})$$

Additionally, requiring the metric to be invariant under the flow of ξ^a leads to the condition that A , B_A , and q_{AB} depend on the coordinates u and v only via their product uv .

B.3 Quasi-Killing vectors

In addition to the global Killing vector ξ^a , the analysis of section 3.2 deals extensively with quasi-Killing vectors, which satisfy Killing's equation only on the null surface \mathcal{N} . A generic such vector parallel to the horizon generator l^a can be expressed in terms of a function $\tau(v, y^A)$ as

$$\zeta_\tau^a = \tau l^a + u \nabla^a \tau. \quad (\text{B.14})$$

The Lie derivative of the metric on \mathcal{N} can then be written as

$$\mathcal{L}_{\zeta_\tau} g_{ab} \stackrel{\mathcal{N}}{=} \tau \mathcal{L}_l g_{ab} + 2 \nabla_{(a} \tau l_{b)} + 2 \nabla_{(a} u \nabla_{b)} \tau \stackrel{\mathcal{N}}{=} \tau \mathcal{L}_l g_{ab}, \quad (\text{B.15})$$

where we have dropped terms proportional to u . Hence we need only verify that $\mathcal{L}_l g_{ab} = 0$. Note using the GNuC expressions (B.11), one finds

$$\mathcal{L}_l l_a = (i_l dl)_a + \nabla_a (l \cdot l) \stackrel{\mathcal{N}}{=} 0 \quad (\text{B.16})$$

$$\mathcal{L}_l n_a = (i_l dn)_a + \nabla_a (l \cdot n) = 0. \quad (\text{B.17})$$

Then from the decomposition of the metric $g_{ab} = q_{ab} + n_a l_b + l_a n_b$, we find that

$$\mathcal{L}_l g_{ab} = \mathcal{L}_l q_{ab} = \sigma_{ab} + \frac{1}{d-2} \Theta q_{ab} = 0, \quad (\text{B.18})$$

where the last equality uses that a Killing horizon has vanishing expansion and shear. This then verifies that ζ_τ^a is a quasi-Killing vector.

An additional restriction on the quasi-Killing vector is that it preserve the gauge conditions (C.7) employed in the canonical analysis for the null surface. The third condition there is nontrivial, since it involves derivatives transverse to the null surface. To check whether the vector ζ_τ^a preserves this condition, we evaluate

$$\begin{aligned} \frac{1}{2} n^a \nabla_a (l^b l^c \mathcal{L}_{\zeta_\tau} g_{bc}) &= n^a \nabla_a \left(l^b l^c \nabla_b (\tau l_c + u \nabla_c \tau) \right) \\ &= n^a \nabla_a \left(l^b \nabla_b \tau (l \cdot l) + \tau l^b l^c \nabla_b l_c + l^b \nabla_b u l^c \nabla_c \tau + u l^b l^c \nabla_b \nabla_c \tau \right) \\ &= l^b l^c \nabla_b \nabla_c \tau + \mathcal{O}(u) = \mathcal{L}_l \mathcal{L}_l \tau + \mathcal{O}(u). \end{aligned} \quad (\text{B.19})$$

For this to vanish, the function τ must be at most linear in the coordinate v . This agrees with the set of null surface symmetries derived in [67].

It is also useful to derive expressions involving the derivatives of the vectors ζ_τ^a , which can be applied when evaluating gravitational charges in appendix C. These expressions only rely on the form of the vector (B.14) in GNuC, and do not rely on \mathcal{N} being a Killing horizon. We have

$$l^a \nabla_a \zeta_\tau^b = l^b l^a \nabla_a \tau + \tau l^a \nabla_a l^b + l^a \nabla_a u \nabla^b \tau + u l^a \nabla_a \nabla^b \tau \stackrel{\mathcal{N}}{=} l^b l^a \nabla_a \tau \quad (\text{B.20})$$

$$l_b \nabla_a \zeta_\tau^b = (l \cdot l) \nabla_a \tau + \frac{1}{2} \tau \nabla_a (l \cdot l) + \nabla_a u l^b \nabla_b \tau + u l_b \nabla_a \nabla^b \tau \stackrel{\mathcal{N}}{=} -l_a l^b \nabla_b \tau \quad (\text{B.21})$$

$$\nabla_a \zeta_\tau^a = l^a \nabla_a \tau + \tau \Theta + \nabla_a u \nabla^a \tau + u \nabla_a \nabla^a \tau \stackrel{\mathcal{N}}{=} \tau \Theta. \quad (\text{B.22})$$

C Canonical formalism and gravitational charges

In this appendix, we provide additional details on the canonical formulation of general relativity on null hypersurfaces needed to arrive at the identities discussed in section 3.2. We will largely follow the conventions and notation of [68], see also [67, 69–73] for related recent treatments of the canonical analysis of general relativity on null hypersurfaces. This canonical analysis is done in the framework of the covariant phase space, see [137–140] for overviews of this formalism.

The Lagrangian for the theory consists of a sum of a gravitational and matter contributions, $L = L^g + L^\psi$, with the gravitational Lagrangian comprised of an Einstein-Hilbert term and cosmological constant,

$$L^g = \frac{1}{16\pi G_N} \epsilon (R - 2\Lambda). \quad (\text{C.1})$$

The covariant symplectic potential $\theta = \theta^g + \theta^\psi$ arises from the boundary term appearing in the variation of the Lagrangian, $\delta L = E_\phi \cdot \delta \phi + d\theta$, where $\phi = (g_{ab}, \psi)$ collectively denotes the full set of dynamical fields, and $E_\phi = 0$ are the equations of motion. The gravitational piece is given by

$$\theta^g = \frac{1}{16\pi G_N} \epsilon_{a\dots} \left(g^{bc} \delta \Gamma_{bc}^a - g^{ac} \delta \Gamma_{bc}^b \right). \quad (\text{C.2})$$

When pulled back to a null Cauchy surface \mathcal{N} , θ^g may be decomposed as [68]

$$\underline{\theta^g} \stackrel{\mathcal{N}}{=} -\delta \ell + d\beta + \mathcal{E}^g \quad (\text{C.3})$$

where the underline denotes the pullback, and

$$\ell = -\frac{1}{16\pi G_N} \eta (\Theta + 2k) \quad (\text{C.4})$$

$$\beta = \frac{1}{32\pi G_N} g^{ab} \delta g_{ab\mu} \quad (\text{C.5})$$

$$\mathcal{E}^g = \frac{1}{16\pi G_N} \eta \left[\sigma^{ab} \delta q_{ab} - \left(k + \frac{d-3}{d-2} \Theta \right) q^{ab} \delta q_{ab} - 2(\Theta n_a + \varpi_a) \delta l^a \right]. \quad (\text{C.6})$$

For the present applications, it is helpful to perform a partial gauge-fixing with respect to diffeomorphisms tangent to \mathcal{N} . The relevant gauge-conditions were considered by Chandrasekaran, Flanagan, and Prabhu (CFP) [67], who showed that one can always impose the conditions that l_a and l^a are fixed, l^a remains null, and the inaffinity of l^a vanishes, $k = 0$. These translate to the following conditions on the variations:

$$\delta l_a \stackrel{\mathcal{N}}{=} 0; \quad g_{ac}\delta l^c = -l^b\delta g_{ab} \stackrel{\mathcal{N}}{=} 0; \quad \delta k = \frac{1}{2}n^b\nabla_b(l^a l^c \delta g_{ac}) \stackrel{\mathcal{N}}{=} 0. \quad (\text{C.7})$$

Note that we can satisfy these conditions by gauge fixing to Gaussian null coordinates and imposing that the only variations of the metric come from variations of A , B_A and q_{AB} appearing in (B.11). With these gauge conditions, the terms appearing in the decomposition (C.3) simplify to

$$\ell = -\frac{1}{16\pi G_N}\eta\Theta \quad (\text{C.8})$$

$$\beta = \frac{1}{32\pi G_N}q^{ab}\delta g_{ab\mu} \quad (\text{C.9})$$

$$\mathcal{E}^g = \frac{1}{16\pi G_N}\eta \left[\sigma^{ab}\delta q_{ab} - \frac{d-3}{d-2}\Theta q^{ab}\delta q_{ab} \right]. \quad (\text{C.10})$$

An additional consequence of the above gauge fixing is that it allows us to define the location of a horizon cut in a diffeomorphism invariant manner: in Gaussian null coordinates, a surface at $v = \lambda(y^A)$ can simply be interpreted as a cut located a given affine distance away from the bifurcation surface.

The quantity $\mathcal{E} = \mathcal{E}^g + \mathcal{E}^\psi$ serves as a symplectic potential current on the null surface, and hence the integral of its variation defines the symplectic form

$$\Omega = \int_{\mathcal{N}} \delta\mathcal{E}, \quad (\text{C.11})$$

where we are treating δ as an exterior derivative on field space, so that in terms of individual variations, $\delta\mathcal{E}[\delta_1\phi, \delta_2\phi] = \delta_1\mathcal{E}[\delta_2\phi] - \delta_2\mathcal{E}[\delta_1\phi]$. Note that $\delta\mathcal{E}$ differs from the symplectic form constructed from the spacetime-covariant current $\omega = \delta\theta$ by a total derivative $-d\delta\beta$. We can determine the charge associated with a diffeomorphism generated by a vector ξ^a parallel to \mathcal{N} by evaluating [68, 69]

$$\Omega[\delta\phi, \mathcal{L}_\xi\phi] = \int_{\mathcal{N}} d(\delta M_\xi - M_{\delta\xi} - i_\xi\mathcal{E}) + \int_{\mathcal{N}} (\delta C_\xi - C_{\delta\xi}) \quad (\text{C.12})$$

where

$$M_\xi = Q_\xi + i_\xi\ell - \beta[\mathcal{L}_\xi\phi], \quad (\text{C.13})$$

Q_ξ is the covariant Noether potential, which for general relativity evaluates to [139]

$$Q_\xi = -\frac{1}{16\pi G_N}\epsilon^a{}_{b\dots}\nabla_a\xi^b, \quad (\text{C.14})$$

and C_ξ are combinations of the equations of motion comprising the constraints of the theory.

To satisfy Hamilton's equation, the right hand side of (C.12) must take the form of a total variation. The possible obstructions arise either from field-dependence of the generator ξ^a in the terms $M_{\delta\xi}$ and $C_{\delta\xi}$ or from the flux term $i_\xi\mathcal{E}$ evaluated at the asymptotic boundaries of the null surface $\partial\mathcal{N}$. We verify below that the field dependence for quasi-Killing vectors on a null surface has vanishing contributions to the fluxes. The remaining obstruction to integrability comes from $i_\xi\mathcal{E}$. To handle this term, we impose boundary conditions on the future horizon that the matter field fluxes go to zero and that the shear is asymptotically vanishing. Additionally, the requirement that \mathcal{N} is an event horizon imposes that the expansion vanishes in the far future, so examining the form of \mathcal{E}^g in (C.10) shows that it vanishes on the future boundary, and hence the full obstruction to integrability vanishes here.

Integrability at the past boundary of \mathcal{N} is more subtle since Θ is constrained to satisfy (3.30), and hence cannot vanish at both the past and future boundaries of \mathcal{N} unless there is zero null energy flux. However, when working perturbatively in $\varkappa = \sqrt{32\pi G_N}$, the expansion itself is $\mathcal{O}(\varkappa^2)$, and since the metric variation δq_{ab} is $\mathcal{O}(\varkappa)$, we find that the flux term coming from the expansion in \mathcal{E}^g is suppressed. Thus, in the $G_N \rightarrow 0$ limit, the expansion term does not obstruct integrability of the charges, and assuming the shear and matter flux fall off at the past boundary of \mathcal{N} , the full contribution from $i_\xi\mathcal{E}$ vanishes there.

Hence, we can define the Hamiltonian

$$H_\xi = \int_{\partial\mathcal{N}} M_\xi + \int_{\mathcal{N}} C_\xi, \quad (\text{C.15})$$

which, according to the discussion of integrability above, satisfies

$$\delta H_\xi = \Omega[\delta\phi, \mathcal{L}_\xi\phi]. \quad (\text{C.16})$$

The Hamiltonian can also be usefully expressed in terms of a local integral on \mathcal{N} by noting the relation

$$dM_\xi + C_\xi \stackrel{\mathcal{N}}{=} \mathcal{E}[\mathcal{L}_\xi\phi], \quad (\text{C.17})$$

which holds when all forms are pulled back to \mathcal{N} . In general this equation contains corrections if the vector ξ^a is not tangent to \mathcal{N} or does not act covariantly on ℓ , as discussed in detail in [68, 134], but one can show that there is no contribution from the anomalous transformation of ℓ when the vector field ξ^a preserves the gauge-fixing conditions (C.7). According to equation (B.19), this will hold in particular for quasi-Killing vectors ζ_τ^a , with $\tau(v, y^A)$ containing only linear and constant terms in v . This results in the equivalent expression for the Hamiltonian

$$H_\xi = \int_{\mathcal{N}} \mathcal{E}[\mathcal{L}_\xi\phi]. \quad (\text{C.18})$$

The gravitational contribution to H_ξ follows directly from the form of \mathcal{E}^g given in (C.10). The matter contribution in the simplest cases is directly related to the matter stress tensor. To arrive at this conclusion, we assume that the matter symplectic potential θ^ψ does not contain nontrivial terms involving the metric in its decomposition on \mathcal{N} , and hence $\theta^\psi = \mathcal{E}^\psi$. We can then consider the matter contribution to the Noether current

$$J_\xi^\psi = \theta^\psi[\mathcal{L}_\xi\phi] - i_\xi L \stackrel{\mathcal{N}}{=} \mathcal{E}^\psi[\mathcal{L}_\xi\phi] \quad (\text{C.19})$$

where the second equality holds upon pulling back to \mathcal{N} and using that ξ^a is tangent to \mathcal{N} . The divergence of the Noether current satisfies

$$dJ_\xi^\psi = -E_\psi \cdot \mathcal{L}_\xi \psi - \epsilon T_{(\psi)}^{ab} \nabla_a \xi_b = dC_\xi^\psi + N_\xi^\psi \quad (\text{C.20})$$

where both C_ξ^ψ and N_ξ^ψ depend algebraically on ξ^a . The Noether identities $N_\xi^\psi = 0$ hold identically due to diffeomorphism invariance, and hence $J_\xi^\psi = C_\xi^\psi + dQ_\xi^\psi$. Again, for most standard matter theories, $Q_\xi^\psi = 0$, and hence the Noether current agrees with the contribution of the matter terms to the constraint. Examining (C.20), we see that

$$C_\xi^\psi = -\epsilon_{a\dots}(T_{(\psi)})^a{}_b \xi^b + (E_\psi \text{ terms}). \quad (\text{C.21})$$

Together with (C.19) and (C.18), this shows that the matter contribution to the full Hamiltonian is given simply by the integral of the matter stress tensor, up to terms that vanish on shell. Note that for the most general matter theories, one must allow for nonzero values of $\delta\ell^\psi$ and β^ψ in the decomposition of θ^ψ , and also allow for a nonzero value of Q_ξ^ψ . These can be straightforwardly incorporated into the above discussion, but for the remainder of this work we do not consider these generalizations.

We now would like to evaluate the gravitational charge potential M_{ζ_τ} for the vectors defined by (B.14), which are quasi-Killing vectors of the background spacetime. Using that $\epsilon = l \wedge n \wedge \mu$ and the identities (B.20) and (B.21), the Noether potential (C.14) pulled back to \mathcal{N} evaluates to

$$Q_{\zeta_\tau} \stackrel{\mathcal{N}}{=} -\frac{1}{16\pi G_N} (l^a n_b - n^a l_b) \nabla_a \zeta_\tau^b \mu = \frac{1}{8\pi G_N} \mu l^a \nabla_a \tau. \quad (\text{C.22})$$

Additionally, from the definitions (C.8) and (C.9) and the identity (B.22), the remaining terms in M_{ζ_τ} are given by

$$i_{\zeta_\tau} \ell \stackrel{\mathcal{N}}{=} -\frac{1}{16\pi G_N} i_{\zeta_\tau} \eta \Theta = -\frac{1}{16\pi G_N} \mu \tau \Theta \quad (\text{C.23})$$

$$-\beta [\mathcal{L}_{\zeta_\tau} g_{ab}] \stackrel{\mathcal{N}}{=} -\frac{1}{16\pi G_N} \nabla_a \zeta_\tau^a \mu = -\frac{1}{16\pi G_N} \mu \tau \Theta. \quad (\text{C.24})$$

This results in the expression

$$M_{\zeta_\tau} \stackrel{\mathcal{N}}{=} \frac{1}{8\pi G_N} \mu \left(l^a \nabla_a \tau - \tau \Theta \right), \quad (\text{C.25})$$

consistent with the expressions in CFP [67].

Finally, we must also verify that the field-dependence in the vector ζ_τ^a does not spoil integrability of the charge. Given the expression (B.14) for the quasi-Killing vector, we require that $\delta\tau = 0$, and by gauge fixing to Gaussian null coordinates we can ensure that $\delta l^a = 0$ and $\delta u = 0$ in a neighborhood of \mathcal{N} . This leads to

$$\delta \zeta_\tau^a = -u \delta g_{bc} g^{ba} \nabla^c \tau. \quad (\text{C.26})$$

Hence $\delta\zeta^a$ vanishes on \mathcal{N} , and its covariant derivative there does as well, since

$$\nabla_a \delta\zeta^b \stackrel{\mathcal{N}}{=} -\nabla_a u \delta g_{bc} g^{ab} \nabla^c \tau = l^b \delta g_{bc} \nabla^c \tau = 0, \quad (\text{C.27})$$

where we have used (B.12) and (C.7) evaluated at \mathcal{N} . Since the possible obstructions to integrability $M_{\delta\zeta_\tau}$ and $C_{\delta\zeta_\tau}$ appearing in (C.12) involve only $\delta\zeta^a$ and $\nabla_a \delta\zeta^b$ at \mathcal{N} , we see that these terms vanish.

References

- [1] J. D. Bekenstein, *Black holes and the second law*, *Lett. Nuovo Cimento* **4** (1972) 737–740.
- [2] J. D. Bekenstein, *Black Holes and Entropy*, *Phys. Rev. D* **7** (1973) 2333–2346.
- [3] R. D. Sorkin, *1983 paper on entanglement entropy: “On the Entropy of the Vacuum outside a Horizon”*, [[arXiv:1402.3589](#)].
- [4] L. Bombelli, R. K. Koul, J. Lee and R. D. Sorkin, *Quantum source of entropy for black holes*, *Phys. Rev. D* **34** (1986) 373–383.
- [5] M. Srednicki, *Entropy and area*, *Physical Review Letters* **71** (1993) 666–669.
- [6] V. Frolov and I. Novikov, *Dynamical origin of the entropy of a black hole*, *Physical Review D* **48** (1993) 4545–4551.
- [7] L. Susskind and J. Uglum, *Black hole entropy in canonical quantum gravity and superstring theory*, *Phys. Rev. D* **50** (1994) 2700–2711. [[arXiv:hep-th/9401070](#)].
- [8] T. Jacobson, *Black hole entropy and induced gravity*, [[arXiv:gr-qc/9404039](#)].
- [9] F. Larsen and F. Wilczek, *Renormalization of black hole entropy and of the gravitational coupling constant*, *Nucl. Phys. B* **458** (1996) 249–266. [[arXiv:hep-th/9506066](#)].
- [10] S. N. Solodukhin, *Entanglement entropy of black holes*, *Living Rev. Rel.* **14** (2011) 8. [[arXiv:1104.3712](#)].
- [11] J. H. Cooperman and M. A. Luty, *Renormalization of Entanglement Entropy and the Gravitational Effective Action*, *JHEP* **12** (2014) 045. [[arXiv:1302.1878](#)].
- [12] R. Bousso, Z. Fisher, S. Leichenauer and A. C. Wall, *Quantum focusing conjecture*, *Phys. Rev. D* **93** (2016) 064044. [[arXiv:1506.02669](#)].
- [13] E. Bianchi and R. C. Myers, *On the Architecture of Spacetime Geometry*, *Class. Quant. Grav.* **31** (2014) 214002. [[arXiv:1212.5183](#)].

- [14] R. C. Myers, R. Pourhasan and M. Smolkin, *On Spacetime Entanglement*, *JHEP* **06** (2013) 013. [[arXiv:1304.2030](#)].
- [15] S. W. Hawking, *Gravitational Radiation from Colliding Black Holes*, *Phys. Rev. Lett.* **26** (1971) 1344–1346.
- [16] S. W. Hawking, *Black holes in general relativity*, *Commun. Math. Phys.* **25** (1972) 152–166.
- [17] A. C. Wall, *A Survey of Black Hole Thermodynamics*, [[arXiv:1804.10610](#)].
- [18] R. Bousso, Z. Fisher, J. Koeller, S. Leichenauer and A. C. Wall, *Proof of the Quantum Null Energy Condition*, *Phys. Rev. D* **93** (2016) 024017. [[arXiv:1509.02542](#)].
- [19] S. Balakrishnan, T. Faulkner, Z. U. Khandker and H. Wang, *A General Proof of the Quantum Null Energy Condition*, *JHEP* **09** (2019) 020. [[arXiv:1706.09432](#)].
- [20] F. Ceyhan and T. Faulkner, *Recovering the QNEC from the ANEC*, *Commun. Math. Phys.* **377** (2020) 999–1045. [[arXiv:1812.04683](#)].
- [21] S. Ryu and T. Takayanagi, *Aspects of holographic entanglement entropy*, *J. High Energy Phys.* **2006** (2006) 045.
- [22] S. Ryu and T. Takayanagi, *Holographic derivation of entanglement entropy from AdS/CFT*, *Phys. Rev. Lett.* **96** (2006) 181602. [[arXiv:hep-th/0603001](#)].
- [23] V. E. Hubeny, M. Rangamani and T. Takayanagi, *A Covariant holographic entanglement entropy proposal*, *JHEP* **07** (2007) 062. [[arXiv:0705.0016](#)].
- [24] A. Lewkowycz and J. Maldacena, *Generalized gravitational entropy*, *JHEP* **08** (2013) 090. [[arXiv:1304.4926](#)].
- [25] T. Faulkner, A. Lewkowycz and J. Maldacena, *Quantum corrections to holographic entanglement entropy*, *JHEP* **11** (2013) 074. [[arXiv:1307.2892](#)].
- [26] N. Engelhardt and A. C. Wall, *Quantum Extremal Surfaces: Holographic Entanglement Entropy beyond the Classical Regime*, *JHEP* **01** (2015) 073. [[arXiv:1408.3203](#)].
- [27] S. Leutheusser and H. Liu, *Causal connectability between quantum systems and the black hole interior in holographic duality*, *Phys. Rev. D* **108** (2023) 086019. [[arXiv:2110.05497](#)].
- [28] S. Leutheusser and H. Liu, *Emergent Times in Holographic Duality*, *Phys. Rev. D* **108** (2023) 086020. [[arXiv:2112.12156](#)].
- [29] E. Witten, *Gravity and the crossed product*, *JHEP* **10** (2022) 008. [[arXiv:2112.12828](#)].

- [30] V. Chandrasekaran, G. Penington and E. Witten, *Large N algebras and generalized entropy*, *JHEP* **04** (2023) 009. [[arXiv:2209.10454](#)].
- [31] V. Chandrasekaran, R. Longo, G. Penington and E. Witten, *An algebra of observables for de Sitter space*, *JHEP* **02** (2023) 082. [[arXiv:2206.10780](#)].
- [32] J. Kudler-Flam, S. Leutheusser and G. Satishchandran, *Generalized Black Hole Entropy is von Neumann Entropy*, [[arXiv:2309.15897](#)].
- [33] K. Jensen, J. Sorce and A. J. Speranza, *Generalized entropy for general subregions in quantum gravity*, *JHEP* **12** (2023) 020. [[arXiv:2306.01837](#)].
- [34] S. Leutheusser and H. Liu, *Subalgebra-subregion duality: emergence of space and time in holography*, [[arXiv:2212.13266](#)].
- [35] E. Bahiru, A. Belin, K. Papadodimas, G. Sarosi and N. Vardian, *State-dressed local operators in AdS/CFT*, [[arXiv:2209.06845](#)].
- [36] E. Bahiru, A. Belin, K. Papadodimas, G. Sarosi and N. Vardian, *Holography and Localization of Information in Quantum Gravity*, [[arXiv:2301.08753](#)].
- [37] T. Faulkner and M. Li, *Asymptotically isometric codes for holography*, [[arXiv:2211.12439](#)].
- [38] A. C. Wall, *A proof of the generalized second law for rapidly changing fields and arbitrary horizon slices*, *Phys. Rev. D* **85** (2012) 104049. [[arXiv:1105.3445](#)].
- [39] H. Casini, E. Teste and G. Torroba, *Modular Hamiltonians on the null plane and the Markov property of the vacuum state*, *J. Phys. A* **50** (2017) 364001. [[arXiv:1703.10656](#)].
- [40] E. Witten, *Princeton physics 539, lecture 19*, November 17, 2022. <https://phy.princeton.edu/academics/graduate-program/graduate-course-recordings>.
- [41] H. J. Borchers, *The CPT-theorem in two-dimensional theories of local observables*, *Communications in Mathematical Physics* **143** (1992) 315–332.
- [42] H.-J. Borchers, *On revolutionizing quantum field theory with Tomita’s modular theory*, *Journal of Mathematical Physics* **41** (2000) 3604–3673.
- [43] H. W. Wiesbrock, *Half sided modular inclusions of von Neumann algebras*, *Commun. Math. Phys.* **157** (1993) 83–92.
- [44] U. Haagerup, *Operator-valued weights in von Neumann algebras. I*, *J. Functional Analysis* **32** (1979) 175–206.
- [45] U. Haagerup, *Operator-valued weights in von Neumann algebras. II*, *J. Functional Analysis* **33** (1979) 339–361.

- [46] M. Ali and V. Suneeta, *A local Generalized second law in crossed product constructions*, [[arXiv:2404.00718](#)].
- [47] K. Fredenhagen, *On the modular structure of local algebras of observables*, *Quantum Field Theory: A Selection of Papers in Memoriam Kurt Symanzik* (1985) 79–89.
- [48] D. Buchholz and R. Verch, *Scaling algebras and renormalization group in algebraic quantum field theory*, *Reviews in Mathematical Physics* **7** (1995) 1195–1239.
- [49] R. Haag, *Local quantum physics: Fields, particles, algebras*. Springer-Verlag, Berlin, 1992.
- [50] R. Bousso, H. Casini, Z. Fisher and J. Maldacena, *Entropy on a null surface for interacting quantum field theories and the Bousso bound*, *Phys. Rev. D* **91** (2015) 084030. [[arXiv:1406.4545](#)].
- [51] K. Schmüdgen, *Unbounded Self-adjoint Operators on Hilbert Space*. Graduate Texts in Mathematics. Springer, Dordrecht, 2012.
- [52] R. Haag, *Bemerkungen zum nahwirkungsprinzip in der quantenphysik*, *Annalen der Physik* **466** (1963) 29–34.
- [53] K. Fredenhagen and J. Hertel, *Local algebras of observables and pointlike localized fields*, *Communications in Mathematical Physics* **80** (1981) 555 – 561.
- [54] T. A. Roman, *Quantum Stress Energy Tensors and the Weak Energy Condition*, *Phys. Rev. D* **33** (1986) 3526–3533.
- [55] A. Borde, *Geodesic focusing, energy conditions and singularities*, *Class. Quant. Grav.* **4** (1987) 343–356.
- [56] T. A. Roman, *On the ‘Averaged Weak Energy Condition’ and Penrose’s Singularity Theorem*, *Phys. Rev. D* **37** (1988) 546–548.
- [57] T. Faulkner, R. G. Leigh, O. Parrikar and H. Wang, *Modular Hamiltonians for Deformed Half-Spaces and the Averaged Null Energy Condition*, *JHEP* **09** (2016) 038. [[arXiv:1605.08072](#)].
- [58] T. Hartman, S. Kundu and A. Tajdini, *Averaged Null Energy Condition from Causality*, *JHEP* **07** (2017) 066. [[arXiv:1610.05308](#)].
- [59] P. Kravchuk and D. Simmons-Duffin, *Light-ray operators in conformal field theory*, *JHEP* **11** (2018) 102. [[arXiv:1805.00098](#)].
- [60] R. M. Wald and U. Yurtsever, *General proof of the averaged null energy condition for a massless scalar field in two-dimensional curved space-time*, *Phys. Rev. D* **44** (1991) 403–416.
- [61] R. Penrose, R. D. Sorkin and E. Woolgar, *A Positive mass theorem based on the focusing and retardation of null geodesics*, [[arXiv:gr-qc/9301015](#)].

- [62] N. Graham and K. D. Olum, *Achronal averaged null energy condition*, *Phys. Rev. D* **76** (2007) 064001. [[arXiv:0705.3193](#)].
- [63] E. Witten, *Light Rays, Singularities, and All That*, *Rev. Mod. Phys.* **92** (2020) 045004. [[arXiv:1901.03928](#)].
- [64] E.-A. Kontou and K. Sanders, *Energy conditions in general relativity and quantum field theory*, *Class. Quant. Grav.* **37** (2020) 193001. [[arXiv:2003.01815](#)].
- [65] S. J. Summers and R. Verch, *Modular inclusion, the Hawking temperature and quantum field theory in curved space-time*, *Lett. Math. Phys.* **37** (1996) 145–158.
- [66] G. L. Sewell, *Quantum fields on manifolds: PCT and gravitationally induced thermal states*, *Annals Phys.* **141** (1982) 201–224.
- [67] V. Chandrasekaran, E. E. Flanagan and K. Prabhu, *Symmetries and charges of general relativity at null boundaries*, *JHEP* **11** (2018) 125. [[arXiv:1807.11499](#)].
- [68] V. Chandrasekaran and A. J. Speranza, *Anomalies in gravitational charge algebras of null boundaries and black hole entropy*, *JHEP* **01** (2021) 137. [[arXiv:2009.10739](#)].
- [69] V. Chandrasekaran, E. E. Flanagan, I. Shehzad and A. J. Speranza, *A general framework for gravitational charges and holographic renormalization*, *Int. J. Mod. Phys. A* **37** (2022) 2250105. [[arXiv:2111.11974](#)].
- [70] L. Ciambelli, L. Freidel and R. G. Leigh, *Null Raychaudhuri: canonical structure and the dressing time*, *JHEP* **01** (2024) 166. [[arXiv:2309.03932](#)].
- [71] G. Odak, A. Rignon-Bret and S. Speziale, *General gravitational charges on null hypersurfaces*, *JHEP* **12** (2023) 038. [[arXiv:2309.03854](#)].
- [72] V. Chandrasekaran and E. E. Flanagan, *Horizon phase spaces in general relativity*, [[arXiv:2309.03871](#)].
- [73] H. Adami, D. Grumiller, M. M. Sheikh-Jabbari, V. Taghiloo, H. Yavartanoo and C. Zwickel, *Null boundary phase space: slicings, news & memory*, *JHEP* **11** (2021) 155. [[arXiv:2110.04218](#)].
- [74] S. Hollands, R. M. Wald and V. G. Zhang, *The Entropy of Dynamical Black Holes*, [[arXiv:2402.00818](#)].
- [75] S. Hollands and R. M. Wald, *Stability of Black Holes and Black Branes*, *Commun. Math. Phys.* **321** (2013) 629–680. [[arXiv:1201.0463](#)].
- [76] M. R. Visser and Z. Yan, *Properties of Dynamical Black Hole Entropy*, [[arXiv:2403.07140](#)].
- [77] C. P. Burgess, *Quantum Gravity in Everyday Life: General Relativity as an Effective Field Theory*, *Living Reviews in Relativity* **7** (2004) 5. [[arXiv:gr-qc/0311082](#)].

- [78] J. F. Donoghue, *The effective field theory treatment of quantum gravity*, *AIP Conf. Proc.* **1483** (2012) 73–94. [[arXiv:1209.3511](#)].
- [79] H. Araki, *Relative entropy of states of von Neumann algebras*, *Publ. Res. Inst. Math. Sci.* **11** (1975/76) 809–833.
- [80] A. Uhlmann, *Relative entropy and the Wigner-Yanase-Dyson-Lieb concavity in an interpolation theory*, *Comm. Math. Phys.* **54** (1977) 21–32.
- [81] M. Ohya and D. Petz, *Quantum entropy and its use*. Springer-Verlag Berlin, 1993.
- [82] C. Dappiaggi, V. Moretti and N. Pinamonti, *Rigorous construction and Hadamard property of the Unruh state in Schwarzschild spacetime*, *Adv. Theor. Math. Phys.* **15** (2011) 355–447.
- [83] H.-J. Borchers, *Field operators as C^∞ functions in spacelike directions*, *Nuovo Cimento (10)* **33** (1964) 1600–1613.
- [84] E. Witten, *Algebras, Regions, and Observers*, [[arXiv:2303.02837](#)].
- [85] H. Borchers, *Half-sided translations in connection with modular groups as a tool in quantum field theory*, *Preprint ESI 446* (1997) .
- [86] M. Takesaki, *Conditional expectations in von Neumann algebras*, *J. Functional Analysis* **9** (1972) 306–321.
- [87] M. Takesaki, *Theory of operator algebras. II*, vol. 125 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2003.
- [88] R. Longo, *Solution of the factorial Stone-Weierstrass conjecture. An application of the theory of standard split W^* -inclusions*, *Invent. Math.* **76** (1984) 145–155.
- [89] J. Haláček and T. Ledvinka, *The analytic conformal compactification of the Schwarzschild spacetime*, *Class. Quant. Grav.* **31** (2014) 015007. [[arXiv:1401.1337](#)].
- [90] R. M. Wald, *General Relativity*. Chicago Univ. Pr., Chicago, USA, 1984.
- [91] M. O. Fumio Hiai and M. Tsukada, *Sufficiency, kms condition and relative entropy in von neumann algebras*, *Pac. J. Math.* **96** (1981) 99–109.
- [92] D. Petz, *Properties of the relative entropy of states of von neumann algebras*, *Acta. Math. Hung.* **47** (1986) 65–72.
- [93] D. Petz, *On certain properties of the relative entropy of states of operator algebras*, *Math. Z.* **206** (1991) 351–361.
- [94] J. B. Hartle and S. W. Hawking, *Path-integral derivation of black-hole radiance*, *Phys. Rev. D* **13** (1976) 2188–2203.

- [95] A. Ashtekar and M. Streubel, *Symplectic Geometry of Radiative Modes and Conserved Quantities at Null Infinity*, *Proceedings of the Royal Society of London Series A* **376** (1981) 585–607.
- [96] R. M. Wald and A. Zoupas, *A General definition of ‘conserved quantities’ in general relativity and other theories of gravity*, *Phys. Rev. D* **61** (2000) 084027. [[arXiv:gr-qc/9911095](#)].
- [97] A. M. Grant, K. Prabhu and I. Shehzad, *The Wald–Zoupas prescription for asymptotic charges at null infinity in general relativity*, *Class. Quant. Grav.* **39** (2022) 085002. [[arXiv:2105.05919](#)].
- [98] H. Bondi, M. G. J. van der Burg and A. W. K. Metzner, *Gravitational waves in general relativity. 7. Waves from axisymmetric isolated systems*, *Proc. Roy. Soc. Lond. A* **269** (1962) 21–52.
- [99] R. K. Sachs, *Gravitational waves in general relativity. 8. Waves in asymptotically flat space-times*, *Proc. Roy. Soc. Lond. A* **270** (1962) 103–126.
- [100] A. Ashtekar and A. Magnon-Ashtekar, *Energy-Momentum in General Relativity*, *Phys. Rev. Lett.* **43** (1979) 181–184.
- [101] K. Prabhu, G. Satishchandran and R. M. Wald, *Infrared finite scattering theory in quantum field theory and quantum gravity*, *Phys. Rev. D* **106** (2022) 066005. [[arXiv:2203.14334](#)].
- [102] R. Longo and E. Witten, *A note on continuous entropy*, *Pure Appl. Math. Quart.* **19** (2023) 2501–2523. [[arXiv:2202.03357](#)].
- [103] D. L. Jafferis and L. Lamprou, *Inside the hologram: reconstructing the bulk observer’s experience*, *JHEP* **03** (2022) 084. [[arXiv:2009.04476](#)].
- [104] J. de Boer, D. L. Jafferis and L. Lamprou, *On black hole interior reconstruction, singularities and the emergence of time*, [[arXiv:2211.16512](#)].
- [105] R. C. Myers and M. J. Perry, *Black Holes in Higher Dimensional Space-Times*, *Annals Phys.* **172** (1986) 304.
- [106] S. Hollands, A. Ishibashi and R. M. Wald, *A Higher dimensional stationary rotating black hole must be axisymmetric*, *Commun. Math. Phys.* **271** (2007) 699–722. [[arXiv:gr-qc/0605106](#)].
- [107] S. W. Hawking, *Particle Creation by Black Holes*, *Commun. Math. Phys.* **43** (1975) 199–220.
- [108] R. M. Wald, *On Particle Creation by Black Holes*, *Commun. Math. Phys.* **45** (1975) 9–34.

- [109] B. S. Kay and R. M. Wald, *Theorems on the Uniqueness and Thermal Properties of Stationary, Nonsingular, Quasifree States on Space-Times with a Bifurcate Killing Horizon*, *Phys. Rept.* **207** (1991) 49–136.
- [110] V. P. Frolov and K. S. Thorne, *Renormalized Stress - Energy Tensor Near the Horizon of a Slowly Evolving, Rotating Black Hole*, *Phys. Rev. D* **39** (1989) 2125–2154.
- [111] W. G. Unruh, *Notes on black hole evaporation*, *Phys. Rev. D* **14** (1976) 870.
- [112] R. M. Wald, *Quantum Field Theory in Curved Space-Time and Black Hole Thermodynamics*. Chicago Lectures in Physics. University of Chicago Press, Chicago, IL, 1995.
- [113] H. Araki and L. Zsidó, *Extension of the structure theorem of Borchers and its application to half-sided modular inclusions*, *Rev. Math. Phys.* **17** (2005) 491–543.
- [114] J. Kudler-Flam, S. Leutheusser, A. A. Rahman, G. Satishchandran and A. J. Speranza, *A covariant regulator for entanglement entropy: proofs of the Bekenstein bound and QNEC*, [[arXiv:2312.07646](https://arxiv.org/abs/2312.07646)].
- [115] M. Junge, R. Renner, D. Sutter, M. M. Wilde and A. Winter, *Universal Recovery Maps and Approximate Sufficiency of Quantum Relative Entropy*, *Annales Henri Poincaré* **19** (2018) 2955–2978. [[arXiv:1509.07127](https://arxiv.org/abs/1509.07127)].
- [116] T. Faulkner, S. Hollands, B. Swingle and Y. Wang, *Approximate Recovery and Relative Entropy I: General von Neumann Subalgebras*, *Commun. Math. Phys.* **389** (2022) 349–397. [[arXiv:2006.08002](https://arxiv.org/abs/2006.08002)].
- [117] T. Faulkner and S. Hollands, *Approximate recoverability and relative entropy II: 2-positive channels of general von Neumann algebras*, *Lett. Math. Phys.* **112** (2022) 26. [[arXiv:2010.05513](https://arxiv.org/abs/2010.05513)].
- [118] L. Accardi and C. Cecchini, *Conditional expectations in von neumann algebras and a theorem of takesaki*, *Journal of Functional Analysis* **45** (1982) 245–273.
- [119] D. Petz, *A dual in von Neumann algebras with weights*, *The Quarterly Journal of Mathematics* **35** (1984) 475–483.
- [120] B. Simon, *Loewner’s theorem on monotone matrix functions*. Springer, 2019.
- [121] F. Kubo and T. Ando, *Means of positive linear operators*, *Mathematische Annalen* **246** (1980) 205–224.
- [122] K. Furuya, N. Lashkari and S. Ouseph, *Monotonic multi-state quantum f -divergences*, *J. Math. Phys.* **64** (2023) 042203. [[arXiv:2103.09893](https://arxiv.org/abs/2103.09893)].
- [123] S. Doplicher and R. Longo, *Standard and split inclusions of von Neumann algebras*, *Invent. Math.* **75** (1984) 493–536.

- [124] R. Longo, Y. Tanimoto and Y. Ueda, *Free products in AQFT*, *Annales Inst. Fourier* **69** (2019) 1229–1258. [[arXiv:1706.06070](#)].
- [125] A. Connes, *Une classification des facteurs de type III*, *Ann. Sci. École Norm. Sup. (4)* **6** (1973) 133–252.
- [126] I. E. Segal, *A note on the concept of entropy*, *Journal of Mathematics and Mechanics* **9** (1960) 623–629.
- [127] A. Rignon-Bret, *Second law from the Noether current on null hypersurfaces*, *Phys. Rev. D* **108** (2023) 044069. [[arXiv:2303.07262](#)].
- [128] N. Engelhardt and A. C. Wall, *Decoding the Apparent Horizon: Coarse-Grained Holographic Entropy*, *Phys. Rev. Lett.* **121** (2018) 211301. [[arXiv:1706.02038](#)].
- [129] N. Engelhardt and A. C. Wall, *Coarse Graining Holographic Black Holes*, *JHEP* **05** (2019) 160. [[arXiv:1806.01281](#)].
- [130] B. Freivogel and D. Krommydas, *The Smeared Null Energy Condition*, *JHEP* **12** (2018) 067. [[arXiv:1807.03808](#)].
- [131] J. R. Fliss and B. Freivogel, *Semi-local Bounds on Null Energy in QFT*, *SciPost Phys.* **12** (2022) 084. [[arXiv:2108.06068](#)].
- [132] J. R. Fliss, B. Freivogel and E.-A. Kontou, *The double smeared null energy condition*, *SciPost Phys.* **14** (2023) 024. [[arXiv:2111.05772](#)].
- [133] E. Witten, *APS Medal for Exceptional Achievement in Research: Invited article on entanglement properties of quantum field theory*, *Rev. Mod. Phys.* **90** (2018) 045003. [[arXiv:1803.04993](#)].
- [134] V. Chandrasekaran, E. E. Flanagan, I. Shehzad and A. J. Speranza, *Brown-York charges at null boundaries*, *JHEP* **01** (2022) 029. [[arXiv:2109.11567](#)].
- [135] E. Gourgoulhon and J. L. Jaramillo, *A 3+1 perspective on null hypersurfaces and isolated horizons*, *Phys. Rept.* **423** (2006) 159–294. [[arXiv:gr-qc/0503113](#)].
- [136] R. Penrose, *Techniques of differential topology in relativity*. Society for Industrial and Applied Mathematics, 1972.
- [137] C. Crnkovic and E. Witten, *Covariant description of canonical formalism in geometrical theories*, in *Three Hundred Years of Gravitation* (S. W. Hawking and W. Israel, eds.), ch. 16, pp. 676–684. Cambridge University Press, 1987.
- [138] J. Lee and R. M. Wald, *Local symmetries and constraints*, *J. Math. Phys.* **31** (1990) 725–743.
- [139] V. Iyer and R. M. Wald, *Some properties of Noether charge and a proposal for dynamical black hole entropy*, *Phys. Rev. D* **50** (1994) 846–864. [[arXiv:gr-qc/9403028](#)].

- [140] D. Harlow and J.-Q. Wu, *Covariant phase space with boundaries*, *JHEP* **10** (2020) 146. [[arXiv:1906.08616](#)].