

Optimal anisotropies for p -Laplace type operators in the plane ^{*†}

Raul Fernandes Horta [‡]

*Departamento de Matemática, Universidade Federal de Minas Gerais,
Caixa Postal 702, 30123-970, Belo Horizonte, MG, Brazil*

Marcos Montenegro [§]

*Departamento de Matemática, Universidade Federal de Minas Gerais,
Caixa Postal 702, 30123-970, Belo Horizonte, MG, Brazil*

Abstract

Sharp lower and upper uniform estimates are obtained for fundamental frequencies of p -Laplace type operators generated by quadratic forms. Optimal constants are exhibited, rigidity of the upper estimate is proved, anisotropic attainability of the lower estimate is derived as well as characterization of anisotropic extremizers for circular and rectangular membranes. Sharp quantitative anisotropic inequalities associated with lower constants are also established, providing as a by-product information on anisotropic stability. When the uniform ellipticity condition is relaxed, we show that the optimal lower constant remains positive, while anisotropic extremizers no longer exist. Our sharp lower estimate can be viewed as an isoanisotropic counterpart of the Faber-Krahn isoperimetric inequality in the plane.

1 Introduction

A field of research that has aroused great interest in the mathematical community for several decades is the study of problems linking shape of domains and elliptic operators to the corresponding spectra. Most of their solutions demand techniques in areas as calculus of variations, elliptic PDEs, geometry and spectral analysis. A well-known prototype of the first group of problems is the celebrated Faber-Krahn geometric inequality which states that the first Dirichlet eigenvalue $\lambda_1(\Omega)$ of the Laplace operator in a bounded domain $\Omega \subset \mathbb{R}^n$, also called the fundamental frequency of Ω , satisfies the isoperimetric property

$$\lambda_1(\Omega) \geq \lambda_1(B) \tag{1}$$

for every domain Ω with $|\Omega| = |B|$, where B denotes the unit n -Euclidean ball and $|\cdot|$ stands for the Lebesgue measure of a measurable subset of \mathbb{R}^n . Moreover, equality holds in (1) if, and only if, Ω is equal to B , up to a translation and a set of zero capacity. Inequality (1), as its name suggests, was independently

^{*}2020 Mathematics Subject Classification: 49J20, 49J30, 47J20

[†]Key words: Anisotropic optimization, sharp anisotropic estimates, anisotropic rigidity, anisotropic extremizers

[‡]*E-mail addresses:* raul.fernandes.horta@gmail.com (R. F. Horta)

[§]*E-mail addresses:* montene@mat.ufmg.br (M. Montenegro)

proved by Faber [31] and Krahn [40] for any dimension $n \geq 2$ using Schwarz symmetrization technique. A number of variants of it has been established along of the history. For instance, related inequalities involving the first nonzero eigenvalue of the Laplace operator under different boundary conditions are: Bossel-Daners inequality [9, 14, 23, 24] (Robin condition) (see also [25, 26]), Szegő-Weinberger inequality [49, 54] (Neumann condition) (see also [16]) and Brock-Weinstock inequality [12, 55] (Steklov condition) (being the first one of lower kind, while other two ones are upper). Different quantitative forms and stability of geometric inequalities have also been widely discussed for the spectrum of elliptic operators; among an extensive bibliography, we refer for example to [8, 11, 15, 33] for some quantitative inequalities and [32, 35, 43] concerning stability. Besides, extensions of (1) to more general operators have been of interest, see [1, 7, 10, 20] for some of these developments. For an overview on a variety of problems, improvements and open questions on several related topics, we quote the excellent monographs: the survey [47], the collection of contributions [38] and the books [5, 6, 19, 37].

A particularly important counterpart of (1), closely connected to the present work, was established by Belloni, Ferone and Kawohl [4] in the anisotropic context. In a precise manner, let $H : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function of C^1 class in $\mathbb{R}^n \setminus \{0\}$ satisfying the homogeneity and strict positivity assumptions:

$$H(t\xi) = |t|H(\xi), \quad t \in \mathbb{R}, \quad \forall \xi \in \mathbb{R}^n \quad (\text{H})$$

and

$$\kappa_1|\xi| \leq H(\xi) \leq \kappa_2|\xi|, \quad \forall \xi \in \mathbb{R}^n, \quad (\text{P})$$

where $\kappa_1 = \kappa_1(H)$ and $\kappa_2 = \kappa_2(H)$ are positive constants depending of H .

Given any fixed number $p > 1$, consider the anisotropic operator in divergence form associated to p and H :

$$\Delta_p^H u := -\operatorname{div}(H^{p-1}(\nabla u) \nabla H(\nabla u)).$$

It arises naturally from derivation of the energy functional on $W_0^{1,p}(\Omega)$ defined by

$$\mathcal{E}_{p,H}(u) = \int_{\Omega} H^p(\nabla u) \, dx,$$

since (p, H) -harmonic functions (*i.e.* weak solutions of $\Delta_p^H u = 0$) are critical points of $\mathcal{E}_{p,H}$.

The first Dirichlet eigenvalue $\lambda_{1,p}^H(\Omega)$ of the operator Δ_p^H , also called the H -anisotropic fundamental p -frequency of Ω , is variationally characterized as

$$\lambda_{1,p}^H(\Omega) = \inf \left\{ \mathcal{E}_{p,H}(u) : u \in W_0^{1,p}(\Omega), \|u\|_{L^p(\Omega)} = 1 \right\}.$$

From the conditions (H) and (P), it follows that $\lambda_{1,p}^H(\Omega)$ is positive and Δ_p^H is uniformly elliptic. The latter fact along with (H) and C^2 regularity of H outside the origin imply that $\lambda_{1,p}^H(\Omega)$ is the unique principal eigenvalue of Δ_p^H , which is also simple. In addition, all eigenfunctions of Δ_p^H in $W_0^{1,p}(\Omega)$ belong to $C^{1,\beta}(\Omega)$ and extends $C^{1,\beta}$ at points on $C^{1,\alpha}$ parts of the boundary of Ω where $\alpha > \beta$. For the part of regularity we refer to [51, 52] and references therein.

Consider the unit ball $B_H = \{\xi \in \mathbb{R}^n : H(\xi) \leq 1\}$ and its polar body defined by

$$B_H^\circ = \{\xi \in \mathbb{R}^n : \langle \xi, \eta \rangle \leq 1, \quad \eta \in B_H\}.$$

The anisotropic Faber-Krahn inequality established in [4] states that, for any bounded domain Ω with $|\Omega| = |B_H^\circ|$,

$$\lambda_{1,p}^H(\Omega) \geq \lambda_{1,p}^H(B_H^\circ) \quad (2)$$

and, in addition, equality holds in (2) if, and only if, Ω is equal to B_H° , module a translation and a set of zero capacity.

On the other hand, it deserves to be remarked that the Faber-Krahn inequality (2) provides a sharp lower uniform estimate of $\lambda_{1,p}^H(\Omega)$ in terms of the measure $|\Omega|$, namely

$$\lambda_{1,p}^H(\Omega) \geq |B_H^\circ|^{\frac{p}{n}} \lambda_{1,p}^H(B_H^\circ) |\Omega|^{-\frac{p}{n}}.$$

Within this spirit, considerable advances have been achieved in optimization problems that consist in finding, given a bounded domain Ω_0 , uniform estimates of eigenvalues with respect to the coefficients of elliptic operators and/or other involved parameters, with optimal lower and upper constants being explicitly computed in some cases. More precisely, this question has been addressed to Dirichlet eigenvalues $\lambda_k(V)$ of the Schrödinger operator

$$\mathcal{L}_S u = -\hbar \Delta u + V(x)u \quad \text{in } \Omega_0,$$

where \hbar denotes the Planck's constant and V is a potential function subject to the restrictions

$$\|V\|_{L^p(\Omega_0)} \leq \kappa \quad \text{and} \quad \kappa_1 \leq V \leq \kappa_2 \quad \text{in } \Omega_0$$

for a priori fixed constants κ , κ_1 and κ_2 , we refer to [21, 28, 30, 39, 41, 45, 50, 56] for the one-dimensional case and [2, 13, 17, 18, 27, 36, 44] and Chapter 9 of [38] for higher dimensions.

Sharp estimates have been also obtained for Dirichlet eigenvalues $\lambda_k(\sigma)$ of the elliptic operator

$$\mathcal{L}_C u = -\operatorname{div}(\sigma(x)\nabla u) \quad \text{in } \Omega$$

for conductivity functions σ normalized simultaneously by the uniform and L^p constraints above with constants $\kappa_1, \kappa_2 > 0$, see [3, 22, 29, 53] for some developments.

Inequality as (1) and (2) are of isoperimetric nature once one treats sharp lower estimate of the first Dirichlet eigenvalue of elliptic operators on bounded domains with prescribed Lebesgue measure. A natural query to ask is whether there are isoanisotropic counterparts of (1) and (2). A positive answer to this question relies on finding some prescribed “measure” condition of functions H . Once an appropriate normalization has been defined, we are faced with two fundamental issues on a fixed bounded domain Ω . More precisely, we have:

(A) Are there any positive explicit optimal constants $\lambda_{1,p}^{\min}(\Omega)$ and $\lambda_{1,p}^{\max}(\Omega)$ such that

$$\lambda_{1,p}^{\min}(\Omega) \leq \lambda_{1,p}^H(\Omega) \leq \lambda_{1,p}^{\max}(\Omega)$$

for every “normalized” convex function H satisfying (H) and (P)?

(B) Are there functions H yielding any equality in (A)? Is there any characterization of them?

The positivity of $\lambda_{1,p}^{\min}(\Omega)$ and finiteness of $\lambda_{1,p}^{\max}(\Omega)$ must necessarily depend on the set of anisotropic functions to be introduced. Inequalities in (A) display sharp uniform estimates on H for the first eigenvalue of the operator Δ_p^H . Moreover, the first of them is the isoanisotropic parallel of the n -dimensional Faber-Krahn isoperimetric inequality. Functions H solving (B) are called anisotropic extremizers.

As with most eigenvalue optimization problems focused on domains varying with fixed measure, optimization problems that involve the dependence on second order elliptic operators in a fixed domain are usually difficult.

The purpose of this paper is to start the discussion of (A) and (B) in the plane within the class of functions H generated by positive quadratic forms (so that Δ_p^H is uniformly elliptic), namely functions H under the form

$$H(x, y) = (\alpha x^2 + 2\beta xy + \gamma y^2)^{1/2}$$

satisfying (P) with positive constants κ_1 and κ_2 independent of H . Our family of elliptic operators includes the Laplace and p -Laplace operators, since

$$\Delta = \Delta_2^{H_0} \quad \text{and} \quad \Delta_p = \Delta_p^{H_0}$$

for the Euclidean norm $H_0(x, y) = |(x, y)| = (x^2 + y^2)^{1/2}$.

The question (A) will be completely solved for the referred class. Already the answer to the question (B) is divided into three parts: we first prove the rigidity of the second inequality in (A), being attained exactly by H_0 ; we prove the existence of at least one anisotropic extremizer for the first inequality; we characterize all anisotropic extremizers for the first inequality on disks and on a family of rectangles, where it becomes clear that the quantity (multiplicity) of such optimal functions depends on the shape of the domain Ω . We also present sharp quantitative anisotropic estimates related to the lower constant $\lambda_{1,p}^{\min}(\Omega)$. Finally, we discuss (A) and (B) when the uniform ellipticity condition is relaxed, in particular we show that the optimal lower constant remains positive, while anisotropic extremizers no longer exist.

This work seems to be the first to develop a comprehensive optimization theory for a family of quasilinear elliptic operators in dimension $n = 2$. The leading arguments are based on maximum principles, monotonic property of $\lambda_{1,p}^H(\Omega)$ with respect to H and mainly on a key explicit relation stated in Theorem 5.1 which is the bridge that connects the quadratic anisotropic environment to the study of anisotropic stability of optimal lower constants via aforementioned quantitative estimates.

The remainder of paper is organized into four sections. In Section 2 we introduce basic notations and the entire setup of quadratic anisotropic optimization. In Section 3 we present all statements related to (A) and (B). Section 4 is devoted to proofs of the results dealing with anisotropic maximization, while Section 5 is dedicated to all proofs related to anisotropic minimization.

2 The anisotropic optimization setting

We will use from now on the letter Q (and derivatives thereof) in place of H in the notations regarding anisotropic eigenvalues and other ingredients, once the class of functions H to be considered is in correspondence with positive quadratic forms by means of the relation $H = Q^{1/2}$.

The set of all positive quadratic forms on \mathbb{R}^2 is given by

$$\mathcal{Q} := \{Q(x, y) = \alpha x^2 + 2\beta xy + \gamma y^2 : \alpha, \gamma > 0, \beta \geq 0 \text{ and } \beta^2 < \alpha\gamma\}.$$

For each $Q \in \mathcal{Q}$, set

$$Q_{\min} = \min_{|(x,y)|=1} Q(x,y) \quad \text{and} \quad Q_{\max} = \max_{|(x,y)|=1} Q(x,y),$$

where $|\cdot|$ denotes the Euclidean norm.

For $a \in (0, 1]$, we consider the subset of \mathcal{Q} of normalized functions as

$$\mathcal{Q}^a := \{Q \in \mathcal{Q} : Q_{\min} \geq a \quad \text{and} \quad Q_{\max} = 1\}.$$

For any $Q \in \mathcal{Q}^a$, clearly we have

$$a|(x,y)|^2 \leq Q(x,y) \leq |(x,y)|^2, \quad \forall (x,y) \in \mathbb{R}^2. \quad (3)$$

For $a = 0$, we denote by \mathcal{Q}^0 the subset of \mathcal{Q} with relaxed strict coercivity, that is

$$\mathcal{Q}^0 := \{Q \in \mathcal{Q} : Q_{\max} = 1\}.$$

Let $\Omega \subset \mathbb{R}^2$ be a bounded domain and let $p > 1$ be a fixed parameter. The Q -anisotropic fundamental p -frequency is defined for $Q \in \mathcal{Q}$ as

$$\lambda_{1,p}^Q(\Omega) := \inf \left\{ \iint_{\Omega} Q^{\frac{p}{2}}(\nabla u) dA : u \in W_0^{1,p}(\Omega), \|u\|_p = 1 \right\},$$

where dA denotes the usual area element.

As mentioned in the introduction, it is a classical result that the infimum is always attained by a positive function $\varphi_p \in C^{1,\beta}(\Omega)$, which is a principal eigenfunction associated to the principal Dirichlet eigenvalue $\lambda_{1,p}^Q(\Omega)$ of the Q -anisotropic operator

$$\Delta_p^Q u := -\operatorname{div} \left(Q^{\frac{p-1}{2}}(\nabla u) \nabla Q^{\frac{1}{2}}(\nabla u) \right).$$

In addition, φ_p is $C^{1,\beta}$ on $C^{1,\alpha}$ parts of the boundary of Ω with $\alpha > \beta$.

Notice also that the operator Δ_p^Q is uniformly elliptic for each $Q \in \mathcal{Q}^a$ in a uniform sense since the corresponding ellipticity constants depend only on the fixed number a . Taking into account this uniformity, we consider two anisotropic optimization problems for each $a \in [0, 1]$:

$$\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) := \inf_{Q \in \mathcal{Q}^a} \lambda_{1,p}^Q(\Omega) \quad \text{and} \quad \lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega) := \sup_{Q \in \mathcal{Q}^a} \lambda_{1,p}^Q(\Omega).$$

From the definition of \mathcal{Q}^a and (3), one easily checks that

$$a^{p/2} \lambda_{1,p}(\Omega) \leq \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) \leq \lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega) \leq \lambda_{1,p}(\Omega), \quad (4)$$

where $\lambda_{1,p}(\Omega)$ denotes the principal Dirichlet eigenvalue of the p -Laplace operator. In particular, both constants are finite and clearly positive if $a > 0$. Furthermore, when $a = 1$, the definition of \mathcal{Q}^a and inequalities in (4) imply that $\mathcal{Q}^1 = \{|\cdot|^2\}$ and

$$\lambda_{1,p}^{\min}(\mathcal{Q}^1, \Omega) = \lambda_{1,p}^{\max}(\mathcal{Q}^1, \Omega) = \lambda_{1,p}(\Omega).$$

The questions (A) and (B) presented in the introduction can now be rephrased for $a \in [0, 1]$ as

(A1) Can the optimal anisotropic constants $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$ and $\lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega)$ be explicitly determined?

(B1) Are there anisotropic extremizers for $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$ and $\lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega)$? Is it possible to characterize them?

Lastly, for each $a \in (0, 1)$ we will need the set of non-normalized functions by coercivity:

$$\begin{aligned}\mathcal{Q}_{nn}^a &:= \{Q \in \mathcal{Q} : aQ_{\max} \leq Q_{\min}\} \\ &= \{Q \in \mathcal{Q} : Q(x, y) \geq aQ_{\max} |(x, y)|^2, \forall (x, y) \in \mathbb{R}^2\}.\end{aligned}$$

Indeed, remark that $Q^\varepsilon(x, y) = \varepsilon|(x, y)|^2$ belongs to \mathcal{Q}_{nn}^a but not to \mathcal{Q}^a for any $0 < \varepsilon < a$, so that there are anisotropic operators Δ_p^Q associated to $Q \in \mathcal{Q}_{nn}^a$ with arbitrarily small ellipticity constant.

3 Main results on anisotropic optimization

Our first result answers affirmatively both questions (A1) and (B1) for $\lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega)$.

Theorem 3.1 (Rigidity). *For any $a \in [0, 1)$, we have*

$$\lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega) = \lambda_{1,p}(\Omega).$$

Moreover, $Q = |\cdot|^2 \in \mathcal{Q}^a$ is the unique anisotropic maximizer for $\lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega)$ provided that $\partial\Omega$ is $C^{1,\alpha}$ by parts.

Regarding the anisotropic minimization problem we need to introduce a family of elements of \mathcal{Q}^a and of orthogonal 2×2 matrices in order to state the next theorems.

For $\alpha \in [a, 1]$, we consider the anisotropic function $Q_\alpha(x, y) = \alpha x^2 + 2\beta(\alpha)xy + \gamma(\alpha)y^2$ with coefficients

$$\beta(\alpha) = \sqrt{(1-\alpha)(\alpha-a)} \quad \text{and} \quad \gamma(\alpha) = 1 + a - \alpha.$$

As will be shown later, $Q_\alpha \in \mathcal{Q}^a$ and $(Q_\alpha)_{\min} = a$ for every $\alpha \in [a, 1]$.

Let also \mathcal{O} be the set of all rotations in the plane of angle between 0 and $\pi/2$. For $A \in \mathcal{O}$ we set $\Omega_A = A^T(\Omega)$, that is Ω_A is the rotation of the domain Ω by the orthogonal matrix A^T .

The second result completes the solution of the question (A1) and establishes the existence of at least one anisotropic extremizer for $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$ in the case $a \in (0, 1)$.

Theorem 3.2 (Attainability). *For any $a \in (0, 1)$, we have*

$$\begin{aligned}\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) &= \min \left\{ \lambda_{1,p}^{Q_a}(\Omega_A) : \text{for any } A \in \mathcal{O} \right\} \\ &= a^{p/2} \min \left\{ \lambda_{1,p}(\Omega_A^a) : \text{for any } A \in \mathcal{O} \right\},\end{aligned}$$

where $\Omega^a = \{(x, \sqrt{a}y) : (x, y) \in \Omega\}$.

Moreover, all anisotropic minimizers for $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$ belong to $\{Q_\alpha : \alpha \in [a, 1]\}$ provided that $\partial\Omega$ is $C^{1,\alpha}$ by parts.

Theorems 3.1 and 3.2 applied to the normalized function $Q_{\max}^{-1}Q \in \mathcal{Q}^a$ for $Q \in \mathcal{Q}_{nn}^a$ readily yield

Corollary 3.1. *For any $a \in (0, 1)$, the sharp non-normalized anisotropic estimates hold*

$$\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) Q_{\max}^{p/2} \leq \lambda_{1,p}^Q(\Omega) \leq \lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega) Q_{\max}^{p/2}$$

for every $Q \in \mathcal{Q}_{nn}^a$ with explicit optimal constants $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$ and $\lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega)$.

A second consequence that will be easily checked states that

Corollary 3.2. *Assume $\partial\Omega$ is $C^{1,\alpha}$ by parts. For any $a \in (0, 1)$, we have*

$$a^{p/2} \lambda_{1,p}(\Omega) < \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) < \lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega).$$

It is natural to wonder at this point what happens if we relax the condition $Q_{\min} \geq a$ with $a \in (0, 1]$ to $Q_{\min} > 0$, in other words, if we consider the anisotropic minimization problem in the set \mathcal{Q}^0 . It is easily checked that

$$\lambda_{1,p}^{\min}(\mathcal{Q}^0, \Omega) = \inf_{a \in (0,1]} \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega).$$

The next result ensures that $\lambda_{1,p}^{\min}(\mathcal{Q}^0, \Omega)$ remains positive, however, anisotropic extremizer no longer exist, so the condition of strict coercivity for some fixed number $a \in (0, 1]$ is necessary for its existence.

Theorem 3.3. *The optimal lower constant $\lambda_{1,p}^{\min}(\mathcal{Q}^0, \Omega)$ is always positive. In addition, anisotropic minimizers do not exist in \mathcal{Q}^0 provided that $\partial\Omega$ is $C^{1,\alpha}$ by parts.*

Unlike the upper constant $\lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega)$, the lower $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$ depends on the sets \mathcal{Q}^a . In general, we have $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) \leq \lambda_{1,p}^{\min}(\mathcal{Q}^b, \Omega)$ whenever $a \leq b$ since $\mathcal{Q}^b \subset \mathcal{Q}^a$.

Other important topic concerns the stability of $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$ in relation to the sets \mathcal{Q}^a . This follows directly from the quantitative anisotropic inequality:

Theorem 3.4 (Quantitative I). *Let $a \in (0, 1)$ and $b \in [a, 1)$. The upper quantitative anisotropic inequality states that*

$$\frac{\lambda_{1,p}^{\min}(\mathcal{Q}^b, \Omega)}{\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)} - 1 \leq p \sqrt{\frac{b^{p-1}(b-a)}{a^p(1-a)}}.$$

Moreover, it is sharp in the sense that equality implies $b = a$.

We also establish a reverse quantitative anisotropic inequality.

Theorem 3.5 (Quantitative II). *Let $a \in (0, 1)$ and $b \in [a, 1)$. The lower quantitative anisotropic inequality*

$$\lambda_{1,p}^{\min}(\mathcal{Q}^b, \Omega) - \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) \geq C(a, b, p, \Omega)(b - a)$$

holds for

$$C(a, b, p, \Omega) = \begin{cases} \frac{pa^{(p-2)/2}}{2} c_0(p, \Omega), & \text{if } p \geq 2 \\ \frac{pb^{(2-p)/2}}{2} \lambda_{1,p}(\Omega)^{(p-2)/2} c_0(p, \Omega)^{2/p}, & \text{if } 1 < p < 2, \end{cases}$$

where

$$c_0(p, \Omega) := \inf_{A \in \mathcal{O}} \inf_{u \in W_0^{1,p}(\Omega)} \left\{ \iint_{\Omega_A} |D_x u|^p dA : \|u\|_p = 1 \right\}$$

is a positive constant.

Moreover, both are sharp provided that $\partial\Omega$ is $C^{1,\alpha}$.

Regardless of the shape of the membrane Ω , Theorem 3.2 guarantees that $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$ always admits at least one anisotropic minimizer. The final two theorems of this section deal with the exact multiplicity of these extremizers. They seem to point to a close relationship between the numbers of anisotropic minimizers and of certain symmetries of the region Ω . In particular, we give a complete answer to (B1) in the case of disks and a class of rectangles.

Theorem 3.6. *Let $a \in (0, 1)$ and assume Ω is a disk D centered at the origin. Then,*

$$\lambda_{1,p}^{\min}(\mathcal{Q}^a, D) = \lambda_{1,p}^{Q_a}(D) = a^{p/2} \lambda_{1,p}(\mathcal{E}_a),$$

where \mathcal{E}_a denotes the elliptical region $\{(x, \sqrt{a}y) : (x, y) \in D\}$.

Moreover, the set of all anisotropic minimizers is precisely given by $\{Q_\alpha : \alpha \in [a, 1]\}$.

Theorem 3.7. *Let $a \in (0, 1)$ and assume Ω is the rectangle $R_a = [-1, 1] \times [-1/\sqrt{a}, 1/\sqrt{a}]$. Then,*

$$\lambda_{1,p}^{\min}(\mathcal{Q}^a, R_a) = \lambda_{1,p}^{Q_a}(R_a) = a^{p/2} \lambda_{1,p}(R).$$

where R denotes the square $[-1, 1] \times [-1, 1]$.

Moreover, the unique anisotropic minimizers are Q_a and Q_1 .

4 Optimization problems associated to $\lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega)$

We begin with a monotonicity result which will be used in the proof of Theorems 3.1, 3.2 and 3.3 and Corollary 3.2.

Proposition 4.1. *Let $Q_1, Q_2 \in \mathcal{Q}$. It holds that $\lambda_{1,p}^{Q_1}(\Omega) \leq \lambda_{1,p}^{Q_2}(\Omega)$ whenever $Q_1 \leq Q_2$ in \mathbb{R}^2 .*

Moreover, $\lambda_{1,p}^{Q_1}(\Omega) = \lambda_{1,p}^{Q_2}(\Omega)$ only when $Q_1 = Q_2$ in \mathbb{R}^2 provided that $\partial\Omega$ is $C^{1,\alpha}$ by parts.

For the proof of the second part of this statement we recall a simple property about nonnegative quadratic forms.

Lemma 4.1. *Let Q be a nonnegative quadratic form on \mathbb{R}^2 . If $Q \not\equiv 0$, then either $Q > 0$ in $\mathbb{R}^2 \setminus \{(0, 0)\}$ or the kernel of Q is a straight line in \mathbb{R}^2 .*

Proof. Let M be a symmetric matrix that represents Q , that is, $Q(v) = \langle v, Mv \rangle$ with $v = (x, y)$. Using that $Q \geq 0$ we ensure that $Q(v_0) = 0$ if, and only if, $Mv_0 = 0$. Assume $Q(v_0) = 0$ and consider the function $h_v(t) = Q(tv + v_0)$ for $t \in \mathbb{R}$, where v is any nonzero fixed vector in \mathbb{R}^2 . Noting that

$$h_v(t) = t^2 \langle v, Mv \rangle + 2t \langle v, Mv_0 \rangle + \langle v_0, Mv_0 \rangle.$$

and $t = 0$ is a minimum point of $h_v(t)$, we have $h'_v(0) = 0$, or equivalently $\langle v, Mv_0 \rangle = 0$. Since v is an arbitrary vector, it follows that $Mv_0 = 0$. The reciprocal is immediate.

Assume now that $Q \not\equiv 0$. For the conclusion, it suffices to show that there are no two linearly independent vectors in the kernel of Q . Let v_1, v_2 be nonzero vectors such that $Q(v_1) = 0 = Q(v_2)$. Then, from the above claim, we have $Mv_1 = 0 = Mv_2$. If v_1 and v_2 are linearly independent, then they span \mathbb{R}^2 and therefore $M \equiv 0$, so $Q \equiv 0$, contradicting the hypothesis of the statement. \square

Proof of Proposition 4.1. The claim $\lambda_{1,p}^{Q_1}(\Omega) \leq \lambda_{1,p}^{Q_2}(\Omega)$ is immediate once it follows readily of the variational characterization. The part that needs to be carefully checked is that equality $\lambda_{1,p}^{Q_1}(\Omega) = \lambda_{1,p}^{Q_2}(\Omega)$ implies $Q_1 = Q_2$ in \mathbb{R}^2 . Proceeding by contradiction and applying Lemma 4.1 to the nonnegative quadratic form $Q = Q_2 - Q_1$, it follows that $Q_1 < Q_2$ at least in \mathbb{R}^2 except a straight line passing through the origin.

Let $\varphi_p \in W_0^{1,p}(\Omega) \cap C^{1,\beta}(\Omega)$ be a positive minimizer for $\lambda_{1,p}^{Q_1}(\Omega)$ normalized by $\|\varphi_p\|_p = 1$. Using the assumption that $\partial\Omega$ is $C^{1,\alpha}$ by parts, we recall that φ_p extends $C^{1,\beta}$ until the boundary up to a finite number of points. Then, applying the Hopf's Lemma to the quasilinear elliptic operator $\Delta_p^{Q_1}$, we guarantee that $\nabla\varphi_p$ is a nonzero vector on each $C^{1,\alpha}$ part of $\partial\Omega$ pointing inward of Ω . Since Ω is a domain, we can choose a point (x_0, y_0) on some smooth part of $\partial\Omega$ so that $Q_1(\nabla\varphi_p(x_0, y_0)) < Q_2(\nabla\varphi_p(x_0, y_0))$. It is clear that the strict inequality $Q_1(\nabla\varphi_p) < Q_2(\nabla\varphi_p)$ remains in the open $\Omega_\delta = \{(x, y) \in \Omega : |(x - x_0, y - y_0)| < \delta\}$ for $\delta > 0$ small enough. Using the strict inequality in Ω_δ and $Q_1(\nabla\varphi_p) \leq Q_2(\nabla\varphi_p)$ in $\Omega \setminus \Omega_\delta$, we get

$$\begin{aligned} \lambda_{1,p}^{Q_1}(\Omega) &\leq \iint_{\Omega} Q_1^{\frac{p}{2}}(\nabla\varphi_p) dA = \iint_{\Omega_\delta} Q_1^{\frac{p}{2}}(\nabla\varphi_p) dA + \iint_{\Omega \setminus \Omega_\delta} Q_1^{\frac{p}{2}}(\nabla\varphi_p) dA \\ &\leq \iint_{\Omega_\delta} Q_1^{\frac{p}{2}}(\nabla\varphi_p) dA + \iint_{\Omega \setminus \Omega_\delta} Q_2^{\frac{p}{2}}(\nabla\varphi_p) dA \\ &< \iint_{\Omega_\delta} Q_2^{\frac{p}{2}}(\nabla\varphi_p) dA + \iint_{\Omega \setminus \Omega_\delta} Q_2^{\frac{p}{2}}(\nabla\varphi_p) dA = \lambda_{1,p}^{Q_2}(\Omega). \end{aligned}$$

□

Proof of Theorem 3.1. By definition, for any $Q \in \mathcal{Q}^a$, we have $Q(x, y) \leq \overline{Q}(x, y) := x^2 + y^2$ for all $(x, y) \in \mathbb{R}^2$ and so, by Proposition 4.1,

$$\lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega) = \lambda_{1,p}^{\overline{Q}}(\Omega) = \lambda_{1,p}(\Omega)$$

since $\overline{Q} \in \mathcal{Q}^a$. Moreover, by the same proposition, the strict inequality $\lambda_{1,p}^Q(\Omega) < \lambda_{1,p}^{\overline{Q}}(\Omega)$ holds whenever $Q \not\equiv \overline{Q}$, so that \overline{Q} is the unique anisotropic extremizer for $\lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega)$. □

5 Optimization problems associated to $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$

First we develop all the necessary ingredients to prove the existence and shape of anisotropic extremizers for $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$ on \mathcal{Q}^a .

5.1 The structure of the set \mathcal{Q}^a

Let

$$\mathcal{Q}_a := \{Q(x, y) \in \mathcal{Q}^a : Q_{\min} = a\}.$$

Clearly, we have $\mathcal{Q}_a \subset \mathcal{Q}^a$. The next result describes precisely the set \mathcal{Q}_a .

Proposition 5.1. *For each $a \in (0, 1)$, the set \mathcal{Q}_a is precisely given by $\{Q_\alpha : \alpha \in [a, 1]\}$, where Q_α is as defined in Section 3.*

Proof. Let $Q = \alpha x^2 + 2\beta xy + \gamma y^2$ with $\alpha, \gamma > 0$, $\beta \geq 0$ and $\beta^2 < \alpha\gamma$. Consider an orthogonal matrix A diagonalizing the symmetric matrix

$$M = \begin{bmatrix} \alpha & \beta \\ \beta & \gamma \end{bmatrix},$$

into its real eigenvalues μ_1 and μ_2 which can be assumed $\mu_1 \leq \mu_2$. One easily checks that $(Q \circ A)(x, y) = \mu_1 x^2 + \mu_2 y^2$. Hence, $Q \in \mathcal{Q}_a$ if, and only if, $\mu_1 = a$ and $\mu_2 = 1$. But this is equivalent to say that β and γ satisfy the system

$$\begin{cases} (\alpha - a)(\gamma - a) = \beta^2 \\ (\alpha - 1)(\gamma - 1) = \beta^2, \end{cases}$$

whose solution is given for each $\alpha \in [a, 1]$ by

$$\beta(\alpha) = \sqrt{(1 - \alpha)(\alpha - a)} \quad \text{and} \quad \gamma(\alpha) = 1 + a - \alpha.$$

□

A direct consequence of this result is

Corollary 5.1. *Let $a \in (0, 1)$. For any $\alpha \in [a, 1]$, we have $Q_\alpha \circ A_\alpha = Q_a$, where A_α denotes the orthogonal matrix*

$$A_\alpha = \begin{bmatrix} \sqrt{\frac{1-\alpha}{1-a}} & \sqrt{\frac{\alpha-a}{1-a}} \\ -\sqrt{\frac{\alpha-a}{1-a}} & \sqrt{\frac{1-\alpha}{1-a}} \end{bmatrix}.$$

In particular, there is a natural bijection between \mathcal{Q}_a and $\{A_\alpha : \alpha \in [a, 1]\}$ which in turn is in correspondence with the set of rotations \mathcal{O} defined in the introduction.

Proof. It is immediate to see that the columns of A_α are eigenvectors of M associated to a and 1 in this order and also that the set $\{A_\alpha : \alpha \in [a, 1]\}$ coincides with $\mathcal{O} = \{R_\theta : \theta \in [0, \frac{\pi}{2}]\}$, where

$$R_\theta = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}.$$

□

A convenient way to look at the set \mathcal{Q}^a is through the disjoint decomposition $\dot{\bigcup}_{b \in [a, 1]} \mathcal{Q}_b$. Using this viewpoint, we establish the following key relation:

Theorem 5.1. *Let $a \in (0, 1)$. Given any $Q \in \mathcal{Q}^a$ there is a function $Q_\alpha \in \mathcal{Q}_a$ such that*

$$Q(x, y) = \frac{1-b}{1-a} Q_\alpha(x, y) + \frac{b-a}{1-a} (x^2 + y^2), \quad \forall (x, y) \in \mathbb{R}^2,$$

where $b \in [a, 1]$ is the number for which $Q \in \mathcal{Q}_b$.

Particularly, this statement implies that $Q \geq Q_\alpha$ in \mathbb{R}^2 and $Q \equiv Q_\alpha$ if, and only if, $b = a$.

Proof. Let $Q \in \mathcal{Q}^a$ and let $b \in [a, 1]$ be such that $Q \in \mathcal{Q}_b$. Writing $Q(x, y) = \bar{\alpha}x^2 + 2\bar{\beta}xy + \bar{\gamma}y^2$ with $\bar{\alpha}, \bar{\gamma} > 0$, $\bar{\beta} \geq 0$ and $\bar{\beta}^2 < \bar{\alpha}\bar{\gamma}$, by Proposition 5.1, we know that

$$\bar{\alpha} \in [b, 1], \quad \bar{\beta} = \sqrt{(1 - \bar{\alpha})(\bar{\alpha} - b)} \quad \text{and} \quad \bar{\gamma} = 1 + b - \bar{\alpha}.$$

Let us seek a function $Q_\alpha \in \mathcal{Q}_a$ such that

$$Q \geq Q_\alpha \quad \text{in } \mathbb{R}^2. \quad (5)$$

Again by that proposition, we have $Q_\alpha(x, y) = \alpha x^2 + 2\beta xy + \gamma y^2$ where

$$\alpha \in [a, 1], \quad \beta = \sqrt{(1 - \alpha)(\alpha - a)} \quad \text{and} \quad \gamma = 1 + a - \alpha.$$

Notice that the inequality (5) holds if, and only if,

$$(\bar{\beta} - \beta)^2 - (\bar{\alpha} - \alpha)(\bar{\gamma} - \gamma) \leq 0. \quad (6)$$

The idea now is to prove the equivalence between this inequality and the relation

$$\alpha = \frac{1 - a}{1 - b}\bar{\alpha} + \frac{a - b}{1 - b}. \quad (7)$$

Replacing $\bar{\beta}$ and $\bar{\gamma}$ in function of $\bar{\alpha}$ and β and γ in function of α in (6), after some development, we arrive at

$$(1 - \alpha)(\bar{\alpha} - b) + (1 - \bar{\alpha})(\alpha - a) - 2\sqrt{(1 - \alpha)(\bar{\alpha} - b)(1 - \bar{\alpha})(\alpha - a)} \leq 0.$$

On the other hand, this inequality can be rewritten as

$$(\sqrt{(1 - \alpha)(\bar{\alpha} - b)} - \sqrt{(1 - \bar{\alpha})(\alpha - a)})^2 \leq 0.$$

Consequently,

$$(1 - \alpha)(\bar{\alpha} - b) = (1 - \bar{\alpha})(\alpha - a)$$

which yields the value of α as in (7). This equation also allows us to express $\bar{\alpha}$, $\bar{\beta}$ and $\bar{\gamma}$ in terms respectively of α , β and γ . Indeed, it can be placed into three suitable ways:

$$\bar{\alpha} = \frac{1 - b}{1 - a}\alpha + \frac{b - a}{1 - a}, \quad (8)$$

$$1 - \bar{\alpha} = \frac{1 - b}{1 - a}(1 - \alpha) \quad \text{and} \quad \bar{\alpha} - b = \frac{1 - b}{1 - a}(\alpha - a).$$

The last equalities clearly imply that $\alpha \in [a, 1]$ if, and only if, $\beta \in [b, 1]$. Moreover, they produce the relations

$$\bar{\beta} = \sqrt{(1 - \bar{\alpha})(\bar{\alpha} - b)} = \frac{1 - b}{1 - a}\sqrt{(1 - \alpha)(\alpha - a)} = \frac{1 - b}{1 - a}\beta \quad (9)$$

and

$$\bar{\gamma} = 1 + b - \bar{\alpha} = \frac{1 - b}{1 - a}(1 - \alpha) + b = \frac{1 - b}{1 - a}(1 + a - \alpha) + \frac{b - a}{1 - a} = \frac{1 - b}{1 - a}\gamma + \frac{b - a}{1 - a}. \quad (10)$$

Plugging (8), (9) and (10) in Q , we deduce that

$$Q(x, y) = \frac{1-b}{1-a} Q_\alpha(x, y) + \frac{b-a}{1-a} (x^2 + y^2).$$

□

5.2 Proof of Theorem 3.2 and Corollary 3.2

We first prove Theorem 3.2 and after we invoke it in the proof of Corollary 3.2.

Proof of Theorem 3.2. Let $Q \in \mathcal{Q}^a$ and assume that $Q \in \mathcal{Q}_b$ for some $b \in (a, 1]$. By Theorem 5.1, there exists $Q_\alpha \in \mathcal{Q}_a$ such that $Q \geq Q_\alpha$ and $Q \not\equiv Q_\alpha$ since $b > a$. By Proposition 4.1, we have in general that $\lambda_{1,p}^Q(\Omega) \geq \lambda_{1,p}^{Q_\alpha}(\Omega)$. Therefore, it follows that

$$\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) = \inf_{Q \in \mathcal{Q}_a} \lambda_{1,p}^Q(\Omega)$$

and moreover, by assuming $\partial\Omega$ is $C^{1,\alpha}$ by parts, the above monotonicity is strict, so that any anisotropic extremizer, if exist, belongs to \mathcal{Q}_a .

On the other hand, by Corollary 5.1, each function $Q \in \mathcal{Q}_a$ corresponds to an orthogonal matrix $A \in \mathcal{O}$ such that $Q \circ A = Q_a$. Then, using the change of variable $v(X') = u(X)$ for $X' = A^T X \in \Omega_A$ with $X = (x, y)$ and $X' = (x', y')$, we obtain

$$\iint_{\Omega} Q^{\frac{p}{2}}(\nabla u) dA = \iint_{\Omega_A} Q_a^{\frac{p}{2}}(\nabla v) dA \quad \text{and} \quad \iint_{\Omega} |u|^p dA = \iint_{\Omega_A} |v|^p dA,$$

which yields $\lambda_{1,p}^Q(\Omega) = \lambda_{1,p}^{Q_a}(\Omega_A)$. Making now the change $v(x', y') = u(x, y)$ for $x' = x$ and $y' = \sqrt{a}y$ for $(x', y') \in \Omega_A^a$, we get

$$\iint_{\Omega_A} Q_a^{\frac{p}{2}}(\nabla u) dA = a^{(p-1)/2} \iint_{\Omega_A^a} |\nabla v|^p dA \quad \text{and} \quad \iint_{\Omega_A} |u|^p dA = a^{-1/2} \iint_{\Omega_A^a} |v|^p dA,$$

and so $\lambda_{1,p}^{Q_a}(\Omega_A) = a^{p/2} \lambda_{1,p}(\Omega_A^a)$. Consequently, we derive

$$\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) = \inf_{A \in \mathcal{O}} \lambda_{1,p}^{Q_a}(\Omega_A) = a^{p/2} \inf_{A \in \mathcal{O}} \lambda_{1,p}(\Omega_A^a).$$

Notice now that \mathcal{O} is a closed subset of the compact set of all orthogonal 2×2 matrices and the first eigenvalue of the p -Laplace operator $\lambda_{1,p}(\Omega)$ depends continuously on Ω with respect to the Hausdorff topology (see Theorem 3.2 in [42]). Hence, the above infimum is always attained and therefore,

$$\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) = \min_{A \in \mathcal{O}} \lambda_{1,p}^{Q_a}(\Omega_A) = a^{p/2} \min_{A \in \mathcal{O}} \lambda_{1,p}(\Omega_A^a).$$

Moreover, each minimizer $A \in \mathcal{O}$ is associated to the anisotropic extremizer $Q = Q_a \circ A^T \in \mathcal{Q}_a$ for $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$. □

Proof of Corollary 3.2. Let $a \in (0, 1)$. Consider the functions $\underline{Q}(x, y) = a(x^2 + y^2)$ and $\overline{Q}(x, y) = x^2 + y^2$. By Theorems 3.1 and 3.2, we know that

$$\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) = \lambda_{1,p}^{Q_a}(\Omega) \quad \text{and} \quad \lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega) = \lambda_{1,p}^{\overline{Q}}(\Omega)$$

for some $Q_\alpha \in \mathcal{Q}_a$. From the definition of \mathcal{Q}_a , one knows that $\underline{Q} \not\leq Q_\alpha \not\leq \overline{Q}$ in \mathbb{R}^2 . Since $\partial\Omega$ is $C^{1,\alpha}$ by parts, by Proposition 4.1, we get

$$a^{p/2}\lambda_{1,p}(\Omega) = \lambda_{1,p}^{\underline{Q}}(\Omega) < \lambda_{1,p}^{Q_\alpha}(\Omega) < \lambda_{1,p}^{\overline{Q}}(\Omega),$$

so that

$$a^{p/2}\lambda_{1,p}(\Omega) < \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) < \lambda_{1,p}^{\max}(\mathcal{Q}^a, \Omega).$$

□

5.3 Proof of Theorem 3.3

We start proving the positivity of $\lambda_{1,p}^{\min}(\mathcal{Q}^0, \Omega)$. For any $a \in (0, 1]$, due to Theorem 3.2, we have $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) = \lambda_{1,p}^{Q_a}(\Omega_A)$ for some rotation $A \in \mathcal{O}$. If $u_a \in W_0^{1,p}(\Omega)$ with $\|u_a\|_p = 1$ is an eigenfunction associated to $\lambda_{1,p}^{Q_a}(\Omega_A)$, then

$$\begin{aligned} \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) &= \lambda_{1,p}^{Q_a}(\Omega_A) = \iint_{\Omega_A} Q_a^{\frac{p}{2}}(\nabla u_a) dA \\ &= \iint_{\Omega_A} (a|D_x u_a|^2 + |D_y u_a|^2)^{\frac{p}{2}} dA \\ &\geq \iint_{\Omega_A} |D_y u_a|^p dA \\ &\geq d(p, \Omega_A) \iint_{\Omega_A} |u_a|^p dA = d(p, \Omega_A), \end{aligned}$$

where $d(p, \Omega_A) = t_p^p w(\Omega_A, \xi)^{-p}$ is the positive constant given in Lemma 1 of [34] with $\xi = (0, 1)$. Hence, using that \mathcal{O} is compact and $w(\Omega_A, \xi)$ depends continuously on A , one has

$$d_0(p, \Omega) := \inf_{A \in \mathcal{O}} d(p, \Omega_A) = \min_{A \in \mathcal{O}} d(p, \Omega_A) > 0,$$

so that

$$\lambda_{1,p}^{\min}(\mathcal{Q}^0, \Omega) = \inf_{a \in (0, 1]} \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) \geq d_0(p, \Omega) > 0.$$

The second statement is proved by contradiction. Assume that there is $Q \in \mathcal{Q}^0$ such that $\lambda_{1,p}^{\min}(\mathcal{Q}^0, \Omega) = \lambda_{1,p}^Q(\Omega)$. Set $a = Q_{\min} > 0$, so $\lambda_{1,p}^{\min}(\mathcal{Q}^0, \Omega) = \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)$. From Theorem 3.2, we know that $\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) = \lambda_{1,p}^{Q_a}(\Omega_A)$ for some $A \in \mathcal{O}$. Since $Q_a \geq Q_{a/2}$ in \mathbb{R}^2 and $Q_a \not\equiv Q_{a/2}$, by Proposition 4.1, we get

$$\lambda_{1,p}^{\min}(\mathcal{Q}^0, \Omega) = \lambda_{1,p}^{Q_a}(\Omega_A) > \lambda_{1,p}^{Q_{a/2}}(\Omega_A) = \lambda_{1,p}^{\tilde{Q}}(\Omega)$$

with $\tilde{Q} = Q_{a/2} \circ A^T \in \mathcal{Q}^0$, which is a contradiction.

5.4 Proof of Theorems 3.4 and 3.5

Proof of Theorem 3.4. Let $a \in (0, 1)$ and $b \in [a, 1)$. Given $Q \in \mathcal{Q}_b$, by Theorem 5.1, there is a function $Q_\alpha \in \mathcal{Q}_a$ such that

$$Q(x, y) = \frac{1-b}{1-a} Q_\alpha(x, y) + \frac{b-a}{1-a} |(x, y)|^2.$$

Let $\varphi_p \in W_0^{1,p}(\Omega)$ with $\|\varphi_p\|_p = 1$ such that

$$\iint_{\Omega} Q_\alpha^{\frac{p}{2}}(\nabla \varphi_p) dA = \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega).$$

Then using the mean value theorem,

$$\begin{aligned} \lambda_{1,p}^{\min}(\mathcal{Q}^b, \Omega) - \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) &\leq \iint_{\Omega} Q^{\frac{p}{2}}(\nabla \varphi_p) dA - \iint_{\Omega} Q_\alpha^{\frac{p}{2}}(\nabla \varphi_p) dA = \iint_{\Omega} Q^{\frac{p}{2}}(\nabla \varphi_p) - Q_\alpha^{\frac{p}{2}}(\nabla \varphi_p) dA \\ &\leq \iint_{\Omega} p Q^{\frac{p-1}{2}}(\nabla \varphi_p) \left(Q^{\frac{1}{2}}(\nabla \varphi_p) - Q_\alpha^{\frac{1}{2}}(\nabla \varphi_p) \right) dA \\ &\leq p \iint_{\Omega} Q^{\frac{p-1}{2}}(\nabla \varphi_p) \left(\sqrt{\frac{1-b}{1-a}} Q_\alpha^{\frac{1}{2}}(\nabla \varphi_p) + \sqrt{\frac{b-a}{1-a}} |\nabla \varphi_p| - Q_\alpha^{\frac{1}{2}}(\nabla \varphi_p) \right) dA \\ &\leq \frac{p}{\sqrt{1-a}} \iint_{\Omega} Q^{\frac{p-1}{2}}(\nabla \varphi_p) \left((\sqrt{1-b} - \sqrt{1-a}) Q_\alpha^{\frac{1}{2}}(\nabla \varphi_p) + \sqrt{b-a} |\nabla \varphi_p| \right) dA \\ &\leq \frac{p}{\sqrt{1-a}} \iint_{\Omega} Q^{\frac{p-1}{2}}(\nabla \varphi_p) \sqrt{b-a} |\nabla \varphi_p| dA \\ &\leq \frac{p\sqrt{b-a}}{\sqrt{1-a}} \iint_{\Omega} Q^{\frac{p-1}{2}}(\nabla \varphi_p) \frac{Q_\alpha^{\frac{1}{2}}(\nabla \varphi_p)}{\sqrt{a}} dA \\ &= \frac{p\sqrt{b-a}}{\sqrt{a(1-a)}} \iint_{\Omega} \left(\frac{1-b}{1-a} Q_\alpha(\nabla \varphi_p) + \frac{b-a}{1-a} |\nabla \varphi_p|^2 \right)^{\frac{p-1}{2}} Q_\alpha^{\frac{1}{2}}(\nabla \varphi_p) dA \\ &\leq \frac{p\sqrt{b-a}}{\sqrt{a(1-a)^p}} \iint_{\Omega} \left(1-b + \frac{b-a}{a} \right)^{\frac{p-1}{2}} Q_\alpha^{\frac{p}{2}}(\nabla \varphi_p) dA \\ &= p \sqrt{\frac{b^{p-1}(b-a)}{a^p(1-a)}} \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega). \end{aligned}$$

Hence, we derive the upper quantitative inequality

$$\frac{\lambda_{1,p}^{\min}(\mathcal{Q}^b, \Omega)}{\lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega)} - 1 \leq p \sqrt{\frac{b^{p-1}(b-a)}{a^p(1-a)}},$$

which is sharp since equality in the above third inequality only holds when $b = a$ since $b < 1$. \square

Proof of Theorem 3.5. We first get a lower quantitative inequality for $p \geq 2$. Let $Q \in \mathcal{Q}_b$ and choose a positive function $\psi_p \in W_0^{1,p}(\Omega)$ with $\|\psi_p\|_p = 1$ such that

$$\lambda_{1,p}^{\min}(\mathcal{Q}^b, \Omega) = \iint_{\Omega} Q^{\frac{p}{2}}(\nabla \psi_p) dA.$$

Again taking $Q_\alpha \in \mathcal{Q}_a$ as in Theorem 5.1 and applying the mean value theorem, one gets

$$\begin{aligned}
\lambda_{1,p}^{\min}(\mathcal{Q}^b, \Omega) - \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) &\geq \iint_{\Omega} Q^{\frac{p}{2}}(\nabla \psi_p) dA - \iint_{\Omega} Q_{\alpha}^{\frac{p}{2}}(\nabla \psi_p) dA \\
&\geq \iint_{\Omega} \frac{p}{2} Q_{\alpha}^{\frac{p}{2}-1}(\nabla \psi_p) (Q(\nabla \psi_p) - Q_{\alpha}(\nabla \psi_p)) dA \\
&= \frac{p}{2} \iint_{\Omega} Q_{\alpha}^{\frac{p}{2}-1}(\nabla \psi_p) \left(\frac{1-b}{1-a} Q_{\alpha}(\nabla \psi_p) + \frac{b-a}{1-a} |\nabla \psi_p|^2 - Q_{\alpha}(\nabla \psi_p) \right) dA \\
&= \frac{p}{2} \iint_{\Omega} Q_{\alpha}^{\frac{p}{2}-1}(\nabla \psi_p) \frac{b-a}{1-a} (|\nabla \psi_p|^2 - Q_{\alpha}(\nabla \psi_p)) dA \\
&\geq \frac{pa^{(p-2)/2}(b-a)}{2(1-a)} \iint_{\Omega} |\nabla \psi_p|^{p-2} (|\nabla \psi_p|^2 - Q_{\alpha}(\nabla \psi_p)) dA \\
&= \frac{pa^{(p-2)/2}(b-a)}{2(1-a)} \iint_{\Omega_A} |\nabla v_p|^{p-2} (|\nabla v_p|^2 - a|D_x v_p|^2 - |D_y v_p|^2) dA \\
&= \frac{pa^{(p-2)/2}(b-a)}{2} \iint_{\Omega_A} |\nabla v_p|^{p-2} |D_x v_p|^2 dA \\
&\geq \frac{pa^{(p-2)/2}(b-a)}{2} \iint_{\Omega_A} |D_x v_p|^p dA,
\end{aligned}$$

where $A \in \mathcal{O}$ is the matrix so that $Q_{\alpha} \circ A = Q_a$ and $v_p = \psi_p \circ A$. Therefore,

$$\lambda_{1,p}^{\min}(\mathcal{Q}^b, \Omega) - \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) \geq \frac{pa^{(p-2)/2}(b-a)}{2} c_0(p, \Omega),$$

where, as in the statement,

$$c_0(p, \Omega) := \inf_{A \in \mathcal{O}} \inf_{u \in W_0^{1,p}(\Omega)} \left\{ \iint_{\Omega_A} |D_x u|^p dA : \|u\|_p = 1 \right\}.$$

The case $1 < p < 2$ is treated in a similar manner. With the same notation, applying the mean value theorem and then reverse Hölder inequality, we have

$$\begin{aligned}
\lambda_{1,p}^{\min}(\mathcal{Q}^b, \Omega) - \lambda_{1,p}^{\min}(\mathcal{Q}^a, \Omega) &\geq \iint_{\Omega} Q^{\frac{p}{2}}(\nabla \psi_p) dA - \iint_{\Omega} Q_{\alpha}^{\frac{p}{2}}(\nabla u) dA \\
&\geq \iint_{\Omega} \frac{p}{2} Q^{\frac{p}{2}-1}(\nabla \psi_p) (Q(\nabla \psi_p) - Q_{\alpha}(\nabla \psi_p)) dA \\
&= \frac{p}{2} \iint_{\Omega} Q^{\frac{p}{2}-1}(\nabla \psi_p) \left(\frac{1-b}{1-a} Q_{\alpha}(\nabla \psi_p) + \frac{b-a}{1-a} |\nabla \psi_p|^2 - Q_{\alpha}(\nabla \psi_p) \right) dA \\
&= \frac{p}{2} \iint_{\Omega} Q^{\frac{p}{2}-1}(\nabla \psi_p) \frac{b-a}{1-a} (|\nabla \psi_p|^2 - Q_{\alpha}(\nabla \psi_p)) dA \\
&\geq \frac{p(b-a)}{2(1-a)} \iint_{\Omega} |\nabla \psi_p|^{p-2} (|\nabla \psi_p|^2 - Q_{\alpha}(\nabla \psi_p)) dA \\
&= \frac{p(b-a)}{2(1-a)} \iint_{\Omega_A} |\nabla v_p|^{p-2} (|\nabla v_p|^2 - a|D_x v_p|^2 - |D_y v_p|^2) dA \\
&= \frac{p(b-a)}{2} \iint_{\Omega_A} |\nabla v_p|^{p-2} |D_x v_p|^2 dA
\end{aligned}$$

$$\begin{aligned}
&\geq \frac{p(b-a)}{2} \left(\iint_{\Omega_A} |\nabla v_p|^p dA \right)^{\frac{p-2}{p}} \left(\iint_{\Omega_A} |D_x v_p|^p dA \right)^{\frac{2}{p}} \\
&= \frac{p(b-a)}{2} \left(\iint_{\Omega} |\nabla \psi_p|^p dA \right)^{\frac{p-2}{p}} \left(\iint_{\Omega_A} |D_x v_p|^p dA \right)^{\frac{2}{p}} \\
&\geq \frac{p(b-a)}{2} \left(\iint_{\Omega} \frac{Q^{\frac{p}{2}}(\nabla \psi_p)}{b^{p/2}} dA \right)^{\frac{p-2}{p}} \left(\iint_{\Omega_A} |D_x v_p|^p dA \right)^{\frac{2}{p}} \\
&= \frac{pb^{(2-p)/2}(b-a)}{2} (\lambda_{1,p}^{\min}(\mathcal{Q}^b, \Omega))^{\frac{p-2}{2}} \left(\iint_{\Omega_A} |D_x v_p|^p dA \right)^{\frac{2}{p}} \\
&\geq \frac{pb^{(2-p)/2}(b-a)}{2} \lambda_{1,p}(\Omega)^{(p-2)/2} c_0(p, \Omega)^{2/p}.
\end{aligned}$$

Besides, if $u \in W_0^{1,p}(\Omega)$ and $\|u\|_p = 1$, we have

$$\iint_{\Omega_A} |D_x u|^p dA \geq c(p, \Omega_A) \iint_{\Omega_A} |u|^p dA = c(p, \Omega_A),$$

where $c(p, \Omega_A) = t_p^p w(\Omega_A, \xi)^{-p}$ is the positive constant given in Lemma 1 of [34] with $\xi = (1, 0)$. Arguing exactly as in the proof of Theorem 3.3, we deduce that

$$c_0(p, \Omega) \geq \inf_{A \in \mathcal{O}} c(p, \Omega_A) = \min_{A \in \mathcal{O}} c(p, \Omega_A) > 0.$$

Finally, if $\partial\Omega$ is of $C^{1,\alpha}$ class, then $\psi_p \in C^{1,\beta}(\overline{\Omega})$ and, thanks to the Hopf's Lemma, $|\nabla \psi_p|^{-1} \nabla \psi_p$ maps the unity circle \mathbb{S}^1 . Consequently, equality in the second inequality of both cases implies that $Q(\nabla \psi_p) = Q_\alpha(\nabla \psi_p)$, so that $Q \equiv Q_\alpha$ and then $b = a$. \square

5.5 Proof of Theorems 3.6 and 3.7

Proof of Theorem 3.6: When Ω is a disk D centered at the origin, we have that $\Omega_A = D$ and $\Omega_A^a = \mathcal{E}_a$ for every $A \in \mathcal{O}$. Thus, $\lambda_{1,p}^{Q_a}(\Omega_A)$ doesn't depend on A and is equal to the constant value $\lambda_{1,p}^{Q_a}(D) = a^{p/2} \lambda_{1,p}(\mathcal{E}_a)$. Therefore, by Theorem 3.2, it follows that

$$\lambda_{1,p}^{\min}(\mathcal{Q}^a, D) = \lambda_{1,p}^{Q_a}(D) = a^{p/2} \lambda_{1,p}(\mathcal{E}_a)$$

and the corresponding set of extremizers is

$$\mathcal{Q}_a = \{Q_a \circ A^T : A \in \mathcal{O}\} = \{Q_\alpha : \alpha \in [a, 1]\}.$$

\square

The proof of Theorem 3.7 uses an important result on minimization of $\lambda_{1,p}(R)$ over all quadrilaterals R of same area. The Faber-Krahn inequality for quadrilaterals states that the minimum of $\lambda_{1,p}(R)$ is attained only at squares of same area. For $p = 2$, this is a classical theorem due to Pólya and Szegő [48] whose proof is based on Steiner's symmetrization arguments. For $p \neq 2$, the result is also true and its proof is carried out step by step as in [48], see for example the proof of Theorem 1.1 of [46] where the same method is employed in the nonlocal p -Laplace case.

Proof of Theorem 3.7: Assume $\Omega = R_a = [-1, 1] \times [-1/\sqrt{a}, 1/\sqrt{a}]$ for $a \in (0, 1)$. Notice that Ω_A^a is a quadrilateral of area 4 for any $A \in \mathcal{O}$. Recall that the minimizers of $\lambda_{1,p}^{\min}(\mathcal{Q}^a, R_a)$ are given by $Q_a \circ A^T$ where A minimizes $\lambda_{1,p}(\Omega_A^a)$. On the other hand, there are only two matrices $A \in \mathcal{O}$ that transform Ω_A^a into a square, namely, the identity matrix I and

$$A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

Therefore, the unique minimizers of $\lambda_{1,p}^{\min}(\mathcal{Q}^a, R_a)$ are Q_a and $Q_a \circ A^T = Q_1$. Finally, using the identity matrix, we derive

$$\lambda_{1,p}^{\min}(\mathcal{Q}^a, R_a) = \lambda_{1,p}^{Q_a}(R_a) = a^{p/2} \lambda_{1,p}(R),$$

where $R = [-1, 1] \times [-1, 1]$. □

Acknowledgments: The first author was partially supported by CAPES (PROEX 88887.712161/2022-00) and the second author was partially supported by CNPq (PQ 307432/2023-8, Universal 404374/2023-9) and Fapemig (Universal APQ 01598-23).

*On behalf of all authors, the corresponding author states that there is no conflict of interest.

*Data availability: Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

References

- [1] M.S. Ashbaugh - *Isoperimetric and universal inequalities for eigenvalues*, London Math. Soc. Lecture Note Ser., 273, Cambridge University Press, Cambridge, 95-139, 1999.
- [2] M.S. Ashbaugh, E.M. Harrell - *Maximal and minimal eigenvalues and their associated nonlinear equations*, J. Math. Phys., 28, no. 8, 1770-1786, 1987.
- [3] C. Bandle - *Extremal problems for eigenvalues of the Sturm-Liouville type*, General inequalities, 5 (Oberwolfach, 1986), 319-336, Internat. Schriftenreihe Numer. Math., 80, Birkhäuser, Basel, 1987.
- [4] M. Belloni, V. Ferone, and B. Kawohl - *Isoperimetric inequalities, Wulff shape and related questions for strongly nonlinear elliptic operators*, Zeitschrift für Angewandte Mathematik und Physik (ZAMP), 54, no. 5, 771-783, 2003.
- [5] P. Bérard - *Spectral Geometry: Direct and Inverse Problems*, Lecture Notes in Math., 1207, Springer, 1986.
- [6] M. Berger - *Geometry of the spectrum*, in Differential Geometry (Chern, S. S. and Osserman, R., eds.), Proc. Sympos. Pure Math., Vol. 27, Part 2, American Mathematical Society, pp. 129-152, 1975.
- [7] S. Biagi, S. Dipierro, E. Valdinoci, E. Vecchi - *A Faber-Krahn inequality for mixed local and nonlocal operators*, J. Anal. Math. 150, no. 2, 405-448, 2023.

- [8] I. Birindelli, G. Galise, H. Ishii - *A family of degenerate elliptic operators: maximum principle and its consequences*, Ann. Inst. H. Poincaré C Anal. Non Linéaire 35, no. 2, 417-441, 2018.
- [9] M-H. Bossel - *Elastically supported membranes, either nonhomogeneous or on a surface: a new extension of the Rayleigh-Faber-Krahn isoperimetric theorem*, Z. Angew. Math. Phys. 39, no.5, 733-742, 1988.
- [10] L. Brasco, E. Cinti, S. Vita - *A quantitative stability estimate for the fractional Faber-Krahn inequality*, J. Funct. Anal. 279, no. 3, 108560, 49 pp, 2020.
- [11] L. Brasco, G. De Philippis, B. Velichkov - *Faber-Krahn inequalities in sharp quantitative form*, Duke Math. J. 164, no. 9, 1777-1831, 2015.
- [12] F. Brock - *An isoperimetric inequality for eigenvalues of the Stekloff problem*, Z. Angew. Math. Mech. 81, no. 1, 69-71, 2001.
- [13] D. Bucur, G. Buttazzo, B. Velichkov - *Spectral optimization problems for potentials and measures*, SIAM J. Math. Anal. 46, no. 4, 2956-2986, 2014.
- [14] D. Bucur, D. Daners - *An alternative approach to the Faber-Krahn inequality for Robin problems*, Calc. Var. Partial Differential Equations 37, no. 1-2, 75-86, 2010.
- [15] D. Bucur, V. Ferone, C. Nitsch, C. Trombetti - *The quantitative Faber-Krahn inequality for the Robin Laplacian*, J. Differential Equations 264, no. 7, 4488-4503, 2018.
- [16] D. Bucur, A. Henrot - *Maximization of the second non-trivial Neumann eigenvalue*, Acta Math. 222, no. 2, 337-361, 2019.
- [17] G. Buttazzo, A. Gerolin, B. Ruffini, B. Velichkov - *Optimal potentials for Schrödinger operators*, J. Éc. polytech. Math. 1, 71-100, 2014.
- [18] S. Chanillo, D. Grieser, M. Imai, K. Kurata, I. Ohnishi - *Symmetry breaking and other phenomena in the optimization of eigenvalues for composite membranes*, Comm. Math. Phys. 214, no. 2, 315-337, 2000.
- [19] I. Chavel - *Eigenvalues in Riemannian Geometry*, Academic Press, 1994.
- [20] A.M.H. Chorwadwala, R. Mahadevan, F. Toledo - *On the Faber-Krahn inequality for the Dirichlet p -Laplacian*, ESAIM Control Optim. Calc. Var. 21, no. 1, 60-72, 2015.
- [21] J. Chu, G. Meng, Z. Zhang - *Minimizations of positive periodic and Dirichlet eigenvalues for general indefinite Sturm-Liouville problems*, Adv. Math. 432, Paper No. 109272, 38 pp, 2023.
- [22] S.J. Cox, R. Lipton - *Extremal eigenvalue problems for two-phase conductors*, Arch. Rational Mech. Anal., 136, 101-117, 1996.
- [23] D. Daners - *A Faber-Krahn inequality for Robin problems in any space dimension*, Mathematische Annalen 335, no. 4, 767-785, 2006.
- [24] D. Daners, J. Kennedy - *Uniqueness in the Faber-Krahn inequality for Robin problems*, SIAM J. Math. Anal. 39, no. 4, 1191-1207, 2007.

- [25] F. Della Pietra, N. Gavitone - *Faber-Krahn inequality for anisotropic eigenvalue problems with Robin boundary conditions*, Potential Analysis 41, 1147-1166, 2014.
- [26] F. Della Pietra, G. Piscitelli - *Sharp estimates for the first Robin eigenvalue of nonlinear elliptic operators*, J. Differential Equations 386, 269-293, 2024.
- [27] H. Egnell - *Extremal properties of the first eigenvalue of a class of elliptic eigenvalue problems*, Ann. Scuola Norm. Sup. Pisa Cl. Sci., (4) 14, no. 1, 1-48, 1987.
- [28] Y. Egorov, S. Karaa - *Optimisation de la première valeur propre de l'opérateur de Sturm-Liouville*, C. R. Acad. Sci. Paris Série I Math., 319, no. 8, 793-798, 1994.
- [29] Y. Egorov, V. Kondratiev - *On spectral theory of elliptic operators*, Oper. Theory Adv. Appl., 89, Birkhäuser, Basel, 1996.
- [30] M. Essen - *On estimating eigenvalues of a second order linear differential operator*, General inequalities, 5 (Oberwolfach, 1986), 347-366, Internat. Schriftenreihe Numer. Math., 80, Birkhäuser, Basel, 1987.
- [31] G. Faber - *Beweis, dass unter allen homogenen Membranen von gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt*, Sitzungsberichte 8 Bayer Akad. d. Wiss., 1923.
- [32] N. Fusco, F. Maggi, A. Pratelli - *Stability estimates for certain Faber-Krahn, isocapacitary and Cheeger inequalities*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5), 8:51-71, 2009.
- [33] N. Fusco, Y.R. Zhang - *A quantitative form of Faber-Krahn inequality*, Calc. Var. Partial Differential Equations 56, no. 5, 44 pp, 2017.
- [34] J. Haddad, C. H. Jiménez, M. Montenegro - *From affine Poincaré inequalities to affine spectral inequalities*, Adv. Math. 386, 107808, 35 pp, 2021.
- [35] W. Hansen, N. Nadirashvili - *Isoperimetric inequalities in potential theory*, Proceedings from the International Conference on Potential Theory (Amersfoort, 1991). Potential Anal., 3:1-14, 1994.
- [36] E.M. Harrell - *Hamiltonian operators with maximal eigenvalues*, J. Math. Phys., 25 (1984), no. 1, 48-51 and 27 (1986), no. 1, 419.
- [37] A. Henrot - *Extremum problems for eigenvalues of elliptic operators*, Series: Frontiers in Mathematics, Birkhäuser Verlag, Basel, 202 pp, 2006.
- [38] A. Henrot - *Shape optimization and spectral theory*, Edited by Antoine Henrot, De Gruyter Open, Warsaw, 464 pp, 2017.
- [39] S. Karaa - *Sharp estimates for the eigenvalues of some differential equations*, SIAM J. Math. Anal., 29, no. 5, 1279-1300, 1998.
- [40] E. Krahn - *Über eine von Rayleigh formulierte minimaleigenschaft des kreises*, Mathematische Annalen 94, no 1, 97-100, 1925.
- [41] P. Kurasov, Pavel, A. Serio - *Optimal potentials for quantum graphs*, Ann. Henri Poincaré 20, no. 5, 1517-1542, 2019.

- [42] I. Ly - *The first eigenvalue of the p -Laplacian operator*, J. Inequalities Pure Appl. Math. 6, Article 91, 2005.
- [43] A.D. Melas - *The stability of some eigenvalue estimates*, J. Diff. Geom. 36, 19-33, 1992.
- [44] M. Mihailescu, V. Radulescu, D. Repovš, Dušan - *On a non-homogeneous eigenvalue problem involving a potential: an Orlicz-Sobolev space setting*, J. Math. Pures Appl. (9) 93, no. 2, 132-148, 2010.
- [45] L. Notarantonio - *Extremal properties of the first eigenvalue of Schrödinger-type operators*, J. Funct. Anal. 156, no. 2, 333-346, 1998.
- [46] F. Olivares Contador - *The Faber-Krahn inequality for the first eigenvalue of the fractional Dirichlet p -Laplacian for triangles and quadrilaterals*, Pacific J. Math. 288, no. 2, 425-434, 2017.
- [47] R. Osserman - *Isoperimetric inequalities and eigenvalues of the Laplacian*, in Proc. Int. Congress of Math. Helsinki, 1978, and Bull.Amer. Math. Soc. 84, pp. 1182-1238, 1978.
- [48] G. Pólya, G. Szegő - *Isoperimetric Inequalities in Mathematical Physics*, Annals of Mathematics Studies, no. 27, Princeton University Press, Princeton, N. J., 1951.
- [49] G. Szegő - *Inequalities for certain eigenvalues of a membrane of given area*, J. Rational Mech. Anal. 3, 343-356, 1954.
- [50] G. Talenti - *Estimates for eigenvalues of Sturm-Liouville problems*, General inequalities, 4 (Oberwolfach, 1983), 341-350, Internat. Schriftenreihe Numer. Math., 71, Birkhäuser, Basel, 1984.
- [51] P. Tolksdorf - *On the Dirichlet problem for quasilinear equations in domains with conical boundary points*, Comm. in PDE 8, 773-817, 1983.
- [52] P. Tolksdorf - *Regularity for a more general class of quasilinear elliptic equations*, J. Diff. Equations 51, 126-150, 1984.
- [53] B.A. Troesch - *An isoperimetric sloshing problem*, Comm. Pure Appl. Math., 18, 319-338, 1965.
- [54] H.F. Weinberger - *An isoperimetric inequality for the N -dimensional free membrane problem*, J. Rational Mech. Anal. 5, 633-636, 1956.
- [55] R. Weinstock - *Inequalities for a classical eigenvalue problem*, J. Rational Mech. Anal. 3, 745-753, 1954.
- [56] Z. Zhang, X. Wang - *Sharp estimates of lowest positive Neumann eigenvalue for general indefinite Sturm-Liouville problems*, J. Differential Equations 382, 302-320, 2024.