Multiple quantum exceptional, diabolical, and hybrid points in multimode bosonic systems: II. Nonconventional \mathcal{PT} -symmetric dynamics and unidirectional coupling

Jan Peřina Jr.,^{1,*} Kishore Thapliyal,^{1,†} Grzegorz Chimczak,² Anna Kowalewska-Kudłaszyk,² and Adam Miranowicz²

¹Joint Laboratory of Optics, Faculty of Science, Palacký University,

Czech Republic, 17. listopadu 12, 771 46 Olomouc, Czech Republic

²Institute of Spintronics and Quantum Information, Faculty of Physics,

Adam Mickiewicz University, 61-614 Poznań, Poland

We analyze the existence and degeneracies of quantum exceptional, diabolical, and hybrid points of simple bosonic systems, composed of up to six modes with damping and/or amplification and exhibiting nonconventional dynamics. They involve the configurations in which the dynamics typical for PT-symmetric systems is observed only in a subspace of the whole Liouville space of the system states (nonconventional PT-symmetric dynamics) as well as those containing unidirectional coupling. The system dynamics described by quadratic non-Hermitian Hamiltonians is governed by the Heisenberg-Langevin equations. Conditions for the observation of inherited quantum hybrid points with up to sixth-order exceptional and second-order diabolical degeneracies are revealed, though relevant only for short-time dynamics. This raises the question of whether higher-order inherited singularities exist in bosonic systems that exhibit physically meaningful behavior at arbitrary times. On the other hand, for short times, unidirectional coupling of various types enables the concatenation of simple bosonic systems with second- and third-order exceptional degeneracies on demand. This approach allows for the creation of arbitrarily high exceptional degeneracies observed in systems with diverse structures. Methods for numerical identifying the quantum exceptional and hybrid points, and determining their degeneracies are discussed. Rich dynamics of higher-order fieldoperator moments is analyzed from the point of view of the presence of exceptional and diabolical points with their degeneracies in general.

I. INTRODUCTION

Non-Hermitian bosonic \mathcal{PT} -symmetric systems are endowed with interesting physical properties that occur at and around quantum exceptional and hybrid points (QEPs and QHPs). Some studies [1–5] suggest that these points can enable improvements in measurement precision beyond the classical limit. They also allow to enhance effective system nonlinearities that result in the generation of highly nonclassical and entangled states [6, 7]. The strength of these effects depends in many cases on the order of the degeneracy of exceptional points (EPs): The higher-order the degeneracy, the more enhanced the processes become. This lead us in part I of the paper [8] to the analysis of QEPs and QHPs of paritytime (PT) -symmetric bosonic systems with up to five modes considering different configurations. However, this analysis revealed only the systems with QHPs with the second- and third-order exceptional geneneracies (EDs) and second-order diabolical degeneracies (DDs), despite the fact that the bosonic systems with five modes were considered with the promise of observation of QHPs with the fifth-order ED.

For this reason, we extend our previous analysis [8] in two directions to allow for the observation of higher-order EDs for inherited QEPs and QHPs. First, we weaken our requirements for the observation of \mathcal{PT} -symmetric dynamics by considering only subspace(s) of the whole Liouville space of the statistical operators. We note here that, owing to the linearity of quantum mechanics, we can equivalently describe the system dynamics [9, 10] in the Liouville space of the statistical operators and the complete space spanned by the operators of measurable quantities. We also note that the linearity gives the one-to-one correspondence between the subspaces of the above mentioned spaces. When such \mathcal{PT} -symmetric-like behavior is restricted to only a subspace we refer to nonconventional \mathcal{PT} -symmetric system behavior.

Second, we admit in our analysis more general non-Hermitian \mathcal{PT} -symmetric Hamiltonians. We recall here that, in Ref. [8], the non-Hermiticity of the investigated Hamiltonians originated only in the presence of damping and amplification terms, whose non-Hermiticity was 'remedied' by the presence of the Langevin fluctuating operator forces [10, 11]. This guarantees the system evolution preserving the bosonic canonical commutation relations. Here, we consider also the Hamiltonians that describe bosonic systems with unidirectional coupling between the modes. The reason is that unidirectional coupling allows to concatenate two bosonic subsystems such that they keep their original eigenvalues. Moreover the original subspaces belonging to the same eigenvalues merge together which results in the increased EDs. This property gives rise to the method suggested and elaborated in Refs. [12, 13] that provides QEPs with higher-order EDs. However, we note that, apart from a complex experimental realization, unidirectional coupling is highly non-Hermitian and violates reciprocity of physical processes. Nevertheless, up to our best knowledge, this is the only straightforward method for reaching

^{*} jan.perina.jr@upol.cz

[†] kishore.thapliyal@upol.cz

QEPs and QHPs with high-order EDs for open bosonic systems described correctly according to quantum mechanics. Higher-order QEPs using unidirectional coupling were already realized in [14].

We note that when Hamiltonians are directly considered, their higher-order EPs can relatively easily be observed. For example, higher-order EPs were predicted in optomechanical [15] and cavity magnonic systems [16], those described by the Bose-Hubbard model [17], or photonic structures [18, 19]. Higher-order EDs were also studied in Refs. [20, 21]. They play significant role in amplification [22] and sensing [23] as well as speeding up entanglement generation [24].

Increasing complexity of the bosonic systems poses the question about identification of inherited QEPs and QHPs and the determination of their degeneracies. Whereas simple bosonic systems allow for analytical derivation of the eigenvalues and eigenvectors of their dynamical matrices, more complex bosonic systems admit only numerical treatment. In this case, we may numerically decompose a given dynamical matrix into its Jordan form that directly reveals QEPs and QHPs with their degeneracies. Alternatively, we may add a little ϵ perturbation to any element of the dynamical matrix that can remove both EDs and DDs and even allow for distinguishing ED and DD.

Some of the effects related to the presence of QEPs and QHPs with higher-order EDs and DDs are observed also in the behavior of higher-order field-operator moments (FOMs) [9, 11, 25]. We note that we refer to genuine QEPs and QHPs in the case of higher-order FOMs. This originates from the fact that the dynamics of nthorder FOMs is built, in certain sense, as a 'multiplied dynamics' of first-order FOMs with its inherited QEPs and QHPs. This 'multiplied dynamics' then naturally contains the genuine QEPs and QHPs with higher-order EDs and DDs, as it was discussed in [25]. In general, genuine QEPs with ED orders up to nth power of ED orders of inherited QEPs of the first-order FOMs are expected in the dynamics of *n*th-order FOMs. However, the structure of genuine QEPs and QHPs in the dynamics of higherorder FOMs intimately depends on that of the inherited QEPs and QHPs found for the first-order FOMs. For this reason, we have explicitly revealed the structure of genuine QEPs and QHPs belonging to the second-order FOMs for the bosonic systems in tables of Ref. [8]. They explicitly elucidate the relation between EDs and DDs of these genuine QEPs and QHPs and degeneracies of inherited QEPs and QHPs. We note that induced QEPs and QHPs, which were introduced in Ref. [25], have their origin in the existence of identical or similar (related by the field commutation relations) FOMs in the formal construction of higher-order FOMs spaces and they further increase the multiplicity of spectral degeneracies. Nevertheless, they do not enrich the system dynamics. There exist some general properties of EDs and DDs of genuine QEPs and QHPs of bosonic systems independent of their configuration that we address here to complete the

analysis of specific bosonic systems.

The paper is organized as follows. Section II is devoted to bosonic systems exhibiting nonconventional \mathcal{PT} symmetric dynamics, An an example, four-mode systems are analyzed. Section III contains the analysis of bosonic systems with unidirectional coupling in linear configurations involving in turn from two to six modes. The dynamics of the two-mode bosonic system with unidirectional coupling and its applicability are analyzed in Sec. IV. Numerical methods for the identification of QEPs and QHPs and their degeneracies are discussed in Sec. V. A general analysis of genuine and induced QHPs in the dynamics of arbitrary-order FOMs is given in Sec. VI. Section VII brings conclusions. In Appendix A, the properties of the Langevin operator forces are discussed considering the model with unidirectional coupling. Statistical properties of the two-mode bosonic system with unidirectional coupling are described in Appendix B.

II. BOSONIC SYSTEMS WITH BIDIRECTIONAL COUPLING AND NONCONVENTIONAL \mathcal{PT} -SYMMETRIC DYNAMICS

When seeking for QEPs and QHPs in simple bosonic systems, we have observed the situations in which the behavior typical for \mathcal{PT} -symmetric systems occurs only in certain subspaces of the whole space spanned by the field operators and their moments. We speak about nonconventional \mathcal{PT} -symmetric dynamics in these subspaces. We note that we may alternatively specify the corresponding subspaces in the Liouville space of statistical operators [9]. As the conditions for the observation of nonconventional \mathcal{PT} -symmetric dynamics are less restrictive than those required for the usual \mathcal{PT} -symmetric dynamics, we analyze here simple bosonic systems exhibiting this form of dynamics from the point of view of the occurrence of higher-order QEPs and QHPs. In the following, we consider in turn four-mode bosonic systems in their circular and tetrahedral configurations (see Fig. 1).

A. Circular configuration

The Hamiltonian $\hat{H}_{4,c}$ of a four-mode bosonic system in the circular configuration depicted in Fig. 1(a) takes the following form :

$$\hat{H}_{4,c} = \begin{bmatrix} \hbar\epsilon \hat{a}_1^{\dagger}\hat{a}_2 + \hbar\epsilon \hat{a}_2^{\dagger}\hat{a}_3 + \hbar\epsilon \hat{a}_3^{\dagger}\hat{a}_4 + \hbar\epsilon \hat{a}_4^{\dagger}\hat{a}_1 + \hbar\kappa \hat{a}_1\hat{a}_2 \\ + \hbar\kappa \hat{a}_2\hat{a}_3 + \hbar\kappa \hat{a}_3\hat{a}_4 + \hbar\kappa \hat{a}_4\hat{a}_1 \end{bmatrix} + \text{H.c.}$$
(1)

where \hat{a}_j (\hat{a}_j^{\dagger}) for j = 1, ..., 4 denotes the annihilation (creation) operator of the *j*th mode, ϵ (κ) is the linear (nonlinear) coupling strength between the modes [26]. Symbol H.c. replaces the Hermitian-conjugated terms.



FIG. 1. Schematic diagrams of the four-mode bosonic systems in (a) circular and (b) tetrahedral configurations that exhibit QHPs in their nonconventional \mathcal{PT} -symmetric dynamics. Strengths ϵ and κ characterize, respectively, the linear and nonlinear coupling between the modes, while γ , with subscripts indicating the mode number(s), are the damping or amplification rates, and the annihilation operators \hat{a} identify the mode number via their subscripts.

Damping or amplification of mode j is described by the damping (amplification) rate γ_j and the corresponding Langevin stochastic operator forces, \hat{L}_j and \hat{L}_j^{\dagger} that occur in the dynamical Heisenberg-Langevin equations written below in Eq. (2). The Langevin stochastic operator forces are assumed to have the Markovian and Gaussian properties specific to the damping and amplification processes [25]. Their presence in the Heisenberg-Langevin equations guarantees the fulfillment of the fieldoperator commutation relations. Moreover the properties of the Langevin stochastic operator forces are related to the damping (amplification) rates via the fluctuationdissipation theorems [10, 11].

The Heisenberg-Langevin equations corresponding to the Hamiltonian $\hat{H}_{4,c}$ in Eq. (1) are written in the form:

$$\frac{d\hat{a}}{dt} = -iM_{c}^{(4)}\hat{a} + \hat{L}, \qquad (2)$$

where the vectors $\hat{\boldsymbol{a}}$ of field operators and $\hat{\boldsymbol{L}}$ of the Langevin operator forces are given as $\hat{\boldsymbol{a}} = [\hat{\boldsymbol{a}}_1, \hat{\boldsymbol{a}}_2, \hat{\boldsymbol{a}}_3, \hat{\boldsymbol{a}}_4]^T \equiv [\hat{\boldsymbol{a}}_1, \hat{\boldsymbol{a}}_1^{\dagger}, \hat{\boldsymbol{a}}_2, \hat{\boldsymbol{a}}_2^{\dagger}, \hat{\boldsymbol{a}}_3, \hat{\boldsymbol{a}}_3^{\dagger}, \hat{\boldsymbol{a}}_4, \hat{\boldsymbol{a}}_4^{\dagger}]^T$ and $\hat{\boldsymbol{L}} = [\hat{\boldsymbol{L}}_1, \hat{\boldsymbol{L}}_1^{\dagger}, \hat{\boldsymbol{L}}_2, \hat{\boldsymbol{L}}_2^{\dagger}, \hat{\boldsymbol{L}}_3, \hat{\boldsymbol{L}}_3^{\dagger}, \hat{\boldsymbol{L}}_4, \hat{\boldsymbol{L}}_4^{\dagger}]^T$. The dynamical matrix $\boldsymbol{M}_{\mathbf{c}}^{(4)}$ introduced in Eq. (2) is derived in the form

$$M_{\rm c}^{(4)} = \begin{bmatrix} -i\gamma_1 & \xi & 0 & \xi \\ \xi & -i\tilde{\gamma}_2 & \xi & 0 \\ 0 & \xi & -i\tilde{\gamma}_3 & \xi \\ \xi & 0 & \xi & -i\tilde{\gamma}_4 \end{bmatrix}.$$
 (3)

The 2 × 2 submatrices $\tilde{\gamma}_j$, j = 1, ..., 4, and $\boldsymbol{\xi}$ occurring in Eq. (3) are defined as:

$$\tilde{\gamma}_{j} = \begin{bmatrix} \gamma_{j}/2 & 0\\ 0 & \gamma_{j}/2 \end{bmatrix}, \quad \boldsymbol{\xi} = \begin{bmatrix} \epsilon & \kappa\\ -\kappa & -\epsilon \end{bmatrix}, \quad (4)$$

and γ_j stands for the damping or amplification rate of the mode j.

Applying the conditions

$$\gamma_1 = \gamma_3 \equiv \gamma_{13}, \quad \gamma_2 = \gamma_4 \equiv \gamma_{24}, \tag{5}$$

in the dynamical matrix $M_{c}^{(4)}$ in Eq. (3), we reveal its eigenvalues $\lambda_{j}^{M_{c}^{(4)}}$:

$$\lambda_{1}^{M_{c}^{(4)}} = -i\gamma_{13},$$

$$\lambda_{2}^{M_{c}^{(4)}} = -i\gamma_{24},$$

$$\lambda_{3,4}^{M_{c}^{(4)}} = -i\gamma_{+} \mp \beta.$$
(6)

The corresponding eigenvectors are derived as follows:

$$\begin{aligned} \boldsymbol{y_1^{M_c^{(4)}}} &= [-1, 0, 1, 0]^T, \\ \boldsymbol{y_2^{M_c^{(4)}}} &= [0, -1, 0, 1]^T, \\ \boldsymbol{y_3^{M_c^{(4)}}} &= \left[-\frac{2\xi}{\chi^*}, 1, -\frac{2\xi}{\chi^*}, 1 \right]^T, \\ \boldsymbol{y_4^{M_c^{(4)}}} &= \left[\frac{2\xi}{\chi}, 1, \frac{2\xi}{\chi}, 1 \right]^T, \end{aligned}$$
(7)

where $\chi = i\gamma_+ + \beta$, $\beta^2 = 4\xi^2 - \gamma_-^2$, and $4\gamma_{\pm} = \gamma_{13} \pm \gamma_{24}$. If $\beta = 0$ then $\lambda_3^{M_c^{(4)}} = \lambda_4^{M_c^{(4)}}$ and also $\boldsymbol{y_3^{M_c^{(4)}}} = \boldsymbol{y_4^{M_c^{(4)}}}$. We note that the eigenvalues $\lambda_3^{M_c^{(4)}}$ and $\lambda_4^{M_c^{(4)}}$ share also their imaginary parts, which is important for the observa-

tion of \mathcal{PT} -symmetric-like dynamics in a suitable interaction frame [27]. Provided that the system initial conditions are chosen such that only the eigenvalues $\lambda_3^{M_c^{(4)}}$ and $\lambda_4^{M_c^{(4)}}$ determine its dynamics, we observe a second-order QEP. This QEP changes into a QHP with second-order ED and DD when the 8×8 matrix $M_c^{(4)}$ is analyzed. The condition $\beta = 0$ transforms into the formula

$$\frac{\kappa^2}{\epsilon^2} + \frac{\gamma_-^2}{4\epsilon^2} = 1 \tag{8}$$

for an ellipse in the parameter space $(\kappa/\epsilon, \gamma_{-}/\epsilon)$ that identifies the positions of QHPs. Real parts of two eigenvalues $\lambda_{j}^{M_{c}^{(4)}}$, j = 3, 4, that form QHPs are plotted in this space in Fig. 2(a).

The diagonalized 8×8 dynamical matrix $M_c^{(4)}$ is obtained with the help of the eigenvalues and eigenvectors given in Eqs. (6) and (7), respectively, and the eigenvalues $\lambda_{1,2}^{\xi}$ and the eigenvectors $y_{1,2}^{\xi}$ of the matrix ξ :

$$\lambda_{1,2}^{\xi} = \mp \zeta, \tag{9}$$

$$\boldsymbol{y}_{1,2}^{\boldsymbol{\xi}} = \left[-\frac{\epsilon \mp \zeta}{\kappa}, 1\right]^{T}, \qquad (10)$$

where $\zeta = \sqrt{\epsilon^2 - \kappa^2}$. Details can be found in Appendix of Ref. [8]. In the basis with the diagonal 8×8 dynamical matrix $M_c^{(4)}$, the system dynamics is described by



FIG. 2. Real parts $\lambda^{\rm r}$ of the eigenvalues (a) $\lambda_{3,4}^{M_c^{(4)}}$ of the matrix $M_c^{(4)}$, given in Eq. (3), for the four-mode bosonic system in the circular configuration with different damping and/or amplification rates of neighbor modes and (b) $\lambda_{3,4}^{M_t^{(4)}}$ of the matrix $M_t^{(4)}$, given in Eq. (13), for $\xi = \pm \zeta$ for the four-mode bosonic system in the tetrahedral configuration with the same damping and/or amplification rates of neighbor modes are drawn in the parameter space ($\kappa/\epsilon, \gamma_-/\epsilon$). Dashed red curves indicate the positions of the QHPs and are given by Eq. (8).

the new field operators $\hat{\boldsymbol{b}} = \begin{bmatrix} \hat{b}_1, \hat{b}_1^{\dagger}, \hat{b}_2, \hat{b}_2^{\dagger}, \hat{b}_3, \hat{b}_4, \hat{b}_3^{\dagger}, \hat{b}_4^{\dagger} \end{bmatrix}^T$. If the initial conditions allow to describe the complete system dynamics in terms of the field operators \hat{b}_3 , \hat{b}_4 , \hat{b}_3^{\dagger} , and \hat{b}_4^{\dagger} with equal damping or amplification rate γ_+ then the first- and second-order FOMs exhibit in their dynamics QEPs and QHPs as summarized in Tab. I.

B. Tetrahedral configuration

In the tetrahedral configuration depicted in Fig. 1(b), the Hamiltonian $\hat{H}_{4,t}$ of four-mode bosonic system attains the form:

$$\hat{H}_{4,t} = \left[\hbar\epsilon \hat{a}_1^{\dagger}\hat{a}_2 + \hbar\epsilon \hat{a}_1^{\dagger}\hat{a}_3 + \hbar\epsilon \hat{a}_1^{\dagger}\hat{a}_4 + \hbar\epsilon \hat{a}_2^{\dagger}\hat{a}_3 + \hbar\epsilon \hat{a}_2^{\dagger}\hat{a}_4\right]$$

$$+\hbar\epsilon\hat{a}_{3}\dot{a}_{4} + \hbar\kappa\hat{a}_{1}\hat{a}_{2} + \hbar\kappa\hat{a}_{1}\hat{a}_{3} + \hbar\kappa\hat{a}_{1}\hat{a}_{4} + \hbar\kappa\hat{a}_{2}\hat{a}_{3}$$
$$+\hbar\kappa\hat{a}_{2}\hat{a}_{4} + \hbar\kappa\hat{a}_{3}\hat{a}_{4}] + \text{H.c.}$$
(11)

The Heisenberg-Langevin equations corresponding to the Hamiltonian $\hat{H}_{4,t}$ are derived in the form:

$$\frac{d\hat{a}}{dt} = -iM_{t}^{(4)}\hat{a} + \hat{L}$$
(12)

using the following dynamical matrix $M_{\rm t}^{(4)}$:

$$M_{\rm t}^{(4)} = \begin{bmatrix} -i\tilde{\gamma}_1 & \xi & \xi & \xi \\ \xi & -i\tilde{\gamma}_2 & \xi & \xi \\ \xi & \xi & -i\tilde{\gamma}_3 & \xi \\ \xi & \xi & \xi & -i\tilde{\gamma}_4 \end{bmatrix}.$$
 (13)

In seeking QEPs, we assume equal damping and/or amplification rates of modes 1 and 2, and also of modes 3 and 4:

$$\gamma_1 = \gamma_2 \equiv \gamma_{12}, \quad \gamma_3 = \gamma_4 \equiv \gamma_{34}. \tag{14}$$

We note that, due to the symmetry, identical results are obtained when assuming equal damping and/or amplification rates of modes 1 and 3 and also of modes 2 and 4 [compare Eq. (5)].

Under these conditions, diagonalization of the dynamical matrix $M_{\rm t}^{(4)}$ in Eq. (13) leaves us with the following eigenvalues:

$$\lambda_{1}^{M_{t}^{(4)}} = -i\gamma_{12} - \xi,$$

$$\lambda_{2}^{M_{t}^{(4)}} = -i\gamma_{34} - \xi,$$

$$\lambda_{3,4}^{M_{t}^{(4)}} = -i\gamma_{+} + \xi \mp \beta.$$
(15)

The corresponding eigenvectors are written as:

$$\mathbf{y}_{1}^{M_{t}^{(4)}} = [-1, 1, 0, 0]^{T}, \\
 \mathbf{y}_{2}^{M_{t}^{(4)}} = [0, 0, -1, 1]^{T}, \\
 \mathbf{y}_{3}^{M_{t}^{(4)}} = \left[1 - \frac{2i\gamma_{-}}{\chi_{-}}, 1 - \frac{2i\gamma_{-}}{\chi_{-}}, 1, 1\right]^{T}, \\
 \mathbf{y}_{4}^{M_{t}^{(4)}} = \left[1 - \frac{2i\gamma_{-}}{\chi_{+}}, 1 - \frac{2i\gamma_{-}}{\chi_{+}}, 1, 1\right]^{T}, \quad (16)$$

and $\chi_{\pm} = i\gamma_{-} \pm \beta + 2\xi$, $\beta^{2} = 4\xi^{2} - \gamma_{-}^{2}$, and $4\gamma_{\pm} = \gamma_{12} \pm \gamma_{34}$.

Provided that $\beta = 0$, we have $\lambda_3^{M_t^{(4)}} = \lambda_4^{M_t^{(4)}}$ and $y_3^{M_t^{(4)}} = y_4^{M_t^{(4)}}$ as $\chi_- = \chi_+$. As the imaginary parts of eigenvalues $\lambda_{1,2}^{M_t^{(4)}}$ differ from those of $\lambda_{3,4}^{M_t^{(4)}}$, the system can exhibit only the non-conventional \mathcal{PT} -symmetric dynamics: If the system initial conditions are such that only the eigenvalues $\lambda_3^{M_t^{(4)}}$ and $\lambda_4^{M_t^{(4)}}$ suffice in describing its dynamics, we observe a second-order QEP for the 4×4 dynamical matrix $M_t^{(4)}$. As the eigenvalues $\lambda_j^{M_t^{(4)}}$ in

Λ_j^{i}	Λ_j^r	Momer	nts	Moment	Genuine a	nd induced QHPs	Genuine QHPs	
				deg.	Partial	Partial	Partial	Partial
					QDP x	QDP x	QDP x	QDP x
					$\ensuremath{\operatorname{QEP}}$ deg.	QEP deg.	QEP deg.	$\ensuremath{\operatorname{QEP}}$ deg.
γ_+	$\pm\beta$	$\langle \hat{b}_3 angle, \langle \hat{b}_3^\dagger angle$	$\langle \hat{B}_5 angle$	1	1x2	2x2	1x2	2x2
		$\langle \hat{b}_4 \rangle, \langle \hat{b}_4^\dagger \rangle$	$\langle \hat{B}_6 angle$	1	1x2		1x2	
$2\gamma_+$	$\pm 2\beta$	$\langle \hat{b}_3 \hat{b}_4 \rangle, \langle \hat{b}_3^\dagger \hat{b}_4^\dagger \rangle$	$\langle \hat{B}_5 \hat{B}_6 angle$	2	2x4	4x4	1x4	1x4
	$\beta - \beta$	$\langle \hat{b}_3^\dagger \hat{b}_4 angle$		2				+
	$\beta - \beta$	$\langle \hat{b}_3 \hat{b}_4^\dagger angle$		2				
	$\pm 2\beta$	$\langle \hat{b}_3^2 \rangle, \langle \hat{b}_3^{\dagger 2} \rangle$	$\langle \hat{B}_5^2 angle$	1	1x4		1x3	2x3
	$\beta - \beta$	$\langle \hat{b}_3^\dagger \hat{b}_3 angle$		2				
	$\pm 2\beta$	$\langle \hat{b}_4^2 \rangle, \langle \hat{b}_4^{\dagger 2} \rangle$	$\langle \hat{B}_{6}^{2} angle$	1	1x4		1x3	
	$\beta - \beta$	$\langle \hat{b}_4^\dagger \hat{b}_4 angle$		2				

TABLE I. Real and imaginary parts of the complex eigenfrequencies $\Lambda_j^{\rm r} - i\Lambda_j^{\rm i}$ of the matrix $M_{\rm c}^{(4)}$, given in Eq. (3). for the four-mode bosonic system in the circular configuration with different damping and/or amplification rates of neighbor modes valid for the regime of nonconventional \mathcal{PT} -symmetric dynamics and derived from the equations for the FOMs up to second order. The corresponding moments written in the 'diagonalized' field operators involving the operators \hat{b}_3 , \hat{b}_3^{\dagger} , \hat{b}_4 , and \hat{b}_4^{\dagger} are given together with their degeneracies (deg.) coming from different possible relative positions of the field operators. The DDs of QHPs (partial DDs) derived from the indicated FOMs and the EDs of the constituting QEPs are given. Both genuine and induced QEPs and QHPs are considered. The operators \hat{B}_j for j = 5, 6 are defined in the lines written for $\Lambda_i^i = \gamma_+$.

Eq. (15) show the linear dependence on ξ , the diabolical second-order degeneracy of the 8×8 dynamical matrix $M_{\rm c}^{(4)}$, originating in the form of the eigenvalues $\lambda_j^{M_{\rm c}^{(4)}}$ in Eq. (6), is not observed in the tetrahedral configuration. Instead, for $\beta = 0$, we find one second-order QEP for $\xi = \zeta$ and another second-order QEP for $\xi = -\zeta$. These QEPs occur at the positions described in Eq. (8) in the parameter space $(\kappa/\epsilon, \gamma_{-}/\epsilon)$. Real parts of four eigenvalues $\lambda_{3,4}^{M_{\rm t}^{(4)}}$ for $\xi = \pm \zeta$ that build two QEPs are drawn in this space in Fig. 2(b).

In the basis with the diagonal 8×8 dynamical matrix $M_{t}^{(4)}$, the system dynamics is described by the new field operators $\hat{\boldsymbol{b}} = \left[\hat{b}_{1}, \hat{b}_{1}^{\dagger}, \hat{b}_{2}, \hat{b}_{2}^{\dagger}, \hat{b}_{3}, \hat{b}_{4}, \hat{b}_{4}^{\dagger}, \hat{b}_{3}^{\dagger}\right]^{T}$. The corresponding eigenvalues and eigenvectors are discussed in general in Appendix of Ref. [8]. The first- and second-order FOMs exhibit in their dynamics QEPs and QHPs that are summarized in Tab. II.

III. CONCATENATED BOSONIC SYSTEMS WITH UNIDIRECTIONAL COUPLING: HIGHER-ORDER QUANTUM EXCEPTIONAL POINTS ON DEMAND

The previous analysis of the bosonic systems with up to five modes in their linear, circular, tetrahedron, and pyramid configurations revealed only the inherited QEPs and QHPs with second- and third-order EDs. Here, we extend our analysis to more general non-Hermitian Hamiltonians that involve unidirectional coupling between the modes. Whereas the non-Hermiticity of the above discussed systems is given solely by the presence of damping and/or amplification, the Hamiltonians with unidirectional coupling are non-Hermitian per se, while damping and amplification make them even more non-Hermitian. Despite their non-Hermiticity, they have direct physical implementations based upon counter-directional field propagation and mutual scattering [12, 13]. Moreover, it has been shown in Ref. [28] that exponential improvement of measurement precision can be reached in QEPs in systems with unidirectional coupling.

It was shown in Refs. [12, 13] that using a specific unidirectional coupling of two field modes belonging to different \mathcal{PT} -symmetric systems with QEPs, the combined system exhibits a QEP with ED given as the sum of those of the constituting systems. This opens the door for observing inherited QEPs with EDs of orders higher than three.

To demonstrate the method, let us consider two bosonic modes that are unidirectionally coupled via the matrix $\boldsymbol{\xi}$ defined in Eq. (4) [see the scheme in Fig. 3(a)]. As we demonstrate below, this unidirectional coupling creates a QEP. The corresponding Heisenberg-Langevin equations take the form

$$\frac{d\hat{a}}{dt} = -iM_{\mathbf{u}}^{(1+1)}\hat{a} + \hat{L}, \qquad (17)$$

where the vectors $\hat{\boldsymbol{a}}$ of field operators and $\hat{\boldsymbol{L}}$ of the Langevin operator forces are given as $\hat{\boldsymbol{a}} = [\hat{\boldsymbol{a}}_1, \hat{\boldsymbol{a}}_2]^T \equiv \begin{bmatrix} \hat{a}_1, \hat{a}_1, \hat{a}_2, \hat{a}_2^{\dagger} \end{bmatrix}^T$ and $\hat{\boldsymbol{L}} = \begin{bmatrix} \hat{L}_1, \hat{L}_1^{\dagger}, \hat{L}_2, \hat{L}_2^{\dagger} \end{bmatrix}^T$. The properties of the Langevin operator forces are described in detail below, i.e. in Eq. (68).

The dynamical matrix $M_{\rm u}^{(1+1)}$ introduced in Eq. (17) and corresponding to unidirectional coupling of modes is written as

$$M_{\mathbf{u}}^{(1+1)} = \begin{bmatrix} -i\tilde{\gamma}_1 & \mathbf{0} \\ \boldsymbol{\xi} & -i\tilde{\gamma}_2 \end{bmatrix}, \quad (18)$$

Λ_j^{i}	Λ_j^{r}	Momen	nts	Moment	Genuine a	nd induced QHPs	Genuine QHPs	
				deg.	Partial	Partial	Partial	Partial
					QDP x	QDP x	QDP x	QDP x
					QEP deg.	QEP deg.	QEP deg.	QEP deg.
γ_+	$-\zeta \pm \beta$	$\langle \hat{b}_3 \rangle, \langle \hat{b}_4^\dagger \rangle$	$\langle \hat{B}_5 angle$	1	1x2	1x2	1x2	1x2
	$\zeta \pm \beta$	$\langle \hat{b}_4 \rangle, \langle \hat{b}_3^\dagger \rangle$	$\langle \hat{B}_{6} angle$	1	1x2	1x2	1x2	1x2
$2\gamma_+$	$\pm 2\beta$	$\langle \hat{b}_3 \hat{b}_4 \rangle, \langle \hat{b}_4^\dagger \hat{b}_3^\dagger \rangle$	$\langle \hat{B}_5 \hat{B}_6 angle$	2	2x4	2x4	1x4	1x4
	$\pm 2\zeta$	$\langle \hat{b}_3 \hat{b}_3^\dagger \rangle, \langle \hat{b}_4^\dagger \hat{b}_4 \rangle$		2				
	$-2\zeta \pm 2\beta$	$\langle \hat{b}_3^2 \rangle, \langle \hat{b}_4^{\dagger 2} \rangle$	$\langle \hat{B}_5^2 angle$	1	1x4	1x4	1x3	1x3
	-2ζ	$\langle \hat{b}_4^\dagger \hat{b}_3 angle$		2				
	$2\zeta \pm 2\beta$	$\langle \hat{b}_4^2 \rangle, \langle \hat{b}_3^{\dagger 2} \rangle$	$\langle \hat{B}_{6}^{2} angle$	1	1x4	1x4	1x3	1x3
	2ζ	$\langle \hat{b}_3^\dagger \hat{b}_4 angle$		2				

TABLE II. Real and imaginary parts of the complex eigenfrequencies $\Lambda_j^{\rm r} - i\Lambda_j^{\rm i}$ of the matrix $M_t^{(4)}$, given in Eq. (13), for the four-mode bosonic system in the tetrahedral configuration valid for the regime of nonconventional \mathcal{PT} -symmetric dynamics and derived from the equations for the FOMs up to second order. For details, see the caption to Tab. I.



FIG. 3. Schematic diagrams of bosonic systems composed of two subsystems mutually coupled by unidirectional coupling and having: (a) two, (b) three, (c) four, (d) five, and (e) six modes in typical linear configurations that exhibit QHPs. The coupling strengths ϵ and κ characterize both unidirectional and bidirectional coupling between modes, γ , with subscripts indicating the mode number(s), give the damping or amplification rates, and the annihilation operators \hat{a} identify the mode number via their subscripts.

using the damping or amplification 2×2 submatrices $\tilde{\gamma}_j$, j = 1, 2, defined in Eq. (4). The eigenvalues and eigenvectors of the 2×2 matrix $M_{\mathbf{u}}^{(1+1)}$ are determined as

and

$$\lambda_{1,2}^{M_{1}^{(1+1)}} = -i\gamma_{1,2}/2, \tag{19}$$

$$y_{1}^{M_{u}^{(1+1)}} = \left[-\frac{\gamma_{1} - \gamma_{2}}{2\xi}, 1 \right]^{T},$$

$$y_{2}^{M_{u}^{(1+1)}} = \left[0, 1 \right]^{T}.$$
 (20)

Provided that

$$\gamma_1 = \gamma_2, \tag{21}$$

we have $\lambda_1^{M_u^{(1+1)}} = \lambda_2^{M_u^{(1+1)}}$ together with $\boldsymbol{y_1^{M_u^{(1+1)}}} = \boldsymbol{y_2^{M_u^{(1+1)}}}$, and so we observe the creation of a QEP with second-order ED. This means that the 4 × 4 matrix $M_u^{(1+1)}$ obtained after inserting the 2 × 2 submatrices $\tilde{\gamma}_j$, j = 1, 2, and $\boldsymbol{\xi}$ into Eq. (18) exhibits a QHP with second-order ED and DD. It is worth noting that these QHPs occur independently of the values of the coupling strengths ϵ and κ , in strike difference to the QHPs found

in the two-mode bosonic system analyzed in Sec. II of Ref. [8].

In the next step we demonstrate the increase of the order of ED of a QEP by considering a single-mode bosonic system unidirectionally coupled to a two-mode bosonic system with a QEP [see the scheme in Fig. 4(b)]. The dynamical matrix $M_{\rm u}^{(1+2)}$ of the concatenated system is given as:

$$M_{\mathbf{u}}^{(1+2)} = \begin{bmatrix} -i\tilde{\gamma}_1 & \mathbf{0} & \mathbf{0} \\ \boldsymbol{\xi} & -i\tilde{\gamma}_2 & \boldsymbol{\xi} \\ \mathbf{0} & \boldsymbol{\xi} & -i\tilde{\gamma}_3 \end{bmatrix}. \quad (22)$$

Provided that

$$\gamma_1 = (\gamma_2 + \gamma_3)/2 \equiv 2\gamma_+, \tag{23}$$

we arrive at the eigenvalues:

$$\lambda_{1}^{M_{u}^{(1+2)}} = -i\gamma_{+},$$

$$\lambda_{2,3}^{M_{u}^{(1+2)}} = -i\gamma_{+} \mp \beta,$$
 (24)

and the corresponding eigenvectors:

$$\boldsymbol{y_{1}^{M_{u}^{(1+2)}}} = \left[\frac{\beta^{2}}{\xi^{2}}, \frac{i\gamma_{-}}{\xi}, 1\right]^{T},$$
$$\boldsymbol{y_{2,3}^{M_{u}^{(1+2)}}} = \left[0, \frac{-i\gamma_{-} \mp \beta}{\xi}, 1\right]^{T}, \qquad (25)$$

where $\beta^2 = \xi^2 - \gamma_-^2$ and $4\gamma_{\pm} = \gamma_2 \pm \gamma_3$. If $\beta = 0$, i.e., when the condition

$$\frac{\kappa^2}{\epsilon^2} + \frac{\gamma_-^2}{\epsilon^2} = 1 \tag{26}$$

for the constituting two-mode system is fulfilled, we have $\lambda_1^{M_u^{(1+2)}} = \lambda_2^{M_u^{(1+2)}} = \lambda_3^{M_u^{(1+2)}}$, and $y_1^{M_u^{(1+2)}} = y_2^{M_u^{(1+2)}} = y_3^{M_u^{(1+2)}}$. This identifies a QEP with third-order ED that emerged from the original QEP with second-order ED. This means that we have a QHP with third-order ED and second-order DD in the 6×6 matrix $M_u^{(1+2)}$. Introducing the new field operators, in which the dynamical matrix $M_u^{(1+2)}$ attains its diagonal form, we reveal QEPs and QHPs and their degeneracies appropriate to the dynamics of the first- and second-order FOMs. They can be found in Tab. III derived for a general *n*-mode bosonic system exhibiting an inherited QHP with *n*th-order ED and second-order DD for n > 2. We note that Tab. III applies also to the bosonic systems analyzed below.

Now, we construct four-, five-, and six-mode bosonic systems by concatenating the two- and three-mode systems with second- and third-order QEPs analyzed in Secs. II and III of Ref. [8]. We begin with combining two two-mode systems [see the scheme in Fig. 4(c)] whose dynamical matrices denoted as $M^{(2)}(\tilde{\gamma}_1, \tilde{\gamma}_2, \xi)$ and $M^{(2)}(\tilde{\gamma}_3, \tilde{\gamma}_4, \xi)$ are defined as follows [compare Eq. (3) of part I of the paper]:

$$M^{(2)}(\tilde{\gamma}_1, \tilde{\gamma}_2, \xi) = \begin{bmatrix} -i\tilde{\gamma}_1 & \xi \\ \xi & -i\tilde{\gamma}_2 \end{bmatrix}.$$
(27)

The joint dynamical matrix $M_{\rm u}^{(2+2)}$, defined as

$$M_{\rm u}^{(2+2)} = \begin{bmatrix} M^{(2)}(\tilde{\gamma}_1, \tilde{\gamma}_2, \xi) & 0\\ \Upsilon^{(2+2)} & M^{(2)}(\tilde{\gamma}_3, \tilde{\gamma}_4, \xi) \end{bmatrix},$$
(28)

includes unidirectional coupling between modes 2 and 3 described by the submatrix $\Upsilon^{(2+2)}$:

$$\Upsilon^{(2+2)} = \begin{bmatrix} 0 & \xi \\ 0 & 0 \end{bmatrix}.$$
 (29)

Assuming

$$\gamma_1 = \gamma_3 \equiv \gamma_{13}, \quad \gamma_2 = \gamma_4 \equiv \gamma_{24}, \tag{30}$$

we reveal the eigenvalues of the 4×4 matrix $M_{\rm u}^{(2+2)}$ as follows:

$$\lambda_{1,2}^{M_{u}^{(2+2)}} = -i\gamma_{+} \mp \beta, \lambda_{3,4}^{M_{u}^{(2+2)}} = -i\gamma_{+} \mp \beta,$$
(31)

using $\beta^2 = \xi^2 - \gamma_-^2$ and $4\gamma_{\pm} = \gamma_{13} \pm \gamma_{24}$. For $\beta = 0$, a QEP with fourth-order ED is found at the positions in the parameter space given by Eq. (26), i.e., where the constituting two-mode systems form QEPs with second-order EDs. The fourth-order QEP implies a QHP with fourth-order ED and second-order DD of the 8×8 matrix $M_{\rm u}^{(2+2)}$. It is worth noting that the additional unidirectional coupling of modes 1 and 4, i.e. when $\tilde{\Upsilon}^{(2+2)} = \begin{bmatrix} 0 & \xi \\ \xi & 0 \end{bmatrix}$, does not change the eigenvalues in Eq. (30) and also preserves the structure of eigenvectors with the discussed QEPs and QHPs.

The unidirectional coupling of the two- and three-mode systems allows to observe a QEP with fifth-order ED [for the configuration, see Fig. 4(d)]. The corresponding dynamical matrix $M_{\rm u}^{(2+3)}$ combines the two-mode dynamical matrix $M^{(2)}$ in Eq. (27) and the three-mode dynamical matrix $M^{(3)}$ [compare Eq. (19) in [8]],

$$M^{(3)}(\tilde{\gamma}_1, \tilde{\gamma}_2, \tilde{\gamma}_3, \xi) = \begin{bmatrix} -i\tilde{\gamma}_1 & \xi & 0\\ \xi & -i\tilde{\gamma}_2 & \xi\\ 0 & \xi & -i\tilde{\gamma}_3 \end{bmatrix}. (32)$$

The matrix $M_{\mathrm{u}}^{(2+3)}$ is expressed as

$$M_{\rm u}^{(2+3)} = \begin{bmatrix} M^{(2)}(\tilde{\gamma}_1, \tilde{\gamma}_2, \xi) & 0\\ \Upsilon^{(2+3)} & M^{(3)}(\tilde{\gamma}_3, \tilde{\gamma}_4, \tilde{\gamma}_5, \xi) \end{bmatrix},$$
(33)

Λ_{i}^{i}	Λ_{i}^{r}	Moments		Moment	Genuine and induced QHPs		Genuine QHPs	
5	5			deg.	Partial	Partial	Partial	Partial
					QDP x	QDP x	QDP x	QDP x
					QEP deg.	QEP deg.	QEP deg.	QEP deg.
γ_+	$\pm\beta_1,\ldots,\pm\beta_n$	$\langle \hat{b}_1 \rangle, \langle \hat{b}_1^{\dagger} \rangle, \dots, \langle \hat{b}_{n/2} \rangle, \langle \hat{b}_{n/2}^{\dagger} \rangle$	$\langle \hat{B}_1 angle$	1	1xn	2xn	1xn	2xn
		$\langle \hat{b}_{n/2+1} \rangle, \langle \hat{b}_{n/2+1}^{\dagger} \rangle, \dots, \langle \hat{b}_n \rangle, \langle \hat{b}_n^{\dagger} \rangle$	$\langle \hat{B}_{2} angle$	1	1xn		1xn	
$2\gamma_+$	$\pm(\beta_k+\beta_l)$	$\langle \hat{b}_k \hat{b}_{n/2+l} angle, \langle \hat{b}_k^\dagger \hat{b}_{n/2+l}^\dagger angle$	$\langle \hat{B}_1 \hat{B}_2 angle$	2	$2xn^2$	$4xn^2$	$1 \mathbf{x} n^2$	$1 \mathbf{x} n^2$
	$\beta_l - \beta_k$	$\langle \hat{b}_k^{\dagger} \hat{b}_{n/2+l} angle$		2				+
	$\beta_k - \beta_l$	$\langle \hat{b}_k \hat{b}_{n/2+l}^\dagger angle$		2				
		$k, l = 1, \dots n/2$						
	$\pm 2\beta_k$	$\langle \hat{b}_k^2 angle, \langle \hat{b}_k^{\dagger 2} angle$	$\langle \hat{B}_1^2 angle$	1	$1 \mathrm{x} n^2$		$1 \mathrm{x} \ (n+1)n/2$	$2{\rm x}~(n+1)n/2$
		$k = 1, \dots, n/2$						
	$\pm(\beta_k+\beta_l)$	$\langle b_k b_l \rangle, \langle b_k^{\dagger} b_l^{\dagger} \rangle$		2				
	0 0	$k, l = 1, \dots n/2, l < k$						
	$\beta_l - \beta_k$	$\langle b_k b_l \rangle$ k l = 1 n/2		2				
	$\pm 2\beta_k$	$\langle \hat{b}^2 \rangle_{(2,1,1)} \rangle, \langle \hat{b}^{\dagger 2} \rangle_{(2,1,1)} \rangle$	$\langle \hat{B}_2^2 \rangle$	1	$1 \mathrm{x} n^2$		1x (n+1)n/2	
		$k = 1, \dots n/2$	× 2/					
	$\pm(\beta_k+\beta_l)$	$\langle \hat{b}_{n/2+k} \hat{b}_{n/2+l} \rangle, \langle \hat{b}_{n/2+k}^{\dagger} \hat{b}_{n/2+l}^{\dagger} \rangle$		2				
		$k,l=1,\ldots n/2, l< k$						
	$\beta_l - \beta_k$	$\langle \hat{b}_{n/2+k}^{\dagger}\hat{b}_{n/2+l} angle$		2				
		$k, l=1, \dots n/2$						

TABLE III. Real and imaginary parts of the complex eigenfrequencies $\Lambda_j^{\rm r} - i\Lambda_j^{\rm i}$ of the matrix $\boldsymbol{M}_{\rm u}^{(n)}$ for *n*-mode bosonic system (n > 2) with unidirectional coupling having a QHP with *n*th-order ED and second-order DD derived from the equations for the FOMs up to second order. Table is valid for even *n*, where the vector $\hat{\boldsymbol{b}}$ of the diagonalized field operators is written as $\hat{\boldsymbol{b}} = [\hat{b}_1, \hat{b}_2, \hat{b}_1^{\dagger}, \hat{b}_2^{\dagger}, \dots \hat{b}_{n-1}, \hat{b}_n, \hat{b}_{n-1}^{\dagger}, \hat{b}_n^{\dagger}]$. The vector $\hat{\boldsymbol{b}}$ attains the form $\hat{\boldsymbol{b}} =$ $[\hat{b}_1, \hat{b}_1^{\dagger}, \dots, \hat{b}_m, \hat{b}_m^{\dagger}, \dots, \hat{b}_{m+1}, \hat{b}_{m+2}, \hat{b}_{m+1}^{\dagger}, \hat{b}_{m+2}^{\dagger}, \dots \hat{b}_{n-1}, \hat{b}_n, \hat{b}_{n-1}^{\dagger}, \hat{b}_n^{\dagger}]$ if there exist *m* un-paired eigenvalues in the subsystems that compose the analyzed bosonic system [see, e.g., Eqs. (23), (47), and (54) in Ref. [8]]. In such cases, the columns entitled Moments have to be modified accordingly, but all other columns remain valid. For details, see the caption of Tab. I.

assuming the unidirectional coupling between modes 2 and 3:

$$\boldsymbol{\Upsilon}^{(2+3)} = \begin{bmatrix} \mathbf{0} & \boldsymbol{\xi} \\ \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix}.$$
 (34)

Provided that

$$\gamma_1 + \gamma_2 = \gamma_3 + \gamma_5, \qquad 2\gamma_4 = \gamma_3 + \gamma_5, \qquad (35)$$

we determine the eigenvalues of the 5×5 matrix $M_{\rm u}^{(2+3)}$ as follows:

$$\lambda_{1}^{M_{u}^{(2+3)}} = -i\gamma_{+},
\lambda_{2,3}^{M_{u}^{(2+3)}} = -i\gamma_{+} \mp \bar{\beta},
\lambda_{4,5}^{M_{u}^{(2+3)}} = -i\gamma_{+} \mp \beta,$$
(36)

where $\beta^2 = \xi^2 - \gamma_-^2$, $\bar{\beta}^2 = 2\xi^2 - \bar{\gamma}_-^2$, $4\gamma_{\pm} = \gamma_1 \pm \gamma_2$, and $4\bar{\gamma}_- = \gamma_3 - \gamma_5$. Provided that $\beta = 0$ and $\bar{\beta} = 0$, we find a QEP with fifth-order ED. These conditions define the positions, which are specified in Eq. (26) and the following one:

$$\frac{\kappa^2}{\epsilon^2} + \frac{\bar{\gamma}_-^2}{2\epsilon^2} = 1. \tag{37}$$

Both conditions have to be fulfilled simultaneously, which results in the following condition

$$\gamma_{-} = \bar{\gamma}_{-} / \sqrt{2}. \tag{38}$$

Thus, we reveal the QHPs of the 10×10 matrix $M_{\rm u}^{(2+3)}$ under the conditions, given in Eqs. (35) and (38), observed in the space $(\kappa/\epsilon, \gamma_-/\epsilon)$ at the positions obeying Eq. (26).

The last analyzed system is created by unidirectional combining of two three-mode systems with identical parameters in the configuration shown in Fig. 4(e). Its dynamical matrix, denoted as $M_{\rm u}^{(3+3)}$, is composed of the two three-mode dynamical matrices $M^{(3)}$ from Eq. (32) and the coupling matrix $\Upsilon^{(3+3)}$,

$$\Upsilon^{(3+3)} = \begin{bmatrix} 0 & 0 & \xi \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$
 (39)

It is expressed in the form

$$M_{\rm u}^{(3+3)} = \begin{bmatrix} M^{(3)}(\tilde{\gamma}_1, \tilde{\gamma}_2, \tilde{\gamma}_3, \xi) & 0\\ \Upsilon^{(3+3)} & M^{(3)}(\tilde{\gamma}_4, \tilde{\gamma}_5, \tilde{\gamma}_6, \xi) \end{bmatrix},$$
(40)

in which we assume:

$$\gamma_1 = \gamma_4 \equiv \gamma_{14}; \quad \gamma_3 = \gamma_6 \equiv \gamma_{36}, \\ 2\gamma_2 = 2\gamma_4 = \gamma_{14} + \gamma_{36}.$$
 (41)

The eigenvalues of the 6×6 matrix $M_{\rm u}^{(3+3)}$ are determined in the form:

$$\lambda_{1,2}^{M_{\rm u}^{(3+3)}} = -i\gamma_+$$

$$\lambda_{3,4}^{M_{u}^{(3+3)}} = -i\gamma_{+} \mp \beta,$$

$$\lambda_{5,6}^{M_{u}^{(3+3)}} = -i\gamma_{+} \mp \beta,$$
 (42)

where $\beta^2 = 2\xi^2 - \gamma_-^2$ and $4\gamma_{\pm} = \gamma_{14} \pm \gamma_{36}$. It holds $\beta = 0$ at the positions in the parameter space $(\kappa/\epsilon, \gamma_-/\epsilon)$ that fulfil the condition

$$\frac{\kappa^2}{\epsilon^2} + \frac{\gamma_-^2}{2\epsilon^2} = 1. \tag{43}$$

At these positions we have six identical eigenvalues in Eq. (42). They indicate a QEP with sixth-order ED that implies a QHP of sixth-order ED and second-order DD in the dynamical 12×12 matrix $M_{\rm u}^{(3+3)}$.

In the last three analyzed systems, explicit forms for the eigenvectors of the dynamical matrices $M_{\rm u}^{(2+2)}$, $M_{\rm u}^{(2+3)}$, and $M_{\rm u}^{(3+3)}$ were not analyzed because of their complexity. Instead, these dynamical matrices were diagonalized under the conditions at which QEPs are expected and the transformed matrices in the Jordan form with unit elements at the upper diagonal confirmed the presence of QEPs.

The considered forms of unidirectional coupling matrices $\Upsilon^{(2+2)}$, $\Upsilon^{(2+3)}$, and $\Upsilon^{(3+3)}$ can be replaced by those connecting different pairs of modes in the constituting systems. This replacement does not change the eigenvalues as well as the degeneracies of the eigenvectors that give rise to the observed QEPs and QHPs. We note that the form of eigenvectors depends on which pairs of modes are unidirectionally coupled. We can even include unidirectional coupling of several pairs of modes in the constituting systems and this property still holds. The only requirement is that all couplings point out from one subsystem to the other subsystem.

We note that there exist alternative ways to realize unidirectional coupling between the constituting bosonic systems. Contrary to the considered unidirectional coupling that is described directly via the dynamical matrices, we may consider an alternative form of unidirectional coupling characterized by the Hamiltonians

$$\hat{H}_{c-u,1} = \hbar \epsilon \hat{a}_i \hat{a}_j^{\dagger} + \hbar \kappa \hat{a}_i \hat{a}_j, \qquad (44)$$

$$\hat{H}_{c-u,2} = \hbar \epsilon \hat{a}_i \hat{a}_j^{\dagger} + \hbar \kappa \hat{a}_i^{\dagger} \hat{a}_j^{\dagger}.$$

$$\tag{45}$$

They result in the system dynamics similar to that discussed above and lead to the eigenvalues and eigenvectors with characteristic EDs and DDs. This improves the feasibility of practical realizations of such concatenated bosonic systems with higher-order QEPs and QHPs based on unidirectional coupling.

The method is general, allowing for concatenating simple bosonic systems into more complex ones that keep QEPs and QHPs of the original systems using unidirectional coupling of different kinds. Combing the analyzed bosonic systems with second- and third-order EDs together, bosonic systems with arbitrary-order EDs can be achieved.

IV. UNIDIRECTIONAL COUPLING AND ITS APPLICABILITY

In this section, we reveal characteristic features of the bosonic systems with unidirectional coupling and specify the conditions of their applicability. We consider the simplest two-mode bosonic system with unidirectional coupling whose dynamics is described by the Heisenberg-Langevin equations written in Eq. (17) with the dynamical matrix $M_{\rm u}^{(1+1)}$ given in Eq. (18). We assume the most typical configuration of \mathcal{PT} -symmetric systems in which mode 1 is damped and mode 2 is amplified:

$$\gamma_1 = -\gamma_2 \equiv 2\gamma. \tag{46}$$

The corresponding Langevin fluctuating operator forces embedded in the vector \hat{L} are modelled by two independent quantum random Gaussian processes [29–31]. This results in the following correlation functions:

$$\langle \hat{L}_1(t) \rangle = \langle \hat{L}_1^{\dagger}(t) \rangle = 0, \quad \langle \hat{L}_2(t) \rangle = \langle \hat{L}_2^{\dagger}(t) \rangle = 0, \langle \hat{L}_1^{\dagger}(t) \hat{L}_1(t') \rangle = 0, \quad \langle \hat{L}_1(t) \hat{L}_1^{\dagger}(t') \rangle = 2\gamma \delta(t - t'), \langle \hat{L}_2^{\dagger}(t) \hat{L}_2(t') \rangle = 2\gamma \delta(t - t'), \quad \langle \hat{L}_2(t) \hat{L}_2^{\dagger}(t') \rangle = 0.$$
 (47)

The remaining second-order correlation functions are zero. Symbol δ stands for the Dirac function.

The solution to the stochastic linear differential operator equations in Eq. (17) is expressed in the form

$$\hat{a}(t) = P(t,0)\hat{a}(0) + \hat{F}(t)$$
 (48)

using the evolution matrix \boldsymbol{P} defined as

$$\mathbf{P}(t,t') = \exp[-i\mathbf{M}_{\mathbf{u}}^{(1+1)}(t-t')].$$
(49)

The fluctuating operator forces \hat{F} introduced in Eq. (48) are determined along the formula

$$\hat{\boldsymbol{F}}(t) = \int_0^t dt' \boldsymbol{P}(t,t') \hat{\boldsymbol{L}}(t').$$
(50)

It implies the following formula for their second-order correlation functions,

$$\begin{aligned} \langle \hat{\boldsymbol{F}}(t) \hat{\boldsymbol{F}}^{\dagger \boldsymbol{T}}(t) \rangle &= \int_{0}^{t} d\tilde{t} \int_{0}^{t} d\tilde{t}' \\ \boldsymbol{P}(t, \tilde{t}) \langle \hat{\boldsymbol{L}}(\tilde{t}) \hat{\boldsymbol{L}}^{\dagger \boldsymbol{T}}(\tilde{t}') \rangle \boldsymbol{P}^{\dagger \boldsymbol{T}}(t, \tilde{t}'). \end{aligned} (51)$$

The solution in Eq. (48) can be recast into a simpler form written for the annihilation operators \hat{a}_1 and \hat{a}_2 :

$$\begin{bmatrix} \hat{a}_1(t) \\ \hat{a}_2(t) \end{bmatrix} = \boldsymbol{U}(t) \begin{bmatrix} \hat{a}_1(0) \\ \hat{a}_2(0) \end{bmatrix} + \boldsymbol{V}(t) \begin{bmatrix} \hat{a}_1^{\dagger}(0) \\ \hat{a}_2^{\dagger}(0) \end{bmatrix} + \begin{bmatrix} \hat{f}_1(t) \\ \hat{f}_2(t) \end{bmatrix}.$$
(52)

The elements of the matrices U and V are defined as $U_{j,k}(t) = P_{2j-1,2k-1}(t,0)$ and $V_{j,k}(t) = P_{2j-1,2k}(t,0)$ for j, k = 1, 2, and we also have $\hat{f}_j(t) = \hat{F}_{2j-1}(t)$ for j = 1, 2.

10

Using the eigenvalues and eigenvectors written in Eqs. (9), (10), (19), and (20), we arrive at the formulas specific to our model:

$$\boldsymbol{U}(t) = \begin{bmatrix} \mu(t) & 0\\ \frac{-i\epsilon s(t)}{\gamma} & \frac{1}{\mu(t)} \end{bmatrix}, \boldsymbol{V}(t) = \begin{bmatrix} 0 & 0\\ \frac{-i\kappa s(t)}{\gamma} & 0 \end{bmatrix}, (53)$$

$$\langle \hat{\boldsymbol{F}}(t) \hat{\boldsymbol{F}}^{\dagger \boldsymbol{T}}(t) \rangle = \begin{bmatrix} \boldsymbol{F_1}(t) & \boldsymbol{F_{12}}(t) \\ \boldsymbol{F_{12}^{*T}}(t) & \boldsymbol{F_2}(t) \end{bmatrix}, \qquad (54)$$

$$\mathbf{F_1}(t) = \begin{bmatrix} 1 - \mu^2(t) & 0\\ 0 & 0 \end{bmatrix}, \quad \mathbf{F_{12}}(t) = \frac{i\sigma(t)}{2\gamma} \begin{bmatrix} \epsilon & -\kappa\\ 0 & 0 \end{bmatrix},$$
$$\mathbf{F_2}(t) = \frac{s(t) - 2\gamma t}{2\gamma^2} \begin{bmatrix} \epsilon^2 & -\epsilon\kappa\\ -\epsilon\kappa & -\kappa^2 \end{bmatrix} + \left(\frac{1}{\mu^2(t)} - 1\right) \begin{bmatrix} 0 & 0\\ 0 & 1 \end{bmatrix},$$
(55)

where $\mu(t) = \exp(-\gamma t)$, $\sigma(t) = \exp(-2\gamma t) - 1 + 2\gamma t$, and $s(t) = \sinh(2\gamma t)$ using the hyperbolic sinus function.

To check consistency of the model with unidirectional coupling, we determine the mean values of equal time field-operators commutation relations. Compared to the usual canonical commutation relations $\langle [\hat{a}_j(t), \hat{a}_k(t)] \rangle = 0$, $\langle [\hat{a}_j(t), \hat{a}_k^{\dagger}(t)] \rangle = \delta_{jk}$, and $\langle [\hat{a}_j^{\dagger}(t), \hat{a}_k^{\dagger}(t)] \rangle = 0$ for j, k = 1, 2, the following two relations are found:

$$\langle [\hat{a}_2(t), \hat{a}_2^{\dagger}(t)] \rangle = 1 + \frac{(\epsilon^2 - \kappa^2)\phi(t)}{2\gamma^2},$$

$$\langle [\hat{a}_1(t), \hat{a}_2^{\dagger}(t)] \rangle = \frac{-i\epsilon\psi(t)}{2\gamma},$$
(56)

where $\psi(t) = \exp(-2\gamma t) - 1 - 2\gamma t$ and $\phi(t) = \exp(2\gamma t) - 1 - 2\gamma t$. For short times t assuming $t \ll 1/\gamma$ we have $\langle [\hat{a}_2(t), \hat{a}_2^{\dagger}(t)] \rangle = 1 + (\epsilon^2 - \kappa^2)t^2$ and $\langle [\hat{a}_1(t), \hat{a}_2^{\dagger}(t)] \rangle = -2i\epsilon t$. Thus, we additionally require $t \ll 1/\epsilon$ and $t \ll 1/\sqrt{\epsilon^2 - \kappa^2}$. As we usually assume in \mathcal{PT} -symmetric systems that $\kappa \leq \epsilon$, we are left with the following conditions for applicability of the model with unidirectional coupling:

$$t \ll \min\left\{\frac{1}{\gamma}, \frac{1}{\epsilon}\right\}.$$
 (57)

The form of the commutation relations given in Eq. (56) and the ensuing restricted validity of the model poses the question about possible corrections of the model using suitable properties of the reservoir Langevin operator forces. Similarly as it is done when damping and amplification are introduced into the Heisenberg equations (the Wigner–Weisskopf model of damping, see Ref. [11]). However, as discussed in detail in Appendix A, this approach is not successful.

Also, in Appendix B the properties of the modes are analyzed in the framework of the Gaussian states, their nonclassicality depths and logarithmic negativity are determined. These results point out at specific properties of the two-mode bosonic system with unidirectional coupling applicable only under the conditions given in Eq. (57). These results lead us to the conclusion that the method for concatenating simpler bosonic systems with QEPs to arrive at QEPs with higher-order ED has limitations. The question how to obtain higher-order inherited QEPs in bosonic systems without these limitations is open.

V. NUMERICAL IDENTIFICATION OF EXCEPTIONAL POINTS AND THEIR DEGENERACIES

When an analyzed system is more complex and we do not succeed in revealing its eigenvalues and eigenvectors analytically, we can apply two different numerical approaches to arrive at the orders of exceptional degeneracies. We demonstrate both approaches analytically using the simplest two-mode systems with the usual and unidirectional couplings.

A. Jordan canonical form of a general matrix

The Jordan form J_M of a matrix M contains nonzero elements on the diagonal and also the nearest upper diagonal. It has the dimension of the matrix M. Some elements at the nearest upper diagonal equal one if the matrix M is non-diagonalizable. The neighbor elements equal to one form groups. The number of elements in a given group gives the order of ED (equal to the number of elements + 1) related to the corresponding eigenvalue.

For example and considering the dynamical matrix $M^{(2)}$, given in Eq. (27), belonging to the two-mode bosonic system with usual coupling under the condition $\beta = 0$, i.e., where a second-order QEP occurs, the Jordan form $J_{M^{(2)}}$ and the corresponding similarity transformation $S_{M^{(2)}}$ such that

$$M^{(2)} = S_{M^{(2)}} J_{M^{(2)}} S_{M^{(2)}}^{-1}$$
(58)

take the form:

$$\boldsymbol{J}_{\boldsymbol{M}^{(2)}} = \begin{bmatrix} -i\gamma_{+} & 1\\ 0 & -i\gamma_{+} \end{bmatrix}, \quad \boldsymbol{S}_{\boldsymbol{M}^{(2)}} = \begin{bmatrix} i & -1/\gamma_{-}\\ 1 & 0 \end{bmatrix}.$$
(59)

In Eq. (59), the element (1,2) of the matrix $J_{M^{(2)}}$, equal to 1, identifies a second-order QEP.

Similarly, when analyzing the two-mode bosonic system with unidirectional coupling and the dynamical matrix $M_{\rm u}^{(1+1)}$, written in Eq. (18), under the condition $\gamma_1 = \gamma_2$ guaranteeing the existence of a second-order QEP, we arrive at:

$$\boldsymbol{J}_{\boldsymbol{M}^{(1+1)}} = \begin{bmatrix} -i\gamma_2/2 & 1\\ 0 & -i\gamma_2/2 \end{bmatrix}, \quad \boldsymbol{S}_{\boldsymbol{M}^{(1+1)}} = \begin{bmatrix} 0 & 1/\xi\\ 1 & 0 \end{bmatrix}.$$
(60)

B. Perturbation of a dynamical matrix

The second approach is based upon introducing a suitable small perturbation of a dynamical matrix M that can remove both EDs and DDs. In this approach, the eigenvalues of the perturbed dynamical matrix M_{δ} are determined and the degeneracies are removed as the degenerated eigenvalues split and then gradually diverge with the increasing perturbation. Perturbation can also be used for characterizing ED [32–34].

Identifying EDs, we demonstrate different kinds of the influence of perturbation δ on the eigenvalues of the matrix M considering the two-mode systems with the usual and unidirectional couplings and different positions of the perturbation δ inside the matrix M. Specifically, we demonstrate splitting of the eigenvalues in their real and/or imaginary parts and splitting proportional to $\sqrt{\delta}$ and δ when a QEP with second-order ED is disturbed. We quantify the strength of the perturbation by the overlap F of the normalized eigenvectors y_1 and y_2 that become gradually distinguishable as the perturbation δ increases

$$F = \frac{\langle \boldsymbol{y}_2 | \boldsymbol{y}_1 \rangle}{\sqrt{\langle \boldsymbol{y}_1 | \boldsymbol{y}_1 \rangle \langle \boldsymbol{y}_2 | \boldsymbol{y}_2 \rangle}} \tag{61}$$

and symbol $\langle | \rangle$ stands for the scalar product of complex vectors.

1. A two-mode system with usual coupling and $\beta = 0$ described by the perturbed dynamical matrix

$$\boldsymbol{M}_{\boldsymbol{\delta},\mathbf{1}}^{(2)} = \boldsymbol{M}^{(2)} + \begin{bmatrix} \delta & 0\\ 0 & 0 \end{bmatrix}.$$
 (62)

The eigenvalues and the corresponding eigenvectors take, respectively, the form:

$$\lambda_{1,2}^{M_{\delta,1}^{(2)}} = -i\gamma_{+} + \frac{\delta}{2} \mp \sqrt{\delta\left(\frac{\delta}{4} - i\gamma_{-}\right)} \tag{63}$$

and

$$\boldsymbol{y}_{1,2}^{\boldsymbol{M}_{\boldsymbol{\delta},1}^{(2)}} = \left[i - \frac{\delta \mp \sqrt{\delta \left(\delta - 4i\gamma_{-}\right)}}{2\gamma_{-}}, 1\right]^{T}.$$
 (64)

2. A two-mode system with usual coupling and $\beta = 0$ described by the perturbed dynamical matrix

$$\boldsymbol{M}_{\boldsymbol{\delta},\boldsymbol{2}}^{(2)} = \boldsymbol{M}^{(2)} + \begin{bmatrix} 0 & -\delta \\ 0 & 0 \end{bmatrix}.$$
 (65)

The eigenvalues and the corresponding eigenvectors are derived, respectively, as follows:

$$\lambda_{1,2}^{M_{\delta,2}^{(2)}} = -i\gamma_+ \mp \sqrt{\delta\gamma_-} \tag{66}$$

and

$$\boldsymbol{y}_{1,2}^{\boldsymbol{M}_{\delta,2}^{(2)}} = \left[i \pm \sqrt{\frac{\delta}{\gamma_{-}}}, 1\right]^{T}.$$
 (67)

3. A two-mode system with unidirectional coupling and $\gamma_1 = \gamma_2$ described by the perturbed dynamical matrix

$$M_{\mathbf{u},\delta}^{(1+1)} = M_{\mathbf{u}}^{(1+1)} + \begin{bmatrix} \delta & 0\\ 0 & 0 \end{bmatrix}.$$
 (68)

The eigenvalues and the corresponding eigenvectors are attained, respectively, in the form:

$$\lambda_{1}^{M_{u,\delta}^{(1+1)}} = -i\gamma_{2}/2, \lambda_{2}^{M_{u,\delta}^{(1+1)}} = -i\gamma_{2}/2 + \delta$$
(69)

and

$$y_{1}^{M_{u,\delta}^{(1+1)}} = [0,1]^{T}, y_{2}^{M_{u,\delta}^{(1+1)}} = \left[\frac{\delta}{\xi},1\right]^{T}.$$
(70)

The real and imaginary parts of the eigenvalues λ_j , j = 1, 2, from Eqs. (63), (66), and (69) are plotted in Figs. 4(a,b) as they depend on the perturbation δ . The perturbation δ in general disturbs more strongly the system with the usual coupling, as it is apparent both from the graphs of the eigenvalues and the overlap F of eigenvectors shown in Fig. 4(c).

In the above-discussed cases, the second-order DD present in both two-mode systems was not modified by the perturbation δ because of the structure of these systems. However, suitable positioning of the perturbation δ inside a dynamical matrix M may also result in revealing the DDs. As the DD is embedded in the 2×2 matrix $\boldsymbol{\xi}$ given in Eq. (4), the perturbation δ has to affect this matrix. We consider two kinds of perturbation in the twomode system with the dynamical matrix $M^{(2)}$: The first one splits all eigenvalues Λ_j , $j = 1, \ldots, 4$, in their real parts, whereas the second one distinguishes two eigenvalues in their real parts and the remaining two eigenvalues in their imaginary parts. We note that the perturbation δ primarily modifies the eigenvalues of the matrix $\boldsymbol{\xi}$, which removes the DD. Secondarily, as the eigenvalues of $\boldsymbol{\xi}$ determined for nonzero δ differ from those valid for $\delta = 0$. the conditions for having a QEP of the matrix M change and the original setting of the system parameters for a QEP is lost and so also the corresponding ED is lost.

1. A two-mode system with usual coupling and $\beta = 0$ described by the dynamical matrix $M^{(2)}$ where

$$\boldsymbol{\xi}_{\delta,1} = \boldsymbol{\xi} + \begin{bmatrix} \delta & 0\\ 0 & 0 \end{bmatrix}. \tag{71}$$

The eigenvalues and the corresponding eigenvectors of matrix $\boldsymbol{\xi}_{\delta,1}$ are written, respectively, as:

$$\lambda_{1,2}^{\xi_{\delta,1}} = \frac{\delta}{2} \mp \sqrt{\zeta^2 + \delta\left(\epsilon + \frac{\delta}{4}\right)} \tag{72}$$



FIG. 4. (a) Real $\lambda^{\rm r}$ and (b) imaginary $\lambda^{\rm i}$ parts of eigenvalues $\lambda_{1,2}$ of the dynamical matrices $M_{\delta,1}^{(2)}$ ($\gamma_+ = 0, \gamma_- = 0.5$, blue solid curves), $M_{\delta,2}^{(2)}$ ($\gamma_+ = 0, \gamma_- = 0.5$, red dashed curves), and $M_{\mathbf{u},\delta}^{(1+1)}$ ($\gamma_2 = 0$, green dot-dashed curves) given in turn in Eqs. (62), (65), and (68) as they depend on perturbation parameter δ . In (c) the overlap F of the corresponding eigenvectors is plotted. In (a) and (b), the numbers denote the curves of the corresponding eigenvalues.

and

$$\boldsymbol{y}_{1,2}^{\boldsymbol{\xi}\boldsymbol{\delta},1} = \left[-\frac{\epsilon + \lambda_{1,2}^{\boldsymbol{\xi},\boldsymbol{\delta},1}}{\kappa}, 1\right]^{T}.$$
 (73)

2. A two-mode system with usual coupling and $\beta = 0$ described by the dynamical matrix $M^{(2)}$ where

$$\boldsymbol{\xi}_{\delta,2} = \boldsymbol{\xi} + \begin{bmatrix} \delta & \delta \\ 0 & 0 \end{bmatrix}. \tag{74}$$

The eigenvalues and the corresponding eigenvectors of matrix $\boldsymbol{\xi}_{\delta,2}$ are obtained, respectively, as:

$$\lambda_{1,2}^{\xi_{\delta,2}} = \frac{\delta}{2} \mp \sqrt{\zeta^2 + \delta\left(\epsilon - \kappa + \frac{\delta}{4}\right)} \tag{75}$$



FIG. 5. (a) Real $\Lambda^{\rm r}$ and (b) imaginary $\Lambda^{\rm i}$ parts of the eigenvalues $\Lambda_{1,\ldots,4}$ of the dynamical matrix $M^{(2)}$ involving $\boldsymbol{\xi}_{\delta,1}$ (blue solid curves) and $\boldsymbol{\xi}_{\delta,2}$ (red dashed curves) given in Eqs. (71) and (74), respectively, as they depend on perturbation parameter δ ; $\gamma_+ = 0$, $\beta = 0$, $\kappa/\epsilon = 1/2$. In (a) and (b), the numbers denote the curves of the corresponding eigenvalues.

and

$$\boldsymbol{y}_{1,2}^{\boldsymbol{\xi}\boldsymbol{\delta},\boldsymbol{2}} = \left[-\frac{\boldsymbol{\epsilon} + \lambda_{1,2}^{\boldsymbol{\xi}+\boldsymbol{\delta}}}{\kappa}, 1\right]^{T}.$$
 (76)

The looked for eigenvalues and eigenvectors are then reached using the eigenvalues and eigenvectors of the 2×2 matrix $M^{(2)}$ written in Eq. (27):

 $\lambda_{1,2}^{M^{(2)}} = -i\gamma_+ \mp \beta$

and

(77)

 $y_{1,2}^{M^{(2)}} = \left[-\frac{i\gamma_{-}\pm\beta}{\xi}, 1\right]^{-},$ (78)where $4\gamma_{\pm} = \gamma_1 \pm \gamma_2$ and $\beta^2 = \xi^2 - \gamma_-^2$. The real and imaginary parts of the eigenvalues Λ_j , $j = 1, \ldots, 4$, of

the 4×4 matrix $M^{(2)}$ are plotted in Fig. ??.

Finally, we mention a specific case of the perturbation δ that removes DD but keeps one from the two originally diabolically-degenerated QEPs present in the system. The dynamical matrix $M^{(2)}$ of the two-mode bosonic system with usual coupling is perturbed in the following way:

$$\tilde{\boldsymbol{\gamma}}_{\mathbf{1}}^{\delta,3} = \tilde{\boldsymbol{\gamma}}_{\mathbf{1}} + \begin{bmatrix} \delta & 0\\ 0 & 0 \end{bmatrix}.$$
(79)

The eigenvalues $\Lambda^{M_{\delta,3}^{(2)}}$ of the dynamical matrix in Eq. (79) are derived in the form:

$$\Lambda_{1,2}^{M^{(2)}_{\delta,3}} = -i\gamma_+ \mp \beta$$

$$\Lambda_{3,4}^{M_{\delta,3}^{(2)}} = -i\gamma_{+} + \frac{\delta}{2} \mp \sqrt{\beta^2 + \delta\left(\frac{\delta}{4} - i\gamma_{-}\right)}.$$
 (80)

The corresponding eigenvectors are expressed as follows:

$$\mathbf{Y}_{\mathbf{1,2}}^{\boldsymbol{M}_{\boldsymbol{\delta},\mathbf{3}}^{(2)}} = \left[0, \frac{i\gamma_{-}\pm\beta}{\epsilon}, -\frac{\kappa}{\epsilon}, 1 \right]^{T}, \\
 \mathbf{Y}_{\mathbf{3,4}}^{\boldsymbol{M}_{\boldsymbol{\delta},\mathbf{3}}^{(2)}} = \left[\frac{i\gamma_{-}-\Lambda_{\mathbf{3,4}}^{M_{\boldsymbol{\delta},\mathbf{3}}^{(2)}}}{\kappa}, 0, -\frac{\epsilon}{\kappa}, 1 \right]^{T}. \quad (81)$$

According to Eqs. (80) and (81), we observe a single QEP with second-order ED for $\beta = 0$ and $\delta \neq 0$. This contrasts with the observation of a QEP with second-order ED and also second-order DD for $\beta = 0$ and $\delta = 0$.

VI. HIGHER-ORDER HYBRID POINTS REVEALED BY FIELD-OPERATOR MOMENTS

In the last section, we derive general formulas that give us the orders of EDs and DDs of QHPs that occur in the dynamics of higher-order FOMs. We note that the structure of higher-order FOM spaces mapped onto suitable lattices and its influence to system's properties has been analyzed in detail in Ref. [35] for two- and threemode bosonic systems.

Let us consider a bosonic system composed on n modes and described by an appropriate quadratic non-Hermitian Hamiltonian. Such a system is described by the 2n annihilation and creation operators, and the dynamical matrix $M^{(n)}$ of its Heisenberg-Langevin equations has 2n eigenvalues. Let us fix a position in the system parameter space. After diagonalization of the dynamical matrix $M^{(n)}$, we identify the eigenvalues λ_j with different eigenvectors and count their degeneration numbers n_j according to the number of coalescing eigenvectors (from the original maximal Hilbert space). Denoting the number of such eigenvalues by Σ_{λ} , we have

$$\sum_{j=1}^{\Sigma_{\lambda}} n_j = 2n. \tag{82}$$

We assign, to any eigenvalue λ_j , an operator vector \hat{B}_j that encompasses the diagonalized field operators \hat{b}_l and \hat{b}_l^{\dagger} (l = 1, ..., n) that are associated with the coalescing vectors of this eigenvalue.

Now we consider the dynamics of kth-order FOMs. These kth-order FOMs can formally be expressed in their general form using the above-defined operator vectors $\hat{\boldsymbol{B}}_{\boldsymbol{j}}$ as $\langle \prod_{j=1}^{\Sigma_{\lambda}} \hat{\boldsymbol{B}}_{\boldsymbol{j}}^{k_j} \rangle$, where the nonnegative integers k_j obey $\sum_{j=1}^{\Sigma_{\lambda}} k_j = k$. The vector \boldsymbol{k} defined as $(k_1, k_2, \ldots, k_{\Sigma_{\lambda}})$ then points out at a possible QHP with the complex eigenfrequency λ given as

$$\lambda = \sum_{j=1}^{\Sigma_{\lambda}} \lambda_j^{k_j}.$$
(83)

The degrees d_{e}^{k} of ED and d_{d}^{k} of DD of this QHP are given by the following formulas:

$$d_{\rm e}^{\boldsymbol{k}} = \prod_{j=1}^{\Sigma_{\lambda}} n_j^{k_j}, \qquad (84)$$

$$d_{\rm d}^{\boldsymbol{k}} = \frac{k!}{\prod_{j=1}^{\Sigma_{\lambda}} k_j!}.$$
(85)

The number $N_B^{(k)}$ of such QHPs observed for the kthorder FOMs (see the fourth columns of Tabs. I—VI in Ref. [8]) is determined using the combination number as:

$$N_B^{(k)} = \begin{pmatrix} \Sigma_\lambda + k - 1 \\ k \end{pmatrix}.$$
 (86)

Similar formula also gives the number $N_b^{(k)}$ of different kth-order FOMs written in the operators \hat{b}_l and \hat{b}_l^{\dagger} for $l = 1 \dots n$:

$$N_b^{(k)} = \begin{pmatrix} 2n+k-1\\k \end{pmatrix}.$$
 (87)

These FOMs are explicitly written in the third columns of Tabs. I—VI in Ref. [8]. Also, the number $N_b^{(k)}$ of different kth-order FOMs written in the operators \hat{b}_l and \hat{b}_l^{\dagger} and belonging to a fixed **k** is expressed as:

$$N_{b}^{(k)} = \prod_{l=1}^{k} \left(\begin{array}{c} n_{l} + k_{l} - 1 \\ k_{l} \end{array} \right).$$
(88)

To demonstrate the general approach and formulas, we apply them to the four-mode bosonic system in the circular configuration analyzed in Sec. II in the regime of nonconventional PT-symmetric dynamics. Its first- and second-order FOMs and the revealed QHPs with their degeneracies are given in Tab. I. The general approach results in Tab. IV that allows to easily derive the content of Tab. I: Two lines belonging to $\Lambda_{j}^{i} = \gamma_{+}$ are obtained assuming $\mathbf{k} = (0, 0, 0, 0, 1, 0)$ and (0, 0, 0, 0, 0, 1). For $\Lambda_{j}^{i} = 2\gamma_{+}$ the entries to Tab. I are in turn generated assuming $\mathbf{k} = (0, 0, 0, 0, 1, 1), (0, 0, 0, 0, 2, 0),$ and (0, 0, 0, 0, 2). The orders of the corresponding EDs and DDs are derived using Eqs. (84) and (85), respectively.

These results can be used for detailed discussions of genuine and induced QHPs and their degeneracies for FOMs of arbitrary orders and considering different systems with their specific inherited QHPs observed in the dynamics of field operators governed by the Heisenberg-Langevin equations. The occurrence of a QHP with n^k th ED and second-order DD in the dynamics of kth-order FOMs in a bosonic system with an inherited QHP with *n*th-order ED and second-order DD is probably the most valuable result.

VII. CONCLUSIONS

We have continued the analysis of the dynamics of simple bosonic systems described by quadratic non-

λ_j	n_j	\hat{B}_j	k_{j}
γ_{13}	1	$\hat{B}_1\equiv [\hat{b}_1]$	$k_1 = 0$
γ_{13}	1	$oldsymbol{\hat{B}_2}\equiv [\hat{b}_1^\dagger]$	$k_2 = 0$
γ_{24}	1	$\hat{m{B}}_{m{3}}\equiv [\hat{b}_2]$	$k_3 = 0$
γ_{24}	1	$\hat{m{B}}_{4}\equiv [\hat{b}_{2}^{\dagger}]$	$k_4 = 0$
γ_+	2	$\hat{m{B}}_{m{5}}\equiv[\hat{b}_3,\hat{b}_3^\dagger]$	k_5
γ_+	2	$\hat{m{B}}_{m{6}}\equiv[\hat{b}_4,\hat{b}_4^\dagger]$	k_6
$\sum_{j=1}^{\Sigma_{\lambda}} \lambda_j = \lambda$	$\sum_{j=1}^{\Sigma_{\lambda}} n_j = 2n$	$\langle \prod_{j=1}^{\Sigma_{\lambda}} \hat{B}_{j}^{k_{j}} angle$	$\sum_{j=1}^{\Sigma_{\lambda}} k_j = k$

TABLE IV. Eigenvalues λ_j , their degeneracies n_j , the corresponding field-operator vectors \hat{B}_j , and their varying powers k_j for the four-mode bosonic system analyzed in Tab. I. QHPs are uniquely identified by powers of k_5 and k_6 .

Hermitian Hamiltonians from the point of view of the occurrence of quantum exceptional, diabolical, and hybrid points. We have revealed the bosonic systems exhibiting nonconventional \mathcal{PT} -symmetric dynamics characterized by the observation of quantum exceptional and hybrid points only in certain subspace(s) of the whole system Liouville space. We have identified such nonconventional second-order inherited quantum exceptional and hybrid points in four-mode bosonic systems.

Applying the method of concatenating simple bosonic systems via unidirectional coupling, we have found the conditions for the observation of up to sixth-order inherited quantum exceptional and hybrid points. However, by analyzing the behavior of the two-mode bosonic system with unidirectional coupling we have shown that the bosonic systems with unidirectional coupling are applicable only for short times in which they ensure physically consistent behavior. In short times, more complex bosonic systems with diverse structures and arbitraryorder exceptional degeneracies can be built by concatenating simple bosonic systems via unidirectional coupling of several types. Nevertheless, tailoring the properties of the Langevin operator stochastic forces in systems with unidirectional coupling does not enable extending their applicability to arbitrary times, in contrast to how the damping and amplification are consistently described.

Using analytical formulas for two-mode bosonic systems we have elucidated the operation of two numerical methods for the identification of quantum exceptional points and their degeneracies. These include: (1) the transformation of a dynamical matrix into its Jordan form and (2) the introduction of a suitable perturbation δ into the dynamical matrix and its subsequent eigenvalue analysis.

The exceptional and diabolical degeneracies of inherited quantum hybrid points have been used to derive higher-order degeneracies observed in the dynamics of higher-order field-operator moments. The quantum exceptional and hybrid points of second-order field-operator moments have been summarized in tables that evidence a rich dynamics of the field-operator moments.

Numbers of genuine and induced quantum hybrid points and their exceptional and diabolical degeneracies have been expressed as they depend on the order of the field-operator moments in the general form using parameters of the inherited quantum exceptional and hybrid points and their degeneracies.

This analysis considerably broadens the investigations of bosonic systems with \mathcal{PT} -symmetry exhibiting correct physical behavior, though it does not reveal bosonic systems with higher-order exceptional and hybrid singularities found for arbitrarily long times. The bosonic systems with unidirectional coupling enable generating higher-order exceptional and hybrid points only in their short-time dynamics. The systems with nonconventional \mathcal{PT} -symmetric dynamics exhibit only low- (second-) order exceptional and hybrid degeneracies, similarly as their counterparts analyzed in Ref. [8]. This analysis together with that in Ref. [8] show that the observation of higher-order exceptional and hybrid singularities in real, *i.e. physically well-behaved, bosonic systems at arbitrary times is challenging.*

VIII. ACKNOWLEDGEMENTS

The authors thank Ievgen I. Arkhipov for useful discussions. J.P. and K.T. acknowledge support by the project OP JAC CZ.02.01.01/00/22_008/0004596 of the Ministry of Education, Youth, and Sports of the Czech Republic. A.K.-K., G.Ch., and A.M. were supported by the Polish National Science Centre (NCN) under the Maestro Grant No. DEC-2019/34/A/ST2/00081.

Appendix A: Two-mode system with unidirectional coupling and relevant reservoir properties

Considering the approach outlined in Ref. [36], we construct the matrix $\langle \hat{L}^{\mathbf{u}}(t) \hat{L}^{\mathbf{u}\dagger}(t) \rangle$ of the stochastic Langevin operator forces such that the bosonic commutation relations of the field operators are obeyed for an arbitrary time t.

First, we note that the additional, unwanted, terms in the commutation relations in Eq. (56) disappear when we replace the correlation matrix $\langle \hat{F}(t)\hat{F}^{\dagger}(t)\rangle$ of the fluctuating operator forces \hat{F} given in Eq. (54) by the matrix $\langle \hat{F}^{u}(t)\hat{F}^{u\dagger}(t)\rangle$ with the constituting submatrices:

$$\begin{aligned} \mathbf{F_1^u}(t) &= \mathbf{F_1}(t), \\ \mathbf{F_2^u}(t) &= \mathbf{F_2}(t) + \frac{\phi(t)}{2\gamma^2} \begin{bmatrix} -\epsilon^2 & 0\\ 0 & \kappa^2 \end{bmatrix}, \\ \mathbf{F_{12}^u}(t) &= \mathbf{F_{12}}(t) + \frac{i\psi(t)}{2\gamma} \begin{bmatrix} \epsilon & 0\\ 0 & 0 \end{bmatrix}, \end{aligned}$$
(A1)

where the functions ϕ and ψ are defined below Eq. (56).

Inverting Eq. (51) (for details, see Eq. (18) in Ref. [36]) we obtain the correlation matrix $\langle \hat{\boldsymbol{L}^{u}}(t) \hat{\boldsymbol{L}^{u\dagger}}(t') \rangle$ of the Langevin operator forces as follows:

$$\langle \hat{L^{\mathbf{u}}}(t)\hat{L^{\mathbf{u}\dagger}}(t')\rangle = \delta(t-t')$$

$$\times \begin{bmatrix} 2\gamma & 0 & -i\epsilon l_1(t) & 0\\ 0 & 0 & 0 & 0\\ i\epsilon l_1(t) & 0 & -\frac{\epsilon^2 \mu(t)^2 l_2(t)}{\gamma} & \frac{\epsilon \kappa l_1(t) l_2(t)}{2\gamma}\\ 0 & 0 & \frac{\epsilon \kappa l_1(t) l_2(t)}{2\gamma} & 2\gamma - \frac{\kappa^2 l_2(t)}{\gamma} \end{bmatrix},$$
(A2)

where $l_{1,2}(t) = \exp(-2\gamma t) \pm 1$; $\mu(t) = \exp(-\gamma t)$.

However, the correlation matrix $\langle \hat{L}^{\mathbf{u}}(t) L^{\mathbf{u}\dagger}(t') \rangle$ does not represent a physical reservoir, as it has a negative eigenvalue. This can be even analytically confirmed considering t = 0. In this case, the eigenvalues ν are obtained as:

$$\nu_{1,2,3,4} = 0, \ 2\gamma, \ \gamma \pm \sqrt{\gamma^2 + 4\epsilon^2}.$$
 (A3)

This means in general that there does not exist a physical reservoir with properties such that the model with unidirectional coupling could be applied for an arbitrary time t.

Appendix B: Statistical properties of a two-mode bosonic system with unidirectional coupling

We consider modes 1 and 2 being in their initial coherent states α_1 and α_2 , respectively. The modes evolution described in Eq. (52) maintains the state Gaussian form described by the following normal characteristic function [7, 11, 37]:

$$C_{\mathcal{N}}(\beta_1, \beta_2; t) = \exp\left[-B_2(t)|\beta_2|^2 + \{C_2(t)\beta_2^{*2} + \text{c.c.}\}/2\right] \\ \times \exp[\alpha_1^*(t)\beta_1 + \alpha_2^*(t)\beta_2 - D(t)\beta_1^*\beta_2^* + \text{c.c.}],$$
(B1)

where symbol c.c. replaces the complex conjugated terms. The functions $B_2(t)$, $C_2(t)$, and D(t) are given as:

$$B_{2}(t) = \langle \delta \hat{a}_{2}^{\dagger}(t) \delta \hat{a}_{2}(t) \rangle = \frac{\kappa^{2} \psi(t)}{2\gamma^{2}} - \frac{l_{1}(t)}{\mu^{2}(t)},$$

$$C_{2}(t) = \langle [\delta \hat{a}_{2}(t)]^{2} \rangle = -\frac{\epsilon \kappa \psi(t)}{2\gamma^{2}},$$

$$D(t) = \langle \delta \hat{a}_{1}(t) \delta \hat{a}_{2}(t) \rangle = -i\kappa t;$$
(B2)

and $\delta \hat{x} = \hat{x} - \langle \hat{x} \rangle$ for an arbitrary operator \hat{x} . The functions ϕ , ψ , μ , and l_1 are defined below Eqs. (56) and (A2), while the complex modes amplitudes $\alpha_1(t)$ and $\alpha_2(t)$ are derived from their initial values along the relations

$$\begin{bmatrix} \alpha_1(t) \\ \alpha_2(t) \end{bmatrix} = \boldsymbol{U}(t) \begin{bmatrix} \alpha_1(0) \\ \alpha_2(0) \end{bmatrix} + \boldsymbol{V}(t) \begin{bmatrix} \alpha_1^*(0) \\ \alpha_2^*(0) \end{bmatrix}, (B3)$$

where the matrices U(t) and V(t) are given in Eq. (53).

Using the formulas in Eq. (B2), the nonclassicality depth τ_2 of mode 2 [38], determined as

$$\tau_2(t) = \max\{0, |C_2(t)| - B_2(t)\}, \qquad (B4)$$

is vanishing, which means the classical behavior of the marginal field in mode 2. Also the marginal field in mode 1 stays classical, as we have $\tau_1 = 0$.

We quantify the entanglement between the fields in modes 1 and 2 by the logarithmic negativity E_N [39] determined from the simplectic eigenvalues ν_{\pm} of the partially-transposed symmetrically-ordered covariance matrix $\boldsymbol{\sigma}^{\rm PT}$ [40], as given by:

$$\boldsymbol{\sigma}^{\mathrm{PT}}(t) = \begin{bmatrix} \mathbf{1} & \boldsymbol{\sigma}_{12}^{\mathrm{PT}}(t) \\ \begin{bmatrix} \boldsymbol{\sigma}_{12}^{\mathrm{PT}}(t) \end{bmatrix}^{\mathrm{T}} & \boldsymbol{\sigma}_{2}^{\mathrm{PT}}(t) \end{bmatrix},$$
(B5)

$$\sigma_2^{\text{PT}}(t) = \begin{bmatrix} 1 + 2B_2(t) + 2C_2(t) & 0\\ 0 & 1 + 2B_2(t) - 2C_2(t) \end{bmatrix}$$

$$\boldsymbol{\sigma}_{12}^{\mathrm{PT}}(t) = 2 \begin{bmatrix} 0 & -\mathrm{Im}\{D(t)\} \\ \mathrm{Im}\{D(t)\} & 0 \end{bmatrix},$$
(B6)

where **1** stands for the two-dimensional identity matrix. Determining two invariants $\delta = 1 + \text{Det}\{\sigma_2^{\text{PT}}\} + 2\text{Det}\{\sigma_{12}^{\text{PT}}\}$ and $\Delta = \text{Det}\{\sigma^{\text{PT}}\}$ of the matrix σ^{PT} , the simplectic eigenvalues ν_{\pm} are expressed as [40]:

$$2\nu_{\pm}^2 = \delta \pm \sqrt{\delta^2 - 4\Delta}.$$
 (B7)

The logarithmic negativity E_N is then determined along the formula

$$E_N = \max\{0, -\ln(\nu_{-})\}.$$
 (B8)

As the model is applicable only for short times t, fulfilling the conditions in Eq. (57), we derive the logarithmic negativity E_N using the Taylor expansion in t to first order:

$$E_N(t) = -\ln\left[1 + 2\gamma t \left(1 - \sqrt{1 + (\kappa/\gamma)^2}\right)\right].$$
 (B9)

The formula (B9) predicts nonzero negativity E_N for short times t despite the fact that mode 1 remains in a coherent state. If $\alpha_1(0) = (0,0)$ then $\alpha_1(t) = (0,0)$ and mode 1 remains in the vacuum state. However, this state cannot be entangled with mode 2, as the formula (B9) for negativity E_N predicts. This is another manifestation of the limited applicability of the studied model with unidirectional propagation.

- W. Chen, S. K. Özdemir, G. Zhao, J. Wiersig, and L. Yang, Exceptional points enhance sensing in an optical microcavity, Nature (London) 548, 192 (2017).
- [2] Z.-P. Liu, J. Zhang, S. K. Özdemir, B. Peng,

H. Jing, X.-Y. Lü, C.-W. Li, L. Yang, F. Nori, and Y.-X. Liu, Metrology with \mathcal{PT} -symmetric cavities: Enhanced sensitivity near the \mathcal{PT} -phase transition, Phys. Rev. Lett. **117**, 110802 (2016).

- [3] L. Feng, R. El-Ganainy, and L. Ge, Non-Hermitian physics and *PT* symmetry, Nat. Photon. **11**, 752 (2017).
- [4] R. El-Ganainy, M. Khajavikhan, S. Rotter, D. N. Christodoulides, and Ş. K. Özdemir, Non-Hermitian physics and *PT* symmetry, Commun. Phys. 2, 1 (2019).
- [5] M. Parto, Y. G. N. Liu, B. Bahari, M. Khajavikhan, and D. N. Christodoulides, Non-Hermitian and topological photonics: Optics at an exceptional point, Nanophotonics 10, 403 (2021).
- [6] J. Peřina Jr. and A. Lukš, Quantum behavior of a *PT*symmetric two-mode system with cross-Kerr nonlinearity, Symmetry **11**, 1020 (2019).
- [7] J. Peřina Jr., A. Lukš, J. K. Kalaga, W. Leoński, and A. Miranowicz, Nonclassical light at exceptional points of a quantum *PT*-symmetric two-mode system, Phys. Rev. A **100**, 053820 (2019).
- [8] K. Thapliyal, J. Peřina Jr., G. Chimczak, A. Kowalewska-Kudłaszyk, and A. Miranowicz, Multiple quantum exceptional, diabolical, and hybrid points in multimode bosonic systems: I. Inherited and genuine singularities, arxiv (2024).
- [9] J. Peřina Jr., On the equivalence of some projection operator techniques, Physica A 214, 309 (1995).
- [10] W. Vogel and D. G. Welsch, Quantum Optics, 3rd ed. (Wiley-VCH, Weinheim, 2006).
- [11] J. Peřina, Quantum Statistics of Linear and Nonlinear Optical Phenomena (Kluwer, Dordrecht, 1991).
- [12] Q. Zhong, J. Kou, S. Özdemir, and R. El-Ganainy, Hierarchical construction of higher-order exceptional points, Phys. Rev. Lett. **125**, 203602 (2020).
- [13] J. Wiersig, Revisiting the hierarchical construction of higher-order exceptional points, Phys. Rev. A 106, 063526 (2022).
- [14] S. Wang, B. Hou, W. Lu, Y. Chen, Z. Zhang, and C. T. Chan, Arbitrary order exceptional point induced by photonic spin–orbit interaction in coupled resonators, Nature Commun. 10, 832 (2019).
- [15] H. Jing, S. K. Özdemir, H. Lu, and F. Nori, High-order exceptional points in optomechanics, Sci. Rep. 7, 3386 (2017).
- [16] G.-Q. Zhang and J. You, Higher-order exceptional point in a cavity magnonics system, Phys. Rev. B 99, 054404 (2019).
- [17] E. M. Graefe, U. Günther, H. J. Korsch, and A. E. Niederle, A non-Hermitian symmetric Bose–Hubbard model: Eigenvalue rings from unfolding higher-order exceptional points, J. Phys. A: Math. Theor. 41, 255206 (2008).
- [18] M. Teimourpour, R. El-Ganainy, A. Eisfeld, A. Szameit, and D. N. Christodoulides, Light transport in ptinvariant photonic structures with hidden symmetries, Phys. Rev. A **90**, 053817 (2014).
- [19] M. Znojil, Complex symmetric Hamiltonians and exceptional points of order four and five, Phys. Rev. A 98, 032109 (2018).
- [20] I. Mandal and E. J. Bergholtz, Symmetry and higherorder exceptional points, Phys. Rev. Lett. 127, 186601 (2021).
- [21] P. Delplace, T. Yoshida, and Y. Hatsugai, Symmetryprotected multifold exceptional points and their topo-

logical characterization, Phys. Rev. Lett. **127**, 186602 (2021).

- [22] Q. Zhong, D. N. Christodoulides, M. Khajavikhan, K. Makris, and R. El-Ganainy, Power-law scaling of extreme dynamics near higher-order exceptional points, Phys. Rev. A 97, 020105 (2018).
- [23] H. Hodaei, A. U. Hassan, S. Wittek, H. Garcia-Gracia, R. El-Ganainy, D. N. Christodoulides, and M. Khajavikhan, Enhanced sensitivity at higher-order exceptional points, Nature (London) 548, 187 (2017).
- [24] Z.-Z. Li, W. Chen, M. Abbasi, K. W. Murch, and K. B. Whaley, Speeding up entanglement generation by proximity to higher-order exceptional points, Phys. Rev. Lett. 131, 100202 (2023).
- [25] J. Peřina Jr., A. Miranowicz, G. Chimczak, and A. Kowalewska-Kudłaszyk, Quantum Liouvillian exceptional and diabolical points for bosonic fields with quadratic Hamiltonians: The Heisenberg-Langevin equation approach, Quantum 6, 883 (2022).
- [26] R. W. Boyd, Nonlinear Optics, 2nd edition (Academic Press, New York, 2003).
- [27] G. Chimczak, A. Kowalewska-Kudłaszyk, E. Lange, K. Bartkiewicz, and J. Peřina Jr., The effect of thermal photons on exceptional points in coupled resonators, Sci. Rep. 13, 5859 (2023).
- [28] A. McDonald and A. A. Clerk, Exponentially-enhanced quantum sensing with non-hermitian lattice dynamics, Nature Communications 11, 5382 (2020).
- [29] P. Meystre and M. Sargent III, *Elements of Quantum Optics, 4nd edition* (Springer, Berlin, 2007).
- [30] G. S. Agarwal and K. Qu, Spontaneous generation of photons in transmission of quantum fields in *PT*-symmetric optical systems, Phys. Rev. A 85, 031802(R) (2012).
- [31] V. Peřinová, A. Lukš, and J. Křepelka, Quantum description of a *PT*-symmetric nonlinear directional coupler, J. Opt. Soc. Am. B **36**, 855 (2019).
- [32] M. Znojil, Unitarity corridors to exceptional points, Phys. Rev. A 100, 032124 (2019).
- [33] J. Wiersig, Response strengths of open systems at exceptional points, Phys. Rev. Res. 4, 023121 (2022).
- [34] J. Wiersig, Distance between exceptional points and diabolic points and its implication for the response strength of non-Hermitian systems, Phys. Rev. Res. 4, 033179 (2022).
- [35] I. I. Arkhipov, A. Miranowicz, F. Nori, S. K. Özdemir, and F. Minganti, Fully solvable finite simplex lattices with open boundaries in arbitrary dimensions, Phys. Rev. Res. 5, 043092 (2023).
- [36] J. Peřina Jr., A. Miranowicz, J. K. Kalaga, and W. Leoňski, Unavoidability of nonclassicality loss in *PT*symmetric systems, Phys. Rev. A **108**, 033512 (2023).
- [37] J. Peřina Jr. and J. Peřina, Quantum statistics of nonlinear optical couplers, in *Progress in Optics, Vol.* 41, edited by E. Wolf (Elsevier, Amsterdam, 2000) pp. 361—419.
- [38] C. T. Lee, Measure of the nonclassicality of nonclassical states, Phys. Rev. A 44, R2775 (1991).
- [39] S. Hill and W. K. Wootters, Computable entanglement, Phys. Rev. Lett. 78, 5022 (1997).
- [40] G. Adesso and F. Illuminati, Entanglement in continuous variable systems: Recent advances and current perspectives, J. Phys. A: Math. Theor. 40, 7821 (2007).