Thermodynamics of Non-Hermitian Josephson junctions with exceptional points

D. Michel Pino,¹ Yigal Meir,² and Ramón Aguado¹

¹Instituto de Ciencia de Materiales de Madrid (ICMM), Consejo Superior de Investigaciones Científicas (CSIC), Sor Juana Inés de la Cruz 3, 28049 Madrid, Spain

²Department of Physics, Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel

We present an analytical formulation of the thermodynamics, free energy and entropy, of any generic Bogoliubov de Genes model which develops exceptional point (EP) bifurcations in its complex spectrum when coupled to reservoirs. We apply our formalism to a non-Hermitian Josephson junction where, despite recent claims, the supercurrent does not exhibit any divergences at EPs. The entropy, on the contrary, shows a universal jump of 1/2 log 2 which can be linked to the emergence of Majorana zero modes (MZMs) at EPs. Our method allows us to obtain precise analytical boundaries for the temperatures at which such Majorana entropy steps appear. We propose a generalized Maxwell relation linking supercurrents and entropy which could pave the way towards the direct experimental observation of such steps in e.g. quantum-dot based minimal Kitaev chains.

Introduction-At weak coupling, an external environment only induces broadening and small shifts to the levels of a quantum system. In contrast, the strong coupling limit is highly nontrivial and gives rise to many interesting concepts in e.g. quantum dissipation [1], quantum information science [2] or quantum thermodynamics [3], just to name a few. An interesting example is the emergence of spectral degeneracies in the complex spectrum (resulting from integrating out the environment), also known as exceptional point (EP) bifurcations, where eigenvalues and eigenvectors coalesce [4]. During the last few years, a great deal of research is being developed in so-called non-Hermitian (NH) systems with EPs, in various contexts including open photonic systems [5], Dirac [6], Weyl [7] and topological matter in general [8–11]. The role of NH physics and EP bifurcations in systems with Bogoliubov-de Gennes (BdG) symmetry has hitherto remained unexplored, until recently. Specifically, there is an ongoing debate on how to correctly calculate the free energy in open BdG systems, a question relevant in e.g Josephson junctions coupled to external electron reservoirs. Depending on different approximations, such "NH junctions" have been predicted to exhibit exotic effects including imaginary persistent currents [13, 14] and supercurrents [15, 16] or various transport anomalies at EPs [17]. If one instead uses the biorthogonal basis associated with the NH problem [18], or an extension of scattering theory to include external electron reservoirs [19], the supercurrents are real and exhibit no anomalies. At the heart of this debate is whether a straightforward use of the complex spectrum plugged into textbook definitions of thermodynamic functions leads to meaningful results or whether, on the contrary, NH physics needs to be treated with care when calculating the free energy.

We here present a well-defined procedure, valid for arbitrary coupling and temperature, which allows us to calculate the free energy, Eq. (4), without any divergences at EPs. Derivatives of this free energy, allow us to calculate physical observables such as entropy (6) or supercurrents (7). Interestingly, entropy changes of $\log 2/2$ can be connected to emergent Majorana zero modes (MZMs) at EPs [20–22]. While such fractional entropy steps were predicted before in seemingly different contexts [23–25], our analysis in terms of EPs allows us to obtain precise analytical boundaries for the temperatures at which they appear. We propose a novel Maxwell relation connecting supercurrents and entropy, Eq. (12), which would allow the experimental detection of the effects predicted here.

Exceptional points in open BdG models– The starting point of our analysis is the description of the open quantum system in terms of a Green's function

$$G^{\text{eff}}(\omega) = [\omega - H_{\text{eff}}(\omega)]^{-1}, \qquad (1)$$

where $H_{\text{eff}}(\omega) = H_Q + \Sigma^r(\omega)$ is an effective NH Hamiltonian which takes into account how the quantum system H_{Q} is coupled to an external environment through the retarded self-energy $\Sigma^{r}(\omega)$. In what follows, we consider the case where an electron reservoir induces a tunneling rate [39], such that the complex poles of Eq. (1) have a well-defined physical interpretation in terms of quasibound states. If H_{eff} is a BdG Hamiltonian, electron-hole symmetry can be satisfied in two non-equivalent ways: (i) one can have pairs of poles with opposite real parts and with the same imaginary part $\epsilon^{\pm} = \pm E - i\gamma = -(\epsilon^{\mp})^*$; or, alternatively, (ii) two independent and purely imaginary poles $\epsilon^{\pm} = -i\gamma^{\pm} = -(\epsilon^{\pm})^*$. A bifurcation of the former, corresponding to standard finite-energy BdG modes with an equal decay to the reservoir γ , into the latter, two MZMs with different decay rates [20–22], defines an EP (Fig. 1a, inset).

NH Free energy–Calculating the free energy of an open quantum system is nontrivial since a direct substitution of a complex spectrum $\epsilon_j = E_j - i\gamma_j$ in the standard expression $F = -\frac{1}{\beta} \log Z = -\frac{1}{\beta} \sum_j \log \left(1 + e^{-\beta(\epsilon_j - \mu)}\right)$, can lead to complex results and divergences after an EP [15, 16]. To avoid inconsistencies, one possibility is to use the occupation $\langle N \rangle = \int_{-\infty}^{\infty} d\omega G^{<}(\omega) = -\frac{\partial F}{\partial \mu}$, which is a



Figure 1: Top (inset): Formation of an exceptional point. The complex eigenvalues of a NH BdG Hamiltonian evolve as a function of some parameter until they coalesce at a so-called EP and then bifurcate into two purely imaginary eigenvalues with different decay rates to the reservoir γ^{\pm} , (quasi-bound MZMs). Top (main): Entropy development after an exceptional point. Before the EP, both eigenvalues have identical absolute values by particlehole symmetry, $\epsilon^{\pm} = \pm E - i\gamma$, and thus both poles contribute to the entropy in (6) at the same temperature until the EP is reached $(T^{\text{EP}} = |\epsilon^{\pm}|/2 = \gamma/2)$, blue curve). After the EP, E = 0 and their imaginary parts are no longer identical, $\epsilon^{\pm} = -i\gamma^{\pm}$. Hence, their contributions to the entropy come at different temperatures $(T^{\pm} = \gamma^{\pm}/2)$, orange curve), giving rise to a fractional plateau $S = \log(2)/2$ of width $\gamma_{\rm asym} = (\gamma^- - \gamma^+)/2$. Bottom: Free energy changes in the temperature dependence (by more than six decades) of Fare barely distinguishable before and after an EP, as opposed to the entropy above $S = -\partial F / \partial T$, which illustrates the subtlety of the calculations presented here. The inclusion of a finite cutoff leads to a convergent low-temperature asymptotic limit $F_{T\to 0} = \frac{1}{2} \sum_{j} \frac{\gamma_j}{\pi} (\log \frac{\gamma_j}{D} - 1)$ see Eq. (5) for $E_j = \mu = 0$, which cures a divergence $F_{T\to 0} = \frac{1}{2} \sum_{j} \frac{\gamma_j}{\pi} (\log \frac{\gamma_j}{2\pi T} - 1)$ [12].

well-defined quantity in open quantum systems, even in non-equilibrium situations where the so-called Keldysh lesser Green's function $G^{<}(\omega)$ can be generalized beyond the fluctuation-dissipation theorem [26]. In BdG language, $\langle N \rangle$ can be written as

$$\langle N \rangle = \frac{1}{2} \int d\omega \,\Omega(D) \left[\rho^p(\omega) f(\omega - \mu) + \rho^h(\omega) f(-\omega - \mu) \right],$$
(2)

where $f(\omega)$ is the Fermi-Dirac function and we have ex-

plicitly separated the total spectral function

$$\rho(\omega) = -\frac{1}{2\pi} \operatorname{Im} \operatorname{Tr} G^{\text{eff}}(\omega) = -\frac{1}{2\pi} \operatorname{Im} \sum_{j} \frac{1}{\omega - \epsilon_{j}} , \quad (3)$$

in its particle ($\operatorname{Re}(\epsilon_j) > 0$) and hole ($\operatorname{Re}(\epsilon_j) < 0$) branches, $\rho^p(\omega)$ and $\rho^h(\omega)$, respectively [40]. We have also added a Lorentzian cutoff $\Omega(D) = D^2/(D^2 + \omega^2)$ to avoid divergences in the thermodynamic quantities as $T \to 0$ (see Supplemental Material [12]). The integral in Eq. (2) can be analytically solved by residues [12] and then be used to obtain F as:

$$F = -\int d\mu \langle N \rangle = \frac{1}{2} \sum_{j} \left[2T \operatorname{Re} \log \Gamma \left(\frac{1}{2} + \frac{i\gamma_j - E_j + \mu}{2i\pi T} \right) - \frac{2T\gamma_j}{D} \operatorname{Re} \log \Gamma \left(\frac{1}{2} + \frac{D + i\mu}{2\pi T} \right) + h(T, E_j, \gamma_j) \right],$$
(4)

with $\log \Gamma(z)$ being the log-gamma function and $h(T, E_j, \gamma_j)$ a generic function coming from the integration. We now perform the limits $\gamma_j = 0$ and $T \to 0$ of the previous expression,

$$F_{\gamma=0} = \frac{1}{2} \sum_{j} \left[T \log(2\pi) - T \log\left(2 \cosh\frac{E_{j} - \mu}{2T}\right) - \frac{\mu}{2} + h(T, E_{j}, 0) \right] ,$$

$$F_{T \to 0} = \frac{1}{2} \sum_{j} \left[\frac{\gamma_{j}}{\pi} \log\frac{\sqrt{\gamma_{j}^{2} + (E_{j} - \mu)^{2}}}{D} - \frac{\mu}{2} - \frac{E_{j} - \mu}{\pi} \arctan\frac{E_{j} - \mu}{\gamma_{j}} + h(0, E_{j}, \gamma_{j}) \right] ,$$
(5)

which, by comparison with well-known limits [27, 28], give $h(T, E_j, \gamma_j) = -\gamma_j/\pi - T \log(2\pi)$ [41]. From now on, we fix $\mu = 0$ but the complete derivation with full expressions, including $\mu \neq 0$, can be found in [12]. Derivatives of Eq. (4) allow us to obtain relevant thermodynamic quantities, as we discuss now.

Entropy steps from EPs- Using the free energy in Eq. (4), the entropy, defined as $S = -\partial F/\partial T$, reads:

$$S = \frac{1}{2} \sum_{j} \left[\log(2\pi) - 2 \operatorname{Re} \log \Gamma \left(\frac{1}{2} + \frac{i\gamma_{j} - E_{j}}{2i\pi T} \right) + \frac{2\gamma_{j}}{D} \log \Gamma \left(\frac{1}{2} + \frac{D}{2\pi T} \right) + \frac{\gamma_{j}}{\pi T} \operatorname{Re} \psi \left(\frac{1}{2} + \frac{i\gamma_{j} - E_{j}}{2i\pi T} \right) - \frac{E_{j}}{\pi T} \operatorname{Im} \psi \left(\frac{1}{2} + \frac{i\gamma_{j} - E_{j}}{2i\pi T} \right) - \frac{\gamma_{j}}{\pi T} \psi \left(\frac{1}{2} + \frac{D}{2\pi T} \right) \right],$$
(6)

where $\psi(z)$ is the digamma function. From Eq. (6), we can define a critical temperature $T_j = |\epsilon_j|/2$ as the inflection point when the eigenvalue ϵ_j begins to have a non-zero contribution $(S_j = \log(2)/4)$ to the total entropy, Fig. 1 (top). Hence, two standard BdG poles $\epsilon^{\pm} = \pm E - i\gamma$ will contribute at the same temperature to the entropy of the system since their absolute values are equal, $|\epsilon^{\pm}| = \sqrt{E^2 + \gamma^2}$, and thus a single plateau of $S = \log(2)$ can be measured. On the contrary, after an EP at $\epsilon^+ = \epsilon^- = -i\gamma$, the poles bifurcate taking zero real parts and different imaginary parts, $\epsilon^+ = -i\gamma^+$ and $\epsilon^- = -i\gamma^-$, and separating from each other a distance $\gamma^- - \gamma^+$. Then, each pole will contribute to the entropy at a different temperature $T^{\pm} = |\epsilon^{\pm}|/2 = \gamma^{\pm}/2$, giving rise to a fractional plateau $S = \log(2)/2$ of width $T^- - T^+ = (\gamma^- - \gamma^+)/2$. Here it is very important to point out that this nontrivial behavior of the entropy seems absent in the free energy that we used for the calculation of $S = -\partial F/\partial T$. Indeed, F is seemingly insensitive to any EP bifurcation even when varying the temperature over six decades, Fig. 1 (bottom).

Non-Hermitian Josephson junction with EPs– Our method also allows to calculate the supercurrent of any generic NH Josephson junction (or similarly the persistent current through a normal ring [18]) by just considering a phase-dependent spectrum $\epsilon_j(\phi)$ and taking phase derivatives of Eq. (4) as $I = \frac{\partial F}{\partial \Phi} = \frac{2e}{\hbar} \frac{\partial F}{\partial \phi}$, which gives

$$I(\phi) = \frac{e}{\hbar} \sum_{j} \left[-\frac{1}{\pi} \operatorname{Im} \psi \left(\frac{1}{2} + \frac{i\gamma_{j} - E_{j}}{2i\pi T} \right) \frac{\partial E_{j}}{\partial \phi} + \frac{1}{\pi} \operatorname{Re} \psi \left(\frac{1}{2} + \frac{i\gamma_{j} - E_{j}}{2i\pi T} \right) \frac{\partial \gamma_{j}}{\partial \phi} - \frac{1}{\pi} \frac{\partial \gamma_{j}}{\partial \phi} \qquad (7)$$
$$- \frac{2T}{D} \log \Gamma \left(\frac{1}{2} + \frac{D}{2\pi T} \right) \frac{\partial \gamma_{j}}{\partial \phi} \right].$$

As $T \to 0$, the supercurrent carried by a pair of BdG poles simply becomes [42]

before EP:
$$I_{T\to 0} = -\frac{e}{\hbar} \frac{2}{\pi} \arctan\left(\frac{E}{\gamma}\right) \frac{\partial E}{\partial \phi}$$
,
after EP: $I_{T\to 0} = \frac{e}{2\hbar} \frac{2}{\pi} \log\left(\frac{\gamma^+}{\gamma^-}\right) \frac{\partial \gamma^+}{\partial \phi}$, (8)

which, for example, allows us to calculate the supercurrent carried by Andreev levels in a short junction coupled to an electron reservoir almost straightforwardly [12]. Note that, although $I(\phi)$ has a cutoff-dependent term, it cancels by the particle-hole symmetry of the problem [12]. Eqs. (8) strongly differ from a calculation using directly the complex spectrum [15, 16]

$$I^{\text{alt}}(\phi) = \frac{e}{\hbar} \left(\frac{\partial E}{\partial \phi} - i \frac{\partial \gamma}{\partial \phi} \right) . \tag{9}$$

Eqs. (4), (6) and (7) are the main results of this paper and allow to calculate thermodynamics from generic open BdG models (arbitrary coupling and temperature) that can be written in terms of complex poles (Eq. (1)).

Non-Hermitian minimal Kitaev Josephson junction-As an application we now consider a quantum dot (QD)



Figure 2: Top: Schematic illustration of the four-Majorana Josephson junction device. Each segment comprises two quantum dots connected via a middle superconductor in a so-called minimal Kitaev geometry. In the low-energy regime, only the Majorana modes located at the edges are considered in the model. The two inner modes are connected through a weak link, coupling t_{23} , which defines a Josephson junction. Its superconducting phase difference ϕ can be controlled by the magnetic flux $\Phi = \frac{\hbar}{2e}\phi$ through a SQUID loop connecting the superconductors, with the cross denoting an ancillary junction, see e.g. [29, 30]. The internal QDS are coupled to reservoirs with rates $\gamma_2^0 \neq \gamma_3^0$. Center: Energy spectrum of the junction showing an **EP near phase** $\phi = \pi$. Lightblue/yellow (darkblue/red) lines correspond to real/imaginary parts of $\epsilon_{\text{outer}}^{\pm}$ ($\epsilon_{\text{inner}}^{\pm}$). Innermost states bifurcate around $\phi = \pi$, presenting two different topological phases (pink/green) divided by a pair of EPs. Gray lines correspond to the closed system ($\gamma_i^0 = 0$). Bottom: Phase dependence of supercurrent. Colored curves correspond to different temperatures (see legend). The black curve is associated with the Hermitian (closed) analog at $T = 10^{-8}$, assuming equilibrium occupations (hence no 4π Josephson effect) which shows the typical sawtooth-like profile for a perfectly transparent Andreev level [12]. The dashed line shows the calculation using the real part of Eq. (9). System parameters are fixed as $t_{12} = \Delta_{12}$, $t_{34} = \Delta_{34}$, $t_{23} = \gamma_2^0 = \Delta$ and $\gamma_3^0 = 0$.

array in a so-called minimal Kitaev model

$$H_{\rm QD} = -\sum_{i} \mu_i c_i^{\dagger} c_i - t_{i,i+1} c_i^{\dagger} c_{i+1} + \Delta_{i,i+1} c_i c_{i+1} + \text{H.c.} ,$$
(10)

where c_i^{\dagger} (c_i) denote creation (annihilation) operators

on each QD with a chemical potential μ_i . The QDs couple via a common superconductor that allows for crossed Andreev reflection and single-electron elastic cotunneling, with coupling strengths $\Delta_{i,i+1}$ and $t_{i,i+1}$, respectively. Remarkably, only two QDs are enough to host two localized MZMs η_1 and η_2 when a so-called sweet spot is reached with $\Delta_{1,2} = t_{1,2}$. This theoretical prediction [31] has recently been experimentally implemented [32, 33]. Let consider now a second double QD array (Majoranas η_3 and η_4) that forms a Josephson junction with the former array with a coupling $H_{\rm JJ} = -t_{2,3}e^{i\frac{\phi}{2}}c_2^{\dagger}c_3 + {\rm H.c.}, \text{ with } \phi \text{ being the supercon-}$ ducting phase difference between both arrays and $t_{2,3}$ the tunneling coupling between inner QDs. If, additionally, the two inner QDs are coupled to normal reservoirs with rates γ_2^0 and γ_3^0 this system is a realization of a Non-Hermitian Josephson junction containing Majorana modes (see the sketch in Fig. 2 top). In the low-energy regime, this model can be described in terms of four Majorana modes [34]. Assuming $\mu_i = 0 \ \forall i$ and $\Delta_{12} = t_{12}$ and $\Delta_{34} = t_{34}$, only the inner Majoranas η_2 and η_3 are coupled and lead to BdG fermionic modes of energy $\epsilon_{\text{inner}}^{\pm} = -\frac{i}{2}\gamma_0 \pm \frac{1}{2}\Lambda(\phi)$, where $\gamma_0 =$ $\gamma_2^0 + \gamma_3^0$, $\delta_0 = \gamma_2^0 - \gamma_3^0$ and $\Lambda(\phi) = \sqrt{2t_{23}^2(1 + \cos \phi) - \delta_0^2}$, while the outer modes remain completely decoupled and $\epsilon_{\text{outer}}^{\pm} = 0$. For $\delta_0 = 0$, one recovers the standard Majorana Josephson term: $\epsilon_{\text{inner}}^{\pm} = -\frac{i}{2}\gamma_0 \pm t_{23}\cos(\phi/2)$. When $\delta_0 \neq 0$, the spectrum develops EPs at phases $\phi_{\text{EP}} = \arccos\left[\frac{\delta_0^2}{2t_{23}^2} - 1\right]$. For an example of the resulting supercurrents and a comparison against Eq. (9), see Fig. 2 bottom. Similarly, when $\Delta_{12} = \Delta_{34} = \Delta$, $t_{12} = t_{34} = t$ but $t \neq \Delta$, an additional pair of EPs appears at [43] $\phi_{\rm EP} = \arccos\left[\frac{\delta_0^2 - [4(\Delta - t) - \gamma_0]^2}{2t_{23}^2} - 1\right].$

Using these analytics, the critical temperatures $T_j = |\epsilon_j|/2$ read

before EP:
$$T_{\text{inner}}^{\pm} = \frac{1}{4} \sqrt{\gamma_0^2 + \Lambda^2(\phi)}$$
,
after EP:
$$\begin{cases} T_{\text{inner}}^+ = \frac{\gamma_0 - \sqrt{\delta_0^2 - 2t_{23}^2(1 + \cos\phi)}}{4} & (11) \\ T_{\text{inner}}^+ = \frac{\gamma_0 + \sqrt{\delta_0^2 - 2t_{23}^2(1 + \cos\phi)}}{4} \end{cases}$$

To illustrate their physical meaning, we plot a full calculation of the entropy $S(T, \phi)$ using Eq. (6), Fig. 3 top, together with the analytical expressions in Eq. (11) (solid lines). This plot demonstrates that changes in entropy can be understood from EPs, a claim that is even clearer by analyzing cuts at fixed phase (Fig. 3 center). Interestingly, a universal entropy change of $1/2 \log 2$ can be linked to the emergence of MZMs at phase $\phi = \pi$ as $T \rightarrow 0$. Alternatively, the entropy loss due to Majoranas can be seen in phase-dependent cuts taken at different temperatures, Fig. 3 bottom, which show as an interesting behavior where a $S = 2 \log 2$ plateau at large temperatures becomes an emergent narrow resonance, centered at $\phi = \pi$ and of height $S = 1.5 \log 2$, as T is lowered.



Figure 3: Entropy of a Majorana Josephson junction with EPs. Top: Entropy as a function of ϕ and T. Two EPs[±] are marked in white at ϕ_{EP} and $2\pi - \phi_{\text{EP}}$ (??). Black-/red curves correspond to T_{inner}^{\pm} , agreeing with the jumps in entropy. Center: Entropy as a function of T for different phases. We have also marked the position of T_{inner}^{\pm} for each curve. Bottom: Entropy as a function of ϕ for different temperatures. Gray intermediate steps follow the trajectory in between such colored curves. System parameters are fixed as $t_{12} = \Delta_{12}$, $t_{34} = \Delta_{34}$, $t_{23} = \gamma_2^0 = \Delta$ and $\gamma_3^0 = 0$.

Experimental detection of fractional entropy- recently it has been demonstrated that one can measure entropies of mesoscopic systems, either via Maxwell relations [35, 36] or via thermopower [37, 38]. The Maxwell relation method relies on continuously changing a parameter x (e.g. chemical potential or magnetic field) while measuring its conjugate variable y (e.g. electron number or magnetization, respectively) such that $y = \partial F/\partial x$. Then, the Maxwell relation yields $\partial S/dx = -\partial y/dT$. Here we propose a novel application of this procedure, employing the Josephson current $I(\phi)$, which gives

$$\Delta S_{\phi_1 \to \phi_2} = -\int_{\phi_1}^{\phi_2} \frac{\partial I(\phi)}{\partial T} \, d\phi \;. \tag{12}$$

Since the phase difference on the Josephson junction can be controlled by e.g. embedding it in a SQUID loop [29, 30], Fig. 2 top, one can integrate dI/dT between $\phi_1 = 0$ and $\phi_2 = \pi$. From Fig. 3 (bottom) we expect ΔS to change from zero at high-T to $\log 2/2$ at low-T, an unequivocal signature of Majorana zero modes in the junction.

Acknowledgements–DMP and RA acknowledge financial support from the Horizon Europe Framework Program of the European Commission through the European Innovation Council Pathfinder Grant No. 101115315 (QuKiT), the Spanish Ministry of Science through Grants No. PID2021-125343NB-I00 and No. TED2021-130292B-C43 funded by

- U. Weiss, *Quantum Dissipative Systems* (World Scientific Pub Co Inc; 5th edition, 2021).
- [2] P. M. Harrington, E. J. Mueller, and K. W. Murch, Nature Reviews Physics 4, 660 (2022), URL https: //doi.org/10.1038/s42254-022-00494-8.
- [3] A. Rivas, Phys. Rev. Lett. **124**, 160601 (2020), URL https://link.aps.org/doi/10.1103/PhysRevLett. 124.160601.
- M. V. Berry, Czechoslovak Journal of Physics 54, 1039 (2004), URL https://doi.org/10.1023/B:CJ0P. 0000044002.05657.04.
- [5] J. Doppler, A. A. Mailybaev, J. Böhm, U. Kuhl, A. Girschik, F. Libisch, T. J. Milburn, P. Rabl, N. Moiseyev, and S. Rotter, Nature 537, 76 (2016).
- [6] B. Zhen, C. W. Hsu, Y. Igarashi, L. Lu, I. Kaminer, A. Pick, S.-L. Chua, J. D. Joannopoulos, and M. Soljačić, Nature 525, 354 (2015).
- [7] A. Cerjan, S. Huang, M. Wang, K. P. Chen, Y. Chong, and M. C. Rechtsman, Nature Photonics 13, 623 (2019).
- [8] H. Shen, B. Zhen, and L. Fu, Phys. Rev. Lett. 120, 146402 (2018), URL https://link.aps.org/doi/10. 1103/PhysRevLett.120.146402.
- [9] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, Phys. Rev. X 8, 031079 (2018), URL https://link.aps.org/doi/10. 1103/PhysRevX.8.031079.
- [10] K. Kawabata, K. Shiozaki, M. Ueda, and M. Sato, Phys. Rev. X 9, 041015 (2019), URL https://link.aps.org/ doi/10.1103/PhysRevX.9.041015.
- [11] E. J. Bergholtz, J. C. Budich, and F. K. Kunst, Rev. Mod. Phys. 93, 015005 (2021), URL https://link.aps. org/doi/10.1103/RevModPhys.93.015005.
- [12] See supplemental material at http://link.aps.org/ supplemental/ for further details on.
- [13] S.-B. Zhang, M. M. Denner, T. c. v. Bzdušek, M. A. Sentef, and T. Neupert, Phys. Rev. B 106,

MCIN/AEI/10.13039/501100011033, "ERDF A way of making Europe" and European Union Next Generation EU/PRTR, as well as the CSIC Interdisciplinary Thematic Platform (PTI+) on Quantum Technologies (PTI-QTEP+). YM acknowledges support from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme under Grant Agreement No. 951541 and the Israel Science Foundation Grant No. 154/19.

Note added While finishing this manuscript, two recent preprints in Arxiv [18] and [19] also pointed out the subtleties of calculating the free energy in a NH Josephson junction. Eq. (9) in Ref. [18] for the supercurrent agrees with our Eq. (7) in the limit without cutoffs $D \to \infty$. Moreover, Eq. (16) in Ref. [19] agrees with our Eq. (7) in the regime without EPs. This latter case, in particular, results in a reduction factor $\frac{2}{\pi} \arctan\left(\frac{E}{\gamma}\right)$ in the $T \to 0$ supercurrent, see Eq. (8), owing to the coupling with the reservoir.

L121102 (2022), URL https://link.aps.org/doi/10. 1103/PhysRevB.106.L121102.

- [14] Q. Li, J.-J. Liu, and Y.-T. Zhang, Phys. Rev. B 103, 035415 (2021), URL https://link.aps.org/doi/10. 1103/PhysRevB.103.035415.
- [15] J. Cayao and M. Sato (2023), 2307.15472.
- [16] C.-A. Li, H.-P. Sun, and B. Trauzettel (2024), 2307.04789.
- [17] V. Kornich, Phys. Rev. Lett. **131**, 116001 (2023), URL https://link.aps.org/doi/10.1103/PhysRevLett. 131.116001.
- [18] P.-X. Shen, Z. Lu, J. L. Lado, and M. Trif (2024), 2403.09569.
- [19] C. W. J. Beenakker (2024), 2404.13976.
- [20] D. I. Pikulin and Y. V. Nazarov, Phys. Rev. B 87, 235421 (2013), URL https://link.aps.org/doi/10. 1103/PhysRevB.87.235421.
- [21] P. San-Jose, J. Cayao, E. Prada, and R. Aguado, Scientific Reports 6, 21427 (2016).
- [22] J. Avila, F. Peñaranda, E. Prada, P. San-Jose, and R. Aguado, Communications Physics 2, 133 (2019), ISSN 2399-3650, URL https://doi.org/10. 1038/s42005-019-0231-8.
- [23] S. Smirnov, Phys. Rev. B 92, 195312 (2015).
- [24] E. Sela, Y. Oreg, S. Plugge, N. Hartman, S. Lüscher, and J. Folk, Phys. Rev. Lett. **123**, 147702 (2019), URL https://link.aps.org/doi/10.1103/PhysRevLett. 123.147702.
- [25] V. F. Becerra, M. Trif, and T. Hyart, Phys. Rev. Lett. 130, 237002 (2023), URL https://link.aps.org/doi/ 10.1103/PhysRevLett.130.237002.
- [26] Y. Meir and N. S. Wingreen, Phys. Rev. Lett. 68, 2512 (1992), URL https://link.aps.org/doi/10. 1103/PhysRevLett.68.2512.
- [27] A. Zagoskin, Quantum Theory of Many-Body Systems: Techniques and Applications (Springer, 2014).

- [28] A. C. Hewson, *The Kondo Problem to Heavy Fermions*, Cambridge Studies in Magnetism (Cambridge University Press, 1993).
- [29] A. Bargerbos, M. Pita-Vidal, R. Žitko, L. J. Splitthoff, L. Grünhaupt, J. J. Wesdorp, Y. Liu, L. P. Kouwenhoven, R. Aguado, C. K. Andersen, et al., Phys. Rev. Lett. 131, 097001 (2023), URL https://link.aps.org/ doi/10.1103/PhysRevLett.131.097001.
- [30] M. Pita-Vidal, A. Bargerbos, R. Žitko, L. J. Splitthoff, L. Grünhaupt, J. J. Wesdorp, Y. Liu, L. P. Kouwenhoven, R. Aguado, B. van Heck, et al., Nature Physics 19, 1110 (2023).
- [31] M. Leijnse and K. Flensberg, Phys. Rev. B 86, 134528 (2012).
- [32] T. Dvir, G. Wang, N. van Loo, C.-X. Liu, G. P. Mazur, A. Bordin, S. L. D. ten Haaf, J.-Y. Wang, D. van Driel, F. Zatelli, et al., Nature 614, 445 (2023).
- [33] S. L. D. ten Haaf, Q. Wang, A. M. Bozkurt, C.-X. Liu, I. Kulesh, P. Kim, D. Xiao, C. Thomas, M. J. Manfra, T. Dvir, et al. (2024), 2311.03208.
- [34] D. M. Pino, R. S. Souto, and R. Aguado, Phys. Rev. B 109, 075101 (2024), URL https://link.aps.org/doi/ 10.1103/PhysRevB.109.075101.
- [35] N. Hartman, C. Olsen, S. Lüscher, M. Samani, S. Fallahi, G. C. Gardner, M. Manfra, and J. Folk, Nat. Phys. 14, 1083 (2018).
- [36] T. Child, O. Sheekey, S. Lüscher, S. Fallahi, G. C. Gardner, M. Manfra, A. Mitchell, E. Sela, Y. Kleeorin, Y. Meir, et al., Phys. Rev. Lett. 129,

227702 (2022), URL https://link.aps.org/doi/10. 1103/PhysRevLett.129.227702.

- [37] Y. Kleeorin, H. Thierschmann, H. Buhmann, A. Georges, L. W. Molenkamp, and Y. Meir, Nat. Commun. 10, 5801 (2019).
- [38] E. Pyurbeeva, C. Hsu, D. Vogel, C. Wegeberg, M. Mayor, H. van der Zant, J. A. Mol, and P. Gehring, Nano Letters 21, 9715 (2021), pMID: 34766782, https://doi.org/10.1021/acs.nanolett.1c03591, URL https://doi.org/10.1021/acs.nanolett.1c03591.
- [39] We consider the simplest case where this selfenergy originates from a tunneling coupling to an external electron reservoir in the so-called wideband approximation where both the tunneling amplitudes and the density of states in the reservoir are energy independent, which allows to write the tunneling couplings as constant [26].
- [40] A global factor of 1/2 arises from the duplicity of dimensions in BdG formalism.
- [41] Interestingly, this term is relevant for the entropy since it adds a physical contribution $\log(2\pi)$. In contrast, the term $-\gamma_j/\pi$ cancels out in the supercurrent [12].
- [42] For a single pair of BdG poles, γ is independent/dependent of the phase before/after the EP and the other way around for E [12].
- [43] Analytical expressions of eigenvalues, a similar calculation for a junction containing Andreev levels, as well as detailed discussions can be found in the [12].