

# Time-dependent Aharonov-Bohm type topological effects on dipoles

H. O. Cildiroglu

<sup>a</sup>*Boston University, Physics Department, Boston, 02100, MA, USA*

<sup>b</sup>*Ankara University, Department of Physics Engineering, 06100, Ankara, Türkiye*

---

## Abstract

Exploring the time-dependent characteristics of AB-type effects holds significant importance in contemporary physics and its practical applications. Here, we delve into the investigation of time-dependent topological effects emerging in AB-type experimental setups. We first analyze the topological effects on magnetic dipoles moving in closed trajectories around the time-varying magnetic field source solenoid, then on electrical dipoles around a time-varying electric field source in 2+1 dimensions without any approximation. Last, we discuss the characteristics of the topological effects by considering the identity and dualities between phases from an integrated perspective.

**Keywords:** Time-Dependent Aharonov Bohm Effect, Aharonov-Casher Effect, He-McKellar-Wilkens Phase, Topological Phases

---

## 1. Introduction

Quantum mechanical topological effects are elucidated as the introduction of the vector or vector potential-like physical quantities as complex phase factors into the wave functions of particles moving along the closed trajectories around singularities created by electromagnetic field sources without the effects of the classical forces. One of the pioneering and first examples is the Aharonov-Bohm (AB) phase, which demonstrates the significance of electromagnetic potentials in quantum theory [1–3]. This complex phase exerts a measurable impact on the interference pattern of wave functions of electric charges reaching a screen via two distinct trajectories around an infinitely long solenoid as a static magnetic field source. Subsequently, various topological effects have emerged by considering Maxwell’s dualities. These include the Aharonov-Casher (AC) effect arises from the motion of magnetic dipoles around a linear electric charge distribution [4; 5], the He-McKellar-Wilkens (HMW) effect from the motion of electric dipoles around a linear magnetic charge distribution [6–8], and the dual AB effect (DAB) from the movement of magnetic charges around an electric field tube [9; 10].

In recent years, studies on AB-type effects have focused on investigating the time dependence of the effect. Singleton et al. discussed two covariant generalizations of the AB effect with time-dependent flux, noting that the AB phase shift is canceled by the phase shift of the external electric field associated with the Lorentz force [11], Bright et al. explored the time-dependent (TD) AB effect for non-Abelian gauge fields revealing cancellations between phase shifts from non-Abelian electromagnetic fields [12], Ababekri et al. examined the non-relativistic behavior of particles with electric dipoles on noncommutative space uncovering quantum phase corrections [13], Singleton et al. developed a covariant expression for the AC phase, investigating its interaction with electromagnetic fields [14], Jing et al. re-examined the AB effect in the

background of a time-dependent vector potential, highlighting alterations in interference patterns [15], Ma et al. explored noncommutative corrections to the TD-AB effect by revealing three types of corrections and proposing dimensionless quantities for parameter extraction based on measured phase shifts [16], Choudhury et al. discovered a frequency-dependent AB phase shift [17], Jing et al. revisited the TD-AB effects in noncommutative space-time, finding no noncommutative corrections to the AB effects for both cases up to the first order of the noncommutative parameter [18], Wang et al. investigated the TD-HMW effect in noncommutative space by confirming gauge symmetry, and time-dependent AC effect and its corrections due to spatial noncommutativity on noncommutative space [19; 20], Saldanha proposed an electrodynamic AB scheme challenging the topological nature of the phase [21], and Wakamatsu et al. analyzed the AB effect’s interaction energy and its gauge invariance [22].

In this letter, we study the TD-AB type topological effects of time-varying electromagnetic fields on dipoles in 2+1 dimensions. We begin with a TD solenoidal magnetic field oriented in the  $z$  direction, and the resulting rotational electric field appearing in the  $xy$  plane. To ensure that particles are not subject to classical forces, we use magnetic dipole carrier chargeless particles. Accounting for the relativistic electromagnetic effects arising from the motion of the particles, we give direct calculations for appearance of a vector-potential-like physical quantity which enters as a complex observable phase in the wave functions of the particles. We conclude that the resulting phase is topological. Accordingly, we emphasize that, although there is no direct interaction, the phase involves the time-rate change of the energy term, and the interference pattern can be controlled by means of a changing magnetic field (i.e. flux). Last, we consider the fully dual of this problem and show the topological effects on electric dipole moment carrier chargeless particles moving around a time-varying electric field.

## 2. Time-Dependent AB-type Effect on Magnetic Dipoles

In an AB-type experimental setup, as a time-dependent magnetic field source solenoid oriented in the  $z$ -direction is placed just behind the double-slits (See Fig. 1). The solenoid with the radius of  $a$ , extends to infinity to prevent the presence of magnetic fields in the region where the particles are allowed to move. In accordance with the nature of AB-type effects and to ensure that the particles are not subject to a classical Lorentz force, we use magnetic dipole moment carrier chargeless particles. In this scenario, the particles are expected to remain unaffected, as there is no magnetic field in the region where the particles are allowed to move ( $\nabla \times \mathbf{E}_{r>a} = 0$ ) from the classical perspective. Nevertheless, as the particles are in motion, relativistic electromagnetic effects arise, and the wave functions of the particles acquire a measurable geometric/topological phase. Accordingly, it is appropriate to start with the relativistic Lagrangian describing the motion of magnetic dipoles under electromagnetic fields,

$$L = \bar{\psi} \left( i\gamma^\mu \partial_\mu - m + \frac{\mu}{2} \sigma^{\mu\nu} F_{\mu\nu} \right) \psi \quad (1)$$

and the corresponding equations of motion derived from this Lagrangian:

$$\left( i\gamma^\mu \partial_\mu - m + \frac{\mu}{2} \sigma^{\mu\nu} F_{\mu\nu} \right) \psi = 0 \quad (2)$$

Due to the symmetry of the problem, there is invariance under spatial translations along the  $z$ -direction, making an examination in 2+1 dimensions sufficient for the emergence of quantum mechanical effects. In 2+1 dimensional spacetime (+, -, -), gamma matrices change from their well-known four-dimensional representations. Therefore, the  $\sigma^{\mu\nu}$  term in equation (2) needs to be redefined. With the selection of the free Dirac Hamiltonian  $H_D$ ,

$$H_D = \alpha_x p_x + \alpha_y p_y + m\beta \quad (3)$$

the gamma matrices can be defined in two-dimensional space in two different representations satisfying the Clifford algebra ( $s = \pm 1$ ) for the up and down spin polarization states:

$$\begin{aligned} \alpha_x &= \sigma_x & \beta &= \gamma_0 = \sigma_z & \gamma_1 &= \beta\alpha_x = i\sigma_y \\ \alpha_y &= s\sigma_y & \{\gamma^0, \gamma^i\} &= 0 & \gamma_2 &= \beta\alpha_y = -is\sigma_x \end{aligned} \quad (4)$$

Hence, by considering the magnetic field on the plane is  $\mathbf{B} = 0$ , and the transformation  $F_{\mu\nu} = \eta_{\mu\alpha} F^{\alpha\beta} \eta_{\beta\nu}$ , the  $\sigma_{\mu\nu} F^{\mu\nu}$  term can be written,

$$\sigma^{\mu\nu} F_{\mu\nu} = 2\sigma^{0i} F_{0i} + \sigma^{ij} F_{ij} = 2i\alpha_i E_i = 2i(\sigma_x E_x + s\sigma_y E_y) \quad (5)$$

Here,  $\sigma^{0i} = i\alpha_i$ ,  $\sigma^{ij} = \sigma^k$ . Now, one can multiply (2) by  $\gamma^0$  from the left after separating the space and time components of the covariant derivative, it becomes,

$$\left[ m\beta + \alpha_x p_x + \alpha_y p_y + s\mu(\alpha_y E_x - \alpha_x E_y) \right] \psi = i\partial_t \psi \quad (6)$$




Figure 1: Schematic representation of moving magnetic dipoles around a time-dependent magnetic field source solenoid with the radius  $a$  in AB-type setup.

In this case, it is evident that the expression inside the parentheses on the right-hand side is the total Hamiltonian ( $H$ ) of the system.

$$H = \left[ m\beta + \alpha_x p_x + \alpha_y p_y + s\mu(\alpha_y E_x - \alpha_x E_y) \right] \quad (7)$$

Thus, by considering (3), the interaction Hamiltonian  $\Delta H$  is as follows,

$$\Delta H = s\mu(\alpha_y E_x - \alpha_x E_y) \quad (8)$$

According to the current configuration, in order to ensure computational simplicity and confining the electric field to two dimensions, we define a new electric field  $\tilde{\mathbf{E}}(\mathbf{r}, t) = \mathbf{E} \times \hat{\mathbf{z}} = \hat{\mathbf{x}}E_y - \hat{\mathbf{y}}E_x$ . In this case, confined electric field vector can be written in terms of its components in the form  $(\tilde{E}_x, \tilde{E}_y) = (E_y, -E_x)$ . Thus, the interaction Hamiltonian can be rearranged as in the form of,

$$\Delta H = -s\mu(\alpha_x \tilde{E}_x + \alpha_y \tilde{E}_y) = -s\mu\alpha_\perp \cdot \tilde{\mathbf{E}} \quad (9)$$

The current stage is crucial as it allows the total Hamiltonian to be expressed as the sum of the free and interaction terms:

$$[H_D + \Delta H] \psi = i\partial_t \psi \quad (10)$$

In order to eliminate the interaction term  $\Delta H$ , a series of preparations are required. By using the definition  $\mathbf{p} = -i\nabla$ , we obtain:

$$\left[ m\beta - i\alpha_\perp \cdot (\nabla - is\mu\tilde{\mathbf{E}}) \right] \psi = i\partial_t \psi \quad (11)$$

Here, the term  $s\mu\tilde{\mathbf{E}}$  is vector potential-like quantity for magnetic dipoles ( $\tilde{\mathbf{A}} = s\mu\tilde{\mathbf{E}}$ ) [23–25]. Furthermore, in the Lorentz gauge ( $-\nabla \cdot \tilde{\mathbf{A}} = \partial_t \tilde{\phi}$ ), it gives rise to a scalar potential-like term ( $\tilde{\phi} = s\tilde{\mu} \cdot \mathbf{B}(\mathbf{t})$ ). In this context, considering the motion of chargeless particles in an electromagnetic field, by incorporating the transformations of the time ( $p^0$ ) and spatial ( $\mathbf{p}$ ) components of momentum in 2+1 dimensional spacetime:

$$\begin{aligned} p^0 &\longrightarrow p^0 - s\tilde{\mu} \cdot \mathbf{B} \\ \mathbf{p} &\longrightarrow \mathbf{p} - s\mu\tilde{\mathbf{E}} \end{aligned} \quad (12)$$

we can achieve the covariant derivatives under this configuration:

$$\begin{aligned} i\partial_t &\longrightarrow i\partial_t - s\vec{\mu} \cdot \mathbf{B} = i[\partial_t + is\vec{\mu} \cdot \mathbf{B}] = i\tilde{D}_t \\ -i\nabla &\longrightarrow -i\nabla - s\mu\tilde{\mathbf{E}} = -i[\nabla - is\mu\tilde{\mathbf{E}}] = -i\tilde{\mathbf{D}} \end{aligned} \quad (13)$$

Thus, there is no magnetic field presence in the region where the particles are allowed to move, following expression is obtained for moving dipoles,

$$[m\beta - i\alpha_\perp \cdot \tilde{\mathbf{D}}]\psi = i\tilde{D}_t\psi \quad (14)$$

The form covariance of the Dirac equation requires the invariance of  $\psi$ :

$$[m\beta - i\alpha_\perp \cdot \tilde{\mathbf{D}}']\psi' = i\tilde{D}_t'\psi' \quad (15)$$

Therefore,  $\psi$  undergoes a unitary transformation as  $\psi' = U\psi$ , the covariant derivative defined in (13) must transforms,

$$\begin{aligned} D_t\psi &\longrightarrow (D_t\psi)' = D_t'\psi' = U(D_t\psi) \\ \mathbf{D}\psi &\longrightarrow (\mathbf{D}\psi)' = \mathbf{D}'\psi' = U(\mathbf{D}\psi) \end{aligned} \quad (16)$$

In order to ensure unitarity, the transformation will be in the form  $U = e^{i\chi(\mathbf{r},t)}$ , where  $\chi(\mathbf{r},t)$  is either real or Hermitian. Now, it is explicit that for (16) to be satisfied, gauge transformations needs to be made to scalar and vector potential-like physical quantities. Taking into account the transformed forms of the time and spatial components of the covariant derivative, we arrive at the expression  $\tilde{\mathbf{A}}' = \tilde{\mathbf{A}} + \nabla\chi(\mathbf{r},t)$ . Finally, after the gauge-fixing  $\tilde{\mathbf{A}}' = 0$ , it becomes apparent that the gauge function ensuring this change must satisfy  $-\nabla\chi(\mathbf{r},t) = s\mu\tilde{\mathbf{E}}$ , and the function  $\chi(\mathbf{r},t)$  is obtained:

$$\chi(\mathbf{r},t) = -s\mu \int^x \tilde{\mathbf{E}}(\mathbf{r},t) \cdot d\mathbf{l} \quad (17)$$

Hence, the transformed Dirac spinor is,

$$\psi' = e^{is\mu\partial_t\Phi_B(t)}\psi \quad (18)$$

This results clearly demonstrate how the variation of the magnetic field affects the interference pattern, although there is no direct interaction of the moving dipoles with the magnetic field for  $r > a$ . Moreover, equations (17) and (18) clearly indicate that the phase shares the same characteristics as other topological effects, albeit with its own distinct features. If the integral on the right-hand side of (17) yields a linear charge distribution, the resulting effect would be an AC phase. However, incorporating the time variation of the field term indicates it would be a magnetic flux, thus described as an AB phase performed with magnetic dipoles. Here, as an illustration, let's choose the magnetic field as  $\mathbf{B}(t) = B_0 \sin \omega t \hat{\mathbf{z}}$ . Since the magnetic field is time-dependent, it induces a rotational electric field in the xy-plane:  $\mathbf{E}(\mathbf{r},t) = \frac{\pi a^2 \omega}{2r} B_0 \cos \omega t (-\hat{\mathbf{x}} \sin \theta + \hat{\mathbf{y}} \cos \theta)$ . In this case, the transformed Dirac spinor can be obtained explicitly as  $\psi' = e^{is\mu(\pi a^2 \omega B_0 \cos \omega t)}\psi$ .

### 3. Time-Dependent AB-type Effect on Electric Dipoles

In this section, as a complete dual of the discussion in the previous section, we consider electric dipole carrier chargeless particles moving around a time-varying electric field oriented in the z direction in an infinitely long flux tube with radius  $a$  (See Fig. 2). In the region where particles are permitted to move, the presence of a time-varying electric field induces a magnetic field, while no electric field is present. Given that the particles lack charge, they remain unaffected by classical forces. Nevertheless, relativistic electromagnetic effects manifest, leading to interaction between the moving dipoles and the magnetic field, as the field source moves within the inertial reference frame of the particles. Accordingly, it is appropriate to start with the relativistic Lagrangian describing the motion of electrical dipoles under electromagnetic fields:

$$L = \bar{\psi} \left( i\gamma^\mu \partial_\mu - m - \frac{i}{2} d \sigma^{\mu\nu} \gamma^5 F_{\mu\nu} \right) \psi \quad (19)$$

Here,  $m$  and  $d$  are the mass and the electric dipole moment of the chargeless particles, and the  $\gamma^5$  is a representation formed by products of other gamma matrices defined in Clifford algebra. The equations of motion that can be derived from equation (19) are in the form,

$$\left( i\gamma^\mu \partial_\mu - m - \frac{i}{2} d \sigma^{\mu\nu} \gamma^5 F_{\mu\nu} \right) \psi = 0. \quad (20)$$

The last term in parenthesis, using  $\tilde{F}^{\mu\nu} = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} F_{\alpha\beta}$ , can be rearranged as,

$$\frac{i}{2} d (\sigma^{\mu\nu} \gamma^5) F_{\mu\nu} = -\frac{d}{2} \sigma_{\alpha\beta} \tilde{F}^{\alpha\beta}. \quad (21)$$

Similar to the previous section, it is evident that the redefinition of the term  $\sigma_{\alpha\beta} \tilde{F}^{\alpha\beta}$  is necessary when the discussion is carried out in 2+1 dimensions. Thus, by definition  $\Sigma^k = \begin{pmatrix} \sigma^k & 0 \\ 0 & \sigma^k \end{pmatrix}$ , and in the region where  $\mathbf{E} = 0$  is allowed for the particle motions, the term  $\sigma_{\alpha\beta} \tilde{F}^{\alpha\beta}$  can be explicitly written in the form,

$$\sigma_{\alpha\beta} \tilde{F}^{\alpha\beta} = 2i\alpha_i \tilde{F}^{0i} + \epsilon_{ijk} \Sigma^k \tilde{F}^{ij} = 2i\alpha_\perp \cdot \mathbf{B} \quad (22)$$

Therefore, commencing with the initial Hamiltonian in (3) and its representation in (4), when (22) is multiplied by  $\gamma^0$  from the left, it becomes,

$$i\partial_t\psi = [m\beta + \alpha_x p_x + \alpha_y p_y + d(\sigma_y B_x - \sigma_x B_y)]\psi \quad (23)$$

The expression within the parentheses on the right-hand side represents the total Hamiltonian of the system. For ease of computation and to confine the magnetic field to the plane, we define  $\tilde{\mathbf{B}} = \mathbf{B} \times \hat{\mathbf{z}}$  and  $(\tilde{B}_x, \tilde{B}_y) = (B_y, -B_x)$ , thereby obtaining the interaction term as follows:

$$\Delta H = -sd(\alpha_x \tilde{B}_x + \alpha_y \tilde{B}_y) = -sd\alpha_\perp \cdot \tilde{\mathbf{B}} \quad (24)$$

It is clear that this term is fully dual to expression (9). Thus, the total Hamiltonian of the system is in the form




Figure 2: Schematic representation of moving electric dipoles around a time-dependent electric field source with the radius of  $a$ .

$(H = H_D + \Delta H_{VHMW})$ . At this stage, steps (10)- (17) are repeated with the electric-magnetic duality transformations  $\mu \rightarrow d$ ,  $E \rightarrow B$  and  $B \rightarrow -E$ , the phase and the transformed Dirac spinor are reached:

$$\xi(\mathbf{r}, t) = -sd \int^x \tilde{\mathbf{B}}(\mathbf{r}, t) \cdot d\mathbf{l}$$

$$\psi' = r^{isd\partial_t \Phi_E(t)} \psi. \quad (25)$$

The topological phase of an electric dipole under varying electric field is revealed. The electric field variation without any interaction has entered the wave function of the particles and has measurable effects on the interference pattern. The equation (25) exhibits a complete duality with (19). It also shows dependence on the orientation of the spin polarizations of the particles. If the integral for the function  $\xi(\mathbf{r}, t)$  given by (25) were to give a magnetic charge distribution, then the HMW phase would appear. However, this topological phase is completely dual to the AB-type phase introduced in the first section. In this respect, a review of the identity duality relations between the phases is left for the last section.

#### 4. Discussion and Conclusion

AB-type effects can be addressed in 2+1 dimensions due to the symmetry of the problems. Such an investigation, conducted without approximations, illuminates the duality and identity relationships between the phases clearly. Accordingly, in two-dimensional space, the AB phase can be thought of as emerging from the motion of electrons along closed trajectories around polarized magnetic dipoles, since a solenoid can be regarded as a coherent array of magnetic dipoles (See Fig. 3). Similarly, since the electric and magnetic linear charge distributions can be considered as electric and magnetic charges in two dimensions, the AC and HMW phases emerge as a result of the motion of unpolarized magnetic dipoles around electrons and unpolarized electric dipoles around magnetic charges in closed orbits, respectively. From this perspective, one can describe an




Figure 3: Schematic representations of topological phases in 2+1 dimensions. Here, the moving dipoles do not need to be polarized. This notation is utilized to reinforce the notation, and the polarization of the particles can be controlled by  $s$ . The above four phases are the four topological phases known in the literature as AB-type effects. There is a kind of identity between the AB-AC and DAB-HMW phases, provided that the moving dipoles are polarized. There is a duality relation between the left and right groups. These relations between the phases are explicit in the Hamiltonians obtained in 2+1 dimensions. Two setups given in the lower part represent schematizations of the topological phases proposed in this work. Unlike the other phases, in the condition of time dependence of the field sources, it arises as a result of the motion of the dipoles around the dipoles. Again, there is a complete duality between the two phases.

identity - rather than a duality - that emerges in the stationary reference frames of moving electrons (or polarized dipoles) between the AB and AC phases (the same relationship holds true between the DAB and HMW). However, it is clear that the AB and AC phases grouped on the left are fully dual to the DAB and HMW phases. On the other hand, AC and HMW effects depend on the polarization directions of the moving dipoles and have the same character. The AB and DAB effects do not depend on the spin orientation of the incoming electrons. All these relations can also be clearly seen from the interaction Hamiltonians describing the dynamics of the systems.

In this context, considering the scenarios proposed in the preceding sections, it becomes apparent that they correspond to topological phase structures consistent with the characteristics outlined for AB-type effects. Accordingly, the first phase, which is a combination of the AB and AC effects, is described by the motion of the moving magnetic dipoles around a variable magnetic field source. Moving particles are not required to be polarized. Their polarization can be controlled by using  $s$ , derived from representations of the Clifford algebra in 2+1 dimensions. It enters the phase factor ( $\tilde{A} = s\mu\tilde{E}$ ) as a vector potential-like physical quantity through the electric field present region where the particles are allowed to move and the resulting relativistic effects. Thus, although there is no direct interaction, the interference pattern is measurably affected by the

time-dependent variation in magnetic flux. In complete duality, a similar effect arises if electric dipoles are moved around time-dependent electric field source. Hence, the topological phase that emerges in the scenario discussed in the third section depends on the polarization states of moving particles and can be considered a combination of the HMW-DAB phases. Resultant phases align with the characteristics of AB-type effects, exhibiting complete duality between them. Furthermore, the phases are contingent upon the polarization state of the moving dipoles, and the interaction Hamiltonians take the form of AB-type effects. This opens up avenues for future studies, such as investigating the effects on entangled quantum states through the determination of instantaneous eigenvectors using matrix formalism [26].

In conclusion, the results obtained by studying the time-dependent topological effects on dipoles, in accordance with the characteristics of AB-type effects, support the complex interactions of the systems considered in the 2+1 dimension and the experimental and theoretical investigation of these interactions.

## Acknowledgements

The author would like to thank late Prof. Namık Kemal Pak for continuing to enlighten his path, Prof. Müge Boz for her invaluable contributions and precious efforts, and dear Ian Bouche for his contribution to the preliminary work.

## References

- [1] Y. Aharonov, D. Bohm, Significance of electromagnetic potentials in the quantum theory, *Phys. Rev.* 115 (1959) 485–491. doi:10.1103/PhysRev.115.485.
- [2] R. G. Chambers, Shift of an electron interference pattern by enclosed magnetic flux, *Phys. Rev. Lett.* 5 (1960) 3–5. doi:10.1103/PhysRevLett.5.3.
- [3] A. Tonomura, N. Osakabe, T. Matsuda, T. Kawasaki, J. Endo, S. Yano, H. Yamada, Evidence for Aharonov-Bohm effect with magnetic field completely shielded from electron wave, *Phys. Rev. Lett.* 56 (1986) 792–795. doi:10.1103/PhysRevLett.56.792.
- [4] Y. Aharonov, A. Casher, Topological quantum effects for neutral particles, *Phys. Rev. Lett.* 53 (1984) 319–321. doi:10.1103/PhysRevLett.53.319.
- [5] A. Cimmino, G. I. Opat, A. G. Klein, H. Kaiser, S. A. Werner, M. Arif, R. Clothier, Observation of the topological Aharonov-Casher phase shift by neutron interferometry, *Phys. Rev. Lett.* 63 (1989) 380–383. doi:10.1103/PhysRevLett.63.380.
- [6] X.-G. He, B. H. J. McKellar, Topological phase due to electric dipole moment and magnetic monopole interaction, *Phys. Rev. A* 47 (1993) 3424–3425. doi:10.1103/PhysRevA.47.3424.
- [7] M. Wilkens, Quantum phase of a moving dipole, *Phys. Rev. Lett.* 72 (1994) 5–8. doi:10.1103/PhysRevLett.72.5.
- [8] J. Gillot, S. Lepoutre, A. Gauguier, M. Büchner, J. Vigué, Measurement of the He-McKellar-Wilkens topological phase by atom interferometry and test of its independence with atom velocity, *Phys. Rev. Lett.* 111 (2013) 030401. doi:10.1103/PhysRevLett.111.030401.
- [9] J. P. Dowling, C. P. Williams, J. D. Franson, Maxwell duality, lorentz invariance, and topological phase, *Phys. Rev. Lett.* 83 (1999) 2486–2489. doi:10.1103/PhysRevLett.83.2486.
- [10] H. O. Cildiroglu, A. U. Yilmazer, Topological effects of Aharonov-Bohm type and the role of the quantum entanglement, *AIP Conference Proceedings* 2042 (1) (2018) 020045. doi:10.1063/1.5078917.
- [11] E. C. V. Douglas Singleton, The covariant, time-dependent aharonov-bohm effect, *Phys. Lett. B* 723 (2013) 241–244. doi:10.1016/j.physletb.2013.05.014. URL <https://doi.org/10.1016/j.physletb.2013.05.014>.
- [12] M. Bright, D. Singleton, Time-dependent non-abelian aharonov-bohm effect, *Phys. Rev. D* 91 (2015) 085010. doi:10.1103/PhysRevD.91.085010.
- [13] M. Ababekri, A. Anwar, M. Hekim, R. Rashidin, Aharonov-Bohm phase for an electric dipole on a noncommutative space, *Frontiers in Physics* 4 (2016) 22. doi:https://doi.org/10.3389/fphy.2016.00022.
- [14] J. U. Douglas Singleton, The time-dependent aharonov-casher effect, *Phys. Lett. B* 753 (2016) 91–96. doi:10.1016/j.physletb.2015.12.002. URL <https://doi.org/10.1016/j.physletb.2015.12.002>.
- [15] J. Jing, K. Wang, Z.-W. Long, S.-H. Dong, On the time-dependent Aharonov-Bohm effect, *Phys. Lett. B* 774 (2017) 87–90. doi:10.1016/j.physletb.2017.09.041. URL <https://doi.org/10.1016/j.physletb.2017.09.041>.
- [16] K. Ma, J.-H. Wang, Probing the noncommutative effects of phase space in the time-dependent Aharonov-Bohm effect, *Annals of Physics* 383 (2017) 120–129. doi:10.1016/j.aop.2017.05.005. URL <https://doi.org/10.1016/j.aop.2017.05.005>.
- [17] S. R. Choudhury, S. Mahajan, Direct calculation of time varying Aharonov-Bohm effect, *Physics Letters A* 383 (21) (2019) 2467–2471. doi:https://doi.org/10.1016/j.physleta.2019.05.008.
- [18] J. Jing, L.-B. Kong, Q. Wang, S.-H. Dong, On the noncommutative aharonov-bohm effects, *Physics Letters B* 808 (135660) (2020). doi:10.1016/j.physletb.2020.135660. URL <https://doi.org/10.1016/j.physletb.2020.135660>.
- [19] T. Wang, K. Ma, Time-dependent Aharonov-Casher effect on noncommutative space, *Communications in Theoretical Physics* 75 (1) (2022) 015203.
- [20] T. Wang, K. Ma, Time-dependent He-McKellar-Wilkens effect on noncommutative space, *International Journal of Modern Physics A* 38 (2) (2023) 2350011. doi:10.1142/S0217751X23500112.
- [21] P. L. Saldanha, Electrodynamical Aharonov-Bohm effect, *Phys. Rev. A* 108 (2023) 062218. doi:10.1103/PhysRevA.108.062218. URL <https://link.aps.org/doi/10.1103/PhysRevA.108.062218>.
- [22] M. Wakamatsu, Is the Aharonov-Bohm phase shift for a non-closed path a measurable quantity?, *The European Physical Journal Plus* 139 (2) (2024) 1–19. doi:https://doi.org/10.1140/epjp/s13360-024-04922-9.
- [23] H. O. Cildiroglu, A. U. Yilmazer, Dual scalar Aharonov-Bohm phase, *AIP Conference Proceedings* 2178 (1) (2019) 030047. doi:10.1063/1.5135445.
- [24] M. V. Berry, Quantal phase factors accompanying adiabatic changes, *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences* 392 (1984) 45–57. doi:10.1098/rspa.1984.0023.
- [25] R. Mignani, Aharonov-Casher effect and geometrical phases, *Journal of Physics A: Mathematical and General* 24 (8) (1991) L421. doi:10.1088/0305-4470/24/8/006.
- [26] H. Cildiroglu, A. Yilmazer, Investigation of the Aharonov-Bohm and Aharonov-Casher topological phases for quantum entangled states, *Physics Letters A* 420 (2021) 127753. doi:https://doi.org/10.1016/j.physleta.2021.127753.