# RKHS, ODZIJEWICZ, BEREZIN AND FEDOSOV-TYPE QUANTIZATIONS ON SMOOTH COMPACT MANIFOLDS

RUKMINI DEY

I.C.T.S.-T.I.F.R., Bangalore, India.

Key words: Berezin quantization, coherent states, reproducing kernel MSC: 81-XX, 53-XX, 43-XX, 46-XX, 47-XX.

ABSTRACT. In this article we define Odzijewicz, Berezin and Fedosov-type quantization on compact smooth manifolds. The method is as follows. We embed the smooth manifold of real dimension n into  $\mathbb{C}P^n$  (and in the Fedosov quantization case embed into any real 2n dimensional symplectic manifold). The pullback coherent states are defined in the usuall way. In the Odzijewicztype, Berezin-type quantization the Hilbert space of geometric quantization of  $\mathbb{C}P^n$ . In the Berezin case, the operators that are quantized are those induced from the ambient space. The Berezin-type quantization exhibited here is a generalization of an earlier work of the author and Kohinoor Ghosh (where we had needed totally real embedding). The Fedosov-type quantization is carried out by restriction to the submanifold given by the embedding.

#### 1. Introduction

Some quantum systems donot come from quantizing classical systems (which are expected to have a symplectic structure) but there is a semi-classical limit of the quantum system, spin being such a system ([14]). We wish to include systems which donot have symplectic structure (or group action) and study if they are a semi-classical limit of some quantum system as  $\hbar$  goes to zero. A Poisson structure is needed but it is induced from another Poisson manifold where the manifold is embedded, in this case  $\mathbb{C}P^n$  or  $\mathbb{C}^n$ . The quantization is induced from  $\mathbb{C}P^n$  or  $\mathbb{C}^n$  (depending on whether we expect a finite or an infinite dimensional Hilbert space). In this paper we will concentrate on embeddings into  $\mathbb{C}P^n$ .

The other motivation of the work is that sometimes the Hilbert space of the problem turns out to be different from what the actual parameter space should prescribe. The Hilbert space could be just be obtained from geometric quantization of  $\mathbb{C}^n$ , or  $\mathbb{C}P^n$ . Roughly speaking in these two cases the Hilbert space correspond to multinomails. In some situations this could be at least a good approximation, for example the Quantum Hall Effect (where polynomials suffice for lowest Landau levels [16]).

In this paper we show Odzijewicz, Berezin and Fedosov-type quantizations on smooth compact manifolds. The first two use coherent states in a very essential way. The literature on coherent states is vast, see for instance a review by Ali,

#### RUKMINI DEY

Gazeau, Antoine and Mueller [1], Gazeau and Monceau [9], proceedings by editors Strasburger, Ali, Antoine, Gazeau, Odzijewicz [2]. After Berezin's orginal work [3], Berezin quantization has been generalized to many domains and manifolds, see Englis [7] for an example.

We embed a compact smooth manifold into  $\mathbb{C}P^n$  (using Whitney embedding or any other embedding) and pull back coherent states from  $\mathbb{C}P^n$  [4]. Role of  $\mathbb{C}P^n$  or  $\mathbb{C}^n$  can taken by other appropriate manifolds too. For instance, we can embed the compact smooth manifold in a symplectic manifold of twice the dimension which has a reproducing kernel Hilbert space (namely coherent states).

In [11] Odzijewicz studied the quantization of a Kähler manifold using coherent state embedding and showed that the calculation of the Feynman path integral is equivalent to finding the reproducing kernel function. We generalize this to symplectic manifolds and even to smooth compact manifolds, as we can now talk of reproducing kernel Hilbert spaces pulled back from  $\mathbb{C}P^n$ .

In [3] Berezin gave a way of defining a star product on the symbol of bounded linear operators acting on a Hilbert space (with a reproducing kernel) on a Kähler manifold under certain conditions. There is a parameter in the theory (namely  $\hbar$ ) such that in the limit  $\hbar \to 0$  the star product tends to usual product and the commutator of the star product is proportional to the Poisson bracket upto first order. This is called the correspondence principle.

In this paper we embed a compact smooth manifold into  $\mathbb{C}P^n$  and pull back the reproducing kernel Hilbert space. Pullback coherent states give symbols of bounded linear operators induced from those corresponding to  $\mathbb{C}P^n$  and it is trivial to see that they satisfy the correspondence principle.

In this context we recall that in [4] Dey and Ghosh had considered pull back coherent states and totally real submanifolds of  $\mathbb{C}P^n$  and defined pull back operators and their  $\mathbb{C}P^n$ -symbols and showed that they satisfied the correspondence principle. In other words we had shown Berezin-type quantization of certain operators for manifolds which can be embedded in  $\mathbb{C}P^n$  as totally real submanifolds. This was part of Ghosh's thesis [10]. Our present work is a generalization of this, as we donot need the condition totally real submanifolds. The condition of totally real submanifolds can give topological obstructions. We have no conditions on the smooth compact manifold.

Dey and Ghosh had also defined in [5] a Berezin-type quantization on even dimensional compact manifolds (of real dimension 2d) by removing a set of measure zero and embedding it into  $\mathbb{C}P^d$ . This was part of Ghosh's thesis [10]. In this article our approach is slightly different. We embed a d real dimensional manifold into  $\mathbb{C}P^d$ .

In [8] Fedosov constructs a deformation quantization on symplectic manifolds. We embed a smooth compact manifold of real dimension d into any symplectic manifold of real dimension 2d by the Whitney embedding and induce the Fedosov quantization on the submanifold. This is done simply by restriction to the embedded submanifold.

All the quantizations depend on the embedding.

## 2. Review of Geometric Quantization and Coherent states on $\mathbb{C}P^n$

This review follows [3] and [10] closely.

Let  $U_0 \subset \mathbb{C}P^n$  given by  $U_0 = \{\mu_0 \neq 0\}$  where  $[\mu_0, ..., \mu_n]$  are homogeneous coordinates on  $\mathbb{C}P^n$ . Let  $(\mu_1, \mu_2, \dots, \mu_n)$  be inhomogenous coordinates on  $U_0 \equiv \mathbb{C}^n$ such that  $[1, \mu_1, \mu_2, ..., \mu_n] \in U_0$ .

The Fubini-Study form is given by  $\Omega_{FS} = \sum_{i,j=1}^{n} \Omega_{ij}^{FS} d\mu_i \wedge d\bar{\mu}_j$ , where the Kähler metric G and the Kähler form  $\Omega_{FS}$  are related by  $\Omega_{FS}(X,Y) = G(IX,Y)$ .

The Poisson bracket of two functions t and s:

 $\{t,s\}_{FS} = \sum_{i,j=1}^{n} \Omega_{FS}^{ij} \left( \frac{\partial t}{\partial \bar{\mu}_j} \frac{\partial s}{\partial \mu_i} - \frac{\partial s}{\partial \bar{\mu}_i} \frac{\partial t}{\partial \mu_j} \right)$  where  $(\Omega_{FS}^{ij})$  are the matrix coefficients of the inverse of the matrix  $(\Omega_{ij}^{FS})$  of the Fubini-Study form.

Let  $H^{\otimes m}$  be the *m*-th tensor product of the hyperplane section bundle H on  $CP^n$ . Recall that  $m\Omega_{FS}$  is its curvature form and  $m\Phi_{FS}$  is a local Kähler potential where  $e^{m\Phi_{FS}(\mu,\bar{\mu})} = (1+|\mu|^2)^m$ . Let  $\{\phi_i\}_{i=1}^N$  be an orthonormal basis for the space of holomorphic sections.

On  $U_0$  the sections of  $H^{\otimes m}$  are functions since the bundle is trivial when restricted to  $U_0$ . They can be identified with polynomials in  $\{\mu_i\}_{i=1}^n$  of degree at most m.

Let  $\hbar = \frac{1}{m}$  be a parameter. Then  $\{\phi_i\}$  depend on  $\hbar$ .

We define  $dV(\mu) = |\Omega_{FS}^n(\mu)|_{U_0}| = \mathcal{G}(\mu)\Pi_{i=1}^n |d\mu_i \wedge d\bar{\mu}_i| = \mathcal{G}(\mu)|d\mu \wedge d\bar{\mu}| =$  $\frac{|d\mu\wedge d\bar{\mu}|}{(1+|\mu|^2)^{n+1}} \text{ to be a volume form on } \mathbb{C}^n, \text{ where } \mathcal{G} = \det[g^{ij}|_{U_0}].$ 

Then  $V = \int_{\mathbb{C}^n} dV = \int_{\mathbb{C}^n} \frac{|d\mu \wedge d\bar{\mu}|}{(1+|\mu|^2)^{n+1}} < \infty.$ Let  $(c(m))^{-1} = \int_{U_0} \frac{1}{(1+|\nu|^2)^m} dV(\nu) = \int_{U_0} e^{-m\Phi_{FS}(\nu,\bar{\nu})} dV(\nu)$ Let an innerproduct on the space of functions on  $U_0$  be defined as

$$\langle f,g \rangle = c(m) \int_{U_0} \frac{\overline{f(\nu)}g(\nu)}{(1+|\nu|^2)^m} dV(\nu) = c(m) \int_{U_0} \overline{f(\nu)}g(\nu)e^{-m\Phi_{FS}(\nu,\bar{\nu})} dV(\nu).$$

Also,  $D_{(q_1,q_2,...,q_n;q)} = c(m) \int_{U_0} \frac{|\nu_1|^{2q_1}...|\nu_n|^{2q_n}}{(1+|\nu|^2)^m} dV(\nu)$ , where  $q'_i s$  are all possible positive integers such that  $q_1 + ... + q_n = q; q = 0, ..., m$ . Let  $\Phi_{(q_1,q_2,...,q_n;q)}(\mu) = \frac{1}{\sqrt{D_{(q_1,...,q_n;q)}}} \mu_1^{q_1}...\mu_n^{q_n}$  where  $q_1 + ... + q_n = q; q = 0, ..., m$ .

It is easy to check that  $\{\Phi_{(q_1,\ldots,q_n;q)}\}$  are orthonormal in  $\mathbb{C}^n$  with respect to the inner product defined as above and are restriction of a basis for sections of  $H^{\otimes m}$ to  $U_0$ . The span of these form a Hilbert space with the above inner product.

**Definition:** The Rawnsley-type coherent states [14], [15] are given on  $U_0$  by  $\psi_{\mu}$ reading as follows:

$$\psi_{\mu}(\nu) := \sum_{q_1+q_2+\ldots+q_n=q; q=0,1,\ldots,m} \overline{\Phi_{(q_1,q_2,\ldots,q_n;q)}(\mu)} \Phi_{(q_1,q_2,\ldots,q_n;q)}(\nu).$$

In short hand notation  $\psi_{\mu} := \sum_{I} \Phi_{I}(\mu) \Phi_{I}$  where the multi-index  $I = (q_{1}, ..., q_{n}; q)$ runs over the set  $q_1 + ... + q_n = q; q = 0, ..., m$ .

This is a reproducing kernel in the sense below.

**Proposition 2.1.** Reproducing kernel property. If  $\Psi$  is any other section, then  $\langle \psi_{\mu}, \Psi \rangle = \Psi(\mu)$ . In particular,  $\langle \psi_{\mu}, \psi_{\nu} \rangle = \psi_{\nu}(\mu)$ .

Proposition 2.2. Resolution of identity property:

$$c(m)\int_{U_0} \langle \Psi_1, \psi_\mu \rangle \langle \psi_\mu, \Psi_2 \rangle e^{-m\Phi_{FS}(\mu,\bar{\mu})} dV(\mu) = \langle \Psi_1, \Psi_2 \rangle.$$

In particular.

$$c(m) \int_{U_0} \langle \psi_{\nu}, \psi_{\mu} \rangle \langle \psi_{\mu}, \psi_{\nu} \rangle e^{-m\Phi_{FS}(\mu,\bar{\mu})} dV(\mu) = \langle \psi_{\nu}, \psi_{\nu} \rangle.$$

The proofs of these are easy and can be found for instance in [5]. It is in general true of Rawnsley type coherent states.

# 3. A reproducing kernel Hilbert space on a compact smooth manifolds and coherent states

In this section we construct a reproducing kernel Hilbert space and coherent states on a compact smooth manifolds by emedding it into  $\mathbb{C}P^d$ . This generalizes a result in [4]. We proceed similar to [4], but we do not need the "totally real" condition. We use the Whitney embedding of any compact smooth manifold. Any other smooth embedding will also work.

Let  $M^d$  be a compact smooth manifold of real dimension d. Let  $\epsilon: M \to \mathbb{R}^{2d}$ be any embedding (for instance Whitney embedding). Let  $i: \mathbb{R}^{2d} \to \mathbb{C}P^d$  be the inclusion such that  $R^{2d}$  is identified with  $U_0 \subset \mathbb{C}P^d$  and  $\chi = i \circ \epsilon$ . It is clear that  $\chi: M \to \mathbb{C}P^d$  is an embedding and that  $\epsilon: M \to \epsilon(M)$  is a diffeomorphism. Let  $\Sigma = \epsilon(M).$ 

Let  $\mathcal{H}_m$  be the sections of  $H^{\otimes m}$  with norm denoted for short as  $||s||_{\mathbb{C}P^d}$ . Let  $\Psi_{(q_1,q_2,...,q_n;q)}(\mu) = \frac{1}{\sqrt{D_{(q_1,...,q_n;q)}}} \mu_1^{q_1} ... \mu_n^{q_n}$  where  $q_1 + ... + q_n = q; q = 0, ..., m$  be an orthonormal basis for it as mentioned in the previous section.

Let  $\mathcal{H}_{1m} = i^*(\mathcal{H}_m)$  be the pullback Hilbert space on  $\Sigma = \epsilon(M)$ . The norm on

 $\mathcal{H}_{1m}$  is given by  $||\tilde{s}||_{\Sigma} = \min_{s \in \mathcal{H}_m} \{ ||s||_{\mathbb{CP}^d} : \tilde{s} = s \circ i \}.$ Let  $\mathcal{H}_{2m} = \chi^*(\mathcal{H}_m)$  be the pullback Hilbert space on M. Thus if  $\tilde{s} \in \mathcal{H}_{2m}$ , it is of the form  $\tilde{s} = s \circ \chi$ . The norm on  $\mathcal{H}_{2m}$  is given by  $||\tilde{s}||_M = \min_{s \in \mathcal{H}_m} \{||s||_{\mathbb{CP}^d}$ :  $\tilde{\mathbf{s}} = \mathbf{s} \circ \chi \}.$ 

Let  $\epsilon^{-1}: \Sigma \to M$  be the inverse of  $\epsilon$  on  $\Sigma$ .

One can show that  $||\tilde{s}||_M = ||\epsilon^{-1*}\tilde{s}||_{\Sigma}$ . This is because

$$\begin{split} \tilde{s}||_{M} &= \min_{\mathbf{s} \in \mathcal{H}_{m}} \{ ||\mathbf{s}||_{\mathbb{C}\mathbf{P}^{d}} : \tilde{\mathbf{s}} = \mathbf{s} \circ \epsilon \} \\ &= \{ ||s||_{\mathbb{C}P^{d}} : s = \tilde{s} \circ \epsilon^{-1} \} \end{split}$$

**Proposition 3.1.**  $\mathcal{H}_{1m}$  and  $\mathcal{H}_{2m}$  are Hilbert spaces in the respective norms.

*Proof.* For proof see for instance [12].

Let  $\eta_I$  be an orthonormal basis for  $\mathcal{H}_{2m}$ . **Definition:** 

The Rawnsley-type coherent states on M are defined locally as  $\psi_p = \sum_{k=1}^{l} \overline{\eta_{I_k}(p)} \eta_{I_k}$ where  $p \in M$ .

As before they are overcomplete, have reproducing kernel property, resolution of identity property.

### 4. Odzijewicz-type quantization on compact smooth manifolds

4.1. Review of Odzijewicz quantization on  $\mathbb{C}P^n$ . The Odzijewicz quantization goes through verbatim as in his paper [11] when one takes the Kähler manifold to be  $\mathbb{C}P^n$ .

Take  $\mathcal{M}$  to be holomorphic sections of  $H^{\otimes m}$  where H is the hyperplane section bundle and m is a large positive integer.

In this case  $\mathcal{M}$  is sufficiently ample and the coherent state embedding is given by

 $\mu \in \mathbb{C}P^n \mapsto \psi_\mu \in \mathbb{C}P(\mathcal{M}).$ 

We will henceforth use Odzijewicz's notation – and write  $\psi_{\mu}(\cdot) = K_0(\bar{\mu}, \cdot)$  on  $U_0$ . Let  $[1, w_1, ..., w_n]$  be coordinates on  $U_0 \subset \mathbb{C}P^n$ .

We need to solve the Monge-Ampere equation (2.22) in [11] given by (when  $\frac{(-i)^n \omega}{K}$  is positive)

$$det[\frac{\partial^2 log\rho(w)}{\partial w_j \partial \bar{w}_k}] = C(-1)^{\frac{n(n+1)}{2}} \frac{1}{n!} \rho(w) K_0(\bar{w}, w),$$

Here C > 0 when  $\frac{(-i)^n \omega}{K_0}$  is positive and C < 0 when  $\frac{(-i)^n \omega}{K_{00}}$  is positive. We take  $\rho(w) = \frac{1}{(1+|w|^2)^N} = \exp(-N(\log(1+|w|^2)))$ . We know from previous section  $K(\bar{w}, w) = (1+|w|^2)^m$ .

Left hand side of the Monger-Ampere equation is  $\frac{-N}{(1+|w|^2)^{n+1}}$ .

$$\frac{-N}{(1+|w|^2)^{n+1}} = C(-1)^{n(n+1)/2} \frac{1}{n!} \frac{1}{(1+|w|^2)^{N-m}}.$$

Then N = n + m + 1 and  $C = \pm Nn!$  depending on whether  $(-1)^{\frac{n(n+1)}{2}+1}$  is positive or negative (i.e.  $\frac{(-i)^n \omega}{K}$  is positive or negative). One can thus have a coherent state embedding  $w \in \mathbb{C}P^n \mapsto K(\bar{w}, \cdot) \in \mathbb{C}P(\mathcal{M})$ 

One can thus have a coherent state embedding  $w \in \mathbb{C}P^n \mapsto K(\bar{w}, \cdot) \in \mathbb{C}P(\mathcal{M})$ One can define path integral as in (2.28) [11] and show that it is related to the pull back metric on  $\mathbb{C}P^n$  of the Fubini-Study metric on  $\mathbb{C}P(\mathcal{M})$ .

We have from (2.30)

$$d_{\mathbb{C}P^n}(z_1, z_2) = \sqrt{2}(1 - |a_{0\beta}(z_1, z_2)|)^{\frac{1}{2}}.$$

where by (2.20)  $a_{0\beta}(\bar{v}, w) = \frac{K_{0\beta}(\bar{v}, w)}{K_{0\beta}(\bar{v}, v)^{\frac{1}{2}}K_{0\beta}(\bar{w}, w)^{\frac{1}{2}}}$ , where  $v \in U_0, w \in U_\beta, \beta = 0, ..., n$ . Note: Equation (2.29) in [11] originates from the pullback of the Fubini-Study

Note: Equation (2.29) in [11] originates from the pullback of the Fubini-Study metric on  $\mathbb{C}P(\mathcal{M})$  to  $\mathcal{M}$ .

4.2. Induced Odzijewicz-type quantization on compact smooth manifolds. Let M be a compact smooth manfield of real dimension d. Let n = dand  $U_0$  be the open subset of  $\mathbb{C}P^n$  as mentioned in section 2. We know as a topological space  $U_0 \equiv \mathbb{R}^{2d}$ . Let  $\epsilon : M \mapsto U_0 \subset \mathbb{C}P^n$  be an embedding (for example Whitney emebdding).

Let  $p, q \in M$  and  $v = \epsilon(p)$  and  $z = \epsilon(q)$ . Then  $v, z \in U_0 \subset \mathbb{C}P^n$ .

We can define the transition probability between p, q on M to be

$$A(p,q) \equiv a_{00}(\bar{v},z) = \frac{K_{00}(\bar{v},z)}{K_{00}(\bar{z},z)^{1/2}K_{00}(\bar{v},v))^{1/2}}.$$

where  $a_{00}$  is the transition amplitude of points in  $U_0$ .

Once again it is easy to show that

$$d_M(p,q) = d_{\mathbb{C}P^n}(z_1, z_2) = \sqrt{2}(1 - |a_{00}(z_1, z_2)|)^{\frac{1}{2}},$$

where  $z_1 = \epsilon(p), z_2 = \epsilon(q)$ .

#### RUKMINI DEY

#### 5. Induced Berezin-type quantization on compact smooth manifolds

Let M be a compact smooth manifold.

Let  $\chi: M \mapsto \mathbb{C}P^n$  as in section 3.

 $\mathbf{6}$ 

Let us continue on  $\mathbb{C}P^n$  and recall the Berezin quantization on it.

**Notation:** As in [3], we denote by  $\mathcal{L}_m(\mu, \bar{\mu}) = \langle \psi_\mu, \psi_\mu \rangle = \psi_\mu(\mu), \mathcal{L}_m(\mu, \bar{\nu}) =$  $\langle \psi_{\mu}, \psi_{\nu} \rangle = \psi_{\nu}(\mu).$ 

Let  $\hat{A}$  be a bounded linear operator acting on  $\mathcal{H}$ . Then, as in [3], one can define a symbol of the operator as

$$A(\nu,\bar{\mu}) = \frac{\left\langle \psi_{\nu}, \hat{A}\psi_{\mu}, \right\rangle}{\left\langle \psi_{\nu}, \psi_{\mu} \right\rangle}.$$

One can show that one can recover the operator from the symbol by a formula [3].

Let  $\hat{A}_1, \hat{A}_2$  be two such operators and let  $\hat{A}_1 \circ \hat{A}_2$  be their composition.

Then the symbol of  $\hat{A}_1 \circ \hat{A}_2$  will be given by the star product defined as in [3]:

$$(A_{1} * A_{2})(\mu, \bar{\mu}) = c(m) \int_{U_{0}} A_{1}(\mu, \bar{\nu}) A_{2}(\nu, \bar{\mu}) \frac{\mathcal{L}_{m}(\mu, \bar{\nu})\mathcal{L}_{m}(\nu, \bar{\mu})}{\mathcal{L}_{m}(\mu, \bar{\mu})\mathcal{L}_{m}(\nu, \bar{\nu})} \mathcal{L}_{m}(\nu, \bar{\nu}) e^{-m\tilde{\Phi}(\nu, \bar{\nu})} dV(\nu),$$

where recall  $\frac{1}{c(m)} = \int_{U_0} e^{-m\Phi_{FS}(\nu,\bar{\nu})} dV(\nu).$ 

This is the symbol of  $\hat{A}_1 \circ \hat{A}_2$ .

One can show the following is the reproducing kernel [5], [10].

**Proposition 5.1.**  $\psi_{\mu}(\nu) = (1 + \overline{\mu} \cdot \nu)^m$ .

#### **Theorem 5.2** (Berezin). Let $\mu \in \mathbb{C}^n$ .

The star product satisfies the correspondence principle:

- 1.  $\lim_{m\to\infty} (A_1 \star A_2)(\mu, \bar{\mu}) = A_1(\mu, \bar{\mu})A_2(\mu, \bar{\mu}),$
- 2.  $\lim_{m \to \infty} m(A_1 \star A_2 A_2 \star A_1)(\mu, \bar{\mu}) = i\{A_1, A_2\}_{FS}(\mu, \bar{\mu}).$

See [3], [5] for proof.

5.1. Induced operators and correspondence principle. Let A be a bounded linear operator from  $\mathcal{H}_m$  to itself. Now we turn to M.

Let  $\hat{B} = \chi^*(\hat{A})$  be a bounded linear operator from  $\mathcal{H}_{2m} = \chi^*(\mathcal{H}_m)$  to itself. It is defined by  $\chi^*(\hat{A})(\chi^*(s)) = \chi^*(\hat{A}s)$ . Given  $\hat{B}, \hat{A}$  is not unique. Suppose we have  $\hat{B} = \chi^*(\hat{A}_1) = \chi^*(\hat{A}_2)$ . Let  $\hat{A}$  be the one of lowest norm i.e. if  $\hat{B} = \chi^*(\hat{A}) = \chi^*(\hat{A}_1)$ , then  $||\hat{A}_1|| \ge ||\hat{A}||$ .

**Definition:** Let  $B: M \times M \to \mathbb{C}$  be the symbol of  $\hat{B}$  in the coherent states  $\psi_p$ , i.e.  $B(p,p) = \frac{\langle \psi_p, \hat{B}(\psi_p) \rangle_M}{\langle \psi_p, \psi_p \rangle_M}$  and  $B(p,q) = \frac{\langle \psi_p, \hat{B}(\psi_q) \rangle_M}{\langle \psi_p, \psi_q \rangle_M}$ Here the norm is defined as in section 3.

**Proposition 5.3.**  $B(p,q) = \frac{\langle s_{\chi(p)}, \hat{A}(s_{\chi(q)}) \rangle_{CP^d}}{\langle s_{\chi(p)}, s_{\chi(q)} \rangle_{CP^d}} = A(\chi(p), \overline{\chi(q)}), \text{ where } \hat{B} = \chi^*(\hat{A})$ and  $\hat{A}$  is the one of lowest norm over all  $\hat{C}$  such that  $\hat{B} = \chi^*(\hat{C}).$ 

*Proof.*  $\psi_p = \chi^*(s_{\chi(p)})$ , where  $s_{\chi(p)} = \sum_{k=1}^l \overline{\Phi_{I_k}(\chi(p))} \Phi_{I_k}$  is the coherent state in  $\mathcal{H}_m$ . This is because  $\eta_{I_k} = \chi^*(\Phi_{I_k})$ .

$$B(p,q) = \frac{\hat{B}(\psi_q)(p)}{\psi_q(p)} = \frac{\hat{A}(s_{\chi(q)})(\chi(p))}{s_{\chi(q)}(\chi(p))} = \frac{\langle s_{\chi(p)}, \hat{A}(s_{\chi(q)}) \rangle_{\mathbb{C}P^d}}{\langle s_{\chi(p)}, s_{\chi(q)} \rangle_{\mathbb{C}P^d}}.$$

### **Definition:**

Let  $B_1$  and  $B_2$  be symbols of  $\hat{B}_1$  and  $\hat{B}_2$  bounded linear operators on  $\mathcal{H}_{2m}$ . Then  $B_1 * B_2$  is the symbol of  $\hat{B}_1 \circ \hat{B}_2$ , [3].

One sees that  $\hat{B}_1 \circ \hat{B}_2 = \chi^*(\hat{A}_1 \circ \hat{A}_2)$ , where we can take  $\hat{A}_1 \circ \hat{A}_2$  is the one of the least norm.

Then,  $B_1 * B_2 = A_1 * A_2$  where the second one is  $\mathbb{C}P^d$  star product. Let  $\{B_1, B_2\}_{FS}(p, p) = \{A_1, A_2\}_{FS}(\chi(p), \overline{\chi(p)}).$ 

**Theorem 5.4.** The star product on the symbol of bounded linear operators on  $\mathcal{H}_{2m}$  satisfies the correspondence principle:

(1)  $\lim_{m\to\infty} (B_1 * B_1)(p,p) = B_1(p,p)B_2(p,p).$ 

(2)  $\lim_{m\to\infty} m(B_1 * B_2 - B_2 * B_1)(p,p) = i\{B_1, B_2\}_{FS}(p,p).$ 

*Proof.* This follows from the fact that  $A_1 * A_2$  satisfy the correspondence principle, see Theorem 5.2 (Berezin).

#### 6. Induced Fedosov-type quantization on compact smooth manifolds

Let M be a compact smooth manifold of real dimension d. Let  $\epsilon_1 : M \mapsto \mathbb{R}^{2d}$ is an embedding (say, Whitney embedding). Let N be a symplectic manifold of dimension 2d such that  $N \setminus N_0 \equiv \mathbb{R}^{2d}$  where  $N_0$  is a lower dimensional skeleton [6]. Thus there is an embedding  $\epsilon : M \mapsto N \setminus N_0$ .

Let  $X = \epsilon(M)$ .

In this section let us recall the Fedosov deformation quantization [8] on N and induce the same on X.

Let  $x, y \in X$ . Since  $\epsilon$  is an embedding there exists a special chart V in N such that if  $x = (x_1, ..., x_{2d})$  then in  $V \cap X$ ,  $x_{d+1} = x_{d+2} = ... = x_{2d} = 0$ . Similarly, let  $y = (y_1, ..., y_{2d})$  and there is a special chart W in N such that in  $W \cap X$ ,  $y_{d+1} = y_{d+2} = ... = y_{2d} = 0$ .

Thus on  $X = \epsilon(M)$  the local expression for  $a \in W \cap X$  in [8] (2.4) is

$$a = \sum h^k a_{k,i_1\dots i_p,j_1\dots j_q(x)} y^{i_1}\dots y^{i_p} dx^{j_1} \wedge \dots \wedge dx^{j_q}$$

where  $i_1, ..., i_p \in \{1, ..., d\}$  and  $j_1, ..., j_p \in \{1, ..., d\}$ . Let  $b \in W \cap X$ . Then  $a \circ b$  is the composition defined on X. Since definitions of  $\delta$  and  $\delta^*$  is local, one can define it on X. We also have  $\delta^2 = \delta^{*2} = 0$  and for monomials  $\delta\delta^* + \delta^*\delta = (p+q)$ id on X as in Lemma (2.2). Once again since the definitions are local, we can define on X,  $\delta^{-1} = \frac{\delta^*}{p+q}$  for p+q > 0 and  $\delta^{-1} = 0$  for p+q = 0 and  $a = \delta\delta^{-1}a + \delta^{-1}\delta a + a_{00}$ . All definitions and notations are the same as in [8].

Let  $\omega$  be the symplectic form on N. The symplectic form restricts to a 2-form  $\tau = \omega|_X$  on X. A symplectic connection  $\partial$  is defined in Definition (2.3) in [8]. In a Darboux chart  $U \subset N$  one has  $\Gamma_{ijk} = \omega_{im} \Gamma_{jk}^m$  where  $\Gamma_{jk}^m$  are Christofel symbols and  $\Gamma_{ijk}$  are completely symmetric with respect to the indices i, j, k. Even though  $\tau$  maynot be symplectic we can define the restriction of  $\partial$  to X. On the special chart V in a Darboux neighbourhood of  $x \in X$ , i.e. on  $U \cap V$ , where  $x_{d+1} = \ldots = x_{2d} = 0$ , we can define as in [8] (2.9),  $\partial a = dx^i \wedge \partial_i a$  where  $i = 1, \ldots, d$ . (2.10) also holds at  $x \in U \cap V \subset X$ . One can restrict  $\gamma$  to X and define Da accordingly. Lemma (2.4)

#### RUKMINI DEY

is local and holds verbatim.  $\Omega$  also restricts to X and lemma (2.6) holds since the proof is local. Construction of the abelian connection carries through.

In [8] Fedosov has defined  $W_D \subset W$  constrained of flat connections Da = 0.

We can show that non-trivial flat sections exists on X i.e.  $W_D$  restricted to X is not empty. This is because Fedosov iteratively constructs the section  $a \in W_D$  by its symbol  $a_0 = \sigma(a)$ :

$$a = a_0 + \partial_i a_0 y^i + \frac{1}{2} \partial_i \partial_j a_0 y^i y^j + \frac{1}{6} \partial_i \partial_j \partial_k a_0 y^i y^j y^k - \frac{1}{24} R_{ijkl} \omega^{lm} \partial_m a_0 y^i y^j y^k + \dots,$$

and each of there terms restrict to X. One can take  $a_0$  to be non identically zero on X. So  $W_D$  is non-empty.

Then we can show an equivalent theorem to theorem (3.3) on X since the proof is local. Thus  $\sigma: W_D \mapsto Z$  is invertible on X as well.

On X we can define for  $a, b \in Z$   $a * b = \sigma(\sigma^{-1}(a) \circ \sigma^{-1}(b))$ .

This will satisfy all conditions (i) - (iii) formulated in the introduction of [8] with all quantities restricted to X. For instance (iii) reads as

$$[a,b] \equiv a * b - b * a = -ih\{a_0, b_0\}|_X + \dots$$

#### 7. Acknowledgement

Rukmini Dey acknowledges support from the project RTI4001, Department of Atomic Energy, Government of India.

#### References

- S.T. Ali, J.-P. Antoine J. P. Gazeau and U.A. Mueller: Coherent States and their Generalizations: a mathematical overview, *Reviews in Mathematical Physics*, 7, 1013-1104 (1995).
- [2] Quantization, Coherent States and Poisson Structures, Proceedings of the XIV Workshop on Geometric Methods in Physics, A. Strasburger, S.T. Ali, J.-P. Antoine and A. Odzijewicz eds., PWN, Warszawa 1998.
- [3] F. A. Berezin: Quantization, Math USSR Izvestija 8, 5, 1109-1165 (1974).
- [4] R. Dey and K. Ghosh: Pull back coherent states and squeezed states and quantization, Symmetry, Integrability and Geometry: Methods and Applications (SIGMA),18, 028, 1-14 (2022), arxiv: 2108.08082
- [5] R. Dey and K. Ghosh: Berezin-type quantization on even-dimensional compact manifolds, J. Phys.: Conf. Ser. 2667 012003 (2023), arXiv:2210.08814v5
- [6] P. H. Doyle and J. G. Hocking: A Decomposition Theorem for n-dimensional manifolds, Proc. Amer. Math. Soc. 13 469 (1962)
- [7] M. Englis: Berezin Quantization and Reproducing Kernel on Complex Domains Trans. Amer. Math. Soc. 348 411 (1996).
- [8] B.V. Fedosov: A Simple Geometrical Construction of Deformation Quantization, J. Diff. Geom, 40 213-238 (1994).
- [9] J. P. Gazeau, P. Monceau: Generalized Coherent States for Arbitrary Quantum Systems in Mathematical Physics Studies book series, vol. 21/22, MPST, Springer 2000.
- [10] K. Ghosh: Berezin-type quantization of even-dimensional manifids and pullback coherent states, *Thesis ICTS-TIFR* https://thesis.icts.res.in/ (2023).
- [11] A. Odzijewicz: On Reproducing Kernels and Quantization of States, Commun. Math. Phys. 114, 577–597 (1988).
- [12] V. I. Paulsen and M. Raghupathi: An Introduction to the Theory of Reproducing Kernel Hilbert Spaces in Cambridge Studies in Advanced Mathematics vol. 152, Cambridge University Press 2016.
- [13] J. M. Radcliffe: Some Properties of Coherent Spin States, J. Phys. A: Gen. Phys. 4 313 (1971).
- [14] J. H. Rawnsley: Coherent States and Kahler Manifolds, Quart. J. Math., 28 403-415 (1977).

- [15] M. Spera: On Kählerian Coherent States, Proceedings of the International Conference on Geometry, Integrability and Quantization 241-256, Bulgarian Academy of Sciences, 2004.
- [16] D. Tong: The Quantum Hall Effect Tata Infosys Lectures http://www.damtp.cam.ac.uk/user/tong/qhe.html (2016).

RUKMINI DEY, EMAIL:RUKMINI@ICTS.RES.IN