

Longitudinal Momentum Spectra of pair created in a pulsed electric field at finite times: Are Oscillations “Real”¹

Deepak Sah^{*,1,2}, Manoranjan P. Singh^{1,2}

¹*Homi Bhabha National Institute, Training School Complex, Anushakti Nagar, Mumbai 400094, India* ²*Theory and Simulations Lab, Raja Ramanna Centre for Advanced Technology, Indore-452013, India*

*deepakk@rrcat.gov.in, dsah129@gmail.com

Abstract

The relativistic Dirac theory laid the foundation for understanding antiparticles and the potential for particle creation and annihilation within the realm of quantum field theory (QFT). This theory laid the groundwork for understanding the fundamental behavior of particles in the context of QFT. Moreover, the dynamic nature of gravitational fields, particularly in an expanding Universe, is pivotal in enabling the spontaneous generation of particles from the vacuum. While conceptually straightforward, this process leads to the transformation of creation operators into superpositions of creation and annihilation operators, with profound implications for our understanding of phenomena such as black holes, the early Universe, and quantum aspects of gravitational physics. Additionally, similar behavior is observed when dealing with time-varying electric fields, as discussed in reference [5]. This behavior effectively recovers the Schwinger effect, particularly in the case of a constant electric field. This phenomenon exemplifies the intricate interplay between quantum fields and varying background fields. QFT grapples with the profound and unresolved challenge of describing quantum systems undergoing time-dependent and far-from-equilibrium transformations. Traditional particle concepts become inadequate in such scenarios, necessitating the introduction of quasi-particles to characterize the evolving states better. Perturbative methods, typically employed in near-equilibrium conditions, are the cornerstone of QFT. However, this study delves into the semi-classical aspects of QFT, exploring the universal principles of asymptotic expansions to dissect the behavior of quantum systems as they evolve under the influence of time-dependent forces far from equilibrium. The research motivation behind this study stems from fundamental inquiries. The objective is to investigate whether real-time particle production observations are feasible and devise methodologies for addressing the non-equilibrium consequences arising from the back-reaction in particle generation mechanisms. In the specific context of Schwinger particle production, developing a real-time framework for tracking particle numbers carries significant promise. Such an approach can potentially lower the critical electric field threshold required for observing the Schwinger effect and offer precise control over this phenomenon. These insights are not confined to theoretical exploration but extend to practical applications, especially in the realm of high-intensity laser physics.

Furthermore, a real-time framework may unlock a deeper understanding of the time-dependent process of pair recombination, a pivotal phenomenon involving the annihilation of particle-antiparticle pairs. This understanding is crucial for unraveling the back-reaction effects that

¹Based on talk given on 12th September 2023 in 5th Conference on Extremely High Intensity Laser Physics (ExHILP 2023) 12-15 September 2023 at Center for Relativistic Laser Science (CoReLS), Institute for Basic Science (IBS), Gwangju Korea

influence particle production. Such insights enrich our comprehension of the Schwinger Effect and other particle production mechanisms. The significance of these developments extends beyond high-energy physics, reaching into the domain of cosmology. A deeper understanding of particle production and related processes may illuminate the fundamental dynamics of the universe, shedding light on the early moments of the cosmos and its subsequent evolution. This research serves as a bridge between the quantum realm of particle physics and the macroscopic universe, offering a more profound grasp of the physical processes underlying our universe's existence.


Pair creation can be envisaged as transforming a quantum system, traversing from an initial state of equilibrium to a final equilibrium state through an intermediate phase of non-equilibrium instigated by a potent external field. During these intermediate non-equilibrium phases, as matter fields interact with a time-dependent external field, the classical Hamiltonian loses its time translation symmetry. Consequently, various options for annihilation and creation operators (and, consequently, the vacuum) emerge within Fock quantization, thereby engendering complexities in depicting the evolution of the vacuum and generating particle-antiparticle pairs over time. Numerous approaches to vacuum selection have been explored in the literature, their choice contingent upon the specific system attributes under scrutiny and the adopted quantum theory. Adiabatic vacua have found wide applicability, especially in cosmological and Schwinger effect contexts. These adiabatic modes are constructed through semiclassical WKB-type approximations, often employing reference mode functions in the form of plane waves, particularly effective when the external background field evolves gradually over time. An alternative to this traditional approach is the utilization of adiabatic vacua, an idea initially proposed by Parker and subsequently formalized by Lueders and Roberts. Within this established framework, the asymptotic analysis of particle states in the distant past (in-states, prior to the initiation of the external field) and the distant future (out-states, long after the interaction with the external field has concluded) is well-understood. This relies on the expression of the quantum field operator in terms of creation and annihilation operators, linked to one-particle states in both the present and the future. By exploiting the interrelationship between these sets of operators in the past and the future, we can derive an equation for the S-matrix of the process and quantify the number of particles generated throughout the process.


We investigate the production of electron-positron pairs from the vacuum in a time-varying, spatially uniform pulsed electric field given by $E(t) = E_0 \text{sech}^2(t/\tau)$, with height of E_0 and width of τ . Such background field has received extensive attention in the literature [1,2,3], with a focus on its asymptotic behavior, particularly the probability of pair production. However, the problem of pair production after a finite evolution time has rarely been discussed [4,5]. This raises questions about the instantaneous appearance of particles in pair production and their behavior at intermediate times when using a formalism that involves solving an evolution equation for a dynamical quantity. Is it possible to make general statements about this behavior? To address these questions, we analytically compute the probability of (e^+e^-) pair production in momentum space using the exact solution of the one-particle time-dependent Dirac equation, and we compare the result with quantum kinetic theory (QKT), which is rigorously derived from QED by canonical quantization of the Dirac field and subsequent Bogoliubov transformation to a quasi-particle representation [6]. Both approaches allow us to study the particle momentum spectrum at any instant of time and can potentially unveil valuable information regarding quantum non-equilibrium physics. We analyze the Longitudinal Momentum Spectrum (LMS) of the created particles at finite times in both approaches. We observe oscillatory


structure in the LMS at time $t > \tau$, and this oscillation pattern continuously changes from $t > \tau$ up to $t < 6\tau$. After that, this oscillation slowly fades away, and we see smooth Gaussian-shape spectra for late times, reproducing Schwinger's non-perturbative signature. This oscillation behavior at finite time clearly illustrates the quantum interference effects associated with particle production, as explained in recent work [7,9]. It is worth noting that both approaches exhibit quantum interference patterns at finite times, manifested as oscillations observed in the LMS. This clearly reveals that these oscillations are not due to transient excitations and basis-dependent signatures [8]. We again emphasize that the oscillations seen in the LMS from both approaches are not artifacts but rather possess significant physical relevance. Therefore, comparing the two approaches sheds light on pair production at finite times and provides more physical information on the non-equilibrium state.


References:


- [1] A. I. Nikishov, *Soviet Physics JETP* **30** (1970), 660–662.
- [2] A. B. Balantekin et al., *Int. J. Mod. Phys. A* **6** (1991), 695–725.
- [3] S. P. Gavrilov and D. M. Gitman, *Phys. Rev. D* **53** (1996), 7162–7175.
- [4] J. Hallin and P. Liljenberg, *Phys. Rev. D* **52** (1995), 1150–1164.
- [5] A. Di Piazza, *Phys. Rev. D* **70** (2004), 053013.
- [6] Y. Kluger et al., *Phys. Rev. Lett.* **67**, 2427 (1991).
- [7] D. Sah and M. P. Singh, [arXiv:2301.06545 [hep-ph]].
- [8] R. Dabrowski and G. V. Dunne, *Phys. Rev. D* **90**, 025021 (2014).
- [9] D. Sah and M. P. Singh, [arXiv:2309.12079 [hep-ph]].




































This figure "2.png" is available in "png" format from:


<http://arxiv.org/ps/2405.02947v2>










































This figure "ACCE_PHY.png" is available in "png" format from:


<http://arxiv.org/ps/2405.02947v2>






























This figure "acc._1.png" is available in "png" format from:

<http://arxiv.org/ps/2405.02947v2>

This figure "acc_decc.png" is available in "png" format from:

<http://arxiv.org/ps/2405.02947v2>



This figure "apprx_ftime.png" is available in "png" format from:

<http://arxiv.org/ps/2405.02947v2>

This figure "arxived_soon.png" is available in "png" format from:

<http://arxiv.org/ps/2405.02947v2>

This figure "arxived_soon_a.png" is available in "png" format from:

<http://arxiv.org/ps/2405.02947v2>

This figure "arxived_soon_b.png" is available in "png" format from:

<http://arxiv.org/ps/2405.02947v2>

This figure "asytm_pp.png" is available in "png" format from:

<http://arxiv.org/ps/2405.02947v2>

