

Arrival Times Versus Detection Times

Sheldon Goldstein*, Roderich Tumulka[†] and Nino Zanghì[‡]

May 7, 2024

Abstract

How to compute the probability distribution of a detection time, i.e., of the time which a detector registers as the arrival time of a quantum particle, is a long-debated problem. In this regard, Bohmian mechanics provides in a straightforward way the distribution of the time at which the particle actually does arrive at a given surface in 3-space in the absence of detectors. However, as we discuss here, since the presence of detectors can change the evolution of the wave function and thus the particle trajectories, it cannot be taken for granted that the arrival time of the Bohmian trajectories in the absence of detectors agrees with the one in the presence of detectors, and even less with the detection time. In particular, we explain why certain distributions that Das and Dürr [4] presented as the distribution of the detection time in a case with spin, based on assuming that all three times mentioned coincide, is actually not what Bohmian mechanics predicts.

Key words: Bohmian mechanics; POVM.

1 Introduction

In a time-of-arrival experiment, one places detectors along a surface Σ in 3-space and asks for the time T_D and place \mathbf{Q}_D recorded for the detection of a particle. In Bohmian mechanics, a particle has a well defined position $\mathbf{Q}(t)$ at any time t , so there is a fact even in the absence of detectors about when and where the particle first arrives at Σ . However, the presence of a detector can change the Bohmian trajectories, even if no detector clicks. After all, as we discuss, the detector may effectively collapse away parts of the wave function that would have influenced the trajectory. That is, possibly $\mathbf{Q}_{WOD}(t) \neq \mathbf{Q}_{WID}(t)$, where *WOD* means *without detector*, *WID* means *with detector*. As a consequence, possibly

$$T_{WOD} \neq T_{WID} \tag{1}$$

*Departments of Mathematics and Physics, Rutgers University, 110 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA. E-mail: oldstein@math.rutgers.edu

[†]Fachbereich Mathematik, Eberhard-Karls-Universität Tübingen, Auf der Morgenstelle 10, 72076 Tübingen, Germany. E-mail: roderich.tumulka@uni-tuebingen.de

[‡]INFN sezione di Genova and Dipartimento di Fisica, Università di Genova, via Dodecaneso 33, 16146 Genova, Italy. E-mail: zanghi@ge.infn.it

for the first arrival time without detectors,

$$T_{WOD} \equiv \inf\{t \geq 0 : \mathbf{Q}_{WOD}(t) \in \Sigma\} \quad (2)$$

and likewise with detectors. We write \mathcal{P}_{WOD} (\mathcal{P}_{WID} , \mathcal{P}_D) for the probability distribution of T_{WOD} (T_{WID} , T_D).

Das and Dürr [4] computed \mathcal{P}_{WOD} for a particular setup involving a spin-1/2 particle in an axially symmetric wave guide. Assuming that the presence of the detector would be a “mild disturbance” and thus that $T_{WOD} = T_{WID} = T_D$ up to negligible errors, they suggested that the prediction of Bohmian mechanics for \mathcal{P}_D agrees with \mathcal{P}_{WOD} .

Recently, we proved in [14] that the specific distribution \mathcal{P}_{WOD} computed in [4] cannot be the prediction of Bohmian mechanics for \mathcal{P}_D because it is not given by a positive-operator-valued measure (POVM), not even approximately.¹ Here, we give more detail about the proof and related considerations, including a more detailed analysis of the distributions compatible with a POVM. We provide as well further, alternative arguments for $\mathcal{P}_{WOD} \neq \mathcal{P}_D$, including some discussion of the relations among T_{WOD} , T_{WID} , and T_D . In particular, we suggest that even $\mathbf{Q}_{WID} = \mathbf{Q}_D$ and $T_{WID} = T_D$ can’t be taken for granted.

To be sure, there are situations in which the presence of detectors does not significantly influence the trajectories, for example in far-field detection (i.e., as $t \rightarrow \infty$, at great distances from the support of the initial wave function) as considered in scattering theory. In that case,

$$T_{WOD} = T_{WID} = T_D, \quad (3)$$

at least approximately, and \mathcal{P}_D can be computed by computing \mathcal{P}_{WOD} . Specifically, for Σ the sphere around the origin of large radius R , the distribution of T_{WOD} is (with small relative error) given by a POVM, in fact a projection-valued-measure (PVM), namely corresponding to the quantum observable $mR/|\hat{\mathbf{p}}|$ with $\hat{\mathbf{p}} = (\hat{p}_x, \hat{p}_y, \hat{p}_z)$ the momentum operator and m the particle’s mass.² But in near-field detection, there is no self-adjoint operator that could serve as the observable for T_D , which is the origin of the perennial debate about how to compute \mathcal{P}_D . This situation has led in orthodox quantum mechanics to a multitude of guesses about \mathcal{P}_D , as well as to premature claims

¹A POVM, or Positive-Operator-Valued Measure, is a generalization of the concept of observable in quantum mechanics. In traditional quantum mechanics, observables are associated with collections of orthogonal projectors E_i , each corresponding to a possible outcome of a measurement. The family of such projectors is called a projection-valued measure (PVM) and, via the spectral theorem, is in correspondence with the usual notion of self-adjoint-operator-as-observable. However, a POVM allows for more flexibility by allowing E_i to be positive operators that are not necessarily projectors, but are such that $\sum_i E_i = I$, where I is the identity operator. Then the standard probabilistic rules of quantum mechanics extend to these more general quantum observables: for a system in the state ψ and measurement associated with the POVM $E = \{E_i\}$, the probability of outcome i is given by $\langle \psi | E_i | \psi \rangle$. Indeed, it can be shown that the probabilities of *any* quantum experiment can be expressed in terms of a POVM. What has just been described for observables with discrete values can be easily extended to observables with continuous values, replacing E_i by $E(dt)$.

²Moreover, for large R , the joint distribution of $(T_{WOD}, \mathbf{Q}_{WOD})$ is given by the observable $(mR|\hat{\mathbf{p}}|^{-1}, R|\hat{\mathbf{p}}|^{-1}\hat{\mathbf{p}})$.

that it is impossible to determine \mathcal{P}_D for a detecting surface because of the quantum Zeno effect.

A moral from our discussion is that *detection time is not the same thing as arrival time*. It turns out to be wise to keep in mind the advice often emphasized in orthodox quantum mechanics that we should not take measurement for granted, that the apparatus often plays an active role that must be taken into account. Note, however, that the difference between detection time and arrival time is clearly meaningful and visible in a precise theory such as Bohmian mechanics, whereas in the framework of orthodox quantum mechanics what is happening on the microscopic level is too vague for this to be so.

The remainder of this paper is organized as follows. In Section 2, we recall the equations of Bohmian mechanics, describe the setup considered by Das and Dürr in [4], and discuss some reasons for expecting that $\mathcal{P}_{WOD} \neq \mathcal{P}_D$. In Section 3, we analyze POVMs on \mathbb{C}^2 , provide alternative proofs of the no-go theorem of [14], and discuss why superluminal signaling would be possible if $\mathcal{P}_{WOD} = \mathcal{P}_D$. In Section 4, we consider different equivariant equations of motion and discuss their relevance to the question whether $T_D = T_{WOD}$, as well as whether $T_D = T_{WID}$. In Section 5, we conclude.

2 Arrival Times

2.1 Bohmian Arrival Times

Bohmian mechanics, a one-parameter family. We consider a 1-particle wave function $\Psi = \Psi_t(\mathbf{q})$ that is spinor-valued and obeys Schrödinger's equation

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \Delta \Psi + V \Psi, \quad (4)$$

where $V(\mathbf{q})$ is a Hermitian 2×2 matrix for every $\mathbf{q} \in \mathbb{R}^3$. As the Bohmian equation of motion for the particle position $\mathbf{Q} = \mathbf{Q}(t)$, two different possibilities are sometimes considered, corresponding to $\lambda = 1$ or $\lambda = 0$ in the general form

$$\frac{d\mathbf{Q}}{dt} = \frac{\hbar}{m} \operatorname{Im} \frac{\Psi^\dagger \nabla \Psi}{\Psi^\dagger \Psi}(\mathbf{Q}) + \lambda \frac{\hbar}{2m} \frac{\nabla \times (\Psi^\dagger \boldsymbol{\sigma} \Psi)}{\Psi^\dagger \Psi}(\mathbf{Q}) \quad (5)$$

$$= \frac{\hbar}{m} \operatorname{Im} \frac{\Psi^\dagger (I_2 - i\lambda \boldsymbol{\sigma} \times) \nabla \Psi}{\Psi^\dagger \Psi}(\mathbf{Q}), \quad (6)$$

where $\boldsymbol{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ is the triple of the Pauli matrices, $\Psi^\dagger \Phi$ denotes the scalar product in the spinor space \mathbb{C}^2 , \times the cross product in 3d, and I_2 the 2×2 identity matrix.³ In [4], $\lambda = 1$ was considered, which arises in the non-relativistic limit of the Bohmian equation of motion associated with the Dirac equation. In fact, $|\Psi|^2$ is equivariant for every $\lambda \in \mathbb{R}$. T_{WOD} can be computed for every initial position \mathbf{Q}_0 by solving (4), (5),

³Equation (5) has perhaps been considered so far only for $\lambda = 0$ or $\lambda = 1$. But for the purpose of our analysis we shall allow other values as well.

and (2). Since \mathbf{Q}_0 is random with distribution $|\Psi_0|^2$, T_{WOD} is random, too, and its distribution can in principle be computed.

Note that even if $V(\mathbf{q})$ is real-valued (i.e., a multiple of I_2), so that the Schrödinger equation (4) does not even couple to spin, then the equation of motion (5) still couples to spin for $\lambda \neq 0$, leading to a spin dependence of T_{WOD} and the question whether T_D might depend on the spin as well.

Setup. Das and Dürr [4] considered the real-valued potential V given by

$$V(x, y, z) \equiv \begin{cases} \infty & \text{if } z < 0 \\ \frac{m}{2}\omega^2(x^2 + y^2) & \text{if } z \geq 0, \end{cases} \quad (7)$$

where $\omega > 0$ is a constant. This potential does two things: it provides a reflecting wall at $z = 0$ and acts as a wave guide that will keep the wave function near the z -axis.

For any unit vector \mathbf{n} in three-dimensional space, consider the spinor $|\mathbf{n}\rangle$ in \mathbb{C}^2 defined uniquely up to a phase θ by the condition $\mathbf{n} = \langle \mathbf{n} | \boldsymbol{\sigma} | \mathbf{n} \rangle$. Explicitly,

$$|\mathbf{n}\rangle = e^{i\theta} \begin{pmatrix} \cos(\alpha/2) \\ \sin(\alpha/2) e^{-i\beta} \end{pmatrix} \quad (8)$$

with α and β spherical coordinates of \mathbf{n} , i.e., $\alpha \in [0, \pi]$ the angle between \mathbf{n} and the z -axis and $\beta \in [0, 2\pi)$ the angle between (n_x, n_y) and the x -axis (azimuthal angle).

The initial wave function is taken to factorize,

$$\Psi_0 = |\mathbf{n}\rangle \otimes \psi_0 \quad (9)$$

with arbitrary unit spinor $|\mathbf{n}\rangle$ and a certain fixed choice of ψ_0 . The surface considered is

$$\Sigma = \{(x, y, z) \in \mathbb{R}^3 : z = L\} \quad (10)$$

for some $L > 0$. It follows that $\Psi_t = |\mathbf{n}\rangle \otimes \psi_t$ with ψ_t the solution of the corresponding scalar Schrödinger equation, and that $\psi_t(x, y, z)$ vanishes for $z < 0$ and all t (but can be non-zero for $z > L$).

Findings for \mathcal{P}_{WOD} . Das and Dürr found that the distribution $\mathcal{P}_{WOD, \mathbf{n}}$ of T_{WOD} for given \mathbf{n} , with spin-dependent law of motion (5) with $\lambda = 1$, depends strongly on the choice of the spinor $|\mathbf{n}\rangle$. This is visible in Figure 1 as the difference between the red and the blue curve, and in Figure 2 as the non-constancy of either the red or the black curve. We will derive below that \mathcal{P}_D (and its mean) can depend on \mathbf{n} only in very limited ways that are incompatible with the \mathbf{n} -dependence of $\mathcal{P}_{WOD} = \mathcal{P}_{WOD, \mathbf{n}}$. We also note for comparison that for the spin-independent law of motion (5) with $\lambda = 0$, the trajectories and therefore T_{WOD} are of course independent of \mathbf{n} .

2.2 The Decoupling Argument

Before we discuss the no-go theorem of [14] in Section 3, we describe another, much simpler argument to the effect that $T_{WOD} \neq T_D$ in the setting of [4]. We will call it the *decoupling argument*:




Figure 1: \mathcal{P}_{WOD} in the setting of [4], here denoted by $\Pi_{\text{Bohm}}^{\alpha|\beta}$, for two different spinors $|\mathbf{n}\rangle$; the blue curve ($\alpha = 0$) corresponds to \mathbf{n} parallel to the z -axis, the red curve ($\alpha = \frac{\pi}{2}$) to \mathbf{n} perpendicular to the z -axis. (Reproduced from [4]. The inset shows a detail of the red curve in magnification.)




Figure 2: The red and black curves show, for two values of the parameter ω from (7), the expected arrival time $\mathbb{E}T_{WOD}$, here denoted by $\langle \tau \rangle$, in the setting of [4] as a function of α , the angle between \mathbf{n} and the z -axis. (Reproduced from [4]. The red and black curves are not related to the red and black curves in Figure 1.)

Assume that the initial wave function Ψ_0 of the particle is of product form (9) with any unit spinor $|\mathbf{n}\rangle$ and any normalized scalar function ψ_0 . One of the remarkable features of \mathcal{P}_{WOD} is its dependence on \mathbf{n} . Although the Hamiltonian H_p of the particle does not couple to the spin, i.e., is of the form

$$H_p = I_2 \otimes \tilde{H}_p \quad (11)$$

relative to the factorization of Hilbert space into spin and position degrees of freedom, the Bohmian equation of motion (5) does couple to the spin, which is why the trajectories (and hence T_{WOD}) depend on $|\mathbf{n}\rangle$. Now we regard the detectors as a quantum system coupled to the particle by means of an interaction Hamiltonian H_{int} . If H_{int} does not couple to the spin of the particle, i.e., if

$$H_{\text{int}} = I_2 \otimes \tilde{H}_{\text{int}}, \quad (12)$$

then also the full wave function Φ of particle and detectors together at any (late) time will be of the form $|\mathbf{n}\rangle \otimes \text{something}$, where the “something” does not depend upon \mathbf{n} . Thus the $|\Phi|^2$ distribution of the apparatus’s pointer position or records will be independent of \mathbf{n} , so that \mathcal{P}_D will as well. This shows that the Bohmian prediction for \mathcal{P}_D differs from that for \mathcal{P}_{WOD} , so $T_{WOD} \neq T_D$, provided (12) holds. This completes the argument. The mathematical content is that

$$\text{assuming (11) and (12), } \mathcal{P}_D \text{ cannot depend on } \mathbf{n}. \quad (13)$$

The decoupling argument shows two things: First, that (3) is not an innocent assumption, as it is provably false whenever (11) and (12) are true. Second, that it is reasonable to expect that the observed distribution in the experiment is independent of \mathbf{n} when (11) holds, as (12) sounds like a reasonable assumption, one that would be satisfied, at least to a high degree of approximation, for many apparently reasonable detection procedures. However, since observations in actual experiments include relativistic effects in principle, and since the relativistic Hamiltonian of an electron (i.e., the Dirac Hamiltonian) does not decouple from spin as in (11) (and not either as $I_4 \otimes \tilde{H}_p$ for \mathbb{C}^4 as the spin space), the relativistic (and thus also the observed) distribution for T_D should not (exactly) be independent of \mathbf{n} . On the other hand, the no-go theorem of [14] does not assume (11) or (12), which is why it provides a relevant piece of information that goes beyond the decoupling argument.

2.3 How the Presence of Detectors Changes Trajectories

Let us flesh out our statement that the presence of a detector can change the Bohmian trajectories, even if it does not click.

Consider a double-slit experiment with a detector in the left slit. It will effectively collapse the 1-particle wave function to a wave packet that passed either through the left or the right slit, depending on whether the detector clicked or not. A wave that passed through one slit only will lead to different Bohmian trajectories (no interference fringes, possibility to cross the symmetry plane between the slits) than a wave that

passed through both. In this example, the presence of the detector perhaps has only a minor effect on the arrival *time* at the screen, but a major effect on the arrival *place* at the screen. But there are three aspects to note: first, that the effect on the trajectories is not a mild disturbance, but a drastic change of the trajectory; second, that using better detectors will not make the change less drastic; and third, once we realize that the presence of detectors can drastically change the trajectories, we have to expect that in another case also the arrival *time* would be affected.

Since in the double-slit experiment with a detector in a slit, the Bohmian trajectory is changed *beyond* the plane containing the detector and the two slits, one might wonder whether, in general, the trajectory is not changed *before* the surface Σ containing detectors. This cannot be taken for granted: even neglecting that a part of the wave might be reflected from a detecting surface, we notice that in the free evolution of ψ in the absence of detectors on Σ , parts of ψ beyond Σ can in general also propagate back to the region before Σ and contribute there. Thus, if parts of ψ get collapsed away when passing Σ , ψ_t in the region before Σ will be different from what it would have been in the free evolution. As a consequence, also trajectories will be different from what they would have been in the free evolution.

3 POVMs

In this section, we discuss several ways of proving the no-go theorem mentioned above, which asserts [14] that *there is no possible experiment (for which the experimental design does not depend upon \mathbf{n}) that would, when applied to a particle in the setup of [4] with $\Psi_0 = |\mathbf{n}\rangle \otimes \psi_0$, have outcome with distribution $\mathcal{P}_{WOD,\mathbf{n}}$ for every \mathbf{n}* . The theorem means, in other words, that there can be no experiment or procedure that yields the arrival time T_{WOD} unless the procedure itself directly incorporates information about the direction \mathbf{n} of the spin state; it shows in particular that if the experiment consists of putting on Σ the kind of device commonly regarded as a detector, however it might work, the distribution \mathcal{P}_D of its outcome T_D will not in general agree with \mathcal{P}_{WOD} .

By experiment we shall henceforth always mean one whose design is independent of \mathbf{n} (respectively Ψ_0). By way of contrast, suppose we were to measure the position of the particle at the initial time and then use our knowledge of the quantum state of the particle to compute (what would have been) its trajectory and thereby its arrival time T_{WOD} . Such a procedure would not count as an experiment according to this stipulation.

We start from the known fact that for every experiment that can be carried out on a system with arbitrary initial wave function Ψ_0 , there is a POVM $E_0(\cdot)$ such that the outcome statistics are given by $\langle \Psi_0 | E_0(\cdot) | \Psi_0 \rangle$. This fact is a theorem in Bohmian mechanics [8], [20, Sec. 5.1]. In particular, \mathcal{P}_D must be given by some POVM $E_0(dt)$. But, as we will show, \mathcal{P}_{WOD} is not given by a POVM.

3.1 Spin POVM

Since Ψ_0 factorizes according to (9) into a spinor $|\mathbf{n}\rangle$ and a scalar function ψ_0 , if we regard ψ_0 as fixed and $|\mathbf{n}\rangle$ as variable, then the spin dependence of the outcome statistics is captured by the POVM

$$E(dt) = \langle \psi_0 | E_0(dt) | \psi_0 \rangle, \quad (14)$$

where the inner product is a partial inner product taken in the position degrees of freedom but not those for the spin. Thus $E(dt)$ is a *spin POVM*, a POVM on \mathbb{R} acting on \mathbb{C}^2 ; as such it is given by 2×2 matrices. It follows that for $\Psi_0 = |\mathbf{n}\rangle \otimes \psi_0$, the distribution of the detection time is

$$P_{\mathbf{n}}(dt) \equiv \langle \mathbf{n} | E(dt) | \mathbf{n} \rangle. \quad (15)$$

For example, the most familiar spin POVM is for the measurement of the z -component σ_z of spin for a spin-1/2 particle, which is given by

$$E = \frac{1}{2} \left(\delta(t-1) + \delta(t+1) \right) I_2 + \frac{1}{2} \left(\delta(t-1) - \delta(t+1) \right) \sigma_z \quad (16)$$

with I_2 the 2×2 identity.

In the following, we point out several properties of distributions arising from spin POVMs that are incompatible with the \mathbf{n} -dependence of \mathcal{P}_{WOD} found in [4]; each of them proves the no-go theorem asserting that $\mathcal{P}_{WOD} \neq \mathcal{P}_D$. Such a property was first pointed out in [14].

3.2 Properties of Distributions from POVMs

Trace property. Since $|\mathbf{n}\rangle$ and $|\mathbf{-n}\rangle$ form an orthonormal basis of spin space \mathbb{C}^2 , and since the trace of an operator is the same in any basis, we have that for any spin POVM $E(\cdot)$,

$$\langle \mathbf{-n} | E(\cdot) | \mathbf{-n} \rangle + \langle \mathbf{n} | E(\cdot) | \mathbf{n} \rangle = \text{Tr } E(\cdot) \text{ is independent of } \mathbf{n}. \quad (17)$$

In particular,

$$P_{\mathbf{n}}(dt) + P_{\mathbf{-n}}(dt) \text{ is independent of } \mathbf{n}. \quad (18)$$

Now let $\mathbf{x}, \mathbf{y}, \mathbf{z}$ be the unit vectors in the x, y, z direction. As a consequence of the axial symmetry of the setup of Das and Dürr, $\mathcal{P}_{WOD, \mathbf{n}}$ is independent of β ; thus

$$\mathcal{P}_{WOD, \mathbf{-x}} = \mathcal{P}_{WOD, \mathbf{x}}. \quad (19)$$

They also proved that

$$\mathcal{P}_{WOD, \mathbf{-z}} = \mathcal{P}_{WOD, \mathbf{z}}. \quad (20)$$

If \mathcal{P}_{WOD} of [4] came from a POVM E , then we would have that $\mathcal{P}_{WOD, \mathbf{n}} = P_{\mathbf{n}}$ for all \mathbf{n} , so that $P_{\mathbf{-x}} = P_{\mathbf{x}}$ and $P_{\mathbf{-z}} = P_{\mathbf{z}}$. But then, by (18),

$$P_{\mathbf{x}} = \frac{1}{2}(P_{\mathbf{x}} + P_{\mathbf{-x}}) = \frac{1}{2}(P_{\mathbf{z}} + P_{\mathbf{-z}}) = P_{\mathbf{z}} \quad (21)$$

and hence $\mathcal{P}_{WOD,\mathbf{x}} = \mathcal{P}_{WOD,\mathbf{z}}$, in contradiction with the strong difference between them found in [4] (clearly visible as the difference between the red and blue curves in Figure 1). This is our first proof of the no-go theorem.

In fact, there cannot even exist a spin POVM E that agrees with $\mathcal{P}_{WOD,\mathbf{n}}$ for merely the three directions $\mathbf{n} = \mathbf{x}, -\mathbf{x}$ and \mathbf{z} . This is so because in this case the trace property would imply that $P_{\mathbf{z}} \leq 2P_{\mathbf{x}}$, which is clearly incompatible with what we see in Fig. 1. This is our second proof of the no-go theorem.

Remark. Apart from arrival time questions, Eq. (17) can be used to show that the spin state $|\mathbf{n}\rangle$ itself (or the associated direction \mathbf{n}) is not measurable; that is, there is no experiment that could, upon the input of a particle with arbitrary spin state $|\mathbf{n}\rangle$, yield \mathbf{n} as the result. Indeed, if such an experiment existed, it would be associated with a POVM $F(d\mathbf{u})$ on the unit sphere $\{\mathbf{u} \in \mathbb{R}^3 : |\mathbf{u}| = 1\}$ acting on \mathbb{C}^2 such that $\langle \mathbf{n} | F(d\mathbf{u}) | \mathbf{n} \rangle = \delta(\mathbf{u} - \mathbf{n}) d\mathbf{u}$. But $\delta(\mathbf{u} - \mathbf{n}) + \delta(\mathbf{u} + \mathbf{n})$ clearly depends upon \mathbf{n} , so that by (17) no such POVM exists.

Relation to Pauli matrices. Any Hermitian 2×2 matrix M can be written in the form

$$M = m_0 I_2 + \mathbf{m} \cdot \boldsymbol{\sigma} \quad (22)$$

with some real m_0 and some real 3-vector \mathbf{m} . We thus have that

$$E(dt) = e_0(dt) I_2 + \mathbf{e}(dt) \cdot \boldsymbol{\sigma} \quad (23)$$

and thus

$$P_{\mathbf{n}}(dt) = e_0(dt) + \mathbf{e}(dt) \cdot \mathbf{n} \quad (24)$$

for suitable e_0 and $\mathbf{e} = (e_x, e_y, e_z)$.

Note that by (24) $e_0(dt) = \frac{1}{2}(P_{\mathbf{n}}(dt) + P_{-\mathbf{n}}(dt))$ (yielding (18) once again). Thus e_0 must be a probability measure. As a consequence, the signed measures e_ν (for $\nu = x, y, z$) must have vanishing total mass, $\int_{-\infty}^{\infty} e_\nu(dt) = 0$. Moreover, any such e_0 and \mathbf{e} correspond via (23) to a spin POVM if and only if $\mathbf{e}^2 \leq e_0^2$, since, as it is not hard to see, this amounts to positivity.

Axial symmetry. Now take the axial symmetry of the setup into account. Then were \mathcal{P}_{WOD} given by a POVM E , E itself would have to be axially symmetric, so that its dependence on $\boldsymbol{\sigma}$ would be via only σ_z . Thus it would be of the form

$$E(dt) = e_0(dt) I_2 + e_z(dt) \sigma_z, \quad (25)$$

which leads to

$$P_{\mathbf{n}}(dt) = e_0(dt) + e_z(dt) \cos \alpha. \quad (26)$$

Setting $\alpha = \pi/2$ we would thus have that $e_0 = P_{\mathbf{x}}$; setting $\alpha = 0$ and $\alpha = \pi$ and adding we would obtain that $2e_0 = P_{\mathbf{z}} + P_{-\mathbf{z}}$. But this—or (18)—implies that $P_{\mathbf{z}} + P_{-\mathbf{z}} = 2P_{\mathbf{x}}$, and hence that $\mathcal{P}_{WOD,\mathbf{z}} + \mathcal{P}_{WOD,-\mathbf{z}} = 2\mathcal{P}_{WOD,\mathbf{x}}$, in contradiction with what we see in Figure 1. This is our third proof of the no-go theorem. (Note that the positivity condition for an axially symmetric POVM (26) is that $|e_z| \leq e_0$.)

Condition for spin independence. It follows from (24) that for any unit vector \mathbf{a} , $P_{\mathbf{a}} - P_{-\mathbf{a}} = 2\mathbf{e} \cdot \mathbf{a}$, so the vector $\mathbf{e} = \mathbf{e}(dt)$ can be expressed for any orthonormal basis $\mathbf{a}, \mathbf{b}, \mathbf{c}$ as

$$\mathbf{e} = \frac{1}{2}(P_{\mathbf{a}} - P_{-\mathbf{a}})\mathbf{a} + \frac{1}{2}(P_{\mathbf{b}} - P_{-\mathbf{b}})\mathbf{b} + \frac{1}{2}(P_{\mathbf{c}} - P_{-\mathbf{c}})\mathbf{c}. \quad (27)$$

Therefore,

$$\text{if } P_{-\mathbf{a}} = P_{\mathbf{a}}, P_{-\mathbf{b}} = P_{\mathbf{b}}, \text{ and } P_{-\mathbf{c}} = P_{\mathbf{c}} \text{ for orthonormal} \\ \text{vectors } \mathbf{a}, \mathbf{b}, \mathbf{c}, \text{ then } \mathbf{e}(dt) = \mathbf{0}, \text{ and } P_{\mathbf{n}} \text{ is independent of } \mathbf{n}. \quad (28)$$

Now use (19) and the fact that, by axial symmetry, $\mathcal{P}_{WOD,-\mathbf{y}} = \mathcal{P}_{WOD,\mathbf{y}}$, as well as (20); if \mathcal{P}_{WOD} came from a POVM, then $\mathcal{P}_{WOD,\mathbf{n}} = P_{\mathbf{n}}$ for all \mathbf{n} , so $P_{-\mathbf{x}} = P_{\mathbf{x}}$, $P_{-\mathbf{y}} = P_{\mathbf{y}}$, and $P_{-\mathbf{z}} = P_{\mathbf{z}}$. By (28), $P_{\mathbf{n}}$ must then be independent of \mathbf{n} , so $\mathcal{P}_{WOD,\mathbf{n}}$ is then independent of \mathbf{n} , but we know it is not. This is our fourth proof of the no-go theorem.

Sinusoidal dependence. Let us go back to (24). In terms of the spherical coordinates α, β mentioned in Section 2, (24) becomes a combination of the first (and 0-th) spherical harmonics:

$$P_{\mathbf{n}}(dt) = e_0(dt) + e_x(dt) \sin \alpha \cos \beta + e_y(dt) \sin \alpha \sin \beta + e_z(dt) \cos \alpha. \quad (29)$$

If this is to agree with $\mathcal{P}_{WOD,\mathbf{n}}$, which is axially symmetric, it must be independent of β and thus of the form

$$P_{\mathbf{n}}(dt) = e_0(dt) + e_z(dt) \cos \alpha, \quad (30)$$

as we argued earlier.⁴ But the dependence of $\mathcal{P}_{WOD,\mathbf{n}}(dt)$ on α is not of the form (30). This can perhaps be seen most easily by considering the expected detection time.

Expected detection time. It follows from (30) that were $\mathcal{P}_{WOD,\mathbf{n}}$ given by a POVM, also the expectation $\mathbb{E}T_D$ of T_D would depend on α in a sinusoidal way: We would have that

$$\mathbb{E}T_D = \int_0^\infty t P_{\mathbf{n}}(dt) \quad (34)$$

$$= \int_0^\infty t e_0(dt) + \cos \alpha \int_0^\infty t e_z(dt) \quad (35)$$

$$= \tau_0 + \tau_z \cos \alpha \quad (36)$$

⁴A perhaps slightly more convenient formula for $P_{\mathbf{n}}$ in the case of axial symmetry is obtained by expanding the POVM E (25) in terms of $I_2 - \sigma_z$ and σ_z instead of I_2 and σ_z :

$$E(dt) = A(dt)(I_2 - \sigma_z) + B(dt)\sigma_z, \quad (31)$$

so that

$$P_{\mathbf{n}}(dt) = A(dt)(1 - \cos \alpha) + B(dt) \cos \alpha. \quad (32)$$

Setting $\alpha = 0$ and $\pi/2$ we see that $A = P_{\mathbf{x}}$ and $B = P_{\mathbf{z}}$, and (32) becomes

$$P_{\mathbf{n}}(dt) = P_{\mathbf{x}}(dt)(1 - \cos \alpha) + P_{\mathbf{z}}(dt) \cos \alpha. \quad (33)$$

with $\tau_\nu \equiv \int_0^\infty t e_\nu(dt)$ (for $\nu = 0, z$). Compare this to the graph of the dependence of $\mathbb{E}T_{WOD}$ on α found in [4], reproduced here as Figure 2. They are clearly incompatible. This is our fifth proof of the no-go theorem.

The sinusoidal dependence of $\mathbb{E}T_D$ on α can also be obtained from (25):

$$\mathbb{E}T_D = \langle \mathbf{n} | M | \mathbf{n} \rangle, \quad (37)$$

where the matrix M is the mean of the POVM $E(dt)$,

$$M = \int_0^\infty t E(dt) = \tau_0 I_2 + \tau_z \sigma_z, \quad (38)$$

yielding (36).

Approximate measurement. We've seen that there is no POVM that will make $P_{\mathbf{n}}$ coincide with $\mathcal{P}_{WOD,\mathbf{n}}$ even for at least the 3 directions $\mathbf{z}, \mathbf{x}, -\mathbf{x}$ and certainly not for all 4 directions $\mathbf{z}, -\mathbf{z}, \mathbf{x}, -\mathbf{x}$. This remains true when we replace “coincide” with “coincide approximately”:

For every spin POVM E there is a direction \mathbf{n} among $\mathbf{z}, -\mathbf{z}, \mathbf{x}, -\mathbf{x}$ such that (39)

$$\|\mathcal{P}_{WOD,\mathbf{n}} - P_{\mathbf{n}}\| \geq \|\mathcal{P}_{WOD,\mathbf{x}} - \mathcal{P}_{WOD,\mathbf{z}}\|/2.$$

Since the right-hand side as computed in [4] is a non-negligible quantity, this relation means that $\mathcal{P}_{WOD,\mathbf{n}}$ cannot be close to $P_{\mathbf{n}}$ for all 4 directions.

To verify (39), consider the triangle inequality in the version

$$\|D_{\mathbf{z}} + D_{-\mathbf{z}} - D_{\mathbf{x}} - D_{-\mathbf{x}}\| \leq \|D_{\mathbf{z}}\| + \|D_{-\mathbf{z}}\| + \|D_{\mathbf{x}}\| + \|D_{-\mathbf{x}}\| \quad (40)$$

for $D_{\mathbf{n}} = \mathcal{P}_{WOD,\mathbf{n}} - P_{\mathbf{n}}$. By (18), the contributions from P cancel on the left-hand side. By (19) and (20), the left-hand side equals $2\|\mathcal{P}_{WOD,\mathbf{x}} - \mathcal{P}_{WOD,\mathbf{z}}\|$. Finally, it can't be the case that each of the 4 summands on the right-hand side is less than a quarter of the left-hand side, which completes the proof of (39). (A similar statement, involving the positive part of $\mathcal{P}_{WOD,\mathbf{z}} - 2\mathcal{P}_{WOD,\mathbf{x}}$, could be made for just the 3 directions.)

3.3 Previous Results

To put the no-go theorem into context, we mention a couple of similar previous results.

First, in [8], it was shown that, according to Bohmian mechanics, there is no experiment that could measure the velocity $d\mathbf{Q}/dt$ of a given single particle.⁵ One method of proof is to show that the distribution of $d\mathbf{Q}/dt$ is not given by a POVM; the situation is different with the *asymptotic* (long-time average) velocity, which is measurable.

⁵Measuring its position and then using the guiding equation to compute the velocity requires knowledge of the quantum state and hence does not count as an experiment according to our stipulation.

Second, Vona, Hinrichs, and Dürr [22] already studied whether \mathcal{P}_{WOD} is given by a POVM. Their considerations are not limited to the setting of [4], and many of them can be applied to rather arbitrary V , Σ , and Ψ_0 (evolving in \mathbb{R}^d , with or without spin). They considered the set S of initial wave functions Ψ_0 obeying the so-called current positivity condition

$$\boldsymbol{\nu}(\mathbf{q}) \cdot \mathbf{j}(t, \mathbf{q}) \geq 0 \quad \forall \mathbf{q} \in \Sigma \quad \forall t \geq 0, \quad (41)$$

where $\boldsymbol{\nu}(\mathbf{q})$ is the outward unit normal vector on Σ at \mathbf{q} and \mathbf{j} is given by $\Psi^\dagger \Psi$ times the right-hand side of (5) or (6). (They considered $\lambda = 0$ but any fixed λ could be considered instead.) From this condition it follows that the distribution of $(T_{WOD}, \mathbf{Q}_{WOD})$ has density $\boldsymbol{\nu} \cdot \mathbf{j}$ and is thus given by an operator-valued measure. However, the operators need not be positive, since linear combinations of wave functions in S may fail to be in S . Of course, an *operator-valued measure* is not sufficient, as the outcome of an experiment always has distribution given by a *positive-operator-valued measure*. Vona et al. established the following result: Let $(V$ and) Σ be given, and let S_0 be a set of normalized wave functions satisfying (41). Suppose there are $\Psi, \Phi \in S_0$ such that, with $\Psi_\pm := (\Psi \pm \Phi)/\|\Psi \pm \Phi\|$, we have that $\Psi_+ \in S_0$ while Ψ_- satisfies

$$\int_{\Sigma} d^2\mathbf{q} \boldsymbol{\nu}(\mathbf{q}) \cdot \mathbf{j}_{\Psi_-}(t_0, \mathbf{q}) < 0 \quad (42)$$

for some $t_0 \geq 0$. Then there is no POVM E_0 such that $\langle \Psi_0 | E_0(\cdot) | \Psi_0 \rangle$ agrees with $\mathcal{P}_{WOD, \Psi_0}$ for all $\Psi_0 \in S_0$. Moreover there are, at least for $V = 0$ and Σ a plane, Ψ and Φ in S as just described, thus establishing that there is at least one $\Psi_0 \in S$ for which $T_{WOD} \neq T_D$.

Our no-go result and that of Vona et al. both concern the non-existence of a POVM for \mathcal{P}_{WOD} , but for different classes of wave functions. Vona et al.'s is for wave functions obeying the current positivity condition (41) while ours concerns wave functions of the form $|\mathbf{n}\rangle \otimes \psi_0$ with any \mathbf{n} , most of which don't obey (41). The two results thus complement each other.

3.4 POVMs and Measurability

When investigating whether in Bohmian mechanics a quantity X (such as the first arrival time at Σ) is measurable, i.e., whether there is an experiment whose outcome is X —an experiment that measures X —we have used that it is a necessary condition that the probability distribution of X be given by a POVM. Is this condition also sufficient? No. Here is why. Suppose there is a POVM E such that we have, for the probability distribution of X when the system is in state Ψ , that $\text{Prob}_\Psi(X \in B) = \langle \Psi | E(B) | \Psi \rangle$ for all wave functions Ψ and all sets B . There then remain two possible obstacles to the existence of an experiment that measures X :

1. Maybe no appropriate apparatus leading to E can be built. To be sure, any POVM E on a system Hilbert space can be expressed in the form

$$E(B) = \langle \Phi | U^\dagger P(\zeta^{-1}(B)) U | \Phi \rangle, \quad (43)$$

where the inner product is a partial inner product in suitable apparatus variables, Φ is a suitable ready state of the apparatus, U a unitary time evolution of system and apparatus together, P the position PVM, and ζ the function that yields the outcome from the final configuration. Nonetheless, it may not be possible to physically realize such U, Φ, ζ .

2. The other obstacle is that, even if the POVM of some experiment equals E , the individual outcome Z of the experiment need not agree with X . After all, it is only guaranteed that the random variables X and Z have the same distribution, but that does not entail they are equal.

For example, consider a simple harmonic oscillator in $d \geq 2$ space dimensions, with Hamiltonian H . Then, since all eigenvalues of H are integer multiples of the ground state energy ε_0 for $d = 1$, the time evolution operator $\exp(-iHt/\hbar)$ yields the identity for $t = t_0 := 2\pi\hbar/\varepsilon_0$. Thus we have that the wave function Ψ_{t_0} of the system at time t_0 is the same as the wave function Ψ_0 at time 0, and hence similarly for the distributions of the (Bohmian) positions $\mathbf{Q}(t_0)$ and \mathbf{Q}_0 at those times. Thus the distribution of $X = \mathbf{Q}(t_0)$ is given by a POVM, the position POVM (given by the spectral projections for the position operator). Moreover, a measurement of \mathbf{Q}_0 is an experiment the distribution of whose result $Z (= \mathbf{Q}_0)$ is given by this POVM. But, nonetheless, this experiment measures, not $\mathbf{Q}(t_0)$, but \mathbf{Q}_0 instead. Indeed, if $d > 1$, the Bohmian position for the harmonic oscillator at time t_0 is in general different from the Bohmian position at time 0, even though the wave function at these times is the same.

On top of that, here is a trivial example, involving a POVM acting on a one-dimensional subspace: Regardless of whether or not X is measurable, and whatever the distribution \mathcal{P}_X of X for Ψ_0 , there trivially exists a POVM E , namely $E = \mathcal{P}_X I$, on the 1d subspace $\mathbb{C}\Psi_0$, that reproduces \mathcal{P}_X ; and there trivially exists an experiment with this POVM, viz., running a random number generator that creates a number Z with distribution \mathcal{P}_X . Since Z is generated independently of X , the two will in general not be equal.

3.5 Superluminal Signaling

It is a consequence of the spin dependence of \mathcal{P}_{WOD} in the setup of [4] that the hypothesis $\mathcal{P}_D = \mathcal{P}_{WOD}$ would imply the possibility of superluminal signaling. Here is how.

Procedure. For sending a bit $X \in \{0, 1\}$ from Alice to Bob at spacelike separation, a pair of particles is prepared (as for a Bell experiment) in the singlet state $\propto |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$, then one particle is transported to Alice's lab and one to Bob's. Depending on whether $X = 0$ or $X = 1$, Alice carries out a Stern-Gerlach experiment on her particle in either the z - or the x -direction. Because of the rotational invariance of and the perfect anti-correlations in the singlet state, this will result in a collapsed wave function of Bob's particle that is $|\pm \mathbf{z}\rangle$ (where each sign has probability 1/2) if $X = 0$ or $|\pm \mathbf{x}\rangle$ if $X = 1$.

Now Bob carries out an arrival-time experiment in the setup of [4] on his particle. As mentioned (see (19) and (20)), \mathcal{P}_{WOD} does not depend on the random sign; it does, however, depend on whether the spin direction is \mathbf{z} or \mathbf{x} , as shown by the blue and red curves in Figure 1. So, if \mathcal{P}_D were equal to \mathcal{P}_{WOD} , then Bob would obtain probabilistic information about X ; if Alice and Bob repeat the procedure 100 times with the same value of X , Bob could decide correctly with high probability whether his observations are distributed according to the blue or the red curve, and thus determine the message X that Alice wanted to send.

When one realizes that a hypothesis implies the possibility of superluminal signaling, one should become very skeptical of the hypothesis, not just because nobody has been able to experimentally achieve it, but also because there is a general proof that it is impossible in Bohmian mechanics [19, 1, 11, 12], [20, Sec.s 5.5.9 and 7.6.2]. On top of that, we can analyze the procedure in terms of POVMs; this is the content of the next paragraph.

Does It Work? We can easily specify the necessary and sufficient condition for the impossibility of superluminal signaling by means of an experiment of the sort just described, one allowing for the choice of any pair \mathbf{n}_1 and \mathbf{n}_2 of directions and not just \mathbf{z} and \mathbf{x} : It is that

$$\mathcal{P}_{D,\mathbf{n}_1} + \mathcal{P}_{D,-\mathbf{n}_1} = \mathcal{P}_{D,\mathbf{n}_2} + \mathcal{P}_{D,-\mathbf{n}_2} \quad (44)$$

for any pair of unit vectors \mathbf{n}_1 and \mathbf{n}_2 . After all, $\frac{1}{2}(\mathcal{P}_{D,\mathbf{n}} + \mathcal{P}_{D,-\mathbf{n}})$ is the detection time distribution that Bob will observe if Alice uses the direction \mathbf{n} . As we have noted already in (18), (44) must hold because \mathcal{P}_D must be given by a spin POVM.

4 T_{WID} versus T_D

It may sound innocent to assume that a detector should click when and where the particle actually arrives, which would imply that

$$T_D = T_{WID}. \quad (45)$$

And in classical mechanics, this is the case because the interaction between particle and detector is local. But in Bohmian mechanics, the structure is more involved because, although the interaction term in the Hamiltonian is still local, it directly affects only the (entangled) wave function, which in turn affects the motion of the particles; so it is not obvious whether (45) is correct. More precisely, it is not clear whether there exists a device D —a potential detector—for which (45) holds. It might well be that for any potential detector D , T_{WID} is like the spin state $|\mathbf{n}\rangle$, or the wave function, in that it is measurable, neither by D itself, so that (45) is violated, nor by any other device. After all, T_{WID} represents the Bohmian arrival time of a particle interacting with a certain environment (the detector), while T_D is the outcome, registered by the device, produced by the interaction, which is conceptually and physically quite different. Therefore, the comparison is similar to the earlier comparison between T_{WOD} and T_D , with the distinction between the comparisons being the presence or absence of a certain

environment. (Nonetheless, it could be the case that there is a fundamental difference between measurement of T_{WID} and T_{WOD} . The latter can in general not be measured even approximately, whereas for the former, though there may be limitations, it might well be that approximate measurement is always possible; see also [7, 21].)

4.1 Comparison to Position Measurements

It should be observed that this state of affairs is not confined to time measurements alone; even when it comes to position measurements, a discrepancy may arise between the actual position in the presence of a detector and the position recorded by the detector. As emphasized in the introduction, even the assumption that $\mathbf{Q}_{WID} = \mathbf{Q}_D$ should not be accepted without scrutiny. In Bohmian mechanics there is no reason for exempting position or time from adhering to the principle that quantum measurement outcomes are the joint result of the combined behaviors of both the system and the apparatus, a principle which is indeed a direct consequence of the Bohmian analysis of quantum measurements [8].

We should also note that for most potential measurements of a quantity X , there is no difference between X_{WOD} ($= X$) and X_{WID} , so that the measurability of X boils down to whether $X_{WID} = X_D$ (for some device D). This will be so whenever the time evolution plays no role in the definition of X , unlike for the case of arrival times. For example, this is the case when X is the position of a particle, in quantum state Ψ , at a given time t , say, time $t = 0$. The presence or absence of a detector is not relevant to the meaning of X , though it is of course relevant to the position of the particle at later times.

So let's consider the simple case when X is the position of a particle at a given time. It is generally believed, with good reason, that this can be measured with unlimited precision. But even in this case, one could take a step further and argue that the measurability of position is a contingent feature of the non-relativistic framework within which the theory is formulated. Consider, for instance, a relativistic scenario, specifically involving the measurement of the position of a Dirac electron. Suppose, as is widely believed, that in the one-particle Hilbert space of the electron, comprising exclusively positive energy states, there is an inherent limitation on how narrow the wave function can be. This restriction is intuitively understandable since a delta function cannot be exclusively formed with positive energy wave functions. Then the minimal width of the wave function (and thus, by Born's rule, the minimal possible inaccuracy of an experimenter's knowledge of the electron's position) would occur on a scale approximately corresponding to its Compton wavelength. Consequently, any putative detector probing below that scale would register a specific position that typically deviates systematically from the actual position of the electron.

4.2 Different Equations of Motion

It has long been known that different equations of motion, regarded here as different possible laws of nature, can all leave the Born distribution $|\Psi|^2$ equivariant while leading to different trajectories. Any such law of motion, as part of a universal theory, leads

to the same probability distribution as in Bohmian mechanics and orthodox quantum mechanics for any experimental outcome. That is because we should apply this law of motion to the joint configuration of both the observed system and the apparatus to conclude that the joint configuration of the apparatus and the system at any time τ after the experiment has begun, say at time 0, will be $|\Psi_\tau|^2$ distributed; since it can be arranged that the outcome can be read off from the apparatus configuration for some $\tau > 0$, the distribution of the outcome must agree with the one obtained from Bohmian mechanics or orthodox quantum mechanics. That is, the theory with this law of motion is empirically equivalent to Bohmian mechanics and orthodox quantum mechanics (even though it need not lead to the same individual outcome as Bohmian mechanics when starting from the same initial configuration).

Of course, this consideration includes the distribution \mathcal{P}_D of the detection time of a particle in an arrival-time experiment. But when applied to the particle alone, a different law of motion leads to generically different arrival times T_{WOD} on the surface Σ , and in fact to different distributions \mathcal{P}_{WOD} . Thus, for most of such laws of motion, $\mathcal{P}_D \neq \mathcal{P}_{WOD}$.

4.2.1 Examples

Here are four different examples of such laws of motion:

- For the Bohmian equation of motion for the position $\mathbf{Q}(t)$ of a non-relativistic spin- $\frac{1}{2}$ particle, there are the possibilities indexed by $\lambda \in \mathbb{R}$ in (5). As mentioned already in Section 2, \mathcal{P}_{WOD} is independent of \mathbf{n} for $\lambda = 0$ but not for $\lambda \neq 0$.
- Nelson’s stochastic mechanics [16, 13] posits a stochastic particle motion given by a diffusion process. The process belongs to a 1-parameter family indexed by the diffusion constant $\sigma \geq 0$ [20, App. A.3], and Bohmian mechanics is included in the family at $\sigma = 0$. The theory is implausible for large σ , but \mathcal{P}_{WOD} will already change significantly for rather small σ .
- Deotto and Ghirardi [6] described further ODEs as alternatives to Bohm’s. The formulas would not seem convincing as laws of nature, but they do ensure equivariance.
- Colin, Wiseman, and Struyve [3, 18, 15] considered a $|\Psi|^2$ distributed Markov process for Dirac wave functions Ψ , called the “zig-zag process” and based on introducing a further hidden variable in $\{+1, -1\}$ for each particle representing “handedness.” In the non-relativistic limit for a wave function factorizing into a spinor and a space part, $\Psi = |\mathbf{n}\rangle \otimes \psi$, the particle carries out, on top of a slow Bohm-like motion, a fast (speed c), random, back-and-forth motion in direction $\pm \mathbf{n}$ (see [15] for pictures).

4.2.2 Implications

The existence of different equivariant laws of motion shows, in fact, that (45) can fail: While the joint wave function Φ_τ of particle and detector (and records) at a late

time τ fixes \mathcal{P}_D , different equivariant laws of motion will in general lead to different trajectories $Q(t)$ of particle and detector together, and we see no reason why in general these different trajectories should have equal T_{WID} or \mathcal{P}_{WID} . For the zig-zag process, Nelson's stochastic mechanics, and its extensions with other diffusion constants σ , it seems rather obvious from the stochastic fluctuations that \mathcal{P}_{WID} will be different from what it is in Bohmian mechanics, for example allowing for many much smaller arrival times.

5 Conclusion

Das and Dürr hoped that the experiment they studied could be realized and would yield outcome statistics in agreement with the \mathcal{P}_{WOD} that they computed. Our results show that Bohmian mechanics predicts otherwise. We have thus shown that the quantity T_{WOD} is not measurable, that is, that there is no experiment whose outcome conveys the time at which the particle would have reached the surface Σ in the absence of detectors. The upshot of our considerations is that arrival times are less accessible than some might imagine.

Acknowledgments. We thank Dustin Lazarovici and Ward Struyve for useful discussions.

References

- [1] J.S. Bell: The theory of local beables. *Epistemological Letters* **9**: 11 (1976)
Reprinted as chapter 7 of [2].
- [2] J.S. Bell: *Speakable and unspeakable in quantum mechanics*. Cambridge University Press (1987)
- [3] S. Colin and H.M. Wiseman: The zig-zag road to reality. *Journal of Physics A: Mathematical and Theoretical* **44**: 345304 (2011) <http://arxiv.org/abs/1107.4909>
- [4] S. Das and D. Dürr: Arrival Time Distributions of Spin-1/2 Particles. *Scientific Reports* **9**: 2242 (2019) <http://arxiv.org/abs/1802.07141>
- [5] M. Daumer, D. Dürr, S. Goldstein, and N. Zanghì: On the quantum probability flux through surfaces. *Journal of Statistical Physics* **88**: 967–977 (1997). Reprinted in [9]. <http://arxiv.org/abs/quant-ph/9512016>
- [6] E. Deotto and G.C. Ghirardi: Bohmian Mechanics Revisited. *Foundations of Physics* **28**: 1–30 (1998) <http://arxiv.org/abs/quant-ph/9704021>
- [7] V. Dubey, C. Bernardin, and A. Dhar: Quantum dynamics under continuous projective measurements: non-Hermitian description and the continuum space limit. *Physical Review A* **103**: 032221 (2021) <http://arxiv.org/abs/2012.01196>

- [8] D. Dürr, S. Goldstein, and N. Zanghì: Quantum Equilibrium and the Role of Operators as Observables in Quantum Theory. *Journal of Statistical Physics* **116**: 959–1055 (2004) Reprinted in [9]. <http://arxiv.org/abs/quant-ph/0308038>
- [9] D. Dürr, S. Goldstein, and N. Zanghì: *Quantum Physics Without Quantum Philosophy*. Heidelberg: Springer-Verlag (2013)
- [10] D. Dürr and S. Teufel: *Bohmian Mechanics*. Heidelberg: Springer (2009)
- [11] P.H. Eberhard: Bell’s theorem and the different concepts of locality. *Il Nuovo Cimento B* **46**: 392–419 (1978)
- [12] G.C. Ghirardi, R. Grassi, A. Rimini, and T. Weber: Experiments of the EPR Type Involving CP-Violation Do not Allow Faster-than-Light Communication between Distant Observers. *Europhysics Letters* **6**: 95–100 (1988)
- [13] S. Goldstein: Stochastic Mechanics and Quantum Theory. *Journal of Statistical Physics* **47**: 645–667 (1987)
- [14] S. Goldstein, R. Tumulka, and N. Zanghì: On the Spin Dependence of Detection Times and the Nonmeasurability of Arrival Times. *Scientific Reports* **14**: 3775 (2024) <http://arxiv.org/abs/2309.11835>
- [15] C. Maes, K. Meerts, and W. Struyve: Diffraction and interference with run-and-tumble particles. *Physica A* **598**: 127323 (2022) <http://arxiv.org/abs/2106.09568>
- [16] E. Nelson: *Quantum Fluctuations*. Princeton University Press (1985)
- [17] S. Römer, D. Dürr, and T. Moser: Asymptotic Behavior of Bohmian Trajectories in Scattering Situations. *Journal of Physics A: Mathematical and General* **38**: 8421–8443 (2005) <http://arxiv.org/abs/math-ph/0505074>
- [18] W. Struyve: On the zig-zag pilot-wave approach for fermions. *Journal of Physics A: Mathematical and Theoretical* **45**: 195307 (2012) <http://arxiv.org/abs/1201.4169>
- [19] K.S. Tausk: *A Medida na Mecânica Quântica*. Ph.D. dissertation, Faculdade de Filosofia, Ciências e Letras, Universidade de São Paulo (1967)
- [20] R. Tumulka: *Foundations of Quantum Mechanics*. Heidelberg: Springer (2022)
- [21] R. Tumulka: On a Derivation of the Absorbing Boundary Rule. *Physics Letters A* **494**: 129286 (2024) <http://arxiv.org/abs/2310.01343>
- [22] N. Vona, G. Hinrichs, and D. Dürr: What does one measure when one measures the arrival time of a quantum particle? *Physical Review Letters* **111**: 220404 (2013) <http://arxiv.org/abs/1307.4366>