

Enhanced Lieb-Robinson bounds for a class of Bose-Hubbard type Hamiltonians

Tomotaka Kuwahara^{*1,2} and Marius Lemm^{†3}

¹Analytical quantum complexity RIKEN Hakubi Research Team, RIKEN Cluster for Pioneering Research (CPR)/ RIKEN Center for Quantum Computing (RQC), Wako, Saitama 351-0198, Japan

²PRESTO, Japan Science and Technology (JST), Kawaguchi, Saitama 332-0012, Japan

³Department of Mathematics, University of Tübingen, 72076 Tübingen, Germany

May 7, 2024

Abstract

Several recent works have considered Lieb-Robinson bounds (LRBs) for Bose-Hubbard-type Hamiltonians. For certain special classes of initial states (e.g., states with particle-free regions or perturbations of stationary states), the velocity of information propagation was bounded by a constant in time, $v \leq C$, similarly to quantum spin systems. However, for the more general class of bounded-density initial states, the first-named author together with Vu and Saito derived the velocity bound $v \leq Ct^{D-1}$, where D is the spatial lattice dimension. For $D \geq 2$, this bound allows for accelerated information propagation. It has been known since the work of Eisert and Gross that some systems of lattice bosons are capable of accelerated information propagation. It is therefore a central question to understand under what conditions the bound $v \leq Ct^{D-1}$ can be enhanced. Here, we prove that additional physical constraints, translation-invariance and a p -body repulsion of the form n_x^p with $p > D + 1$, lead to a LRB with $v \leq Ct^{\frac{D}{p-D-1}}$ for any initial state of bounded energy density. We also identify examples of quantum states which show that no further enhancement is possible without using additional dynamical constraints.

1 Introduction

Lieb-Robinson bounds (LRBs) [LR72] provide a speed limit on the propagation of quantum information in a quantum many-body lattice system with local interactions. The standard LRBs, see [LR72, Has04, NS06], concern two local bounded operators (“observables”) O_1, O_2 and are of the form

$$\|[O_1(t), O_2]\| \leq C \|O_1\| \|O_2\| e^{C(vt-d(O_1, O_2))}, \quad (1.1)$$

where $O_1(t) = e^{itH} O_1 e^{-itH}$ is the Heisenberg time evolution generated by the Hamiltonian H . Here, $d(O_1, O_2) = \text{dist}(\text{supp } O_1, \text{supp } O_2)$ and the support of an observable O is defined as the smallest lattice region X such that O acts as the identity on X^c , i.e., $O = O_X \otimes \text{Id}_{X^c}$.

^{*}tomotaka.kuwahara@riken.jp

[†]marius.lemm@uni-tuebingen.de

The commutator version of the LRB (1.1) is equivalent to the following locality bound on the spreading of a single local operator O with $\text{supp } O = X_0$, under the Heisenberg dynamics,

$$\|O(t) - O(t, X_0[R])\| \leq C\|O\|e^{C(vt-R)}, \quad t, R \geq 1. \quad (1.2)$$

Here, $O(t, X_0[R])$ is a local operator, supported on the R -enlargement of the initial support $X_0 = \text{supp } O$

$$X_0[R] := \{i \in \Lambda \mid \text{dist}(i, X_0) \leq R\}, \quad R > 0. \quad (1.3)$$

Thus, (1.2) says that the time evolution $O(t)$, which is generally supported on the entire lattice, is well-approximated by the local observable $O(t, R)$, which is supported only on $X_0[R]$, in the space-time region where $R > vt$. The locality version (1.2) of the LRB is the one that is relevant for most, but not all, applications. A common explicit choice of the local approximation $O(t, X_0[R])$ is to take the time evolution with respect to the restricted subregion Hamiltonian^{*1}

$$O(t, X_0[R]) = e^{itH_{X_0[R]}} O e^{-itH_{X_0[R]}}. \quad (1.4)$$

In either formulation of the LRB — commutator bound (1.1) or locality bound (1.2) — the key quantity that characterizes the quantum information propagation is $v \geq 0$. This bound is called the Lieb-Robinson velocity. In the standard LRB, which holds for many-body systems with *bounded* interactions, v is proportional to the operator norm of the local interaction and independent of the time t . Consequently, the standard LRB [LR72, Has04, NS06] is only meaningful for systems with bounded interactions such as quantum spin systems (and also lattice fermions [NSY18]).

LRBs have blossomed into a rich subject that connects mathematical physics with condensed matter physics, quantum information science, and high-energy physics. This success story started in the early 2000s [Has04, HW05, BHV06, NS06, NOS06, Has07a, Has07b, NS07, BMNS12, KGK⁺14], kicked-off by the highly influential realization of Hastings that LRBs, which are dynamical bounds, are surprisingly effective tools to study equilibrium properties, e.g., clustering of correlations [Has04, NS06], the area law for entanglement entropy [Has07a] and topological quantum phases [HW05, BMNS12]. Since then, it has been realized that LRBs are among the few robust, effective, and versatile tools for analyzing quantum many-body systems in and out of equilibrium. More recently, they have also been used to limit information scrambling in the theory of quantum many-body chaos which is relevant for black hole physics and quantum information science [LSH⁺13, RS16, CL19, KS21a]. For further background on LRBs, we refer to the reviews [NS10, GE16, CLY23].

1.1 Bosonic propagation bounds

Since the standard LRBs require bounded interactions, they do not give meaningful bounds for lattice bosons or systems in continuous space.

Special bosonic systems such as perturbations of harmonic oscillator systems have been considered [CSE08, NRSS09, WCP15, WP16]. However, the results covering the paradigmatic *Bose-Hubbard Hamiltonian* [SHOE11, WH20] were quite restricted, and proving robust LRBs for this Hamiltonian, in particular for a broad class of initial states, was a major methodological barrier. In the last three years, this barrier was broken and bosonic LRBs for Bose-Hubbard Hamiltonians have

^{*1}Another possibility due to [BHV06] is to take $O(t, X_0[R])$ as the partial trace (a.k.a. conditional expectation) over the complement region $X_0[R]^c$.

been derived by three different research groups by three different methods [KS21b, FLS22b, YL22, KVS24, SZ22, LRS⁺23], which we summarize below. Similar bounds for systems in continuous space have also recently been studied by related methods [GNRS20, APSS21, BFLS22, BFL⁺23, HLS23].

Part of the recent interest from the physics community in LRBs for Bose-Hubbard type Hamiltonians stems from the fact that the latter can be experimentally realized, finely tuned, and measured with high fidelity, in tabletop experiments with ultracold quantum gases in optical lattices [DSH⁺02, BDZ08, GF08]. These lattice bosons provide some of the early examples of quantum simulators [Jak04, BDN12, LSA12, GB17, YSO⁺20, KN21, SSH⁺23]. In this context, bosonic LRBs have been experimentally observed since about 10 years ago [BPCK12, CBP⁺12, Che22] and their validity can provide important performance bounds on the use of quantum simulation and quantum information protocols [BHV06, KGE14, EW17, FLS22b, KVS24, SZ22, LRS⁺23].

We briefly summarize the state-of-the-art bosonic LRBs for Bose-Hubbard type Hamiltonians of the form

$$H = H_0 + V, \quad H_0 = \sum_{i,j \in \Lambda} J_{i,j} (b_i^\dagger b_j + b_j^\dagger b_i), \quad V = \sum_{i,j \in \Lambda} \tilde{v}(n_i, n_j), \quad (1.5)$$

where $\Lambda \subset \mathbb{Z}^D$ is a finite set. The term H_0 represents boson hopping and V represents boson-boson density-dependent interaction. The prototypical case is that of nearest-neighbor hopping $J_{i,j} = \delta_{\langle i,j \rangle}$ (where $\langle i,j \rangle$ means i and j are nearest neighbors) and quadratic repulsion $V = \sum_{i \in \Lambda} (n_i(n_i - 1) - \mu n_i)$.

Since lattice bosons involve unbounded interactions, it is natural that the bosonic LRBs do not involve operator norms (which are, after all, worst-case quantities) and instead control quantum information propagation for certain classes of “well-behaved” initial states ρ . In this vein, the following types of initial states have been considered with qualitatively different LRBs. The works [SHOE11, KS21b, YL22, FLS22b, KVS24, LRS⁺23] proved a bosonic LRB with v independent of time for various types of special initial states. More precisely, [SHOE11] covered states with all particles in a fixed finite region, [KS21b] covered perturbations of stationary states, [YL22] covered states $\rho = e^{-\mu N}$, and [FLS22b, SZ22, LRS⁺23] covered states with general particle-free subregions and long-range interactions. Most relevant for our purposes is the work [KVS24] which proved a bosonic LRB with $v \sim t^{D-1}$ for initial states with suitably bounded particle density. (As one would expect, all these bounds have in common that the LR velocity v is independent of system size, which is necessary to have a meaningful bound in the thermodynamic limit. Even this arguably basic requirement is far from trivial for bosons [WH20].)

Any LRB is merely an upper bound on information transport. It is always of interest to complement it by a qualitatively similar lower bound, i.e., to specify situations when the transfer is indeed of a prescribed speed. This question is particularly pressing in the situation considered in [KVS24] because there, for $D \geq 2$, the velocity bound $v \sim t^{D-1}$ grows with time. This bound in principle allows for *information acceleration* by which we mean that any LRB, say, $\text{Tr}(\rho(O(t) - O(t, R))) \lesssim f(R - vt)$, with f decaying at large arguments, must have v *growing* with time. It is unclear if and when this information acceleration truly happens for Bose-Hubbard Hamiltonians in bounded-density initial states, which is a glaring gap in our understanding of bosonic transport even at a qualitative level. In view of this, we put forward the following question which is open, fundamental, and pressing.

Question 1: Consider a time-independent Bose-Hubbard type Hamiltonian such as (1.5), prepared in a bounded-density initial state ρ . Does it display a finite speed of information propagation?

This question is also open for the case of the standard Bose-Hubbard Hamiltonian where the hopping is nearest-neighbor and the interaction is quadratic and on-site.

Naive intuition would suggest that information acceleration cannot occur in a closed quantum system governed by unitary dynamics that is not subjected to any external forcing. However, this naive intuition is provably incorrect. Indeed, there exist *explicit examples of simple, time-independent, and translation-invariant bosonic Hamiltonians which rigorously display information acceleration*, meaning that no LRB can hold with v independent of time. Indeed, Eisert and Gross [EG09] exhibited a one-dimensional, translation-invariant, time-independent lattice boson Hamiltonian where any LRB must have $v \sim e^{Ct}$, i.e., information accelerates exponentially. While this Hamiltonian is *not* of Bose-Hubbard type (1.5), it impressively shows that information acceleration for bosons is a realistic possibility that has to be confronted in a model-dependent way, unlike for quantum spin systems. In the words of [EG09]: “Since we cannot rule out the presence of accelerating excitations by imposing natural assumptions, one must not take the existence of a finite speed of sound in any bosonic model for granted.”

The work [KVS24] also exhibits a protocol for a time-dependent Bose-Hubbard type Hamiltonian which rigorously propagates quantum information with $v \sim t^{D-1}$. This has also sparked heuristic arguments that the scaling $v \sim t^{D-1}$ could be optimal for time-independent Bose-Hubbard type Hamiltonians as well. Hence, a different, weaker variant of Question 1 is to ask to what extent the LRB for bounded initial states in [KVS24] is in fact optimal.

Question 1’: Consider a time-independent Bose-Hubbard type Hamiltonian such as (1.5), prepared in a bounded-density initial state ρ . Can the LRB from [KVS24], which gives $v \leq Ct^{D-1}$, be enhanced under physically meaningful assumptions?

A particularly natural condition of interest is the translation invariance of the system Hamiltonian. This intuitively limits the possibility of the boson concentration on some particular regions, which is a central puzzle piece in the explicit protocol displaying the accelerating information transfer. In fact, naive intuition might suggest that the translation invariance immediately leads to bounded local particle numbers and thus a finite Lieb-Robinson velocity: since a translation-invariant state with large boson occupation at a site must also have an equal chance of having large boson occupation at all other sites which should lead to a combinatorial suppression of the possibility of such problematic states. Unfortunately, this naive combinatorial picture is incorrect, because quantum mechanics allows for the phenomenon of “superpositions of ‘bad’ states”. In Appendix A, we construct an explicit family of translation-invariant states which have significant boson concentration (while also satisfying the energetic bounds we use here). These examples show that there are limitations to use only certain dynamical constraints (including translation-invariance).

In this paper, we provide a positive answer to Question 1’. We work under two additional physical assumptions: (i) strong on-site repulsion and (ii) translation-invariance. We emphasize that translation-invariance alone is not sufficient to obtain the improvement. We view our result as another puzzle piece in the quest to understand the surprising richness of bosonic transport behavior, a richness that was to our knowledge first emphasized by Eisert and Gross in [EG09].

1.2 Statement of result

We now state the main example of the kind of result we prove. It establishes two enhanced LRBs which are qualitatively different depending on how the approximation error $O(t) - O(t, R)$ is measured when tested with respect to the initial state ρ ; either in trace norm, $\|A\|_1 = \text{tr}|A|$, or in expectation. We write $\mathcal{S}(\mathcal{H})$ for the set of quantum states on a Hilbert space \mathcal{H} , i.e., $\rho \in \mathcal{S}(\mathcal{H})$ iff ρ is a positive semi-definite trace class operator of trace equal to 1. Recall Definition (1.4) of the localized time evolution $O(t, R)$ and Definition (1.3) of the enlarged set $X[R]$. We denote the local particle number on set X by $n_X = \sum_{i \in \Lambda} n_i$. Below, we introduce suitable time-dependent velocities $v_1(t)$ and $v_{\text{ex}}(t)$ in order to compare with other results.

Theorem 1.1 (Main example of results). *Consider a nearest-neighbor, translation invariant Bose-Hubbard Hamiltonian (1.5) with $J_{i,j} = \bar{J}\delta_{i \sim j}$ on the discrete D -dimensional torus $\Lambda_L = ((-L/2, L/2] \cap \mathbb{Z})^d$.*

Let $p \in (1, \infty)$ and consider a local interaction of the form

$$V = \sum_{i \in \Lambda} [n_i^p + v(n_i)]. \quad (1.6)$$

We assume that there exist $c_V, \epsilon > 0$ such that $|v(n)| \leq c_V(n+1)^{p-\epsilon}$ for all $n \geq 0$.

Let $\rho \in \mathcal{S}(\mathcal{F}(\ell^2(\Lambda_L)))$ be a translation-invariant quantum state that has bounded expected energy density, i.e.,

$$\frac{\text{tr}(\rho H)}{|\Lambda_L|} = E_\rho < \infty. \quad (1.7)$$

Let $X_0 \subset \Lambda$. There exists a constant $C = C(D, J, c_V, \epsilon, E_\rho, X_0)$ such that the following holds for every bounded observable O on $\mathcal{F}(\ell^2(\Lambda_L))$ with $\text{supp } O = X_0 \subset \Lambda_L$ that commutes with n_{X_0} .

(I) *Suppose that $p > 2D + 2$ and set $v_1(t) = t^{\frac{D}{p/2-D-1}}$. Then*

$$\|\rho[O(t) - O(t, X_0[R])]\|_1 \leq C\|O\| \left(\frac{v_1(t)t}{R}\right)^{p/2-D-1}, \quad t, R \geq 1. \quad (1.8)$$

(II) *Suppose that $p > D + 1$ and set $v_{\text{ex}}(t) = t^{\frac{D}{p-D-1}}$. Then*

$$|\text{tr}[\rho(O(t) - O(t, X_0[R]))]| \leq C\|O\| \left(\frac{v_{\text{ex}}(t)t}{R}\right)^{p-D-1}, \quad t, R \geq 1. \quad (1.9)$$

As usual in statistical physics, the key point is that the constant C is independent of system the size parameter L . Theorem 1.1 is applicable to any initial quantum state whose energy density per volume E_ρ is bounded uniformly in ρ . This is a fairly basic requirement.

Theorem 1.1 stems from the more general Theorems 2.4 and 2.5. These theorems offer results that are applicable to a wider variety of Bose-Hubbard Hamiltonians (1.5) (which don't necessarily have to be uniform across space) as long as they involve interactions that are limited in range and adhere to a specific condition over time: $\text{tr}[\rho_0(t)n_i^p] \leq C_p$ for all $i \in \Lambda$. We then demonstrate that this condition is satisfied when two physical criteria are met: (i) translation-invariance and (ii) strong on-site repulsion, which then imply Theorem 1.1 as a consequence. This also makes it clear that the assumption in Theorem 1.1 that the initial observable O is particle-number preserving can

be substantially relaxed — it suffices to assume control on how many particles can be created by O , cf. Definition 2.3.

We compare and contrast our results to those of [KVS24].

- As in [KVS24], we are able to cover a very broad class of initial states, namely those of finite particle density. This includes the experimentally most relevant class of Mott states [BDZ08, CBP⁺12, Che22].
- Theorem 1.1 establishes an LRB which bounds the velocity of information propagation by $v_1(t) = t^{\frac{D}{D+1-p/2}}$ respectively $v_{\text{ex}}(t) = t^{\frac{D}{D+1-p}}$ depending on how the size of the deviation $O(t) - O(t, X_0[R])$ is measured (either in trace norm or in quantum expectation). Note that $\lim_{p \rightarrow \infty} v_1(t) = \lim_{p \rightarrow \infty} v_{\text{ex}}(t) = 1$, so for sufficiently large p , we obtain an approximately constant maximal velocity bound, closer to the case of quantum spin systems. This is a manifestation of the fact that the system gets increasingly rigid as p grows.
- The smallest p -values such that the velocities $v_1(t)$ and $v_{\text{ex}}(t)$ scale more favorably than the bound $v \sim t^{D-1}$ from [KVS24] can be formulated as the conditions $p > 2D + 2 + \frac{2D}{D-1}$ (for the trace norm) or $p > D + 1 + \frac{D}{D-1}$ (for the expectation). For the physical dimensions $D = 2$ and $D = 3$, the smallest integer value of p for which our result yields an improved scaling compared to [KVS24] is $p = 6$. That is, in physical terms, we prove that six-body boson-boson repulsion yields qualitatively slower information transport than what is expected from [KVS24]. This also shows that the “accumulation scenario” in Figure 4a of [KVS24], which, as we recall, produces information propagation with $v \sim t^{D-1}$, is excluded in our setting. (We emphasize that our result is not in any contradiction to the rigorous protocol in [KVS24]. The reason is that the protocol requires a special time-dependent Bose-Hubbard Hamiltonian that neither enjoys strong repulsion nor translation-invariance.)

Under the assumptions of Theorem 1.1, we also obtain an improvement of the particle propagation bound for arbitrary moments (Main result 1 in [KVS24]) by an interpolation argument; see Proposition 2.7.

1.3 Discussion

We comment on the role of assumptions of (i) strong repulsion and (ii) translation invariance and the connection of our work to [KVS24] at a conceptual and technical level.

Firstly, local interactions expressed as n_i^p with an integer exponent p greater than 2 are physically derived from the local repulsion between multiple particles, approximated through the tight-binding model. You can find more details in sources like [MKM20]. The most significant case for physical applications is when $p = 2$, representing two-body interactions in the typical Bose-Hubbard Hamiltonian. Our findings do not extend to this case. On the one hand, to further improve our result to include $p = 2$, we anticipate the need for a more intricate analysis of how energy flows locally and how higher-order particle correlations develop dynamically. Dealing with these aspects of local dynamics will necessitate the introduction of new insights and ideas. On the other hand, it is by now well-established that bosonic information propagation is subtle and can produce rapidly accelerating information propagation in some cases [EG09, KVS24], it is not clear at all, even at a physical level, if the LRB from [KVS24] can be improved for the original Bose-Hubbard model with $p = 2$ in the most interesting regime of $D \leq 3$ for a broad class of initial states.

Second, concerning *translation-invariance*, we remark that naive combinatorial heuristics suggest that translation-invariance by itself could be enough to obtain an LRB with $v \sim 1$. These heuristics go as follows: Note first that large speed v comes from local boson accumulation. Now consider a time-evolved state that is “bad” in the sense that it exhibits extreme boson accumulation at a given site. By translation-invariance, it must be accompanied by many other bad states, so that when looking at a fixed site the total state has a small chance of having boson accumulation nearby, which should in turn yield $v \sim 1$. Unfortunately, this combinatorial heuristic is false. This can be seen by constructing translation-invariant states by superposing states with periodic boson accumulation on highly occupied lines. This shows that, *unfortunately, translation-invariance is not as useful as one might naively hope based on classical probabilistic intuition*. We give examples of such states in Appendix A. This is an instance of the more general phenomenon that “superpositions of bad states can behave well”. This genuinely quantum phenomenon is at the heart of the challenge of using combinatorial-probabilistic arguments in the quantum many-body context. In fact, excluding a similar “superpositions of bad states,” kind of scenario also constitutes the main technical work in [KVS24]. There, the main dynamical constraint used is that the speed of particle transport is uniformly bounded in time (see also [FLS22a, VKS23, LRS⁺23, LRZ23] for more general bounds of this type, e.g., for long-range hopping). More precisely, one shows that starting from a bounded-density initial state, within time t the number of particles on a $1D$ subregion is at most t^{D-1} and this is ultimately the reason for $v \sim t^{D-1}$. By considering this scenario, one sees that particle propagation bounds alone are insufficient to improve over $v \sim t^{D-1}$. Our result shows that translation-invariance and global energy conservation, for a sufficiently strong repulsion, are sufficient for an improvement. On a technical level, we replace the exponential but time-dependent bound on the local particle number in [KVS24] with a polynomial time-independent bound that we derive from translation-invariance and energy conservation. Then we draw on the idea of concatenating and short-time LRBs from [KS21b] and certain estimates on comparison with truncated dynamics from [KVS24]. For this, we have to take extra care to handle the slower decay which is polynomial in our case. In addition, we observe that the bound on the expectation value can be improved compared by re-ordering certain expansion terms in the Duhamel expansion of the truncated dynamics.

1.4 Organization of the paper

In Section 2, we introduce the formal setup and state the main results, Theorems 2.4 and 2.5 which imply Theorem 1.1 as an easy corollary. We also state an improved bound on the particle propagation, Proposition 2.7, that may be of independent interest.

Section 3 contains the overall proof strategy for Theorems 2.4 and 2.5. It shows how these theorems are reduced to suitable short-time LRBs (Subtheorems 3.1 and 3.2) by a suitable concatenation procedure. Subtheorem 3.1 is proved in Section 4 and Subtheorem 3.2 is proved in Section 5. We summarize our findings and give an outlook in Section 6.

In Appendix A, we construct translation-invariant quantum states with bounded energy density and significant boson concentration. These examples serve to show that the propagation bounds we prove are sharp conditional on the kinds of dynamical constraints that we use (translation-invariance and conservation of the first, even of the second, moment of the Hamiltonian). Further improvements will therefore need to leverage additional dynamical constraints and likely a much finer understanding of the energy flow on microscopic length scales that is generated by the many-body dynamics.

We heavily rely on several technical results from [KS21b, KVS24] as important ingredients to prove our enhanced LRB. This is in some sense natural given that we seek to enhance an already rather non-trivial bound. Whenever we use results from these works, we include their precise statement, but we do not give the proofs in order to keep the paper at a moderate length.

2 Setup and main results

2.1 Notation and setup

Let Λ be a finite graph. We also write Λ for the set of total sites and for an arbitrary partial set $X \subseteq \Lambda$, we denote its cardinality by $|X|$. We also denote the complementary subset and the surface subset of X by $X^c := \Lambda \setminus X$ and $\partial X := \{i \in X | d_{i, X^c} = 1\}$, respectively.

For arbitrary subsets $X, Y \subseteq \Lambda$, we define $d_{X, Y}$ as the shortest path length on the graph that connects X and Y and if $X \cap Y \neq \emptyset$, we set $d_{X, Y} = 0$. When X is composed of only one element (i.e., $X = \{i\}$), we denote $d_{\{i\}, Y}$ by $d_{i, Y}$ for simplicity. We also define the diameter $\text{diam}(X)$ in a non-standard, but convenient way as follows:

$$\text{diam}(X) := 1 + \max_{i, j \in X} (d_{i, j}). \quad (2.1)$$

For a subset $X \subseteq \Lambda$, recall that we define the extended subset $X[r] := \{i \in \Lambda | d_{X, i} \leq r\}$ where $X[0] = X$ and $r \in \mathbb{R}^+$. We write \preceq for operator inequalities.

Assumption 2.1 (*D-dimensional graph*). *We assume that the graph Λ is D-dimensional, in the sense that its surface growth is controlled via a lattice parameter $\gamma \geq 1$ by*^{*2}

$$|\partial(i[\ell])| \leq \gamma \ell^{D-1}, \quad \text{for all } \ell \geq 0. \quad (2.2)$$

We remark that the surface growth bound (2.2) implies the volume growth bound $|i[\ell]| \leq (\gamma + 1)\ell^D$. The reverse implication does not hold in general; see, e.g., [Tes07, Section 2.3].

We consider the bosonic Fock space $\mathcal{F}(\ell^2(\Lambda))$ with bosonic creation and annihilation operators $\{b_i^\dagger, b_i\}_{i \in \Lambda}$ satisfying the canonical commutation relations. We write $n_i = b_i^\dagger b_i$ for the particle number operator at site $i \in \Lambda$ and $\mathcal{N} = \sum_{i \in \Lambda} n_i$ for the total particle number operator.

Let $p \in (1, \infty)$. On $\mathcal{F}(\ell^2(\Lambda))$, we consider general Bose-Hubbard-type Hamiltonians of the form

$$\begin{aligned} H &= H_0 + V, \\ \text{where } H_0 &:= \sum_{\langle i, j \rangle} J_{i, j} (b_i b_j^\dagger + b_i^\dagger b_j), \quad |J_{i, j}| \leq \bar{J}, \\ V &:= f(\{n_i\}_{i \in \Lambda}), \end{aligned} \quad (2.3)$$

where we recall that $\langle i, j \rangle$ means i and j are nearest neighbors.

Assumption 2.2 (*Finite-range interaction V*). *We assume that the interaction V is finite-range, i.e., there exists a constant $k \geq 1$ such that*

$$V = \sum_{\substack{X \subseteq \Lambda: \\ \text{diam}(X) \leq k}} f_X(\{n_i\}_{i \in X}). \quad (2.4)$$

^{*2}It is trivial that such a constant $\gamma < \infty$ exists for any finite graph for any D , but the point is that the bounds we prove will only depend on the graph Λ through the constant γ . This means that the bounds are stable in the limit where Λ is large and γ stays fixed (which is the case in most practical situations, e.g., for large boxes or tori in \mathbb{Z}^D).

Notice that H commutes with the total particle number operator \mathcal{N} and can therefore be block-diagonalized as $H = \bigotimes_{N \geq 0} H_N$. It is self-adjoint in the domain

$$\mathcal{D}(H) = \left\{ \varphi = (\varphi)_{N \geq 0} \in \mathcal{F}(\ell^2(\Lambda)) : \sum_{N \geq 0} \|H_N \varphi_N\|^2 < \infty \right\},$$

see, e.g., [FLS22b, Appendix A] for a proof. By Stone's theorem, e^{-itH} is a strongly continuous one-parameter group of unitaries and we can thus define the Heisenberg time evolution of any bounded observable O by $O(t) = e^{itH} O e^{-itH}$. Any restriction of the Hamiltonian $H_{X_0[R]}$ is also self-adjoint for the same reasons and so we can similarly define $O(t, X_0[R])$ by (1.4).

Our main results apply to operators O that are block-diagonal in the local particle number basis. The size of the block is captured by the parameter q_0 which physically represents the maximal change of local particle number that can be achieved by the operator O under consideration. Here and in the following, given a self-adjoint operator A and $\alpha \in \mathbb{R}$, we write $\Pi_{A > \alpha}$ for the spectral projector onto the spectral subspace where $A > \alpha$, and we define $\Pi_{A < \alpha}$, $\Pi_{A = \alpha}$, etc., analogously.

Definition 2.3 (Maximal change of local particle number). *Let $X_0 \subset \Lambda$ be bounded. Let O be an observable supported on X_0 . We say that O changes the local particle number by at most $q_0 \geq 0$, if*

$$\Pi_{n_{X_0} > m + q_0} O_{X_0} \Pi_{n_{X_0} = m} = 0, \quad \text{and} \quad \Pi_{n_{X_0} < m - q_0} O_{X_0} \Pi_{n_{X_0} = m} = 0, \quad \forall m > 0. \quad (2.5)$$

Of course, any particle-number conserving observable on X_0 satisfies Definition 2.3 with $q_0 = 0$. We shall derive the LRB for observables O that satisfy this definition for some $q_0 < \infty$.

2.2 Main results

Our main results are LRBs of the form (I) and (II) appearing in Theorem 1.1 but for much more general Bose-Hubbard type Hamiltonians (2.3) and assuming the moment bound condition (2.6). Our main example where the moment bound (2.6) can then be verified is in the presence of (i) strong local repulsion and (ii) translation invariance. Indeed, we confirm in Section 2.3 below that Theorem 1.1 follows from these results in a straightforward way.

We write ρ for the initial quantum state, which is a positive semidefinite operator of $\text{tr}(\rho) = 1$. Recall that $v_1(t) = t^{\frac{D-1}{p/2-D-1}}$ as has been defined in Theorem 1.1.

Theorem 2.4 (Enhanced LRB for trace norms). *Assume that there exist $T \in [0, \infty]$ and $p > 2D+2$ such that*

$$\text{tr}[\rho n_i^p(t)] \leq C_p, \quad \text{for all } t \in [0, T]. \quad (2.6)$$

Fix $X_0 \subset \Lambda$ and $q_0 \geq 0$.

Then, there exists a constant $C = C(D, \gamma, \bar{J}, k, p, C_p, X_0, q_0) > 0$ such that for all bounded observables O supported on X_0 satisfying Definition 2.3, we have

$$\|\rho [O(t) - O(t, R)]\|_1 \leq C \|O\| \left(\frac{v_1(t)t}{R} \right)^{p/2-D-1} \quad (2.7)$$

for all $R \geq 1$ and $t \in [1, T)$.

The constant C depends on the initial state only through p and C_p . We obtain Theorem 1.1 shows by bounding these constants uniformly in the initial state for translation-invariant Hamiltonian with strong local repulsion (with the same value of p).

For quantum-mechanical expectation values $\text{tr}[\rho(\cdot)]$, we require the moment bound (2.6) only under the weaker assumption $p > D + 1$ and we obtain a further enhanced velocity $v_{ex}(t) = t^{\frac{D}{p-D-1}}$.

Theorem 2.5 (Further-enhanced LRB for expectation values). *Assume that there exist $T \in [0, \infty]$ and $p > D + 1$ such that (2.6) holds. Fix $X_0 \subset \Lambda$ and $q_0 \geq 0$.*

Then, there exists a constant $C = C(D, \gamma, \bar{J}, k, p, C_p, X_0, q_0) > 0$ such that for all bounded observables O supported on X_0 satisfying Definition 2.3, we have

$$\|\rho[O(t) - O(t, R)]\|_1 \leq C \|O\| \left(\frac{v_1(t)t}{R} \right)^{p-D-1} \quad (2.8)$$

for all $R \geq 1$ and $t \in [1, T)$.

As the next section shows, Theorem 1.1 is now a straightforward application of these results with $T = \infty$. It is obtained by verifying the moment bounds in the presence of (i) strong repulsion and (ii) translation invariance.

2.3 Proof of Theorem 1.1 assuming Theorems 2.4 and 2.5

We recall the setting is that of a system of bosons on the discrete torus of side length $L \geq 1$,

$$\Lambda_L = ((-L/2, L/2] \cap \mathbb{Z})^D.$$

We work on the bosonic Fock space $\mathcal{F}(\ell^2(\Lambda_L))$ and consider a Bose-Hubbard type Hamiltonian of the form

$$H = H_0 + V, \quad H_0 = -J \sum_{\langle i, j \rangle} (b_i^\dagger b_j + b_j^\dagger b_i), \quad V = \sum_{i \in \Lambda_L} [n_i^p + v(n_i)]. \quad (2.9)$$

There exist $c_V, \epsilon > 0$ such that $|v(n)| \leq c_V(n+1)^{p-\epsilon}$ for all $n \geq 0$.

We consider either the case $p > 2D + 2$ or $p > D + 1$. Thanks to Theorems 2.4 and 2.5, applied with $T = \infty$, in either case, it suffices to prove the moment bound $\text{tr}[\rho n_i^p(t)] \leq C_p$ for all $t \geq 0$. Therefore, the proof of Theorem 1.1 is completed by the following lemma.

Lemma 2.6 (Moment bound). *Let $p > 1$ and let ρ be a translation-invariant state satisfying*

$$\frac{\text{tr}(\rho H)}{|\Lambda_L|} = E_\rho < \infty.$$

There exists $C = C(J, p, D, \epsilon, c_v)$ such that for any $L \geq 1$ and any $i \in \Lambda_L$, we have

$$\text{tr}(\rho n_i^p(t)) \leq 2(E_\rho + C), \quad \text{for all } t \geq 0. \quad (2.10)$$

Explicitly,

$$C(J, p, D, \epsilon, c_v) = \max_{n \in \mathbb{N}} \left(-\frac{n^p}{2} + c_V(n+1)^{p-\epsilon} + 4JDn \right). \quad (2.11)$$

Proof of Lemma 2.6. We first control the hopping term H_0 by the repulsion. By the Cauchy-Schwarz inequality for operators, we have

$$H_0 \succeq -J \sum_{\langle i,j \rangle} (n_i + n_j) = -4JDn,$$

where $O_1 \succeq O_2$ implies that the $O_1 - O_2$ is positive semidefinite. Hence, using also that $|v(n)| \leq c_V(n+1)^{p-\epsilon}$, we have

$$\begin{aligned} H &= H_0 + V \succeq \sum_{j \in \Lambda_L} \left[n_j^p + v(n_j) - 4JDn_j \right] \\ &\succeq \sum_{j \in \Lambda_L} \left[n_j^p - c_V(n_j + 1)^{p-\epsilon} - 4JDn_j \right]. \end{aligned}$$

By energy conservation (more precisely, the fact that the Heisenberg dynamics preserves the Hamiltonian) and cyclicity of the trace, we have

$$\mathrm{tr}(\rho H) = \mathrm{tr}(\rho(-t)H) \geq \sum_{j \in \Lambda_L} \mathrm{tr} \left\{ \rho(-t) \left[n_j^p - c_V(n_j + 1)^{p-\epsilon} - 4JDn_j \right] \right\}.$$

Fix a point $i \in \Lambda$. Since H commutes with translations, the time-evolved state $\rho(-t) = e^{-itH} \rho e^{itH}$ is also translation-invariant. This yields

$$\sum_{i \in \Lambda_L} \mathrm{tr} \left\{ \rho(-t) \left[n_j^p - c_V(n_j + 1)^{p-\epsilon} - 4JDn_j \right] \right\} = |\Lambda_L| \mathrm{tr} \left\{ \rho(-t) \left[n_i^p - c_V(n_i + 1)^{p-\epsilon} - 4JDn_i \right] \right\}.$$

Since $p > 1$, a simple case distinction shows that

$$n^p - c_V(n+1)^{p-\epsilon} - 4JDn \geq \frac{n^p}{2} - C, \quad \text{for all } n \geq 0,$$

with $C = C(J, p, D, \epsilon, c_v)$. By cyclicity of the trace and $\mathrm{tr}(\rho) = 1$, this yields

$$\mathrm{tr}(\rho H) \geq |\Lambda_L| \mathrm{tr} \left\{ \rho(-t) \left[\frac{1}{2} n_i^p - C \right] \right\} = |\Lambda_L| \left\{ \frac{1}{2} \mathrm{tr} [\rho n_i(t)^p] - C \right\}.$$

This proves Lemma 2.6 and therefore also Theorem 1.1. \square

2.4 An improved particle propagation bound for all moments

The main result 1 of [KVS24] is the particle propagation bound

$$\mathrm{tr} [\rho n_i^q(t)]^{1/q} \leq C \left(t^D + q + t \right), \quad (2.12)$$

for any $q \geq 1$ and $i \in \Lambda$, under the assumption that the initial state satisfies $\mathrm{tr} [\rho n_i^q]^{1/q} \leq Cq$.

It turns out that it is possible to improve this bound for general moments q by suitably interpolating between (2.12) and Lemma 2.6. This shows that the rigidity effect we leverage also constrains transport properties of higher moments of the local particle number.

Proposition 2.7 (Improved particle propagation bound). *Let $q \geq p$ and suppose that $\text{tr}[\rho n_i^q]^{1/q} \leq Cq$. Under the assumption of Theorem 1.1, there exists $C = C(J, p, q, D, \epsilon, c_v, E_\rho)$ such that we have the bound*

$$(\text{tr}(\rho n_i(t)^q))^{1/q} \leq Ct^{D(1-p/q)}, \quad t \geq 1. \quad (2.13)$$

Compared to (2.12), this has the additional negative term $-Dp/q$. The improvement becomes progressively weaker as $q \rightarrow \infty$ and the bound continuously turns into Lemma 2.6 at the endpoint $q = p$. These features are expected, because we prove Proposition 2.7 by interpolation.

Proof of Proposition 2.7. We use the Riesz-Thorin interpolation theorem for a suitably chosen classical probability space. For $q \geq p$, let $q_1 \geq 1$ and $\theta \in [0, 1]$ be such that

$$\frac{1}{q} = \frac{1-\theta}{p} + \frac{\theta}{q_1}.$$

Then, we claim that

$$\text{tr}(\rho n_i(t)^q)^{1/q} \leq C \left(t^D + q_1 + t \right)^\theta. \quad (2.14)$$

Given the state $\rho(-t)$, consider its spectral decomposition

$$\rho(-t) = \sum_j \lambda_j \Pi_{\psi_j},$$

where Π_{ψ_j} project onto the eigenvector ψ_j corresponding to the eigenvalue λ_j . We allow for degeneracy and choose the $\{\psi_j\}_j$ orthonormal.

Fix a site $i \in \Lambda$. Define a classical probability distribution $\mathbf{p} : \mathbb{N} \rightarrow [0, 1]$ by

$$\mathbf{p}(n) = \sum_j \lambda_j \|\Pi_{n_i=n} \psi_j\|^2$$

(Notice that indeed $\sum_{n \geq 0} \mathbf{p}(n) = \sum_j \lambda_j = 1$).

Then expectation values of powers of n_i can be viewed as L^q -norms over this probability space

$$(\text{tr}(\rho n_i^q))^{1/q} = \mathbb{E}[X^q]^{1/q} \equiv \|X\|_q,$$

where X denotes the corresponding \mathbb{N} -valued classical random variable.

In this setting, we can apply the Riesz-Thorin interpolation theorem for general L^q -measure spaces, which says that

$$\|X\|_{q_\theta} \leq \|X\|_2^{1-\theta} \|X\|_q^\theta$$

with θ, q_θ as above. Writing this out in terms of the quantum state $\rho(-t)$ and applying Lemma 2.6 and (2.12) afterwards yields

$$\text{tr}(\rho n_i^{q_\theta}(t))^{1/q_\theta} = \text{tr}(\rho(-t) n_i^{q_\theta})^{1/q_\theta} \leq \text{tr}(\rho n_i^2(t))^{\frac{1-\theta}{2}} \text{tr}(\rho n_i^q(t))^{\frac{\theta}{q}} \leq C \left(t^D + q \right)^\theta.$$

This proves (2.14).

It remains to choose the parameters suitably. The bound (2.14) clearly improves over (2.12) for every q because the RHS of (2.14) behaves as $t^{D\theta}$ for $t \gg 1$ and $\theta < 1$. To make the improvement quantitative, we fix $q > p$ and find a pair of $\theta \in [0, 1]$ and $q_1 > q$ such that $\frac{1}{q} = \frac{1-\theta}{p} + \frac{\theta}{q_1}$. The minimal value of the factor θ is achieved for $q_1 \rightarrow \infty$ (in which case $\theta \rightarrow 1 - p/q$). However, this

choice makes the RHS of (2.14) infinitely large. We can remedy this by making the (effectively very large) choice $q_1 = t^D$. This gives

$$\theta = \frac{1 - p/q}{1 - p/t^D} = (1 - p/q_*) \left(1 + \frac{p}{t^D - p}\right), \quad (2.15)$$

and so (2.14) becomes

$$\text{tr}(\rho n_i(t)^q) \leq Ct^{D(q-p)} \left(1 + \frac{p}{t^D - p}\right) \leq Ct^{D(q-p)}, \quad (2.16)$$

where the second bounds holds because $t^{D(q-p)} \frac{p}{t^D - p} = \exp\left[D(q-p)p \frac{\log(t)}{t^D - p}\right] = \mathcal{O}(1)$ as $t \rightarrow \infty$. This proves Proposition 2.7. \square

The reader may wonder if Proposition 2.7 can be used in place of Lemma 2.6 in the proof of the Lieb-Robinson bound to further improve the light cone beyond what we establish in Theorems 2.5 and 2.4. This, however, is not the case. Implementing this strategy in the arguments detailed below would ultimately have the effect that the scaling of the parameter \tilde{C}_p in Eq. (3.20) changes to $C_{p,t}^{(2)} \propto t^{D(q-p)}$, and hence the light cone shape is given by

$$\left\| \left[O_X(t) - O_{i_0[R]}^{(\tilde{m})} \right] \rho_0 \right\|_1 \lesssim t^{D(q-p)/2} \frac{t^{q/2-1}}{R^{q/2-D-1}}. \quad (2.17)$$

In the average case, we have

$$\left| \text{tr} \left\{ \left[O_X(t) - O_{i_0[R]}^{(\tilde{m})} \right] \rho_0 \right\} \right| \lesssim t^{D(q-p)} \frac{t^{q-1}}{R^{q-D-1}}. \quad (2.18)$$

By optimizing these bounds in $q \geq p$, one finds that depending on the value of p , these bounds reproduce either the velocity bound $v(t) \sim t^D$ from [KVS24] or $v_1(t)$ for (2.17) and $v_{\text{ex}}(t)$ for (2.18) that we prove in Theorem 1.1. The reason for this is that the light cone exponent is a ratio of linear functions in q and therefore monotonic, which means it reaches its minimum always at one of the endpoints $q = p$ or $q = \infty$. In summary, the additional interpolation does not pay off from the perspective of light cone scaling, only for particle transport.

3 Proof strategy for Theorems 2.4 and 2.5

Terminology. In the following, we refer to all constants that are independent of R, t , and the number of vertices or edges in Λ as “ $\mathcal{O}(1)$ constants”. Examples of such $\mathcal{O}(1)$ constants are γ, \bar{J}, k and the dimension D .

3.1 Setting up the proof

For an arbitrary subset $X \subseteq \Lambda$, we define the subset Hamiltonians $H_{0,X}, V_X$ and H_X as follows:

$$\begin{aligned} H_X &= H_{0,X} + V_X, \\ H_{0,X} &:= \sum_{i,j \in X} J_{i,j} (b_i b_j^\dagger + \text{h.c.}), \quad V_X := \sum_{Z \subseteq X} v_Z. \end{aligned} \quad (3.1)$$

Note that they are supported on the subset X . Also, we define the boundary hopping terms on the region X as follows:

$$\partial h_X := H_0 - H_{0,X} - H_{0,X^c} = \sum_{i \in \partial X} \sum_{j \in X^c: d_{i,j}=1} J_{i,j} (b_i b_j^\dagger + \text{h.c.}). \quad (3.2)$$

For the proof, we, first of all, rely on the unitary-concatenation technique that was employed in Ref. [Kuw16, KS21a, KS21b, KVS24]. This technique allows us to reduce the LRB to short-time LRBs (see Subtheorem 3.1 below) by a concatenation procedure that we explain now.

For the remainder of this section, we work under the assumptions of Theorem 2.4. The proof involves various parameter choices, which can be made relatively explicitly. To this end, we introduce the following $\mathcal{O}(1)$ constants in the sense defined above.

The argument can be adapted to the case when this assumption does not hold, but the explicit parameter choices become more cumbersome.

Given two parameters $\eta, \tau > 0$, which we will later choose as suitable $\mathcal{O}(1)$ constants, we define

$$c_{\tau,1} := 40e^{4\gamma\bar{J}\tau}, \quad (3.3)$$

$$c_{\tau,2} := e^{4\gamma\bar{J}\tau+1} (1 + 8\gamma\bar{J}\tau + \tau), \quad (3.4)$$

$$c_3 := 4\bar{J}\eta\gamma(2k)^D d_G, \quad (3.5)$$

$$c'_3 := 16ekc_3\gamma(2k)^D. \quad (3.6)$$

Here, d_G is the maximum degree of the graph, which is also upper-bounded by γ , but may be smaller.

We also define

$$\mu := C_p^{-1/p} (4q_0^3 c_{\tau,1} + c_{\tau,2} p) + c_{\tau,1} + 2^{D+1} D! c_{\tau,1}, \quad (3.7)$$

and

$$\tilde{C}_p := 16C_p(2\mu/\eta)^p (1 + 2\gamma\bar{J}\eta 2^{p/2})^2. \quad (3.8)$$

The parameters $\eta, \tau > 0$ will be a suitably chosen $\mathcal{O}(1)$ constants and so $c_{\tau,1}$, $c_{\tau,2}$, c_3 , c'_3 , μ , \tilde{C}_p , will be $\mathcal{O}(1)$ constants as well. The fact that we are able to choose μ as an $\mathcal{O}(1)$ parameter, as opposed to the time-dependent choice in [KVS24], requires using the polynomial error bounds and it is one of the key differences in the proof to [KVS24]. Another parameter in the proof is $r > 0$. This parameter plays a different role and it is not $\mathcal{O}(1)$ in the sense defined above. Roughly speaking, $r \approx R/t$ which makes it large outside the light cone.

3.2 The short-time LRB and how it implies Theorem 2.4

Subtheorem 3.1 (Short-time LRB). *Fix a region X and parameters $\eta, \tau, r > 0$ subject to the constraints*

$$\tau \leq \frac{1}{ec'_3}, \quad e^{-r/2}|X| \leq 1. \quad (3.9)$$

Let $\tilde{X} = X[r] \setminus X$. There exists a unitary operator $U_{X[r]}$ supported on $X[r]$ such that the following bound holds for any quantum state ρ satisfying the condition (2.6),

$$\left\| \left[O_X(\tau) - U_{X[r]}^\dagger O_X U_{X[r]} \right] \rho \right\|_1 \leq e^3 c_3 \tau |\partial(X[r])| r e^{-r/(4k)} + \tilde{C}_p^{1/2} |\tilde{X}| r^{-p/2+1}, \quad (3.10)$$

In essence, the statement is that $U_{X[r]}$ functions as an appropriate local approximation of the time evolution $e^{-iH\Delta t}$. We prove Subtheorem 3.1 in Section 4. The explicit form of $U_{X[r]}$ is not important for what follows; it suffices that it exists and functions as a local approximation of the time evolution in the sense of Subtheorem 3.1. We obtain its existence through Proposition 4.1 which is taken from [KS21b], from which an explicit can be obtained if desired.

Proof of Theorem 2.4 assuming Subtheorem 3.1. Fix $t \in [1, T)$ and $R \geq 1$. Notice that the claim of Theorem 2.4 is trivial when $R/(v_1(t)t) = R/t^\zeta \leq 1$ where we set $\zeta = 1 + \frac{D-1}{p/2-D-1} > 1$. Hence, we may assume without loss of generality that $R > t^\zeta$. The idea is to apply Subtheorem 3.1 various times and concatenate the resulting short-time LRBs.

Step 1: Decomposition and short-time LRBs. We decompose the time t into $\bar{m} := t/\Delta t$ equal time steps of size Δt , an $\mathcal{O}(1)$ constant to be determined:

$$t := \bar{m}\Delta t. \quad (3.11)$$

It is convenient to measure distance in a radial way. For this, we find a lattice site i_0 and radius r_0 such that the initial region $X_0 \subset i_0[r_0]$, i.e., it is contained in the ball of radius r_0 centered at i_0 .

Then, for each $0 \leq j \leq \bar{m}$, we define the subset X_j as follows:

$$X_j := i_0[r_0 + j\Delta r], \quad \Delta r = \left\lfloor \frac{R - r_0}{\bar{m}} \right\rfloor = \left\lfloor \frac{R'}{\bar{m}} \right\rfloor \quad (R' := R - r_0), \quad (3.12)$$

where, in particular, $X_{\bar{m}} \subseteq i_0[R]$.

Now we apply Subtheorem 3.1 several times to obtain unitaries $\{U_{X_j}\}_{j=1}^{\bar{m}}$. We choose the parameters suitably and check the conditions. We take $\tau = \Delta t = 1/(ec'_3)$ which is an $\mathcal{O}(1)$ constant such that the first condition in (3.9) is satisfied. Moreover, we take $r = \Delta r$ and $X = X_{j-1}$ and $\rho = \rho_0(\bar{m}\Delta t - j\Delta t)$. The second condition in (3.9) is satisfied for all j as long as $e^{-R\Delta t/(2t)}|i_0[R]| \leq 1$, which by our geometric assumptions is implied by

$$\gamma e^{-R\Delta t/(2t)} R^D \leq 1.$$

To verify this condition, recall that we assumed without loss of generality that $R > t^\zeta$ with $\zeta > 1$. From this, we obtain

$$\gamma e^{-R\Delta t/(2t)} R^D \leq \gamma e^{-R^{1-1/\zeta}\Delta t} R^D.$$

This holds for R sufficiently large. Notice that we can also assume that R is sufficiently large without loss of generality since Theorem 2.4 holds trivially if $R \leq t^\zeta$. This can be seen by using $t \geq 1$ and lower-bounding the RHS of (2.7) as $\left(\frac{v_1(t)t}{R}\right)^{p/2-D-1} \geq 1$.

We have verified the conditions of Subtheorem 3.1 with the above parameter choices. Hence, we apply Subtheorem 3.1 to obtain unitaries $\{U_{X_j}\}_{j=1}^{\bar{m}}$ such that

$$\left\| \left[O_{X_{j-1}}(\Delta t) - U_{X_j}^\dagger O_{X_{j-1}} U_{X_j} \right] \rho_0(\bar{m}\Delta t - j\Delta t) \right\|_1. \quad (3.13)$$

Step 2: Concatenation of short-time LRBs. Recall that we consider an arbitrary observable O_{X_0} . We iteratively define local approximations $\{O_{X_j}^{(j)}\}_{j=1}^{\bar{m}}$ via

$$O_{X_j}^{(j)} = U_{X_j}^\dagger O_{X_{j-1}}^{(j-1)} U_{X_j} \quad \text{for } 1 \leq j \leq \bar{m}. \quad (3.14)$$

Moreover, we use $X_{\bar{m}} \subseteq i_0[R]$ to set $O_{i_0[R]} = O_{X_{\bar{m}}}^{(\bar{m})}$.

By following the unitary telescoping argument in Sec. IX B in Ref. [KVS24] and using (3.13), we obtain

$$\begin{aligned}
\left\| \left[O_X(t) - O_{i_0[R]}^{(\bar{m})} \right] \rho_0 \right\|_1 &\leq \sum_{j=1}^{\bar{m}} \left\| \left[O_{X_{j-1}}^{(j-1)}(\Delta t) - O_{X_j}^{(j)} \right] \rho_0(\bar{m}\Delta t - j\Delta t) \right\|_1 \\
&\leq \bar{m} \left[e^3 c_3 \Delta t |\partial(i_0[R])| \Delta r e^{-\Delta r/(4k)} + \tilde{C}_p^{1/2} \Delta r |\partial(i_0[R])| (\Delta r)^{-p/2+1} \right] \\
&\leq \frac{t}{\Delta t} \left[e^3 c_3 \gamma \frac{\Delta t^2 R^D}{t} e^{-\Delta t R/(4kt)} + \tilde{C}_p^{1/2} \gamma R^{D-1} \left(\frac{\Delta t R}{t} \right)^{-p/2+2} \right] \\
&= e^3 c_3 \gamma \Delta t R^D e^{-\Delta t R/(4kt)} + \tilde{C}_p^{1/2} \gamma (\Delta t)^{-p/2+1} \frac{R^{D+1-p/2}}{t^{-p/2+1}}, \tag{3.16}
\end{aligned}$$

where we use that $\tilde{X} = X_j \setminus (X_{j-1}[\Delta r/2])$ always satisfies $|\tilde{X}| \leq \Delta r |\partial X_{\bar{m}}| = \Delta r |\partial(i_0[R])|$. This implies Theorem 2.4 with the right-hand side replaced by

$$C \|O\| \left[R^D e^{-R/(4kt)} + \left(\frac{v_1(t)t}{R} \right)^{p/2-D-1} \right].$$

It remains to remove the first term $R^D e^{-R/(4kt)}$. To this end, we argue similarly as above, recalling that we may assume $R/t^\zeta > 1$ with $\zeta > 1$. Using this as well as $t \geq 1$, we obtain

$$R^D e^{-R/(4kt)} \leq R^D e^{-R^{1-1/\zeta}/(4k)} \leq (t^\zeta/R)^{p-D-1} \underbrace{R^{2D+1-p} e^{-(R^{1-1/\zeta})/(4k)}}_{\leq C'=C'(D,p,k)} \leq C' (t^\zeta/R)^{p-D-1}.$$

This completes the proof of Theorem 2.4. \square

3.3 Short-time LRB for expectation values and how it implies Theorem 2.5

The overall proof strategy is the same as that for Theorem 2.4. That is, we concatenate suitable short-time LRBs which now control the average. That is, our main goal is to derive an analogous statement to Subtheorem 3.1 for the expectation value. We introduce

$$\tilde{C}_p := 4C_p \left(\frac{2\mu}{\eta} \right)^p \left(1 + \frac{\tau \bar{J} \gamma \eta}{2} \sqrt{\gamma 2^D \left(\frac{2^p}{C_p \mu^p} + 1 \right)} \right). \tag{3.17}$$

Subtheorem 3.2 (Short-time LRB for the expectation). *Under the same setup as in Subtheorem 3.1, there exists a unitary $U_{X[r]}$ supported on $X[r]$ such that*

$$\left| \text{tr} \left[\left(O_X(\tau) - U_{X[r]}^\dagger O_X U_{X[r]} \right) \rho \right] \right| \leq e^3 c_3 \tau |\partial(X[r])| r e^{-r/(4k)} + \tilde{C}_p |\tilde{X}| r^{-p+1}, \tag{3.18}$$

where c_3 has been defined in Proposition 4.1, and τ and r should satisfy the conditions in (3.9). Also, \tilde{C}_p is defined in Eq. (3.17) below.

We prove Subtheorem 3.2 in Section 5.

Proof of Theorem 2.5 assuming Subtheorem 3.2. The inequality (3.15) is replaced with

$$\left| \text{tr} \left[\left(O_{X_0}(t) - O_{i_0[R]}^{(\bar{m})} \right) \rho_0 \right] \right| \leq \sum_{j=1}^{\bar{m}} \left| \text{tr} \left\{ \left[O_{X_{j-1}}^{(j-1)}(\Delta t) - O_{X_j}^{(j)} \right] \rho_0 (\bar{m} \Delta t - j \Delta t) \right\} \right|. \quad (3.19)$$

Now we can use Subtheorem 3.2, we prove the main inequality (2.8) by similar calculations as in (3.15):

$$\begin{aligned} \left| \text{tr} \left[\left(O_X(t) - O_{i_0[R]}^{(\bar{m})} \right) \rho_0 \right] \right| &\leq \bar{m} \left[e^3 c_3 \Delta t |\partial(i_0[R])| \Delta r e^{-\Delta r/(4k)} + \tilde{C}_p \Delta r |\partial(i_0[R])| (\Delta r)^{-p+1} \right] \\ &\leq \frac{t}{\Delta t} \left[e^3 c_3 \gamma \frac{\Delta t^2 R^D}{t} e^{-\Delta t R/(4kt)} + \tilde{C}_p \gamma R^{D-1} \left(\frac{\Delta t R}{t} \right)^{-p+2} \right] \\ &= e^3 c_3 \gamma \Delta t R^D e^{-\Delta t R/(4kt)} + \tilde{C}_p \gamma (\Delta t)^{-p} \frac{R^{D+1-p}}{t^{-p+1}} \\ &= e^3 c_3 \gamma \Delta t R^D e^{-\Delta t R/(4kt)} + \tilde{C}_p \gamma (\Delta t)^{-p} \left(\frac{t^{\frac{p-1}{p-D-1}}}{R} \right)^{p-D-1}. \end{aligned} \quad (3.20)$$

The exponential can be removed by the same considerations as in Theorem 2.4. We thus complete the proof of Theorem 2.5. \square

4 Proof of Subtheorem 3.1

4.1 Overview of the proof

The basic strategy is quite similar to the ones in Ref. [KS21b], which treated a scenario in which a steady state is perturbed. The rough idea is to approximate the dynamics with truncated dynamics in which the local boson number is cut off at level \bar{q} , which has to be suitably chosen. The approximation error is controlled by Markov's inequality, and the truncated dynamics effectively describes a quantum spin system with bounded interactions, depending on \bar{q} , and therefore enjoys Lieb-Robinson bounds depending on \bar{q} . Compared to previous works, including Ref. [KVS24], the technical novelty here is that the probability decay is only bounded polynomially as the moment upper bound is assumed only up to p th order. This point forces us to more carefully analyze the error estimation due to the truncation of the boson number at each of the sites. In the proof, we also rely on several technical results from Ref. [KVS24] which we quote as needed. Our presentation is focused on the steps that require novel ideas.

In the remainder of this section, we work in the setting of Subtheorem 3.1. In particular, we fix a region X and parameters $\eta, \tau, r > 0$ subject to the constraints (3.9). We then define the effective Hamiltonian $\tilde{H}[L, q]$ with $L \subseteq \Lambda$ and $q \in \mathbb{N}$ as follows

$$\begin{aligned} \tilde{H}[L, q] &:= \tilde{H}_0[L, q] + \tilde{V}[L, q], \\ \tilde{H}_0[L, q] &:= \bar{\Pi}_{L,q} H_0 \bar{\Pi}_{L,q}, \quad V[L, q] := \bar{\Pi}_{L,q} V \bar{\Pi}_{L,q}, \end{aligned} \quad (4.1)$$

where $\bar{\Pi}_{L,q}$ ($L \subseteq \Lambda$) is defined as the projection onto the boson-number space such that for an arbitrary $i \in L$ the boson number n_i is truncated up to q :

$$\bar{\Pi}_{L,q} := \prod_{i \in L} \Pi_{n_i \leq q}. \quad (4.2)$$




Figure 1: Schematic of the construction of the effective Hamiltonian $\tilde{H}[\tilde{X}, \bar{q}] = \bar{\Pi}_{\tilde{X}, \bar{q}} H \bar{\Pi}_{\tilde{X}, \bar{q}}$. The spectral projection $\bar{\Pi}_{\tilde{X}, \bar{q}}$ implements the truncation of the local boson number at level \bar{q} in the region of $\tilde{X} = X[r] \setminus X[r/2]$.

In the proof, we consider the subset \tilde{X} defined as

$$\tilde{X} := X[r] \setminus X[r/2]. \quad (4.3)$$

Our choice for the boson number truncation will be called \bar{q} and we will take $\bar{q} = \eta \ell_0$.

Roughly speaking, our goal is to approximate the dynamics $O_X(\tau)\rho_0(t)$ in two steps as follows:

$$O_X(\tau)\rho_0(t) \approx O_X(\tilde{H}[\tilde{X}, \bar{q}], \tau)\rho_0(t), \quad (4.4)$$

and

$$O_X(\tilde{H}[\tilde{X}, \bar{q}], \tau)\rho_0(t) \approx U_{X[r]}^\dagger O_X U_{X[r]}\rho_0(t), \quad (4.5)$$

where $U_{X[r]}$ is an appropriately chosen unitary operator.

For the second approximation (4.5), we can directly utilize the previous analyses as follows:

Proposition 4.1 (Proposition 6 and (S.97) in Ref. [KS21b]). *Choose \bar{q} as $\bar{q} = \eta \ell_0$ with $\ell_0 := r/2$. Suppose that $\tau \leq \frac{1}{ec_3}$.*

Then, we have

$$\left\| O_X(\tilde{H}[\tilde{X}, \eta \ell_0], \tau) - U_{X[r]}^\dagger O_X U_{X[r]} \right\|_1 \leq 2e^3 c_3 \tau |\partial(X[r])| \ell_0 e^{-\ell_0/(2k)}. \quad (4.6)$$

We remark that the precise construction of the unitary operator $U_{X[r]}$ is given in Ref. [KS21b, Section S.V.C]

The remaining problem is thus to estimate the approximation error for (4.4), which is given by the following proposition.

Proposition 4.2. *Let ρ be an arbitrary quantum state satisfying the moment upper bound as*

$$\text{tr}(n_i^p \rho) \leq C_p. \quad (4.7)$$

Then, under the condition that

$$e^{-r/2}|X| \leq 1, \quad (4.8)$$

we obtain

$$\left\| O_X(H, \tau)\rho - O_X(\bar{\Pi}_{\tilde{X}, \bar{q}} H \bar{\Pi}_{\tilde{X}, \bar{q}}, \tau)\rho \right\|_1 \leq 4|\tilde{X}|^{1/2} C_p^{1/2} \left(\frac{\mu}{\bar{q}} \right)^{p/2} \left(1 + 4\tau\gamma\bar{J}\bar{q}|\tilde{X}|^{1/2} 2^{p/2} \right). \quad (4.9)$$

Proof of Subtheorem 3.1 assuming Proposition 4.2. By setting $\bar{q} = \eta r/2$ to the upper bound (4.9), we reduce it to

$$\begin{aligned} \left\| O_X(H, \tau)\rho - O_X(\bar{\Pi}_{\tilde{X}, \bar{q}} H \bar{\Pi}_{\tilde{X}, \bar{q}}, \tau)\rho \right\|_1 &\leq 4|\tilde{X}|^{1/2} C_p^{1/2} \left(\frac{2\mu}{\eta r} \right)^{p/2} \left(1 + 2\tau\gamma\bar{J}\eta r|\tilde{X}|^{1/2} 2^{p/2} \right) \\ &\leq \tilde{C}_p^{1/2} |\tilde{X}| r^{-p/2+1} \end{aligned} \quad (4.10)$$

with \tilde{C}_p chosen as in (3.8).

Finally, by combining the two upper bounds of (4.6) and (4.10), we obtain the main inequality (3.10). This completes the proof of Subtheorem 3.1. \square

In the remainder of this section, we prove Proposition 4.2.

4.2 Proof of Proposition 4.2

We first state the two lemmas that we combine to obtain Proposition 4.2.

The first lemma is a general, abstract bound on the difference between the full Heisenberg dynamics and the dynamics generated by an effective projected Hamiltonian that we quote from [KVS24].

Lemma 4.3 (Corollary 26 in Ref. [KVS24]). *Let ρ be an arbitrary quantum state. Then, for an arbitrary projection operator $\bar{\Pi}$, the dynamics of the effective Hamiltonian $\bar{\Pi}H\bar{\Pi}$ is deviated from the true dynamics by the error of*

$$\begin{aligned} \left\| O(H, \tau)\rho - O(\bar{\Pi}H\bar{\Pi}, \tau)\rho \right\|_1 &\leq \left\| \bar{\Pi}^c O(\tau)\sqrt{\rho} \right\|_F + \left\| \bar{\Pi}^c O\sqrt{\rho(\tau)} \right\|_F + \left\| \bar{\Pi}^c \sqrt{\rho} \right\|_F + \left\| \bar{\Pi}^c \sqrt{\rho(\tau)} \right\|_F \\ &\quad + \int_0^\tau \left\| \bar{\Pi}H_0\bar{\Pi}^c O(\tau - \tau_1)\sqrt{\rho(\tau_1)} \right\|_F d\tau_1 + \int_0^\tau \left\| \bar{\Pi}H_0\bar{\Pi}^c \sqrt{\rho(\tau_1)} \right\|_F d\tau_1, \end{aligned} \quad (4.11)$$

where $\|\dots\|_F$ is the Frobenius norm, i.e., $\|O\|_F = \sqrt{\text{tr}(OO^\dagger)}$.

In the following, \tilde{X}, \bar{q} remain fixed and we abbreviate

$$\bar{\Pi} \equiv \bar{\Pi}_{\tilde{X}, \bar{q}}$$

in (4.11).

All the terms in the RHS of the inequality (4.11) can be expressed through (integrals of) norms of the form

$$\left\| \bar{\Pi}_{\tilde{X}, \bar{q}}^c O_X(\tau_2)\sqrt{\rho(\tau_1)} \right\|_F, \quad (4.12)$$

and

$$\left\| \bar{\Pi}_{\tilde{X}, \bar{q}} H_0 \bar{\Pi}_{\tilde{X}, \bar{q}}^c O_X(\tau_2) \sqrt{\rho(\tau_1)} \right\|_F. \quad (4.13)$$

for suitable choices of $\tau_2 = \mathcal{O}(1)$.

The second lemma bounds expressions of the form (4.12) and (4.13).

Lemma 4.4. *Suppose the assumption of Proposition 4.2 hold. For every $\tau_1, \tau_2 \leq \tau$, we have the bounds*

$$\left\| \bar{\Pi}_{\tilde{X}, \bar{q}}^c O_X(\tau_2) \sqrt{\rho(\tau_1)} \right\|_F \leq |\tilde{X}|^{1/2} C_p^{1/2} \left(\frac{\mu}{\bar{q}} \right)^{p/2}, \quad (4.14)$$

and

$$\left\| \bar{\Pi}_{\tilde{X}, \bar{q}} H_0 \bar{\Pi}_{\tilde{X}, \bar{q}}^c O_X(\tau_2) \sqrt{\rho(\tau_1)} \right\|_F \leq 2\gamma \bar{J} \bar{q} |\tilde{X}| \left(\frac{2\mu}{\bar{q}} \right)^{p/2} C_p^{1/2}. \quad (4.15)$$

This result is a polynomial analog of Proposition 29 in Ref. [KVS24].

Proof of Proposition 4.2. We apply Lemma 4.3. Next, we use using the above inequalities with appropriate τ_1 and τ_2 on the RHS in (4.11). This gives

$$\left\| O_X(H, \tau) \rho - O_X(\bar{\Pi}_{\tilde{X}, \bar{q}} H \bar{\Pi}_{\tilde{X}, \bar{q}}^c, \tau) \rho \right\|_1 \leq 4 |\tilde{X}|^{1/2} C_p^{1/2} \left(\frac{\mu}{\bar{q}} \right)^{p/2} \left[1 + 4\tau\gamma \bar{J} \bar{q} |\tilde{X}|^{1/2} 2^{p/2} \right], \quad (4.16)$$

which gives the desired inequality (4.9). \square

It therefore remains for us to prove Lemma 4.4.

4.2.1 Generalized moment bounds

We will control the spectral projector in Lemma 4.4 by moments through Markov's inequality which are then controlled by the following generalized moment bounds. These bounds control the moments of the local particle number with respect to the state $O_X(\tau_2) \rho(\tau_1) [O_X(\tau_2)]^\dagger$, i.e.,

$$\text{tr} \left[O_X(\tau_2) \rho(\tau_1) [O_X(\tau_2)]^\dagger n_i^p \right] = \text{tr} \left[e^{iH\tau_2} O_X \rho(\tau_1 - \tau_2) O_X^\dagger e^{-iH\tau_2} n_i^p \right]. \quad (4.17)$$

We will prove the following Lemma:

Lemma 4.5. *Let ρ be an arbitrary quantum state satisfying the moment upper bound as*

$$\text{tr} (n_i^p \rho) \leq C_p. \quad (4.18)$$

Then,

$$\text{tr} \left[e^{iH\tau_2} O_X \rho(\tau_1 - \tau_2) O_X^\dagger e^{-iH\tau_1} n_i^p \right] \leq \mu_{d_i, X}^p C_p, \quad (4.19)$$

where $\mu_{d_i, X}$ is defined by

$$\mu_{d_i, X} := C_p^{-1/p} \left(4q_0^3 c_{\tau, 1} e^{-d_i, X} + c_{\tau, 2} p \right) + c_{\tau, 1} e^{-d_i, X} |X| + 2^{D+1} D! c_{\tau, 1} \quad (4.20)$$

using the definitions in Eqs. (3.3) and (3.4) for $c_{\tau, 1}$ and $c_{\tau, 2}$.

We will apply this for $i \in \tilde{X}$. Then the standing assumption $e^{-r/2}|X| \leq 1$ will imply that $\mu_{d_i, X} \leq \mu$ with μ given by (3.7).

Remark 4.6. We note that the lemma can be extended to the effective Hamiltonian \tilde{H} as $\tilde{H} := \bar{\Pi}_{\tilde{X}, \bar{q}} H \bar{\Pi}_{\tilde{X}, \bar{q}}$. That is, we have

$$\mathrm{tr} \left(e^{i\tilde{H}\tau_2} O_X \rho(\tilde{H}, \tau_1 - \tau_2) O_X^\dagger e^{-i\tilde{H}\tau_1} n_i^p \right) \leq \mu_{d_i, X}^p C_p. \quad (4.21)$$

This generalization is utilized in considering the error in terms of the average instead of the trace norm; see Section 5 below.

4.2.2 Proof of Lemma 4.5

To make the notation simple, we denote $\rho(\tau_1 - \tau_2)$ and τ_2 by ρ and τ in this proof. We then consider

$$\begin{aligned} \mathrm{tr} \left(e^{iH\tau} O_X \rho O_X^\dagger e^{-iH\tau} n_i^p \right) &= \mathrm{tr} \left\{ O_X \rho O_X^\dagger [n_i(-\tau)]^p \right\} \\ &\leq \mathrm{tr} \left[O_X \rho O_X^\dagger \left(n_i + c_{\tau,1} \hat{\mathcal{D}}_i + c_{\tau,2p} \right)^p \right], \end{aligned} \quad (4.22)$$

where we use the Subtheorem 1 in Ref. [KVS24]. Here, τ is assumed to be smaller than $1/(4\gamma\bar{J})$ ($\tau \leq 1/(4\gamma\bar{J})$), and define $\hat{\mathcal{D}}_i$ as

$$\hat{\mathcal{D}}_i := \sum_{j \in \Lambda} e^{-d_{i,j}} n_j. \quad (4.23)$$

We second use Lemma 3 in Ref. [KVS24], which proves the following statement:

Lemma 4.7 (Lemma 3 in Ref. [KVS24]). *Let O_{q_0} be an arbitrary operator satisfying Definition 2.3 with q_0 . Note that O_{q_0} is not necessarily supported on the subset L . Then, we obtain*

$$O_{q_0}^\dagger \left(\sum_{j \in \Lambda} \nu_j n_j \right)^m O_{q_0} \preceq 4^m \|O_{q_0}\|^2 \left(4\bar{\nu} q_0^3 + \bar{\nu} n_L + \sum_{j \notin L} \nu_j n_j \right)^m, \quad (4.24)$$

where we define $\bar{\nu} := \max_{j \in L} (\nu_j)$ with $\nu_j > 0$. Here, for arbitrary operators O_1 and O_2 , the notation $O_1 \preceq O_2$ means that $O_2 - O_1$ is a semi-positive definite operator.

We apply the lemma with the choices

$$O_q \rightarrow O_X, \quad \nu_j \rightarrow c_{\tau,1} e^{-d_{i,j}}, \quad m \rightarrow p, \quad \bar{\nu} \rightarrow c_{\tau,1} e^{-d_{i,X}}, \quad L \rightarrow X, \quad (4.25)$$

and we obtain

$$O_X^\dagger \left(n_i + c_{\tau,1} \hat{\mathcal{D}}_i + c_{\tau,2p} \right)^p O_X \preceq 4^p \left(4q_0^3 c_{\tau,1} e^{-d_{i,X}} + c_{\tau,1} e^{-d_{i,X}} n_X + c_{\tau,1} \sum_{j \notin X} e^{-d_{i,j}} n_j + c_{\tau,2p} \right)^p. \quad (4.26)$$

Let $m \leq p$. By Hölder's inequality and the moment upper bound (4.18),

$$\mathrm{tr} \left(\rho \prod_{s=1}^m n_s \right) \leq \prod_{s=1}^m \mathrm{tr} (\rho n_s^m)^{1/m} \leq \left(C_m^{1/m} \right)^m = C_m. \quad (4.27)$$

By Jensen's inequality, $C_m^{1/m} \leq C_p^{1/p}$ for any $m \leq p$ and so we have

$$\begin{aligned} & \text{tr} \left[\rho \left(4q_0^3 c_{\tau,1} e^{-d_{i,X}} + c_{\tau,1} e^{-d_{i,X}} n_X + c_{\tau,1} \sum_{j \notin X} e^{-d_{i,j}} n_j + c_{\tau,2p} \right)^p \right] \\ & \leq \left(4q_0^3 c_{\tau,1} e^{-d_{i,X}} + c_{\tau,1} e^{-d_{i,X}} |X| C_p^{1/p} + 2^{D+1} D! c_{\tau,1} C_p^{1/p} + c_{\tau,2p} \right)^p \\ & = C_p \left[C_p^{-1/p} \left(4q_0^3 c_{\tau,1} e^{-d_{i,X}} + c_{\tau,2p} \right) + c_{\tau,1} e^{-d_{i,X}} |X| + 2^{D+1} D! c_{\tau,1} \right]^p, \end{aligned} \quad (4.28)$$

where we use

$$\sum_{j \notin X} e^{-d_{i,j}} \leq 1 + \sum_{s=1}^{\infty} \sum_{j: j \in \partial i[s]} e^{-s} \leq 1 + \sum_{s=1}^{\infty} \gamma s^{D-1} e^{-s} \leq 1 + \frac{1 + 2^D D!}{e} \leq 2^{D+1} D!. \quad (4.29)$$

In the third inequality above, we utilize an elementary bound proved in [KVS24, Lemma 1]. Therefore, by adopting the definition of $\mu_{d_{i,X}}$ as in Eq. (4.20), we prove the main inequality (4.19) by combining the inequalities (4.22), (4.26) and (4.28). This completes the proof. \square

4.2.3 Proof of Lemma 4.4

The proof is a polynomial analog of the one in Ref. [KVS24, Proposition 29]. We include the details for the convenience of the reader. First of all, we choose r such that $e^{-r/2} |X| \leq 1$. We then upper-bound $\mu_{d_{i,X}}$ ($i \in \tilde{X}$) by μ . Hence, using the inequality (4.19), we obtain

$$\text{tr} \left[O_X(\tau_2) \rho(\tau_1) O_X(\tau_2)^\dagger n_i^p \Pi_{n_i \geq x} \right] \leq \mu^p C_p, \quad (4.30)$$

which implies

$$\text{tr} \left[n_i^s \Pi_{n_i \geq x} O_X(\tau_2) \rho(\tau_1) O_X(\tau_2)^\dagger \right] = \text{tr} \left[n_i^{s-p} \Pi_{n_i \geq x} n_i^p O_X(\tau_2) \rho(\tau_1) O_X(\tau_2)^\dagger \right] \leq \mu^p C_p \bar{q}^{-p+s}. \quad (4.31)$$

By using the above inequality with $s = 0$, we immediately prove the first inequality (4.14) as follows:

$$\begin{aligned} \left\| \bar{\Pi}_{\tilde{X}, \bar{q}}^c O_X(\tau_2) \sqrt{\rho(\tau_1)} \right\|_F^2 &= \text{tr} \left[\bar{\Pi}_{\tilde{X}, \bar{q}}^c O_X(\tau_2) \rho(\tau_1) O_X(\tau_2)^\dagger \bar{\Pi}_{\tilde{X}, \bar{q}}^c \right] \leq \sum_{i \in \tilde{X}} \text{tr} \left[\Pi_{i, > \bar{q}} O_X(\tau_2) \rho(\tau_1) O_X(\tau_2)^\dagger \right] \\ &\leq |\tilde{X}| \mu^p C_p \bar{q}^{-p}, \end{aligned} \quad (4.32)$$

which yields the first main inequality (4.14).

Second, to prove the inequality (4.15), we first decompose

$$\bar{\Pi}_{\tilde{X}, \bar{q}} H_0 \bar{\Pi}_{\tilde{X}, \bar{q}}^c = \bar{\Pi}_{\tilde{X}, \bar{q}} \left(H_{0, \tilde{X}} + \partial h_{\tilde{X}} \right) \bar{\Pi}_{\tilde{X}, \bar{q}}^c, \quad (4.33)$$

where we use the definition $\partial h_{\tilde{X}} = H_0 - H_{0, \tilde{X}} - H_{0, \tilde{X}^c}$ for $\partial h_{\tilde{X}}$ and $\bar{\Pi}_{\tilde{X}, \bar{q}} H_{0, \tilde{X}^c} \bar{\Pi}_{\tilde{X}, \bar{q}}^c = H_{0, \tilde{X}^c} \bar{\Pi}_{\tilde{X}, \bar{q}} \bar{\Pi}_{\tilde{X}, \bar{q}}^c = 0$. The operator $\partial h_{\tilde{X}}$ is the summation of the hopping operator between the regions L and L^c . Then, by using

$$|b_i b_j^\dagger| = \sqrt{(b_i b_j^\dagger)^\dagger b_i b_j^\dagger} = \sqrt{n_i (n_j + 1)} \leq n_i + n_j, \quad (4.34)$$

and

$$\bar{\Pi}_{\tilde{X},\bar{q}} b_i b_j^\dagger \bar{\Pi}_{\tilde{X},\bar{q}}^c = \bar{\Pi}_{\tilde{X},\bar{q}} b_i b_j^\dagger \bar{\Pi}_{i,j,\bar{q}}^c, \quad \bar{\Pi}_{i,j,\bar{q}} := \begin{cases} \prod_{n_i \leq \bar{q}} \prod_{n_j \leq \bar{q}} & \text{for } i, j \in \tilde{X}, \\ \prod_{n_i \leq \bar{q}} & \text{for } i \in \tilde{X}, j \in \tilde{X}^c, \end{cases} \quad (4.35)$$

we obtain

$$\begin{aligned} \left\| \bar{\Pi}_{\tilde{X},\bar{q}} H_0 \bar{\Pi}_{\tilde{X},\bar{q}}^c O_X(\tau_2) \sqrt{\rho(\tau_1)} \right\|_F &\leq \bar{J} \sum_{i \in \tilde{X}} \sum_{j: d_{i,j}=1} \left\| \bar{\Pi}_{\tilde{X},\bar{q}} (b_i b_j^\dagger + \text{h.c.}) \bar{\Pi}_{i,j,\bar{q}}^c O_X(\tau_2) \sqrt{\rho(\tau_1)} \right\|_F \\ &\leq \bar{J} \sum_{i \in \tilde{X}} \sum_{j: d_{i,j}=1} \left\| (|b_i b_j^\dagger| + |b_j b_i^\dagger|) \bar{\Pi}_{i,j,\bar{q}}^c O_X(\tau_2) \sqrt{\rho(\tau_1)} \right\|_F \\ &\leq 2\bar{J} \sum_{i \in \tilde{X}} \sum_{j: d_{i,j}=1} \left\| (n_i + n_j) \bar{\Pi}_{i,j,\bar{q}}^c O_X(\tau_2) \sqrt{\rho(\tau_1)} \right\|_F. \end{aligned} \quad (4.36)$$

By slightly changing the inequality (4.31), we have

$$\begin{aligned} \text{tr} \left[(n_i + n_j)^2 \bar{\Pi}_{i,j,\bar{q}}^c O_X(\tau_2) \rho(\tau_1) O_X(\tau_2)^\dagger \right] &= \text{tr} \left[(n_i + n_j)^{2-p} \bar{\Pi}_{i,j,\bar{q}}^c (n_i + n_j)^p O_X(\tau_2) \rho(\tau_1) O_X(\tau_2)^\dagger \right] \\ &\leq \bar{q}^{2-p} \text{tr} \left[(n_i + n_j)^p O_X(\tau_2) \rho(\tau_1) O_X(\tau_2)^\dagger \right] \\ &\leq \bar{q}^{2-p} (2\mu)^p C_p, \end{aligned} \quad (4.37)$$

where we use $\left\| \rho n_i^s n_j^{p-s} \right\|_1 \leq \left\| \rho n_i^p \right\|_1^{s/p} \left\| \rho n_j^p \right\|_1^{1-s/p}$ for an arbitrary quantum state ρ .

By combining the inequalities (4.36) and (4.37), we arrive at the inequality (4.15) as follows:

$$\begin{aligned} \left\| \bar{\Pi}_{\tilde{X},\bar{q}} H_0 \bar{\Pi}_{\tilde{X},\bar{q}}^c O_X(\tau_2) \sqrt{\rho(\tau_1)} \right\|_F &\leq 2\bar{J} \sum_{i \in \tilde{X}} \sum_{j: d_{i,j}=1} \bar{q} \left(\frac{2\mu}{\bar{q}} \right)^{p/2} C_p^{1/2} \\ &\leq 2\gamma \bar{J} \bar{q} |\tilde{X}| \left(\frac{2\mu}{\bar{q}} \right)^{p/2} C_p^{1/2}, \end{aligned} \quad (4.38)$$

where we use $\sum_{j: d_{i,j}=1} 1 \leq |i[1]| \leq \gamma$ for $\forall i \in \Lambda$. We thus obtain the second main inequality (4.15). This completes the proof. \square

5 Proof of Subtheorem 3.2

5.1 Overview of the proof

For the proof of Subtheorem 3.2 compared to Subtheorem 3.1, the only difference arises from the estimation of the error using the effective Hamiltonian $\bar{\Pi}_{\tilde{X},\bar{q}} H \bar{\Pi}_{\tilde{X},\bar{q}}$, which was previously given by Proposition 4.2. For the expectation, we first prove the following statement.

Lemma 5.1. *Let ρ be an arbitrary quantum state. Then, for an arbitrary projection operator $\bar{\Pi}$, the dynamics of the effective Hamiltonian $\bar{\Pi} H \bar{\Pi}$ is deviated from the true dynamics by the error of*

$$\begin{aligned} &\text{tr} \left\{ \left[O(H, \tau) \rho - O(\bar{\Pi} H \bar{\Pi}, \tau) \right] \rho \right\} \\ &\leq \left\| \bar{\Pi}^c \sqrt{\rho} \right\|_F \left(\left\| \bar{\Pi}^c e^{iH\tau} O e^{-iH\tau} \sqrt{\rho} \right\|_F + \left\| \bar{\Pi}^c O e^{-iH\tau} \sqrt{\rho} \right\|_F + 2 \left\| \bar{\Pi}^c O e^{-i\tilde{H}\tau} \sqrt{\rho} \right\|_F \right) \\ &+ \int_0^\tau \left\| \bar{\Pi}^c H_0 \bar{\Pi} e^{-i\tilde{H}\tau_1} \sqrt{\rho} \right\|_F \left\| \bar{\Pi}^c e^{iH(\tau-\tau_1)} O e^{-iH\tau} \sqrt{\rho} \right\|_F + \left\| \bar{\Pi}^c H_0 \bar{\Pi} e^{-iH\tau_1} \sqrt{\rho} \right\|_F \left\| \bar{\Pi}^c O e^{-i\tilde{H}\tau} \sqrt{\rho} \right\|_F d\tau_1, \end{aligned} \quad (5.1)$$

where $\|\cdot\|_F$ is the Frobenius norm, i.e., $\|O\|_F = \sqrt{\text{tr}(OO^\dagger)}$.

This result is similar to Lemma 4.3 but it leverages cyclicity of the trace to perform Cauchy-Schwarz in a slightly different way that will turn out to be beneficial: Under the integral, the term containing H_0 is separated from the term with the observable O . This will allow us to improve the bound (4.15) to the following Lemma 5.2.

Lemma 5.2. *Under the assumptions of Subtheorem 3.2, we have*

$$\begin{aligned} \left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c H_0 \bar{\Pi}_{\tilde{X},\tilde{q}} e^{-i\tilde{H}\tau_1} \sqrt{\rho} \right\|_F &\leq 4^p \cdot 2\bar{J}\gamma\tilde{q}^{-1-p/2} \sqrt{|\tilde{X}|\gamma 2^D (2^p + \mu^p C_p)} \\ &= 4^p \cdot 2\bar{J}\gamma\tilde{q} |\tilde{X}|^{1/2} C_p^{1/2} \left(\frac{\mu}{\tilde{q}}\right)^{p/2} \sqrt{\gamma 2^D \left(\frac{2^p}{C_p \mu^p} + 1\right)}. \end{aligned} \quad (5.2)$$

The same inequality for the norm of $\left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c H_0 \bar{\Pi}_{\tilde{X},\tilde{q}} e^{-iH\tau_1} \sqrt{\rho} \right\|_F$.

We postpone the proofs of Lemmas 5.1 and 5.2 for now.

Proof of Subtheorem 3.2 assuming Lemmas 5.1 and 5.2. We apply Lemma 5.1. We then need to estimate the terms that appear in the RHS of (5.1). We have already proved in Lemma 4.4 the following inequality:

$$\left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c O_X(\tau_2) \sqrt{\rho(\tau_1)} \right\|_F \leq |\tilde{X}|^{1/2} C_p^{1/2} \left(\frac{\mu}{\tilde{q}}\right)^{p/2}. \quad (5.3)$$

Note that Lemma 4.5 also holds for the time evolution with respect to the effective Hamiltonian $\tilde{H} = \bar{\Pi}_{\tilde{X},\tilde{q}} H \bar{\Pi}_{\tilde{X},\tilde{q}}$. That is, we also have

$$\left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c O_X(\tau_2, \bar{\Pi}_{\tilde{X},\tilde{q}} H \bar{\Pi}_{\tilde{X},\tilde{q}}) \sqrt{\rho(\tau_1)} \right\|_F \leq |\tilde{X}|^{1/2} C_p^{1/2} \left(\frac{\mu}{\tilde{q}}\right)^{p/2}. \quad (5.4)$$

By applying these inequalities to (5.1), we obtain

$$\begin{aligned} &\left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c \sqrt{\rho} \right\|_F \left(\left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c e^{iH\tau} O e^{-iH\tau} \sqrt{\rho} \right\|_F + \left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c O e^{-iH\tau} \sqrt{\rho} \right\|_F + 2 \left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c O e^{-i\tilde{H}\tau} \sqrt{\rho} \right\|_F \right) \\ &\leq 4|\tilde{X}| C_p \left(\frac{\mu}{\tilde{q}}\right)^p, \end{aligned} \quad (5.5)$$

and

$$\left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c H_0 \bar{\Pi}_{\tilde{X},\tilde{q}} e^{-i\tilde{H}\tau_1} \sqrt{\rho} \right\|_F \left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c e^{iH(\tau-\tau_1)} O e^{-iH\tau} \sqrt{\rho} \right\|_F \quad (5.6)$$

$$\begin{aligned} &+ \left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c H_0 \bar{\Pi}_{\tilde{X},\tilde{q}} e^{-iH\tau_1} \sqrt{\rho} \right\|_F \left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c O e^{-i\tilde{H}\tau} \sqrt{\rho} \right\|_F \\ &\leq |\tilde{X}|^{1/2} C_p^{1/2} \left(\frac{\mu}{\tilde{q}}\right)^{p/2} \left(\left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c H_0 \bar{\Pi}_{\tilde{X},\tilde{q}} e^{-i\tilde{H}\tau_1} \sqrt{\rho} \right\|_F + \left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c H_0 \bar{\Pi}_{\tilde{X},\tilde{q}} e^{-iH\tau_1} \sqrt{\rho} \right\|_F \right). \end{aligned} \quad (5.7)$$

Thus, the remaining problem is to estimate the terms of

$$\left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c H_0 \bar{\Pi}_{\tilde{X},\tilde{q}} e^{-i\tilde{H}\tau_1} \sqrt{\rho} \right\|_F, \quad \left\| \bar{\Pi}_{\tilde{X},\tilde{q}}^c H_0 \bar{\Pi}_{\tilde{X},\tilde{q}} e^{-iH\tau_1} \sqrt{\rho} \right\|_F. \quad (5.8)$$

For these, we use Lemma 5.2. Hence, from inequality (5.6), we conclude

$$\begin{aligned} & \left\| \bar{\Pi}_{\tilde{X}, \bar{q}}^c H_0 \bar{\Pi} e^{-i\tilde{H}\tau_1} \sqrt{\rho} \right\|_F \left\| \bar{\Pi}_{\tilde{X}, \bar{q}}^c e^{iH(\tau-\tau_1)} O e^{-iH\tau} \sqrt{\rho} \right\|_F + \left\| \bar{\Pi}_{\tilde{X}, \bar{q}}^c H_0 \bar{\Pi}_{\tilde{X}, \bar{q}} e^{-iH\tau_1} \sqrt{\rho} \right\|_F \left\| \bar{\Pi}_{\tilde{X}, \bar{q}}^c O e^{-i\tilde{H}\tau} \sqrt{\rho} \right\|_F \\ & \leq 4^{p+1} \bar{J} \gamma \bar{q} |\tilde{X}| C_p \left(\frac{\mu}{\bar{q}} \right)^p \sqrt{\gamma^{2D} \left(\frac{2^p}{C_p \mu^p} + 1 \right)}. \end{aligned} \quad (5.9)$$

By combining the inequalities (5.5) and (5.9), we obtain the error of the dynamics by the effective Hamiltonian:

$$\text{tr} \left\{ \left[O(H, \tau) \rho - O(\bar{\Pi}_{\tilde{X}, \bar{q}} H \bar{\Pi}_{\tilde{X}, \bar{q}}, \tau) \right] \rho \right\} \leq 4 |\tilde{X}| C_p \left(\frac{\mu}{\bar{q}} \right)^p \left[1 + \tau \bar{J} \gamma \bar{q} \sqrt{\gamma^{2D} \left(\frac{2^p}{C_p \mu^p} + 1 \right)} \right]. \quad (5.10)$$

By applying the choice of $\bar{q} = \eta r / 2$ (see Proposition 4.1), we have

$$4 |\tilde{X}| C_p \left(\frac{\mu}{\bar{q}} \right)^p \left[1 + \tau \bar{J} \gamma \bar{q} \sqrt{\gamma^{2D} \left(\frac{2^p}{C_p \mu^p} + 1 \right)} \right] \quad (5.11)$$

$$\begin{aligned} & = 4 |\tilde{X}| C_p \left(\frac{2\mu}{\eta r} \right)^p \left[1 + \frac{\tau \bar{J} \gamma \eta r}{2} \sqrt{\gamma^{2D} \left(\frac{2^p}{C_p \mu^p} + 1 \right)} \right] \\ & \leq \tilde{C}_p |\tilde{X}| r^{-p+1}. \end{aligned} \quad (5.12)$$

We thus prove the main inequality (3.18) in Suthorem 3.2. This completes the proof. \square

5.1.1 Proof of Lemma 5.1

We first denote

$$\bar{\Pi}^c = 1 - \bar{\Pi}, \quad \tilde{H} = \bar{\Pi} H \bar{\Pi} \quad (5.13)$$

for simplicity. We start from the elementary bound

$$\begin{aligned} & \left| \text{tr} \left\{ \left[O(H, \tau) - O(\tilde{H}, \tau) \right] \rho \right\} \right| \\ & \leq \left| \text{tr} \left[\left(e^{iH\tau} O e^{-iH\tau} - e^{i\tilde{H}\tau} O e^{-iH\tau} \right) \rho \right] \right| + \left| \text{tr} \left[\left(e^{i\tilde{H}\tau} O e^{-i\tilde{H}\tau} - e^{iH\tau} O e^{-i\tilde{H}\tau} \right) \rho \right] \right|, \end{aligned} \quad (5.14)$$

where we take the complex conjugate for the second term in the last inequality. In order to estimate the RHS in the inequality above, we need to upper-bound

$$\left| \text{tr} \left[\left(e^{iH\tau} O e^{-iH\tau} - e^{i\tilde{H}\tau} O e^{-iH\tau} \right) \rho \right] \right| \quad (5.15)$$

for an arbitrary O . We note that by replacing H and \tilde{H} the second term in the RHS of (5.14) also has the form of (5.15). For the estimation of (5.15), we begin with the inequality of

$$\begin{aligned} & \left| \text{tr} \left[\left(e^{iH\tau} O e^{-iH\tau} - e^{i\tilde{H}\tau} O e^{-iH\tau} \right) \rho \right] \right| \\ & \leq \left| \text{tr} \left[\bar{\Pi}^c \left(e^{iH\tau} O e^{-iH\tau} - e^{i\tilde{H}\tau} O e^{-iH\tau} \right) \rho \bar{\Pi}^c \right] \right| + \left| \text{tr} \left[\bar{\Pi} \left(e^{iH\tau} O e^{-iH\tau} - e^{i\tilde{H}\tau} O e^{-iH\tau} \right) \rho \right] \right| \\ & \leq \left\| \bar{\Pi}^c \sqrt{\rho} \right\|_F \left(\left\| \bar{\Pi}^c e^{iH\tau} O e^{-iH\tau} \sqrt{\rho} \right\|_F + \left\| \bar{\Pi}^c O e^{-iH\tau} \sqrt{\rho} \right\|_F \right) + \left| \text{tr} \left[\bar{\Pi} \left(e^{iH\tau} O - e^{i\tilde{H}\tau} O \right) e^{-iH\tau} \rho \right] \right|, \end{aligned} \quad (5.16)$$

where, in the second inequality, we use the Cauchy-Schwarz inequality as $\text{tr}(A^\dagger B) \leq \|A\|_F \|B\|_F$ and

$$\left\| \bar{\Pi}^c e^{i\tilde{H}\tau} O e^{-iH\tau} |\psi\rangle \right\| = \left\| \bar{\Pi}^c O e^{-iH\tau} |\psi\rangle \right\| \quad (5.17)$$

from the relation of $\bar{\Pi}^c \tilde{H} = \bar{\Pi}^c \bar{\Pi} H \bar{\Pi} = 0$.

The third term in the RHS of (5.16) is treated as follows. We start from

$$e^{iH\tau} O - e^{i\tilde{H}\tau} O = -i \int_0^\tau e^{i\tilde{H}\tau_1} (\tilde{H} - H) e^{iH(\tau-\tau_1)} O d\tau_1. \quad (5.18)$$

Furthermore, we decompose $\bar{\Pi} (\tilde{H} - H)$ as

$$\bar{\Pi} (\tilde{H} - H) = \bar{\Pi} H \bar{\Pi} - \bar{\Pi} H (\bar{\Pi} + 1 - \bar{\Pi}) = -\bar{\Pi} H \bar{\Pi}^c = -\bar{\Pi} H_0 \bar{\Pi}^c, \quad (5.19)$$

where we use $[\bar{\Pi}, \tilde{H}] = 0$ and $[V, \bar{\Pi}] = 0$ which is derived from $[n_i, \bar{\Pi}] = 0$ for $\forall i \in \Lambda$. We recall that $\bar{\Pi}^c$ has been defined as $\bar{\Pi}^c := 1 - \bar{\Pi}$ as in Eq. (5.13). Therefore, from Eq. (5.18), we have

$$\begin{aligned} \left| \text{tr} \left[\bar{\Pi} \left(e^{iH\tau} O - e^{i\tilde{H}\tau} O \right) e^{-iH\tau} \rho \right] \right| &= \left| \text{tr} \left(\int_0^\tau e^{i\tilde{H}\tau_1} \bar{\Pi} H_0 \bar{\Pi}^c e^{iH(\tau-\tau_1)} O e^{-iH\tau} \rho \right) d\tau_1 \right| \\ &\leq \int_0^\tau \left| \text{tr} \left(e^{i\tilde{H}\tau_1} \bar{\Pi} H_0 \bar{\Pi}^c e^{iH(\tau-\tau_1)} O e^{-iH\tau} \rho \right) \right| d\tau_1 \\ &\leq \int_0^\tau \left\| \sqrt{\rho} e^{i\tilde{H}\tau_1} \bar{\Pi} H_0 \bar{\Pi}^c \right\|_F \left\| \bar{\Pi}^c e^{iH(\tau-\tau_1)} O e^{-iH\tau} \sqrt{\rho} \right\|_F d\tau_1, \end{aligned} \quad (5.20)$$

which reduces the inequality (5.16) to

$$\begin{aligned} \left| \text{tr} \left[\left(e^{iH\tau} O e^{-iH\tau} - e^{i\tilde{H}\tau} O e^{-i\tilde{H}\tau} \right) \rho \right] \right| &\leq \left\| \bar{\Pi}^c \sqrt{\rho} \right\|_F \left(\left\| \bar{\Pi}^c e^{iH\tau} O e^{-iH\tau} \sqrt{\rho} \right\|_F + \left\| \bar{\Pi}^c O e^{-iH\tau} \sqrt{\rho} \right\|_F \right) \\ &\quad + \int_0^\tau \left\| \bar{\Pi}^c H_0 \bar{\Pi} e^{-i\tilde{H}\tau_1} \sqrt{\rho} \right\|_F \left\| \bar{\Pi}^c e^{iH(\tau-\tau_1)} O e^{-iH\tau} \sqrt{\rho} \right\|_F d\tau_1. \end{aligned} \quad (5.21)$$

In the same way, we can derive

$$\begin{aligned} \left| \text{tr} \left[\left(e^{i\tilde{H}\tau} O e^{-i\tilde{H}\tau} - e^{iH\tau} O e^{-iH\tau} \right) \rho \right] \right| &\leq 2 \left\| \bar{\Pi}^c \sqrt{\rho} \right\|_F \left\| \bar{\Pi}^c O e^{-i\tilde{H}\tau} \sqrt{\rho} \right\|_F \\ &\quad + \int_0^\tau \left\| \bar{\Pi}^c H_0 \bar{\Pi} e^{-iH\tau_1} \sqrt{\rho} \right\|_F \left\| \bar{\Pi}^c O e^{-i\tilde{H}\tau} \sqrt{\rho} \right\|_F d\tau_1 \end{aligned} \quad (5.22)$$

by replacing H and \tilde{H} in the inequality (5.21), where we use $[\bar{\Pi}, \tilde{H}] = 0$. Therefore, by applying the inequalities (5.21) and (5.22) to (5.14), we obtain the desired inequality (5.1). This completes the proof. \square

5.1.2 Proof of Lemma 5.2

Our task is to estimate the following quantity:

$$\left\| \bar{\Pi}_{\tilde{X}, \tilde{q}}^c H_0 \bar{\Pi}_{\tilde{X}, \tilde{q}} e^{-i\tilde{H}\tau_1} \sqrt{\rho} \right\|_F^2 = \text{tr} \left[\bar{\Pi}_{\tilde{X}, \tilde{q}} H_0 \bar{\Pi}_{\tilde{X}, \tilde{q}}^c H_0 \bar{\Pi}_{\tilde{X}, \tilde{q}} \rho(\tilde{H}, \tau_1) \right]. \quad (5.23)$$

By using the relation (4.35), i.e.,

$$\bar{\Pi}_{\tilde{X},\bar{q}} b_i b_j^\dagger \bar{\Pi}_{\tilde{X},\bar{q}}^c = \bar{\Pi}_{\tilde{X},\bar{q}} b_i b_j^\dagger \bar{\Pi}_{i,j,\bar{q}}^c, \quad \bar{\Pi}_{i,j,\bar{q}} := \begin{cases} \prod_{n_i \leq \bar{q}} \prod_{n_j \leq \bar{q}} & \text{for } i, j \in \tilde{X}, \\ \prod_{n_i \leq \bar{q}} & \text{for } i \in \tilde{X}, j \in \tilde{X}^c, \end{cases} \quad (5.24)$$

we obtain

$$\begin{aligned} \bar{\Pi}_{\tilde{X},\bar{q}} H_0 \bar{\Pi}_{\tilde{X},\bar{q}}^c H_0 \bar{\Pi}_{\tilde{X},\bar{q}} &= \sum_{i,j \in \tilde{X}} \sum_{k,l \in \tilde{X}} J_{i,j} J_{k,l} \bar{\Pi}_{\tilde{X},\bar{q}} (b_i b_j^\dagger + \text{h.c.}) \bar{\Pi}_{\tilde{X},\bar{q}}^c \cdot \bar{\Pi}_{\tilde{X},\bar{q}}^c (b_k b_l^\dagger + \text{h.c.}) \bar{\Pi}_{\tilde{X},\bar{q}} \\ &= \sum_{i,j \in \tilde{X}} \sum_{k,l \in \tilde{X}} J_{i,j} J_{k,l} \bar{\Pi}_{\tilde{X},\bar{q}} (b_i b_j^\dagger + \text{h.c.}) \bar{\Pi}_{i,j,\bar{q}}^c \cdot \bar{\Pi}_{k,l,\bar{q}}^c (b_k b_l^\dagger + \text{h.c.}) \bar{\Pi}_{\tilde{X},\bar{q}} \\ &= \sum_{\substack{i,j,k,l \\ i \in \tilde{X}, k \in \tilde{X}, \{i,j\} \cap \{k,l\} \neq \emptyset}} J_{i,j} J_{k,l} \bar{\Pi}_{\tilde{X},\bar{q}} (b_i b_j^\dagger + \text{h.c.}) \bar{\Pi}_{i,j,\bar{q}}^c \cdot \bar{\Pi}_{k,l,\bar{q}}^c (b_k b_l^\dagger + \text{h.c.}) \bar{\Pi}_{\tilde{X},\bar{q}}, \end{aligned} \quad (5.25)$$

where we use $[\bar{\Pi}_{i,j,\bar{q}}^c, \bar{\Pi}_{k,l,\bar{q}}^c] = 0$ and $[b_i b_j^\dagger, \bar{\Pi}_{k,l,\bar{q}}^c] = 0$ for $\{i, j\} \cap \{k, l\} = \emptyset$. Note that $\bar{\Pi}_{\tilde{X},\bar{q}} \bar{\Pi}_{k,l,\bar{q}}^c = 0$.

Each of the terms in the summation of the RHS of (5.25) is upper-bounded by

$$\begin{aligned} &\left| \text{tr} \left\{ \left[J_{i,j} J_{k,l} \bar{\Pi}_{\tilde{X},\bar{q}} (b_i b_j^\dagger + \text{h.c.}) \bar{\Pi}_{i,j,\bar{q}}^c \cdot \bar{\Pi}_{k,l,\bar{q}}^c (b_k b_l^\dagger + \text{h.c.}) \bar{\Pi}_{\tilde{X},\bar{q}} \right] \rho(\tilde{H}, \tau_1) \right\} \right| \\ &\leq \bar{J}^2 \left\| \bar{\Pi}_{i,j,\bar{q}}^c (b_i b_j^\dagger + \text{h.c.}) \sqrt{\rho(\tilde{H}, \tau_1)} \right\|_F \cdot \left\| \bar{\Pi}_{k,l,\bar{q}}^c (b_k b_l^\dagger + \text{h.c.}) \sqrt{\rho(\tilde{H}, \tau_1)} \right\|_F. \end{aligned} \quad (5.26)$$

We then consider

$$\begin{aligned} &\left\| \bar{\Pi}_{i,j,\bar{q}}^c b_i b_j^\dagger \sqrt{\rho(\tilde{H}, \tau_1)} \right\|_F^2 = \text{tr} \left(\bar{\Pi}_{i,j,\bar{q}}^c b_i b_j^\dagger \rho(\tilde{H}, \tau_1) b_i^\dagger b_j \right) \\ &= \text{tr} \left[(n_i + n_j)^{-p+2} \bar{\Pi}_{i,j,\bar{q}}^c (n_i + n_j)^p \frac{1}{n_i + n_j} b_i b_j^\dagger \rho(\tilde{H}, \tau_1) b_i^\dagger b_j \frac{1}{n_i + n_j} \right] \\ &\leq \bar{q}^{2-p} \text{tr} \left[(n_i + n_j)^p \left(\frac{1}{n_i + n_j} b_i b_j^\dagger \right) \rho(\tilde{H}, \tau_1) \left(\frac{1}{n_i + n_j} b_i b_j^\dagger \right)^\dagger \right], \end{aligned} \quad (5.27)$$

where we can ensure

$$\left\| \frac{1}{n_i + n_j} b_i b_j^\dagger \right\| \leq 1 \quad (5.28)$$

because of $|b_i b_j^\dagger| \leq n_i + n_j$ from the inequality (4.34). Using Lemma 4.7 by letting

$$O_q \rightarrow \left(\frac{1}{n_i + n_j} b_i b_j^\dagger \right)^\dagger, \quad q \rightarrow 1, \quad \sum_{j \in \Lambda} \nu_j n_j \rightarrow n_i + n_j, \quad \bar{\nu} \rightarrow 1, \quad L \rightarrow \{i, j\} \quad (5.29)$$

in the inequality (4.24), we obtain

$$\begin{aligned} \text{tr} \left[(n_i + n_j)^p \left(\frac{1}{n_i + n_j} b_i b_j^\dagger \right) \rho(\tilde{H}, \tau_1) \left(\frac{1}{n_i + n_j} b_i b_j^\dagger \right)^\dagger \right] &\leq 4^p \text{tr} \left[(4 + n_i + n_j)^p \rho(\tilde{H}, \tau_1) \right] \\ &\leq 8^p \text{tr} \left\{ [4^p + (n_i + n_j)^p] \rho(\tilde{H}, \tau_1) \right\} \\ &\leq 8^p [4^p + (2\mu)^p C_p], \end{aligned} \quad (5.30)$$

where we use (4.37) in the last inequality. By combining the inequalities (5.27) and (5.30), we obtain

$$\left\| \bar{\Pi}_{i,j,\bar{q}}^c b_i b_j^\dagger \sqrt{\rho(\tilde{H}, \tau_1)} \right\|_F \leq \bar{q}^{1-p/2} 8^{p/2} [4^p + (2\mu)^p C_p]^{1/2}, \quad (5.31)$$

which reduces the inequality (5.26)

$$\left| \text{tr} \left\{ \left[J_{i,j} J_{k,l} \bar{\Pi}_{\tilde{X},\bar{q}} (b_i b_j^\dagger + \text{h.c.}) \bar{\Pi}_{i,j,\bar{q}}^c \cdot \bar{\Pi}_{k,l,\bar{q}}^c (b_k b_l^\dagger + \text{h.c.}) \bar{\Pi}_{\tilde{X},\bar{q}} \right] \rho(\tilde{H}, \tau_1) \right\} \right| \leq 4\bar{J}^2 8^p [4^p + (2\mu)^p C_p] \bar{q}^{2-p}. \quad (5.32)$$

Finally, by using the assumption on the graph geometry, cf. (2.2), we have

$$\sum_{\substack{i,j,k,l \\ i \in \tilde{X}, k \in \tilde{X}, \{i,j\} \cap \{k,l\} \neq \emptyset}} 1 \leq \sum_{i \in \tilde{X}} \sum_{j: d_{i,j}=1} \sum_{k: d_{i,k} \leq 2} \sum_{l: d_{k,l}=1} 1 \leq |\tilde{X}| \gamma^3 2^D, \quad (5.33)$$

which yields

$$\left\| \bar{\Pi}_{\tilde{X},\bar{q}}^c H_0 \bar{\Pi}_{\tilde{X},\bar{q}} e^{-i\tilde{H}\tau_1} \sqrt{\rho} \right\|_F^2 = \text{tr} \left[\bar{\Pi}_{\tilde{X},\bar{q}} H_0 \bar{\Pi}_{\tilde{X},\bar{q}}^c H_0 \bar{\Pi}_{\tilde{X},\bar{q}} \rho(\tilde{H}, \tau_1) \right] \quad (5.34)$$

$$\leq |\tilde{X}| \gamma^3 2^D \cdot 4\bar{J}^2 8^p [4^p + (2\mu)^p C_p] \bar{q}^{2-p}, \quad (5.35)$$

We have thus proved the main inequality (5.2). This completes the proof. \square

6 Conclusions

This work consider a class of Bose-Hubbard Hamiltonians with local interactions of the form n_i^p with $p > 2$ sufficiently large. We prove that general initial states that are translation-invariant and of uniformly bounded energy density satisfy an enhanced LRB compared to [KVS24]. This excludes scenarios for rapid information transport through boson accumulation for our class of Hamiltonians, cf. Figure 4a in [KVS24].

We recall that interactions of the form n_i^p with integer $p > 2$ arise physically from local p -body repulsion through the tight-binding approximation [MKM20]. Of course, the physically most relevant case is $p = 2$, which corresponds to two-body interactions in the standard Bose-Hubbard Hamiltonian, and our result does not cover this case. Addressing Questions 1 and 1' raised in the introduction for $p = 2$ will require a much finer dynamical analysis of the local energy flow and, relatedly, the dynamical build-up of higher particle-particle correlations for suitable initial states. Controlling these other local dynamical quantities will require new insights.

A different feature of our result is that the LRB on the expectation is appreciably stronger than the one on the trace norm. For long-range quantum spin systems, a similar phenomenon is observed [TCE⁺20]. Of course, the expectation value is bounded by the trace norm, but it could indeed be substantially smaller due to cancellations. While we do not expect such cancellations to occur generically, it is an interesting question under what circumstances the different t -scaling truly occurs dynamically, i.e., what kind of lower bounds can be proved. If this phenomenon persists at the level of lower bounds, this would add another unexpected wrinkle to the rich and subtle nature of bosonic information propagation, which would require refining Questions 1 and 1' above based on which measure of size is used.

Finally, looking beyond the specific model, our result places into center focus the significant qualitative rigidity effect that dynamical constraints, specifically symmetries (here, translation-invariance) and energy conservation, have for quantum dynamics of strongly interacting bosons. Our result is a rigorous manifestation of the associated rigidity concepts in the many-boson context. It would be interesting to explore the effect of different types of dynamical constraints, for instance, dipole moment conservation, which was recently studied in the Bose-Hubbard context in [LHS22].

Acknowledgments

The authors thank Jingxuan Zhang for useful comments on a draft version of the manuscript.

T.K. acknowledges the Hakubi projects of RIKEN. T.K. is supported by JST PRESTO (Grant No. JPMJPR2116), ERATO (Grant No. JPMJER2302), and JSPS Grants-in-Aid for Scientific Research (No. JP24H00071), Japan. The research of M.L. is partially supported by the DFG through the grant TRR 352 – Project-ID 470903074.

A Examples of translation-invariant quantum states with bounded energy density and significant boson concentration

We recall that, in proving Theorem 1.1, a key input was the polynomial tail bound on the local boson number distribution

$$\mathrm{tr} [\Pi_{n_i \geq q} \rho_0(t)] \lesssim \frac{1}{q^p}, \quad \text{for all } t. \quad (\text{A.1})$$

which followed via Markov’s inequality from the moment bound

$$\mathrm{tr} [n_i^p \rho_0(t)] \leq C, \quad \text{for all } t. \quad (\text{A.2})$$

These bounds are derived by using translation-invariance and energy conservation $\mathrm{tr}(\rho H) = \mathrm{tr}[\rho(t)H]$, cf. Lemma 2.6.

Any improvement of the tail bound (A.1) would yield an improved LR velocity. It is therefore a natural question whether our approach is sharp, i.e., whether (A.2) can be improved. (For short times, Proposition 2.7 yields a stronger upper bound, but as noted thereafter, this is insufficient for an improved LR bound.)

- Question 1: Is the current proof via Lemma 2.6 sharp, i.e., can we derive a stronger bound than (A.2) assuming translation-invariance and conservation of the first moment of the energy $\mathrm{tr}(\rho H) = \mathrm{tr}(\rho(t)H)$?
- Question 2: Can (A.2) be improved by also using conservation of the second moment of the Hamiltonian $\mathrm{tr}(\rho(t)H^2) = \mathrm{tr}(\rho H^2)$?

In this Appendix, we construct explicit quantum states which show that the answer to Question 1 is “Yes” and the answer to Question 2 is “No.” In this sense, the particular method we develop in this paper is sharp; i.e., it will be necessary to leverage other dynamical constraints to obtain further improvements to the bosonic LRB for translation-invariant initial states.

A.1 Construction of high boson occupation translation-invariant quantum state

In Proposition A.1 below, we construct an explicit translation-invariant quantum state which saturates the bound (A.2) while, at the same time, both $\frac{\text{tr}(\rho H)}{|\Lambda_R|}$ and $\frac{\text{tr}(\rho H^2)}{|\Lambda_R|^2}$ are arbitrarily small.

At the same time, there are many translation-invariant initial states of small particle density for which $\frac{\text{tr}(\rho H)}{|\Lambda_R|}$ and $\frac{\text{tr}(\rho H^2)}{|\Lambda_R|^2}$ are $\mathcal{O}(1)$ (We give a simple, explicit example after Proposition A.1 for completeness).

Therefore, using only translation-invariance and moment conservation does not allow to exclude that a well-behaved (i.e., one of small local particle density) initial quantum state time-evolves into a badly behaved quantum state (i.e., one that saturates (A.2)). In particular, the construction shows that *translation-invariance by itself* is certainly not enough for controlling the boson concentration better than in (A.2).

For simplicity, we work in two dimensions $D = 2$, but the construction generalizes to any dimension in an obvious way.

Proposition A.1. *Let $p \geq 2$ and let $\Lambda_R \subset \mathbb{Z}^2$ be a discrete torus of side length R . We choose the Hamiltonian H as a Bose-Hubbard Hamiltonian of the form*

$$H = J \sum_{\langle i,j \rangle} (b_i b_j^\dagger + \text{h.c.}) + U \sum_{i \in \Lambda_R} (n_i - 1)^p \quad (\text{A.3})$$

with periodic boundary conditions.

Then, there exists a constant $C > 0$ such that the following holds in the limits where $R, q \rightarrow \infty$ with $R \geq q^p$. There exists a bosonic quantum state ρ with the following properties.

(I) *The state ρ is translation invariant and has total particle number $|\Lambda_R|$ (i.e., the global particle density is $= 1$).*

(II) *The state ρ satisfies,*

$$\frac{\text{tr}(\rho H)}{|\Lambda_R|} = \mathcal{O}(q^{-p}), \quad \frac{\text{tr}(\rho H^2)}{|\Lambda_R|^2} = \mathcal{O}(q^{-p}), \quad \text{for } q \rightarrow \infty. \quad (\text{A.4})$$

(III) *For any site $i \in \Lambda_R$, we have*

$$\text{tr}(\rho \Pi_{n_i \geq q}) \geq \frac{C}{q^p}. \quad (\text{A.5})$$

A few remarks are in order.

(i) In assertion (II), the moments of H are normalized in a natural way to obtain finite numbers in the thermodynamic limits $R \rightarrow \infty$.

(ii) In the statement, we consider the boson-boson interaction in the form of $(n_i - 1)^p$, but this is just for convenience and the construction generalizes to many other forms. In fact, if we consider the Bose-Hubbard type interactions, e.g., $n_i^{p-1}(n_i - 1)$, we are able to improve the second order moment bound to $\text{tr}(\rho H^2) - [\text{tr}(\rho H)]^2 = \mathcal{O}(q^{-p})|\Lambda_R|$.

(iii) The error terms $\mathcal{O}(q^{-p})$ are mainly due to finite-size & number-theoretic effects. It is possible to obtain vanishing errors as $R \rightarrow \infty$ by a slightly modified, more involved construction.

(iv) The proof gives the following stronger version of assertion (III): Let $q = \ell^{1/p}$. Fix any site $i \in \Lambda_R$. For sufficiently large R , with probability larger than $1/\ell$, this site belongs to a one-dimensional path of length R with boson number $\sim \ell^{1/p}$ at each of the sites in the path. The fact that the state we construct not only has large occupation on individual sites but on entire one-dimensional paths shows that no improvement can be expected from using the kind of alternative refined LRB for bounded interactions derived in [KVS24] which takes into account the average occupation along a one-dimensional path.

For completeness, we give an explicit example of a well-behaved translation-invariant quantum state $|m\rangle$ (i.e., one of small local particle density) with bounded first two energy moments.

We define the subset

$$L_x = \{(x, y) \in \Lambda_R : y \in \mathbb{Z}\}. \quad (\text{A.6})$$

We use Dirac's bra-ket notation. We assume R is even and introduce the following Mott states

$$|m_{ev}\rangle = \bigotimes_{1 \leq x \leq R} |0\rangle_{L_{2x-1}} \otimes |2\rangle_{L_{2x}}, \quad |m_{odd}\rangle = \bigotimes_{1 \leq x \leq R} |2\rangle_{L_{2x-1}} \otimes |0\rangle_{L_{2x}}.$$

which both consist of alternating vertical strips of occupation numbers = 0 and = 2. Then we consider the state

$$|m\rangle = \frac{|m_{ev}\rangle + |m_{odd}\rangle}{\sqrt{2}}.$$

The state $|m\rangle$ is translation-invariant and has total particle number $|\Lambda_R|$. Moreover, it satisfies the moment bounds^{*3}

$$\frac{\langle m|H|m\rangle}{|\Lambda_R|} = U, \quad \frac{\langle m|H^2|m\rangle}{|\Lambda_R|^2} = U^2 + \frac{4J^2}{|\Lambda_R|}.$$

By Proposition A.1, using only translation-invariance and bounds on the first two moments, it is not possible to exclude that the time-evolved state $|m(t)\rangle$ has significantly accumulated bosons in the sense of (A.5).

A.2 Proof of Proposition A.1.

We first construct the preliminary state $|\psi_0\rangle$. Recall (A.6) and define the rectangles $L_{x:x'}$ by $\bigcup_{x_1 \in [x, x']} L_{x_1}$ (see also Fig. 2). Fix $\ell \geq 1$.

Let $R \geq \ell \geq 3$. We construct the preliminary quantum state $|\psi_0\rangle$ by alternating high-occupancy lines with the low-energy states from Lemma A.2 (see Fig. 2). Let $\gamma_0 \in (0, 1)$ and $q \geq 2$ to be determined later. For simplicity, we suppose that R is a multiple of ℓ . We set

$$|\psi_0\rangle = \bigotimes_{s=0}^{R/\ell-1} |qL_{s\ell}\rangle \otimes |\phi_{\gamma_0, L_{s\ell+1:(s+1)\ell-1}}\rangle, \quad (\text{A.9})$$

^{*3}Here, the second equality holds by

$$\begin{aligned} \langle m|H^2|m\rangle &= J^2 \sum_{\langle i,j \rangle} \langle m|(b_i b_j^\dagger + \text{h.c.})^2|m\rangle + U^2 R^4 \\ &= J^2 \sum_{\langle i,j \rangle} \langle m|n_i(n_j+1) + n_j(n_i+1)|m\rangle + U^2 R^4 \end{aligned} \quad (\text{A.7})$$

$$= 4J^2 R^2 + U^2 R^4. \quad (\text{A.8})$$




Figure 2: Schematic description of the construction of the preliminary quantum state $|\psi_0\rangle$ in Eq. (A.9). In $|\psi_0\rangle$, one-dimensional paths with high-boson density q periodically appear with spacing ℓ . Between these high-density paths, the quantum state is given by a low-energy state with small boson density $\gamma_0 < 1$. The state $|\psi_0\rangle$ is constructed so that the conditions in Eq. (A.4) are satisfied, but it is not translation invariant. Therefore we average it over translations through Eq. (A.13) so that ρ to obtain a translation-invariant state.

Here $|q_{L_{s\ell}}\rangle$ is the Mott state on the region L with the boson number q at the site of $i \in L_0$. The states on the intermediate rectangles $|\phi_{\gamma_0, L_{s\ell+1:(s+1)\ell-1}}\rangle$ are constructed via the following lemma.

Lemma A.2 (Low-energy states). *Let $\gamma_0 \in (0, 1)$. The following holds as $R \geq \ell \rightarrow \infty$. There exist $e_1, e_2 > 0$ and a quantum state $|\phi_{\gamma_0, L_{1:\ell-1}}\rangle$ supported on the subset $L_{1:\ell-1}$ with boson density $\gamma_0 + \mathcal{O}(R^{-1})$ such that the following energetic conditions are satisfied:*

- (a) $|\phi_{\gamma_0, L_{1:\ell-1}}\rangle$ commutes with translations in the vertical direction.
- (b) First-moment condition.

$$\frac{\langle \phi_{\gamma_0, L_{1:\ell-1}} | H_{L_{1:\ell-1}} | \phi_{\gamma_0, L_{1:\ell-1}} \rangle}{|L_{1:\ell-1}|} = -\gamma_0 e_1 + (1 - \gamma_0) e_2 + \mathcal{O}(\ell^{-1}). \quad (\text{A.10})$$

- (c) Second moment condition.

$$\frac{\langle \phi_{\gamma_0, L_{1:\ell-1}} | H_{L_{1:\ell-1}}^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle}{|L_{1:\ell-1}|^2} \leq [-\gamma_0 e_1 + (1 - \gamma_0) e_2]^2 + \mathcal{O}(\ell^{-1}). \quad (\text{A.11})$$

Below, we will choose $\gamma_0 \approx 1$ so that the state has negative energy. The proof idea is as follows: First, we construct a trial state of density-1 of strictly negative energy. Second, we consider the corresponding ground state and deplete it by adding particle-free regions to obtain a state satisfying the condition (a).

For now, we assume Lemma A.2 to hold and continue the proof of Proposition A.1. We link the two parameters q and γ_0 in the state $|\psi_0\rangle$ by the condition that the total boson density equals 1, i.e.,

$$q|L_0| + [\gamma_0 + \mathcal{O}(R^{-1})] |L_{1:\ell-1}| = |L_{0:\ell-1}|,$$

which we can rephrase as

$$q = (1 - \gamma_0)(\ell - 1) + 1 + \mathcal{O}(R^{-1}). \quad (\text{A.12})$$

We construct from $|\psi_0\rangle$ a translation invariant state by averaging it over translations up to order $\ell - 1$,

$$\rho := \frac{1}{\ell} \sum_{x=0}^{\ell-1} |\psi_x\rangle \langle \psi_x|, \quad |\psi_x\rangle := \bigotimes_{s=0}^{R/\ell-1} |q_{L_{x+s\ell}}\rangle \otimes |\phi_{\gamma_0, x+L_{s\ell+1}:x+(s+1)\ell-1}\rangle. \quad (\text{A.13})$$

By construction, the state is periodic in the horizontal direction, and by property (a) of Lemma A.2, it is periodic in the vertical direction. Therefore, it satisfies assertion (I) in Proposition A.1.

In the quantum state $|\psi\rangle$, the reduced density matrix on the line L_0 is given by

$$\rho_{L_0} = \frac{1}{\ell} |q_{L_0}\rangle \langle q_{L_0}| + \frac{\ell - 1}{\ell} \tilde{\rho}_{L_0}, \quad (\text{A.14})$$

where $\tilde{\rho}_{L_0}$ is constructed from the reduced density matrix of $|\phi_{\gamma_0, x+L_{s\ell+1}:x+(s+1)\ell-1}\rangle$. In particular, we see that assertion (III) holds^{*4} if we ensure that $q \geq C\ell^{1/p}$. Using (A.12), this can be rephrased as the constraint

$$(1 - \gamma_0)(\ell - 1) + 1 + \mathcal{O}(R^{-1}) \geq C\ell^{1/p}. \quad (\text{A.15})$$

For large ℓ and $\gamma_0 \approx 1$, this constraint is asymptotic to $\gamma_0 \gtrsim 1 - C\ell^{\frac{1-p}{p}}$.

Hence, to prove Proposition A.1, it remains to find a choice of ℓ so that simultaneously (A.15) holds and the energetic relations in (A.4) from assertion (II) are satisfied.

First moment. By translation-invariance, the average energy is given by

$$\text{tr}(\rho H) = \langle \psi_0 | H | \psi_0 \rangle. \quad (\text{A.16})$$

We decompose H as

$$H = \sum_{s=0}^{R/\ell-1} \left(H_{L_{s\ell+1:(s+1)\ell-1}} + \partial h_{L_{s\ell}} + V_{L_{s\ell}} \right), \quad (\text{A.17})$$

where $\partial h_{L_{s\ell}}$ are the hopping terms that act on the line $L_{s\ell}$. Note that $\langle \psi_0 | \partial h_{L_{s\ell}} | \psi_0 \rangle = 0$. We have

$$\langle \psi_0 | H | \psi_0 \rangle = \frac{R}{\ell} \left[(-\gamma_0 e_1 + (1 - \gamma_0) e_2 + \mathcal{O}(\ell^{-1}) \right] |L_{1:\ell-1}| + (q - 1)^p |L_0|. \quad (\text{A.18})$$

We introduce the following constraint on the parameter γ_0

$$[-\gamma_0 e_1 + (1 - \gamma_0) e_2] |L_{1:\ell-1}| + (q - 1)^p |L_0| = 0$$

Substituting for q by using the previous constraint (A.12), we rephrase this as

$$\gamma_0 e_1 \ell = (1 - \gamma_0)^p (\ell - 1)^p + [(1 - \gamma_0) e_2] \ell + \mathcal{O}(\ell^{-1}). \quad (\text{A.19})$$

^{*4}In fact, recalling the definition of $|\psi\rangle$, we see that $q \gtrsim \ell^{1/p}$ implies the stronger version of assertion (III) that the site belongs to an entire path of boson numbers $\gtrsim \ell^{1/p}$.

For large ℓ and $\gamma_0 \approx 1$, this amounts to $e_1 \ell \approx (1-\gamma_0)^p \ell^p$ or equivalently $\gamma_0 \approx 1 - e_1^{1/p} \ell^{\frac{1-p}{p}}$. Absorbing the case of small ℓ in the constant, we can ensure that (A.15) and (A.19) hold simultaneously. This establishes

$$\text{tr}(\rho H) = \mathcal{O}\left(\frac{R^2}{\ell}\right).$$

Second moment. By translation-invariance,

$$\text{tr}(\rho H^2) = \langle \psi_0 | H^2 | \psi_0 \rangle. \quad (\text{A.20})$$

For the state $|\psi_0\rangle$, we have

$$H |\psi_0\rangle = \sum_{s=0}^{R/\ell-1} \left(H_{L_{s\ell+1:(s+1)\ell-1}} + q^p |L_{s\ell}| \right) |\psi_0\rangle + \partial h_{L_{s\ell}} |\psi_0\rangle. \quad (\text{A.21})$$

Using this and $\langle \psi_0 | H_{L_{s\ell+1:(s+1)\ell-1}} \partial h_{L_{s\ell}} | \psi_0 \rangle = 0$, we find

$$\langle \psi_0 | H^2 | \psi_0 \rangle \quad (\text{A.22})$$

$$= \langle \psi_0 | H^2 | \psi_0 \rangle - \langle \psi_0 | H | \psi_0 \rangle^2 + \mathcal{O}\left(\frac{R^4}{\ell^2}\right) \quad (\text{A.23})$$

$$= \sum_{s=0}^{R/\ell-1} \left(\langle \phi_{\gamma_0, L_{s\ell+1:(s+1)\ell-1}} | H_{L_{s\ell+1:(s+1)\ell-1}}^2 | \phi_{\gamma_0, L_{s\ell+1:(s+1)\ell-1}} \rangle \right) \quad (\text{A.24})$$

$$+ 2q^p |L_{s\ell}| \cdot (-\gamma_0 e_1 + (1-\gamma_0)e_2 + \mathcal{O}(\ell^{-1})) + (q^p |L_{s\ell}|)^2 + \langle \psi_0 | \partial h_{L_{s\ell}}^2 | \psi_0 \rangle \Big) + \mathcal{O}\left(\frac{R^4}{\ell^2}\right)$$

$$= \frac{R}{\ell} \left(\langle \phi_{\gamma_0, L_{1:\ell-1}} | \Delta H_{L_{1:\ell-1}}^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle + \langle \psi_0 | \partial h_{L_\ell}^2 | \psi_0 \rangle \right) + \mathcal{O}\left(\frac{R^4}{\ell^2}\right), \quad (\text{A.25})$$

where we introduced

$$\Delta H_{L_{1:\ell-1}}^2 = H_{L_{1:\ell-1}}^2 - 2q^p |L_0| \cdot (-\gamma_0 e_1 + (1-\gamma_0)e_2) |L_{1:\ell-1}| + (q^p |L_0|)^2.$$

We estimate $\langle \phi_{\gamma_0, L_{1:\ell-1}} | \Delta H_{L_{1:\ell-1}}^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle$ via (A.11) in part (b) of Lemma A.2. By our choice (A.12) of q (i.e., $q \propto \ell^{1/p}$), the order $|L_{1:\ell-1}|^2$ terms cancel and so

$$\langle \phi_{\gamma_0, L_{1:\ell-1}} | \Delta H_{L_{1:\ell-1}}^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle = \mathcal{O}(R^2 \ell).$$

By Cauchy-Schwarz,

$$\begin{aligned} \langle \psi_0 | \partial h_{L_0}^2 | \psi_0 \rangle &\leq J^2 \sum_{\langle i,j \rangle | \{i,j\} \cap L_0 \neq \emptyset} \langle \psi_0 | n_i(n_j+1) + n_j(n_i+1) | \psi_0 \rangle \\ &\leq 3J^2 q(q+1) |L_0| \leq C J^2 \ell^{2/p} R. \end{aligned} \quad (\text{A.26})$$

It follows that $\langle \psi_0 | H^2 | \psi_0 \rangle \leq \mathcal{O}(R^3) + \mathcal{O}\left(\frac{R^4}{\ell^2}\right)$. This completes the proof of Proposition A.1. \square

A.3 Proof of Lemma A.2

The proof is comprised of two steps. First, we construct an explicit trial state of particle density 1 and negative energy expectation. Then we consider the density-1 ground state and lower its density to γ_0 by “dilution”, i.e., tensoring it with zero-occupancy regions in a suitable way. For technical

reasons (to control certain boundary terms via the number of particles on the boundary), we need translation-invariance of these subsystem states and for this reason, we use a periodic subsystem Hamiltonian, which again introduces other boundary terms; see Step 2 for the details.

Step 1. Let $\ell_0 \geq 4$. We consider the following trial state $|\varphi\rangle$ of particle density 1:

$$|\varphi\rangle := \bigotimes_{k=1}^R \left(\bigotimes_{j=1}^{\lfloor \ell_0/2 \rfloor} |\varphi_{(2j-1,k),(2j,k)}\rangle \otimes |1\rangle^{\ell_0/2 - \lfloor \ell_0/2 \rfloor} \right),$$

where k denotes the vertical coordinate and $|\varphi_{(2j-1,k),(2j,k)}\rangle$ is given in the particle occupation basis by

$$|\varphi_{(2j-1,k),(2j,k)}\rangle = \lambda_1 |1, 1\rangle - \lambda_2 \text{sign}(J)(|2, 0\rangle + |0, 2\rangle),$$

with $\lambda_1, \lambda_2 \geq 0$ and $\lambda_1^2 + 2\lambda_2^2 = 1$.

We consider the periodized Hamiltonian

$$H_{L_{1:\ell_0-1}}^{\text{per}} := H_{L_{1:\ell_0-1}} + \partial h_{\ell_0 \leftrightarrow 1}.$$

We calculate $\langle \varphi | H_{L_{1:\ell_0-1}} | \varphi \rangle$ and show that it is strictly negative and on the order of the volume $|L_{1:\ell_0-1}|$ for sufficiently large ℓ_0 .

For the potential energy, we have

$$\langle \varphi | U \sum_{x \in L_{1:\ell_0-1}} (n_x - 1)^2 | \varphi \rangle = 2UR\ell\lambda_2^2.$$

For the kinetic energy, we distinguish cases. Denote $h_{(j,k),(j',k')} = J(b_{(j,k)}^\dagger b_{(j',k')} + \text{h.c.})$. Straight-forward calculations in the bulk ($1 \leq j < \ell/2 - 1$) gives

$$\langle \varphi | h_{(2j,k),(2j+1,k)} | \varphi \rangle = \langle \varphi | h_{(2j,k),(2j,k+1)} | \varphi \rangle = 0.$$

That is, the only non-vanishing hopping terms are those “within” a single tensor factor $|\varphi_{(2j-1,k),(2j,k)}\rangle$. In the bulk, i.e., for $1 \leq j < \ell/2 - 1$, these are given by

$$\langle \varphi | h_{(2j-1,k),(2j,k)} | \varphi \rangle = -4|J|\lambda_1\lambda_2.$$

We estimate the hopping terms at the boundary by Cauchy-Schwarz and obtain

$$\begin{aligned} \langle \varphi | H_{L_{1:\ell_0-1}}^{\text{per}} | \varphi \rangle &\leq R\ell_0(2U\lambda_2^2 - 2|J|\lambda_1\lambda_2) + c(J,U)R \\ &= 2UR\ell_0(\lambda_2^2 - T_1\lambda_1\lambda_2) + c(J,U)R, \quad \text{with } T_1 = \frac{|J|}{U}. \end{aligned}$$

Minimizing over pairs (λ_1, λ_2) satisfying the constraint $\lambda_1^2 + 2\lambda_2^2 = 1$, we obtain

$$\min_{\lambda_1, \lambda_2} \langle \varphi | H_{L_{1:\ell_0-1}}^{\text{per}} | \varphi \rangle = \frac{UR\ell_0}{2} \left(1 - \sqrt{1 + 2T_1^2} \right) + c(J,U)R.$$

Hence, for all sufficiently large ℓ_0 , there exists $\tilde{\epsilon}_1 > 0$ such that

$$\inf \text{spec } H_{L_{1:\ell_0-1}}^{\text{per}} \leq -\tilde{\epsilon}_1 |L_{1:\ell_0-1}|.$$

The Hamiltonian $H_{L_{1:\ell_0-1}}^{\text{per}}$ commutes with $n_{L_{1:\ell_0-1}}$, the particle number on $L_{1:\ell_0-1}$ and so we can block-diagonalize it. Since the trial state $|\varphi\rangle$ satisfies $n_{L_{1:\ell_0-1}}|\varphi\rangle = |L_{1:\ell_0-1}||\varphi\rangle$ (it has global particle density 1), we may conclude

$$\inf \text{spec}(\Pi_{n_{L_{1:\ell_0-1}}=|L_{1:\ell_0-1}|} H_{L_{1:\ell_0-1}}^{\text{per}} \Pi_{n_{L_{1:\ell_0-1}}=|L_{1:\ell_0-1}|}) \leq -\tilde{e}_1 |L_{1:\ell_0-1}|. \quad (\text{A.27})$$

where $\Pi_{n_{L_{1:\ell_0-1}}=|L_{1:\ell_0-1}|}$ projects onto the spectral subspace on which $n_{L_{1:\ell_0-1}} = |L_{1:\ell_0-1}|$.

Step 2. Fix $\gamma_0 \in (0, 1)$. We set $\ell_0 = \lfloor \gamma_0 \ell \rfloor < \ell$. We assume that ℓ is large enough so that (A.27) holds. Using (A.27), the spectral theorem implies that exists a normalized ground state $|\xi_{1:\ell_0-1}\rangle$ of global particle density 1 and energy per volume $-e_1 \leq -\tilde{e}_1 < 0$, i.e.,

$$H_{L_{1:\ell_0-1}}^{\text{per}} |\xi_{1:\ell_0-1}\rangle = -e_1 |L_{1:\ell_0-1}| |\xi_{1:\ell_0-1}\rangle.$$

By block-diagonalizing $H_{L_{1:\ell_0-1}}^{\text{per}}$ with respect to translations in the vertical direction, we may assume that $|\xi_{1:\ell_0-1}\rangle$ is invariant under translations in the vertical direction up to a phase.

To construct $|\phi_{\gamma_0, 0:\ell-1}\rangle$, we dilute $|\xi_{1:\ell_0-1}\rangle$ with particle-free regions as follows.

$$|\phi_{\gamma_0, 0:\ell-1}\rangle := |\xi_{1:\ell_0-1}\rangle \otimes |0\rangle_{\ell_0:\ell-1}, \quad \text{with } |0\rangle_{\ell_0:\ell-1} := \bigotimes_{\ell_0 \leq j \leq \ell-1} \bigotimes_{1 \leq k \leq R} |0_{j,k}\rangle.$$

First, we note that the state $|\phi_{1:\ell}\rangle$ is invariant under translations in the vertical direction up to a phase, which ensures property (a) of Lemma A.2. Moreover, it has particle density $\frac{\ell_0}{\ell} = \gamma_0 + \mathcal{O}(\ell^{-1})$ as desired.

We check property (b). We abbreviate $H_{a:b} \equiv H_{L_{a:b}}$ and $H_{a:b}^{\text{per}} \equiv H_{L_{a:b}}^{\text{per}}$. The building blocks of $|\phi_{1:\ell}\rangle$ are all individually eigenstates of suitable local Hamiltonians,

$$\begin{aligned} H_{1:\ell_0-1}^{\text{per}} |\xi_{1:\ell_0-1}\rangle &= -e_1 \ell_0 R |\xi_{1:\ell_0-1}\rangle, \\ H_{\ell_0:\ell-2} |0\rangle_{\ell_0:\ell-1} &= U |0\rangle_{\ell_0:\ell-1}. \end{aligned} \quad (\text{A.28})$$

We write $H_{1:\ell-1} = H_{1:\ell_0-1}^{\text{per}} + H_{\ell_0:\ell-1} + \partial h_{\ell_0-1 \leftrightarrow \ell_0} - \partial h_{\ell_0-1 \leftrightarrow \ell_0+1}$ and use (A.28) to calculate

$$\begin{aligned} & \frac{\langle \phi_{\gamma_0, L_{1:\ell-1}} | H_{1:\ell-1} | \phi_{\gamma_0, L_{1:\ell-1}} \rangle}{|L_{1:\ell-1}|} \\ &= -\frac{\ell_0 R}{|L_{1:\ell-1}|} e_1 + \frac{|L_{1:\ell-1}| - \ell_0 R}{|L_{1:\ell-1}|} U + \frac{\langle \phi_{\gamma_0, L_{1:\ell-1}} | \partial h_{\ell_0-1 \leftrightarrow \ell_0} - \partial h_{\ell_0-1 \leftrightarrow \ell_0+1} | \phi_{\gamma_0, L_{1:\ell-1}} \rangle}{|L_{1:\ell-1}|} \\ &= -\gamma_0 e_1 + (1 - \gamma_0) U + \mathcal{O}(\ell^{-1}) + \frac{\langle \phi_{\gamma_0, L_{1:\ell-1}} | \partial h_{\ell_0-1 \leftrightarrow \ell_0+1} | \phi_{\gamma_0, L_{1:\ell-1}} \rangle}{|L_{1:\ell-1}|}. \end{aligned} \quad (\text{A.29})$$

In the last step, we used $\frac{\ell_0}{\ell} = \gamma_0 + \mathcal{O}(\ell^{-1})$ and the fact that $\langle \phi_{\gamma_0, L_{1:\ell-1}} | \partial h_{\ell_0-1 \leftrightarrow \ell_0} | \phi_{\gamma_0, L_{1:\ell-1}} \rangle = 0$. By Cauchy-Schwarz,

$$\begin{aligned} \langle \phi_{\gamma_0, L_{1:\ell-1}} | \partial h_{\ell_0-1 \leftrightarrow \ell_0+1} | \phi_{\gamma_0, L_{1:\ell-1}} \rangle &\leq C J \langle \phi_{\gamma_0, L_{1:\ell-1}} | (n_{L_{\ell_0-1}} + n_{L_1}) | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \\ &= \frac{2CJ}{\ell_0 - 1} \langle \xi_{1:\ell_0-1} | n_{L_{1:\ell_0-1}} | \xi_{1:\ell_0-1} \rangle \\ &= 2CJR \end{aligned} \quad (\text{A.30})$$

for an explicit constant $C > 0$ that may change from line to line that does not depend on any parameters, where we use $\langle \xi_{1:\ell_0-1} | n_{L_{1:\ell_0-1}} | \xi_{1:\ell_0-1} \rangle = R |L_{1:\ell_0-1}|$. The first equality in (A.30) holds by the periodicity of $|\xi_{1:\ell_0-1}\rangle$ in the horizontal direction. Returning to (A.29) and using that $\frac{R}{|L_{1:\ell-1}|} = \mathcal{O}(\ell^{-1})$, this proves that property (b) holds with $e_2 := U$.

We come to property (c). Using the eigenvalue equations (A.28), the only non-trivial contributions again come from the hopping term along the boundaries:

$$\begin{aligned}
& \langle \phi_{\gamma_0, L_{1:\ell-1}} | H_{L_{1:\ell-1}}^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \\
&= \langle \phi_{\gamma_0, L_{1:\ell-1}} | (H_{1:\ell_0-1}^{\text{per}} + H_{\ell_0:\ell-1} + \partial h_{\ell_0-1 \leftrightarrow \ell_0} - \partial h_{\ell_0-1 \leftrightarrow 1})^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \\
&= \langle \phi_{\gamma_0, L_{1:\ell-1}} | (H_{1:\ell_0-1}^{\text{per}} + H_{\ell_0:\ell-1})^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \\
&\quad + \langle \phi_{\gamma_0, L_{1:\ell-1}} | \{ H_{1:\ell_0-1}^{\text{per}} + H_{\ell_0:\ell-1}, \partial h_{\ell_0-1 \leftrightarrow \ell_0} - \partial h_{\ell_0-1 \leftrightarrow 1} \} | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \\
&\quad + \langle \phi_{\gamma_0, L_{1:\ell-1}} | (\partial h_{\ell_0-1 \leftrightarrow \ell_0} - \partial h_{\ell_0-1 \leftrightarrow 1})^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \\
&= (-\gamma_0 e_1 + (1 - \gamma_0) e_2 + \mathcal{O}(\ell^{-1}))^2 |L_{0:\ell-1}|^2 \\
&\quad + 2(-\gamma_0 e_1 + (1 - \gamma_0) e_2 + \mathcal{O}(\ell^{-1})) |L_{0:\ell-1}| \langle \phi_{\gamma_0, L_{1:\ell-1}} | (\partial h_{\ell_0-1 \leftrightarrow \ell_0} - \partial h_{\ell_0-1 \leftrightarrow 1}) | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \\
&\quad + \langle \phi_{\gamma_0, L_{1:\ell-1}} | (\partial h_{\ell_0-1 \leftrightarrow \ell_0} - \partial h_{\ell_0-1 \leftrightarrow 1})^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle,
\end{aligned}$$

where we use Eq. (A.28) in the last equation. The boundary terms can be controlled by an extension of the argument in (A.30). For the first-order boundary terms, this is immediate,

$$\begin{aligned}
| \langle \phi_{\gamma_0, L_{1:\ell-1}} | (\partial h_{\ell_0-1 \leftrightarrow \ell_0} - \partial h_{\ell_0-1 \leftrightarrow 1}) | \phi_{\gamma_0, L_{1:\ell-1}} \rangle | &\leq CJ | \langle \phi_{\gamma_0, L_{1:\ell-1}} | (n_{L_1} + n_{L_{\ell_0-1}}) | \phi_{\gamma_0, L_{1:\ell-1}} \rangle | \\
&\leq 2CJR.
\end{aligned}$$

Together with the factor of $|L_{0:\ell-1}|$ this gives a subleading term $\mathcal{O}(\ell^{-1})$ compared to the leading $|L_{0:\ell-1}|^2$ term.

For the squared boundary terms $\langle \phi_{\gamma_0, L_{1:\ell-1}} | (\partial h_{\ell_0-1 \leftrightarrow \ell_0} - \partial h_{\ell_0-1 \leftrightarrow 1})^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle$, we need to use an energy bound. First, multiple applications of Cauchy-Schwarz give

$$\langle \phi_{\gamma_0, L_{1:\ell-1}} | (\partial h_{\ell_0-1 \leftrightarrow \ell_0} - \partial h_{\ell_0-1 \leftrightarrow 1})^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \leq 32CRJ^2 \sum_{x \in L_1 \cup L_{\ell_0-1}} \langle \phi_{\gamma_0, L_{1:\ell-1}} | n_x^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle,$$

with $C > 0$ a universal constant. For instance, denoting $\tilde{n}_x = n_x + 1$,

$$\begin{aligned}
& \langle \phi_{\gamma_0, L_{1:\ell-1}} | \partial h_{\ell_0-1 \leftrightarrow 1}^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \\
&= J^2 \sum_{1 \leq k, k' \leq R} \langle \phi_{\gamma_0, L_{1:\ell-1}} | (b_{1,k}^\dagger b_{\ell_0-1,k} + b_{1,k} b_{\ell_0-1,k}^\dagger) (b_{1,k'}^\dagger b_{\ell_0-1,k'} + b_{1,k'} b_{\ell_0-1,k'}^\dagger) | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \\
&\leq CJ^2 \sum_{1 \leq k < k' \leq R} \langle \phi_{\gamma_0, L_{1:\ell-1}} | (\tilde{n}_{1,k} \tilde{n}_{1,k'} + \tilde{n}_{1,k} \tilde{n}_{\ell_0-1,k'} + \tilde{n}_{\ell_0-1,k} \tilde{n}_{1,k'} + \tilde{n}_{\ell_0-1,k} \tilde{n}_{\ell_0-1,k'}) | \phi_{\gamma_0, L_{1:\ell-1}} \rangle \\
&\leq 2CJR^2 \sum_{x \in L_1 \cup L_{\ell_0-1}} \langle \phi_{\gamma_0, L_{1:\ell-1}} | (n_x + 1)^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle. \\
&\leq 8CJR^2 \sum_{x \in L_1 \cup L_{\ell_0-1}} \langle \phi_{\gamma_0, L_{1:\ell-1}} | n_x^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle.
\end{aligned}$$

The other boundary terms are estimated analogously. To control $R \sum_{x \in L_j} \langle \phi_{\gamma_0, L_{1:\ell-1}} | n_x^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle$, we first recall periodicity of $|\xi_{1:\ell_0-1}\rangle$ in the horizontal direction,

$$R \sum_{x \in L_j} \langle \phi_{\gamma_0, L_{1:\ell-1}} | n_x^2 | \phi_{\gamma_0, L_{1:\ell-1}} \rangle = \frac{R}{\ell_0} \langle \xi_{1:\ell_0-1} | \sum_{x \in L_{1:\ell_0-1}} n_x^2 | \xi_{1:\ell_0-1} \rangle.$$

We control $\sum_{x \in L_{0:\ell_0-1}} n_x^2$ by the energy of $|\xi_{1:\ell_0-1}\rangle$ up to lower order terms that are linear in particle number. Indeed, by Cauchy-Schwarz, we have the operator inequality

$$\sum_{x \in L_{1:\ell_0-1}} n_x^2 \leq \frac{1}{U} \left(H_{1:\ell_0-1}^{\text{per}} + c_1(J, U) n_{L_{1:\ell_0-1}} + c_2(J, U) |L_{1:\ell_0-1}| \right)$$

for suitable constants $c_1(J, U), c_2(J, U) > 0$. This implies

$$\frac{R}{\ell_0} \langle \xi_{1:\ell_0-1} | \sum_{x \in L_{1:\ell_0-1}} n_x^2 | \xi_{1:\ell_0-1} \rangle \leq \frac{R|L_{1:\ell_0-1}|}{\ell_0 U} (c_1(J, U) + c_2(J, U) - e_1) =: c(J, U) R^2.$$

Since $R^2 = |L_{1:\ell_0-1}|^2 \mathcal{O}(\ell_0^{-2}) = |L_{1:\ell_0-1}|^2 \mathcal{O}(\ell^{-2})$ from $\ell_0 = \lfloor \gamma_0 \ell \rfloor \propto \ell$, this establishes property (c) and completes the proof of Lemma A.2. \square

References

- [APSS21] Jack Arbunich, Fabio Pusateri, Israel Michael Sigal, and Avraham Soffer. Maximal speed of quantum propagation. *Letters in Mathematical Physics*, 111:1–16, 2021.
- [BDN12] Immanuel Bloch, Jean Dalibard, and Sylvain Nascimbene. Quantum simulations with ultracold quantum gases. *Nature Physics*, 8(4):267–276, 2012.
- [BDZ08] Immanuel Bloch, Jean Dalibard, and Wilhelm Zwerger. Many-body physics with ultracold gases. *Reviews of Modern Physics*, 80(3):885, 2008.
- [BFL⁺23] Sébastien Breteaux, Jérémy Faupin, Marius Lemm, Dong Hao Ou Yang, Israel Michael Sigal, and Jingxuan Zhang. Light cones for open quantum systems. *arXiv preprint arXiv:2303.08921*, 2023.
- [BFLS22] Sébastien Breteaux, Jérémy Faupin, Marius Lemm, and Israel Michael Sigal. Maximal speed of propagation in open quantum systems. In Rupert L. Frank et al., editor, *The Physics and Mathematics of Elliott Lieb, The 90th Anniversary*, volume I. EMS Press, Berlin, 2022.
- [BHV06] Sergey Bravyi, Matthew B Hastings, and Frank Verstraete. Lieb-Robinson bounds and the generation of correlations and topological quantum order. *Physical Review Letters*, 97(5):050401, 2006.
- [BMNS12] Sven Bachmann, Spyridon Michalakis, Bruno Nachtergaele, and Robert Sims. Automorphic equivalence within gapped phases of quantum lattice systems. *Communications in Mathematical Physics*, 309(3):835–871, 2012.
- [BPCK12] Peter Barmettler, Dario Poletti, Marc Cheneau, and Corinna Kollath. Propagation front of correlations in an interacting Bose gas. *Physical Review A*, 85(5):053625, 2012.
- [CBP⁺12] Marc Cheneau, Peter Barmettler, Dario Poletti, Manuel Endres, Peter Schauß, Takeshi Fukuhara, Christian Gross, Immanuel Bloch, Corinna Kollath, and Stefan Kuhr. Light-cone-like spreading of correlations in a quantum many-body system. *Nature*, 481(7382):484–487, 2012.
- [Che22] Marc Cheneau. Experimental tests of Lieb-Robinson bounds. In Rupert L. Frank et al., editor, *The Physics and Mathematics of Elliott Lieb, The 90th Anniversary*, volume I. EMS Press, Berlin, 2022.
- [CL19] Chi-Fang Chen and Andrew Lucas. Finite speed of quantum scrambling with long range interactions. *Physical Review Letters*, 123(25):250605, 2019.
- [CLY23] Chi-Fang Chen, Andrew Lucas, and Chao Yin. Speed limits and locality in many-body quantum dynamics. *Reports on Progress in Physics*, 86:116001, 2023.

- [CSE08] Marcus Cramer, Alessio Serafini, and Jens Eisert. Locality of dynamics in general harmonic quantum systems. *CRM Series, Scuola Normale Superiore*, 8:51–72, 2008.
- [DSH⁺02] Johannes Hecker Denschlag, Jesse E Simsarian, Hartmut Häffner, C McKenzie, Antoine Browaeys, Donghyun Cho, Kristian Helmerson, Steve L Rolston, and William D Phillips. A Bose-Einstein condensate in an optical lattice. *Journal of Physics B: Atomic, Molecular and Optical Physics*, 35(14):3095, 2002.
- [EG09] Jens Eisert and David Gross. Supersonic quantum communication. *Physical Review Letters*, 102(24):240501, 2009.
- [EW17] Jeffrey M Epstein and K Birgitta Whaley. Quantum speed limits for quantum-information-processing tasks. *Physical Review A*, 95(4):042314, 2017.
- [FLS22a] Jérémy Faupin, Marius Lemm, and Israel Michael Sigal. Maximal speed for macroscopic particle transport in the Bose-Hubbard model. *Physical Review Letters*, 128(15):150602, 2022.
- [FLS22b] Jérémy Faupin, Marius Lemm, and Israel Michael Sigal. On Lieb–Robinson Bounds for the Bose–Hubbard Model. *Communications in Mathematical Physics*, 394(3):1011–1037, 2022.
- [GB17] Christian Gross and Immanuel Bloch. Quantum simulations with ultracold atoms in optical lattices. *Science*, 357(6355):995–1001, 2017.
- [GE16] Christian Gogolin and Jens Eisert. Equilibration, thermalisation, and the emergence of statistical mechanics in closed quantum systems. *Reports on Progress in Physics*, 79(5):056001, 2016.
- [GF08] Markus Greiner and Simon Fölling. Optical lattices. *Nature*, 453(7196):736–738, 2008.
- [GNRS20] Martin Gebert, Bruno Nachtergaele, Jake Reschke, and Robert Sims. Lieb–Robinson Bounds and Strongly Continuous Dynamics for a Class of Many-Body Fermion Systems in \mathbb{R}^d . *Annales Henri Poincaré*, 21(11):3609–3637, 2020.
- [Has04] Matthew B Hastings. Decay of correlations in fermi systems at nonzero temperature. *Physical Review Letters*, 93(12):126402, 2004.
- [Has07a] Matthew B Hastings. An area law for one-dimensional quantum systems. *Journal of Statistical Mechanics: Theory and Experiment*, 2007(08):P08024, 2007.
- [Has07b] Matthew B Hastings. Quantum belief propagation: An algorithm for thermal quantum systems. *Physical Review B*, 76(20):201102, 2007.
- [HLS23] Benjamin Hinrichs, Marius Lemm, and Oliver Siebert. On Lieb-Robinson bounds for a class of continuum fermions. *arXiv preprint arXiv:2310.17736*, 2023.
- [HW05] Matthew B Hastings and Xiao-Gang Wen. Quasiadiabatic continuation of quantum states: The stability of topological ground-state degeneracy and emergent gauge invariance. *Physical Review B*, 72(4):045141, 2005.
- [Jak04] Dieter Jaksch. Optical lattices, ultracold atoms and quantum information processing. *Contemporary Physics*, 45(5):367–381, 2004.
- [KGE14] Martin Kliesch, Christian Gogolin, and Jens Eisert. Lieb-Robinson bounds and the simulation of time-evolution of local observables in lattice systems. *Many-Electron Approaches in Physics, Chemistry and Mathematics: A Multidisciplinary View*, pages 301–318, 2014.
- [KGK⁺14] Martin Kliesch, Christian Gogolin, MJ Kastoryano, Arnau Riera, and Jens Eisert. Locality of temperature. *Physical Review X*, 4(3):031019, 2014.
- [KN21] Adam M Kaufman and Kang-Kuen Ni. Quantum science with optical tweezer arrays of ultracold atoms and molecules. *Nature Physics*, 17(12):1324–1333, 2021.

- [KS21a] Tomotaka Kuwahara and Keiji Saito. Absence of fast scrambling in thermodynamically stable long-range interacting systems. *Physical review letters*, 126(3):030604, 2021.
- [KS21b] Tomotaka Kuwahara and Keiji Saito. Lieb-robinson bound and almost-linear light cone in interacting boson systems. *Physical Review Letters*, 127(7):070403, 2021.
- [Kuw16] Tomotaka Kuwahara. Exponential bound on information spreading induced by quantum many-body dynamics with long-range interactions. *New Journal of Physics*, 18(5):053034, 2016.
- [KVS24] Tomotaka Kuwahara, Tan Van Vu, and Keiji Saito. Effective light cone and digital quantum simulation of interacting bosons. *Nature Communications*, 15(1):2520, Mar 2024.
- [LHS22] Ethan Lake, Michael Hermele, and T Senthil. Dipolar bose-hubbard model. *Physical Review B*, 106(6):064511, 2022.
- [LR72] Elliott H Lieb and Derek W Robinson. The finite group velocity of quantum spin systems. *Communications in Mathematical Physics*, 28(3):251–257, 1972.
- [LRS⁺23] Marius Lemm, Carla Rubiliani, Israel Michael Sigal, Jingxuan Zhang, et al. Information propagation in long-range quantum many-body systems. *Physical Review A*, 108(6):L060401, 2023.
- [LRZ23] Marius Lemm, Carla Rubiliani, and Jingxuan Zhang. Microscopic propagation bounds for long-range bosons. *arXiv preprint arXiv:2310.14896*, 2023.
- [LSA12] Maciej Lewenstein, Anna Sanpera, and Veronica Ahufinger. *Ultracold Atoms in Optical Lattices: Simulating quantum many-body systems*. OUP Oxford, 2012.
- [LSH⁺13] Nima Lashkari, Douglas Stanford, Matthew Hastings, Tobias Osborne, and Patrick Hayden. Towards the fast scrambling conjecture. *Journal of High Energy Physics*, 2013(4):1–33, 2013.
- [MKM20] Suman Mondal, Augustine Kshetrimayum, and Tapan Mishra. Two-body repulsive bound pairs in a multibody interacting bose-hubbard model. *Physical Review A*, 102(2):023312, 2020.
- [NOS06] Bruno Nachtergaele, Yoshiko Ogata, and Robert Sims. Propagation of correlations in quantum lattice systems. *Journal of Statistical Physics*, 124:1–13, 2006.
- [NRSS09] Bruno Nachtergaele, Hillel Raz, Benjamin Schlein, and Robert Sims. Lieb-robinson bounds for harmonic and anharmonic lattice systems. *Communications in Mathematical Physics*, 286:1073–1098, 2009.
- [NS06] Bruno Nachtergaele and Robert Sims. Lieb-Robinson bounds and the exponential clustering theorem. *Communications in Mathematical Physics*, 265:119–130, 2006.
- [NS07] Bruno Nachtergaele and Robert Sims. A multi-dimensional Lieb-Schultz-Mattis theorem. *Communications in Mathematical Physics*, 276:437–472, 2007.
- [NS10] Bruno Nachtergaele and Robert Sims. Lieb-Robinson bounds in quantum many-body physics. *Contemporary Mathematics*, 529:141–176, 2010.
- [NSY18] Bruno Nachtergaele, Robert Sims, and Amanda Young. Lieb-Robinson bounds, the spectral flow, and stability of the spectral gap for lattice fermion systems. *Mathematical Problems in Quantum Physics*, 717:93–115, 2018.
- [RS16] Daniel A Roberts and Brian Swingle. Lieb-Robinson bound and the butterfly effect in quantum field theories. *Physical Review Letters*, 117(9):091602, 2016.
- [SHOE11] Norbert Schuch, Sarah K Harrison, Tobias J Osborne, and Jens Eisert. Information propagation for interacting-particle systems. *Physical Review A*, 84(3):032309, 2011.
- [SSH⁺23] Guo-Xian Su, Hui Sun, Ana Hudomal, Jean-Yves Desaulles, Zhao-Yu Zhou, Bing Yang, Jad C Halimeh, Zhen-Sheng Yuan, Zlatko Papić, and Jian-Wei Pan. Observation of many-body scarring in a Bose-Hubbard quantum simulator. *Physical Review Research*, 5(2):023010, 2023.

- [SZ22] Israel Michael Sigal and Jingxuan Zhang. On propagation of information in quantum many-body systems. *arXiv preprint arXiv:2212.14472*, 2022.
- [TCE⁺20] Minh C Tran, Chi-Fang Chen, Adam Ehrenberg, Andrew Y Guo, Abhinav Deshpande, Yifan Hong, Zhe-Xuan Gong, Alexey V Gorshkov, and Andrew Lucas. Hierarchy of linear light cones with long-range interactions. *Physical Review X*, 10(3):031009, 2020.
- [Tes07] Romain Tessera. Volume of spheres in doubling metric measured spaces and in groups of polynomial growth. *Bulletin de la Société mathématique de France*, 135(1):47–64, 2007.
- [VKS23] Tan Van Vu, Tomotaka Kuwahara, and Keiji Saito. Optimal form of light cones for bosonic transport in long-range systems. *arXiv preprint arXiv:2307.01059*, 2023.
- [WCP15] Mischa P Woods, M Cramer, and Martin B Plenio. Simulating bosonic baths with error bars. *Physical Review Letters*, 115(13):130401, 2015.
- [WH20] Zhiyuan Wang and Kaden RA Hazzard. Tightening the Lieb-Robinson bound in locally interacting systems. *PRX Quantum*, 1(1):010303, 2020.
- [WP16] Mischa P Woods and Martin B Plenio. Dynamical error bounds for continuum discretisation via Gauss quadrature rules—A Lieb-Robinson bound approach. *Journal of Mathematical Physics*, 57(2), 2016.
- [YL22] Chao Yin and Andrew Lucas. Finite speed of quantum information in models of interacting bosons at finite density. *Physical Review X*, 12(2):021039, 2022.
- [YSO⁺20] Bing Yang, Hui Sun, Robert Ott, Han-Yi Wang, Torsten V Zache, Jad C Halimeh, Zhen-Sheng Yuan, Philipp Hauke, and Jian-Wei Pan. Observation of gauge invariance in a 71-site Bose–Hubbard quantum simulator. *Nature*, 587(7834):392–396, 2020.