

Higher Berry Phase from Projected Entangled Pair States in (2+1) dimensions

Shuhei Ohyama^{1,*} and Shinsei Ryu^{2,†}

¹*RIKEN Center for Emergent Matter Science, Wako, Saitama, 351-0198, Japan*

²*Department of Physics, Princeton University, Princeton, New Jersey 08544, USA*

(Dated: May 10, 2024)

We consider families of invertible many-body quantum states in d spatial dimensions that are parameterized over some parameter space X . The space of such families is expected to have topologically distinct sectors classified by the cohomology group $H^{d+2}(X; \mathbb{Z})$. These topological sectors are distinguished by a topological invariant built from a generalization of the Berry phase, called the higher Berry phase. In the previous work [1], we introduced a generalized inner product for three one-dimensional many-body quantum states, (“triple inner product”). The higher Berry phase for one-dimensional invertible states can be introduced through the triple inner product and furthermore the topological invariant, which takes its value in $H^3(X; \mathbb{Z})$, can be extracted. In this paper, we introduce an inner product of four two-dimensional invertible quantum many-body states. We use it to measure the topological nontriviality of parameterized families of 2d invertible states. In particular, we define a topological invariant of such families that takes values in $H^4(X; \mathbb{Z})$. Our formalism uses projected entangled pair states (PEPS). We also construct a specific example of non-trivial parameterized families of 2d invertible states parameterized over $\mathbb{R}P^4$ and demonstrate the use of our formula. Applications for symmetry-protected topological phases are also discussed.

CONTENTS

I. Introduction	2
A. Physical Intuition	3
II. The Higher Berry Phase for (2+1)d Systems	5
A. Preliminaries	5
1. MPO	6
2. Semi-injective PEPS	6
3. Key Propositions and the Unitarity Assumption	7
4. The Definition of the Quadruple Inner Product	8
B. The Construction of the Higher Berry Phase	13
1. Reduction	13
2. Fixed Points of the 2- and 3-Transfer Matrices	14
3. Fixed Points of the 4-Transfer Matrix	16
4. Proof of Props. 3 and 4	19
III. The CZX Model Protected by $\mathbb{Z}/2\mathbb{Z}$ Symmetry	21
IV. The Model Parameterized Over $\mathbb{R}P^4$	23
A. The $\mathbb{R}P^4$ -Parametrized Model	23
B. Transition MPU	24
C. Integration of the Higher Berry Phase	24
V. Geometric Interpretation: 2-Gerbe	28
VI. Summary and Discussion	29

* shuhei.ohyama@riken.jp

† shinseir@princeton.edu

Acknowledgments	29
References	30

I. INTRODUCTION

Invertible states are states realized as the unique ground states of gapped Hamiltonians. A notable subclass of invertible states is symmetry-protected topological (SPT) phases. In the absence of symmetry conditions, these states are trivial and are continuously deformable to, e.g., a product state. On the other hand, when subjected to a set of symmetry conditions, they acquire a distinct topological nature and are separated from trivial states by a quantum phase transition [2–10].

More generally, one can consider a family of invertible states that depend continuously on some parameter(s) x , $\{|\psi(x)\rangle : \text{invertible state } |x \in X\}$, where X represents the parameter space. We refer to such a family as invertible states over X or invertible states parameterized by X . In simple cases, a parameterized family of invertible states can be viewed as an adiabatic process, that can be topologically non-trivial. A classic example is the Thouless pump [11], which is a topologically non-trivial adiabatic process that transports quantized charges.

Considering families of invertible states is also useful in developing a systematic classification of invertible states. According to Kitaev’s conjecture, the classification of such states is provided by a generalized cohomology theory [12]. From general and mathematical considerations using relativistic field theories, the classification of interacting invertible states is expected to be given by a generalized cohomology known as the Anderson dual of bordism theory. Partial proofs supporting this idea have been provided [13]. However, it is worth noting that the classification of invertible states realized as lattice systems does not necessarily have to coincide with the classification expected from quantum field theory. A recent work [14] has pointed out the existence of classes that deviate from the classification by the Anderson dual of bordism theory in $2 + 1$ -dimensional fermion systems.

This paper focuses on topologically non-trivial families of invertible states in dimensions higher than two. In particular, we aim to construct a systematic method for computing the topological invariants of such families. The space of families of d -dimensional invertible states parameterized by X has topologically distinct sectors classified by $H^{d+2}(X; \mathbb{Z})$. This class is distinguished by the higher Berry phase, or more precisely the topological invariants built from the higher Berry phase, which has been actively studied recently as a generalization of the regular Berry phase – see, e.g., [15–27]. However, the explicit construction of the higher Berry phase, in terms of many-body wave functions in particular, is not known completely. In our previous paper [1], we introduced a generalized inner product (or multi-wave function overlap) of three 1d many-body states to measure the nontriviality and construct the invariant which takes its value in $H^3(X; \mathbb{Z})$. In this paper, we introduce an inner product of four 2d states and try to measure the nontriviality in $H^4(X; \mathbb{Z})$.

We do so by using tensor networks, more specifically, projected entangled pair states (PEPS). Tensor networks are a versatile framework to study various problems in many-body quantum physics [10]. Recent works also demonstrate that tensor networks can capture higher geometric structures, such as the higher Berry phase, encoded in many-body quantum wavefunction [1, 22, 28–31]. In our previous work [1], we used the matrix product state (MPS) representation of (1+1)-dimensional invertible states and introduced the triple inner product. The triple inner product assigns a complex number for three invertible states (MPS) in (1+1) dimensions and allows us to extract the higher Berry phase. Furthermore, as a mathematical structure that underlies the higher Berry phase, we constructed and identified a gerbe on X for a parameterized family of (1+1)-dimensional invertible states. Here, a gerbe can be thought of as a generalization of a complex line bundle, which underlies the description of the regular Berry phase. See Ref. [30] that also associates a gerbe to a parameterized family of MPS over X . The subsequent work [29] provides a formulation for the higher Berry curvature using MPS, and also demonstrated numerical evaluations of the Berry curvature using the density matrix renormalization group (DMRG).

Specifically, following Ref. [1], we introduce a generalized inner product for four PEPS, which we refer to as “quadruple inner product”, and demonstrate that we can extract the higher Berry phase. Additionally, we will argue (with some caveats) that the underlying mathematical structure is a 2-gerbe – similarly to how gerbes are higher generalizations of complex line bundles, 2-gerbes are generalizations of gerbes. While our




FIG. 1. (a) Taking the triple inner product of three matrix product states $\{A_\alpha^i\}$, $\{A_\beta^i\}$ and $\{A_\gamma^i\}$. (b) The triple inner product in which the virtual legs at spatial infinities are contracted with the left and right eigenvectors (fixed points) $\Lambda_{\alpha\beta}^L$, $\Lambda_{\alpha\gamma}^R$ and $\Lambda_{\gamma\beta}^R$ of the mixed transfer matrices.

primary focus will be on (2+1)-dimensional PEPS, we anticipate that generalizations to higher dimensions should be possible.

The paper is organized as follows: the remaining part of this section (Sec. IA) provides an overview of the construction of the triple inner product of MPS and outline its higher-dimensional generalization, serving as an instructive guide for the subsequent detailed construction. Section II delves into a specific class of PEPS, the semi-injective PEPS, to implement the ideas sketched in Sec. IA. Our formula for the quadruple inner product and the higher Berry phase, written in terms of various fixed point tensors, is summarized in Eq. (22) in Sec. II A. Also, at the end of Sec. II A, we outline two applications; the topological invariants of SPT phases and the higher Berry phase associated with parameterized invertible states. Specifically, when applied to (2+1)d SPT phases protected by symmetry G , we can use the quadruple inner product to extract the topological invariants of SPT phases valued in $H^3(G, U(1))$. In the SPT context, our formulation is closely related to [32] and other prior works [33–39]. In the subsequent section Sec. IIB, we provide the rationale behind our formula and prove various claims that are necessary to establish the formula. In Sec. III, we discuss concrete models that realize non-trivial SPT phases. For the example, we show that the higher Berry phase is given by the group cohomology phase (3-cocycle) of the SPT phases. In Sec. IV, we present a concrete model parametrized over $\mathbb{R}P^4$ and show, using the quadruple inner product, that it is characterized by a non-trivial topological invariant. Finally in Sec. V, we identify a proper mathematical structure that describes parameterized (2+1)d invertible states and the higher Berry phase. We conclude in Sec. VI.

A. Physical Intuition

Previous works studied the higher Berry phase for (1+1)d many-body states using MPS. In this paper, we consider (2+1)d many-body states and their higher Berry phase using PEPS. The motivation and intuition behind our construction of the quadruple inner product and the higher Berry phase can be best described by reviewing their (1+1)d counterparts.

To set the stage, we consider a family of MPS with MPS matrices $\{A^i(x)\}_{i=1,\dots,D}$ parameterized by $x \in X$, where X is the parameter space. (Here, $i = 1, \dots, D$ represents physical degrees of freedom and indices for the auxiliary (internal) are implicit here – see Ref. [1] for our notations.) We assume that our MPS tensors are normal, $\sum_i A^i A^{i\dagger} = 1$, and work with the right canonical gauge. As shown in Ref. [1], the higher Berry phase for the family, and the topological invariant (the Dixmier-Douady class) can be extracted from a certain overlap of three MPS as follows. We consider an open covering of X , $\{U_\alpha\}$. On each patch, we have MPS, $\{A_\alpha^i(x)\}$. When two patches overlap, two MPS $\{A_\alpha^i(x)\}$ and $\{A_\beta^i(x)\}$ represent the same physical state. The fundamental theorem of injective MPS then asserts that these two MPS are related to each other by a gauge transformation, $A_\alpha^i(x) = g_{\alpha\beta}(x) A_\beta^i(x) g_{\alpha\beta}^\dagger(x)$. Now, at the triple intersection, $U_{\alpha\beta\gamma} := U_\alpha \cap U_\beta \cap U_\gamma$, we consider three MPS, $\{A_\alpha^i(x)\}$, $\{A_\beta^i(x)\}$ and $\{A_\gamma^i(x)\}$, representing the same physical state, and consider the triple inner product, defined schematically in Fig. 1(a). To make the definition precise, however, we need to address two issues. First, we need to specify how virtual bonds at the ends of the chain are contracted, i.e., there are remaining legs at the interface between the two states. Otherwise, Fig. 1 represents an operator rather than a number. Second, we must address a conceptual concern: as we are interested in intrinsic




FIG. 2. (a) A PEPS tensor with one physical and four auxiliary bonds. (b) The PEPS representation of a $(2+1)d$ invertible state. (c) A transition MPO is an eigen MPO for the mixed transfer matrix made of two PEPS A_α^i and A_β^i . Here, the two PEPS are contracted by their physical bonds (not shown explicitly).




FIG. 3. The quadruple inner product for four PEPSs (here represented as 2d "sheets") defined by gluing four PEPSs as illustrated above. (a) Four PEPSs from different patches at the quad intersection $U_{\alpha\beta\gamma\delta}$. The physical indices (not shown explicitly) are contracted such that the four PEPSs are glued with each other. (b) For a pair of PEPS, we can define an eigen MPO $\mathcal{O}_{\alpha\beta}$, that can be used to contract indices of PEPS as in (c). (d) The three-leg tensor can contract indices at the intersection where three MPO meet.

quantities to many-body quantum states, the higher generalization of the inner product and the Berry phase must make sense in the thermodynamic limit. These can be addressed simultaneously by contracting the virtual legs using the left or right fixed point of the mixed transfer matrix [Fig. 1(b)]. Using the eigenvalue equation, we can regard this diagram as the overlap of three infinite systems. This quantity is called the triple inner product, and using it, we can compute the higher Berry phase for $(1+1)d$ systems. Recalling that the regular Berry phase for $(0+1)d$ quantum systems can be computed from the inner product of two states, the above generalization for $(1+1)d$ systems can be considered extremely natural.

Following this line of thought, we expect that the higher Berry phase in general dimensions can be computed by considering d -fold inner products of $(d+1)d$ quantum states. Let us now turn to the $(2+1)d$ case and consider PEPS $\{A^i(x)\}$ parameterized over X . Throughout the paper, we consider PEPS tensors with one physical bond and four auxiliary bonds (Fig. 2). Paralleling the $(1+1)d$ case, it is natural to consider the overlap of four physically equivalent states in different gauges at an intersection $U_{\alpha\beta\gamma\delta} := U_\alpha \cap U_\beta \cap U_\gamma \cap U_\delta$ as in Fig. 3. There, as in the $(1+1)d$ case, the physical indices of the PEPS are contracted "partially" with others. I.e., the physical indices of $\{A_\alpha^i\}$ in the first quadrant are contracted with the physical indices of $\{A_\beta^i\}$ in the third quadrant, etc. As in the $(1+1)d$ case, there are remaining legs (bonds) at the boundaries of the PEPS networks. They need to be properly contracted such that the inner product is well-defined for the many-body states (PEPS) in the thermodynamic limit.

First, let us consider the dangling bonds at the interface where two PEPS networks meet. Much the same way with the $(1+1)d$ case, we consider the mixed transfer matrix at the intersection $U_\alpha \cap U_\beta = U_{\alpha\beta}$ as depicted in Fig. 2(c). Here, we assume the transfer matrix has a unique maximal eigenvalue 1 and the corresponding eigenvector is given as a matrix product operator (MPO). Moreover, we assume that MPO tensors do not depend on the number of sites in the interface direction ¹. This MPO is a higher-dimensional

¹ These are naive guesses intended to give an explanation of the idea for constructing invariants, and we should mention that they are too strong assumptions. When we actually construct the invariants in Sec. ??, we will relax the eigenvalue equations related to the transfer matrix and avoid these difficulties.




FIG. 4. (a) The fixed point condition for the three-leg tensor and (b) the higher Berry phase.

analog of the transition functions, denoted as $g_{\alpha\beta}(x)$, in (1+1)d case. We will call it a transition MPO and denote it as $\mathcal{O}_{\alpha\beta}$. By using the property of the fixed point, we can implement the thermodynamic limit in the bulk direction, i.e., the direction orthogonal to the interface.

In addition to the interface (the orange part) between the two PEPS networks, it is evident that there is "an interface of the interface" (the blue part in Fig. 3). At this point, the three virtual legs of the MPO, $\mathcal{O}_{\alpha\beta}$, $\mathcal{O}_{\beta\gamma}$ and $\mathcal{O}_{\alpha\gamma}$, say, meet with each other. We need an appropriate tensor so that we can contract the legs. We thus postulate the existence of a three-leg tensor $\Lambda_{\alpha\beta\gamma}$ for triple intersection $U_{\alpha\beta\gamma}$. Furthermore, similarly to the transition MPO $\mathcal{O}_{\alpha\beta}$ that satisfies the fixed point condition, it is natural to require the three-leg tensor to satisfy a fixed point condition such that we can take the thermodynamic limit in the directions along the double interfaces between two PEPS networks. Specifically, we require the "higher eigen equation" as in Fig. 4(a). We will adopt this as the defining condition of the three-leg tensor. (More generically, the eigenvalue can be a complex number of modulus one. The phase basically is the Dixmier-Douady class that appeared above when we considered three MPS. For now, we assume such "lower invariant" vanishes for simplicity.)

By using these tensors, we can contract all the remaining legs in Fig. 3. Furthermore, we can use the eigenequations (fixed point conditions) to annihilate physical sites, and then reduce the diagram Fig. 3(d) to the tetrahedral diagram made of four 3-leg tensors as Fig. 4(b). This completes the sketch of the construction of the higher Berry phase (and the topological invariant built from the higher Berry phase) from PEPS. Note however that in the above construction, we assume the existence of various tensors. It is necessary to clarify these points. In the next section, we will describe our construction using semi-injective PEPS [32] to concretely implement this idea.²

Finally, to see why these are natural objects to consider, and how these objects need to be combined, it is good to compare with the (1+1)d case. In (1+1)d, we assign an MPS $|\{A_\alpha^i\}\rangle$ for U_α , $|\{\Lambda_{\alpha\beta}\}\rangle$ for $U_{\alpha\beta}$, and a U(1) phase $c_{\alpha\beta\gamma}$ for $U_{\alpha\beta\gamma}$, respectively. Here, $|\{\Lambda_{\alpha\beta}\}\rangle$ or simply $\Lambda_{\alpha\beta}$ represents "mixed gauge" MPS. In mathematical terms, $|\{\Lambda_{\alpha\beta}\}\rangle$ is a line bundle (one-dimensional Hilbert space) assigned for the intersection $U_{\alpha\beta}$. The U(1) phase $c_{\alpha\beta\gamma}$ is the higher Berry phase. In (2+1)d, we need to consider a "higher" version. We assign a PEPS $|\{A_\alpha^i\}\rangle$ for U_α , an MPO $\mathcal{O}_{\alpha\beta}$ for $U_{\alpha\beta}$, a three-leg tensor $\Lambda_{\alpha\beta\gamma}$ for $U_{\alpha\beta\gamma}$, and a U(1) phase $c_{\alpha\beta\gamma\delta}$ for $U_{\alpha\beta\gamma\delta}$. Roughly, the three-leg tensor $\Lambda_{\alpha\beta\gamma}$ is an analog of $\Lambda_{\alpha\beta}$ in (2+1)d, and the U(1) phase $c_{\alpha\beta\gamma\delta}$ is the higher Berry phase we are looking for. Note that at the double intersection $U_{\alpha\beta}$ we assign an MPO, which can be thought of as an MPS over $U_{\alpha\beta}$, and if so, we are essentially putting a gerbe at the intersection $U_{\alpha\beta}$. This is a generalization of a gerbe, just like a gerbe is a generalization of a line bundle. We will discuss the mathematical structure behind the construction of our invariant in Sec. V.

II. THE HIGHER BERRY PHASE FOR (2+1)D SYSTEMS

A. Preliminaries

In this section, we will materialize the idea sketched above by using a class of PEPS, the so-called semi-injective PEPS [32]. We begin by reviewing necessary backgrounds and notations; MPO, semi-injective PEPS, and transfer matrices.

² In this section and in this figure, we denote the three-leg tensor simply by $\Lambda_{\alpha\beta\gamma}$. However, as we will see in the later sections, we need various types of three-leg tensors for the precise construction.




FIG. 5. The setup for semi-injective PEPS. (a) The unit cell; (b) the invertible operator O ; (c) the full-rank state $|\phi\rangle$; (d) the semi-injective PEPS $(|\phi\rangle, O)$.

1. MPO

Matrix product operators (MPO) are central objects in our discussions. They are defined by a set of matrices (tensors) $\{B^{ij}\}_{i,j=1}^D$. Here, $i, j = 1, \dots, D$ are “physical” indices representing, e.g., physical spin degrees of freedom. In addition, the tensors $\{B^{ij}\}_{i,j=1}^D$ carry “virtual” indices (not shown explicitly). The dimension of the virtual space (“bond dimension”) is denoted by χ . We use $\mathcal{O}_L[\{B^{ij}\}]$ to denote the MPO of length L that is generated by the MPO tensors $\{B^{ij}\}_{i,j=1}^D$. To be concrete, we consider periodic boundary conditions, although it is possible to discuss other boundary conditions. When there is no confusion, we shall call both $\{B^{ij}\}_{i,j=1}^D$ and $\mathcal{O}_L[\{B^{ij}\}]$ MPO. We recall that MPS (MPO) matrices are said to be normal when the corresponding transfer matrix has a unique largest eigenvalue. We also note that any injective matrices are proportional to normal matrices. Conversely, any normal matrices become injective after blocking.

2. Semi-injective PEPS

Semi-injective PEPS, introduced in Ref. [32], is a type of PEPS that can represent a wide class of (2+1)d invertible states including ground states of symmetry-protected topological (SPT) phases. To introduce semi-injective PEPS, let’s consider a square lattice, where each unit cell consists of four sites [Fig. 5(a)]. At each site within a unit cell, we assign a physical quantum mechanical degree of freedom (“spin”) with local Hilbert space. The dimensions of the local Hilbert space can be different from site to site. The total Hilbert space is constructed from the local Hilbert spaces, by arranging the local degrees of freedom in a translationally symmetric manner. With this setup, a semi-injective PEPS is defined by two data $(|\phi\rangle, O)$. Here, $|\phi\rangle$ is a state defined on each 4-site system as in Fig. 5(c) with full rank reduced density matrix. O is an invertible operator that acts on the four corners of four adjacent squares, as in Fig. 5(b). By using these data, a semi-injective PEPS generated by $(|\phi\rangle, O)$ is defined as in Fig. 5(d).

What is important for us is the fundamental theorem of semi-injective PEPS. Let us now consider two semi-injective PEPS, $(|\phi_\alpha\rangle, O_\alpha)$ and $(|\phi_\beta\rangle, O_\beta)$, that generate the same physical state. Then, the fundamental theorem of semi-injective PEPS states that they are related by an invertible MPO acting on their auxiliary indices [32]. We shall call the MPO a transition MPO. (Recall also Fig. 2 (c).) In the following, we denote the MPO matrices (tensors) that generate the transition MPO as $\{B_{\alpha\beta}^{ij}\}_{i,j=1}^D$. Also, $\mathcal{O}_L[\{B_{\alpha\beta}^{ij}\}]$ denotes the corresponding MPO of length L that is generated by the MPO tensors $\{B_{\alpha\beta}^{ij}\}_{i,j=1}^D$. Furthermore, without losing generality, the fundamental theorem asserts that we can take $\{B_{\alpha\beta}^{ij}\}_{i,j=1}^D$ as normal matrices (tensors).

3. Key Propositions and the Unitarity Assumption

In the construction of the quadruple inner product, various quantities are realized as fixed points of the transfer matrix. The following claim proves useful in this context³.

Proposition 1.

Let $\{B_1^{ij}\}_{i,j=1}^D, \dots, \{B_k^{ij}\}_{i,j=1}^D$ be normal MPO matrices and assume that the composition of all the MPO is proportional to the identity MPO. That is, there exists an invertible complex number $c \in \mathbb{C}^\times$ such that

$$\mathcal{O}_L[\{\sum_{j_2, \dots, j_k} B_1^{j_1 j_2} \otimes \dots \otimes B_k^{j_k j_{k+1}}\}] = c^L \mathcal{O}_L[\{\delta^{j_1 j_{k+1}}\}] \quad (1)$$

for all $L \in \mathbb{N}$. Then, the spectrum of the k -transfer matrix

$$T^{(k)}[B_1, \dots, B_k] := \frac{1}{c^D} \sum_{j_1, \dots, j_k} B_1^{j_1 j_2} \otimes \dots \otimes B_k^{j_k j_1}, \quad (2)$$

acting on $M \in \mathbb{C}^{\chi_1 \cdots \chi_k}$ as

$$\begin{array}{c} \text{---} \\ | \\ \text{---} B_1 \\ | \\ \text{---} B_2 \\ | \\ \vdots \\ | \\ \text{---} B_k \\ | \\ \text{---} \end{array} \cdot \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \\ | \\ \vdots \\ | \\ \text{---} \\ | \\ \text{---} \end{array} M := \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \\ | \\ \vdots \\ | \\ \text{---} \\ | \\ \text{---} \end{array}, \quad (3)$$

is $\{0, 1\}$ and the eigenvalue 1 is nondegenerate. In this diagram, black dots at the ends of vertical bonds mean that the physical indices at the top and the bottom are contracted together.

Proof. By taking the vertical trace of Eq. (1), we obtain $\text{Tr}[(T^{(k)})^L] = 1$. Here, Tr is the horizontal trace originating from PBC. Let $\{\xi_i\}$ be the spectrum of $T^{(k)}$. Then, the above equation turns out to be $\sum_i \xi_i^L = 1$ for any L . This equation implies that one of the ξ_i , say ξ_1 , is 1 and the others are 0. \square

In the following, we often use this proposition for $k = 2$ and 3.

As mentioned in Sec. II A 2, while a transition MPO is guaranteed to be normal (and hence invertible), it is not necessarily an MPU. In our construction of the higher Berry phase below, however, this invertibility, which holds when an MPO is an MPU, plays a crucial role. Therefore, we will further assume transition MPO are MPU.⁴ This can be rationalized by recalling the (1+1)d case. There, the transition functions $(g_{\alpha\beta})$ defined for physically equivalent normal MPS matrices are invertible but not necessarily unitary. However, it is possible to make the transition functions unitary by performing a partial gauge fixing known as the canonical form [40]. While the existence of such a canonical form for PEPS is not known at present, it is expected that a similar gauge fixing exists, allowing us to transform the transition MPO into an MPU. We also note that in the context of SPT phases, their classification does not change upon restricting ourselves to the case where the transition MPO is an MPU.⁵ For these reasons, we will assume henceforth that transition MPO are MPU.

For an MPO that is an MPU, the following proposition holds true:

³ We would like to thank Norbert Schuch for teaching us the proposition and its proof.

⁴ When an MPO is an MPU, it is guaranteed that its transfer matrix has an invertible fixed point.

⁵ In fact, when considering the classification of SPT phases, the invariants using MPO take values in $H^3(G; \mathbb{C}^\times)$, and when they have an MPU, they take values in $H^3(G; \mathbb{U}(1))$, but the two cohomologies are known to be mathematically isomorphic.

Proposition 2.

Let $\{B_1^{ij}\}_{i,j=1}^{\mathbb{D}}, \dots, \{B_k^{ij}\}_{i,j=1}^{\mathbb{D}}$ be normal MPU matrices and assume that the composition of $\{B_1^{ij}\}_{i,j=1}^{\mathbb{D}}, \dots, \{B_{k-1}^{ij}\}_{i,j=1}^{\mathbb{D}}$ is proportional to the MPO generated by $\{B_k^{ij}\}_{i,j=1}^{\mathbb{D}}$. That is, there exists an invertible complex number $c \in \mathbb{C}^\times$ such that


$$\mathcal{O}_L[\{\sum_{j_2, \dots, j_{k-1}} B_1^{j_1 j_2} \otimes \dots \otimes B_{k-1}^{j_{k-1} j_k}\}] = c^L \mathcal{O}_L[\{B_k^{j_1 j_k}\}] \quad (4)$$

for all $L \in \mathbb{N}$. Then,

(i) the spectrum of the $(k-1, 1)$ -transfer matrix

$$T^{(k-1,1)}[B_1, \dots, B_{k-1}; B_k] := \frac{1}{c^{\mathbb{D}}} \sum_{j_1, \dots, j_k} B_1^{j_1 j_2} \otimes \dots \otimes B_{k-1}^{j_{k-1} j_k} \otimes B_k^{j_1, j_k \dagger} \quad (5)$$

with respect to the right action on $M \in \mathbb{C}^{\chi_1 \cdots \chi_{k-1}} \otimes \mathbb{C}^{\chi_k^*}$ defined as




is $\{0, 1\}$ and the eigenvalue 1 is nondegenerate. Here, $B^{ij \dagger}$ represents the hermitian conjugation of $(B^{ij})_{ab}$ with respect to the virtual legs a, b , $(B^{ij \dagger})_{ab} = (B^{ij})_{ba}^*$.

(ii) the spectrum of the $(k-1, 1)$ -transfer matrix

$$T^{(k-1,1)}[B_1, \dots, B_{k-1}; B_k] := \frac{1}{c^{\mathbb{D}}} \sum_{j_1, \dots, j_k} B_k^{j_1, j_k \dagger} \otimes B_1^{j_1 j_2} \otimes \dots \otimes B_{k-1}^{j_{k-1} j_k} \quad (7)$$

with respect to the left action on $M \in \mathbb{C}^{\chi_1 \cdots \chi_{k-1}} \otimes \mathbb{C}^{\chi_k^*}$ defined as



is $\{0, 1\}$ and the eigenvalue 1 is nondegenerate.

Since $T^{(1,1)}[B_1; B_2]$ is nothing but the usual mixed transfer matrix [41] for the normal MPU matrices $\{B_1^{ij}\}$ and $\{B_2^{ij}\}$, the fixed point of $T^{(1,1)}[B_1; B_2]$ is invertible.

4. The Definition of the Quadruple Inner Product

With these preliminaries, our definition of the quadruple inner product of four quantum states can be summarized as follows in terms of the fixed points of 2- and 3-transfer matrices. Let $|\psi_\alpha\rangle, |\psi_\beta\rangle, |\psi_\gamma\rangle$ and $|\psi_\delta\rangle$ be physically identical states and assume that they admit the semi-injective PEPS representation. For each pair of these states, we obtain the set of transition MPO tensors $\{B_{\alpha\beta}^{ij}\}$. As we mentioned in Sec. II A 3,

we assume that the MPO $\{\mathcal{O}_L[\{B_{\alpha\beta}^{ij}\}]\}$ are MPU. By using this assumption, $\mathcal{O}_L[\{B_{\alpha\beta}^{ij}\}]^{-1} = \mathcal{O}_L[\{B_{\alpha\beta}^{ij}\}]^\dagger = \mathcal{O}_L[\{B_{\alpha\beta}^{ji^*}\}]$.

For each pair of these states, $|\psi_\alpha\rangle$ and $|\psi_\beta\rangle$, say, we have two transition MPU generated by $\{B_{\alpha\beta}^{ij}\}$ and $\{B_{\beta\alpha}^{ij}\}$. By the fundamental theorem for semi-injective PEPS, we can show that

$$\mathcal{O}_L[\{\sum_k B_{\alpha\beta}^{ik} \otimes B_{\beta\alpha}^{kj}\}] = (c_2)^L \mathcal{O}_L[\{\delta^{ij}\}], \quad (9)$$

for some $c_2 \in \mathbb{C}^\times$. This equation implies that

$$\mathcal{O}_L[\{B_{\beta\alpha}^{ij}\}] = (c_{\alpha\beta})^L \mathcal{O}_L[\{B_{\alpha\beta}^{ij}\}]^{-1} = (c_{\alpha\beta})^L \mathcal{O}_L[\{B_{\alpha\beta}^{ji^*}\}], \quad (10)$$

that is, $\{B_{\beta\alpha}^{ij}\}$ and $\{B_{\alpha\beta}^{ji^*}\}$ are gauge equivalent. Therefore, we can take the gauge in which the MPO matrices satisfy

$$B_{\beta\alpha}^{ij} = B_{\alpha\beta}^{ji^*}. \quad (11)$$

Since Eq. (9) holds, we can define the 2-transfer matrix

$$T_{\alpha\beta}^{(2)} := T^{(2)}[B_{\alpha\beta}, B_{\beta\alpha}], \quad (12)$$

and apply Prop. 1. Let $\Lambda_{\alpha\beta}^{R(2)}$ (resp. $\Lambda_{\alpha\beta}^{L(2)}$) denote the right (resp. left) fixed point of $T_{\alpha\beta}^{(2)}$.

For each triplet of these states, say $|\psi_\alpha\rangle$, $|\psi_\beta\rangle$ and $|\psi_\gamma\rangle$, we have three transition MPU generated by $\{B_{\alpha\beta}^{ij}\}$, $\{B_{\beta\gamma}^{ij}\}$ and $\{B_{\gamma\alpha}^{ij}\}$. By the fundamental theorem for semi-injective PEPS, we can show that

$$\mathcal{O}_L[\{\sum_k B_{\alpha\beta}^{ik} \otimes B_{\beta\gamma}^{kj}\}] = c_3^L \mathcal{O}_L[\{B_{\alpha\gamma}^{ij}\}], \quad (13)$$

for some $c_3 \in \mathbb{C}^\times$. Thus we can define the (2,1)-transfer matrix

$$T_{\alpha\beta\gamma}^{R(2,1)} := T^{(2,1)}[B_{\alpha\beta}, B_{\beta\gamma}; \Lambda_{\alpha\gamma}^R B_{\alpha\gamma} (\Lambda_{\alpha\gamma}^R)^{-1}], \quad (14)$$

and apply Prop. 2. Here, $\Lambda_{\alpha\gamma}^R$ is the unique right fixed point of the transfer matrix $T^{(1,1)}[B_{\alpha\gamma}; B_{\alpha\gamma}]$. Note that although the phase of $\Lambda_{\alpha\gamma}^R$ is indeterminate from the fix point equation, $T_{\alpha\beta\gamma}^{R(2,1)}$ does not depend on the choice of the phase factor. Let $\Lambda_{\alpha\beta\gamma}^{R(3)}$ denote the right fixed point of $T_{\alpha\beta\gamma}^{R(2,1)}$. Similarly, we can define the left (2,1)-transfer matrix

$$T_{\alpha\beta\gamma}^{L(2,1)} := T^{(2,1)}[B_{\alpha\beta}, B_{\beta\gamma}; (\Lambda_{\alpha\gamma}^L)^{-1} B_{\alpha\gamma} \Lambda_{\alpha\gamma}^L], \quad (15)$$

and apply the Prop. 2. Here, $\Lambda_{\alpha\gamma}^L$ is a unique left fixed point of the transfer matrix $T^{(1,1)}[B_{\alpha\gamma}; B_{\alpha\gamma}]$. Note that although the phase of $\Lambda_{\alpha\gamma}^L$ is indeterminate from the fix point equation, $T_{\alpha\beta\gamma}^{L(2,1)}$ does not depend on the choice of the phase factor. Let $\Lambda_{\alpha\beta\gamma}^{L(3)}$ denote the left fixed point of $T_{\alpha\beta\gamma}^{L(2,1)}$.

Now, we have the fixed point tensors $\{\Lambda_{\alpha\beta}^{R(2)}, \Lambda_{\alpha\beta}^{L(2)}\}_{\alpha\beta}$ for each pair, and $\{\Lambda_{\alpha\beta\gamma}^{R(3)}, \Lambda_{\alpha\beta\gamma}^{L(3)}\}_{\alpha\beta\gamma}$ for each triplet. Since these tensors are constructed as the fixed points of the transfer matrices, there is an ambiguity as to how the phases are assigned. In particular, the phases of $\Lambda_{\alpha\beta}^{L(2)}$ and $\Lambda_{\alpha\beta}^{R(2)}$ (as well as those of $\Lambda_{\alpha\beta\gamma}^{L(3)}$ and $\Lambda_{\alpha\beta\gamma}^{R(3)}$) can be determined independently. However, we note the following proposition: ⁶

⁶ In the following, we will denote the fixed point tensor as a white box.

Proposition 3.

We define the inner product of $\Lambda_{\alpha\beta}^{L(2)}$ and $\Lambda_{\alpha\beta}^{R(2)}$ as

$$(\Lambda_{\alpha\beta}^{L(2)}, \Lambda_{\alpha\beta}^{R(2)}) := \Lambda_{\alpha\beta}^{L(2)} \boxed{\phantom{\Lambda_{\alpha\beta}^{R(2)}}} \Lambda_{\alpha\beta}^{R(2)} \quad (16)$$

and the inner product of $\Lambda_{\alpha\beta\gamma}^{L(3)}$ and $\Lambda_{\alpha\beta\gamma}^{R(3)}$ as

$$(\Lambda_{\alpha\beta\gamma}^{L(3)}, \Lambda_{\alpha\beta\gamma}^{R(3)}) := \Lambda_{\alpha\beta\gamma}^{L(2,1)} \boxed{\phantom{\Lambda_{\alpha\beta\gamma}^{R(2,1)}}} \Lambda_{\alpha\beta\gamma}^{R(2,1)} . \quad (17)$$

Then, they are non-zero complex numbers.

Using Prop. 3, once we chose the phase of the right fixed points, we can determine the phase of the left fixed point such that the inner product is 1. We will show this proposition in Sec. II B

For the quartet of states $|\psi_\alpha\rangle, |\psi_\beta\rangle, |\psi_\gamma\rangle$ and $|\psi_\delta\rangle$, we can also show that

$$\mathcal{O}_L[\{\sum_{k,l,m} B_{\alpha\beta}^{ik} \otimes B_{\beta\gamma}^{kl} \otimes B_{\gamma\delta}^{lm} \otimes B_{\delta\alpha}^{mj}\}] = c_4^L \mathcal{O}_L[\{\delta^{ij}\}] \quad (18)$$

for some complex number $c_4 \in \mathbb{C}^\times$. Thus we can define a 4-transfer matrix

$$T_{\alpha\beta\gamma\delta}^{(4)} := T^{(4)}[B_{\alpha\beta}, B_{\beta\gamma}, B_{\gamma\delta}, B_{\delta\alpha}]. \quad (19)$$

Finally, by using these tensors, we define the quadruple inner product as follows:

Definition 1.

For physically identical four states $|\psi_\alpha\rangle$, $|\psi_\beta\rangle$, $|\psi_\gamma\rangle$ and $|\psi_\delta\rangle$, (i) we define $N_{\alpha\beta\gamma\delta}^1 \in \mathbb{N}$ as a minimal natural number such that

$$\left(\begin{array}{c} | \\ | \\ | \\ | \end{array} - \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \Big| \delta^{ij} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right)^{N_{\alpha\beta\gamma\delta}^1} = 0, \quad (20)$$

and define $N_{\alpha\beta\gamma\delta}^2 \in \mathbb{N}$ as a minimal natural number such that

$$\left(\begin{array}{c} | \\ | \\ | \\ | \end{array} - \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \Big| \delta^{ij} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right)^{N_{\alpha\beta\gamma\delta}^2} = 0. \quad (21)$$

(ii) we define the quadruple inner product of them as

$$c_{\alpha\beta\gamma\delta} = \begin{array}{c} \Lambda_{\alpha\beta\delta}^{L(3)} \quad \dots \quad \Lambda_{\alpha\beta\gamma}^{R(3)} \\ \text{---} \\ \Lambda_{\beta\gamma\delta}^{L(3)} \quad \text{---} \quad \Lambda_{\alpha\gamma\delta}^{R(3)} \\ \Lambda_{\alpha\delta}^{L(2)} \quad \text{---} \quad \Lambda_{\alpha\gamma}^{R(2)} \end{array}, \quad (22)$$

where the number of vertical lines is $\min(N_{\alpha\beta\gamma\delta}^1, N_{\alpha\beta\gamma\delta}^2)$.

Viewing the three-leg tensors as vertices, Eq. (22) can be regarded as a tetrahedron as in Fig. 4(b) with some additional tensors. For Eq. (22) to serve as a valid definition of the quadruple inner product and the higher Berry phase, it must have certain properties. In fact, we can show:

Proposition 4.

Let $|\psi_\alpha\rangle$, $|\psi_\beta\rangle$, $|\psi_\gamma\rangle$ and $|\psi_\delta\rangle$ be physically identical four states, and let $c_{\alpha\beta\gamma\delta}$ be the quadruple inner product of them. Then,

(i) $N_{\alpha\beta\gamma\delta}^1$ and $N_{\alpha\beta\gamma\delta}^2$ are finite natural number.

(ii) $c_{\alpha\beta\gamma\delta}$ is a non-zero complex number.

(iii) For physically identical five states $|\psi_\alpha\rangle$, $|\psi_\beta\rangle$, $|\psi_\gamma\rangle$, $|\psi_\delta\rangle$ and $|\psi_\eta\rangle$, $\{c_{\alpha\beta\gamma\delta}\}$ satisfies the cocycle condition

$$c_{\alpha\beta\gamma\delta} c_{\alpha\beta\gamma\eta}^{-1} c_{\alpha\beta\delta\eta} c_{\alpha\gamma\delta\eta}^{-1} c_{\beta\gamma\delta\eta} = 1. \quad (23)$$

(iv) Under the redefinition of the phase

$$\Lambda_{\alpha\beta\gamma}^{R(3)} \mapsto \lambda_{\alpha\beta\gamma} \Lambda_{\alpha\beta\gamma}^{R(3)}, \quad \Lambda_{\alpha\beta\gamma}^{L(3)} \mapsto \lambda_{\alpha\beta\gamma}^{-1} \Lambda_{\alpha\beta\gamma}^{L(3)}, \quad (24)$$

the quadruple inner product $c_{\alpha\beta\gamma\delta}$ changes as

$$c_{\alpha\beta\gamma\delta} \mapsto c_{\alpha\beta\gamma\delta} \lambda_{\alpha\beta\gamma} \lambda_{\alpha\beta\delta}^{-1} \lambda_{\alpha\gamma\delta} \lambda_{\beta\gamma\delta}^{-1}. \quad (25)$$

We will show this proposition in Sec. II B.

This concludes our construction of the quadruple inner product and the higher Berry phase. Furthermore, by using this proposition, we can construct a topological invariant of (families of) invertible states. We provide an outline of this construction and conclude this section.

Application 1: SPT invariant

Let's consider the case where a semi-injective PEPS $|A\rangle = (|\phi\rangle_A, O_A)$ is invariant under an onsite unitary G -symmetry $U(g)_{g \in G}$ for some finite group G . If the state is defined on a closed surface (e.g. 2-dimensional torus), it is completely identical before and after the action of the symmetry. However, G action has a non-trivial effect on the virtual Hilbert space in general. Let $|A^g\rangle$ be a semi-injective PEPS. Let's denote the state obtained by acting the symmetry $U(g)$ on $|A\rangle$ by $|A^g\rangle$, that is, $|A^g\rangle := (|\phi\rangle_A, U(g)O_A)$. Because of the symmetry, $\{|A^g\rangle\}_{g \in G}$ represent identical states and we can evaluate the quadruple inner product for any quadruple states taken from $\{|A^g\rangle\}_{g \in G}$:

$$c_{ghkl} := \text{quadruple inner product of } |A^g\rangle, |A^h\rangle, |A^k\rangle \text{ and } |A^l\rangle. \quad (26)$$

From Prop. 4, we see that c_{ghkl} defines an element $[c_{ghkl}]$ of the group cohomology $H^3(G; \mathbb{C}^\times)$. It is known that the classification of 2-dimensional SPT phases with G symmetry is given by $H^3(G; \mathbb{C}^\times)$ [4, 6] and the quadruple inner product is an invariant for SPT phases.

Application 2: higher Berry phase

Let's consider a family of gapped Hamiltonians over a parameter space X and assume that the ground states admit semi-injective PEPS representations. Though the parent Hamiltonian is parametrized over X , in general, we cannot have global parameterization of ground states. For example, if the ordinary Berry phase is nontrivial, we cannot take a global phase fixing of the state. Similarly, the quadruple inner product measures the higher obstruction to take a global PEPS expression of the family. In the next section, we will explain this point using a concrete model.

Let $\{U_\alpha\}$ be an open covering of the parameter space X , and we take a semi-injective PEPS representation $|A_\alpha(x)\rangle$ ($x \in U_\alpha$) for each patch. Then, for each point of quadruple intersection $x \in U_{\alpha\beta\gamma\delta} := U_\alpha \cap U_\beta \cap U_\gamma \cap U_\delta$, we have physically identical semi-injective PEPS, $|A_\alpha\rangle, |A_\beta\rangle, |A_\gamma\rangle$ and $|A_\delta\rangle$. Thus, we can evaluate the quadruple inner product of them:

$$c_{\alpha\beta\gamma\delta} := \text{quadruple inner product of } |A_\alpha\rangle, |A_\beta\rangle, |A_\gamma\rangle \text{ and } |A_\delta\rangle. \quad (27)$$

From Prop. 4, we see that $c_{\alpha\beta\gamma\delta}$ defines an element $[c_{\alpha\beta\gamma\delta}]$ of the Čech cohomology $H^3(X; \text{sh}(\mathbb{C}^\times)) \simeq H^4(X; \mathbb{Z})$. It is known that the higher Berry phase of 2-dimensional invertible states takes its value in $H^3(X; \text{sh}(\mathbb{C}^\times)) \simeq H^4(X; \mathbb{Z})$, and this implies that the quadruple inner product detects the higher Berry phase. We will carry out an explicit computation of the invariant in Sec. IV for the model parametrized by $\mathbb{R}P^4$.

These two applications of the quadruple inner product are related as follows. The basic idea is that SPT ground states (Hamiltonians) may be realized at particular, special locations of some parameter space X . While generic invertible states in a given family may not respect any symmetry, symmetry enhancement can happen at these special points. In the field theory context, it is known that various SPT phases can be realized as a gapped phase of a quantum non-linear sigma model [34, 42, 43].

In the SPT context, the gauge degrees of freedom of PEPS tensors we discussed are closely related to the symmetry protecting SPT phases. While the symmetry leaves SPT ground states invariant, PEPS tensors undergo a gauge transformation. Thus, for the SPT models, invertible states in different gauge choices can be obtained by acting with the symmetry. More precisely, while the symmetry action does not change the state in the bulk, it can be implemented non-trivially on virtual bonds and on a boundary as non-onsite symmetry; on the boundary, the symmetry operator is given by an MPU. Furthermore, this means that the eigen MPO of the mixed transfer matrix discussed above are given by boundary symmetry unitary of the SPT phase.

Ref. [32] extracted a 3-cocycle from G -symmetric semi-injective PEPS, and showed that it is an SPT invariant. In terms of the detection of SPT phases, the quadruple inner product and the invariants in Ref. [32] are exactly the same quantity. Therefore, the quadruple inner product can be used as an invariant for SPT phases with general G symmetry.

Using the equation $X^i = WY^iV + N^i$ once again, expand the second term of the above equation. Each time we pick WY^iV in the expansion (even once), the corresponding contribution is zero because of Eq. (32). If we choose the term N^i from all factors, then that term becomes zero also since m is greater than the nilpotency length N_0 . Therefore, the second term of Eq. (34) is zero, and Eq. (33) holds.⁷ \square

Therefore, if there is a sufficient number of X^i 's, we can partially "zip" the tensors by the reduction. In particular, when the nilpotency length N_0 is zero, Eq. (33) is called the zipper condition, e.g. [44, 45].

2. Fixed Points of the 2- and 3-Transfer Matrices

We can construct the fixed-point tensors of the 2- and 3-transfer matrices as concatenations of some reduction tensors as follows.

⁷ In the following, we will denote a reduction tensor as a gray box to distinguish it from the fixed point tensor.

Lemma 1.

(i) Let $(V_{\alpha\beta}, W_{\alpha\beta})$ be a reduction from $\{\sum_j B_{\alpha\beta}^{ij} \otimes B_{\beta\alpha}^{jk}\}$ to $\{\delta^{ik}\}$. Then, the right fixed point $\Lambda_{\alpha\beta}^{R(2)}$ of the 2-transfer matrix can be written as

$$\Lambda_{\alpha\beta}^{R(2)} = [T_{\alpha\beta}^{(2)}]^{N_{\alpha\beta}} \cdot W_{\alpha\beta} = \begin{array}{c} \vdots \\ \vdots \end{array} \left[\begin{array}{c} \vdots \\ \vdots \end{array} \right] W_{\alpha\beta}, \quad (35)$$

where $N_{\alpha\beta}$ is the nilpotency length of the reduction $(V_{\alpha\beta}, W_{\alpha\beta})$. Similarly, the left fixed point $\Lambda_{\alpha\beta}^{L(2)}$ of the 2-transfer matrix $T_{\alpha\beta}^{(2)}$ can be written as

$$\Lambda_{\alpha\beta}^{L(2)} = V_{\alpha\beta} \cdot [T_{\alpha\beta}^{(2)}]^{N_{\alpha\beta}} = V_{\alpha\beta} \left[\begin{array}{c} \vdots \\ \vdots \end{array} \right]. \quad (36)$$

(ii) Let $(V_{\alpha\beta\gamma}, W_{\alpha\beta\gamma})$ be a reduction from $\{\sum_j B_{\alpha\beta}^{ij} \otimes B_{\beta\gamma}^{jk}\}$ to $\{B_{\alpha\gamma}^{ik}\}$. Then, the right fixed point $\Lambda_{\alpha\beta\gamma}^{R(2,1)}$ of the right (2,1)-transfer matrix $T_{\alpha\beta\gamma}^{R(2,1)}$ can be written as

$$\Lambda_{\alpha\beta\gamma}^{R(2,1)} = [T_{\alpha\beta\gamma}^{R(2,1)}]^{N_{\alpha\beta\gamma}} \cdot W_{\alpha\beta\gamma} = \begin{array}{c} \vdots \\ \vdots \end{array} \left(\begin{array}{c} \vdots \\ \vdots \end{array} \right)^{(T_{\alpha\beta\gamma}^{R(2,1)})^{N_{\alpha\beta\gamma}}} W_{\alpha\beta\gamma}, \quad (37)$$

where $N_{\alpha\beta\gamma}$ is the nilpotency length of $T_{\alpha\beta\gamma}^{R(2,1)}$. Similarly, the left fixed point $\Lambda_{\alpha\beta\gamma}^{L(2,1)}$ of the left (2,1)-transfer matrix $T_{\alpha\beta\gamma}^{L(2,1)}$ can be written as

$$\Lambda_{\alpha\beta\gamma}^{L(2,1)} = V_{\alpha\beta\gamma} \cdot [T_{\alpha\beta\gamma}^{L(2,1)}]^{N_{\alpha\beta\gamma}} = \begin{array}{c} \vdots \\ \vdots \end{array} \left(\begin{array}{c} \vdots \\ \vdots \end{array} \right)^{(T_{\alpha\beta\gamma}^{L(2,1)})^{N_{\alpha\beta\gamma}}} V_{\alpha\beta\gamma}. \quad (38)$$

Proof. (i) By multiplying the transfer matrix $T_{\alpha\beta}^{(2)}$ to the right-hand side of Eq. (35),

$$T_{\alpha\beta}^{(2)} \cdot [T_{\alpha\beta}^{(2)}]^{N_{\alpha\beta}} \cdot W_{\alpha\beta} = \begin{array}{c} \vdots \\ \vdots \end{array} \left[\begin{array}{c} \vdots \\ \vdots \end{array} \right] W_{\alpha\beta} = \begin{array}{c} \vdots \\ \vdots \end{array} \left[\begin{array}{c} \vdots \\ \vdots \end{array} \right] = [T_{\alpha\beta}^{(2)}]^{N_{\alpha\beta}} \cdot W_{\alpha\beta}. \quad (39)$$

Here, we used the broken zipper equation Eq. (33). Thus, the tensor is the fixed point unless it is the zero tensor. By multiplying left reduction $V_{\alpha\beta}$ to the right-hand side of Eq. (35), the value of the diagram turns out to be the size of the matrices $\{B_{\alpha\beta}^{ij}\}$:

$$V_{\alpha\beta} \Lambda_{\alpha\beta}^{R(2)} = V_{\alpha\beta} \left[\begin{array}{c} \vdots \\ \vdots \end{array} \right] W_{\alpha\beta} = \left[\begin{array}{c} \vdots \\ \vdots \end{array} \right] \left[\begin{array}{c} \vdots \\ \vdots \end{array} \right] \cdots = 1. \quad (40)$$

This implies that the tensor is nonzero. Therefore, the right-hand side of Eq. (35) is the fixed point of the transfer matrix $T_{\alpha\beta}^{(2)}$. Similarly, we can show that the right-hand side of Eq. (36) is the left fixed point of the transfer matrix $T_{\alpha\beta}^{(2)}$.

(ii) By multiplying the transfer matrix $T_{\alpha\beta\gamma}^{R(2,1)}$ to the right-hand side of Eq. (37),

$$\begin{aligned}
T_{\alpha\beta\gamma}^{R(2,1)} \cdot [T_{\alpha\beta\gamma}^{R(2,1)}]^{N_{\alpha\beta\gamma}} \cdot W_{\alpha\beta\gamma} &= \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \\ \text{Diagram 4} \end{array} \\
&= [T_{\alpha\beta\gamma}^{R(2,1)}]^{N_{\alpha\beta\gamma}} \cdot W_{\alpha\beta\gamma}. \tag{41}
\end{aligned}$$

Here, we used the broken zipper equation Eq. (33). Thus, the tensor is the fixed point unless it is the zero tensor. By multiplying left reduction $V_{\alpha\beta\gamma}$ to the right-hand side of Eq. (37), the value of the diagram turns out to be the identity matrix:

$$V_{\alpha\beta\gamma} \Lambda_{\alpha\beta\gamma}^{R(2,1)} = \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \end{array} = \text{Diagram 4} = \text{Diagram 5}. \tag{42}$$

This implies that the tensor is nonzero. Therefore, the right-hand side of Eq. (37) is the fixed point of the transfer matrix $T_{\alpha\beta\gamma}^{R(2,1)}$. Similarly, we can show that the right-hand side of Eq. (38) is the fixed point of the transfer matrix $T_{\alpha\beta\gamma}^{L(2,1)}$. \square

Remark that, as a consequence of Lem. 1, the fixed point $(\Lambda_{\alpha\beta\gamma}^{L(2,1)}, \Lambda_{\alpha\beta\gamma}^{R(2,1)})$ itself is a reduction pair.

3. Fixed Points of the 4-Transfer Matrix

We can construct the fixed-point tensor of the 4-transfer matrix as a concatenation of the fixed points of the 2- and 3-transfer matrices:

Lemma 2.

(i) Let $\Lambda_{\alpha\beta\gamma\delta}^{R(4)}$ be a 4-leg tensor defined by

$$\Lambda_{\alpha\beta\gamma\delta}^{R(4)} = [T_{\alpha\beta\gamma}^{(4)}]^{N_{\alpha\beta\gamma\delta}} \Lambda_{\alpha\beta\gamma}^{R(2,1)} \Lambda_{\alpha\gamma\delta}^{R(2,1)} \Lambda_{\alpha\delta}^R = \text{Diagram}, \quad (43)$$

where $N_{\alpha\beta\gamma\delta}$ is the nilpotency length of the reduction. Then, $\Lambda_{\alpha\beta\gamma\delta}^{R(4)}$ is the right fixed point of the 4-transfer matrix $T_{\alpha\beta\gamma\delta}^{(4)}$.

(ii) Let $\Lambda_{\alpha\beta\gamma\delta}^{L(4)}$ be a 4-leg tensor defined by

$$\Lambda_{\alpha\beta\gamma\delta}^{L(4)} = \Lambda_{\alpha\delta}^L \Lambda_{\alpha\gamma\delta}^{L(2,1)} \Lambda_{\alpha\beta\gamma}^{L(2,1)} [T_{\alpha\beta\gamma}^{(4)}]^{N_{\alpha\beta\gamma\delta}} = \text{Diagram} \quad (44)$$

Then, $\Lambda_{\alpha\beta\gamma\delta}^{L(4)}$ is the left fixed point of the 4-transfer matrix $T_{\alpha\beta\gamma\delta}^{(4)}$.

Proof. (i) By multiplying the 4-transfer matrix $T_{\alpha\beta\gamma\delta}^{(4)}$ to the right-hand side of Eq. (51),

$$\begin{aligned} \text{Diagram} &= \text{Diagram} = \text{Diagram} \\ &= \text{Diagram} = \text{Diagram} \\ &= \text{Diagram} = \text{Diagram}. \end{aligned} \quad (45)$$

Here, we used the broken zipper equation Eq. (33). Therefore, we can zip the top fixed point tensor until the number of remaining transfer matrices becomes $N_{\alpha\beta\gamma}$:

$$\text{Diagram} = \text{Diagram} \quad (46)$$

Similarly, we can zip the middle fixed point tensor until the number of the remaining transfer matrices between the top and middle tensors becomes $N_{\alpha\gamma\delta}$:

$$\text{Diagram} = \text{Diagram} \quad (47)$$

Since there are $N_{\alpha\beta}$ transfer matrices remaining between the middle and the rightmost fixed point tensor, it is possible to zip the rightmost transfer matrix:

$$\text{Diagrammatic equation (48)} \tag{48}$$

Consequently, the tensor is the fixed point unless it is the zero tensor. By multiplying left reduction tensors to the right-hand side of Eq. (51), the value of the diagram turns out to be the size of the matrices $\{B_{\alpha\beta}^{ij}\}$:

$$\text{Diagrammatic equation (49)} \tag{49}$$

Similarly, we can zip the second and the third reduction tensors. Consequently, the left-hand side turns out to be 1:

$$\text{Diagrammatic equation (50)} = 1. \tag{50}$$

This implies that the tensor is nonzero. Therefore, the right-hand side of Eq. (51) is the right fixed point of the transfer matrix $T_{\alpha\beta\gamma\delta}^{(4)}$.

(ii) Similarly, we can show that the right-hand side of Eq. (52) is the left fixed point of the transfer matrix $T_{\alpha\beta\gamma\delta}^{(4)}$. \square

By using the fixed point tensors, we can construct the fixed point of the 4-transfer matrix $T_{\alpha\beta\gamma\delta}^{(4)}$ in a different manner:

Lemma 3.

(i) Let $\tilde{\Lambda}_{\alpha\beta\gamma\delta}^{R(4)}$ be a 4-leg tensor defined by

$$\tilde{\Lambda}_{\alpha\beta\gamma\delta}^{R(4)} := T_{\alpha\beta\gamma}^{(4)} \Lambda_{\beta\gamma\delta}^{R(2,1)} \Lambda_{\alpha\beta\delta}^{R(2,1)} \Lambda_{\alpha\delta}^R = \text{Diagram} \quad (51)$$

Then, $\tilde{\Lambda}_{\alpha\beta\gamma\delta}^{R(4)}$ is the right fixed point of the 4-transfer matrix $T_{\alpha\beta\gamma\delta}^{(4)}$.

(ii) Let $\tilde{\Lambda}_{\alpha\beta\gamma\delta}^{L(4)}$ be a 4-leg tensor defined by

$$\tilde{\Lambda}_{\alpha\beta\gamma\delta}^{L(4)} = \Lambda_{\alpha\delta}^L \Lambda_{\alpha\gamma\delta}^{L(2,1)} \Lambda_{\alpha\beta\gamma}^{L(2,1)} T_{\alpha\beta\gamma}^{(4)} = \text{Diagram} \quad (52)$$

Then, $\tilde{\Lambda}_{\alpha\beta\gamma\delta}^{L(4)}$ is the left fixed point of the 4-transfer matrix $T_{\alpha\beta\gamma\delta}^{(4)}$.

The proof is the same as that of Lem. 2.

Since the fixed point of the 4-transfer matrix $T_{\alpha\beta\gamma\delta}^{(4)}$ is unique, $\Lambda_{\alpha\beta\gamma\delta}^{R(4)}$ and $\tilde{\Lambda}_{\alpha\beta\gamma\delta}^{R(4)}$ are proportional to each other. We denote the proportionality constant as $c_{\alpha\beta\gamma\delta}$:

$$\Lambda_{\alpha\beta\gamma\delta}^{R(4)} = c_{\alpha\beta\gamma\delta} \tilde{\Lambda}_{\alpha\beta\gamma\delta}^{R(4)} \iff \text{Diagram} = c_{\alpha\beta\gamma\delta} \times \text{Diagram} \quad (53)$$

By multiplying the left fixed point $\tilde{\Lambda}_{\alpha\beta\gamma\delta}^{L(4)}$ from the left,

$$\text{Diagram} = c_{\alpha\beta\gamma\delta} \times \text{Diagram} \quad (54)$$

The diagram included on the right-hand side becomes 1 due to the normalization condition. We thus obtain Eq. (22).

4. Proof of Props. 3 and 4

We are now ready to proceed to the proof of Props. 3 and 4. First, we show Prop. 3. As we pointed out at the end of Sec. IIB 2, $(\Lambda_{\alpha\beta}^{L(2)}, \Lambda_{\alpha\beta}^{R(2)})$ and $(\Lambda_{\alpha\beta\gamma}^{L(3)}, \Lambda_{\alpha\beta\gamma}^{R(3)})$ are reduction tensors. Thus, their inner product, defined by

$$(\Lambda_{\alpha\beta}^{L(2)}, \Lambda_{\alpha\beta}^{R(2)}) = \Lambda_{\alpha\beta}^{L(2)} \boxed{\phantom{\Lambda_{\alpha\beta}^{R(2)}}} \Lambda_{\alpha\beta}^{R(2)} \quad (55)$$

and

$$(\Lambda_{\alpha\beta\gamma}^{L(3)}, \Lambda_{\alpha\beta\gamma}^{R(3)}) = \Lambda_{\alpha\beta\gamma}^{L(2,1)} \boxed{\phantom{\Lambda_{\alpha\beta\gamma}^{R(2,1)}}} \Lambda_{\alpha\beta\gamma}^{R(2,1)} \quad (56)$$

III. THE CZX MODEL PROTECTED BY $\mathbb{Z}/2\mathbb{Z}$ SYMMETRY

Having gone through the generalities of the quadruple inner product, we will now discuss some applications and examples outlined at the end of Sec. II A. Here, we will use (2+1)d SPT phases to illustrate the use of the quadruple inner product.

As a particular example of SPT phases, let us consider the CZX model [33] – this is an example of (2+1)d SPT protected by $\mathbb{Z}/2\mathbb{Z}$ on-site unitary symmetry. Let us consider the square lattice with 2^4 -dimensional local Hilbert space (i.e., four copies of spin 1/2) for each site. Let p denote the plaquette label, and i represent the site label. The local Hamiltonian labeled by p is defined as an operator that acts on the plaquette p and its four neighboring edges. To define this, we label the top, bottom, left, and right edges adjacent to plaquette p as $u(p)$, $d(p)$, $l(p)$, and $r(p)$ respectively. The state of the 2 sites belonging to an edge is written as $\{|i, j\rangle_{k(p)} \mid i, j = \uparrow, \downarrow, k = u, d, l, r\}$, where the order inside the ket is arranged in a clockwise manner. For instance, the state where the upper site is \uparrow and the lower site is \downarrow on the edge adjacent to the right of plaquette p is expressed as $|\uparrow\downarrow\rangle_{r(p)}$. The sites belonging to site i will be labeled in a counterclockwise manner starting from the top-left site as $i(1)$, $i(2)$, $i(3)$, and $i(4)$.

The local Hamiltonian for a plaquette p is defined by

$$h_p := -X_p^{(4)} \otimes \bigotimes_{k=u,d,l,r} P_{k(p)}^{(2)}, \quad (62)$$

$$\begin{aligned} \text{where } X_p^{(4)} &:= |\uparrow\uparrow\uparrow\rangle \langle\downarrow\downarrow\downarrow|_p + |\downarrow\downarrow\downarrow\rangle \langle\uparrow\uparrow\uparrow|_p, \\ P_{k(p)}^{(2)} &:= |\uparrow\uparrow\rangle \langle\uparrow\uparrow|_{k(p)} + |\downarrow\downarrow\rangle \langle\downarrow\downarrow|_{k(p)} \quad (k = u, d, l, r). \end{aligned} \quad (63)$$

The total Hamiltonian is defined by the sum of the local Hamiltonians, $H_{\text{CZX}} := \sum_p h_p$. This Hamiltonian commutes with the CZX symmetry $U_{\text{CZX}} := \bigotimes_i u_{\text{CZX},i}$, where

$$u_{\text{CZX},i} := \left(\bigotimes_{k=1}^4 X_{i(k)} \right) \otimes CZ_{i(1)i(2)} \otimes CZ_{i(2)i(3)} \otimes CZ_{i(3)i(4)} \otimes CZ_{i(4)i(1)}. \quad (64)$$

Here, $CZ_{ij} := \sum_{ab} (-1)^{ab} |a_i b_j\rangle \langle a_i b_j|$ is the CZ-gate operator acting on sites i, j . U_{CZX} generates onsite $\mathbb{Z}/2\mathbb{Z}$ symmetry. It is known that the ground state on the closed system is unique and given by

$$|G.S.\rangle := \bigotimes_p \frac{1}{\sqrt{2}} (|\uparrow\uparrow\uparrow\rangle_p + |\downarrow\downarrow\downarrow\rangle_p). \quad (65)$$

This is an example of (2+1)d SPT phases protected by $\mathbb{Z}/2\mathbb{Z}$ symmetry [33]. The ground state of the CZX model can be expressed in terms of a semi-injective PEPS [32].

As mentioned at the end of Sec. II A, what is important for our purposes is the boundary symmetry operators. When making a boundary, there appear free degrees of freedom on the plaquettes that are cut open by the boundary. The boundary symmetry operators are obtained by projecting the (bulk) symmetry operators to the space of the free degrees of freedom on the boundary. Explicitly, they are given by

$$\mathcal{O}_I = \mathbf{1}, \quad \mathcal{O}_Z = \bigotimes_i CZ_{i,i+1} \bigotimes_i X_i, \quad (66)$$

where i labels boundary sites. The boundary symmetry operators can be represented as an MPU with

$$B_I^{01} = B_I^{10} = 0, \quad B_I^{00} = B_I^{11} = 1, \quad (67)$$

and

$$B_Z^{01} = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}, \quad B_Z^{10} = \begin{pmatrix} 0 & 0 \\ 1 & -1 \end{pmatrix}, \quad B_Z^{00} = B_Z^{11} = 0. \quad (68)$$

Remark that these tensors are in the right canonical form.

Now, let's consider the three-leg tensors $\Lambda_{ZZI}^{R(2,1)}$, $\Lambda_{ZIZ}^{R(2,1)}$ and $\Lambda_{IZZ}^{R(2,1)}$. As $\Lambda_{ZIZ}^{R(2,1)}$ and $\Lambda_{IZZ}^{R(2,1)}$ intertwine \mathcal{O}_Z and \mathcal{O}_Z , they are the identity matrix. $\Lambda_{ZZI}^{R(2,1)}$ is the only nontrivial one and is represented as 4×1 tensor. We need to compute the composed tensor $B_{ZZ}^{ij} := \sum_k B_Z^{ik} \otimes B_Z^{kj}$:

$$B_{ZZ}^{00} = \sum_k B_Z^{0k} \otimes B_Z^{k0} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & -1 & 1 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (69)$$

$$B_{ZZ}^{11} = \sum_k B_Z^{1k} \otimes B_Z^{k1} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 1 & -1 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (70)$$

$$B_{ZZ}^{01} = B_{ZZ}^{10} = 0. \quad (71)$$

Here, given 2-dimensional vector spaces V and W with orthonormal bases v_0, v_1 and w_0, w_1 respectively, the basis for the tensor product space is fixed in the order $v_0 \otimes w_0, v_0 \otimes w_1, v_1 \otimes w_0$, and $v_1 \otimes w_1$. As illustrated in [46] and App. A.4 of [10], the reduction pair (V_{ZZI}, W_{ZZI}) from B_{ZZ}^{ij} to B_I^{ij} is given by

$$V_{ZZI} = \frac{1}{\sqrt{2}} (0, 1, -1, 0) \iff (V_{ZZI})_0^{ab} = \frac{(-1)^b}{\sqrt{2}} (1 - \delta_{ab}), \quad (72)$$

$$W_{ZZI} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ -1 \\ 0 \end{pmatrix} \iff (W_{ZZI})_{ab}^0 = \frac{(-1)^b}{\sqrt{2}} (1 - \delta_{ab}), \quad (73)$$

where $a, b = 0, 1$ and δ_{ab} is the Kronecker delta. Therefore, the reduced MPU matrices $\tilde{B}_I^{ij} := (V_{ZZI})^\dagger B_{ZZ}^{ij} W_{ZZI}$ are given by

$$\tilde{B}_I^{00} = \tilde{B}_I^{00} = -1, \quad \tilde{B}_I^{01} = \tilde{B}_I^{10} = 0. \quad (74)$$

Thus, the relative phase between B_I^{ij} and \tilde{B}_I^{ij} is $c_{ZZI} = -1$. This implies that the phase c in Prop. 1 is -1 . Therefore, the normalized transfer matrix

$$T_{ZZI}^{R(2,1)} = -\frac{1}{2} \sum_{ij} B_{ZZ}^{ij} \otimes B_I^{ij\dagger} \quad (75)$$

has a unique fixed point tensor. One can check that the right reduction tensor itself satisfies the higher eigenequation. In fact, when calculating the left-hand side of the higher eigenequation, we find

$$-\frac{1}{2} \sum_{ij} B_{ZZ}^{ij} W_{ZZI} B_I^{ij} = c_{ZZ}^I W_{ZZI}. \quad (76)$$

Therefore, we can take $\Lambda_{ZZI}^{R(2,1)} = W_{ZZI}$. Similarly, the only nontrivial left fixed point is $\Lambda_{ZZI}^{L(2,1)}$. One can check that the left reduction V_{ZZI} satisfies the fixed point equation of the left transfer matrix $T_{ZZI}^{L(2,1)}$. Therefore, we can take $\Lambda_{ZZI}^{L(2,1)} = V_{ZZI}$. Note that the normalization condition Eq.(17) holds. In this model, the nilpotency length $N_{\alpha\beta\gamma\delta}$ in Prop. 2 is zero. Thus, we do not need to insert the vertical lines in Eq. (22).

By using these tensors Λ_{ZZI} , Λ_{ZIZ} and Λ_{IZZ} , let us now compute the quadruple inner product. As a demonstration, the quadruple inner product c_{IZIZ} can be computed as

$$c_{IZIZ} = \sum_{ab} (\Lambda_{ZZI})_{ab}^0 (\Lambda_{ZZI}^\dagger)_{ba}^0 = -\frac{1}{\sqrt{2}} \times \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \times \frac{1}{\sqrt{2}} = -1. \quad (77)$$

After a similar calculation, we can readily check the following result:

$$c_{g_1 g_2 g_3 g_4} = \begin{cases} -1 & (g_1, g_2, g_3, g_4) = (I, Z, I, Z), (Z, I, Z, I) \\ 1 & \text{otherwise.} \end{cases} \quad (78)$$

This is a nontrivial cocycle in $H^3(\mathbb{Z}/2\mathbb{Z}; U(1))$.

IV. THE MODEL PARAMETERIZED OVER $\mathbb{R}P^4$

In this section, we will discuss a concrete model (a family of parameterized (2+1)d Hamiltonians or (2+1)d invertible states) characterized by the non-trivial higher Berry phase. The model is parameterized by $X = \mathbb{R}P^4$. We will show that the model is nontrivial in $H^4(\mathbb{R}P^4; \mathbb{Z}/2\mathbb{Z}) \simeq \mathbb{Z}/2\mathbb{Z}$.

A. The $\mathbb{R}P^4$ -Parametrized Model

The CZX model has the $\mathbb{Z}/2\mathbb{Z}$ symmetry. By interpolating this symmetry, we can construct a family of 2-d systems. To this end, we embed the local Hilbert space \mathbb{C}^2 to \mathbb{C}^4 and take the following orthonormal basis:

$$|+\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad |-\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \quad |\perp_1\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \quad |\perp_2\rangle = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}. \quad (79)$$

Let $\{\vec{z} = (z_1, z_2, z_3) \mid z_{1,2,3} \in \mathbb{C}, |\vec{z}|^2 = 1\}$ be a coordinate of S^5 and we define $|+(\vec{z})\rangle$ and $|-(\vec{z})\rangle$ by

$$|+(\vec{z})\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad |-(\vec{z})\rangle = \begin{pmatrix} 0 \\ z_1 \\ z_2 \\ z_3 \end{pmatrix}. \quad (80)$$

The parameterized basis is obtained from the basis (79) by a unitary transformation,

$$(|+(\vec{z})\rangle, |-(\vec{z})\rangle, |\perp_1(\vec{z})\rangle, |\perp_2(\vec{z})\rangle) = V(\vec{z})(|+\rangle, |-\rangle, |\perp_1\rangle, |\perp_2\rangle),$$

$$\text{where } V(\vec{z}) = \begin{pmatrix} 1 & & & \\ & z_1 & a & b \\ & z_2 & c & d \\ & z_3 & e & f \end{pmatrix} \quad (81)$$

for suitable $a, b, c, d \in \mathbb{C}$. Since $SU(3)$ is the nontrivial S^3 bundle over S^5 , taking the global families of unitary transformations over S^5 continuously should not be possible. However, since we will only use the 2-dimensional subspace of \mathbb{C}^4 spanned by $|+(\vec{z})\rangle, |-(\vec{z})\rangle$ to define a Hamiltonian, our model will be parametrized by S^5 globally.

Let $\sigma^x(\vec{z})$ and $\sigma^z(\vec{z})$ be parametrized Pauli matrices defined by

$$\sigma^x(\vec{z}) = |+(\vec{z})\rangle \langle +(\vec{z})| - |-(\vec{z})\rangle \langle -(\vec{z})|, \quad (82)$$

$$\sigma^z(\vec{z}) = |+(\vec{z})\rangle \langle -(\vec{z})| + |-(\vec{z})\rangle \langle +(\vec{z})|. \quad (83)$$

We introduce the polar coordinate:

$$z_1 = \cos(\theta_1) + i \sin(\theta_1) \cos(\theta_2), \quad (84)$$

$$z_2 = \sin(\theta_1) \sin(\theta_2) \cos(\theta_3) + i \sin(\theta_1) \sin(\theta_2) \sin(\theta_3) \cos(\theta_4), \quad (85)$$

$$z_3 = \sin(\theta_1) \sin(\theta_2) \sin(\theta_3) \sin(\theta_4) \sin(\theta_5) + i \sin(\theta_1) \sin(\theta_2) \sin(\theta_3) \sin(\theta_4) \cos(\theta_5), \quad (86)$$

element of the 4-th Deligne cohomology $\hat{H}^4(X; \mathbb{Z})$ [47]. A representative element of $\hat{H}^4(\mathbb{R}P^4; \mathbb{Z})$ is given by \mathbb{C}^\times -valued function $w_{\alpha\beta\gamma\delta}^{(0)}$ on $U_{\alpha\beta\gamma\delta}$, 1-form $w_{\alpha\beta\gamma}^{(1)}$ on $U_{\alpha\beta\gamma}$, 2-form $w_{\alpha\beta}^{(2)}$ on $U_{\alpha\beta}$, and 3-form $w_\alpha^{(3)}$ on U_α with suitable consistency conditions. We refer to $w^{(1)}$, $w^{(2)}$ and $w^{(3)}$ as higher connections. In our context, $w_{\alpha\beta\gamma\delta}^{(0)}$ is the higher Berry phase. As mentioned in Sec. IV B, the 3-leg tensor does not depend on \vec{z} . This means that $c_{\alpha\beta\gamma\delta}$ admits the trivial higher connections, i.e., $(w_\alpha^{(3)} = 0, w_{\alpha\beta}^{(2)} = 0, w_{\alpha\beta\gamma}^{(1)} = 0, w_{\alpha\beta\gamma\delta}^{(0)} = c_{\alpha\beta\gamma\delta})$ defines an element of the 4-th Deligne cohomology $\hat{H}^4(\mathbb{R}P^4; \mathbb{Z})$. In particular, the higher Berry phase is quantized to $\mathbb{Z}/2\mathbb{Z}$.

Following the integration theory of the Deligne cohomology [48–51], we evaluate the cocycle $(w_\alpha^{(3)} = 0, w_{\alpha\beta}^{(2)} = 0, w_{\alpha\beta\gamma}^{(1)} = 0, c_{\alpha\beta\gamma\delta})$ ⁸. First, we need to take a nontrivial homology class of $H_3(\mathbb{R}P^4; \mathbb{Z}) \simeq \mathbb{Z}/2\mathbb{Z}$ as an integration surface. This is generated by $\mathbb{R}P^3 \subset \mathbb{R}P^4$. Let's consider the open covering of $\mathbb{R}P^3$. We will illustrate $\mathbb{R}P^3$ by considering a cube and identifying antipodal points of the surface. For this cube, we take balls centered on the face-centered cubic lattice and the body center. Due to the identification of antipodal points, the number of such balls is eight. Furthermore, we consider balls centered at the midpoint of each edge. Again, due to the identification of antipodal points, the number of these balls is six. We adjust the radii of these balls appropriately to cover the entire cube.

To simplify the illustration, we flatten these intersections of two patches into planes and illustrate them as surfaces Fig. 6. Fortunately, as already mentioned, the transition MPU in the example is constant on the intersections. Therefore, we can flatten them without loss of information.




FIG. 6. An open covering of $\mathbb{R}P^3$.

All edges correspond to the parts where exactly three patches overlap. Therefore, a dual lattice provides a triangulation of $\mathbb{R}P^3$. We introduce a branching structure, namely, an orientation that does not include closed loops as follows:




FIG. 7. Branching structure of the triangulation

Here, to avoid complexity by incorporating all edges into a single figure, the edges were divided into three groups for illustration.

To compute the higher Berry phase, we need to assign the transfer MPU to each face, assign the fixed point tensors to each edge, and evaluate the quadruple inner product on each vertex. In the following, we explain the systematic assignment procedure.

⁸ See [22] as an example of the explicit computation of the integration.

Step 1: Transfer MPUs on faces To assign the transition MPU to each face systematically, we introduce the base points of each patch [22], which are illustrated by orange dots in Fig. 8. By using the base points, we define transition functions on the intersection $U_{12} = U_1 \cap U_2$ under the following rules:

1. Take a path starting from the base point of the patch U_1 and passing through U_{12} and terminating at the base point of U_2 .
2. If the path is through the surface of the cube, we give $\mathcal{O}\{B_Z^{ij}\}$ on the surface.

Under the rules, the configuration of the transition MPU of the model is illustrated in Fig. 9.




FIG. 8. The configuration of basepoints of $\mathbb{R}P^3$.




FIG. 9. The configuration of transition MPU. Blue faces represent intersections for which a non-trivial transition MPU is assigned.

Step 2: Fixed-point tensors on edges A triangulation is provided by taking the dual lattice. Recall that each triangle was a triple overlap $U_{\alpha\beta\gamma}$ in the original open-covering picture. Therefore, a 3-leg tensor, which is a fixed point of the transfer matrix (e.g. Fig. 4), is assigned to each triangle of the triangulation. Note that the 3-leg tensor has two incoming legs and one outgoing leg (or the opposite assignment). To distinguish these legs, we utilize a branching structure. As shown in Fig. 7, draw the dual lattice of the triangle and rotate the direction of the edges determined by the branching structure by 90 degrees clockwise. Depending on whether the resulting 3-leg tensor has two incoming legs or two outgoing legs, assign $\Lambda_{ZZI}^{R(2,1)}$ or $\Lambda_{ZZI}^{L(2,1)}$, respectively.

In this model, the 3-leg tensors are non-trivial only when both incoming legs are $\{B_Z^{ij}\}$ or both outgoing legs are $\{B_Z^{ij}\}$; otherwise, the 3-leg tensors are trivial, i.e., the Kronecker delta. We show the edges to which non-trivial 3-leg tensors are assigned in Fig. 10.

Step 3: Quadruple inner product on vertices Recall that each triangle is a triple overlap $U_{\alpha\beta\gamma\delta}$ in the original open-covering picture. Therefore, the quadruple inner product is assigned to each tetrahedron of the triangulation. Using the procedure introduced in Step 2, a 3-leg tensor is defined for each face of the tetrahedron, thus providing four 3-leg tensors in total. By gluing them together naturally, we can evaluate the quadruple inner product.

As an example, we evaluate the quadruple inner product of three vertices in Fig. 7. In the triangulation, the tetrahedra, which correspond to these vertices, are

(a)

(b)

(c)

(d)

(92)




FIG. 10. The configuration of nontrivial 3-leg tensors. Red edges represent 3-intersections for which a non-trivial 3-leg tensor is assigned.

where the edges with blue color imply that the transition MPU assigned to the edges are nontrivial. Let's compute the tetrahedra (b) in Eq. (92). For instance, focusing on the right-side triangle, according to the procedure explained in Step 2, the 3-leg tensor with 2-in 1-out configuration is assigned to this face, where the incoming legs are labeled with nontrivial elements of $\mathbb{Z}/2\mathbb{Z}$. Similarly, focusing on the left-side triangle, the 3-leg tensor with 1-in 2-out configuration is assigned to this face, where the outgoing legs are labeled with nontrivial elements of $\mathbb{Z}/2\mathbb{Z}$. We can easily check that the trivial 3-leg tensors are assigned to the other edges. When these 3-leg tensors are glued along the edges, the diagram for the quadruple inner product is

$$\begin{array}{c} \text{tetrahedron (b)} \end{array} = \sum_{ab} (\Lambda_{ZZI}^{R(2,1)})_{ab}{}^0 (\Lambda_{ZZI}^{L(2,1)})_0{}^{ab} = 1. \quad (93)$$

Generally, as with the tetrahedron (b) and (c) in Eq. (92), the quadruple inner product turns out to be trivial specifically in cases where exactly three edges have non-trivial transition functions. It can be confirmed that among the tetrahedra included in the triangulation, only the ones labeled (a) and (d) have four nontrivial edges. Therefore, it is sufficient to evaluate the quadruple inner products corresponding to these two tetrahedra. For the tetrahedron (d) in Eq. (92), the quadruple inner product is

$$\begin{array}{c} \text{tetrahedron (d)} \end{array} = \sum_{ab} (\Lambda_{ZZI}^{R(2,1)})_{ab}{}^0 (\Lambda_{ZZI}^{L(2,1)})_0{}^{ab} = 1. \quad (94)$$

On the other hand, for the tetrahedron (a) in Eq. (92), the quadruple inner product is

$$\begin{array}{c} \text{tetrahedron (a)} \end{array} = \sum_{ab} (\Lambda_{ZZI}^{R(2,1)})_{ab}{}^0 (\Lambda_{ZZI}^{L(2,1)})_0{}^{ba} = -1. \quad (95)$$

Therefore, the total higher Berry phase is -1 in $H^4(\mathbb{R}P^4; \mathbb{Z}) \simeq \mathbb{Z}/2\mathbb{Z}$, and this implies the nontriviality as a family.

V. GEOMETRIC INTERPRETATION: 2-GERBE

In this section, we provide a further discussion on our construction of the higher Berry phase and the topological invariant, from the perspective of a gerbe. In particular, we provide a geometric interpretation, called a 2-gerbe [52], of the family of semi-injective PEPS. We recall that the underlying gerbe structure of the higher Berry phase for (1+1)d many-body states (MPS) are discussed in Refs. [1] and [25]. To address the nontriviality of a family of PEPSs, we need to consider the interface of the interface and put the zipper tensor on it. The zipper tensor can be regarded as the redundancy of the transition MPU, and the transition MPU is the redundancy of the PEPS tensor. In this sense, the zipper tensor is the redundancy of the redundancy. Now, 2-gerbe is a sophisticated tool to handle the redundancy of the redundancy⁹. We note that Ref. [25] points out a 2-gerbe structure for (2+1)-dimensional PEPS using the bundle gerbe.

A 1-gerbe, or simply a gerbe, is a categorification of a line bundle, and a 2-gerbe is a categorification of a 1-gerbe. Therefore, we will first recall a line bundle and explain the relationship between line bundles and gerbes. Consider a (0 + 1)-dimensional unique gapped Hamiltonian parametrized by X . We fix an open covering $\{U_\alpha\}$ of X . We can take a family of normalized ground states $|\psi_\alpha(x)\rangle$ over each open set U_α . Note that the phase of the ground state can freely be changed. In other words, to take a family of the ground states $|\psi_\alpha(x)\rangle$ can be regarded as a gauge fixing of the phase redundancy of the states. At each point of the intersection $U_{\alpha\beta}$, we have two identical states in different gauges. Thus, in particular, it is possible to determine the phase difference between the two states, $\langle\psi_\alpha(x)|\psi_\beta(x)\rangle$. This is a physical realization of a line bundle.

Next, consider a (1 + 1)-dimensional unique gapped Hamiltonian parametrized by X . We assume that we can take a normal MPS representation $\{A^i(x)\}$ of the ground state at each point of X . As in the (0 + 1) dimensional case, this involves gauge fixing with respect to the redundancy of the MPS representation. Using the fundamental theorem of MPS, at each point of the two intersection $U_{\alpha\beta}$, we can take the unique largest eigenvector (or reduction tensor) $\Lambda_{\alpha\beta}$ of the mixed transfer matrix, and at the three intersection $U_{\alpha\beta\gamma}$, a triple product can be taken to obtain a U(1) function. That is, a line bundle spanned by $\Lambda_{\alpha\beta}$ is defined on the 2-intersection, and we can extract a U(1)-valued function on the three intersections. In the case of (0 + 1) dimensions, a family of one-dimensional vector spaces is defined on each open set, and a U(1)-valued function is provided at the two intersections. In contrast, in the case of (1 + 1) dimensions, a family of one-dimensional vector spaces is established at the two intersections, and a U(1)-valued function is provided at the three intersections. In this sense, the degree of intersection has increased by one. Such objects are mathematically referred to as U(1)-gerbes. The type of gerbe obtained by considering a family of normal MPS is called an MPS gerbe [1, 30].

Let's now consider a (2 + 1)-dimensional unique gapped Hamiltonian parametrized by X . We assume that we can take a semi-injective PEPS representation $\{A_\alpha^i(x)\}$ of the ground state at each point of U_α . Using the fundamental theorem of semi-injective, at each point of the two intersection $U_{\alpha\beta}$, we obtain an injective MPO $\mathcal{O}_{\alpha\beta}$ between $\{A_\alpha^i(x)\}$ and $\{A_\beta^i(x)\}$. If we assume the unitarity of the MPO, on the three intersection $U_{\alpha\beta\gamma}$, we have a unique fixed point $\Lambda_{\alpha\beta\gamma}$ of transfer matrix $T_{\alpha\beta\gamma}$, and we can extract a U(1)-valued function on the four intersection. That is, a line bundle spanned by $\Lambda_{\alpha\beta\gamma}$ is defined on the three intersection, and we can extract a U(1)-valued function on the four intersection. Such objects are mathematically referred to as Tor(U(1))-gerbes [52]. However, to extract the U(1)-valued function, we need to use the information of the transition MPU in general, and in such a case, this is not a 2-gerbe. In the case of the example in Sec. IV, we need not to insert the transition MPU. Therefore, in this case, 2-gerbe over $\mathbb{R}P^4$ is constructed from a family of semi-injective PEPS.

If the zipper condition holds, we do not need to insert the transition MPU. Thus, the zipper condition is a sufficient condition for the 2-gerbe description. More abstractly, a 2-gerbe over X is a 2-stack over X such that every 1-morphism is invertible up to 2-morphism and every 2-morphism is exactly invertible [52]. In the PEPS description, 1-morphisms correspond to the transition MPU $\mathcal{O}_{\alpha\beta}$ and 2-morphisms correspond to the fixed point tensors $\Lambda_{\alpha\beta\gamma}$. Since $\mathcal{O}_{\alpha\beta}$ is injective, we can take the inverse $\mathcal{O}_{\alpha\beta}^{-1}$ and reduction pairs from $\mathcal{O}_{\alpha\beta} \cdot \mathcal{O}_{\alpha\beta}^{-1}$ to the identity and from $\mathcal{O}_{\alpha\beta}^{-1} \cdot \mathcal{O}_{\alpha\beta}$ to the identity. This is the invertibility of 1-morphisms up

⁹ In general, n -category has information about the n th-redundancy of objects.

agreed to coordinate submissions of our papers to arXiv. This research was supported in part by Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development and by the Province of Ontario through the Ministry of Colleges and Universities. We are grateful to the long-term workshop YITP-T-23-01 held at YITP, Kyoto University, where a part of this work was done. S.O. is supported by JSPS KAKENHI Grant Number 23KJ1252 and 24K00522. S.R. is supported by a Simons Investigator Grant from the Simons Foundation (Award No. 566116). This work is supported by the Gordon and Betty Moore Foundation through Grant GBMF8685 toward the Princeton theory program.

-
- [1] S. Ohyama and S. Ryu, Higher structures in matrix product states (2023), [arXiv:2304.05356](#).
 - [2] Z.-C. Gu and X.-G. Wen, Tensor-entanglement-filtering renormalization approach and symmetry-protected topological order, *Physical Review B* **80**, [10.1103/physrevb.80.155131](#) (2009).
 - [3] F. Pollmann, E. Berg, A. M. Turner, and M. Oshikawa, Symmetry protection of topological phases in one-dimensional quantum spin systems, *Physical Review B* **85**, [10.1103/physrevb.85.075125](#) (2012).
 - [4] X. Chen, Z.-X. Liu, and X.-G. Wen, Two-dimensional symmetry-protected topological orders and their protected gapless edge excitations, *Physical Review B* **84**, [10.1103/physrevb.84.235141](#) (2011).
 - [5] N. Schuch, D. Pérez-García, and I. Cirac, Classifying quantum phases using matrix product states and projected entangled pair states, *Physical Review B* **84**, [10.1103/physrevb.84.165139](#) (2011).
 - [6] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, Symmetry protected topological orders and the group cohomology of their symmetry group, *Physical Review B* **87**, [10.1103/physrevb.87.155114](#) (2013).
 - [7] T. Senthil, Symmetry-protected topological phases of quantum matter, *Annual Review of Condensed Matter Physics* **6**, 299 (2015).
 - [8] C.-K. Chiu, J. C. Teo, A. P. Schnyder, and S. Ryu, Classification of topological quantum matter with symmetries, *Reviews of Modern Physics* **88**, [10.1103/revmodphys.88.035005](#) (2016).
 - [9] B. Zeng, X. Chen, D.-L. Zhou, and X.-G. Wen, Quantum information meets quantum matter – from quantum entanglement to topological phase in many-body systems (2018), [arXiv:1508.02595 \[cond-mat.str-el\]](#).
 - [10] J. I. Cirac, D. Pérez-García, N. Schuch, and F. Verstraete, Matrix product states and projected entangled pair states: Concepts, symmetries, theorems, *Reviews of Modern Physics* **93**, [10.1103/revmodphys.93.045003](#) (2021).
 - [11] D. J. Thouless, Quantization of particle transport, *Phys. Rev. B* **27**, 6083 (1983).
 - [12] A. Y. Kitaev, On the classification of short-range entangled states, *CSGP Program: Topological Phases of Matter* (2013).
 - [13] D. S. Freed and M. J. Hopkins, Reflection positivity and invertible topological phases, *Geometry and Topology* **25**, 1165–1330 (2021).
 - [14] Y. Ogata, An invariant of symmetry protected topological phases with on-site finite group symmetry for two-dimensional fermion systems, *Communications in Mathematical Physics* **395**, 405 (2022).
 - [15] A. Kapustin and L. Spodyneiko, Higher-dimensional generalizations of berry curvature, *Phys. Rev. B* **101**, 235130 (2020).
 - [16] A. Kapustin and L. Spodyneiko, Higher-dimensional generalizations of the thouless charge pump (2020), [arXiv:2003.09519](#).
 - [17] P.-S. Hsin, A. Kapustin, and R. Thorngren, Berry phase in quantum field theory: Diabolical points and boundary phenomena, *Physical Review B* **102**, [10.1103/physrevb.102.245113](#) (2020).
 - [18] K. Shiozaki, Adiabatic cycles of quantum spin systems, *Phys. Rev. B* **106**, 125108 (2022).
 - [19] Y. Choi and K. Ohmori, Higher berry phase of fermions and index theorem, *Journal of High Energy Physics* **2022**, [10.1007/jhep09\(2022\)022](#) (2022).
 - [20] C. Cordova, D. Freed, H. T. Lam, and N. Seiberg, Anomalies in the space of coupling constants and their dynamical applications I, *SciPost Physics* **8**, [10.21468/scipostphys.8.1.001](#) (2020).
 - [21] C. Cordova, D. Freed, H. T. Lam, and N. Seiberg, Anomalies in the space of coupling constants and their dynamical applications II, *SciPost Physics* **8**, [10.21468/scipostphys.8.1.002](#) (2020).
 - [22] S. Ohyama, Y. Terashima, and K. Shiozaki, Discrete higher berry phases and matrix product states (2023), [arXiv:2303.04252](#).
 - [23] A. Beaudry, M. Hermele, J. Moreno, M. Pflaum, M. Qi, and D. Spiegel, Homotopical foundations of parametrized quantum spin systems (2023), [arXiv:2303.07431 \[math-ph\]](#).
 - [24] X. Wen, M. Qi, A. Beaudry, J. Moreno, M. J. Pflaum, D. Spiegel, A. Vishwanath, and M. Hermele, Flow of higher berry curvature and bulk-boundary correspondence in parametrized quantum systems, *Physical Review B* **108**, [10.1103/physrevb.108.125147](#) (2023).

- [25] M. Qi, D. T. Stephen, X. Wen, D. Spiegel, M. J. Pflaum, A. Beaudry, and M. Hermele, Charting the space of ground states with tensor networks (2023), [arXiv:2305.07700 \[cond-mat.str-el\]](#).
- [26] K. Shiozaki, N. Heinsdorf, and S. Ohyama, Higher berry curvature from matrix product states (2023), [arXiv:2305.08109 \[quant-ph\]](#).
- [27] A. Artymowicz, A. Kapustin, and N. Sopenko, Quantization of the higher berry curvature and the higher thouless pump (2023), [arXiv:2305.06399 \[math-ph\]](#).
- [28] S. Ohyama, K. Shiozaki, and M. Sato, Generalized thouless pumps in $(1 + 1)$ -dimensional interacting fermionic systems, *Phys. Rev. B* **106**, 165115 (2022).
- [29] K. Shiozaki, N. Heinsdorf, and S. Ohyama, Higher berry curvature from matrix product states (2023), [arXiv:2305.08109 \[quant-ph\]](#).
- [30] M. Qi, D. T. Stephen, X. Wen, D. Spiegel, M. J. Pflaum, A. Beaudry, and M. Hermele, Charting the space of ground states with tensor networks (2023), [arXiv:2305.07700 \[cond-mat.str-el\]](#).
- [31] D. D. Spiegel, A c^* -algebraic approach to parametrized quantum spin systems and their phases in one spatial dimension (2023), [arXiv:2305.07951 \[math-ph\]](#).
- [32] A. Molnar, Y. Ge, N. Schuch, and J. I. Cirac, A generalization of the injectivity condition for projected entangled pair states, *Journal of Mathematical Physics* **59**, 021902 (2018).
- [33] X. Chen, Z.-X. Liu, and X.-G. Wen, Two-dimensional symmetry-protected topological orders and their protected gapless edge excitations, *Physical Review B* **84**, 10.1103/physrevb.84.235141 (2011).
- [34] D. V. Else and C. Nayak, Classifying symmetry-protected topological phases through the anomalous action of the symmetry on the edge, *Physical Review B* **90**, 10.1103/physrevb.90.235137 (2014).
- [35] K. Kawagoe and M. Levin, Anomalies in bosonic symmetry-protected topological edge theories: Connection to f-symbols and a method of calculation, *Physical Review B* **104**, 10.1103/physrevb.104.115156 (2021).
- [36] A. Kapustin and N. Sopenko, Anomalous symmetries of quantum spin chains and a generalization of the liebschultz-mattis theorem (2024), [arXiv:2401.02533 \[math-ph\]](#).
- [37] J. G. Rubio, A. Molnar, and Y. Ogata, Classifying symmetric and symmetry-broken spin chain phases with anomalous group actions (2024), [arXiv:2403.18573 \[quant-ph\]](#).
- [38] J. Garre-Rubio, L. Lootens, and A. Molnár, Classifying phases protected by matrix product operator symmetries using matrix product states, *Quantum* **7**, 927 (2023).
- [39] J. G. Rubio and N. Schuch, Fractional domain wall statistics in spin chains with anomalous symmetries (2024), [arXiv:2405.00439 \[cond-mat.str-el\]](#).
- [40] D. Perez-García, F. Verstraete, M. M. Wolf, and J. I. Cirac, Matrix product state representations, *Quantum Info. Comput.* **7**, 401–430 (2007).
- [41] D. Pérez-García, M. M. Wolf, M. Sanz, F. Verstraete, and J. I. Cirac, String order and symmetries in quantum spin lattices, *Phys. Rev. Lett.* **100**, 167202 (2008).
- [42] C. Xu, Three-dimensional symmetry-protected topological phase close to antiferromagnetic né el order, *Physical Review B* **87**, 10.1103/physrevb.87.144421 (2013).
- [43] Z. Bi, A. Rasmussen, K. Slagle, and C. Xu, Classification and description of bosonic symmetry protected topological phases with semiclassical nonlinear sigma models, *Physical Review B* **91**, 10.1103/physrevb.91.134404 (2015).
- [44] D. J. Williamson, N. Bultinck, and F. Verstraete, Symmetry-enriched topological order in tensor networks: Defects, gauging and anyon condensation (2017), [arXiv:1711.07982 \[quant-ph\]](#).
- [45] J. Garre-Rubio, L. Lootens, and A. Molnár, Classifying phases protected by matrix product operator symmetries using matrix product states, *Quantum* **7**, 927 (2023).
- [46] J. Haegeman and F. Verstraete, Diagonalizing transfer matrices and matrix product operators: A medley of exact and computational methods, *Annual Review of Condensed Matter Physics* **8**, 355 (2017).
- [47] K. Gawedzki, Topological actions in two-dimensional quantum field theories, *Nonperturbative Quantum Field Theory. Nato Science Series B*, Springer, New York, NY. **185**, 101 (1988).
- [48] K. Gomi and Y. Terashima, A fiber integration formula for the smooth Deligne cohomology, *International Mathematics Research Notices* **2000**, 699 (2000).
- [49] K. Gomi and Y. Terashima, Higher-dimensional parallel transports, *Math. Res. Lett.* **8** **2000**, 25 (2001).
- [50] A. L. Carey, S. Johnson, and M. K. Murray, Holonomy on D-branes, *Journal of Geometry and Physics* **52**, 186 (2004).
- [51] K. Gomi and Y. Terashima, Discrete torsion phases as topological actions, *Comm. Math. Phys.* **287**, 889 (2009).
- [52] L. Breen, [On the classification of 2-gerbes and 2-stacks](#), Astérisque No. 225 (Société mathématique de France, 1994).
- [53] L. Borsten, M. J. Farahani, B. Jurco, H. Kim, J. Narozny, D. Rist, C. Saemann, and M. Wolf, Higher gauge theory (2024), [arXiv:2401.05275 \[hep-th\]](#).