MAPS BETWEEN SPHERICAL GROUP RINGS

SHACHAR CARMELI, THOMAS NIKOLAUS, AND ALLEN YUAN

ABSTRACT. We prove that for finitely generated abelian groups A and B, the space of \mathbb{E}_{∞} -ring maps between the spherical groups rings $\mathbb{S}[A] \to \mathbb{S}[B]$ is equivalent to the discrete set of group homomorphisms $A \to B$. We also prove generalizations where the sphere is replaced by other ring spectra, e.g. we give a formula for the strict units in group rings of the form R[A] for A a finite *p*-group and R *p*-completely chromatically complete.

CONTENTS

| 1. | Introduction | 1 |
|------------|---|----|
| 2. | Rank 1 units in δ -rings | 5 |
| 3. | Tautological units, rigidity, and staticity | 19 |
| 4. | Rigidity for finite p -groups | 25 |
| 5. | Rigidity for free abelian groups | 30 |
| 6. | Strict units and maps of spherical group algebras | 38 |
| References | | 40 |

1. INTRODUCTION

This paper deals with a question about commutative ring spectra, also known as \mathbb{E}_{∞} -rings. In recent years, many concepts from the ordinary algebra of rings have been generalized to the higher algebra of ring spectra, leading to significant breakthroughs in many classical areas, such as algebraic *K*-theory [BHM93, HM03] and *p*-adic Hodge theory [BMS19]. However, some very basic questions in the area remain open — most notably, understanding maps between ring spectra is generally very hard. For an abelian group *A*, we consider the commutative ring spectrum

$$\mathbb{S}[A] := \Sigma^{\infty}_{+} A,$$

called the *spherical group ring of* A, which is a lift of the usual group ring $\mathbb{Z}[A]$ from the integers \mathbb{Z} to the sphere spectrum \mathbb{S} . The main question that we address in this paper is the question of which maps exist between these ring spectra. Our answer can be summarized as follows.

Theorem 1.1. The spherical group ring functor $A \mapsto S[A]$ from the category of finitely generated abelian groups to the ∞ -category of commutative ring spectra is fully faithful.

Date: May 13, 2024.

In other words: for finitely generated abelian groups A and B, the space of \mathbb{E}_{∞} -ring maps $\mathbb{S}[A] \to \mathbb{S}[B]$ is discrete and in bijection with the set of group homomorphisms $A \to B$.

This result is very surprising, at least to the authors. Indeed, the analogous result in ordinary algebra is completely false: there are many maps $\mathbb{Z}[A] \to \mathbb{Z}[B]$ which do not arise from group homomorphisms $A \to B$. For instance, when $A = \mathbb{Z}$, such maps are simply units in $\mathbb{Z}[B]$, and the question of units in group rings has been extensively studied [Hig40, Kar83, JM96, Gar21]. In addition to the unit -1, which does not come from an element of B, we have for instance the *Bass cyclic units*, see, e.g., [JM96, §2].

Theorem 1.1 asserts that none of these units lift to S, implying that much of the number theory underlying these units, such as the theory of cyclotomic fields and cyclotomic units, does not extend to the sphere. The theorem can also be seen as a strong rigidity result about group rings over S. In fact, it says more — the discreteness of the mapping spaces asserts that the maps that do lift, lift in a unique way. More precisely, the fibers of the map of spaces

(1)
$$\operatorname{Map}_{\operatorname{CAlg}(\operatorname{Sp})}(\mathbb{S}[A], \mathbb{S}[B]) \to \operatorname{Map}_{\operatorname{Ring}}(\mathbb{Z}[A], \mathbb{Z}[B])$$

given by base-change along $\mathbb{S} \to \mathbb{Z}$ (or equivalently by taking π_0) are empty or contractible.

1.1. Background and motivation. There is a slick way to see that the image of (1) consists only of maps induced by group homomorphisms $A \rightarrow B$. The idea is that such maps must be compatible with certain natural operations acting on the homotopy groups of commutative ring spectra.

Namely, suppose R is an ordinary ring and we have a lift \mathbb{S}_R to a commutative ring spectrum such that $\mathbb{S}_R \otimes_{\mathbb{S}} \mathbb{Z} \simeq R$ (i.e., a Moore spectrum)¹. Then, the Tate-valued Frobenius map of [NS18] naturally endows R_p^{\wedge} with a (p-)Frobenius lift for every prime p. In fact, R acquires a natural δ_p -ring structure on each of its p-completions – together, we refer to this as a $\hat{\delta}$ -ring structure.²

In the special case of group rings $\mathbb{Z}[A]$, the $\hat{\delta}$ -ring structure arising from the spherical lift $\mathbb{S}[A]$ is simple: the corresponding Frobenius lifts are induced from the group homomorphisms $A \xrightarrow{p} A$ for every prime p. While there can be many interesting ring homomorphisms $\mathbb{Z}[A] \to \mathbb{Z}[B]$, one can show by elementary algebra that the only ones compatible with this $\hat{\delta}$ -ring structure are the ones induced by group homomorphisms $A \to B$.

Summarizing this discussion, the map (1) naturally lands in morphisms of $\hat{\delta}$ -rings and our main result is equivalent to the assertion that the resulting map

 $\operatorname{Map}_{\operatorname{CAlg}(\operatorname{Sp})}(\mathbb{S}[A], \mathbb{S}[B]) \to \operatorname{Map}_{\hat{\delta}\operatorname{-Ring}}(\mathbb{Z}[A], \mathbb{Z}[B])$

is an equivalence. We view this result as answering an important special case of the following general question:

Question 1. For which $\hat{\delta}$ -rings R does there exist an \mathbb{E}_{∞} -lift to the sphere \mathbb{S}_R , and for which pairs of lifts $\mathbb{S}_R, \mathbb{S}_{R'}$ is the induced map

(2)
$$\operatorname{Map}_{\operatorname{CAlg}(\operatorname{Sp})}(\mathbb{S}_R, \mathbb{S}_{R'}) \to \operatorname{Map}_{\hat{\delta}\operatorname{-Ring}}(R, R')$$

an equivalence?

¹Note that the underlying spectrum of \mathbb{S}_R is determined by R, so this is really about the \mathbb{E}_{∞} -structure. ²We warn the reader that this is non-standard terminology and there are some subtleties about derived completion

hidden.

Example 1.2. If R is a Q-algebra, then each p-completion of R vanishes and thus a δ -ring structure is no additional data. Moreover, we have $\mathbb{S}_R = R$, so (2) is tautologically an equivalence.

On the other hand, a *p*-complete δ -ring is just a δ_p -ring, as all other completions vanish. In this case, Question 1 is understood to have a positive answer in formally étale situations, such as the spherical Witt vectors $\mathbb{S}_{W(\kappa)}$ of $R = W(\kappa)$ by [Lur18, Example 5.2.7]. A variant of our main theorem extends this to the case of *p*-complete group rings:

Theorem 1.3 (Corollary 6.2). Suppose that S, S' are \mathbb{E}_{∞} -rings of the form $\mathbb{S}_{W(\kappa)}[M]_p^{\wedge}$ for some perfect ring κ of characteristic p and some finitely generated abelian group M. Then the natural map

 $\pi_0 : \operatorname{Map}_{\operatorname{CAlg}(\operatorname{Sp})}(S, S') \to \operatorname{Map}_{\delta_p - \operatorname{Ring}}(\pi_0(S), \pi_0(S'))$

is an equivalence (cf. Construction 2.3 for the δ_p -structures in the target).

We point out that unlike in the integral case, the equivalent mapping spaces in Theorem 1.3 are *not* in general equivalent to the corresponding set of group homomorphisms, unless the groups involved have no prime to p torsion (Theorem 2.30).

We believe the above results to be the first systematic computations of mapping spaces between such flat S-algebras that are not (formally) étale extensions of the sphere spectrum or completions thereof. Yet, some fundamental cases of Question 1 remain open: for instance, one can study spherical *monoid* rings, rather than group rings. In fact, to the knowledge of the authors, even the case of polynomial rings is open in all nontrivial cases:

Question 2. What is the space of maps $\operatorname{Map}_{\operatorname{CAlg}(\operatorname{Sp})}(\mathbb{S}[\mathbb{N}^i], \mathbb{S}[\mathbb{N}^j])$ for any $i, j \geq 1$?

Example 1.4. The map (2) is not an equivalence in general. For example, if $\mathbb{S}_R = \mathbb{S} \oplus \mathbb{S}$ is a square zero extension of \mathbb{S} by itself, then the space of \mathbb{E}_{∞} -maps $\mathbb{S}_R \to \mathbb{S}_R$ contains $\Omega^{\infty}\mathbb{S}$ is a retract and hence it is not discrete.

As for the first part of Question 1, the lifting question is equivalent to the question of highly structured multiplications on Moore spectra and has been extensively studied [Oka84, Bur22, Bha22], though we do not know many interesting examples of spherical lifts in the \mathbb{E}_{∞} case beyond the spherical Witt vectors of [Lur18, Example 5.2.7]. For instance, we believe the following is open:

Question 3. Does the free δ_p -ring on one generator lift to \mathbb{S} ?

Our interest in studying Question 1 arose in work of the second author joint with B. Antieau and A. Krause on prismatic cohomology: for a given δ_p -ring R and any R-algebra S, there is prismatic cohomology $\Delta_{S/R} \in \mathcal{D}(\mathbb{Z})$, an important invariant in p-adic geometry generalizing crystalline cohomology to mixed characteristic [BMS19, BS22, AKN23]. The development of this theory was inspired by topological periodic homology and one can generally ask if $\Delta_{S/R}$ admits a topological refinement, i.e. if it is the associated graded of a filtration on topological periodic homology of a cyclotomic spectrum³. It will be shown in forthcoming work of the second author with Antieau and Krause that this is indeed the case if R admits a lift \mathbb{S}_R to the sphere with the cyclotomic spectrum in question given by THH(S/\mathbb{S}_R). The cyclotomic spectrum THH(S/\mathbb{S}_R) is clearly functorial in the lift \mathbb{S}_R . This relates Question 1 to a question of P. Scholze about which prisms can be realized by cyclotomic spectra and how functorially this can be done [Sch].

 $^{^{3}}$ We are suppressing Breuil-Kisin twists, Frobenius twists and Nygaard completions for simplicity.

1.2. Outline and further results. The core content of the paper is Theorem 1.3 in the case $S = \mathbb{S}[\mathbb{Z}]_p^{\wedge}$ and $S' = \mathbb{S}[M]_p^{\wedge}$, i.e. proving the equivalence

$$\operatorname{Map}_{\operatorname{CAlg}(\operatorname{Sp})}(\mathbb{S}[\mathbb{Z}]_p^{\wedge}, \mathbb{S}[M]_p^{\wedge}) \xrightarrow{\sim} \operatorname{Map}_{\delta_p - \operatorname{Ring}}(\mathbb{Z}[\mathbb{Z}]_p^{\wedge}, \mathbb{Z}[M]_p^{\wedge}).$$

Our approach is to compute both sides. The right-hand side is purely algebraic, and we start by completely understanding it in the self-contained §2. The mapping space on the left-hand side is known as the *strict units* of $\mathbb{S}[M]_p^{\wedge}$, denoted by $\mathbb{G}_m(\mathbb{S}[M]_p^{\wedge})$.

Remark 1.5. Strict units of commutative ring spectra are of significant interest in homotopy theory [Fun20], as they are closely related to questions of adjoining roots to ring spectra [Law20] and multiplicative twists of cohomology theories by ordinary cohomology classes [MS06, ABG18, SW15]. Despite this, the difficulty of computing maps between commutative ring spectra has allowed for only a few computations so far — notably the cases of Lubin–Tate theories (unpublished work of Rezk and Hopkins–Lurie, cf. also [BSY22]) and the sphere and its completions [Car22].

Using the augmentation of $\mathbb{S}[M]_p^{\wedge}$, we obtain a splitting

$$\mathbb{G}_m(\mathbb{S}[M]_p^{\wedge}) \simeq \mathbb{G}_m^{\mathrm{red}}(\mathbb{S}[M]_p^{\wedge}) \oplus \mathbb{G}_m(\mathbb{S}_p)$$

into $\mathbb{G}_m(\mathbb{S}_p)$, which has been computed in [Car22], and what we call the *reduced units* $\mathbb{G}_m^{\mathrm{red}}(\mathbb{S}[M]_p^{\wedge})$. We study these via the "obvious" map $\theta: M \to \mathbb{G}_m^{\mathrm{red}}(\mathbb{S}[M])$ given informally by $m \mapsto [m]$. A key idea is that θ is a special case of a map

$$\Theta_R : \underline{M}(R) \to \mathbb{G}_m^{\mathrm{red}}(R[M])$$

which is natural in the commutative ring spectrum R and, critically, often an equivalence.⁴ This property, which we call *rigidity* ("R is M-*rigid*"), is an organizing principle in this paper, as we can analyze its closure under changing R and M. After compiling some basic techniques for proving rigidity in §3, the remainder of the paper proceeds as follows:

§4: When R is *p*-complete and M is a finite abelian *p*-group. We show:

Theorem 1.6. Assume that R is p-complete, R/p is chromatically complete, and M is a finite abelian p-group. Then R is M-rigid.

In fact, we prove a more general form of this theorem as Corollary 4.13. We start with the case where R is instead a T(n)-local ring spectrum that admits a primitive higher root of unity in the sense of [CSY21] (e.g., any algebra over a height n Lubin–Tate theory). Here, we can verify Theorem 1.6 directly by computation using the chromatic Fourier equivalence [BCSY22, HL13a]

$$R[M] \simeq R^{B^n M}$$

Then, closure under limits allows us to bootstrap to the general setting.

§5: Extending to torsion free $M = \mathbb{Z}^m$. Here, we specialize to $R = \mathbb{S}_{W(\kappa)}$, the spherical Witt vectors of a perfect ring κ (though intermediate results are proven in greater generality). We show that the short exact sequence $M \xrightarrow{p} M \to M/p$ induces a cofiber sequence of spectra

(3)
$$\mathbb{G}_m^{\mathrm{red}}(\mathbb{S}_{W(\kappa)}[M]_p^{\wedge}) \to \mathbb{G}_m^{\mathrm{red}}(\mathbb{S}_{W(\kappa)}[M]_p^{\wedge}) \to \mathbb{G}_m^{\mathrm{red}}(\mathbb{S}_{W(\kappa)}[M/p]_p^{\wedge})$$

It turns out that the first map can be identified with the map induced by the Tate-valued Frobenius, which for general reasons induces multiplication by p on strict units. This allows us to bootstrap our knowledge of $\mathbb{G}_m^{\mathrm{red}}(\mathbb{S}_{W(\kappa)}[M/p]_p^{\wedge})$ from Theorem 1.6 to obtain:

⁴Here, <u>M(R)</u> denotes locally constant functions $\operatorname{Spec}(\pi_0(R)) \to M$.

Theorem 1.7. Let κ be a perfect \mathbb{F}_p -algebra and M be a finitely generated abelian group. Then $\mathbb{S}_{W(\kappa)}$ is M-rigid.

Combined with Theorem 1.6, we obtain an analogous result when M is additionally allowed to have p-power torsion part (Corollary 5.19).

§6: Assembling the pieces. We first add in the prime to p torsion in M, in particular proving Theorem 1.3. Then, we use an arithmetic fracture square argument to combine the p-complete results of the previous sections to prove our main theorem about R = S.

1.3. Acknowledgements. The authors would like to thank Zhouhang Mao, Tomer Schlank, and the entire Copenhagen homotopy theory seminar group for useful discussions related to this material, as well as Achim Krause for sharing his ideas related to §2.4. We would also like to thank Edith Hübner and Robert Szafarczyk for helpful discussions and suggestions concerning Section 2. The first author was partially supported by the Danish National Research Foundation through the Copenhagen Centre for Geometry and Topology (DNRF151). The second author was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project-ID 427320536 - SFB 1442, as well as under Germany's Excellence Strategy EXC 2044 390685587, Mathematics Münster: Dynamics-Geometry-Structure. The third author was supported in part by NSF grant DMS-2002029.

2. Rank 1 units in δ -rings

In this section, we consider the algebraic analogue of the main results of this paper. We start by reviewing the relevant structures to set notation.

Definition 2.1. For a prime p, a δ_p -ring⁵ is a commutative ring R together with an operation $\delta_p : R \to R$ satisfying the identities:

$$\delta_p(0) = \delta_p(1) = 0,$$

$$\delta_p(x+y) = \delta_p(x) + \delta_p(y) + \frac{1}{p}(x^p + y^p - (x+y)^p),$$

$$\delta_p(xy) = x^p \delta_p(y) + y^p \delta_p(x) + p \delta_p(x) \delta_p(y).$$

We denote the category of these by δ_p -Ring.

The identities are rigged so that the operation $\psi(x) = x^p + p\delta_p(x)$ is a Frobenius lift. In fact, a δ_p -structure on a torsion-free ring is simply a Frobenius lift. This becomes even cleaner in the animated setting.

Definition 2.2. Let ACR denote the category of animated commutative rings (cf. e.g. [CS19, $\S1.2$]). Then an animated δ_p -ring is an animated commutative ring R together with a lift

$$\begin{array}{ccc} R & & & R \\ & & & \downarrow \\ R /\!\!/ p & & & \varphi_{R /\!\!/ p} \\ \end{array} \end{array} \xrightarrow{\varphi_{R /\!\!/ p}} R /\!\!/ p.$$

Here, $R/\!\!/p$ denotes the derived reduction mod p, and $\varphi_{R/\!\!/p}$ denotes the animated Frobenius. We let ACR^{δ_p} denote the category of animated δ_p -rings.

⁵These are often simply called δ -rings, but we add the p to the notation to allow for different primes.

This definition is compatible with Definition 2.1 in the sense that the objects in ACR^{δ_p} with discrete underlying ring are exactly δ_p -rings, and ACR^{δ_p} is the animation of δ_p -Ring [Hol23, Thm. 2.4.4]. We will be particularly interested in the following δ_p -rings:

Construction 2.3. Let M be an abelian group. Since the group ring $\mathbb{Z}[M]$ is torsion free, a δ_p -ring structure on it is the same as a lift of Frobenius. Accordingly, we equip the group rings $\mathbb{Z}[M]$ with the δ_p -structures corresponding to the lift of Frobenius

$$\varphi(\sum_{m \in M} a_m \cdot [m]) = \sum_{m \in M} a_m \cdot [pm].$$

More generally, for a δ_p -ring R, we endow $R[M] = R \otimes \mathbb{Z}[M]$ with the tensor product δ_p -ring structure, i.e. $\varphi_{R[M]} = \varphi_R \otimes \varphi_{\mathbb{Z}[M]}$.

The analogue of strict units in this setting are the following:

Definition 2.4. A rank 1 unit in a δ_p -ring A is a morphism of δ_p -rings

$$\mathbb{Z}[\mathbb{Z}] =: \mathbb{Z}[t^{\pm 1}] \to A$$

We denote by $\mathbb{G}_m^{\delta_p}(A)$ the set of rank 1 units, which can be described as the units $t \in \mathbb{G}_m(A)$ such that $\delta_p(t) = 0$. In fact, the identities for δ_p immediately imply that $\mathbb{G}_m^{\delta_p}(A)$ is a subgroup of $\mathbb{G}_m(A)$.

Remark 2.5. The functor $A \mapsto \mathbb{G}_m^{\delta_p}(A)$ from δ_p -rings to abelian groups is right adjoint to the group algebra functor of Construction 2.3.

Example 2.6 (Witt vectors, cf. e.g. [Bha]). For any perfect ring κ , the ring of *p*-typical Witt vectors $W(\kappa)$ admits a unique δ_p -structure, with Frobenius lift induced by the Frobenius map $\kappa \to \kappa$. In fact, these are precisely the perfect δ_p -rings, i.e. those with invertible lift of Frobenius. In this case, the multiplicative lift and the projection $W(\kappa) \to \kappa$ induce isomorphisms

$$\kappa^{\times} \xrightarrow{\sim} \mathbb{G}_m^{\delta_p}(W(\kappa)) \xrightarrow{\sim} \kappa^{\times}$$

Our first goal in this section is to compute the rank 1 units of group algebras. The techniques are quite different from those in the S-linear situation of the later sections – they rely on the deformation theory of the functor $\mathbb{G}_m^{\delta_p}(-)$, which we study in §2.1. Applying this in §2.2, we show:

Theorem 2.7. Let A be a p-complete δ_p -ring and M be a finitely generated abelian group whose torsion part is p-power torsion. Then there is a natural isomorphism

$$\mathbb{G}_m^{\delta_p}(A[M]_p^{\wedge}) \cong \mathbb{G}_m^{\delta_p}(A) \oplus \underline{M}(A).$$

Here, $\underline{M}(A)$ denotes the abelian group of locally constant M-valued functions on $\operatorname{Spec}(A)$.

We finish with some material connecting this story to the remainder of the paper. In §2.3, we globalize these results (across primes p) using the notion of a $\hat{\delta}$ -ring, which is essentially a ring with a δ_p -structure on its p-completion for each p. Finally, in §2.4, we relate this discussion back to the topological story by showing that if R is a commutative ring spectrum such that $R \otimes \mathbb{Z}$ is discrete (i.e., a *Moore spectrum*), then $\pi_0(R)$ acquires a natural $\hat{\delta}$ -ring structure.

2.1. Deformation theory of rank 1 units. We study rank 1 units of group algebras by deformation theory, starting in §2.1.1 with the case of a square-zero extension of δ_p -rings. Here, by explicit analysis, we obtain an exact sequence

(4)
$$0 \to I^{\delta_p = 1} \to \mathbb{G}_m^{\delta_p}(A) \to \mathbb{G}_m^{\delta_p}(A/I).$$

The key ingredient to going further is the observation that units of rank 1 satisfy a certain "square nil-invariance." That is, under appropriate completeness conditions on an ideal $I \subset A$, we prove in §2.1.2 an isomorphism $\mathbb{G}_m^{\delta_p}(A) \cong \mathbb{G}_m^{\delta_p}(A/I^2)$. This gives us (4) more generally.

The final ingredient we need is to identify the fiber term in the case where A is a group algebra. It turns out that this can be done with a form of Artin–Schreier theory for δ_p -rings, which we discuss in §2.1.3.

2.1.1. δ_p -ideals and square-zero extensions.

Definition 2.8. We say an ideal I in a δ_p -ring A is a δ_p -ideal if $\delta_p(I) \subseteq I$.

 δ_p -ideals are particularly easy to analyze in the case where I is square-zero, that is, $I^2 = 0$.

Proposition 2.9. Let A be a δ_p -ring and let $I \subset A$ be a square zero δ_p -ideal. Then:

(1) The map $\delta_p: I \to I$ is a φ -semilinear morphism of A-modules. Namely, it is a group homomorphism and satisfies

$$\delta_p(xa) = \varphi(x)\delta_p(a), \quad \forall x \in A, a \in I$$

(2) For $x \in A$ and $a \in I$ we have

$$\delta_p(x+a) = \delta_p(x) + \delta_p(a) - x^{p-1}a.$$

Proof. Both statements follow immediately from the additive and multiplicative identities of δ_p , using the fact that I is a square zero ideal. Specifically:

$$\delta_p(a+b) = \delta_p(a) + \delta_p(b) + \frac{a^p + b^p - (a+b)^p}{p} = \delta_p(a) + \delta_p(b) \quad \forall a, b \in I,$$

$$\delta_p(xa) = x^p \delta_p(a) + a^p \delta_p(x) + p \delta_p(x) \delta_p(a) = (x^p + p \delta_p(x)) \delta_p(a) = \varphi(x) \delta_p(a) \quad \forall x \in A, a \in I,$$

and

$$\delta_p(x+a) = \delta_p(x) + \delta_p(a) + \frac{x^p + a^p - (x+a)^p}{p} = \delta_p(x) + \delta_p(a) - x^{p-1}a \quad \forall x \in A, a \in I.$$

This gives us the following statement about how $\mathbb{G}_m^{\delta_p}(-)$ changes in square-zero extensions.

Proposition 2.10. Let A be a δ_p -ring and let I be a square-zero δ_p -ideal in A. Then, we have an exact sequence

$$0 \longrightarrow I^{\delta_p = 1} \longrightarrow \mathbb{G}_m^{\delta_p}(A) \longrightarrow \mathbb{G}_m^{\delta_p}(A/I)$$

Proof. Let $1 + x \in \mathbb{G}_m^{\delta_p}(A)$ be an element of $\ker(\mathbb{G}_m^{\delta_p}(A) \to \mathbb{G}_m^{\delta_p}(A/I))$, where $x \in I$. Then we have (using Proposition 2.9(2))

$$0 = \delta_p(1+x) = \delta_p(1) + \delta_p(x) - x.$$

Noting that $\delta_p(1) = 0$, we obtain the conclusion.

Remark 2.11. As we analyze the behavior of $\mathbb{G}_m^{\delta_p}(-)$ along deformations, we focus on the case when the relevant ideals are closed in the *p*-adic topology. However, even if $I \subseteq A$ is closed, its powers I^n may not be, so we let $\overline{I^n}$ denote their closures in the *p*-adic topology. (In any case, for our applications, this distinction will not be important as our ideals will be finitely generated.)

The following lemma ensures that these operations play well with the δ_p -structure:

Lemma 2.12. Let $I \subset A$ be a δ_p -ideal. Then:

- (1) For $n \ge 0$, I^n is a δ_p -ideal.
- (2) The ideal $\overline{I} \subset A$ is a δ_p -ideal.

Proof. The first statement is immediate from the addition and multiplication formulas for δ_p , and the latter follows from the fact that δ_p is *p*-adically continuous.

2.1.2. Square nil-invariance. While the functor $\mathbb{G}_m^{\delta_p}(-)$ is not nil-invariant in general, we will show here that it is invariant under quotient by the square of an ideal with respect to which a δ_p -ring is complete. The key input is the following lemma:

Lemma 2.13. Let A be a δ_p -ring and $I \subset A$ be a δ_p -ideal. Then, for $m \geq 2$, the images of the induced maps

$$\begin{split} \delta_p &: I^m / I^{m+1} \to I^m / I^{m+1} \\ \delta_p &: \overline{I^m} / \overline{I^{m+1}} \to \overline{I^m} / \overline{I^{m+1}} \end{split}$$

are contained in $p \cdot I^m / I^{m+1}$ and $p \cdot \overline{I^m} / \overline{I^{m+1}}$ respectively.

Proof. The second statement is immediate from the first because δ_p is *p*-adically continuous. To see the first statement, we note that by the additivity statement of Proposition 2.9, it suffices to consider elements of the form $ab \in I^m$ for $a \in I$ and $b \in I^{m-1}$. Then we have

$$\begin{split} \delta_p(ab) &= a^p \delta_p(b) + b^p \delta_p(a) + p \delta_p(a) \delta_p(b) \\ &\equiv p \delta_p(a) \delta_p(b) \pmod{I^{m+1}}. \end{split}$$

Given this, we have our main statement:

Proposition 2.14. Let A be a δ_p -ring and let I be a δ_p -ideal such that A is classically (I, p)complete. Then, the map

$$A \to A/\overline{I^2}$$

induces an isomorphism

$$\mathbb{G}_m^{\delta_p}(A) \xrightarrow{\sim} \mathbb{G}_m^{\delta_p}(A/\overline{I^2}).$$

Proof. Our assumption that A is (I, p)-complete implies that

$$A \simeq \lim A / \overline{I^m}$$

To see this, note that the map from A to the limit is clearly surjective by assumption, and so it is enough to check that $\cap \overline{I^m} = \{0\}$. But if $x \in \cap \overline{I^m}$, then for all $j \in \mathbb{N}$, $x \in \cap I^m$ modulo p^j , and hence $x \equiv 0 \mod p^j$ since A/p^j is *I*-complete. Since A is *p*-complete this implies that x = 0. It therefore suffices to show that the induced map

$$\mathbb{G}_m^{\delta_p}(A/\overline{I^{m+1}}) \to \mathbb{G}_m^{\delta_p}(A/\overline{I^m})$$

is an isomorphism when $m \geq 2$. For this, let $x \in \mathbb{G}_m^{\delta_p}(A/\overline{I^m})$ and let \tilde{x} be a lift of x to $A/\overline{I^{m+1}}$. We will show there is a unique $a \in \overline{I^m}/\overline{I^{m+1}}$ such that $\tilde{x} + a \in \mathbb{G}_m^{\delta_p}(A/\overline{I^{m+1}})$. As A is I-complete, $\tilde{x} + a$ is certainly a unit, so $\tilde{x} + a \in \mathbb{G}_m^{\delta_p}(A/\overline{I^{m+1}})$ if and only if $\delta_p(\tilde{x} + a) = 0$. Since the ideal $\overline{I^m}/\overline{I^{m+1}} \subset A/\overline{I^{m+1}}$ is square-zero, we have by Proposition 2.9(2)

$$\delta_p(\tilde{x}+a) = \delta_p(\tilde{x}) + \delta_p(a) - \tilde{x}^{p-1}a$$

Since $x \in \mathbb{G}_m^{\delta_p}(A/\overline{I^m})$, the first term is in $\overline{I^m}/\overline{I^{m+1}}$. It therefore suffices to show that the function

$$\frac{\overline{I^m}}{\overline{I^m+1}} \to \overline{I^m}/\overline{I^{m+1}}$$
$$a \mapsto \delta_p(a) - \tilde{x}^{p-1}a$$

is a bijection. But \tilde{x} is a unit, so it's enough to show that $\operatorname{id} - \frac{1}{\tilde{x}^{p-1}}\delta_p$ is an isomorphism. But this has inverse given by

$$b \mapsto \sum_{j=0}^{\infty} \left(\frac{1}{\tilde{x}^{p-1}} \delta_p\right)^{(j)}(b),$$

which is a well-defined element of $\overline{I^m}/\overline{I^{m+1}}$ because the *j*-th term of this series is divisible by p^j by Lemma 2.13.

Combining this with Proposition 2.14, we have the following extension of Proposition 2.10:

Proposition 2.15. Let A be a δ_p -ring and let I be a δ_p -ideal in A such that A is classically (I, p)complete. Then $A \to A/\overline{I^2}$ induces an isomorphism on $\mathbb{G}_m^{\delta_p}(-)$ and we obtain an exact sequence

$$0 \longrightarrow (\overline{I}/\overline{I^2})^{\delta_p=1} \longrightarrow \mathbb{G}_m^{\delta_p}(A) \longrightarrow \mathbb{G}_m^{\delta_p}(A/\overline{I}) .$$

Here, the left term is the kernel of the homomorphism $\delta_p - \mathrm{id} \colon \overline{I}/\overline{I^2} \to \overline{I}/\overline{I^2}$.

Remark 2.16. In the situation of Proposition 2.15, the map $\mathbb{G}_m^{\delta_p}(A) \to \mathbb{G}_m^{\delta_p}(A/\overline{I})$ is not generally surjective. However, it will be in our cases of interest, as we take $A \to A/\overline{I}$ to be the augmentation of an augmented algebra (the group ring), which has a section.

2.1.3. Artin–Schreier theory. Another general result regarding δ_p -rings that we need is Artin–Schreier theory. In this sub-section, for an animated commutative ring $R \in ACR$, we regard the derived scheme Spec(R) as endowed with the étale topology on its small étale site (see, e.g., $[\text{CS19}, \S5.2.3]$). We refer to sheaves on Spec(R) simply as étale sheaves, and unless stated otherwise consider spectral coefficients. For an abelian group A, we denote by $\underline{A}^{\text{et}}$ the constant étale sheaf on Spec(R).

Definition 2.17. Let $R \in ACR_{\mathbb{Z}/p^r}$ and let $\overline{R} := R \otimes_{\mathbb{Z}/p^r} \mathbb{Z}/p$. A *lift of Frobenius* for R is a map $\varphi_R \colon R \to R$ which lifts the derived Frobenius map $\varphi_{\overline{R}} \colon \overline{R} \to \overline{R}$ along the reduction map $R \twoheadrightarrow \overline{R}$.

More precisely, it is the data of a commutative square of the form:

$$\begin{array}{c} R & \stackrel{\varphi}{\longrightarrow} & R \\ \downarrow & & \downarrow \\ R & \stackrel{\overline{\varphi}}{\longrightarrow} & \overline{R} \end{array}$$


Our main example for lifts of Frobenius is the following.

Example 2.18. Let A be a δ_p -ring, and $\varphi \colon A \to A$ the corresponding Frobenius map. Then, by definition, the map $A/\!\!/ p^r \to A/\!\!/ p^r$ induced from φ by reducing mod p^r extends canonically to a lift of Frobenius. Moreover, if A is derived p-complete, we may view the δ_p -ring structure on A as a compatible choice of such lifts of Frobenius.

It turns out that a lift of Frobenius for R automatically extends to all étale R-algebras.

Lemma 2.19. Let R be an animated commutative \mathbb{Z}/p^r -algebra endowed with a lift of Frobenius $\varphi_R \colon R \to R$. Then, every étale R-algebra S admits a unique lift of Frobenius $\varphi_S \colon S \to S$ compatible with φ_R .

Proof. Let $\overline{S} := S \otimes_{\mathbb{Z}/p^r} \mathbb{Z}/p$, so that we have a reduction mod p map $S \to \overline{S}$. A compatible lift of Frobenius for S is a solution to the lifting problem



Since S is an animated \mathbb{Z}/p^r -algebra as well, the right vertical map is a nil-thickening, and the uniqueness of the lift now follows from the lifting property of étale morphisms of animated commutative rings (i.e. by the vanishing of the cotangent complex $L_{S/B}^{\text{alg}}$).

Corollary 2.20. Let R be an animated commutative \mathbb{Z}/p^r -algebra with a lift of Frobenius $\varphi_R \colon R \to R$. Then, the map φ_R extends uniquely to an endomorphism of sheaves of \mathbb{Z}/p^r -algebras $\varphi \colon \mathcal{O}_R \to \mathcal{O}_R$ on the small étale site of $\operatorname{Spec}(R)$. Here, \mathcal{O}_R denotes the structure sheaf of $\operatorname{Spec}(R)$.

Proof. The data of a morphism $\varphi \colon \mathcal{O}_R \to \mathcal{O}_R$ lifting φ_R is the same as that of a map of sheaves of animated commutative *R*-algebras $\varphi_R^* \mathcal{O}_R \to \mathcal{O}_R$. Note that both the source and target consist of sheaves of étale algebras over *R*.

Since the ∞ -category of étale *R*-algebras is equivalent to the 1-category of étale $\pi_0 R$ -algebras (see, e.g., [CS19, Proposition 5.2.4]), it follows that a natural transformation $\varphi_R^* \mathcal{O}_R \to \mathcal{O}_R$ is determined by its restriction to objects and morphisms in the category of étale *R*-algebras. Individual lifts at the objects exist uniquely by Lemma 2.19, and they are compatible with morphisms immediately by the uniqueness.

Since φ_R is a morphism of sheaves of \mathbb{Z}/p^r -algebras, we have a canonical null sequence of sheaves

$$(\mathbb{Z}/p^r)^{\mathrm{et}} \longrightarrow \mathcal{O}_R \xrightarrow{\varphi - 1} \mathcal{O}_R,$$

where $(\mathbb{Z}/p^r)^{\text{et}}$ is the constant étale sheaf on the value \mathbb{Z}/p^r .

Lemma 2.21. Let R be an animated commutative \mathbb{Z}/p^r -algebra with a lift of Frobenius $\varphi_R \colon R \to R$. For all i > 0, the map φ_R induces a nilpotent endomorphism of $\pi_i R$. In particular, $\varphi_R - 1$ is an isomorphism on $\pi_i R$.

Proof. Let $\overline{R} := R \otimes_{\mathbb{Z}/p^r} \mathbb{Z}/p$. Tensoring the *p*-adic filtration of \mathbb{Z}/p^r with *R*, we obtain a natural finite filtration R_{\bullet} with associated graded

$$\operatorname{gr}(R_{\bullet}) \simeq R \otimes_{\mathbb{Z}/p^r} \operatorname{gr}((\mathbb{Z}/p^r)_{\bullet}) \simeq \overline{R}[t]/t^r.$$

Since the filtration is finite and natural in R, it is stable under φ_R and it suffices to show that φ_R is nilpotent on $\pi_i \operatorname{gr}(R_{\bullet})$; in fact we will show it is zero. Indeed, we have

$$\varphi_R(at^j) = \varphi_{\overline{R}}(a)t^j \quad \forall a \in \pi_i \overline{R},$$

which is 0 because $\varphi_{\overline{R}}$ vanishes in positive degrees (cf. [BS17, Prop 11.6, Rem 11.8]).

Corollary 2.22. Let R be an animated commutative \mathbb{Z}/p^r -algebra with a lift of Frobenius $\varphi_R \colon R \to R$. Then, the square

is a Cartesian square of étale sheaves on $\operatorname{Spec}(R)$.

Proof. It suffices to prove that the square becomes Cartesian after evaluation at Spec(S) for an étale *R*-algebra *S*, namely that

is Cartesian. The fiber of both vertical maps is $\tau_{\geq 1}S$, and $\varphi_S - 1$ is an isomorphism on this fiber by Lemma 2.21. This implies the result.

We are ready to state and prove a version of Artin-Schreier theory for Frobenius lifts.

Proposition 2.23 (Artin-Schreier for Frobenius Lifts). Let $R \in ACR_{\mathbb{Z}/p^r}$ with a lift of Frobenius $\varphi_R \colon R \to R$, and let $\varphi \colon \mathcal{O}_R \to \mathcal{O}_R$ be its unique extension from Lemma 2.19. Then, the null sequence

$$(\underline{\mathbb{Z}/p^r})^{\mathrm{et}} \longrightarrow \mathcal{O}_R \xrightarrow{\varphi - 1} \mathcal{O}_R$$

is a fiber sequence of sheaves of \mathbb{Z}/p^r -module spectra on $\operatorname{Spec}(R)$, and it induces a short exact sequence

$$0 \longrightarrow (\underline{\mathbb{Z}/p^r})^{\text{et}} \longrightarrow \pi_0 \mathcal{O}_R \xrightarrow{\varphi - 1} \pi_0 \mathcal{O}_R \longrightarrow 0$$

of sheaves of abelian groups on $\operatorname{Spec}(R)$.

Proof. First, observe that the first sequence being a fiber sequence and the second sequence being exact are equivalent, by Corollary 2.22. Now, the claim for r = 1 follows from classical Artin-Schreier theory, which says that for a (classical) \mathbb{Z}/p -scheme X the sequence

$$0 \longrightarrow (\underline{\mathbb{Z}/p})^{\mathrm{et}} \longrightarrow \mathcal{O}_X \xrightarrow{\varphi_X - 1} \mathcal{O}_X \longrightarrow 0$$

is a short exact sequence of étale sheaves. For the general case, let $R \in \operatorname{ACR}_{\mathbb{Z}/p^r}$ with lift of Frobenius φ_R , and set $\overline{R} := R \otimes_{\mathbb{Z}/p^r} \mathbb{Z}/p$. Then, via the equivalence of étale sites $\operatorname{Spec}(R) \simeq \operatorname{Spec}(\overline{R})$, the base-change of the null sequence

$$(\underline{\mathbb{Z}/p^r})^{\mathrm{et}} \longrightarrow \mathcal{O}_R \xrightarrow{\varphi - 1} \mathcal{O}_R$$

along the map $\mathbb{Z}/p^r \to \mathbb{Z}/p$ identifies with the sequence

$$(\underline{\mathbb{Z}/p})^{\mathrm{et}} \longrightarrow \mathcal{O}_{\overline{R}} \xrightarrow{\varphi_{\overline{R}}-1} \mathcal{O}_{\overline{R}}$$

Since the latter is a fiber sequence by the case r = 1 and the functor $\mathbb{Z}/p \otimes_{\mathbb{Z}/p^r} (-)$ is exact and conservative, we deduce that the former sequence is a fiber sequence and the result follows. \Box

Corollary 2.24. Let A be a derived p-complete δ_p -ring. For every $r \in \mathbb{N}$ we have an exact sequence

$$0 \longrightarrow \underline{\mathbb{Z}/p^r}(A) \to A/p^r \xrightarrow{\varphi_A - 1} A/p^r.$$

Proof. Let $R = A/\!\!/ p^r$, which is an animated \mathbb{Z}/p^r -algebra with lift of Frobenius. Applying Proposition 2.23 to it, we get an exact sequence

$$0 \longrightarrow (\underline{\mathbb{Z}/p^r})^{\text{et}} \longrightarrow \pi_0 \mathcal{O}_R \xrightarrow{\varphi - 1} \pi_0 \mathcal{O}_R \longrightarrow 0$$

of étale sheaves on Spec(R). Taking the long exact sequence of cohomologies associated with this exact sequence of sheaves, we get an exact sequence

$$0 \longrightarrow H^0_{\text{et}}(\operatorname{Spec}(R); \mathbb{Z}/p^r) \to A/p^r \to A/p^r.$$

Now, the zeroth étale cohomology with constant coefficients is locally constant functions so

$$H^0_{\text{et}}(\operatorname{Spec}(R); \mathbb{Z}/p^r) \simeq \underline{\mathbb{Z}/p^r}(R) = \underline{\mathbb{Z}/p^r}(A/p^r)$$

Finally, the assumption that A is derived p-complete implies that $(A, (p^r))$ is an Henselian pair ([Sta18, Lemma 15.93.10]), so that $A \to A/p^r$ induces a bijection on Zariski-components [Sta18, Lemma 15.11.6], and hence on global sections of the constant sheaf $\underline{\mathbb{Z}/p^r}$. This concludes the proof.

2.2. Rigidity for δ_p -group algebras. In this section, we compute the rank 1 units of a *p*-complete group algebra $A[M]_p^{\wedge}$ in terms of M and the rank 1 units in A. It will be convenient to phrase our results in terms of a slight variant of $\mathbb{G}_m^{\delta_p}(-)$.

Definition 2.25. For an abelian group M, set

$$\mathbb{G}_m^{\delta_p, \mathrm{red}}(A[M]) := \mathrm{Ker}(\mathbb{G}_m^{\delta_p}(A[M]) \to \mathbb{G}_m^{\delta_p}(A)), \\
\mathbb{G}_m^{\delta_p, \mathrm{red}}(A[M]_p^\wedge) := \mathrm{Ker}(\mathbb{G}_m^{\delta_p}(A[M]_p^\wedge) \to \mathbb{G}_m^{\delta_p}(A)).$$

Here, the δ_p -structure on the *p*-completion of a δ_p -ring R is the unique one for which $R \to R_p^{\wedge}$ is a δ_p -ring map, see [BS22, Lemma 2.17]. We refer to these as the *reduced units* of rank 1.

Since the augmentation $A[M]_p^{\wedge} \to A$ is split, Theorem 2.7 is equivalent to:

Theorem 2.26. Let A be a p-complete δ_p -ring and M be a finitely generated abelian group whose torsion part is p-power torsion. Then there is a natural isomorphism

$$\mathbb{G}_m^{\delta_p, \mathrm{red}}(A[M]_p^\wedge) \cong \underline{M}(A).$$

We handle the case where M is a finite abelian p-group in §2.2.1 and the general case in 2.2.2.

2.2.1. The torsion case. Our strategy is to apply the square nil-invariance results to the augmentation ideal in A[M] when M is a finite abelian p-group. To do so, we need the following lemma:

Lemma 2.27. Let A be a p-complete commutative ring, let M be a finite abelian p-group, and let $J \subset A[M]$ denote the kernel of the augmentation $A[M] \to A$. Then A[M] is (p, J)-complete.

Proof. Note that J is generated by the finite set of elements [m] - 1 for $m \in M$. Since A is assumed to be p-complete, it suffices to show that some power of each [m] - 1 is divisible by p. But we have that

$$([m] - 1)^{p^j} \equiv [m]^{p^j} - 1 = [p^j m] - 1 \pmod{p}$$

and so this is 0 (mod p) for p^j larger than the order of torsion in M.

It follows from Proposition 2.15 that we may identify $\mathbb{G}_m^{\delta_p, \text{red}}(A[M])$ with $(J/J^2)^{\delta_p=1}$. We now compute this group.

Lemma 2.28. Let A be a δ_p -ring, let M be a finite abelian p-group, and let $J \subset A[M]$ denote the kernel of the augmentation $A[M] \to A$. Then there is a canonical isomorphism of abelian groups

$$A \otimes M \simeq J/J^2$$
$$a \otimes m \mapsto a([m] - 1) \in J$$

Under this isomorphism, the operation δ_p on the right-hand side is identified with the endomorphism $\varphi_A \otimes id_M$ on the left-hand side.

Proof. To see this isomorphism of abelian groups, we note that J is a free A-module with generators [m] - 1 for $m \in M$, and that J^2 is generated by the relations ([m] - 1)([n] - 1) = 0, that is,

$$([m] - 1) + ([n] - 1) \equiv [m + n] - 1 \mod J^2.$$

Thus, the quotient J/J^2 is visibly the same as the left-hand side, so it remains to identify these two operations $\varphi_A \otimes \operatorname{id}_M$ and δ_p .

Since $\delta_p(a([m]-1)) \equiv \varphi_A(a)\delta_p([m]-1)$ on J/J^2 by Proposition 2.9(1), it suffices to prove the claim for a = 1. In this case we have

$$\delta_p([m]-1) = \frac{\varphi([m]-1) - ([m]-1)^p}{p} = \frac{([m]^p - 1) - ([m]-1)^p}{p}$$

The right-hand side is a polynomial q([m]) with integer coefficients such that q(1) = 0 and q'(1) = 1, and hence

$$q([m]) \equiv [m] - 1 \mod ([m] - 1)^2$$

as desired.

Given this identification, we can use the results of 2.1.3 to prove the desired result:

Proposition 2.29. Let A be a p-complete δ_p -ring and M be a finite abelian p-group. Then there is a natural isomorphism

$$\mathbb{G}_m^{\delta_p, \operatorname{red}}(A[M]_p^{\wedge}) \simeq \underline{M}(A).$$

Proof. By Proposition 2.15 and Lemma 2.28, we may identify $\mathbb{G}_m^{\delta_p, \text{red}}(A[M]_p^{\wedge})$ with the fixed points of $\varphi_A \otimes \text{id}_M$ on $A \otimes M$. Since M is a finite abelian p-group, it is a finite direct sum of quotients of the form \mathbb{Z}/p^r , so we may reduce to the case $M = \mathbb{Z}/p^r$. In this case, we have $A \otimes M = A/p^r$ and Corollary 2.24 identifies the fixed points of $\varphi_A \otimes \text{id}_M$ on this object with $\mathbb{Z}/p^r(A) = \underline{M}(A)$.

2.2.2. The general case.

Theorem 2.30. Let M be a finitely generated abelian group whose torsion part is p-power torsion and let A be a p-complete δ_p -ring. Then

$$\mathbb{G}_m^{\delta_p, \mathrm{red}}(A[M]_p^\wedge) \simeq \underline{M}(A).$$

Proof. Note that there is a canonical map $\underline{M}(A) \to \mathbb{G}_m^{\delta_p, \text{red}}(A[M]_p^{\wedge})$ locally given by $m \mapsto [m]$. We shall refer to its image as the "tautological units". Note further that the assumption on M implies that $\bigcap_r p^r M = \{0\}$. As a result, for every finite subset $T \subseteq M$, there exists r > 0 such that T injects into M/p^r .

Given a reduced unit $f \in \mathbb{G}_m^{\delta_p, \text{red}}(A[M]_p^{\wedge})$ of rank 1, we wish to show that we can find a *finite* decomposition $A \simeq \prod_i A_i$ such that the image of f in $A_i[M]_p^{\wedge}$ is of the form [m] for some $m \in M$. Expand f as a p-adically convergent series

$$f = \sum_{m \in M} x_m[m],$$

and let $f_{r,s}$ be the image of f in $A/p^r[M/p^s]$. In this notation, we will allow $r = \infty$ or $s = \infty$, in which case we write $f_{\infty,s}$ (resp. $f_{r,\infty}$) to mean the image of f in $A[M/p^s]$ (resp. $A/p^r[M]$). Thus, our aim is to show that $f := f_{\infty,\infty}$ is a tautological unit.

Since $f_{\infty,s}$ is a reduced unit of rank 1, by Proposition 2.29 it is a tautological unit for every $s \in \mathbb{N}$. On the other hand, $f_{1,\infty}$ is an element of A/p[M] so it is a sum $f_{1,\infty} = \sum_{m \in T} \overline{x}_m[m]$, for some finite $T \subseteq M$. Choosing $s \in \mathbb{N}$ for which the map $T \to M/p^s$ is injective, we see that the fact that $f_{1,s}$ is tautological implies the same for $f_{1,\infty}$. This implies that there is a finite decomposition $A/p \simeq \prod_i \overline{A}_i$ such that the image of $f_{1,\infty}$ in $\overline{A}_i[M]$ is tautological. By Hensel's lemma, we can lift the mentioned decomposition to a decomposition $A \simeq \prod_i A_i$. Replacing A by one of the A_i 's, we may thus assume that $f_{1,\infty} = [m]$ for some $m \in M$. Multiplying by [-m], we are thus reduced to the case where $f_{1,\infty} = 1$. We claim that in this case, we already have f = 1, and in particular that it is tautological. Indeed, assuming the contrary, let $0 \neq m_0 \in M$ be an element for which $x_{m_0} \neq 0$. Choose r large enough so that $x_m \neq 0 \mod p^r$, and write again

$$f_{r,\infty} = \sum_{m \in T'} x_m[m]$$

for some finite set $T' \subseteq M$. Choose s large enough so that the composition $T' \hookrightarrow M \twoheadrightarrow M/p^s$ is injective. Then, the fact that $x_{m_0} \neq 0 \mod p^r$ implies that

$$f_{r,s} = 1 + \overline{x}_{m_0}[m_0] + \dots,$$

where \overline{x}_{m_0} is the (non-vanishing) reduction of x_{m_0} modulo p^r . Since $f_{r,s}$ is the image of $f_{\infty,s}$ under the map $A \to A/p^r$, it is also tautological, and since the coefficient of $0 \in M$ is invertible, this forces all other coefficients, including \overline{x}_{m_0} , to be 0. This is a contradiction and we are done. 2.3. $\hat{\delta}$ -rings. We now introduce a global variant of δ_p -rings which we call $\hat{\delta}$ -rings. In contrast with the theory of λ -rings, our notion of $\hat{\delta}$ -rings "decouples" different primes.

Definition 2.31. Define the category of $\hat{\delta}$ -rings by the pullback

$$\widehat{\delta} - \operatorname{Ring} \longrightarrow \prod_{p} \operatorname{ACR}^{\delta_{p}} \\ \downarrow \qquad \qquad \downarrow \\ \operatorname{CAlg}_{\mathbb{Z}}^{\heartsuit} \xrightarrow{\prod(-)_{p}^{\wedge}} \prod_{p} \operatorname{ACR}_{p}^{\wedge}.$$

In other words, a $\hat{\delta}$ -ring is a (discrete) commutative ring R with a lift of each derived p-completion R_p^{\wedge} to an animated δ_p -ring.

Although we assume the underlying ring R is discrete, its (derived) *p*-completions may not be, and thus we need to work in the animated context for the above definition. Nevertheless, it is easy to see that the category of $\hat{\delta}$ -rings is (equivalent to) a 1-category. Additionally, we will generally only consider $\hat{\delta}$ -rings with bounded torsion, so this point will not be important.

Remark 2.32. If R is already derived p-complete, then a $\hat{\delta}$ -ring structure on R is exactly given by a δ_p -ring structure, since all other completions vanish. On the other hand, if R is rational, then a $\hat{\delta}$ -ring structure on R is no data at all.

As in Construction 2.3, $\mathbb{Z}[M]$ for an abelian group M acquires a natural $\hat{\delta}$ -ring structure.

Definition 2.33. Define the group of rank 1 $\hat{\delta}$ -units of R by

$$\mathbb{G}_m^{\delta}(R) := \operatorname{Map}_{\widehat{\delta} - \operatorname{Ring}}(\mathbb{Z}[t^{\pm 1}], R).$$

While $\mathbb{G}_{m}^{\widehat{\delta}}(R)$ can in general be non-discrete, we will typically consider R with bounded torsion, in which case $\mathbb{G}_{m}^{\widehat{\delta}}(R)$ can be described as the subgroup of units $t \in R^{\times}$ such that $\delta_{p}(t) = 0$ for all p. This notion of $\widehat{\delta}$ -rings allows us to make an integral analogue of the previous results in the section, which we view as the algebraic analogue of our main theorem (Theorem 1.1).

Theorem 2.34. Let M and N be finitely generated abelian groups. Then the natural map

$$\operatorname{Hom}_{\operatorname{Ab}}(N, M) \to \operatorname{Hom}_{\widehat{\delta}-\operatorname{Ring}}(\mathbb{Z}[N], \mathbb{Z}[M])$$

is an isomorphism. In particular, $\mathbb{G}_m^{\widehat{\delta}}(\mathbb{Z}[M]) \simeq M$.

Proof. Both sides send colimits in N to limits, so it suffices to consider $N = \mathbb{Z}$, where the statement becomes $\mathbb{G}_m^{\widehat{\delta}}(\mathbb{Z}[M]) \simeq M$. Here, write $M = M' \oplus \Lambda$ where Λ is free and M' is finite. Since $\mathbb{Z}[M']$ has connected Zariski spectrum, we have an isomorphism $\mathbb{G}_m(\mathbb{Z}[M'][\Lambda]) \simeq \mathbb{G}_m(\mathbb{Z}[M']) \oplus \Lambda$ (cf. [Kar83, Theorem 1]). It follows that

$$\mathbb{G}_m^{\widehat{\delta}}(\mathbb{Z}[M'][\Lambda]) \simeq \mathbb{G}_m^{\widehat{\delta}}(\mathbb{Z}[M']) \oplus \Lambda.$$

Hence, it suffices to prove the result for M finite. We start by showing that in this case $\mathbb{G}_m^{\hat{\delta}}(\mathbb{Z}[M])$ is finite. Choose a prime p. Since the p-completion map

$$\mathbb{G}_m^{\delta}(\mathbb{Z}[M]) \to \mathbb{G}_m^{\delta_p}(\mathbb{Z}[M]_p^{\wedge})$$

is injective, it suffices to show that $\mathbb{G}_m^{\delta_p}(\mathbb{Z}[M]_p^{\wedge})$ is finite. Writing $M = M_p \oplus M_{\bar{p}}$ where M_p is a *p*-group and $M_{\bar{p}}$ is of order prime to *p*, we have by Proposition 2.29 that

$$\mathbb{G}_m^{\delta_p}(\mathbb{Z}[M]_p^{\wedge}) \simeq M_p(\mathbb{Z}[M_{\bar{p}}]_p^{\wedge}) \oplus \mathbb{G}_m^{\delta_p}(\mathbb{Z}[M_{\bar{p}}]_p^{\wedge}).$$

The first summand is finite because $\operatorname{Spec}(\mathbb{Z}[M_{\bar{p}}]_p^{\wedge})$ has finitely many connected components, and the second is finite because $\mathbb{Z}[M_{\bar{p}}]_p^{\wedge}$ is perfect and hence by Example 2.6 we have

$$\mathbb{G}_m^{\delta_p}(\mathbb{Z}[M_{\bar{p}}]_p^{\wedge}) \simeq \mathbb{G}_m(\mathbb{F}_p[M_{\bar{p}}]).$$

Finally, once we know $\mathbb{G}_m^{\hat{\delta}}(\mathbb{Z}[M])$ is finite, the result follows from [Hig40, Theorem 2], which asserts that all torsion reduced units in the group algebra are in the image of the map $M \to \mathbb{G}_m(\mathbb{Z}[M])$. \Box

2.4. δ -rings and commutative ring spectra. Now we will link the existence of $\hat{\delta}$ -ring structures on R to lifts \mathbb{S}_R to the sphere spectrum. Recall from [NS18, §IV.1] that a commutative ring spectrum R admits two natural maps to the Tate construction R^{tC_p} : the Tate-valued Frobenius

frob: $R \to R^{tC_p}$,

given by the composite of the Tate diagonal $R \to (R^{\otimes p})^{tC_p}$ with multiplication, and the map

triv: $R \to R^{tC_p}$.

given by the composite $R \to R^{hC_p} \to R^{tC_p}$. Note that the latter is defined for an arbitrary spectrum and does not depend on the multiplication, but when R is a commutative ring spectrum then triv is a commutative ring map.

The Segal conjecture supplies a large collection of spectra for which the triv map is an isomorphism.

Definition 2.35. Let X be a spectrum. We say that X is of *finite Tor-amplitude* if X is bounded below and $X \otimes \mathbb{Z}$ is bounded. Equivalently, X admits a cell structure with cells of bounded dimension. We further say that X is a *Moore spectrum* if $X \otimes \mathbb{Z}$ is concentrated in degree 0.

Proposition 2.36 ([BS24, Theorem 1.2, Example 4.4], cf. also [Yua22, Lemma 6.7]). Let X be a spectrum of finite Tor-amplitude. Then, triv : $X \to X^{tC_p}$ exhibits X^{tC_p} as the p-completion of X.

Thus, any commutative ring spectrum R of finite Tor-amplitude admits a canonical map

$$\varphi_R := \operatorname{triv}^{-1} \circ \operatorname{frob} \colon R \to R_n^{\wedge},$$

which we call the *Frobenius map* of R. Our goal for the rest of the section is to show that in the special case where R is a Moore spectrum (i.e., $R \otimes \mathbb{Z}$ is *discrete*), these Frobenius maps for varying p induce a $\hat{\delta}$ -ring structure on $\pi_0 R$. We need a few preliminary lemmas.

Lemma 2.37. Let $F, G : ACR \to C$ be functors and F_0, G_0 denote their restriction to the full subcategory $\operatorname{Poly}_{\mathbb{Z}} \subset ACR$ of (discrete) polynomial algebras. If F commutes with sifted colimits, then there is an isomorphism of spaces of natural transformations

$$\operatorname{Nat}(F,G) \simeq \operatorname{Nat}(F_0,G_0)$$

Proof. This follows from the fact that $ACR = P_{\Sigma}(Poly_{\mathbb{Z}})$ is the the animation of the category of polynomial rings.

Lemma 2.38. Let R be a discrete commutative ring, so that the Frobenius and trivial maps $R \to R^{tC_p}$ factor uniquely through $\tau_{\geq 0}R^{tC_p}$. Then, there is a natural isomorphism $\tau_{[0,1]}R^{tC_p} \simeq R/p$ in $\operatorname{CAlg}_{\mathbb{Z}}^{\operatorname{cn}}$ fitting into the commutative diagrams

$$\begin{array}{cccc} R & \xrightarrow{\mathrm{triv}} & \tau_{\geq 0} R^{tC_p} & & R & \xrightarrow{\mathrm{frob}} & \tau_{\geq 0} R^{tC_p} \\ & & & \downarrow^{\wr} & & \downarrow^{\downarrow} & \\ R /\!\!/ p & \xrightarrow{=} & R /\!\!/ p \simeq \tau_{[0,1]} R^{tC_p} & & R /\!\!/ p \xrightarrow{\varphi_{R /\!\!/ p}} & R /\!\!/ p \simeq \tau_{[0,1]} R^{tC_p}. \end{array}$$

Proof. Note that for $R \in \text{Poly}_{\mathbb{Z}}$, the constructions $R/\!\!/ p$ and $\tau_{[0,1]} R^{tC_p}$ are both given by the discrete ring R/p and thus are naturally identified. By Lemma 2.37, we obtain a natural transformation

$$R/\!\!/ p \xrightarrow{\nu} \tau_{[0,1]} R^{tC_p}$$

which is an isomorphism at least for $R \in \operatorname{Poly}_{\mathbb{Z}}$. Moreover, the diagrams commute because they commute for $R \in \operatorname{Poly}_{\mathbb{Z}}$; this is clear for the first one and for the second, it amounts to the statement that frob induces the usual Frobenius on $\pi_0 R/p$ [NS18, Example IV.1.2].

It remains to check that ν is an isomorphism for *all* discrete rings. Let $G(R) : \operatorname{CAlg}_{\mathbb{Z}}^{\operatorname{an}} \to \operatorname{Mod}_{\mathbb{Z}}$ be the cofiber of the composite

$$R_{hC_p} \xrightarrow{\operatorname{Nm}} R^{hC_p} \xrightarrow{\operatorname{res}} R.$$

Note that we have a natural transformation $\tau_{[0,1]}R^{tC_p} \to \tau_{[0,1]}G(R)$ which is an isomorphism on discrete rings. Therefore, it suffices to show that for R discrete, the induced map

$$\nu': R/\!\!/ p \to \tau_{[0,1]}G(R)$$

is an isomorphism. But the source and target both preserve colimits, as functors to $\operatorname{Mod}_{\mathbb{Z}}^{[0,1]}$, so the conclusion follows from the fact that ν , and therefore ν' , is an isomorphism for $R \in \operatorname{Poly}_{\mathbb{Z}}$.

Lemma 2.39. Let $R, S \in ACR$ and assume that S is 1-truncated. Then

$$\operatorname{Map}_{\operatorname{ACR}}(R, S) \to \operatorname{Map}_{\operatorname{Calg}_{\pi}}(R, S)$$

is an inclusion of connected components, which is an equivalence if $\pi_1(S)$ does not have 2-torsion.

Proof. First, since the condition of the theorem is closed under limits, it suffices to consider the case $R = \mathbb{Z}[x]$ where we consider the map $\Omega^{\infty}S \to \operatorname{Map}_{\operatorname{CAlg}_{\mathbb{Z}}}(\mathbb{Z}[x], S)$. Letting $\mathbb{Z}\{x\} = \bigoplus_{n \geq 0} \mathbb{Z}[B\Sigma_n]$ denote the free commutative \mathbb{Z} -algebra on a degree 0 class, we note that $\pi_0\mathbb{Z}\{x\} = \mathbb{Z}[x]$ and $\pi_1\mathbb{Z}\{x\} = \beta\mathbb{Z}/2[x]$ (for instance because the sign character $\Sigma_n \to C_2$ is the abelianization map). Consequently, defining P by the pushout

$$\mathbb{Z}\{S^1/2\} \longrightarrow \mathbb{Z}$$
$$\downarrow^{\beta} \qquad \qquad \downarrow$$
$$\mathbb{Z}\{x\} \longrightarrow P,$$

the natural map $P \to \mathbb{Z}[x]$ is an isomorphism in degrees 0, 1 and surjective on π_2 . It follows that $\operatorname{Map}(P, S) \simeq \operatorname{Map}(\mathbb{Z}[x], S)$, so it suffices to show that $\Omega^{\infty}S \to \operatorname{Map}_{\operatorname{CAlg}_{\mathbb{Z}}}(P, S)$. The conclusion then follows by considering commutative \mathbb{Z} -algebra maps from the above pushout square into S. \Box

Lemma 2.40. Let X be a bounded below spectrum. Then

$$(X_n^{\wedge}) \otimes \mathbb{Z} = (X \otimes \mathbb{Z})_n^{\wedge}$$


Proof. This follows because $(X_p^{\wedge}) \otimes \mathbb{Z}$ is *p*-complete [BB19, Lemma 3.3] and the natural map $(X_p^{\wedge}) \otimes \mathbb{Z} \to (X \otimes \mathbb{Z})_p^{\wedge}$ becomes an isomorphism after tensoring with \mathbb{S}/p .

We can now prove the main statement about the relationship between commutative ring Moore spectra and $\hat{\delta}$ -rings:

Proposition 2.41. Let $\operatorname{CAlg}^{M} \subset \operatorname{CAlg}(\operatorname{Sp})$ denote the full subcategory of commutative ring Moore spectra. Then for $R \in \operatorname{CAlg}^{M}$, $\pi_{0}(R)$ acquires a natural $\widehat{\delta}$ -ring structure, i.e., $\pi_{0}(-)$ lifts to a functor

$$\pi_0 : \mathrm{CAlg}^\mathrm{M} \to \widehat{\delta} - \mathrm{Ring}.$$

Proof. For each prime p, consider the diagram of commutative ring spectra



where the right vertical map is induced by the animated Frobenius. The commutativity of the diagram is clear other than the right two triangles, which commute by Lemma 2.38. Since the right vertical map is \mathbb{Z} -linear and $\mathbb{Z} \otimes R_p^{\wedge} \simeq (\pi_0 R)_p^{\wedge}$ by Lemma 2.40, the outer square induces a square

$$\begin{array}{ccc} (\pi_0 R)_p^{\wedge} \xrightarrow{\varphi_R \otimes \mathbb{Z}} (\pi_0 R)_p^{\wedge} \\ & & \downarrow \\ & & \downarrow \\ \pi_0 R /\!\!/ p \xrightarrow{\varphi_{\pi_0 R /\!\!/ p}} \pi_0 R /\!\!/ p. \end{array}$$

To see that this induces an animated δ_p -ring structure on $(\pi_0 R)_p^{\wedge}$, it suffices to lift the left vertical map to an animated ring map. But $(\pi_0 R)_p^{\wedge}$ is 1-truncated, and we claim there is no 2-torsion in π_1 in case p = 2. Indeed, if there were, then $\pi_0(R)_2^{\wedge}/2 = \pi_0(R)/2$ would have nontrivial π_2 , which is impossible. Thus, by Lemma 2.39, $\varphi_R \otimes \mathbb{Z}$ lifts uniquely to an animated map, as desired.

Remark 2.42. In fact, A. Krause has shown that δ -ring structures on a torsion-free commutative ring are *equivalent* to H_{∞} -ring structures on the corresponding Moore spectrum [Kra].

3. TAUTOLOGICAL UNITS, RIGIDITY, AND STATICITY

Let R_0 be a (discrete) commutative ring and let M be a finitely generated abelian group. Then there is a natural map of abelian groups

$$\theta: M \to R_0[M]^{\times}$$
$$m \mapsto [m]$$

picking out what we call the *tautological units*. There are at least two reasons why this map fails to be an isomorphism:

- (1) The target also contains units coming from R_0^{\times} . (2) If R_0 splits as a product $R'_0 \times R''_0$, then we have units of the form $([m], [m']) \in (R'_0 \times R''_0[M])^{\times}$.

These obstructions carry over verbatim when R_0 is replaced by a commutative ring spectrum R and units are replaced by strict units. To side-step (1), we introduce the following definition:

Definition 3.1. Let R be a commutative ring spectrum and S be an augmented commutative *R*-algebra (e.g. R[M] or $R[M]_p^{\wedge}$ with their canonical augmentations). Then we denote

$$\mathbb{G}_m^{\mathrm{red}}(S) := \mathrm{fib}(\mathbb{G}_m(S) \to \mathbb{G}_m(R))$$

so that there is a splitting

$$\mathbb{G}_m(S) \simeq \mathbb{G}_m(R) \oplus \mathbb{G}_m^{\mathrm{red}}(S).$$

We refer to $\mathbb{G}_m^{\text{red}}(S)$ as the reduced strict units of S. ⁶

Thus, the question of computing $\mathbb{G}_m(R[M])$ reduces to separately computing $\mathbb{G}_m(R)$ and $\mathbb{G}_m^{\text{red}}(R[M])$. In our case of interest R = S, the former vanishes by [Car22], and so we begin our analysis in §3.1 with the basic properties of reduced units $\mathbb{G}_m^{\mathrm{red}}(-)$.

Next, while the map θ lands in reduced units, (2) still obstructs it from being an isomorphism. Thus, we consider instead a certain natural map

$$\Theta: \underline{M}(R) \to \mathbb{G}_m^{\mathrm{red}}(R[M])$$

where $\underline{M}(R)$ denotes the group of locally constant M-valued functions on $\operatorname{Spec}(R)$. The map Θ is a better approximation to $\mathbb{G}_m^{\mathrm{red}}(R[M])$ than θ , and turns out to be an isomorphism for a large class of commutative ring spectra R. We refer to this feature of (R, M) as rigidity. Much of the paper is dedicated to proving rigidity in various situations, so we set notation more carefully in §3.2.

Finally, in §3.3, we discuss a condition under which rigidity is easy to check. We say that a pair (R, M) is $\mathbb{G}_m^{\mathrm{red}}(-)$ -static if the natural map

$$\mathbb{G}_m^{\mathrm{red}}(R[M]) \to \mathbb{G}_m^{\mathrm{red}}(\pi_0 R[M])$$

is an inclusion of connected components. In this case, rigidity reduces to an essentially algebraic question of checking whether all (reduced) strict units of R[M] are tautological at the level of π_0 (cf. Proposition 3.22).

⁶We warn that the notation is slightly abusive, as $\mathbb{G}_m^{\mathrm{red}}(S)$ depends on both S and its augmentation, but we will clarify whenever there is risk for confusion.

3.1. Strict units, reduced units, and descent. The main object of study in this paper is the following:

Definition 3.2. A strict unit in a commutative ring spectrum R is a map of commutative ring spectra

$$\mathbb{S}[\mathbb{Z}] =: \mathbb{S}[t^{\pm 1}] \to R$$

We denote the spectrum of strict units by $\mathbb{G}_m(R)$, so that

$$\Omega^{\infty} \mathbb{G}_m(R) = \operatorname{Map}_{\operatorname{CAlg}}(\mathbb{S}[t^{\pm 1}], R) \simeq \operatorname{Map}_{\operatorname{Sp}}(\mathbb{Z}, gl_1(R)).$$

In fact, $\mathbb{G}_m(R)$ naturally has a connective \mathbb{Z} -module spectrum structure [Lur18, Cons. 1.6.10].

As mentioned earlier, we will need the following variant of strict units:

Definition 3.3 (Reduced units). Let R be a commutative ring spectrum and S be an augmented commutative R-algebra. Then we denote

$$\mathbb{G}_m^{\mathrm{red}}(S) := \mathrm{fib}(\mathbb{G}_m(S) \to \mathbb{G}_m(R))$$

so that there is a splitting

$$\mathbb{G}_m(S) \simeq \mathbb{G}_m(R) \oplus \mathbb{G}_m^{\mathrm{red}}(S).$$

We refer to $\mathbb{G}_m^{\mathrm{red}}(S)$ as the reduced strict units of S.⁷

For us, we will generally take S to be R[M] or $R[M]_p^{\wedge}$ with their canonical augmentations.

Remark 3.4. The reduced strict units can alternately be interpreted as a space of *augmented* maps:

$$\Omega^{\infty} \mathbb{G}_m^{\mathrm{red}}(R[M]) \simeq \mathrm{Map}_{\mathrm{CAlg}_p^{\mathrm{aug}}}(R[\mathbb{Z}], R[M]).$$

Accordingly, we have for every $R \in CAlg$ an adjunction

$$R[-]: \operatorname{Mod}_{\mathbb{Z}}^{\operatorname{cn}} \rightleftharpoons \operatorname{CAlg}_{B}^{\operatorname{aug}} : \mathbb{G}_{m}^{\operatorname{red}}(-).$$

3.1.1. Descent for reduced units. The construction $\mathbb{G}_m^{\text{red}}(R[M])$ is functorial in both the ring R and the group M. In the ring R, we have the following descent statement:

Proposition 3.5. For $M \in \text{Mod}_{\mathbb{Z}}^{\text{cn}}$, the functors $\mathbb{G}_m(-[M]), \mathbb{G}_m^{\text{red}}(-[M]) \in \text{Fun}(\text{CAlg}, \text{Mod}_{\mathbb{Z}}^{\text{cn}})$ are sheaves for the Zariski topology. If M is additionally finite and discrete, then $\mathbb{G}_m(-[M])$ and $\mathbb{G}_m^{\text{red}}(-[M])$ are moreover affine group schemes (i.e., they are co-representable by commutative Ralgebras).

Proof. Since $\mathbb{G}_m^{\text{red}}-[M]$ is a retract of $\mathbb{G}_m(-[M])$, it suffices to prove the claims for the latter functor. Since the Zariski topology is finitary, the sheaf condition for $\mathbb{G}_m(-[M])$ amounts to the construction $R \mapsto \mathbb{G}_m(R[M])$ commuting with a finite limit, which follows from the facts that $R \mapsto R[M]$ preserves finite limits and \mathbb{G}_m preserves all limits. If M is finite, then $R \mapsto R[M]$ and \mathbb{G}_m are accessible and limit preserving, so the result follows from the Adjoint Functor Theorem. \Box

The behavior of $\mathbb{G}_m^{\text{red}}(R[M])$ as M changes is more subtle, but there are some cases when it interacts with limits or colimits in a predictable way. First, while $\mathbb{G}_m^{\text{red}}(R[-])$ does not preserve direct sums, one can always decompose $\mathbb{G}_m^{\text{red}}(R[M \oplus N])$ into simpler pieces.

⁷We warn that the notation is slightly abusive, as $\mathbb{G}_m^{\mathrm{red}}(S)$ depends on both S and its augmentation, but we will clarify whenever there is risk for confusion.

Observation 3.6. Let R be a commutative ring spectrum and let $M, N \in Mod_{\mathbb{Z}}^{cn}$. Then there are canonical isomorphisms

$$\mathbb{G}_m^{\mathrm{red}}(R[M \oplus N]) \simeq \mathbb{G}_m^{\mathrm{red}}(R[M]) \oplus \mathbb{G}_m^{\mathrm{red}}(R[M][N]),$$
$$\mathbb{G}_m^{\mathrm{red}}(R[M \oplus N]_p^{\wedge}) \simeq \mathbb{G}_m^{\mathrm{red}}(R[M]_p^{\wedge}) \oplus \mathbb{G}_m^{\mathrm{red}}(R[M][N]_p^{\wedge})$$

where we regard R[M][N] as being augmented over R[M] (and similarly for the p-complete version).

We also have the following descent statement along injections $M \to N$ of abelian groups.

Definition 3.7. For a morphism $\alpha: M \to N$ in $Mod_{\mathbb{Z}}$, its *Čech nerve* is the cosimplicial diagram

$$C^{\bullet}(\alpha) := \left(N \Longrightarrow N \oplus_{M} N \Longrightarrow N \oplus_{M} N \oplus_{M} N \oplus_{M} N \cdots \right) \in (\operatorname{Mod}_{\mathbb{Z}}^{\operatorname{cn}})^{\Delta}$$

obtained as an augmented cosimplicial object from α by left Kan extension along $\Delta_{+}^{\leq 0} \hookrightarrow \Delta_{+}$.

Proposition 3.8. Let $\alpha: M \hookrightarrow N$ be an injective map of abelian groups. Then the natural maps

$$\mathbb{G}_m^{\mathrm{red}}(R[M]) \to \varprojlim \mathbb{G}_m^{\mathrm{red}}(R[C^{\bullet}(\alpha)])$$
$$\mathbb{G}_m^{\mathrm{red}}(R[M]_p^{\wedge}) \to \varprojlim \mathbb{G}_m^{\mathrm{red}}(R[C^{\bullet}(\alpha)]_p^{\wedge})$$

are isomorphisms.

Proof. We consider the non *p*-complete case first. Since $\mathbb{G}_m^{\mathrm{red}}(R[M])$ is the fiber of the map $\mathbb{G}_m(R[M]) \to \mathbb{G}_m(R)$ and constant functors preserve contractible limits, it suffices to prove the analogous statement for $\mathbb{G}_m(R[M])$. This follows from the fact that \mathbb{G}_m preserves limits and $R[M] \to R[N]$ is descendable in the sense of [Mat16, Definition 3.18, Proposition 3.20], as it admits a retract. The *p*-complete case follows similarly, using that $R[M]/p \to R[N]/p$ admits a retract.

3.2. **Rigidity.** We begin by constructing the inclusion of the tautological units.

Definition 3.9 (Constant group scheme [Lur18, Exm. 2.5.7]). For $M \in Mod_{\mathbb{Z}}^{cn}$, denote by

$$\underline{M} : \operatorname{CAlg} \to \operatorname{Mod}_{\mathbb{Z}}^{\operatorname{cn}}$$

the Zariski sheafification of the constant functor valued at M.

Remark 3.10. For an abelian group M and commutative ring spectrum R, we have

 $\underline{M}(R) = \{ \text{Locally constant functions } f : |\operatorname{Spec}(R)| \to M \}.$

Moreover, when M is finite, the functor \underline{M} is corepresented by the commutative ring spectrum \mathbb{S}^{M} .

Construction 3.11 (Tautological units). A consequence of Definition 3.9 is that for any Zariski sheaf F on CAlg, we have

$$\operatorname{Nat}(\underline{M}(-), F) = \operatorname{Map}_{\operatorname{Mod}_{-}^{\operatorname{cn}}}(M, F(\mathbb{S})),$$

since the constant functor construction is adjoint to evaluation at the initial object. Applying this to the Zariski sheaf $F = \mathbb{G}_m^{\text{red}}(-[M])$ (Proposition 3.5), we define

$$\Theta_M : \underline{M}(-) \to \mathbb{G}_m^{\mathrm{red}}(-[M])$$

to be the unique natural transformation whose value on \mathbb{S} agrees with the unit map $M \to \mathbb{G}_m^{\mathrm{red}}(\mathbb{S}[M])$ of the adjunction of Remark 3.4.

The bulk of this paper is dedicated to analyzing the pairs $(R \in \text{CAlg}, M \in \text{Ab})$ for which $\Theta_M(R)$ is an isomorphism.

Definition 3.12. Let M be an abelian group and let R be a commutative ring spectrum.

- (1) We say that R is *M*-rigid if the map $\Theta_M(R): \underline{M}(R) \to \mathbb{G}_m^{\mathrm{red}}(R[M])$ is an isomorphism.
- (2) We say that R is *p*-rigid if it is *M*-rigid for every finite abelian *p*-group *M*.

Variant 3.13. If R is additionally p-complete, we say R is p-completely M-rigid or p-rigid if the analogous conditions hold for the composite $\widehat{\Theta}_M(R) : \underline{M}(R) \xrightarrow{\Theta} \mathbb{G}_m^{\mathrm{red}}(R[M]) \to \mathbb{G}_m^{\mathrm{red}}(R[M]_p^{\wedge})$. Note that for M finite, $R[M] \simeq R[M]_p^{\wedge}$ so this coincides with the above definition.

Example 3.14 (see, e.g., [Kar83, Theorem 1]). Let R be a discrete commutative ring and let Λ be a finitely generated free abelian group, i.e. $\Lambda \simeq \mathbb{Z}^k$. A commutative ring R is Λ -rigid if and only if R is *reduced*, that is, contains no non-zero nilpotent elements. More generally, for every connected commutative ring R, the units of $R[\Lambda]$ are the products $r \cdot \lambda_0 \cdot f$ where $r \in R^{\times}$, $\lambda_0 \in \Lambda$, and $f \in R[\Lambda]$ is congruent to 1 modulo the nil-radical of R.

We next discuss closure properties of rigidity with respect to M and R. First, since Θ_M is a morphism of Zariski sheaves, the collection of M-rigid commutative ring spectra R is always closed under Zariski covers. For *finite* M, we have a much stronger closure property.

Proposition 3.15. Let M be a finite abelian group. Then the collection of M-rigid (resp. p-rigid) commutative ring spectra is closed under all limits in CAlg(Sp).

Proof. The second statement clearly follows from the first. For fixed finite M, it suffices to note that $\mathbb{G}_m^{\text{red}}(-[M])$ and $\underline{M}(-)$ are representable (Proposition 3.5, Remark 3.10).

3.3. Staticity. Consider the forgetful map

(5)
$$\mathbb{G}_m(R) \to \mathbb{G}_m(\pi_0(R)) = \pi_0(R)^{\times}.$$

One difficulty in computing strict units is that $\mathbb{G}_m(R)$ may not be discrete, and so while it can be relatively easy to see which units in $\pi_0(R)$ lift to strict units (as we saw in the introduction), there may be a nontrivial space of ways in which they lift. In this paper, we are interested in when this does *not* happen – that is, when (5) is just an inclusion of connected components. We call this phenomenon *staticity*.

Definition 3.16. Suppose $F : \operatorname{CAlg}^{\operatorname{cn}} \to S$ and $R \in \operatorname{CAlg}^{\operatorname{cn}}$. Then we say R is *F*-static (or symmetrically F is R-static) if the map $F(R) \to F(\pi_0(R))$ is an inclusion of spaces.⁸ For functors G taking values in spectra, we say R is G-static if it is $\Omega^{\infty}G$ -static.

Example 3.17. Let M be an abelian group. Then every connective commutative ring spectrum R is <u>M</u>-static.

Staticity exhibits closure with respect to many operations.

Lemma 3.18. Let R be a connective commutative ring spectrum.

(1) If $\{F_i\}_{i \in I}$ is a diagram of functors in Fun(CAlg, S) and F_i is R-static for every *i*, then $\lim F_i$ is R-static.

⁸Note that here, an inclusion (of spaces) means a map which is the inclusion of a union of connected components, or equivalently, a (-1)-truncated map.

(2) If $F_0 \to F_1 \to F_2$ is a fiber sequence in Fun(CAlg, Sp^{cn}) with F_0 and F_2 both R-static, then F_1 is R-static.

Proof. The claims follow by using that limits of functors are computed pointwise and (-1)-truncated maps are closed under limits and passage to total spaces of principal fibrations in S.

The behavior with respect to the rings is more subtle, but at least we have the following:

Lemma 3.19. Let $F : \operatorname{CAlg} \to S$ be representable (equivalently, limit preserving) and let $\{R_i\}_{i \in I}$ be a diagram of connective commutative ring spectra. If each of the R_i is F-static and the coassembly map

$$\pi_0 \lim R_i \to \lim \pi_0 R_i$$

is an isomorphism⁹, then $\lim R_i$ is F-static.

Proof. This follows again because (-1)-truncated maps are closed under limits.

We now study staticity in a few special situations.

3.3.1. Staticity is automatic away from the residual characteristic.

Proposition 3.20. Let R be a p-complete commutative ring spectrum and let \mathbb{G} be an affine abelian group scheme such that the multiplication by p map $\mathbb{G} \xrightarrow{p} \mathbb{G}$ is invertible. Then the map $\mathbb{G}(R) \to \mathbb{G}(\pi_0 R)$ is an isomorphism, and in particular R is \mathbb{G} -static.

Proof. Consider the Postnikov tower

$$R \xrightarrow{\sim} \lim \tau_{\leq m} R \to \cdots \to \tau_{\leq 2} R \to \tau_{\leq 1} R \to \tau_{\leq 0} R = \pi_0(R).$$

Since G is an affine abelian group scheme, it is a limit preserving functor, and hence

$$\mathbb{G}(R) \xrightarrow{\sim} \underline{\lim} \mathbb{G}(\tau_{\leq m} R) \quad \in \quad \mathrm{Mod}_{\mathbb{Z}}^{\mathrm{cn}}$$

It therefore suffices to show that the maps $\tau_{\leq m+1}R \to \tau_{\leq m}R$ become isomorphisms after applying \mathbb{G} . Each of these maps fits into a pullback square

where the bottom right corner is the split square-zero extension of $\pi_0(R)$ by $\Sigma^{m+2}\pi_{m+1}R$. Thus, it suffices to show that the map $\pi_0 R \to \pi_0 R \oplus \Sigma^{m+2}\pi_{m+1}R$ induces an isomorphism upon applying \mathbb{G} . Since this map has a left inverse given by the canonical augmentation $\pi_0(R) \oplus \Sigma^{m+2}\pi_{m+1}(R) \to \pi_0(R)$, it remains to show that the fiber of the map

$$\mathbb{G}(\Sigma^{m+2}\pi_{m+1}(R)\oplus\pi_0(R))\to\mathbb{G}(\pi_0(R))$$

vanishes. Let L be the cotangent fiber of \mathbb{G} at its identity element. Then, by definition, we have

$$\operatorname{fib}(\mathbb{G}(\Sigma^{m+2}\pi_{m+1}(R)\oplus\pi_0(R))\to\mathbb{G}(\pi_0(R)))\simeq\operatorname{hom}_{\operatorname{Sp^{cn}}}(L,\Sigma^{m+2}\pi_{m+1}(R))$$

⁹Here, on the right hand side, the limit is computed in CAlg(Sp), i.e., it is the "derived" limit of a diagram of discrete commutative rings.

Since p is invertible on G, it also acts invertibly on L. Since R is p-complete, so is $\Sigma^{m+2}\pi_{m+1}(R)$. We conclude that

$$\hom_{\mathrm{Sp}^{\mathrm{cn}}}(L, \Sigma^{m+2}\pi_{m+1}(R)) \simeq 0$$

and the result follows.

3.3.2. $\mathbb{G}_m^{\text{red}}(-[M])$ -staticity. We now specialize to functors of the form $\mathbb{G}_m^{\text{red}}(-[M])$. The staticity of these functors is compatible with direct sums in M, in the following sense.

Proposition 3.21. Let M and N be abelian groups and let R be a commutative ring spectrum. If R is $\mathbb{G}_m^{\text{red}}(-[M])$ -static and R[M] is $\mathbb{G}_m^{\text{red}}(-[N])$ -static, then R is $\mathbb{G}_m^{\text{red}}(-[M \oplus N])$ -static. Similarly, if $R = R_p^{\wedge}$ is $\mathbb{G}_m^{\text{red}}(-[M]_p^{\wedge})$ -static and $R[M]_p^{\wedge}$ is $\mathbb{G}_m^{\text{red}}(-[N]_p^{\wedge})$ -static, then R is $\mathbb{G}_m^{\text{red}}(-[M \oplus N]_p^{\wedge})$ -static.

Proof. We prove the non *p*-complete version, as the *p*-complete one is analogous. By Observation 3.6 we can identify the morphism

$$\mathbb{G}_m^{\mathrm{red}}(R[M \oplus N]) \to \mathbb{G}_m^{\mathrm{red}}(\pi_0 R[M \oplus N])$$

with the direct sum of the morphisms

$$\mathbb{G}_m^{\mathrm{red}}(R[M]) \to \mathbb{G}_m^{\mathrm{red}}(\pi_0 R[M]) \qquad \text{and} \qquad \mathbb{G}_m^{\mathrm{red}}(R[M][N]) \to \mathbb{G}_m^{\mathrm{red}}(\pi_0 R[M][N]).$$

where in the latter we regard $R[M][N] \simeq R[M \oplus N]$ as an augmented R[M]-algebra. The result now follows from the fact that inclusions of connective spectra are closed under finite direct sums. \Box

Next, we reformulate the notion of rigidity in terms of staticity with respect to $\mathbb{G}_m^{\mathrm{red}}(-[M])$.

Proposition 3.22. Let M be an abelian group and let R be a commutative ring spectrum. Then, R is M-rigid if and only if it satisfies the following two conditions:

- (1) It is $\mathbb{G}_m^{\mathrm{red}}(-[M])$ -static, that is, the map $\mathbb{G}_m^{\mathrm{red}}(R[M]) \to \mathbb{G}_m^{\mathrm{red}}(\pi_0 R[M])$ is an inclusion, and so we can regard the left-hand side as a subgroup of the right-hand side.
- (2) The subgroup

$$\mathbb{G}_m^{\mathrm{red}}(R[M]) \subseteq \mathbb{G}_m^{\mathrm{red}}(\pi_0 R[M])$$

is contained in the image of the map

$$\Theta_{\pi_0 R} \colon \underline{M}(\pi_0 R) \to \mathbb{G}_m^{\mathrm{red}}(\pi_0 R[M]).$$

In other words, the reduced strict units of R[M] map to tautological units of $\pi_0(R[M])$.

Similarly, if R is p-complete, then R is p-completely M-rigid if and only if it is $\mathbb{G}_m^{\text{red}}(-[M]_p^{\wedge})$ -static and the reduced strict units of $R[M]_p^{\wedge}$ map to tautological units of $\pi_0(R[M]_p^{\wedge})$.

Proof. We show the non-*p*-complete case, as the *p*-complete one is proved identically. Consider the commutative square

arising from the naturality of Θ . In this square, the left vertical map is an isomorphism by Example 3.17, and the lower horizontal map is an inclusion of abelian groups.

Now, assume that R is M-rigid. Then, the upper horizontal map Θ_R is an isomorphism, and hence the right vertical map is an inclusion, with image contained in that of $\Theta_{\pi_0 R}$, as we wanted to show. Conversely, if R satisfies condition (1) then, since inclusions of spaces satisfy cancellation from the right, this forces the upper vertical map to be an inclusion as well. Now, by the commutativity and the fact that the left vertical map is an isomorphism, condition (2) forces it to be also surjective on connected components. Together, this shows that the upper horizontal map Θ_R is an isomorphism, so that R is M-rigid.

4. Rigidity for finite p-groups

In this section, we will show that many commutative ring spectra are p-rigid (i.e., rigid for every finite abelian p-group).

Example 4.1. Consider the following failures of *p*-rigidity for discrete *R*:

- (1) If M is a finite group of exponent p^e and R is a $\mathbb{Z}[1/p]$ -algebra with a primitive p^e -th root of unity, then there is a discrete Fourier isomorphism $R[M] \cong R^{M^*}$ of algebras. This gives an isomorphism $\mathbb{G}_m^{\text{red}}(R[M]) \simeq \mathbb{G}_m(R)^{M^* \setminus \{0\}}$, which is usually larger than M.
- (2) *p*-rigidity usually fails when R is an \mathbb{F}_p -algebra. For example, $\mathbb{F}_p[C_p] \simeq \mathbb{F}_p[t]/t^p$, and the (automatically strict) reduced units in this ring contain all the elements of the form $1 + a_1t + \cdots + a_{p-1}t^{p-1}$ for arbitrary $a_i \in \mathbb{F}_p$.

One can view these classical obstructions through the lens of chromatic homotopy theory:

- (1) originates from the existence of primitive *p*-th roots of unity, a phenomenon that occurs only when *p* is invertible (i.e., in chromatic height 0), cf. [Dev20]. We shall thus restrict our attention to *p*-complete commutative ring spectra.
- (2) is related to \mathbb{F}_p -linearity: any \mathbb{F}_p -algebra has vanishing T(n)-localizations for $n \geq 0$ and is therefore "concentrated in chromatic height ∞ ". We will therefore work with an appropriate category of commutative ring spectra with "no height ∞ part."

It turns out that the above obstructions are essentially the only ones. Most of this section will be dedicated to showing that T(n)-local ring spectra are *p*-rigid (Theorem 4.11). Since this condition is closed under limits, any iterated limit of T(n)-local rings will be *p*-rigid. In particular, we will see that this holds for the *p*-complete sphere \mathbb{S}_p and any algebra of finite Tor-amplitude over it, by chromatic convergence (§4.2).

4.1. T(n)-local rigidity. Throughout this section we fix a prime p and we consider the T(n) localization of p-local spectra for $n \ge 1$, so that all spectra to which we apply this localization are implicitly assumed to be p-local. The aim of this section is to show that every T(n)-local commutative ring spectrum R is p-rigid (Theorem 4.11), i.e. that the map

$$\Theta_M(R) \colon \underline{M}(R) \to \mathbb{G}_m^{\mathrm{red}}(R[M])$$

is an isomorphism for M any finite abelian p-group.

When n = 0, we understood the T(0)-local (i.e. rational) failure of *p*-rigidity in Example 4.1(2) using a Fourier isomorphism $R[M] \simeq R^{M^*}$. Since the right-hand side is a product of rings, its units are easy to compute.

In the case n > 0, there is an analogous isomorphism known as the *chromatic Fourier transform*, which identifies R[M] as cochains on the classifying space $B^n M^*$. We review this in §4.1.1 and use it to identify the target of

$$\Theta_M : \underline{M} \to \mathbb{G}_m^{\mathrm{red}}(-[M])$$

in §4.1.2. More precisely, for each $r \ge 0$, we will show that if we restrict the target to \mathbb{Z}/p^r -modules M, it can be identified with the functor \underline{M} . Then, we finish by using this to show that Θ_M induces an isomorphism in §4.1.3.

Remark 4.2. Let us indicate why the n = 0 case behaves qualitatively differently. The issue is that in the equivalence $R[M] \simeq R^{M^*}$, the right-hand side depends, as a ring, only on the number of elements of M but not on the group structure. This means that $\mathbb{G}_m^{\text{red}}(R[M])$ depends only on the number of elements of M whereas $\underline{M}(R)$ is clearly sensitive to the group structure of M and so

$$\Theta_M(R) : \underline{M}(R) \to \mathbb{G}_m^{\mathrm{red}}(R[M])$$

cannot be an isomorphism in general. At heights n > 0, the chromatic Fourier transform identifies R[M] with $R^{B^n M^*}$, which, in contrast, *does* depend on the group structure on M^* .

4.1.1. Chromatic Fourier transform. We start by briefly recalling the chromatic Fourier transform from [CSY21, BCSY22]. For a T(n)-local commutative ring spectrum R, a higher root of unity of height n and order p^r in R is a map

$$\omega \colon \mathbb{S}[\Sigma^n \mathbb{Z}/p^r] \to R$$

of commutative ring spectra. We denote the space of such higher roots of unity by $\mu_{p^r}^{(n)}(R)$. Each $\omega \in \mu_{p^r}^{(n)}(R)$ determines a natural transformation

$$\mathcal{F}_{\omega}(M) \colon R[M] \to R^{\Omega^{\infty} \Sigma^n M^*} \in \operatorname{CAlg}_R(\operatorname{Sp}_{T(n)})$$

of functors of M, a π -finite connective \mathbb{Z}/p^r -module spectrum, where

 $M^* := \hom_{\mathbb{Z}}(M, \mathbb{Q}/\mathbb{Z}) \simeq \hom_{\mathbb{Z}/p^r}(M, \mathbb{Z}/p^r) \quad \in \quad \operatorname{Mod}_{\mathbb{Z}/p^r}$

denotes the Pontryagin dual of M. We refer to $\mathcal{F}_{\omega}(M)$ as the chromatic Fourier transform.

The natural quotient maps $\mathbb{Z}/p^r \to \mathbb{Z}/p^{r-1}$ induce maps $\mu_{p^{r-1}}^{(n)}(R) \to \mu_{p^r}^{(n)}(R)$ corresponding to the "inclusion of p^{r-1} -roots of unity." A higher root of unity $\omega \in \mu_{p^r}^{(n)}(R)$ is called *primitive* if the base change to $\mu_{p^{r-1}}^{(n)}(R')$ for any non-zero, T(n)-local *R*-algebra R' is not in the image of the corresponding map $\mu_{p^{r-1}}^{(n)}(R') \to \mu_{p^r}^{(n)}(R')$. For primitive roots, the Fourier transform is an isomorphism (i.e., one has a "Fourier inversion"):

Theorem 4.3 (Chromatic Fourier Transform [BCSY22, Theorem A]). Let $n \ge 0$ and let $R \in CAlg(Sp_{T(n)})$. If $\omega \in \mu_{p^r}^{(n)}(R)$ is primitive, then the Fourier transform

$$\mathcal{F}_{\omega} \colon R[M] \xrightarrow{\sim} R^{\Omega^{\infty - n} M^*} \in \operatorname{CAlg}(\operatorname{Sp}_{T(n)})$$

is an isomorphism for every π -finite connective \mathbb{Z}/p^r -module spectrum M.

While not all commutative T(n)-local algebras have primitive roots of order p^r , they do after passing to a finite Galois extension. In fact, primitive higher roots of unity are classified by a finite Galois extension of the T(n)-local sphere $\mathbb{S}_{T(n)}$.

Theorem 4.4 (Higher Cyclotomic Extensions, [CSY21, Proposition 5.2]). The functor that takes $R \in \text{CAlg}(\text{Sp}_{T(n)})$ to the subspace of $\mu_{p^r}^{(n)}(R)$ consisting of primitive roots is co-representable by a T(n)-local \mathbb{Z}/p^r -Galois extension $R_{n,r}^f$ of $\mathbb{S}_{T(n)}$.

These Galois extensions $R_{n,r}^f$ are referred to in [CSY21] as the *chromatic cyclotomic extensions*. The upshot for us will be that, provided we work over $R_{n,r}^f$, there is a natural Fourier isomorphism: **Corollary 4.5.** There is a natural isomorphism of commutative and cocommutative Hopf algebras in $\operatorname{Sp}_{T(n)}$, defined for $R \in \operatorname{CAlg}_{R_n^f}(\operatorname{Sp}_{T(n)})$ and $M \in \operatorname{Mod}_{\mathbb{Z}/p^r}^{\operatorname{cn}}$ with π_*M finite, of the form

$$\mathcal{F}\colon R[M] \xrightarrow{\sim} R^{\Omega^{\infty - n}M^*}$$

Proof. Let $\omega \in \mu_{p^r}^{(n)}(R_{n,r}^f)$ be the universal primitive p^r -th root of unity of height n, and let $\mathcal{F} = \mathcal{F}_{\omega}$. For the Hopf algebra structure, see [BCSY22, §3.3].

Implicit in Corollary 4.5 is that if M is p-power torsion and n-connected, then $R[M] \simeq R$. We will need a torsion-free variant of this fact:

Lemma 4.6. Suppose R is a T(n)-local commutative ring spectrum and suppose j > n + 1. Then the unit map $R \to L_{T(n)}R[\Sigma^j\mathbb{Z}]$ is an isomorphism. In particular, the connective \mathbb{Z} -module $\mathbb{G}_m(R)$ is (n + 1)-truncated.

Proof. The first statement follows from the formula

$$L_{T(n)}R[\Sigma^{j}\mathbb{Z}] = L_{T(n)} \lim_{n \to \infty} R[\Sigma^{j-1}\mathbb{Z}/p^{r}]$$

and the aforementioned fact about R[M] for M p-power torsion and n-connected (cf. also [CSY20, Lemma 2.4.4, Proposition 3.2.1], for instance). The "in particular" follows from considering R-algebra maps into R.

For our applications, we will need one additional fact about the $R_{n,r}^f$:

Lemma 4.7. For $n, r \ge 0$, the commutative ring spectrum $R_{n,r}^f$ is Zariski-connected.

Proof. Suppose $R_{n,r}^f = R_1 \times R_2$ for nonzero rings R_1, R_2 . Let E_n denote a Lubin-Tate theory of height n and consider $L_{K(n)}(R_{n,r}^f \otimes E_n)$ with its action of the Morava stabilizer group \mathbb{G}_n . By [CSY21, Propositions 5.2, 5.4, 5.5], this has Zariski spectrum isomorphic to $(\mathbb{Z}/p^r)^{\times}$, acted on transitively by \mathbb{G}_n through the "higher cyclotomic character."

On the other hand, by the Devinatz-Hopkins-Smith Nilpotence theorem (cf. [CSY18, Corollary 5.1.17]), the functor $L_{K(n)}(-\otimes E_n)$: $\operatorname{Sp}_{T(n)} \to \operatorname{Sp}_{T(n)}$ is nil-conservative, i.e., it does not send any rings to zero. Thus, $L_{K(n)}(R_{n,r}^f \otimes R_i)$ is nonzero for i = 1, 2, and so the Zariski spectrum of $L_{K(n)}(R_{n,r}^f \otimes E_n)$ splits into at least two nontrivial \mathbb{G}_n -orbits, which is a contradiction.

4.1.2. Identifying the Target of Θ . Recall that our goal is to show that $\Theta_M(R)$ is an isomorphism for every $R \in \operatorname{CAlg}(\operatorname{Sp}_{T(n)})$ and finite abelian *p*-group *M*. In this section, we use the chromatic Fourier transform to abstractly identify $\mathbb{G}_m^{\operatorname{red}}(R[M])$ with $\underline{M}(R)$ under the additional assumption that *R* admits a primitive p^r -th root of unity of height *n* and *M* is p^r -torsion. Later, we will show that the existence of such natural isomorphisms forces every T(n)-local commutative ring spectrum to be *p*-rigid.

Proposition 4.8. There is an isomorphism of functors

 $\Psi_M: \mathbb{G}_m^{\mathrm{red}}(-[M]) \xrightarrow{\sim} \underline{M} \quad \in \quad \mathrm{Shv}_{\mathrm{zar}}(R^f_{n,r}; \mathrm{Mod}_{\mathbb{Z}}^{\mathrm{cn}})$

which is natural in the finite discrete \mathbb{Z}/p^r -module M. In particular, for every commutative $R_{n,r}^f$ algebra R and every such M, there is an isomorphism

$$\mathbb{G}_m^{\mathrm{red}}(R[M]) \simeq \underline{M}(R).$$

Proof. By Corollary 4.5, we have for $R \in \operatorname{CAlg}_{R_{n,r}^f}(\operatorname{Sp}_{T(n)})$ and M a finite discrete \mathbb{Z}/p^r -module a natural equivalence of T(n)-local commutative and cocommutative Hopf algebras

$$\mathcal{F}\colon R[M] \xrightarrow{\sim} R^{\Omega^{\infty - n}M^*} \simeq R^{B^n M^*}.$$

This yields natural equivalences of connective spectra:

$$\mathbb{G}_{m}^{\mathrm{red}}(R[M]) := \mathrm{fib}(\mathbb{G}_{m}(R[M]) \to \mathbb{G}_{m}(R)) \\
\simeq \mathrm{fib}(\mathbb{G}_{m}(R^{B^{n}M^{*}}) \to \mathbb{G}_{m}(R)) \\
\simeq \mathrm{fib}(\mathbb{G}_{m}(R)^{B^{n}M^{*}} \to \mathbb{G}_{m}(R)) \\
\simeq \mathrm{Map}_{\mathcal{S}_{*}}(B^{n}M^{*}, \mathbb{G}_{m}(R)),$$

where the last space is the space of pointed maps $B^n M^* \to \mathbb{G}_m(R)$. Now, $\mathbb{G}_m(R)$ is a connective, (n+1)-truncated \mathbb{Z} -module by Lemma 4.6, and hence there is a natural equivalence

$$\operatorname{Map}_{\mathcal{S}_*}(B^n M^*, \mathbb{G}_m(R)) \simeq \operatorname{Map}_{\operatorname{Mod}_{\mathbb{Z}}}(\tau_{\leq n+1} C_*(B^n M^*; \mathbb{Z}), \mathbb{G}_m(R)),$$

where $\widetilde{C}_*(B^n M^*; \mathbb{Z})$ is the \mathbb{Z} -module of *reduced* integral homology chains of $B^n M^*$ (that is, $\mathbb{Z} \otimes \Sigma^{\infty} B^n M^*$). Finally, by a classical computation of the homology of Eilenberg–MacLane spaces, we see that the counit map

$$\widetilde{C}_*(B^n M^*; \mathbb{Z}) \to \Sigma^n M^* \in \operatorname{Mod}_{\mathbb{Z}}^{\operatorname{cn}}$$

induces an isomorphism in homology up to degree n + 1, and hence an isomorphism of the truncations:

$$\tau_{\leq n+1} \tilde{C}_* (B^n M^*; \mathbb{Z}) \simeq \Sigma^n M^* \quad \in \quad \operatorname{Mod}_{\mathbb{Z}}^{\operatorname{cn}}$$

Combining the above, we obtain a natural equivalence

$$\mathbb{G}_m^{\mathrm{red}}(R[M]) \simeq \mathrm{Hom}_{\mathrm{Mod}_{\mathbb{Z}}^{\mathrm{cn}}}(\Sigma^n M^*, \mathbb{G}_m(R)) = \mathrm{Hom}_{\mathrm{CAlg}_{R_{n,r}^f}(\mathrm{Sp}_{T(n)})}(R_{n,r}^f[\Sigma^n M^*], R).$$

Finally, the (inverse) Fourier transform gives an isomorphism

$$\mathcal{F}\colon R^f_{n,r}[\Sigma^n M^*] \xrightarrow{\sim} (R^f_{n,r})^{\Omega^{\infty} M}$$


of commutative and cocommutative Hopf algebras in $\operatorname{Sp}_{T(n)}$ and so we have an equivalence

$$\mathbb{G}_m^{\mathrm{red}}(-[M]) \simeq \underline{M} \in \operatorname{Shv}_{\mathrm{zar}}(R_{n,r}^f; \operatorname{Sp}^{\mathrm{cn}})$$

of Zariski sheaves of connective spectra over $R_{n,r}^f$. Since <u>M</u> (and hence the left-hand side) is discrete and p^r -torsion, this isomorphism admits a unique \mathbb{Z}/p^r -linear structure, and the result follows.

Remark 4.9. The reason that we work with each finite r rather than assuming R has all p^r -th height n roots of unity is that the argument in Section 4.1.3 is by Galois descent. While one has descent from each $R_{n,r}^f$, descent for the maximal higher cyclotomic extension $\varinjlim_r R_{n,r}^f$ fails in heights $n \geq 2$, as shown by Burklund, Hahn, Levy, and Schlank in [BHLS23, §6.4].

On the other hand, if one is satisfied with the K(n)-local version of Theorem 4.11 (which suffices for our applications), one can simply use descent from Lubin-Tate theory where the required Fourier transform is constructed by Hopkins and Lurie and shown to be an isomorphism for all *p*-local π -finite M, not just \mathbb{Z}/p^r -modules for a fixed r (see [HL13b, Corollary 5.3.26]). 4.1.3. Θ is a T(n)-local equivalence. We would like to show that $\Theta_M : \underline{M} \to \mathbb{G}_m^{\mathrm{red}}(-[M])$ is an isomorphism for any finite abelian *p*-group *M*. It suffices to check this in the case when *M* is a \mathbb{Z}/p^r -module, for every $r \geq 0$. Naturally in a finite discrete \mathbb{Z}/p^r -module *M*, we have a diagram of $\mathrm{Mod}_{\mathbb{Z}}^n$ -valued functors



where $\widetilde{\Theta}_M$ denotes the composite $\Psi_M \circ \Theta_M$. By Proposition 4.8, Ψ_M is an isomorphism and thus showing Θ_M is an isomorphism amounts to showing that $\widetilde{\Theta}_M$ is an isomorphism. Note that both the source and target of $\widetilde{\Theta}_M$ are discrete and p^r -torsion, so we may regard $\widetilde{\Theta}_M$ as a natural morphism of sheaves of ordinary \mathbb{Z}/p^r -modules.

Proposition 4.10. For every $r \ge 0$ and every finite \mathbb{Z}/p^r -module M, the morphism

$$\widetilde{\Theta}_M : \underline{M} \to \underline{M} \in \operatorname{Shv}_{\operatorname{zar}}(R_{n,r}^f; \operatorname{Mod}_{\mathbb{Z}}^{\operatorname{cn}})$$

is an equivalence.

Proof. Since (-) is left adjoint to the evaluation at $R_{n,r}^f$, the natural transformation $\tilde{\Theta}_M$ corresponds to a morphism $M \to \underline{M}(R_{n,r}^f)$ of \mathbb{Z}/p^r -modules. Since $R_{n,r}^f$ is connected by Lemma 4.7, this can be identified with a natural transformation $\theta_M : M \to M$ (for instance given by evaluating $\tilde{\Theta}_M$ at $R_{n,r}^f$), and it suffices to show that θ_M is an isomorphism for all M.

Since θ_M is a natural endomorphism of the identity functor on the category of finite discrete \mathbb{Z}/p^r modules, it is given by multiplication by some scalar $a \in \mathbb{Z}/p^r$. It suffices to show that a is invertible. For this, one only needs to show that a is invertible mod p, or equivalently, that $\theta_{C_p} : C_p \to C_p$ is invertible. But θ_{C_p} is a \mathbb{Z}/p -module map, so this amounts to showing θ_{C_p} is non-zero.

Up to isomorphisms, θ_{C_p} is identified with $\Theta_{C_p}(R_{n,r}^f)$, or even $\Theta_{C_p}(\mathbb{S})$. This is non-constant by definition, so θ_{C_p} cannot be zero.

We can now prove the main result of this section.

Theorem 4.11. For $n \ge 1$, every T(n)-local commutative ring spectrum is p-rigid.

Proof. Let R be a T(n)-local commutative ring spectrum and let M be a finite abelian p-group. We wish to show that R is M-rigid. Choose r > 0 for which M is p^r -torsion and let $R \otimes R_{n,r}^f$ be the T(n)-local tensor product of R and $R_{n,r}^f$. By Proposition 4.10, we have that $R \otimes R_{n,r}^f$ is M-rigid. But the unit map $R \to R \otimes R_{n,r}^f$ is a faithful $(\mathbb{Z}/p^r)^{\times}$ -Galois extension in $\operatorname{CAlg}(\operatorname{Sp}_{T(n)})$, and hence

$$R \simeq (R \widehat{\otimes} R_{n,r}^f)^{h(\mathbb{Z}/p^r)^{\times}} \quad \in \quad \operatorname{CAlg}(\operatorname{Sp}_{T(n)}).$$

Since the embedding $\operatorname{CAlg}(\operatorname{Sp}_{T(n)}) \to \operatorname{CAlg}(\operatorname{Sp})$ is limit preserving, the same formula holds in $\operatorname{CAlg}(\operatorname{Sp})$. Thus, since *p*-rigid rings are closed under limits, this implies that *R* is *M*-rigid as well and the result follows.

4.2. Chromatic convergence and *p*-rigidity over \mathbb{S}_p^{\wedge} . Since *p*-rigidity is closed under limits in CAlg(Sp), we immediately deduce from Theorem 4.11 that:

Theorem 4.12. Let $\mathcal{K} \subset \text{Sp}$ be the smallest subcategory which is closed under limits and which contains $\text{Sp}_{T(n)}$ for any n > 0. Then any $R \in \text{CAlg}(\text{Sp})$ whose underlying spectrum is in \mathcal{K} is *p*-rigid.

The class \mathcal{K} contains many spectra of interest.

Corollary 4.13. Suppose $R \in CAlg(Sp)$ is p-complete and satisfies any of the following conditions:

- (1) R is L_n^f -local for some n > 0.
- (2) R/p is chromatically convergent (more generally, a retract of its chromatic tower).
- (3) R is a \mathbb{S}_p -algebra of finite Tor-amplitude (cf. Definition 2.35).

Then R is p-rigid, i.e., there is a canonical isomorphism

$$\mathbb{G}_m(R[M]) \simeq \mathbb{G}_m(R) \oplus \underline{M}(R).$$

for any finite abelian p-group M.

Proof. Claim (1) follows by induction, starting from the case n = 1 (note that L_1^f -local and T(1)-local are the same for *p*-complete spectra) and using the chromatic fracture squares

Claim (2) follows immediately because $\lim L_n R$ is in \mathcal{K} by limit closure (note that $L_n R$ is L_n^f local) and the hypothesis implies $R \to \lim L_n R$ is a mod p isomorphism of p-complete spectra and therefore an isomorphism. The statement about retracts follows because retracts are limits.

Finally, any R satisfying (3) has a finite filtration with quotients of the form $\Sigma^k(\bigoplus_{i \in I} \mathbb{S}_p)_p^{\wedge}$. Thus, by (2), it suffices to see that any spectrum of the form $\Sigma^k(\bigoplus_{i \in I} \mathbb{S}/p)_p^{\wedge}$ is chromatically convergent. Since \mathbb{S}/p is a finite spectrum, it has finite projective BP-dimension in the sense of [Bar16, Definition 2.1] (see, e.g., [Bar16, Proposition 2.4]). Since direct sums of projective resolutions are projective resolutions, this implies that any sum of copies of \mathbb{S}/p also has finite projective BP-dimension, and hence is chromatically complete by [Bar16, Theorem 3.8]¹⁰.

5. RIGIDITY FOR FREE ABELIAN GROUPS

In the previous section, we gave conditions for the maps $\Theta_M(R) \colon \underline{M}(R) \to \mathbb{G}_m^{\mathrm{red}}(R[M])$ to be isomorphisms in the case that M is a finite abelian p-group — this we called p-rigidity. In this section, we give analogous results when $M \simeq \mathbb{Z}^k$ is a free abelian group of finite rank, for a much more restrictive class of commutative ring spectra R. In fact, our main result is a statement just for spherical Witt vectors (though we prove more specific statements along the way).

 $^{^{10}}$ One can also argue directly using that the chromatic convergence on S is pro-constant on each homotopy group.

Theorem 5.1 (Theorem 5.18). Let κ be a perfect ring of characteristic p and let Λ be a free abelian group of finite rank. Then the ring of spherical Witt vectors $\mathbb{S}_{W(\kappa)}$ is p-completely Λ -rigid.¹¹

Proving rigidity for free abelian groups presents new difficulties: first of all, the functor $\mathbb{G}_m^{\text{red}}(-[\Lambda])$ is no longer limit preserving when Λ is infinite, and second, an analogue of the chromatic Fourier transform for infinite abelian groups is not known. We therefore take a different approach: we access the \mathbb{Z} -module spectrum $\mathbb{G}_m^{\text{red}}(R[\Lambda])$ via the quotient map $\mathbb{G}_m^{\text{red}}(R[\Lambda]) \to \mathbb{G}_m^{\text{red}}(R[\Lambda/p])$. Indeed, while Λ is infinite, the group Λ/p is a finite abelian *p*-group and therefore fits into the framework of the previous section. We summarize our strategy as follows:

(1) In §5.1, we show that when R is p-complete of finite Tor-amplitude, the short exact sequence $\Lambda \xrightarrow{[p]} \Lambda \to \Lambda/p$ induces a cofiber sequence

$$\mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge}) \xrightarrow{|p|_*} \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge}) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[\Lambda/p]) \ .$$

- (2) We then show in §5.2 that the multiplication by p map of the \mathbb{Z} -module $\mathbb{G}_m^{\text{red}}(R[\Lambda]_p^{\wedge})$ can be written as the composite of the map $[p]_*$ and the map induced by the Frobenius of R. Thus, when R is *perfect*, that is, has invertible Frobenius, the map $[p]_*$ can be replaced by the multiplication by p map in the above cofiber sequence.
- (3) Finally, in §5.3, we use the above exact sequence and our results on *p*-rigidity to show that any perfect connective R is $\mathbb{G}_m^{\text{red}}(-[\Lambda]_p^{\wedge})$ -static, in the sense of Definition 3.16. In view of Proposition 3.22, this reduces our question to an algebraic problem, which we solve using our analysis of rank 1 units in δ_p -rings from §2, completing the proof.

5.1. Exactness of reduced strict units. We begin by carrying out part (1) of our strategy. More generally, given an exact sequence of abelian groups

$$0 \longrightarrow M \longrightarrow N \longrightarrow L \longrightarrow 0 ,$$

we may ask whether the induced sequence

$$\mathbb{G}_m^{\mathrm{red}}(R[M]_p^\wedge) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[N]_p^\wedge) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[L]_p^\wedge)$$

is a (co)fiber sequence in $\operatorname{Mod}_{\mathbb{Z}}^{\operatorname{cn}}$. This will turn out to be true under mild hypotheses involving *L*-rigidity (Proposition 5.3). We start by analyzing the special case of a *split* exact sequence.

Proposition 5.2. Let R be a commutative ring spectrum and let M and N be abelian groups. If $R[M]_p^{\wedge}$ is p-completely N-rigid and the map $R \to R[M]_p^{\wedge}$ induces a bijection $\pi_0 \operatorname{Spec}(R[M]_p^{\wedge}) \simeq \pi_0 \operatorname{Spec}(R)$, then the assembly map

$$\mathbb{G}_m^{\mathrm{red}}(R[M]_p^{\wedge}) \oplus \mathbb{G}_m^{\mathrm{red}}(R[N]_p^{\wedge}) \to \mathbb{G}_m^{\mathrm{red}}(R[M \oplus N]_p^{\wedge})$$

is an isomorphism. Equivalently, the split exact sequence

 $0 \longrightarrow M \longrightarrow M \oplus N \longrightarrow N \longrightarrow 0 ,$

is taken by $\mathbb{G}_m^{\mathrm{red}}(R[-]_p^{\wedge})$ to a split cofiber sequence.

¹¹We work with *p*-complete rigidity (cf. Variant 3.13) because for Λ infinite, the spectrum $R[\Lambda]$ may not be *p*-complete even if R is.

Proof. By Observation 3.6, the assembly map is identified with the map

$$\mathbb{G}_m^{\mathrm{red}}(R[M]_p^{\wedge}) \oplus \mathbb{G}_m^{\mathrm{red}}(R[N]_p^{\wedge}) \to \mathbb{G}_m^{\mathrm{red}}(R[M]_p^{\wedge}) \oplus \mathbb{G}_m^{\mathrm{red}}(R[M][N]_p^{\wedge})$$

given by the sum of the identity of $\mathbb{G}_m^{\mathrm{red}}(R[M]_p^{\wedge})$ and the map

(*)
$$\mathbb{G}_m^{\mathrm{red}}(R[N]_p^{\wedge}) \to \mathbb{G}_m^{\mathrm{red}}(R[M][N]_p^{\wedge})$$

induced by applying $\mathbb{G}_m^{\text{red}}((-)[N]_p^{\wedge})$ to the unit map $R \to R[M]_p^{\wedge}$. It therefore suffices to show that (*) is an isomorphism. Since R is a retract of $R[M]_p^{\wedge}$, the assumption that $R[M]_p^{\wedge}$ is pcompletely N-rigid implies that R is p-completely N-rigid as well. Thus, we can identify (*) with the corresponding map

$$\underline{N}(R) \to \underline{N}(R[M]_p^{\wedge}).$$

But this is an isomorphism by the assumption that $\pi_0 \operatorname{Spec}(R[M]_p^{\wedge}) \simeq \pi_0 \operatorname{Spec}(R)$.

Consider now a general exact sequence

$$0 \longrightarrow M \xrightarrow{\alpha} N \xrightarrow{\beta} L \longrightarrow 0$$

There is a standard procedure to approximate it by split exact sequences. Namely, by forming the pushout along the map α , we obtain a morphism of exact sequences of the form

in which the bottom sequence is *split* exact¹² and the vertical maps are all injections. The left square in the diagram is the first part of the Čech nerve of the map α , as defined in Definition 3.7. In fact, using the descent result of Proposition 3.8, we can reduce the exactness question for $\mathbb{G}_m^{\text{red}}(R[-]_p^{\wedge})$ to the split case, and obtain the following result.

Proposition 5.3. Let R be a p-complete commutative ring spectrum and let

$$0 \longrightarrow M \xrightarrow{\alpha} N \xrightarrow{\beta} L \longrightarrow 0$$

be an exact sequence of abelian groups. Assume that for every $\ell \in \mathbb{N}$:

- $R[N \oplus L^{\oplus \ell}]_p^{\wedge}$ is L-rigid.
- The unit map $R \to R[N \oplus L^{\oplus \ell}]_p^{\wedge}$ induces a bijection

$$\pi_0 \operatorname{Spec}(R[N \oplus L^{\oplus \ell}]_p^{\wedge}) \simeq \pi_0 \operatorname{Spec}(R).$$

Then, the null sequence

$$(*) \qquad \qquad \mathbb{G}_m^{\mathrm{red}}(R[M]_p^{\wedge}) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[N]_p^{\wedge}) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[L]_p^{\wedge})$$

is a cofiber sequence in $Mod_{\mathbb{Z}}^{cn}$.

¹²A splitting is given by sending $\ell \in L$ to $(n, -n) \in N \oplus_M N$ for any lift $\beta(n) = \ell$.

Proof. We start by showing the weaker claim that this sequence is a *fiber* sequence in $\operatorname{Mod}_{\mathbb{Z}}^{\operatorname{cn}}$. Consider the Čech nerve functor $C^{\bullet} \colon (\operatorname{Mod}_{\mathbb{Z}})^{\Delta^1} \to \operatorname{Mod}_{\mathbb{Z}}^{\Delta}$ from Definition 3.7. Applying the composition

$$\mathbb{G}_m^{\mathrm{red}}(R[C^{\bullet}(-)]_p^{\wedge}) \colon (\mathrm{Mod}_{\mathbb{Z}}^{\mathrm{cn}})^{\Delta^1} \xrightarrow{C^{\bullet}} (\mathrm{Mod}_{\mathbb{Z}}^{\mathrm{cn}})^{\Delta} \xrightarrow{\mathbb{G}_m^{\mathrm{red}}(R[-])} (\mathrm{Mod}_{\mathbb{Z}}^{\mathrm{cn}})^{\Delta}$$

to the exact sequence of morphisms of abelian groups

$$0 \longrightarrow M \xrightarrow{\alpha} N \xrightarrow{\beta} L \longrightarrow 0$$
$$\downarrow^{\alpha} \qquad \downarrow^{\delta_0} \qquad \parallel$$
$$0 \longrightarrow N \xrightarrow{\delta_1} N \oplus_M N \xrightarrow{\beta'} L \longrightarrow 0$$

from above, we obtain a null sequence in $(Mod_{\mathbb{Z}}^{cn})^{\Delta}$

$$(**) \qquad \qquad \mathbb{G}_m^{\mathrm{red}}(R[C^{\bullet}(\alpha)]_p^{\wedge}) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[C^{\bullet}(\delta_0)]_p^{\wedge}) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[C^{\bullet}(\mathrm{id}_L)]_p^{\wedge}).$$

Since the three maps α , δ_0 and id_L are injections, we deduce from Proposition 3.8 that the sequence (*) from the statement of the proposition is the limit of the sequences (**) over Δ . Since the functor

$$\lim_{\Delta} \colon (\mathrm{Mod}^{\mathrm{cn}}_{\mathbb{Z}})^{\Delta} \to \mathrm{Mod}^{\mathrm{cn}}_{\mathbb{Z}}$$

is limit preserving (and in particular preserves fiber sequences), it suffices to show that for every $\ell \in \mathbb{N}$ the sequence

$$(*_{\ell}) \qquad \qquad \mathbb{G}_m^{\mathrm{red}}(R[C^{\ell}(\alpha)]_p^{\wedge}) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[C^{\ell}(\delta_0)]_p^{\wedge}) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[C^{\ell}(\mathrm{id}_L)]_p^{\wedge})$$

is a fiber sequence in $Mod_{\mathbb{Z}}^{cn}$. We do this by verifying the conditions of Proposition 5.2: note that the exact sequences

$$0 \longrightarrow C^{\ell}(\alpha) \longrightarrow C^{\ell}(\delta_0) \longrightarrow C^{\ell}(\mathrm{id}_L) \longrightarrow 0$$

are split, as they are pushouts of the split exact sequence

$$0 \longrightarrow N \longrightarrow N \oplus_M N \longrightarrow L \longrightarrow 0$$

along the injections $N \to C^{\ell}(\alpha)$. In particular, this means that for $\ell = 0, 1, \ldots$, we obtain that $C^{\ell}(\delta_0) \simeq N \oplus L^{\oplus \ell}$, and we clearly have $C^{\ell}(\mathrm{id}_L) \simeq L$. Therefore, our hypotheses exactly check the conditions of Proposition 5.2. It follows that $(*_{\ell})$ are fiber sequences, and so (*) is as well.

To show that (*) is in fact a cofiber sequence, observe that by our assumption that $R[N]_p^{\wedge}$ is *p*-completely *L*-rigid, the same holds for *R* and hence we have an isomorphism

$$\mathbb{G}_m^{\mathrm{red}}(R[L]_p^\wedge) \simeq \underline{L}(R)$$

fitting into a commutative square

$$\underbrace{M(R) \longrightarrow \underline{L}(R)}_{\substack{\downarrow \widehat{\Theta}_{M} \\ \downarrow \widehat{\Theta}_{M} \\ \pi_{0} \mathbb{G}_{m}^{\mathrm{red}}(R[M]_{p}^{\wedge}) \longrightarrow \pi_{0} \mathbb{G}_{m}^{\mathrm{red}}(R[L]_{p}^{\wedge})}_{p}$$

Since the upper map is surjective, we deduce that the same holds for the bottom map, and hence that (*) is a cofiber sequence in $Mod_{\mathbb{Z}}^{cn}$.

We will be particularly interested in Proposition 5.3 for the exact sequence

$$0 \longrightarrow \Lambda \xrightarrow{[p]} \Lambda \longrightarrow \Lambda/p \longrightarrow 0$$

To check the conditions of the proposition, we need the following:

Lemma 5.4. Let R be a p-complete commutative ring spectrum and let M be a finitely generated abelian group which contains no elements of order prime to p. Then, the unit map $R \to R[M]_p^{\wedge}$ induces a bijection

$$\pi_0 \operatorname{Spec}(R[M]_n^{\wedge}) \simeq \pi_0 \operatorname{Spec}(R)$$

Proof. As both sides depend only on π_0 , we can immediately reduce to R discrete. Moreover, the rings are derived p-complete, so we can in fact reduce modulo p and further assume that R is a ring of characteristic p (in which case we can drop the extra p-completion).

Write $M = M' \oplus \mathbb{Z}^k$ where M' is a finite *p*-group. Then R[M] is a nil-thickening of $R[\mathbb{Z}^k]$, so it suffices to consider the case M' = 0, and in fact we can assume k = 1 and R is reduced. But here, one sees by inspection that any idempotent in $R[t^{\pm 1}]$ is inside R, and the conclusion follows. \Box

We are ready to prove the main result of this subsection:

Proposition 5.5. Let R be a p-complete commutative ring spectrum of finite Tor-amplitude, let Λ be a free abelian group of finite rank (i.e. $\Lambda \simeq \mathbb{Z}^k$), and let $[p] \colon \Lambda \to \Lambda$ denote the multiplication by p map. Then, the null sequence

$$\mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge}) \xrightarrow{[p]_*} \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge}) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[\Lambda/p])$$

is a cofiber sequence.

Proof. In this case, $R[\Lambda \oplus (\Lambda/p)^{\oplus \ell}]_p^{\wedge}$ is also of finite Tor-amplitude and hence *p*-rigid by Corollary 4.13. In particular, it is *p*-completely Λ/p -rigid. The result now follows by combining this with Lemma 5.4 and Proposition 5.3.


5.2. The Frobenius on strict units of group algebras. We turn to (2) in our strategy; that is, we relate the map induced by $[p]: \Lambda \to \Lambda$ to the multiplication by p map on the spectrum $\mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge})$. More specifically, for R p-complete of finite Tor-amplitude, we show:

- The Frobenius map (cf. Section 2.4) of the group algebra $R[\Lambda]_p^{\wedge}$ ("absolute Frobenius") is the composite of $[p]_*: R[\Lambda]_p^{\wedge} \to R[\Lambda]_p^{\wedge}$ ("geometric Frobenius") and the Frobenius φ_R of R("arithmetic Frobenius", cf. §2.4).
- The absolute Frobenius map $\varphi_{R[\Lambda]_n^{\wedge}}$ acts on strict units via raising to the *p*-th power.

It will follow that when R is perfect (i.e., φ_R is an isomorphism), the map $[p]_* : \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge}) \to \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge})$ differs from multiplication by p by an isomorphism. We therefore obtain a variant of Proposition 5.5 with $[p]_*$ replaced by $p : \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge}) \to \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge})$.

We begin with a lemma on the compatibility of Frobenius with tensor products. Let $\widehat{\otimes}$ denote the *p*-completed tensor product in $\operatorname{Sp}_p^{\wedge}$. If *R* and *S* have finite Tor-amplitude, then so does $R \widehat{\otimes} S$, so we have a Frobenius $\varphi_{R \widehat{\otimes} S}$.

Lemma 5.6. Let R and S be p-complete commutative ring spectra of finite Tor-amplitude. Then the two maps $R \widehat{\otimes} S \to R \widehat{\otimes} S$ given by $\varphi_R \widehat{\otimes} \varphi_S$ and $\varphi_{R \widehat{\otimes} S}$ are homotopic. *Proof.* Immediate from the diagram (with completions left implicit for clarity)



where the triangles arise from frob and triv being lax monoidal natural transformations [NS18, Section I.3]. $\hfill \Box$

As a result, we obtain a formula for the Frobenius of group algebras.

Proposition 5.7. Let R be a p-complete commutative ring spectrum of finite Tor-amplitude and let M be an abelian group. Let $[p]: R[M]_p^{\wedge} \to R[M]_p^{\wedge}$ be the morphism induced by the multiplication by p endomorphism of M. Then, the Frobenius morphism $\varphi_{R[M]_p^{\wedge}}$ is given by the composite

$$R[M]_p^{\wedge} \xrightarrow{\varphi_R[M]_p^{\wedge}} R[M]_p^{\wedge} \xrightarrow{[p]} R[M]_p^{\wedge}.$$

In particular, on $\pi_*R[M]_p^{\wedge}$ it is given by

$$\varphi_{R[M]_p^{\wedge}}\left(\sum_{m\in M} r_m[m]\right) = \sum_{m\in M} \varphi_R(r_m)[pm].$$

Proof. Since $R[M]_p^{\wedge} \simeq R \widehat{\otimes}(\mathbb{S}_p[M]_p^{\wedge})$, the compatibility of Frobenius with tensor products (Lemma 5.6) reduces the general case to the special case $R = \mathbb{S}_p$. Since $\varphi_{\mathbb{S}_p}$ is the identity ([NS18, Example IV.1.2(ii)]), it suffices to see that $\varphi_{\mathbb{S}_p[M]_p^{\wedge}} \simeq [p]$. This follows, by passing to *p*-completions, from the identification of the Tate diagonal of $\mathbb{S}[M]$ as in [NS18, Lemma IV.1.3].

To identify the effect of these maps on strict units, we use the following result of [Car22].


Proposition 5.8 ([Car22, Proposition 4.5]). Let R be a p-complete commutative ring spectrum of finite Tor-amplitude and let $\varphi := \varphi_R$ be its Frobenius map. Then the map

$$\varphi_* \colon \mathbb{G}_m(R) \to \mathbb{G}_m(R)$$

is homotopic to the multiplication by p map on the \mathbb{Z} -module spectrum $\mathbb{G}_m(R)$.

Combining this with Proposition 5.7, we obtain the following statement for group algebras:

Corollary 5.9. Let R be a commutative ring spectrum of finite Tor-amplitude and let M be an abelian group. Denote by $[p] : R[M]_p^{\wedge} \to R[M]_p^{\wedge}$ the map induced by multiplication by p on M. Then there is a commutative triangle



We will be particularly interested in rings R satisfying the following additional hypothesis:

Definition 5.10. A *p*-complete commutative ring spectrum R is *perfect* if frob : $R \to R^{tC_p}$ is an isomorphism.

Example 5.11. If κ is a (discrete) perfect \mathbb{F}_p -algebra, then the ring of spherical Witt vectors $\mathbb{S}_{W(\kappa)}$ is perfect. More generally, any algebra of the form $(\mathbb{S}_{W(\kappa)})^A$, for a finite complex A, is perfect.

The notion of a perfect commutative ring spectrum was previously defined in [Car22] and [Yua22] in slightly different ways, but the two definitions agree (and agree with this one) for rings of finite Tor-amplitude.

Remark 5.12. If R is perfect and p-completely flat, then $\pi_0 R/p$ is a perfect ring and $R \simeq \mathbb{S}_{W(\pi_0 R/p)}$.

For a perfect commutative ring spectrum R, Corollary 5.9 implies:

Proposition 5.13. Let R be a perfect p-complete commutative ring spectrum of finite Tor-amplitude and let Λ be a free abelian group of finite rank. Then there is a cofiber sequence in $Mod_{\mathbb{Z}}^{cn}$

$$\mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge}) \xrightarrow{p} \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge}) \longrightarrow \mathbb{G}_m^{\mathrm{red}}(R[\Lambda/p]).$$

Proof. Combine Proposition 5.5 with Corollary 5.9 and the fact that $(\varphi_R)_*$ is an isomorphism.

This gives us *p*-completed Λ -rigidity up to *p*-completion:

Corollary 5.14. Let R be a perfect p-complete commutative ring spectrum of finite Tor-amplitude and let Λ be a free abelian group of finite rank. Then the map

$$\widehat{\Theta}_{\Lambda} : \underline{\Lambda}(R) \to \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_n^{\wedge})$$

induces an isomorphism after p-completion.

Proof. By definition of *p*-completion, it suffices to show that the induced map

$$\underline{\Lambda}(R)/p \to \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^\wedge)/p$$

is an isomorphism. But $\underline{\Lambda}(R)/p \simeq \underline{\Lambda}/p(R)$ and by Proposition 5.13, we have $\mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge})/p \simeq \mathbb{G}_m^{\mathrm{red}}(R[\Lambda/p])$, so the statement follows from Corollary 4.13.

5.3. Staticity and rigidity for free abelian groups. We now turn to part (3) of our plan, completing the proof of rigidity for free abelian groups.

Remark 5.15. At this point, we restrict to $R = \mathbb{S}_{W(\kappa)}$ for clarity of proofs, despite the only critical additional hypothesis being that R is connective. While this restriction appears dramatic, we expect that any *p*-complete perfect connective ring R is of the form $\mathbb{S}_{W(\kappa)}$ (though we do not pursue this further here), so we do not really lose any generality.

The main technique is the notion of staticity introduced in §3.3. The following statement asserts that the space of reduced strict units in $\mathbb{S}_{W(\kappa)}[\Lambda]_p^{\wedge}$ is a *subset* of the reduced (strict) units of its π_0 . This will reduce the computation of strict units to the algebraic question of finding rank 1 units in the δ_p -ring $W(\kappa)[\Lambda]_p^{\wedge}$.

Proposition 5.16. Let κ be a perfect \mathbb{F}_p -algebra and Λ be a free abelian group of finite rank. Then $\mathbb{S}_{W(\kappa)}$ is $\mathbb{G}_m^{\mathrm{red}}(-[\Lambda]_p^{\wedge})$ -static.

Proof. For a connective spectrum X, let

$$V_p X \simeq \underline{\lim} (\dots \xrightarrow{p} X \xrightarrow{p} X \xrightarrow{p} X) \in \operatorname{Sp^{cn}},$$

and consider the fiber sequence of connective spectra

$$V_p \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge}) \to \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge}) \to \mathbb{G}_m^{\mathrm{red}}(R[\Lambda]_p^{\wedge})_p^{\wedge}.$$

By Lemma 3.18(2), the statement reduces to the following two facts:

- (1) $\mathbb{S}_{W(\kappa)}$ is $V_p \mathbb{G}_m^{\text{red}}(-[\Lambda]_p^{\wedge})$ -static. Since $\mathbb{G}_m^{\text{red}}(-)$ is a summand of $\mathbb{G}_m(-)$, it suffices to see that $\mathbb{S}_{W(\kappa)}[\Lambda]_p^{\wedge}$ is $V_p \mathbb{G}_m$ -static. But this follows from Proposition 3.20 and the fact that multiplication by p is invertible on $V_p \mathbb{G}_m \simeq \text{Spec}(\mathbb{S}[\mathbb{Z}^{[1/p]}])$.
- (2) $\mathbb{S}_{W(\kappa)}$ is $\mathbb{G}_m^{\mathrm{red}}(-[\Lambda]_p^{\wedge})_p^{\wedge}$ -static. To see this, we consider the commutative square

We wish to show that the right vertical map is an inclusion. The left vertical map is clearly an isomorphism, and by Corollary 5.14, the upper horizontal map is also an isomorphism. Thus, since the bottom horizontal map is an injection of discrete abelian groups (and in particular an inclusion), the right vertical map is an inclusion as well.

We now turn to the main result of this section, regarding rigidity for free abelian groups.

Proposition 5.17. The image of truncation $\mathbb{G}_m^{\mathrm{red}}(\mathbb{S}_{W(\kappa)}[\Lambda]_p^{\wedge}) \to \mathbb{G}_m^{\mathrm{red}}(W(\kappa)[\Lambda]_p^{\wedge})$ is contained in the image of

$$\widehat{\Theta}_{W(\kappa)} : \underline{\Lambda}(W(\kappa)) \to \mathbb{G}_m^{\mathrm{red}}(W(\kappa)[\Lambda]_p^{\wedge}).$$

Proof. Any (reduced) strict unit in $\mathbb{S}_{W(\kappa)}[\Lambda]_p^{\wedge}$ is classified by a map from $\mathbb{S}[t^{\pm 1}]$. By Proposition 2.41, the induced map on π_0 is naturally a map of δ_p -rings and in particular, the image of $t \in \mathbb{G}_m^{\delta_p, \operatorname{red}}(\mathbb{Z}_p[t^{\pm 1}])$ is a reduced rank 1 unit, i.e., an element of $\mathbb{G}_m^{\delta_p, \operatorname{red}}(W(\kappa)[\Lambda]_p^{\wedge})$. The statement then follows from Theorem 2.30.

Theorem 5.18. Let κ be a perfect \mathbb{F}_p -algebra and let Λ be a free abelian group of finite rank. Then $\mathbb{S}_{W(\kappa)}$ is p-completely Λ -rigid.

Proof. Applying Proposition 3.22, the statement follows from Proposition 5.16 and Proposition 5.17. \Box

Combining this with Corollary 4.13 and Proposition 5.2, we obtain:

Corollary 5.19. Let κ be a perfect \mathbb{F}_p -algebra and M be a finitely generated abelian group with no prime to p torsion. Then $\mathbb{S}_{W(\kappa)}$ is M-rigid.

6. STRICT UNITS AND MAPS OF SPHERICAL GROUP ALGEBRAS

In this final section, we finish proving the main results about strict units and maps between spherical group rings. We handle the *p*-complete case in $\S6.1$ and the integral case in $\S6.2$.

6.1. Group algebras over spherical Witt vectors. In the *p*-complete case, we have:

Theorem 6.1. Let M be a finitely generated abelian group and let κ be a perfect ring of characteristic p. Then applying π_0 induces an isomorphism

$$\mathbb{G}_m(\mathbb{S}_{W(\kappa)}[M]_p^{\wedge}) \xrightarrow{\sim} \mathbb{G}_m^{\delta_p}(W(\kappa)[M]_p^{\wedge}).$$

Proof. When M is a finite group of order prime to p, we have $\mathbb{S}_{W(\kappa)}[M]_p^{\wedge} \simeq \mathbb{S}_{W(\kappa[M])}$ and so the result follows from [Car22, Corollary 4.15] and Example 2.6.

In general, we may split M as $M = M_1 \oplus M_2$ where M_1 is finite of order prime to p and M_2 has only p-primary torsion part. We have compatible splittings

$$\mathbb{G}_m(\mathbb{S}_{W(\kappa)}[M]_p^{\wedge}) \simeq \mathbb{G}_m(\mathbb{S}_{W(\kappa)}[M_1]_p^{\wedge}) \oplus \mathbb{G}_m^{\mathrm{red}}(\mathbb{S}_{W(\kappa)}[M_1][M_2]_p^{\wedge})$$
$$\mathbb{G}_m^{\delta_p}(W(\kappa)[M]_p^{\wedge}) \simeq \mathbb{G}_m^{\delta_p}(W(\kappa)[M_1]_p^{\wedge}) \oplus \mathbb{G}_m^{\delta_p,\mathrm{red}}(W(\kappa)[M_1][M_2]_p^{\wedge}),$$

where the second summand denotes the rank 1 units which map to 1 in $W(\kappa)[M_1]^{\times}$. By the previous special case, the π_0 map identifies the first summands, and so it suffices to show that

$$\mathbb{G}_m^{\mathrm{red}}(\mathbb{S}_{W(\kappa)}[M_1][M_2]_p^{\wedge}) \to \mathbb{G}_m^{\delta_p,\mathrm{red}}(W(\kappa)[M_1][M_2]_p^{\wedge})$$

is an isomorphism. Replacing κ by $\kappa[M_1]$, we may further assume without loss of generality that $M_1 = 0$. But here, the result follows by Proposition 2.29, Corollary 5.19, and the evident commutativity of the diagram

Corollary 6.2. Let p be a prime. Then the functor $\pi_0 : \operatorname{CAlg}^M \to \widehat{\delta}$ -Ring of Proposition 2.41 is fully faithful on the subcategory of rings of the form $\mathbb{S}_{W(\kappa)}[M]_p^{\wedge}$, where κ is a perfect ring of characteristic p and M is a finitely generated abelian group.

Proof. Consider two such objects corresponding to (κ, M) and (κ', M') . We have a map of fiber sequences of spaces

$$\begin{split} \operatorname{Map}_{\mathbb{S}_{W(\kappa)}}(\mathbb{S}_{W(\kappa)}[M]_{p}^{\wedge}, \mathbb{S}_{W(\kappa')}[M']_{p}^{\wedge}) & \longrightarrow \operatorname{Map}_{W(\kappa)/}^{\widehat{\delta}}(W(\kappa)[M]_{p}^{\wedge}, W(\kappa')[M']_{p}^{\wedge}) \\ & \downarrow & \downarrow \\ \operatorname{Map}_{\operatorname{CAlg}}(\mathbb{S}_{W(\kappa)}[M]_{p}^{\wedge}, \mathbb{S}_{W(\kappa')}[M']_{p}^{\wedge}) & \longrightarrow \operatorname{Map}^{\widehat{\delta}}(W(\kappa)[M]_{p}^{\wedge}, W(\kappa')[M']_{p}^{\wedge}) \\ & \downarrow & \downarrow \\ \operatorname{Map}_{\operatorname{CAlg}}(\mathbb{S}_{W(\kappa)}, \mathbb{S}_{W(\kappa')}[M']_{p}^{\wedge}) & \longrightarrow \operatorname{Map}^{\widehat{\delta}}(W(\kappa), W(\kappa')[M']_{p}^{\wedge}). \end{split}$$

The bottom horizontal map is an isomorphism because even the composite with the inclusion into $\operatorname{Map}(W(\kappa), W(\kappa')[M']_p^{\wedge})$ is an isomorphism, by the universal property of spherical Witt vectors [Lur18, Example 5.2.7]. But the top horizontal map is also an isomorphism by considering \mathbb{Z} -module maps from M into the equivalence of Theorem 6.1. Therefore, the middle map is an equivalence, and the desired fully faithfulness follows.


6.2. Group algebras over the sphere. Over the sphere spectrum, we can describe the mapping spaces between group algebras in a particularly simple way.

Proposition 6.3. Let M be a finitely generated abelian group. Then $\mathbb{S}[M]$ is \mathbb{G}_m -static, i.e.

$$\mathbb{G}_m(\mathbb{S}[M]) \to \mathbb{G}_m(\mathbb{Z}[M])$$

is an inclusion of connective \mathbb{Z} -module spectra. Consequently, the same holds for $\mathbb{G}_m^{\mathrm{red}}(\mathbb{S}[M])$.

Proof. Since \mathbb{G}_m is representable, the collection of \mathbb{G}_m -static connective commutative ring spectra are closed under limits which are preserved by π_0 by Lemma 3.19. By the arithmetic fracture square



it suffices to show that $\mathbb{S}[M]_p^{\wedge}$ is \mathbb{G}_m -static for all p and that $\mathbb{S}[M] \otimes \mathbb{Q}$ and $(\prod_p \mathbb{S}[M]_p^{\wedge}) \otimes \mathbb{Q}$ are \mathbb{G}_m static. The latter two rings are discrete and hence static, and the staticity of $\mathbb{S}[M]_p^{\wedge}$ is Theorem 6.1. The statement about $\mathbb{G}_m^{\mathrm{red}}(\mathbb{S}[M])$ follows immediately, as $\mathbb{G}_m^{\mathrm{red}}$ is a retract of \mathbb{G}_m .

To prove the rigidity of the sphere in general, it remains to establish the following:

Proposition 6.4. Let M be a finitely generated abelian group. The image of the inclusion $\mathbb{G}_m^{\mathrm{red}}(\mathbb{S}[M]) \to \mathbb{G}_m^{\mathrm{red}}(\mathbb{Z}[M])$ is contained in the image of the map $\Theta_M \colon M \to \mathbb{G}_m^{\mathrm{red}}(\mathbb{Z}[M])$.

Proof. Since $\pi_0 \mathbb{S}[t^{\pm 1}] \simeq \mathbb{Z}[t^{\pm 1}]$ as $\hat{\delta}$ -rings, the image of $\mathbb{G}_m^{\text{red}}(\mathbb{S}[M])$ is contained in the reduced rank 1 units, i.e. in $\mathbb{G}_m^{\hat{\delta}}(\mathbb{Z}[M]) \subseteq \mathbb{G}_m(\mathbb{Z}[M])$. Hence, the result follows from Theorem 2.34.

We are finally ready to prove our main theorem.

Theorem 6.5. Let M be a finitely generated abelian group. Then the natural map

$$\Theta: M \to \mathbb{G}_m(\mathbb{S}[M])$$

is an isomorphism.

Proof. Since by [Car22, Theorem B, Remark 1.3] we have $\mathbb{G}_m(\mathbb{S}) \simeq 0$, it suffices to show that \mathbb{S} is *M*-rigid. By Proposition 6.3, \mathbb{S} is *M*-static, and by Proposition 6.4, the map $\mathbb{G}_m^{\text{red}}(\mathbb{S}[M]) \to \mathbb{G}_m^{\text{red}}(\pi_0 \mathbb{S}[M])$ lands in the image of Θ . Hence, the result follows from Proposition 3.22(2). \Box

Corollary 6.6. The spherical group ring functor S[-]: $Ab^{fg} \to CAlg$ is fully faithful.

Proof. We want to see that for finitely generated abelian groups M and N, the natural map

$$\operatorname{Hom}_{\operatorname{Ab}^{\operatorname{fg}}}(N, M) \to \operatorname{Hom}_{\operatorname{CAlg}}(\mathbb{S}[N], \mathbb{S}[M])$$

is an isomorphism. This follows by taking \mathbb{Z} -module maps from N to the equivalence of Theorem 6.5.

References

- [ABG18] Matthew Ando, Andrew J. Blumberg, and David Gepner. Parametrized spectra, multiplicative Thom spectra and the twisted Umkehr map. *Geom. Topol.*, 22(7):3761–3825, 2018.
- [AKN23] Benjamin Antieau, Achim Krause, and Thomas Nikolaus. Prismatic cohomology relative to δ -rings. arXivpreprint arXiv:2310.12770, 2023.
- [Bar16] Tobias Barthel. Chromatic completion. Proceedings of the American Mathematical Society, 144(5):2263– 2274, 2016.
- [BB19] Tobias Barthel and A Bousfield. On the comparison of stable and unstable *p*-completion. Proceedings of the American Mathematical Society, 147(2):897–908, 2019.
- [BCSY22] Tobias Barthel, Shachar Carmeli, Tomer M Schlank, and Lior Yanovski. The Chromatic Fourier Transform. arXiv preprint arXiv:2210.12822, 2022.
- $[Bha] Bhargav Bhatt. Lecture II: \delta-rings. https://public.websites.umich.edu/ bhattb/teaching/prismatic-columbia/lecture2-delta-rings.pdf.$
- [Bha22] Prasit Bhattacharya. Higher associativity of Moore spectra. Advances in Mathematics, 402:108319, 2022.
- [BHLS23] Robert Burklund, Jeremy Hahn, Ishan Levy, and Tomer M Schlank. *K*-theoretic counterexamples to Ravenel's telescope conjecture. *arXiv preprint arXiv:2310.17459*, 2023.
- [BHM93] M. Bökstedt, W. C. Hsiang, and I. Madsen. The cyclotomic trace and algebraic K-theory of spaces. Invent. Math., 111(3):465–539, 1993.
- [BMS19] Bhargav Bhatt, Matthew Morrow, and Peter Scholze. Topological Hochschild homology and integral p-adic Hodge theory. Publ. Math. Inst. Hautes Études Sci., 129:199–310, 2019.
- [BS17] Bhargav Bhatt and Peter Scholze. Projectivity of the Witt vector affine Grassmannian. Invent. Math., 209(2):329–423, 2017.
- [BS22] Bhargav Bhatt and Peter Scholze. Prisms and prismatic cohomology. Annals of Mathematics, 196(3):1135– 1275, 2022.
- [BS24] Robert Burklund and Vignesh Subramanian. A note on the Segal conjecture for large objects. arXiv preprint arXiv:2403.06724, 2024.
- [BSY22] Robert Burklund, Tomer M Schlank, and Allen Yuan. The Chromatic Nullstellensatz. arXiv preprint arXiv:2207.09929, 2022.
- [Bur22] Robert Burklund. Multiplicative structures on Moore spectra. arXiv preprint arXiv:2203.14787, 2022.
- [Car22] Shachar Carmeli. On the Strict Picard Spectrum of Commutative Ring Spectra. arXiv preprint arXiv:2208.03073, 2022.
- [CS19] Kestutis Cesnavicius and Peter Scholze. Purity for flat cohomology. arXiv preprint arXiv:1912.10932, 2019.
- [CSY18] Shachar Carmeli, Tomer M Schlank, and Lior Yanovski. Ambidexterity in chromatic homotopy theory. arXiv preprint arXiv:1811.02057, 2018.
- [CSY20] Shachar Carmeli, Tomer M Schlank, and Lior Yanovski. Ambidexterity and Height. arXiv preprint arXiv:2007.13089, 2020.
- [CSY21] Shachar Carmeli, Tomer M Schlank, and Lior Yanovski. Chromatic cyclotomic extensions. arXiv preprint arXiv:2103.02471, 2021.
- [Dev20] Sanath Devalapurkar. Roots of unity in K(n)-local rings. Proc. Amer. Math. Soc., 148(7):3187–3194, 2020.
- [Fun20] Jun Hou Fung. Strict units of commutative ring spectra. PhD thesis, Harvard University, 2020.
- [Gar21] Giles Gardam. A counterexample to the unit conjecture for group rings. Ann. of Math. (2), 194(3):967– 979, 2021.
- [Hig40] Graham Higman. The units of group-rings. Proceedings of the London Mathematical Society, 2(1):231–248, 1940.
- [HL13a] Michael Hopkins and Jacob Lurie. Ambidexterity in K(n)-local stable homotopy theory. preprint, available at https://www.math.ias.edu/ lurie/, 2013.

- [HL13b] Michael Hopkins and Jacob Lurie. Ambidexterity in K(n)-local stable homotopy theory. preprint, 2013.
- [HM03] Lars Hesselholt and Ib Madsen. On the K-theory of local fields. Ann. of Math. (2), 158(1):1–113, 2003.
- [Hol23] Adam Holeman. Derived δ -rings and relative prismatic cohomology. arXiv preprint arXiv:2303.17447, 2023.
- [JM96] Eric Jespers and C Polcino Milies. Units of group rings. Journal of Pure and Applied Algebra, 107(2-3):233-251, 1996.
- [Kar83] G Karpilovsky. On finite generation of unit groups of commutative group rings. Archiv der Mathematik, 40:503–508, 1983.
- [Kra] Achim Krause. Flat H_{∞} Moore spectra are δ -rings. In preparation.
- [Law20] Tyler Lawson. Adjoining roots in homotopy theory. arXiv preprint arXiv:2002.01997, 2020.
- [Lur18] Jacob Lurie. Elliptic cohomology II: Orientations. https://www.math.ias.edu/ lurie/papers/Elliptic-II.pdf, 2018.
- [Mat16] Akhil Mathew. The galois group of a stable homotopy theory. Advances in Mathematics, 291:403–541, 2016.
- [MS06] J. P. May and J. Sigurdsson. Parametrized homotopy theory, volume 132 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2006.
- [NS18] Thomas Nikolaus and Peter Scholze. On topological cyclic homology. Acta Mathematica, 221(2):203–409, 2018.
- [Oka84] Shichirô Oka. Multiplications on the Moore spectrum. Mem. Fac. Sci. Kyushu Univ. Ser. A, 38(2):257– 276, 1984.
- [Sch] Peter Scholze. What are the potential applications of perfectoid spaces to homotopy theory? MathOverflow. URL:https://mathoverflow.net/q/386521 (version: 2021-03-15).
- [Sta18] The Stacks Project Authors. Stacks Project. https://stacks.math.columbia.edu, 2018.
- [SW15] Hisham Sati and Craig Westerland. Twisted Morava K-theory and E-theory. J. Topol., 8(4):887–916, 2015.
- [Yua22] Allen Yuan. Integral models for spaces via the higher frobenius. Journal of the American Mathematical Society, 2022.