POINTWISE REGULARITY FOR LOCALLY UNIFORMLY ELLIPTIC EQUATIONS AND APPLICATIONS

YUANYUAN LIAN AND KAI ZHANG

Dedicated to the book Fully Nonlinear Elliptic Equations (by Luis A. Caffarelli and Xavier Cabré).

ABSTRACT. In this paper, we study the regularity for viscosity solutions of locally uniformly elliptic equations and obtain a series of interior pointwise $C^{k,\alpha}$ ($k \ge 1, 0 < \alpha < 1$) regularity with smallness assumptions on the solution and the right-hand term. As applications, we obtain various interior pointwise regularity for several classical elliptic equations, i.e., the prescribed mean curvature equation, the Monge-Ampère equation, the k-Hessian equations, the k-Hessian quotient equations and the Lagrangian mean curvature equation. Moreover, the smallness assumptions are necessary in most cases (see Remark 2.6, Remark 3.5, Remark 4.7, Remark 5.4 and Remark 6.5).

1. INTRODUCTION

In this paper, we study the interior Schauder regularity for viscosity solutions of

(1.1)
$$F(D^2u, Du, u, x) = f \text{ in } B_1,$$

where $B_1 \subset \mathbb{R}^n$ is the unit open ball and F is a *locally uniformly* elliptic operator. Precisely, we use the following notion.

Definition 1.1. Let $\rho > 0$. The $F : S^n \times \mathbb{R}^n \times \mathbb{R} \times B_1 \to \mathbb{R}$ is called locally uniformly elliptic with ρ (or ρ -uniformly elliptic) if there exist constants $0 < \lambda \leq \Lambda$ such that for any $|M|, |N|, |p|, |s| \leq \rho$ and $x \in B_1$,

(1.2)
$$\mathcal{M}^{-}(N,\lambda,\Lambda) \leq F(M+N,p,s,x) - F(M,p,s,x) \leq \mathcal{M}^{+}(N,\lambda,\Lambda),$$

where S^n denotes the set of $n \times n$ symmetric matrices (see Notation 1.14). The $\mathcal{M}^-, \mathcal{M}^+$ are the usual Pucci's extremal operators:

$$\mathcal{M}^+(M,\lambda,\Lambda) = \Lambda \sum_{\lambda_i > 0} \lambda_i + \lambda \sum_{\lambda_i < 0} \lambda_i, \quad \mathcal{M}^-(M,\lambda,\Lambda) = \lambda \sum_{\lambda_i > 0} \lambda_i + \Lambda \sum_{\lambda_i < 0} \lambda_i,$$

where λ_i are the eigenvalues of M.

²⁰²⁰ Mathematics Subject Classification. Primary 35B65, 35D40, 35J60, 35J96, 35J93. Key words and phrases. Fully nonlinear equation, regularity theory, viscosity solution, Schauder estimate, Monge-Ampère equation.

This research has been financially supported by the Project PID2020-118137GB-I00 and PID2020-117868GB-I00 funded by MCIN/AEI /10.13039/501100011033.

Remark 1.2. If $\rho = \infty$, we arrive at the definition of uniformly elliptic operators (see [27, Definition 2.1]). If F is a smooth operator and

$$\lambda I \leq D_M F(M,p,s,x) \leq \Lambda I, \; \forall \; |M| \leq 2\rho, |p|, |s| \leq \rho, x \in B_1,$$

then (1.2) holds by the Lagrange mean value theorem.

If F is ρ -uniformly elliptic and u is a solution of (1.1), then $\tilde{u} = Ku$ is a solution of

$$\tilde{F}(D^2\tilde{u}, D\tilde{u}, \tilde{u}, x) := KF(\frac{1}{K}D^2u, \frac{1}{K}Du, \frac{1}{K}u, x) = f$$

and \tilde{F} is $K\rho$ -uniformly elliptic. That is, the scaling operation changes the uniform ellipticity. Hence, the scaling operation is restricted in deducing the regularity. This is the main obstacle for developing the regularity theory.

The notion of viscosity solution is defined as follows. The main difference from the classical definition is that the solution must be small and the selection of a test function φ is more restricted.

Definition 1.3. Let F be ρ -uniformly elliptic and $f \in C(B_1)$. We say that $u \in C(B_1)$ is a viscosity supersolution of (1.1) if $||u||_{L^{\infty}(B_1)} \leq \rho/2$ and for any $x_0 \in B_1$ and $\varphi \in C^2(B_1)$ with

(1.3)
$$\|\varphi\|_{C^{1,1}(\bar{B}_1)} \le \rho, \quad \varphi(x_0) = u(x_0), \quad \varphi \le u \text{ in } B_1,$$

we have

 $F(D^2\varphi(x_0), D\varphi(x_0), \varphi(x_0), x_0) \le f(x_0).$

Similarly, we can define viscosity subsolution and viscosity solution as usual.

Since we consider the pointwise regularity in this paper, let us recall the definition of pointwise $C^{k,\alpha}$ (see [86, Definition 2.2] for the definition of $C^{k,\omega}$ with a general modulus of continuity).

Definition 1.4. Let $k \ge 0, 0 < \alpha \le 1$. We say that $f : B_1 \to \mathbb{R}$ is $C^{k,\alpha}$ at 0 or $f \in C^{k,\alpha}(0)$ if there exist a polynomial $P \in \mathcal{P}_k$ (see Notation 1.14) and a constant K such that

(1.4)
$$|f(x) - P(x)| \le K|x|^{k+\alpha}, \ \forall \ x \in B_1.$$

We call P the Taylor polynomial of f at 0 and define

$$D^{m}f(0) = D^{m}P(0), \quad ||f||_{C^{k}(0)} = ||P|| = \sum_{m=0}^{k} |D^{m}P(0)|,$$

 $[f]_{C^{k,\alpha}(0)} = \min\left\{K \middle| (1.5) \text{ holds with } K\right\}, \ \|f\|_{C^{k,\alpha}(0)} = \|f\|_{C^{k}(0)} + [f]_{C^{k,\alpha}(0)}.$

In addition, we say that $f \in C^k(0)$ if there exist a polynomial $P \in \mathcal{P}_k$ such that

(1.5)
$$|f(x) - P(x)| \le |x|^k \omega(|x|), \ \forall \ x \in B_1,$$

where ω is a modulus of continuity, i.e., $\omega : \mathbb{R}_+ \to \mathbb{R}$ and $\omega(r) \to 0$ as $r \to 0$.

The regularity theory for uniformly elliptic equations has been well developed. Taking the $C^{k,\alpha}$ regularity for example, we have interior C^{α} regularity ([27, Chapter 4], [78, 79]), interior $C^{1,\alpha}$ regularity ([27, Chapter 8]), interior $C^{2,\alpha}$ regularity ([16, 50, 74, 75], [27, Chapter 8]), interior $C^{k,\alpha}$ ($k \ge 3$) regularity ([86]), boundary C^{α} regularity ([88]), boundary $C^{1,\alpha}$ and $C^{2,\alpha}$ regularity ([75, 87, 107]), boundary $C^{k,\alpha}$ regularity ([86]).

For locally uniformly elliptic equations, the pointwise $C^{2,\alpha}$ regularity for viscosity of (1.1) was first obtained by Savin [100], which was found to have many applications to other problems, e.g.,

- Partial regularity for fully nonlinear equations [6].
- Unique continuation for fully nonlinear equations [5, 7].
- Regularity for singular nonlinear equations [146].
- Regularity for the σ_k -Loewner–Nirenberg problem [85].
- $W^{2,p}$ estimate for the complex Monge-Ampère equation [39].
- Problems from geometry [41, 83, 84, 108].

Later, it was extended to parabolic equations by Wang [135] and nonlocal elliptic equations by Yu [141]. They all considered homogenous equations (i.e., $f \equiv 0$). In this paper, we study the nonhomogeneous equations and derive a series of pointwise $C^{k,\alpha}$ ($k \geq 1$ and $0 < \alpha < 1$) regularity.

The motivation of studying locally uniformly elliptic equations is its wide applications. Besides the applications mentioned above, we can obtain various new pointwise regularity for non-uniformly elliptic equations (see Sections 2-6 for details). Moreover, it indicates that maybe regarding (transforming) non-uniformly elliptic equations as (to) locally uniformly elliptic equations to study the regularity is essential. In another word, based on the regularity theory for locally uniformly elliptic equations, we can obtain regularity for non-uniformly elliptic equations with *necessary* assumptions (see Remark 2.6, Remark 3.5, Remark 4.7, Remark 5.4 and Remark 6.5). Note that these applications are just a few of examples. This technique/viewpoint has potential applications to other equations, boundary value problems, parabolic equations, complex equations and equations on Riemannian manifolds etc.

Next, we state our main results on the locally uniformly elliptic equations. For their applications, we state the theorems and give the proofs in Sections 2-6. First, we consider the pointwise $C^{1,\alpha}$ regularity for the following equation in a special form:

(1.6)
$$F(D^2u, Du, u, x) = A^{ij}(Du, u, x)u_{ij} + B(Du, u, x) = 0$$
 in B_1 ,

where F is ρ -uniformly elliptic, i.e.,

 $\lambda I \le A(p, s, x) \le \Lambda I, \ \forall \ |p|, |s| < \rho, \ x \in B_1.$

If A is continuous, we define the modulus of continuity

$$\omega_A(r) = \sup\{|A(p,s,0) - A(q,t,0)| : |p|, |q|, |s|, |t| \le \rho, |p-q|, |s-t| \le r\}, \ \forall \ r > 0.$$

Theorem 1.5. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of (1.6) where F is ρ -uniformly elliptic and A is continuous with modulus ω_A . Suppose that for some $K_B, b_0 \ge 0$,

(1.7)
$$|B(p,s,x)| \le K_B, \ \forall \ |p|, |s| < \rho, \ x \in B_1,$$

$$(1.8) \quad -b_0|p-q| \le B(p,s,x) - B(q,s,x) \le b_0|p-q|, \ \forall \ |p|, |q|, |s| \le \rho, \ x \in B_1$$

and

$$\|u\|_{L^{\infty}(B_1)} \le \delta$$

where $\delta > 0$ depends only on $n, \lambda, \Lambda, \rho, b_0, \alpha, \omega_A$ and K_B . Then $u \in C^{\bar{1},\alpha}(0)$, i.e., there exists $P \in \mathcal{P}_1$ such that

$$|u(x) - P(x)| \le C|x|^{1+\alpha}, \ \forall \ x \in B_1$$

and

$$\|P\| \le \bar{C}\delta,$$

where \overline{C} depends only on $n, \lambda, \Lambda, \rho, b_0, \alpha$, and C depends also on ω_A .

Remark 1.6. We will apply Theorem 1.5 to the prescribed mean curvature equation in Section 2. In fact, we can prove interior $C^{1,\alpha}$ regularity under the following more general condition:

$$F(tM, p, s, x) = tF(M, p, s, x), \ \forall \ |M|, |p|, |s| \le \rho, \ x \in B_1, \ t > 0.$$

Since we do not expect any application, we only consider the simpler equation (1.6) in this paper.

Next, we consider the $C^{2,\alpha}$ regularity. If F is ρ -uniformly elliptic and $D_M F$ is continuous, we define the modulus of continuity

$$\omega_F(r) = \sup\{|D_M F(M, p, s, 0) - D_M F(N, q, t, 0)| : \\ |M|, |N|, |p|, |q|, |s|, |t| \le \rho, \ |M - N|, |p - q|, |s - t| \le r\}, \ \forall \ r > 0.$$

We also need the following structure condition. For any $|M|, |p|, |q|, |s|, |t| \leq$ ρ and $x \in B_1$,

$$(1.9) \quad -b_0|p-q|-c_0|s-t| \le F(M,p,s,x) - F(M,q,t,x) \le b_0|p-q|+c_0|s-t|,$$

where b_0, c_0 are two nonnegative constants.

The following is the interior pointwise $C^{2,\alpha}$ regularity.

Theorem 1.7. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of (1.1) where F is ρ -uniformly elliptic, $F(0,0,0,x) \equiv 0$ and $D_M F$ is continuous with modulus ω_F . Suppose that (1.9) holds,

$$(1.10) \quad |F(M, p, s, x) - F(M, p, s, 0)| \le \delta |x|^{\alpha}, \ \forall \ |M|, |p|, |s| \le \rho, \ x \in B_1$$

and

(1.11)
$$||u||_{L^{\infty}(B_1)} \leq \delta, \quad ||f||_{C^{\alpha}(0)} \leq \delta,$$

where $\delta > 0$ depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, \alpha$ and ω_F .

Then
$$u \in C^{2,\alpha}(0)$$
, i.e., there exists $P \in \mathcal{P}_2$ such that

$$|u(x) - P(x)| \le C|x|^{2+\alpha}, \ \forall \ x \in B_1$$

and

$$||P|| \le \overline{C}\delta, \quad F(D^2P, DP(0), P(0), 0) = f(0),$$

where \overline{C} depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, \alpha$, and C depends also on ω_F .

Remark 1.8. Theorem 1.7 was first proved by Savin [100] with $f \equiv 0$.

For higher $C^{k,\alpha}$ $(k \ge 3)$ regularity, we have

Theorem 1.9. Let $k \ge 3, 0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of (1.1) where F is ρ -uniformly elliptic. Let $F_0 \in C^{k-1}$ be ρ -uniformly elliptic, $F_0(0,0,0,x) \equiv 0$ and denote

$$K_F = \|F_0\|_{C^{k-1}(\bar{B}_{\rho} \times \bar{B}_1)}, \quad B_{\rho} = \{(M, p, s) : |M|, |p|, |s| < \rho\}.$$

Suppose that (1.9) holds,

(1.12)

$$|F(M, p, s, x) - F_0(M, p, s, x)| \le \delta |x|^{k-2+\alpha}, \ \forall \ |M|, |p|, |s| \le \rho, \ x \in B_1$$

and

$$||u||_{L^{\infty}(B_1)} \le \delta, \quad ||f||_{C^{k-2+\alpha}(0)} \le \delta,$$

where $\delta > 0$ depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, k, \alpha$ and K_F .

Then $u \in C^{k,\alpha}(0)$, i.e., there exists $P \in \mathcal{P}_k$ such that

$$|u(x) - P(x)| \le C|x|^{k+\alpha}, \ \forall \ x \in B_1$$

and

$$||P|| \leq C\delta, \quad |F_0(D^2P(x), DP(x), P(x), x) - P_f(x)| \leq C|x|^{k-1}, \ \forall \ x \in B_1,$$

where C depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, k, \alpha$ and K_F .

Remark 1.10. This higher pointwise regularity is new. Even for the uniformly elliptic equations, it was proved in [86] recently. In fact, the proof of Theorem 1.9 is inspired by [86]. In addition, by similar arguments in [86], we can obtain the pointwise $C^k(0)$ and $C^{k\ln L}(0)$ $(k \ge 2)$ regularity as well.

In fact, in above theorem, we only need that $||f||_{C^{\alpha}(0)}$ is small.

Corollary 1.11. Let $k \geq 3, 0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of (1.1) where F is ρ -uniformly elliptic. Let $F_0 \in C^{k-1}$ be ρ -uniformly elliptic and $F_0(0,0,0,x) \equiv 0$. Suppose that (1.9) holds,

$$|F(M, p, s, x) - F_0(M, p, s, x)| \le \delta |x|^{k-2+\alpha}, \ \forall \ |M|, |p|, |s| \le \rho, \ x \in B_1$$

and

$$\|u\|_{L^{\infty}(B_1)} \leq \delta, \quad f \in C^{k-2+\alpha}(0), \quad \|f\|_{C^{\alpha}(0)} \leq \delta,$$

where $\delta > 0$ depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, \alpha$ and K_F .

Then $u \in C^{k,\alpha}(0)$. That is, there exists $P \in \mathcal{P}_k$ such that

$$|u(x) - P(x)| \le C|x|^{k+\alpha}, \ \forall \ x \in B_1$$

and

 $||P|| \leq C\delta, \quad |F_0(D^2P(x), DP(x), P(x), x) - P_f(x)| \leq C|x|^{k-1}, \ \forall \ x \in B_1,$ where C depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, k, \alpha, K_F$ and $||f||_{C^{k-2+\alpha}(0)}$.

As a special case, we have

Corollary 1.12. Let $k \geq 3, 0 < \alpha < 1$ and $u \in C(\bar{B}_1)$ be a viscosity solution of (1.1) where $F \in C^{k-1}(\bar{B}_{\rho} \times \bar{B}_1)$ is ρ -uniformly elliptic and $F(0,0,0,x) \equiv 0$. Suppose that

$$||u||_{L^{\infty}(B_1)} \le \delta, \quad f \in C^{k-2+\alpha}(0), \quad ||f||_{C^{\alpha}(0)} \le \delta,$$

where $\delta > 0$ depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, \alpha$ and K_F . Then $u \in C^{k,\alpha}(0)$.

Remark 1.13. In this paper, a constant is called universal if it depends only on $n, \lambda, \Lambda, \rho, b_0$ and c_0 .

We use the perturbation technique as in [100] to prove above theorems. The idea is the following. Take $C^{2,\alpha}$ regularity for example. If u and f are small and F is smooth, the equation is close the Laplace equation (by the compactness method, see (9.4)). Hence, there exists $P_1 \in \mathcal{P}_2$ such that

$$||u - P_1||_{L^{\infty}(B_n)} \le \eta^{2+\alpha}$$

where $0 < \eta < 1$. By scaling, we have

$$||u - P_m||_{L^{\infty}(B_{\eta^m})} \le \eta^{m(2+\alpha)}, \ m \ge 1,$$

which means the $C^{2,\alpha}$ regularity. During the scaling, $||P_m||$ are kept small such that the scaled operators are always $\rho/2$ -uniformly elliptic.

The main obstacle is to show the compactness of solutions. Savin [100] proved a Harnack inequality by the technique of sliding paraboloids and then the Hölder regularity follows. In this paper, we follows the idea in [27] to prove a weak Harnack inequality, which leads to the Hölder regularity as well. In fact, we just repeat the argument in [27].

The idea "smallness implies regularity" has been found and used many years ago (e.g. [17, Proposition 2], [46], [55, (2.20)], [128, Theorem 1.3]). Since the equation is regarded as a perturbation of the Laplace equation, we can obtain $C^{k,\alpha}$ ($k \ge 1$) regularity for any $0 < \alpha < 1$. On the contrast, we can only obtain $C^{k,\alpha}$ regularity for some $0 < \alpha < \bar{\alpha}$ for a general fully nonlinear elliptic equation, where $0 < \bar{\alpha} < 1$ is a universal constant (see [27, Chapter 8]). In addition, the proofs of above theorems are relatively simpler than that for uniformly elliptic equations without the smallness assumptions (compare with [86]).

Note that even for uniformly elliptic equations, one usually start the proof by assuming that f is small (see [27, P. 75, Proof of Theorem 8.1]). Then a scaling argument can transform a general f to a small f. However, for locally uniformly elliptic equations, as pointed out in Remark 1.2, the scaling argument is restricted. Hence, we have to make the assumption in the theorem that f is small.

The paper is organized as follows. We first give the applications of above theorems in Sections 2-6. Precisely, we shall prove a series of interior pointwise $C^{k,\alpha}$ regularity for the prescribed mean curvature equation in Section 2, the Monge-Ampère equation in Section 3, the k-Hessian equation in Section 4, the k-Hessian quotient equation in Section 5 and the Lagrangian mean curvature equation in Section 6 respectively.

The proofs of above theorems are postponed to Sections 7-10. In Section 7, we prepare some preliminaries, such as the Alexandrov-Bakel'man-Pucci maximum principle etc. We prove $C^{1,\alpha}$ regularity in Section 8 and Section 9 is devoted the $C^{2,\alpha}$ regularity. Finally, we give the proof of $C^{k,\alpha}$ $(k \geq 3)$ regularity in Section 10.

Notation 1.14.

- (1) \mathbb{R}^n : the *n*-dimensional Euclidean space; \mathcal{S}^n : the set of $n \times n$ symmetric matrices with the standard order.
- (2) $\{e_i\}_{i=1}^n$: the standard basis of \mathbb{R}^n , i.e., $e_i = (0, ..., 0, 1, 0, ..., 0)$.
- (3) $x = (x_1, ..., x_n) = (x', x_n) \in \mathbb{R}^n$.
- (4) $|x| = \left(\sum_{i=1}^{n} x_i^2\right)^{1/2}$ for $x \in \mathbb{R}^n$; |M| = the spectrum radius of $M \in \mathcal{S}^n$.
- (5) I: the unit matrix in \mathcal{S}^n .
- (6) $a^+ = \max(a, 0), a^- = \max(-a, 0).$
- (7) $B_r(x_0) = B(x_0, r) = \{x \in \mathbb{R}^n : |x x_0| < r\}, B_r = B_r(0),$
- (8) Ω^c : the complement of Ω ; $\overline{\Omega}$: the closure of Ω , where $\Omega \subset \mathbb{R}^n$.
- (9) diam(Ω): the diameter of Ω .
- (10) Given $\varphi : \mathbb{R}^n \to \mathbb{R}$, define $\varphi_i = \partial \varphi / \partial x_i, \ \varphi_{ij} = \partial^2 \varphi / \partial x_i \partial x_j$ and we also use similar notations for higher order derivatives.
- (11) $D^0 \varphi = \varphi, \ D\varphi = (\varphi_1, ..., \varphi_n) \text{ and } D^2 \varphi = (\varphi_{ij})_{n \times n} \text{ etc.}$
- (12) We also use the standard multi-index notation. Let $\sigma = (\sigma_1, ..., \sigma_n) \in$ \mathbb{N}^n , i.e., each component σ_i is a nonnegative integer. Define

$$|\sigma| = \sum_{i=1}^{n} \sigma_i, \quad \sigma! = \prod_{i=1}^{n} (\sigma_i!), \quad x^{\sigma} = \prod_{i=1}^{n} x_i^{\sigma_i}, \quad D^{\sigma}\varphi = \frac{\partial^{|\sigma|}\varphi}{\partial x_1^{\sigma_1} \cdots \partial x_n^{\sigma_n}}.$$

(13)
$$|D^k \varphi| = \left(\sum_{|\sigma|=k} |D^\sigma \varphi|^2\right)^{1/2}$$
 for $k \ge 0$.
(14) Given $F: S^n \times \mathbb{R}^n \times \mathbb{R} \times \Omega \to \mathbb{R}$ defin

(14) Given $F: \mathcal{S}^n \times \mathbb{R}^n \times \mathbb{R} \times \Omega \to \mathbb{R}$, define

$$F_{M_{ij}} = \frac{\partial F}{\partial M_{ij}}, \quad F_{p_i} = \frac{\partial F}{\partial p_i}, \quad F_s = \frac{\partial F}{\partial s}, \quad F_{x_i} = \frac{\partial F}{\partial x_i},$$

where $1\leq i,j\leq n.$ Moreover, let $\xi\in\mathbb{N}^{n\times n}$ denote the matrix-valued multi-index. Then define

$$D_M^{\xi}F = \frac{\partial^{|\xi|}F}{\partial M_{ij}^{\xi_{ij}}}, \quad D_M^kF = \left\{\frac{\partial^kF}{\partial M^{\xi}} \colon |\xi| = k\right\}, \quad |D_M^kF| = \left(\sum_{|\xi|=k} \left|\frac{\partial^kF}{\partial M^{\xi}}\right|^2\right)^{1/2}$$

Similarly, we can define $D_p^k F$, $D_s^k F$, $D_x^k F$ and their norms etc.

(15) $\mathcal{P}_k(k \ge 0)$: the set of polynomials of degree less than or equal to k. That is, any $P \in \mathcal{P}_k$ can be written as

$$P(x) = \sum_{|\sigma| \le k} \frac{a_{\sigma}}{\sigma!} x^{\sigma}$$

where a_{σ} are constants. Define for r > 0

$$||P||_r = \sum_{|\sigma| \le k} r^{|\sigma|} |a_{\sigma}|, \quad ||P|| = ||P||_1 = \sum_{|\sigma| \le k} |a_{\sigma}|$$

(16) $\mathcal{HP}_k(k \ge 0)$: the set of homogeneous polynomials of degree k. That is, any $P \in \mathcal{HP}_k$ can be written as

$$P(x) = \sum_{|\sigma|=k} \frac{a_{\sigma}}{\sigma!} x^{\sigma}.$$

2. Prescribed mean curvature equation

In the following sections, we give the applications of the regularity of locally uniformly elliptic equations to several classical non-uniformly elliptic equations.

Since non-uniformly elliptic equations are much more difficult than uniformly elliptic ones, it is natural to study them by assuming the data are good enough and then obtain a priori estimates, existence of smooth solutions and Liouville type theorems etc. This classical strategy has been used widely. For example,

- For the prescribed mean curvature equation, see [9, 14, 52, 69, 73, 80, 103, 111, 133].
- For the Monge-Ampère equation, see [18, 30, 40, 43, 64, 65, 67, 72, 76, 119].
- For the k-Hessian equations, see [19, 59, 65, 66, 68, 77, 113, 131, 134].
- For the k-Hessian quotient equations, see [8, 32, 33, 34, 37, 44, 49, 89, 91, 113]
- For the Lagrangian mean curvature equation, see [10, 11, 19, 36, 90, 127, 137, 139, 143, 147]

In addition, there exist Pogorelov's type interior $C^{1,1}$ estimates for some equations. The smoothness requirements of the boundary can be relaxed. As a compensation, the boundary value must be an affine function or an admissible function in some sense. For example,

- For the Monge-Ampère equation, see [96], [97, P. 73, Chapter 5.3].
- For the k-Hessian equations, see [42].
- For the k-Hessian quotient equations, see [33, 37].

For the prescribed mean curvature equation and the Lagrangian mean curvature equation, we have stronger pure interior $C^{0,1}$ and $C^{1,1}$ estimates (i.e., estimates independent of the boundary information).

In another direction, it is also natural to consider the regularity of solutions. One may introduce weak solutions in some sense and then prove the existence and regularity. In this respect, there are following examples:

- For the prescribed mean curvature equation, see [45, 53, 54, 57, 81, 92, 149].
- For the Monge-Ampère equation, see [3, 21, 22, 23, 24, 25, 38, 47, 48, 62, 99].
- For the k-Hessian equations, see [31, 114, 116, 117, 118, 121, 122].
- For the Lagrangian mean curvature equation, see [12, 13, 35, 142].

As for the Schauder's type regularity, say $C^{2,\alpha}$ regularity under the assumption $f \in C^{\alpha}$, there are few results in this respect (see [140] for the 2-Hessian equation in dimension 3 and [12] for the Lagrangian mean curvature equation) besides the Monge-Ampère equation (see [22, 70, 98, 102, 123, 130] for the interior regularity; see [99] and [119] for the boundary regularity).

In the following sections, based on the regularity theory for locally uniformly elliptic equations, we will develop pointwise $C^{k,\alpha}$ ($k \ge 1$, $0 < \alpha < 1$) regularity for the prescribed mean curvature equation (Section 2), the Monge-Ampère equation (Section 3), the k-Hessian equation (Section 4), the k-Hessian quotient equation (Section 5) and the Lagrangian mean curvature equation (Section 6).

We use the notion of viscosity solution for these equations as well. For the prescribed mean curvature equation and the Lagrangian mean curvature equation, the definition of viscosity solution is exactly the same as the usual (see [27, Chapter 2]). For the Monge-Ampère equation, the test function φ should be a convex function (see [61, Definition 1.3.1]). The φ should be k-admissible if we define a viscosity solution for a k-Hessian equation or k-Hessian quotient equation (see [124, Section 2], [82, Definition 1.1]).

Note that above definitions of viscosity solution are different from Definition 1.3. In fact, in the following sections, we will regard (transform) an equation as (to) a locally uniformly elliptic equation. Then the (transformed) solution will be a viscosity solution in the sense of Definition 1.3.

We first give a general result:

Theorem 2.1. Let $k \ge 2$, $0 < \alpha < 1$ and u be a viscosity solution of

$$F(D^2u, Du, u, x) = f \quad in \ B_1,$$

where F is smooth. Suppose that $u \in C^2(0), f \in C^{k-2,\alpha}(0)$ and

(2.1) $A := D_M F(D^2 u(0), Du(0), u(0), 0) > 0.$

Then $u \in C^{k,\alpha}(0)$.

Proof. By $u \in C^2(0)$, there exist $P \in \mathcal{P}_2$ and a modulus of continuity ω such that

$$|u(x) - P(x)| \le \omega(|x|)|x|^2, \ \forall \ x \in B_1.$$

Since u is a viscosity solution,

$$F(D^{2}u(0), Du(0), u(0), 0) = F(D^{2}P(0), DP(0), P(0), 0) = f(0).$$

For r > 0, let

$$y = \frac{x}{r}, \quad \tilde{u}(y) = \frac{u(x) - P(x)}{r^2}, \quad \tilde{f}(y) = f(x) - F(D^2 P, DP(x), P(x), x).$$

Then \tilde{u} is a solution of

(2.2)
$$\tilde{F}(D^2\tilde{u}, D\tilde{u}, \tilde{u}, y) = \tilde{f} \text{ in } B_1$$

where

$$\tilde{F}(M, p, s, y) := F(M+D^2P, rp+DP(x), r^2s+P(x), x) - F(D^2P, DP(x), P(x), x)$$
.
Then $\tilde{F}(0, 0, 0, y) \equiv 0$ and by (2.1),

$$D_M \tilde{F}(0,0,0,0) = A > 0.$$

Since \tilde{F} is smooth, \tilde{F} is ρ -uniformly elliptic with some $0 < \lambda \leq \Lambda$ and these three constants depends only on n, A, ||P|| and $||F||_{C^{1,1}}$.

In addition, by the definition of \tilde{u} and \tilde{f} ,

$$\|\tilde{u}\|_{L^{\infty}(B_1)} \le \omega(r), \quad \tilde{f}(0) = 0, \quad [f]_{C^{\alpha}(0)} \le Cr^{\alpha}.$$

Thus, we can choose r small enough such that

$$\|\tilde{u}\|_{L^{\infty}(B_1)} \le \delta, \quad \|\tilde{f}\|_{C^{\alpha}(0)} \le \delta,$$

where δ is small enough such that \tilde{u} is a viscosity solution of (2.2) in the sense of Definition 1.3 and we can apply Corollary 1.12 to (2.2). Therefore, $\tilde{u} \in C^{k,\alpha}(0)$ and hence $u \in C^{k,\alpha}(0)$.

Remark 2.2. If u is an appropriate viscosity solution, (2.1) holds automatically. At least, it holds for all equations treated in the following sections. Hence, in general, if $u \in C^2(0)$ and $f \in C^{k-2,\alpha}(0)$, then $u \in C^{k,\alpha}(0)$. In conclusion, we have one rough but interesting assertion: For any elliptic equation with a smooth operator, the pointwise $C^{2,\alpha}$ regularity holds almost everywhere if u is a convex viscosity solution and $f \in C^{\alpha}$ ($0 < \alpha < 1$).

Now, we consider the prescribed mean curvature equation.

Theorem 2.3. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of (2.3)

$$\operatorname{div}\left(\frac{Du}{\sqrt{1+|Du|^2}}\right) = \frac{1}{\sqrt{1+|Du|^2}} \left(\delta^{ij} - \frac{u_i u_j}{1+|Du|^2}\right) u_{ij} = f \quad in \ B_1,$$

where $f \in L^{\infty}(B_1)$. Then $u \in C^{1,\alpha}(0)$ provided one of the following conditions holds:

(i) there exists $P \in \mathcal{P}_1$ such that

$$||u - P||_{L^{\infty}(B_1)} \le \delta,$$

where $0 < \delta < 1$ depends only on $n, \alpha, |DP|$ and $||f||_{L^{\infty}(B_1)}$. (ii) there exists a constant |K| < n-1 such that

$$\|f - K\|_{L^{\infty}(B_1)} \le \delta,$$

where $0 < \delta < 1$ depends only on n, α, K and $||u||_{L^{\infty}(B_1)}$. (*iii*) $u \in C^{0,1}(0)$. (*iv*) there exists $P \in \mathcal{P}_1$ such that

$$u = P \quad on \ \partial B_{\delta},$$

where $\delta > 0$ depends only on $n, \alpha, |DP|$ and $||f||_{L^{\infty}(B_1)}$.

For higher regularity, we have

Theorem 2.4. Let $k \ge 2, 0 < \alpha < 1$, $u \in C(\overline{B}_1)$ be a viscosity solution of (2.3) and $f \in C^{k-2,\alpha}(0)$. Then $u \in C^{k,\alpha}(0)$ provided one of the four conditions in Theorem 2.3 holds.

Remark 2.5. The mean curvature operator is given by

$$F(M,p) = \frac{1}{\sqrt{1+|p|^2}} \left(\delta^{ij} - \frac{p_i p_j}{1+|p|^2}\right) M_{ij}.$$

Hence, F is smooth and it is easy to verify (cf. Remark 1.2) that F is 1uniformly elliptic with $\lambda = \sqrt{2}/4$ and $\Lambda = \sqrt{2}/2$. Moreover, for any $P \in \mathcal{P}_1$, define

$$G(M, p) = F(M, p + DP).$$

Then G is smooth and 1-uniformly elliptic with λ and Λ , which depend only on |DP|. Hence, the regularity theory for locally uniformly elliptic equations are applicable to (2.3).

Remark 2.6. To investigate the regularity of solutions of (2.3), one usually assumes $f \in C^{0,1}$ at least (e.g. [15, Theorem 4.2],[53],[56, Chapter 16],[57]). In fact, if $f \in C^{0,1}$, one can prove the interior gradient bound, which was first obtained for the minimal surface equation by Bombieri, De Giorgi and Miranda [14] (see also [9, 52, 73, 80, 111, 133]). Then the equation becomes uniformly elliptic and the regularity follows.

On the other hand, if $f \notin C^{0,1}$, we cannot obtain the $C^{1,\alpha}$ regularity in general. Consider the following counterexample borrowed from [45, Section 8]:

$$u(x) = \begin{cases} -(1-|x|)^{\theta}, & \text{if } 0 \le r \le 1; \\ (|x|-1)^{\theta}, & \text{if } 1 < r < 2, \end{cases}$$

where $0 < \theta < 1/2$. It can be checked directly that u is a viscosity solution of (2.3) in B_2 with

$$f \in C^{1-2\theta}(\bar{B}_2), \quad [f]_{C^{1-2\theta}(x_0)} \le C\theta^{-2}, \ \forall \ x_0 \in \partial B_1.$$

However, we have only $u \in C^{\theta}$ at ∂B_1 . Hence, the "smallness" assumptions in Theorem 2.3 cannot be removed.

Moreover, if θ is smaller, f is smoother but $[f]_{C^{1-2\theta}}$ is bigger. Correspondingly, u has lower regularity. This phenomenon indicates that the smallness is more important than the smoothness for the regularity in this case (i.e., non-uniformly elliptic equations with lower regularity on f).

Above observation may imply that regarding the prescribed mean curvature equation as a locally uniformly elliptic equation to study the regularity is essential (see also Remark 3.5, Remark 5.4 and Remark 6.5).

Remark 2.7. Roughly speaking, Theorem 2.3 states that the interior $C^{1,\alpha}$ regularity holds for the prescribed mean curvature equation under some "smallness" assumption (except (iii)).

As pointed out in Remark 2.6, to obtain $C^{1,\alpha}$ regularity, one usually prove the interior gradient bound first and then the equation becomes uniformly elliptic. The assumption (iii) can be understood in some sense that the equation is *pointwise* uniformly elliptic at 0. Then we obtain the pointwise $C^{1,\alpha}(0)$ regularity.

Remark 2.8. The $C^{1,\alpha}$ regularity for the minimal surface equation under the assumption that u is small, i.e.,

$$||u||_{L^{\infty}(B_1)} \leq \delta, \ f \equiv 0 \Longrightarrow u \in C^{1,\alpha}(\bar{B}_{1/2})$$

has been proved by De Giorgi [46] as a special case (see also [58, Chapters 6-8], [100, P. 676] and [101, P. 42]). Theorem 2.3 extends this result to

$$||u||_{L^{\infty}(B_1)} \leq \delta, \ f \in L^{\infty}(B_1) \Longrightarrow u \in C^{1,\alpha}(B_{1/2}).$$

We first prove a lemma.

Lemma 2.9. Let u be a viscosity solution of (2.3). Then for any $\delta_1 > 0$, there exists $\delta > 0$ depending only on n, K and $||u||_{L^{\infty}(B_1)}$ such that if

$$||f - K||_{L^{\infty}(B_1)} \le \delta, \quad |K| < n - 1,$$

we have for some $P \in \mathcal{P}_1$

$$\|u - P\|_{L^{\infty}(B_r)} \le \delta_1 r$$

and

$$|DP| \le C$$

where 0 < r < 1/2 and C depend only on n, K and $||u||_{L^{\infty}(B_1)}$.

Proof. Since |K| < n - 1, there exist two solutions $v^{\pm} \in C^{\infty}(B_1) \cap C(\bar{B}_1) \cap W^{1,1}(B_1)$ of the following prescribed mean curvature equations (see [56, Theorem 16.11] and [53, Theorem 1])

$$\begin{cases} \operatorname{div} A(Dv^{\pm}) = K \pm 2\delta & \text{ in } B_1; \\ v^{\pm} = u & \text{ on } \partial B_1 \end{cases}$$

where δ is taken small such that $|K \pm 2\delta| \le n-1$ and

(2.4)
$$A(p) := \frac{p}{\sqrt{1+|p|^2}}, \ \forall \ p \in \mathbb{R}^n.$$

Since v^{\pm} are smooth, by the definition of viscosity solution,

$$(2.5) v^+ \le u \le v^- \quad \text{in } B_1.$$

We claim that for any $\delta_2 > 0$, if δ small enough (depending only on $\delta_2, n, K, \|u\|_{L^{\infty}(B_1)}$),

(2.6)
$$\|v^{\pm} - v\|_{L^{\infty}(B_{1/2})} \le \delta_2,$$

where v is the solution of

$$\begin{cases} \operatorname{div} A(Dv) = K & \text{in } B_1; \\ v = u & \text{on } \partial B_1. \end{cases}$$

We prove the claim by contradiction and the proof is inspired by [110]. Suppose not. Then there exist δ_2 and sequences of u_m, v_m^+, v_m satisfying $||u_m||_{L^{\infty}(B_1)} \leq K$,

(2.7)
$$\begin{cases} \operatorname{div} A(Dv_m^+) = K + \frac{1}{m} & \text{in } B_1; \\ v_m^+ = u_m & \text{on } \partial B_1, \end{cases} \begin{cases} \operatorname{div} A(Dv_m) = K & \text{in } B_1; \\ v_m = u_m & \text{on } \partial B_1 \end{cases}$$

and

(2.8)
$$\|v_m^+ - v_m\|_{L^{\infty}(B_{1/2})} > \delta_2$$

By the interior derivatives estimates (see [56, Corollary 16.7]),

$$\|v_m^+\|_{C^{1,1}(\bar{\Omega}')}, \quad \|v_m\|_{C^{1,1}(\bar{\Omega}')} \le C, \ \forall \ \Omega' \subset \subset B_1,$$

where C depends only on $n, \Omega', ||v_m^+||_{L^{\infty}(B_1)}$ and $||v_m||_{L^{\infty}(B_1)}$. From the Alexandrov-Bakel'man-Pucci maximum principle (see [4, Theorem 6]),

(2.9)
$$\|v_m^+\|_{L^{\infty}(B_1)}, \|v_m\|_{L^{\infty}(B_1)} \le C_{2}$$

where C depends only on n and K. Hence, there exist \bar{v}^+ and \bar{v} such that (up to a subsequence)

(2.10)
$$v_m^+ \to \bar{v}^+, v_m \to \bar{v} \text{ in } C^1(\bar{\Omega}'), \forall \Omega' \subset B_1.$$

In addition, since $||v_m^+||_{W^{1,1}(B_1)}, ||v_m||_{W^{1,1}(B_1)}$ are uniformly bounded (see [110, (16), P. 319]),

(2.11)
$$v_m^+ \to \bar{v}^+, v_m \to \bar{v} \text{ weakly in } W^{1,1}(B_1).$$

Note that $v_m^+ - v_m \in W_0^{1,1}(B_1)$. By using it as the test function in (2.7), we have

$$\int_{B_1} \left(A(Dv_m^+) - A(Dv_m) \right) \left(Dv_m^+ - Dv_m \right) = \frac{1}{m} \int_{B_1} \left(v_m^+ - v_m \right).$$

By (2.9) and

$$(A(p) - A(q)) \cdot (p - q) \ge \frac{|p - q|^2}{(1 + |p|^2 + |q|^2)^{3/2}}, \ \forall \ p, q \in \mathbb{R}^n,$$

we have

$$\int_{B_1} \frac{|Dv_m^+ - Dv_m|^2}{(1 + |Dv_m^+|^2 + |Dv_m|^2)^{3/2}} \le \frac{C}{m}.$$

Let $m \to \infty$ and by the Fatou's lemma,

$$\int_{B_1} \frac{|D\bar{v}^+ - D\bar{v}|^2}{(1+|D\bar{v}^+|^2 + |D\bar{v}|^2)^{3/2}} = 0.$$

By (2.11), $\bar{v}^+ - \bar{v} \in W_0^{1,1}(B_1)$. Hence, $\bar{v}^+ \equiv \bar{v}$ in B_1 , which contradicts with (2.8). Therefore, (2.6) holds.

By combining (2.5) with (2.6),

(2.12)
$$||u - v||_{L^{\infty}(B_1)} \le \delta_2.$$

Since v is smooth, there exists $P \in \mathcal{P}_1$ such that

$$|v(x) - P(x)| \le C|x|^2, \ \forall \ x \in B_{1/2}$$

and

$$|DP| \le C,$$

where C depends only on n and $||u||_{L^{\infty}(B_1)}$.

Take 0 < r < 1/2 small enough such that $Cr < \delta_1/2$. Then

(2.13)
$$||v - P||_{L^{\infty}(B_r)} \le \frac{\delta_1}{2}r$$

In addition, take δ_2 small enough such that $\delta_2 \leq \delta_1 r/2$. Therefore, by (2.12) and (2.13),

$$||u - P||_{L^{\infty}(B_r)} \le \delta_1 r.$$

Now, we give the

Proof of Theorem 2.3. For (i), let $\tilde{u} = u - P$ and \tilde{u} is a solution of

$$G(D^2v, Dv) = F(D^2v, Dv + DP) = f \quad \text{in } B_1,$$

where F, G are defined as in Remark 2.5. Hence, G is smooth and is 1uniformly elliptic with ellipticity constants depending only on |DP|. Then by Theorem 1.5, the conclusion follows.

Next, we prove (ii). Let $\delta_1 > 0$ to be specified later. By Lemma 2.9, there exist 0 < r < 1/2 and $P \in \mathcal{P}_1$ such that

$$||u - P||_{L^{\infty}(B_r)} \le \delta_1 r.$$

Let

$$y = \frac{x}{r}$$
, $\tilde{u}(y) = \frac{u(x) - P(x)}{r}$, $\tilde{f}(y) = rf(x)$.

Then \tilde{u} is a solution of

$$F(D^2\tilde{u}, D\tilde{u} + DP) = \tilde{f}$$
 in B_1

and

$$\|\tilde{u}\|_{L^{\infty}(B_1)} \le \delta_1, \quad \|\tilde{f}\|_{L^{\infty}(B_1)} \le \|f\|_{L^{\infty}(B_1)} \le n-1.$$

Then $\tilde{u} \in C^{1,\alpha}(0)$ by Theorem 1.5 provided δ_1 is small enough, which is guaranteed by taking δ small enough.

Next, we prove (iii). For r > 0, let

(2.14)
$$y = \frac{x}{r}, \quad \tilde{u}(y) = \frac{u(x) - u(0)}{r}, \quad \tilde{f}(y) = rf(x)$$

Then \tilde{u} is a solution of

$$F(D^2\tilde{u}, D\tilde{u}) = \tilde{f}$$
 in B_1 .

By the assumption, \tilde{u} is bounded. We choose r small enough (depending on $||f||_{L^{\infty}(B_1)}$) such that $||\tilde{f}||_{L^{\infty}}$ is small. Then the conclusion follows from (ii). Finally, we prove (iv). Let

$$y = \frac{x}{\delta}, \quad \tilde{u}(y) = \frac{u(x) - P(x)}{\delta}, \quad \tilde{f}(y) = \delta f(x).$$

Then \tilde{u} is a solution of

$$F(D^2\tilde{u}, D\tilde{u} + DP) = \tilde{f}$$
 in B_1 .

Take δ small enough such that $\|\tilde{f}\|_{L^{\infty}(B_1)}$ is small. In addition, since $\tilde{u} = 0$ on ∂B_1 , by the Alexandrov-Bakel'man-Pucci maximum principle (see [4, Theorem 6]), $\|\tilde{u}\|_{L^{\infty}(B_1)}$ is small. Then the conclusion follows from Theorem 1.5.

The next is the

Proof of Theorem 2.4. By Theorem 2.3, $u \in C^{1,\alpha}(0)$. For r > 0, let

$$y = \frac{x}{r}, \quad \tilde{u}(y) = \frac{u(x) - P_u(x)}{r}, \quad \tilde{f}(y) = rf(x).$$

Then \tilde{u} is a solution of

$$F(D^2\tilde{u}, D\tilde{u} + DP_u) = \tilde{f}$$
 in B_1

Since $u \in C^{1,\alpha}(0)$,

$$\|\tilde{u}\|_{L^{\infty}(B_{1})} = r^{-1} \|u - P_{u}\|_{L^{\infty}(B_{r})} \le Cr^{\alpha},$$

$$\|\tilde{f}\|_{C^{k-2,\alpha}(0)} \le \sum_{i=0}^{k-2} r^{i+1} |D^{i}f(0)| + r^{k-1+\alpha} [f]_{C^{k-2,\alpha}(0)}.$$

Hence, we can choose r small enough such that $\|\tilde{u}\|_{L^{\infty}}$ and $\|\tilde{f}\|_{C^{k-2,\alpha}(0)}$ are small. Then the conclusion follows from Theorem 1.7 and Theorem 1.9. \Box

At the end of this section, we give two remarks.

Remark 2.10. Since the prescribed mean curvature equation has the divergence structure, one may consider a weak solution $u \in W^{1,1}(B_1)$ of (2.3) rather than a viscosity solution. We have the conclusion for weak solutions as well. For example, let us assume

$$\|u\|_{L^{\infty}(B_1)} \le \delta, \quad \|f\|_{L^{\infty}(B_1)} \le \delta.$$

where $0 < \delta < 1$ depends only on *n* and α .

Then we can approach the regularity by an approximation argument similar to [110, Proof of Theorem 1]. Take sequences of smooth functions u_m, f_m such that

$$u_m \to u$$
 in $W^{1,1}(B_1)$, $f_m \to f$ in $L^2(B_1)$

and

$$||u_m||_{L^{\infty}(B_1)} \le \delta, \quad ||f_m||_{L^{\infty}(B_1)} \le \delta, \ \forall \ m \ge 1$$

Let $v_m \in C^{\infty}(\bar{B}_1)$ be solutions of (see [56, Theorem 16.10])

$$\begin{cases} \operatorname{div} A(Dv_m) = f_m & \text{in } B_1; \\ v_m = u_m & \text{on } \partial B_1. \end{cases}$$

By the Alexandrov-Bakel'man-Pucci maximum principle,

$$\|v_m\|_{L^{\infty}(B_1)} \le C\delta,$$

where C depends only on n. Then from (i) of Theorem 2.3,

$$\|v_m\|_{C^{1,\alpha}(\bar{B}_{1/2})} \le C,$$

where C depends only on n and α . Hence, there exists $\bar{v} \in C^{1,\alpha}(\bar{B}_{1/2})$ such that (up to a subsequence)

$$v_m \to \bar{v}$$
 in $C^1(\bar{B}_{1/2})$.

As in the proof of Lemma 2.9,

$$\int_{B_1} \left(A(Dv_m) - A(Du) \right) \left(Dv_m - Du_m \right) = \int_{B_1} (f_m - f) \left(v_m - u_m \right).$$

Hence,

$$\begin{split} &\int_{B_1} \frac{|Dv_m - Du|^2}{(1+|Dv_m|^2+|Du|^2)^{3/2}} \leq \int_{B_1} \left(A(Dv_m) - A(Du)\right) \left(Dv_m - Du\right) \\ &= \int_{B_1} \left(f_m - f\right) \left(v_m - u_m\right) + \int_{B_1} \left(A(Dv_m) - A(Du)\right) \left(Du_m - Du\right) \\ &\leq C \|f_m - f\|_{L^2(B_1)} + 2\|Du_m - Du\|_{L^1(B_1)}. \end{split}$$

Let $m \to \infty$ and we have

$$\int_{B_{1/2}} \frac{|D\bar{v} - Du|^2}{(1 + |D\bar{v}|^2 + |Du|^2)^{3/2}} = 0.$$

Thus, for some constant c_0 ,

$$u \equiv \bar{v} + c_0 \quad \text{in } B_{1/2}.$$

Therefore, $u \in C^{1,\alpha}(\bar{B}_{1/2})$.

Remark 2.11. The theory for locally uniformly elliptic equations is not applicable to the *p*-Laplace equations. Indeed, the *p*-Laplace equations are more like uniformly elliptic equations (in particular when $Du \neq 0$). Hence, one may use the uniformly elliptic equations techniques to study the *p*-Laplace equations (e.g. [27, 129]). In addition, the *p*-Laplace operator is not smooth.

3. Monge-Ampère equation

In this section, we consider the Monge-Ampère equation:

$$\det D^2 u = f \quad \text{in } B_1.$$

We have the following observation. For any convex polynomial $P \in \mathcal{P}_2$ with $\det(D^2 P) = 1$, define

$$F(M) = \det(M + D^2 P) - 1, \ \forall \ M \in \mathcal{S}^n.$$

Then $F \in C^{\infty}$, F(0) = 0. Moreover, F is ρ -uniformly elliptic with $\tilde{\lambda}$, $\tilde{\Lambda}$ and these three constants depend only on n and $|D^2P|$. Therefore, the regularity theory for locally uniformly elliptic equations is applicable.

We have the following interior pointwise $C^{k,\alpha}$ regularity.

Theorem 3.1. Let $k \ge 2, 0 < \alpha < 1$ and u be a strictly convex viscosity solution of

$$\det D^2 u = f \quad in \ B_1.$$

Suppose that $0 < \lambda \leq f \leq \Lambda$ and $f \in C^{k-2,\alpha}(0)$. Then $u \in C^{k,\alpha}(0)$.

If the dimension n = 2, the solution is always strictly convex (see [2], [63] [51, Theorem 2.19], [120, Remark 3.2]). Hence, we have

Corollary 3.2. Let $k \ge 2, 0 < \alpha < 1$ and u be a convex viscosity solution of

$$\det D^2 u = f \quad in \ B_1 \subset \mathbb{R}^2.$$

Suppose that $0 < \lambda \leq f \leq \Lambda$ and $f \in C^{k-2,\alpha}(0)$. Then $u \in C^{k,\alpha}(0)$.

By applying above result to the prescribed Gaussian curvature equation, we have

Corollary 3.3. Let $k \ge 2, 0 < \alpha < 1$ and u be a strictly convex viscosity solution of

det
$$D^2 u = K(x) \left(1 + |Du|^2\right)^{\frac{n+2}{2}}$$
 in B_1 .

Suppose that $0 < \lambda \leq K \leq \Lambda$ and $K \in C^{k-2,\alpha}(0)$. Then $u \in C^{k,\alpha}(0)$.

Proof. Since u is convex,

$$||Du||_{L^{\infty}(B_{1/2})} \le C ||u||_{L^{\infty}(B_{1})}.$$

Hence, the right-hand term is bounded between two positive constants. Then from the strict convexity of u, we have $u \in C^{1,\beta}(\bar{B}_{1/2})$ for some $0 < \beta < 1$ (see [23, Theorem 2], [51, Corollary 4.21] and [120, Lemma 3.5]). Therefore, the conclusion follows from Theorem 3.1.

Remark 3.4. The $C^{2,\alpha}$ regularity for the Monge-Ampère equation is wellknown. Sabitov [98] and Schulz [102] proved the $C^{2,\alpha}$ regularity for n = 2. If $f \in C^{0,1}$, the $C^{2,\alpha}$ regularity for general dimensions was derived by Urbas [123]. The $C^{2,\alpha}$ regularity for general dimensions was due to Caffarelli [22] (see also [70]). Trudinger and Wang obtained boundary $C^{2,\alpha}$ regularity [119] and the pointwise version was proved by Savin [99]. Of course, Savin's proof is also applicable to derive interior pointwise $C^{2,\alpha}$ regularity.

As pointed out in Remark 1.10, we can obtain pointwise C^k and $C^{k,\text{LnL}}$ $(k \ge 2)$ regularity as well. In this respect, the C^2 regularity was proved by Wang [130] and $C^{2,\text{LnL}}$ regularity was proved by Jian and Wang [70].

Remark 3.5. If the dimension $n \geq 3$, by the well-known Pogorelov's counterexample (see [97, P. 81-83] and a good explanation on this counterexample in [51, Chapter 3.2]), the condition "*u* is strictly convex" can not be dropped. In addition, " $f \geq \lambda > 0$ " cannot be replaced by " $f \geq 0$ ". Indeed, the best regularity for the latter is $C^{1,1}$ in general (see [115, Theorem 2] and [60] for $C^{1,1}$ regularity; see [20, Example 2], [60, (1.3), P. 88] and [132, Example 3 and Remark 1] for counterexamples).

In conclusion, by transforming the Monge-Ampère equation to a locally uniformly elliptic equation, we can obtain the best expected regularity, which may imply that this viewpoint is essential. That the Monge-Ampère equation can be transformed to a locally uniformly elliptic equation has been noted before (e.g. [135, P. 673, L. 1]). We learned this from a note written by Prof. Chuanqiang Chen.

Remark 3.6. There are several criterions to ensure that u is strictly convex in a general domain Ω , such as

- u is an affine function on $\partial\Omega$ (see [21, Corollary 2], [51, Theorem 4.10] and [120, Lemma 3.4]).
- $u \in C^{1,\alpha}(\partial\Omega)$ ($\alpha > 1 2/n$) (see [21, Corollary 4], [51, Corollary 4.11] and [120, Remark 3.1]).
- $u \in W^{2,p}(\Omega)$ (p > n(n-1)/2) (see [112] and [123]).

Remark 3.7. Another important notion of weak solution is the Alexandrov's generalized solution (see [1] and [51, Definition 2.5]). The right-hand term fis not necessarily continuous if we use this notion. These two definitions are equivalent if f is continuous (see [61, Proposition 1.3.4, Proposition 1.7.1]).

Theorem 3.1 holds for Alexandrov's generalized solutions as well. In the following, we make an explanation. Since u is strictly convex, without loss of generality, we can assume that u is an Alexandrov's generalized solution of

$$\begin{cases} \det D^2 u = f & \text{in } \Omega; \\ u = 0 & \text{on } \partial \Omega \end{cases}$$

where Ω is a convex domain and $B_{1/n} \subset \Omega \subset B_1$. Choose a smooth function φ with compact support in B_1 to mollify f by convolution: (see [150, Lemma 3.5.6])

$$f_{\varepsilon} := f * \varphi_{\varepsilon}, \quad \varphi_{\varepsilon}(x) := \varepsilon^{-n} \varphi(x/\varepsilon), \quad \varepsilon > 0$$

such that

 $P = P * \varphi_{\varepsilon}, \ \forall \ P \in \mathcal{P}_{k-2}, \ \varepsilon > 0.$

Then $f_{\varepsilon} \in C^{k-2,\alpha}(0)$ with $P_{f_{\varepsilon}} \equiv P_f$. Let u_{ε} be Alexandrov's solutions (be viscosity solutions as well) of (see [51, Theorem 2.13])

$$\begin{cases} \det D^2 u_{\varepsilon} = f_{\varepsilon} & \text{in } \Omega; \\ u_{\varepsilon} = 0 & \text{on } \partial \Omega. \end{cases}$$

Since $f_{\varepsilon} \to f$ in $L^1(\Omega)$, we have (see [51, Proposition 2.16])

$$u_{\varepsilon} \to u \quad \text{in } L^{\infty}(\Omega).$$

By Theorem 3.1, $u_{\varepsilon} \in C^{k,\alpha}(0)$. That is, there exist $P_{\varepsilon} \in \mathcal{P}_k$ such that

(3.1)
$$|u_{\varepsilon}(x) - P_{\varepsilon}(x)| \le C|x|^{k+\alpha}, \ \forall \ x \in \Omega$$

and

$$\|P_{\varepsilon}\| \le C,$$

where C is independent of ε . Hence, up to a subsequence, there exists $P \in \mathcal{P}_k$ such that $P_{\varepsilon} \to P$. Let $\varepsilon \to 0$ in (3.1) and then

$$|u(x) - P(x)| \le C|x|^{k+\alpha}, \ \forall \ x \in \Omega.$$

That is, $u \in C^{k,\alpha}(0)$.

We first prove a lemma.

Lemma 3.8. Let $B_{1/n} \subset \Omega \subset B_1$ and u be a viscosity solution of

$$\begin{cases} \det D^2 u = f & \text{ in } \Omega; \\ u = 0 & \text{ on } \partial \Omega \end{cases}$$

Assume that 0 is the minimum point of u.

Then for any $\delta_1 > 0$, there exist $\delta, r > 0$ (depending only on n and δ_1) such that for some $P \in \mathcal{P}_2$ with $D^2 P \ge 0$,

$$|u - P||_{L^{\infty}(B_r)} \le \delta_1 r^2$$

and

(3.2)
$$\det D^2 P = 1, \quad ||P|| \le C,$$

provided

$$\|f-1\|_{L^{\infty}(\Omega)} \le \delta,$$

where C depends only on n.

Proof. Suppose not. Then there exist $\delta_1 > 0$, sequences of u_m, Ω_m and f_m such that $B_{1/n} \subset \Omega_m \subset B_1$,

$$\begin{cases} \det D^2 u_m = f_m & \text{in } \Omega_m; \\ u_m = 0 & \text{on } \partial \Omega_m, \end{cases}$$

and

$$\|f_m - 1\|_{L^{\infty}(\Omega_m)} \le 1/m$$

Moreover, for any convex polynomial $P \in \mathcal{P}_2$ satisfying (3.2), we have

$$(3.3) ||u_m - P||_{L^{\infty}(B_r)} \ge \delta_1 r^2,$$

where r is to be specified later.

By the stability of solutions (see [51, Corollary 2.12]), up to a subsequence, there exist $\bar{u}, \tilde{\Omega}$ such that

$$u_m \to \bar{u}, \quad \Omega_m \to \Omega, \quad B_{1/n} \subset \Omega \subset B_1,$$

Moreover, \bar{u} is the solution of

$$\begin{cases} \det D^2 \bar{u} = 1 & \text{ in } \tilde{\Omega}; \\ \bar{u} = 0 & \text{ on } \partial \tilde{\Omega} \end{cases}$$

Since 0 is the minimum point of u_m , we have (see [51, Proposition 4.4])

$$d(0,\partial\Omega_m) \ge c_0 > 0,$$

where $c_0 > 0$ depends only on *n*. Hence,

$$d(0,\partial\Omega) \ge c_0.$$

By the interior regularity (see [51, Theorem 3.10]), there exists a convex polynomial $P \in \mathcal{P}_2$ such that (3.2) holds and

$$|\bar{u}(x) - P(x)| \le C|x|^3, \ \forall \ x \in B_{c_0/2}.$$

Take r small such that

$$Cr = \frac{\delta_1}{2}.$$

Then

(3.4)
$$\|\bar{u} - P\|_{L^{\infty}(B_r)} \leq \frac{1}{2}\delta_1 r^2.$$

By taking the limit in (3.3), we have

$$\|\bar{u} - P\|_{L^{\infty}(B_{r_0})} \ge \delta_1 r^2,$$

which contradicts with (3.4).

Now, we give the

Proof of Theorem 3.1. By subtracting an affine function, we may assume that $u \ge 0$ and u(0) = 0. Since u is strictly convex, for h > 0 small (to be specified later), $S_h(0) \subset B_1$, where $S_h(0)$ the section of u at 0, i.e.

 $S_h(0) := \{ x \in B_1 : u(x) < h \}.$

With the aid of John's lemma (see [71] and [51, A.3.2]), we normalize the section $S_h(0)$ as follows:

(3.5)
$$y = T_h x$$
, $\tilde{\Omega} = T_h(S_h(0))$, $B_{1/n}(\tilde{y}) \subset \tilde{\Omega} \subset B_1(\tilde{y})$, $\tilde{u}(y) = \frac{u(x)}{h}$,

where $T_h \in \mathcal{S}^n$ and $\tilde{y} \in B_1$. Then \tilde{u} is a solution of

$$\begin{cases} \det D^2 \tilde{u} = \tilde{f} & \text{in } \tilde{\Omega}; \\ \tilde{u} = 1 & \text{on } \partial \tilde{\Omega}, \end{cases}$$

where

$$\tilde{f}(y) = \frac{f(x)}{(\det T_h)^2 h^n}.$$

By the uniform estimate (see [120, Lemma 3.2]),

$$(3.6) C_1 \le (\det T_h)^2 h^n \le C_2,$$

where C_1, C_2 depends only on n, λ, Λ . Without loss of generality, we assume $\tilde{f}(0) = 1$. Since $f \in C^{k-2,\alpha}(0)$,

$$|\tilde{f}(y) - 1| \le C|f(x) - f(0)| \le C|x|^{\alpha} \le C|T_h^{-1}| \cdot |y|^{\alpha}.$$

Since u is strictly convex,

$$(3.7) |T_h^{-1}| \to 0 \text{as } h \to 0.$$

Let $0 < \delta_1 < 1$ to be determined later. By (3.7), we can take h small enough such that

(3.8)
$$C\|T_h^{-1}\| \le \delta_1, \quad \|\tilde{f} - 1\|_{L^{\infty}(\tilde{\Omega})} \le \delta,$$

where $\delta \leq \delta_1$ is chosen small enough such that Lemma 3.8 holds. Then by Lemma 3.8, there exist r > 0 and a convex polynomial $P \in \mathcal{P}_2$ such that

(3.9)
$$\|\tilde{u} - P\|_{L^{\infty}(B_r)} \le \delta_1 r^2$$

and

$$\det D^2 P = 1, \quad \|P\| \le C.$$

Let

$$z = \frac{y}{r}, \quad \hat{u}(z) = \frac{\tilde{u}(y) - P(y)}{r^2}$$

Then \hat{u} is a solution of

(3.10)
$$F(D^2\hat{u}) = \hat{f} \text{ in } B_1,$$

where

$$F(M) = \det(M + D^2 P) - 1, \ \forall \ M \in S^n, \quad \hat{f}(z) = \tilde{f}(y) - 1.$$

Note that $F \in C^{\infty}$, F(0) = 0 and F is ρ -uniformly elliptic with ellipticity constants $\tilde{\lambda}, \tilde{\Lambda}$ and they depend only on n, λ, Λ .

By (3.8), (3.9) and the definition of f,

$$\|\hat{u}\|_{L^{\infty}(B_1)} \le \delta_1, \quad \|\hat{f}\|_{C^{\alpha}(0)} \le \delta \le \delta_1.$$

From Theorem 1.7 and Corollary 1.12 (choosing δ_1 small enough), $\hat{u} \in C^{k,\alpha}(0)$ and hence $u \in C^{k,\alpha}(0)$.

Remark 3.9. Two cornerstone results for Theorem 3.1 are a priori estimates/existence of classical solutions and Pogorelov's type estimate. They are implicitly used in Lemma 3.8. We refer to [51, Theorem 3.10] for details.

Remark 3.10. For the Monge-Ampère equation, we obtain almost the same $C^{k,\alpha}$ regularity as the Poisson's equation. However, for the prescribed mean curvature equation (and other types of equations below), we must assume that u (or f) is small or u is uniformly elliptic at 0 (i.e. $u \in C^{0,1}$). The reason is that for the Monge-Ampère equation, we can make an anisotropic scaling such that the equation is unchanged. This is the unique feature of the Monge-Ampère equation.

For the prescribed mean curvature equation, to make f small and the equation unchanged, we have to make the following scaling (see (2.14)):

(3.11)
$$y = \frac{x}{r}, \quad \tilde{u}(y) = \frac{u(x) - u(0)}{r}$$

Thus, to guarantee that \tilde{u} is bounded, we must assume that $u \in C^{0,1}(0)$.

On the contrast, for the Monge-Ampère equation, to make f small and the equation unchanged, we can make an anisotropic scaling (see (3.5)):

(3.12)
$$y = T_h x, \quad \tilde{\Omega} = T_h(S_h(0)), \quad \tilde{u}(y) = \frac{u(x)}{h},$$

where T_h is anisotropic, i.e., its eigenvalues may be not comparable. Note that $h \simeq r^2$ in the sense of anisotropy (see (3.6)). Hence, the scaling transformation (3.12) is essentially the same to (3.11) except that (3.12) is anisotropic. Maybe this is the reason that there are few pointwise Schauder regularity for non-uniformly equations except the Monge-Ampère equation.

4. *k*-Hessian equations

Next, we consider the k-Hessian equations. For $u \in C^2$, denote its eigenvalues by

$$\lambda(D^2 u) = (\lambda_1, \cdots, \lambda_n) \in \mathbb{R}^n.$$

Define for $1 \le k \le n$

$$\sigma_k(D^2u) = \sigma_k(\lambda(D^2u)) = \sum_{1 \le i_1 < \dots < i_k \le n} \lambda_{i_1} \cdots \lambda_{i_k}$$

We say u is k-admissible if $\lambda(D^2 u) \in \Gamma_k$, where Γ_k is the Gårding convex cone in \mathbb{R}^n defined by

$$\Gamma_k \equiv \left\{ \lambda \in \mathbb{R}^n \middle| \sigma_i(\lambda) > 0, \quad \forall \ 1 \le i \le k \right\}.$$

For more basic knowledge of the k-Hessian equations, we refer to [134]. We consider the k-Hessian equation:

$$\sigma_k(D^2 u) = f \text{ in } B_1.$$

Similar to the Monge-Ampère equation, for any k-admissible polynomial $P \in \mathcal{P}_2$ with $\sigma_k(D^2 P) = 1$, define

$$F(M) = \sigma_k(M + D^2 P) - 1, \ \forall \ M \in \mathcal{S}^n.$$

Then $F \in C^{\infty}, F(0) = 0$ and F is ρ -uniformly elliptic with $\tilde{\lambda}, \tilde{\Lambda}$ which depend only on n, k and $|D^2P|$.

Now, we state the main results in this section. Since the k-Hessian equation reduces to the Monge-Ampère equation if the dimension n = 2, we only consider $n \ge 3$ in this section.

Theorem 4.1. Let $2 \le k \le n-1$, $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of

(4.1)
$$\sigma_k(D^2u) = f \quad in \ B_1$$

where $0 < \lambda \leq f \leq \Lambda$. Then $u \in C^{l,\alpha}(0)$ $(l \geq 2)$ provided one of the following conditions holds :

(i) there exists a k-admissible $P \in \mathcal{P}_2$ such that

$$||u - P||_{L^{\infty}(B_1)} \le \delta, \quad \sigma_k(D^2 P) = f(0), \quad f \in C^{l-2,\alpha}(0), \quad [f]_{C^{\alpha}(0)} \le \delta,$$

where $0 < \delta < 1$ depends only on $n, k, \lambda, \Lambda, \alpha$ and $|D^2P|$. (ii) there exists a k-admissible $P \in \mathcal{P}_2$ such that

$$||u - P||_{L^{\infty}(B_1)} \le \delta, \quad \sigma_k(D^2 P) = f(0), \quad f \in C^{l-2,\alpha}(0),$$

where $0 < \delta < 1$ depends only on $n, k, \lambda, \Lambda, \alpha, |D^2P|$ and $[f]_{C^{\alpha}(0)}$. (iii) $u \in C^2(0)$ and $f \in C^{l-2,\alpha}(0)$.

(iv) there exists p > k(n-1)/2 such that

$$u \in W^{2,p}(B_1), \quad f \in C^{l-2,\alpha}(0), \quad [f]_{C^{\alpha}(0)} \le \delta,$$

where δ depends only on $n, k, \lambda, \Lambda, \alpha$ and $||u||_{W^{2,p}(B_1)}$. (v) there exists a k-admissible $P \in \mathcal{P}_2$ such that

u = P on ∂B_{δ} , $\sigma_k(D^2 P) = f(0)$, $f \in C^{l-2,\alpha}(0)$,

where δ depends only on $n, k, \lambda, \Lambda, \alpha$ and $|D^2P|$.

Chaudhuri and Trudinger [31] proved that if k > n/2, $u \in C^2(x_0)$ for *a.e.* x_0 . Hence, we have the following corollary of (iii).

Corollary 4.2. Let k > n/2, $l \ge 2, 0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of

(4.2)
$$\sigma_k(D^2u) = f \quad in \ B_1,$$

where $0 < \lambda \leq f \leq \Lambda$ and $f \in C^{l-2,\alpha}(\bar{B}_1)$. Then $u \in C^{l,\alpha}(x_0)$ for a.e. $x_0 \in B_1$.

Remark 4.3. We point out that $\sigma_k(D^2P) = f(0)$ is not needed. In fact, with the aid of the definition of viscosity solution, we can modify P such that this equality holds if δ is small enough.

We do not know whether the $C^{2,\alpha}$ regularity holds if we only assume $u \in C^{1,1}(0)$ instead of $u \in C^2(0)$ in (iii). On the other hand, $u \in C^2(0)$ can be relaxed to (by a similar proof to that of Theorem 4.4)

$$K_1|x|^2 \le u - P \le K_1|x|^2$$
 in B_1 ,

where $K_1, K_2 > 0$ and $P \in \mathcal{P}_1$. A similar condition has been used by Wang and Bao [125, Theorem 1.1] to prove the rigidity in \mathbb{R}^n for the k-Hessian equation.

Urbas [122] (see also [112, 121] for similar estimates) proved the interior $C^{1,1}$ estimate if p > k(n-1)/2 and $f \in C^{1,1}$. This is the cornerstone of the regularity under the assumption (iv).

The condition (v) concerns the regularity in a small domain. In this direction, Urbas [124, Theorem 3] proved global C^1 estimate. Tian, Wang and Wang [109] studied the local solvability of the k-Hessian equation in B_{r_0} for r_0 small. Maybe Theorem 4.1 can be applied to study the local solvability.

Since the k-Hessian equations do not possess pure interior $C^{1,1}$ estimate, we do not have a similar regularity by assuming only f - f(0) small as in (ii) Theorem 2.3. Instead, we have the following regularity based on a priori estimates and the Pogorelov's type estimate.

Theorem 4.4. Let $2 \le k \le n-1$, $0 < \alpha < 1$ and $u \in C(\overline{\Omega})$ be a viscosity solution of

$$\begin{cases} \sigma_k(D^2 u) = f & \text{ in } \Omega; \\ u = g & \text{ on } \partial\Omega, \end{cases}$$

where $0 < \lambda \leq f \leq \Lambda$. Then $u \in C^{l,\alpha}(0)$ $(l \geq 2)$ provided one of the following conditions holds : (i) $\partial \Omega \in C^{3,1}$, Ω is (k-1)-convex and

$$||g - g_0||_{L^{\infty}(\partial\Omega)} \le \delta, \ g_0 \in C^{3,1}(\partial\Omega), \ f \in C^{l-2,\alpha}(0), \ [f]_{C^{\alpha}(0)} \le \delta,$$

where $0 < \delta < 1$ depends only on $n, k, \lambda, \Lambda, \alpha, \Omega$ and $||g_0||_{C^{3,1}(\partial\Omega)}$. (ii) Ω is a (k-1)-convex domain and

$$\|g - P\|_{L^{\infty}(\partial\Omega)} \le \delta, \ P \in \mathcal{P}_1, \ f \in C^{l-2,\alpha}(0), \ [f]_{C^{\alpha}(0)} \le \delta,$$

where $0 < \delta < 1$ depends only on $n, k, \lambda, \Lambda, \alpha$ and Ω . (iii)

$$\Omega = B_{\delta}, \ \|g - P\|_{L^{\infty}(\partial B_{\delta})} \le \delta_1 \delta^2, \ P \in \mathcal{P}_1, \ f \in C^{l-2,\alpha}(0),$$

where δ_1 depends only on $n, k, \lambda, \Lambda, \alpha$ and δ depends also on $[f]_{C^{\alpha}(0)}$.

Remark 4.5. The regularity under assumption (i) is based on a priori estimates for sufficient smooth data, which was first proved by Caffarelli, Nirenberg and Spruck [19] (see also [113] and [134, Theorem 3.4]). The regularity under assumption (ii) is based on the Pogorelov's type estimate, established by Chou and Wang [42, Theorem 1.5] (see also [134, Theorem 4.3]). In (ii), $\partial\Omega$ is not necessary to be smooth. The (k - 1)-convexity can be defined in some weak sense (e.g. by an approximation, see [116, P. 226, L. 4]; in the viscosity sense, see [117, P. 580]).

Remark 4.6. There are other types of weak solutions, e.g., the weak solution defined by an approximation (see [114], [121], [122]) and the weak solution based on the Borel measure (see [116, 117, 118]). As before, Theorem 4.1 and Theorem 4.4 hold with these weak solutions by an approximation argument.

Remark 4.7. We do not know any counterexample for the k-Hessian equation to demonstrate the necessity of the "smallness" as in the prescribed mean curvature equation (see Remark 2.6), the k-Hessian quotient equations (see Remark 5.4) and the Lagrangian mean curvature equation (see Remark 6.5). Hence, it is interesting to construct a counterexample, e.g., a strictly convex viscosity solution u of (4.1) with $f \in C^{\alpha}$ but $u \notin C^{2,\alpha}$. Or, can we prove Theorem 4.1 and Theorem 4.4 without the smallness assumptions? Note that the counterexample constructed by Pogorelov and its extensions (see [26, 29, 91], [121, P. 6], [124] etc.) are not strictly convex.

We first prove a lemma.

Lemma 4.8. Let u be a viscosity solution of

 $\sigma_k(D^2u) = f \quad in \ B_1.$

Assume that $u \in W^{2,p}(B_1)$ (p > k(n-1)/2).

Then for any $\delta_1 > 0$ and K > 0, there exist $\delta, r > 0$ (depending only on n, δ_1 and K) such that for some k-admissible $P \in \mathcal{P}_2$,

$$\|u - P\|_{L^{\infty}(B_r)} \le \delta_1 r^2$$

and

(4.3)
$$\sigma_k(D^2P) = 1, \quad ||P|| \le C,$$

provided

$$|u||_{W^{2,p}(B_1)} \le K, \quad ||f-1||_{L^{\infty}(B_1)} \le \delta,$$

where C depends only on n, k, p and K.

Proof. Suppose not. Then there exist $\delta_1, K > 0$, sequences of u_m, f_m such that

$$\sigma_k(D^2 u_m) = f_m \text{ in } B_1$$

and

$$|u_m||_{W^{2,p}(B_1)} \le K, \quad ||f_m - 1||_{L^{\infty}(B_1)} \le \frac{1}{m}.$$

Moreover, for any k-admissible $P \in \mathcal{P}_2$ satisfying (4.3), we have

(4.4)
$$||u_m - P||_{L^{\infty}(B_r)} \ge \delta_1 r^2,$$

where r is to be specified later.

Since $n \ge 3$ and $k \ge 2$, we have p > k(n-1)/2 > n/2 and thus for some $\beta > 0$,

$$\|u_m\|_{C^{\beta}(\bar{B}_1)} \le C \|u_m\|_{W^{2,p}(B_1)} \le CK.$$

Then up to a subsequence, there exist \bar{u} such that

$$u_m \to \bar{u} \quad \text{in } L^{\infty}(B_1), \quad \|\bar{u}\|_{W^{2,p}(B_1)} \le K.$$

Moreover, \bar{u} is a viscosity solution of

$$\sigma_k(D^2\bar{u}) = 1 \text{ in } B_1.$$

Hence, \bar{u} is smooth (see [122, Theorem 1.1] and there exists k-admissible $P \in \mathcal{P}_2$ such that (4.3) holds and

$$|\bar{u}(x) - P(x)| \le C|x|^3, \ \forall \ x \in B_{1/2}.$$

Take r small such that $Cr = \delta_1/2$. Then

$$\|\bar{u} - P\|_{L^{\infty}(B_r)} \le \frac{1}{2}\delta_1 r^2.$$

By taking the limit in (4.4), we have

$$\|\bar{u} - P\|_{L^{\infty}(B_r)} \ge \delta_1 r^2,$$

which is contradiction.

Now, we give the

Proof of Theorem 4.1. The proof is similar to that of the prescribed

mean curvature equation. We assume f(0) = 1 without loss of generality. For (i), let $\tilde{u} = u - P$ and then \tilde{u} is a solution of

$$F(D^2\tilde{u}) = \tilde{f} \quad \text{in } B_1,$$

where

(4.5)
$$F(M) := \sigma_k (M + D^2 P) - 1, \ \forall \ M \in \mathcal{S}^n, \quad \tilde{f} := f - 1.$$

Thus, $F \in C^{\infty}$, F(0) = 0 and F is ρ -uniformly elliptic with ellipticity constants $\tilde{\lambda}, \tilde{\Lambda}$ and they depend only on n, k, λ, Λ and $|D^2P|$. In addition, by assumption (i),

$$\|\tilde{u}\|_{L^{\infty}(B_1)} \le \delta, \quad \|f\|_{C^{\alpha}(0)} \le \delta.$$

Therefore, by Corollary 1.12, $\tilde{u} \in C^{l,\alpha}(0)$ and hence $u \in C^{l,\alpha}(0)$. Next, we prove (iii) For n > 0, let

Next, we prove (ii). For r > 0, let

(4.6)
$$y = \frac{x}{r}, \quad \tilde{u}(y) = \frac{u(x) - P(x)}{r^2}, \quad \tilde{f}(y) = f(x) - 1.$$

Then \tilde{u} is a solution of

$$F(D^2\tilde{u}) = \tilde{f}$$
 in B_1 .

We first choose r small enough such that $\|f\|_{C^{\alpha}}$ is small. Next, take δ small such that $\|\tilde{u}\|_{L^{\infty}}$ is small. Then the conclusion follows from Corollary 1.12.

Note that (iii) is a direct consequence of Theorem 2.1.

As regards (iv), by Lemma 4.8, for any $\delta_1 > 0$, there exist k-admissible $P \in \mathcal{P}_2$ and r > 0 such that,

$$\|u - P\|_{L^{\infty}(B_r)} \le \delta_1 r^2.$$

Then the conclusion follows as above through a transformation like (4.6).

Finally, we prove (v). By the comparison estimate for k-Hessian estimate (see [114, Lemma 2.1]),

$$||u - P||_{L^{\infty}(B_{\delta})} \le C\delta^2 ||f^{1/k} - 1||_{L^{\infty}(B_{\delta})} \le C\delta^{2+\alpha}$$

where C depends only on n, k. Then the conclusion follows as above. \Box

Next, we give the

Proof of Theorem 4.4. We assume f(0) = 1 as before. For (i), since $\partial\Omega$ and g_0 are smooth, there exists a solution $v \in C^3(\overline{\Omega})$ of (see [113, Theorem 1.1], [134, Theorem 3.4])

$$\begin{cases} \sigma_k(D^2 v) = 1 & \text{ in } \Omega; \\ v = g_0 & \text{ on } \partial \Omega \end{cases}$$

By the comparison estimate for k-Hessian estimate (see [114, Lemma 2.1]),

(4.7)
$$||u-v||_{L^{\infty}(\Omega)} \leq ||g-g_0||_{L^{\infty}(\partial\Omega)} + C \operatorname{diam}(\Omega)^2 ||f^{1/k}-1||_{L^{\infty}(\Omega)} \leq C \delta^{\alpha}$$
,
where C depends only on n, k and Ω .

In addition, by the interior C^3 estimate for v, there exists a k-admissible $P \in \mathcal{P}_2$ such that

(4.8)
$$|v(x) - P(x)| \le C|x|^3, \ \forall x \in B_{r_0}$$

and

(4.9)
$$\sigma_k(D^2P) = 1, \quad ||P|| \le C$$

where r_0 depends only on Ω and C depends only on $n, k, \Omega, ||g_0||_{C^{3,1}(\partial\Omega)}$.

Let $0 < \delta_1 < 1$ to be specified later. Take δ and r small enough such that

$$Cr \le \frac{\delta_1}{2}, \quad C\delta^{\alpha} \le \frac{\delta_1}{2}r^2.$$

Then by (4.7) and (4.8),

(4.10)
$$||u - P||_{L^{\infty}(B_r)} \le \delta_1 r^2.$$

Therefore, the conclusion follows as before (cf. the proof of Theorem 4.1).

The proof for (ii) is similar to the above. By subtracting P, we may assume that $|u| \leq \delta$ on $\partial\Omega$. Let $v \in C^3(\Omega) \cap C(\overline{\Omega})$ be the solution of (see [134, Theorem 4.4])

$$\begin{cases} \sigma_k(D^2 v) = 1 & \text{in } \Omega; \\ v = 0 & \text{on } \partial \Omega \end{cases}$$

By the comparison estimate,

(4.11)
$$||u - v||_{L^{\infty}(\Omega)} \le ||u||_{L^{\infty}(\partial\Omega)} + C||f^{1/k} - 1||_{L^{\infty}(\partial\Omega)} \le C\delta^{\alpha}.$$

By the interior C^3 estimate for v, there exist $r_0 > 0$ and a k-admissible $P \in \mathcal{P}_2$ such that

$$|v(x) - P(x)| \le C|x|^3, \ \forall \ x \in B_{r_0}$$

and

$$\sigma_k(D^2P) = 1, \quad ||P|| \le C.$$

Then the conclusion follows as above.

For (iii), with the aid of the transformation

$$y = \frac{x}{\delta}, \quad \tilde{u}(y) = \frac{u(x) - P(x)}{\delta^2},$$

the conclusion follows from (ii).

Since the 2-Hessian equation has pure interior $C^{1,1}$ estimate in some circumstances, we have the following additional regularity for this equation.

Theorem 4.9. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of

(4.12)
$$\sigma_2(D^2u) = f \quad in \ B_1 \subset \mathbb{R}^n.$$

where $0 < \lambda \leq f \leq \Lambda$. Then $u \in C^{l,\alpha}(0)$ $(l \geq 2)$ provided one of the following conditions holds : (i)

$$n = 3, 4, \quad f \in C^{l-2,\alpha}(0), \quad [f]_{C^{\alpha}(0)} \le \delta,$$

28

where $0 < \delta < 1$ depends only on $n, \lambda, \Lambda, \alpha$ and $||u||_{L^{\infty}(B_1)}$. (ii)

$$n = 3, 4, \quad u \in C^{1,1}(0), \quad f \in C^{l-2,\alpha}(0).$$

(iii) there exists $\varepsilon > 0$ such that $u + ((n(n-1)/2)^{-1/2} - \varepsilon) |x|^2$ is convex and

$$f \in C^{l-2,\alpha}(0), \quad [f]_{C^{\alpha}(0)} \le \delta,$$

where $0 < \delta < 1$ depends only on $n, \lambda, \Lambda, \alpha, \varepsilon$ and $||u||_{L^{\infty}(B_1)}$. (iv) there exists $\varepsilon > 0$ such that $u + ((n(n-1)/2)^{-1/2} - \varepsilon) |x|^2$ is convex and

$$u \in C^{1,1}(0), \quad f \in C^{l-2,\alpha}(0).$$

Remark 4.10. The regularity under assumptions (i) and (ii) is based on the pure interior $C^{1,1}$ estimate for the 2-Hessian equation when $f \equiv 1$. For n = 3, it was derived by Warren and Yuan [138]. For n = 4, it was derived by Shankar and Yuan [106].

The regularity under assumptions (iii) and (iv) is based on the pure interior $C^{1,1}$ estimates for almost convex viscosity solutions. Mooney [93] proved the convex case; Shankar and Yuan [105] proved the almost convex case. We also note that the interior $C^{1,1}$ estimate holds as well for *smooth* semi-convex solutions (see [59], [104]). However, we can not obtain a regularity result by an approximation argument based on these estimates. This has been pointed out in [93, P. 2474, L. 2] and [105, P. 2].

Remark 4.11. The regularity under assumptions (i), (iii) (resp. (ii), (iv)) is analogous to (ii) (resp. (iii)) in Theorem 2.3 since we have pure interior $C^{1,1}$ estimates for constant f. A regularity result similar to (i) was proved by Xu [140] in dimension 3. The interior $C^{2,\alpha}$ regularity in dimension 3 with $f \in C^{0,1}$ was obtained by Zhou [148, Theorem 1.4].

We first give a lemma.

Lemma 4.12. Let $u \in C(\overline{B}_1)$ be a viscosity solution of

$$\sigma_2(D^2u) = f \quad in \ B_1.$$

and $||u||_{L^{\infty}(B_1)} \leq K$. Suppose that one of the following two conditions holds: (i) n = 3 or 4;

(ii) there exists $\varepsilon > 0$ such that $u + \left((n(n-1)/2)^{-1/2} - \varepsilon \right) |x|^2$ is convex.

Then for any $\delta_1 > 0$, there exist $\delta, r > 0$ depending only on n, K, ε and δ_1 such that if $||f - 1||_{L^{\infty}(B_1)} \leq \delta$, we have

$$||u - P||_{L^{\infty}(B_r)} \le \delta_1 r^2$$

and

(4.13)
$$\sigma_2(D^2P) = 1, \quad ||P|| \le C,$$

where $P \in \mathcal{P}_2$ is 2-admissible and C depends only on n, K, ε .

Proof. We only give the proof under the condition (ii). Suppose not. Then there exist $K, \delta_1 > 0$ and sequences of u_m, f_m such that

$$\sigma_2(D^2 u_m) = f_m \quad \text{in } B_1,$$

$$||u_m||_{L^{\infty}(B_1)} \le K, \quad ||f_m - 1||_{L^{\infty}(B_1)} \le 1/m$$

and $u_m + ((n(n-1)/2)^{-1/2} - \varepsilon) |x|^2$ is convex. However, for any 2-admissible $P \in \mathcal{P}_2$ satisfying (4.13), we have

(4.14)
$$||u_m - P||_{L^{\infty}(B_r)} > \delta_1 r^2,$$

where r is to be specified later.

By the interior Hölder regularity (see [114, Theorem 4.1], [134, corollary 9.1]), u_m are uniformly bounded and equicontinuous in any compact subset of B_1 . Then up a subsequence, there exists \bar{u} such that

$$u_m \to \bar{u}$$
 in $L^{\infty}_{loc}(B_1)$.

Then $\bar{u} + \left((n(n-1)/2)^{-1/2} - \varepsilon \right) |x|^2$ is convex and \bar{u} is a viscosity solution of

$$\sigma_2(D^2\bar{u}) = 1 \quad \text{in } B_1$$

Thus, \bar{u} is smooth (see [105, Theorem 1.1]) and there exists a 2-admissible $P \in \mathcal{P}_2$ such that (4.13) holds and

$$|\bar{u}(x) - P(x)| \le C|x|^3, \ \forall \ x \in B_{1/2},$$

where C depends only on n and K. Take r small such that $Cr = \delta_1/2$. Then

$$\|\bar{u} - P\|_{L^{\infty}(B_r)} \le \frac{1}{2}\delta_1 r^2.$$

By taking the limit in (4.14), we have

$$\|\bar{u} - P\|_{L^{\infty}(B_r)} \ge \delta_1 r^2,$$

which is a contradiction.

Proof of Theorem 4.9. With the aid of Lemma 4.12, the theorem can be proved in a similar way as before and we omit it. \Box

5. k-Hessian quotient equation

In this section, we consider the k-Hessian quotient equations:

$$S_{k,l}(D^2u) := \frac{\sigma_k(D^2u)}{\sigma_l(D^2u)} = f \text{ in } B_1,$$

where $1 \leq l < k \leq n$. Similar to the k-Hessian equation, for any k-admissible polynomial $P \in \mathcal{P}_2$ with $S_{k,l}(D^2P) = 1$, define

$$F(M) = S_{k,l}(M + D^2 P) - 1, \ \forall \ M \in \mathcal{S}^n.$$

Then $F \in C^{\infty}, F(0) = 0$ and F is ρ -uniformly elliptic with $\tilde{\lambda}, \tilde{\Lambda}$ which depend only on n, k, l and $|D^2P|$.

The main results are the following.

Theorem 5.1. Let $1 \leq l < k \leq n$, $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of

(5.1)
$$S_{k,l}(D^2u) = f \text{ in } B_1$$

where $0 < \lambda \leq f \leq \Lambda$. Then $u \in C^{m,\alpha}(0)$ $(m \geq 2)$ provided one of the following conditions holds :

(i) there exists a k-admissible $P \in \mathcal{P}_2$ such that

$$||u - P||_{L^{\infty}(B_1)} \le \delta, \quad S_{k,l}(D^2 P) = f(0), \quad f \in C^{m-2,\alpha}(0), \quad [f]_{C^{\alpha}(0)} \le \delta,$$

where $0 < \delta < 1$ depends only on $n, k, l, \lambda, \Lambda, \alpha$ and $|D^2P|$. (ii) there exists a k-admissible $P \in \mathcal{P}_2$ such that

$$||u - P||_{L^{\infty}(B_1)} \le \delta, \quad S_{k,l}(D^2 P) = f(0), \quad f \in C^{m-2,\alpha}(0),$$

where $0 < \delta < 1$ depends only on $n, k, l, \lambda, \Lambda, \alpha, |D^2P|$ and $[f]_{C^{\alpha}(0)}$. (iii) $u \in C^2(0)$ and $f \in C^{m-2,\alpha}(0)$.

(iv) there exists a k-admissible $P \in \mathcal{P}_2$ such that

u = P on ∂B_{δ} , $\sigma_k(D^2 P) = f(0)$, $f \in C^{m-2,\alpha}(0)$,

where δ depends only on $n, k, l, \lambda, \Lambda, \alpha$ and $|D^2P|$.

Remark 5.2. Since the proof is quite similar to the that for the k-Hessian equation, we omit it. We point out that we need to apply a comparison estimate when proving (iv) (see the proof of (v) in Theorem 4.1). There is no existing literature to cite. Indeed, the comparison estimate for k-Hessian equation (see [114, Lemma 2.2]) can be extended to the k-Hessian quotient equation with the L^p norm in the estimate replaced by the L^{∞} norm. The key is that the operator $(S_{k,l})^{1/(k-l)}$ is 1-homogenous and concave. Then an inequality similar to [114, (2.4)] holds and the comparison estimate can be proved similarly.

For the k-Hessian quotient equations, there are few pure interior $C^{1,1}$ estimates and Pogorelov's type estimates for *smooth convex* solutions (see [44, 89, 91]) until now. Unfortunately, as before, we can not use these to build regularity. Instead, we have the following regularity based on a priori estimates established by Trudinger [113, Theorem 1.1]. Since the proof is similar to that for the k-Hessian equation and we omit it.

Theorem 5.3. Let $1 \leq l < k \leq n$, $0 < \alpha < 1$ and $u \in C(\overline{\Omega})$ be a viscosity solution of

$$\begin{cases} S_{k,l}(D^2u) = f & \text{ in } \Omega; \\ u = g & \text{ on } \partial \Omega. \end{cases}$$

where $0 < \lambda \leq f \leq \Lambda$. Then $u \in C^{m,\alpha}(0)$ $(m \geq 2)$ provided one of the following conditions holds :

(i) $\partial \Omega \in C^{3,1}$, Ω is (k-1)-convex and

 $\begin{aligned} \|g-g_0\|_{L^{\infty}(\partial\Omega)} &\leq \delta, \ g_0 \in C^{3,1}(\partial\Omega), \ f \in C^{m-2,\alpha}(0), \ [f]_{C^{\alpha}(0)} \leq \delta, \end{aligned}$ where $0 < \delta < 1$ depends only on $n, k, l\lambda, \Lambda, \alpha, \Omega$ and $\|g_0\|_{C^{3,1}(\partial\Omega)}$.
(ii)

$$\Omega = B_{\delta}, \ \|g - P\|_{L^{\infty}(\partial B_{\delta})} \le \delta_1 \delta^2, \ P \in \mathcal{P}_1, \ f \in C^{m-2,\alpha}(0),$$

where δ_1 depends only on $n, k, l, \lambda, \Lambda, \alpha$ and δ depends also on $[f]_{C^{\alpha}(0)}$.

Remark 5.4. Similar to the prescribed mean curvature equation, the $C^{2,\alpha}$ regularity for k-Hessian quotient equation can not hold unconditionally. Consider the following counterexample borrowed from [148, P. 2]. For any $1 \leq l < k \leq n$ and $0 < \theta < 1$, define

$$u(x) = \frac{1}{2}|x'|^2 + \frac{1}{1+\theta}|x_n|^{1+\theta}.$$

Then u is a viscosity solution of (5.1) with $f \in C^{1-\theta}$. Similar to the prescribed mean curvature equation, $[f]_{C^{1-\theta}(0)} \simeq \theta^{-1}$. Hence, if θ is smaller, fis smoother but $[f]_{C^{1-\theta}(0)}$ is bigger. Correspondingly, u has lower regularity.

6. LAGRANGIAN MEAN CURVATURE EQUATION

In this section, we consider the Lagrangian mean curvature equation (called special Lagrangian equation for constant f):

(6.1)
$$F(D^2u) = \sum_{i=1}^{n} \arctan \lambda_i = f \quad \text{in } B_1$$

where λ_i are the eigenvalues of $D^2 u$. The f is called phase function. Obviously, there must hold

$$-n\frac{\pi}{2} < f < n\frac{\pi}{2}.$$

We define

(6.2)
$$\varepsilon_f = \inf_{x \in B_1} \min(n\frac{\pi}{2} - f(x), f(x) + n\frac{\pi}{2}).$$

Similar to the prescribed mean curvature equation, $F \in C^{\infty}$ is 1-uniformly elliptic with $\lambda = 1/5, \Lambda = 1$. Moreover, for any $P \in \mathcal{P}_2$, define

$$G(M) = F(M + D^2P) - F(D^2P).$$

Then G(0) = 0 is ρ -uniformly elliptic with $\tilde{\lambda}, \tilde{\Lambda}$ and they depend only on n and $|D^2P|$.

The phase f is divided into three categories by Yuan [144]: critical $(|f| = (n-2)\pi/2)$, subcritical $(|f| < (n-2)\pi/2)$ and supercritical $(|f| > (n-2)\pi/2)$. For critical/supercritical phases, the level set $\{(\lambda_1, ..., \lambda_n) : \sum_{i=1}^n \arctan \lambda_i = c\}$

is convex and then an extended Evans-Krylov theorem [28] can be applied if we assume $u \in C^{1,1}$ a priori. For more knowledge and historic literature with respect to the special Lagrangian equation, we refer to [136] and [145].

Now, we state the main results in this section. The following theorem is similar to the previous ones and we omit its proof.

Theorem 6.1. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of (6.1). Then $u \in C^{k,\alpha}(0)$ $(k \geq 2)$ provided one of the following conditions holds :

(i) there exists $P \in \mathcal{P}_2$ such that

$$||u - P||_{L^{\infty}(B_1)} \le \delta, \quad F(D^2 P) = f(0), \quad f \in C^{k-2,\alpha}(0), \quad [f]_{C^{\alpha}(0)} \le \delta,$$

where $0 < \delta < 1$ depends only on n, α and $|D^2P|$. (ii) there exists $P \in \mathcal{P}_2$ such that

$$||u - P||_{L^{\infty}(B_1)} \le \delta, \quad F(D^2 P) = f(0), \quad f \in C^{k-2,\alpha}(0),$$

where $0 < \delta < 1$ depends only on $n, \alpha, |D^2P|$ and $[f]_{C^{\alpha}(0)}$. (*iii*) $u \in C^2(0)$ and $f \in C^{k-2,\alpha}(0)$. (iv) there exists $P \in \mathcal{P}_2$ such that

$$u = P$$
 on ∂B_{δ} , $F(D^2 P) = f(0)$, $f \in C^{k-2,\alpha}(0)$,

where δ depends only on n, α and $|D^2P|$.

Remark 6.2. We can not obtain $C^{2,\alpha}$ regularity for the special Lagrangian equation without additional assumptions besides $f \in C^{\alpha}$. If f is subcritical, even a constant f, $C^{2,\alpha}$ regularity can not be expected. Nadirashvili and Vlădut[95] constructed counterexamples for $n \geq 3$ with solutions only belonging to $C^{1,1/3}$ for any constant subcritical phase. In fact, for any $\delta > 0$, there exists a viscosity solution $u \notin C^{1,\delta}$, which was given by Wang and Yuan [126]. In [94], Mooney and Savin provided a counterexample such that the solution $u \in C^{0,1}$ but $u \notin C^1$.

If u is a convex viscosity solution, we have the following regularity.

Theorem 6.3. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a convex viscosity solution of (6.1). Suppose that $f \in C^{k-2,\alpha}(0)$ $(k \ge 2)$. Then $u \in C^{k,\alpha}(0)$ provided $u \in C^{1,1}(0)$ or $[f]_{C^{\alpha}(0)} \leq \delta$, where $0 < \delta < 1$ depends only on n, α, ε_f and $||u||_{L^{\infty}(B_1)}.$

Remark 6.4. If f is a constant, Theorem 6.3 has been proved by Chen, Warren and Yuan [36] (for smooth solutions) and Chen, Shankar and Yuan [35] (for viscosity solutions). For nonconstant f and viscosity solutions, Bhattacharya and Shankar proved the following regularity results:

- $f \in C^{2,\alpha} \Longrightarrow u \in C^{4,\alpha}$ (see [13]). $f \in C^2 \Longrightarrow u \in C^{3,\alpha}$ (see [12]).

• $u \in C^{1,\beta}, f \in C^{\alpha} \Longrightarrow u \in C^{2,\alpha}$, where $\beta > (1+\alpha)^{-1}$ (see [12]).

Remark 6.5. Similar to the prescribed mean curvature equation, the smallness condition in Theorem 6.3 can not be removed. Consider the following counterexample borrowed from [10, Remark 1.3] (see also [12, Remark 1.2]). Take n = 2 and $0 < \theta < 1$. Define

$$u(x_1, x_2) = \frac{1}{1+\theta} |x_1|^{1+\theta} + \frac{1}{2}x_2^2, \quad f(x) = \frac{3\pi}{4} - \arctan(\theta^{-1}|x_1|^{1-\theta}).$$

Then u is a strictly convex viscosity solution and $f \in C^{1-\theta}$ is supercritical. However, $u \in C^{1,\theta}(0)$ only.

On the other hand, note that $[f]_{C^{1-\theta}(0)} \simeq \theta^{-1}$. Thus, if θ is smaller, f has more smoothness but $[f]_{C^{1-\theta}(0)}$ is bigger. Correspondingly, u has less regularity. This demonstrate the assertion "smallness is more important than smoothness" again.

If the phase f is critical and supercritical, we can also deduce regularity. We first consider the supercritical case.

Theorem 6.6. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of (6.1). Suppose that for some $\varepsilon > 0$,

$$|f| \ge (n-2)\frac{\pi}{2} + \varepsilon, \quad f \in C^{k-2,\alpha}(0) \ (k \ge 2).$$

Then $u \in C^{k,\alpha}(0)$ provided $u \in C^{1,1}(0)$ or $[f]_{C^{\alpha}(0)} \leq \delta$, where $0 < \delta < 1$ depends only on $n, \alpha, \varepsilon, \varepsilon_f$ and $||u||_{L^{\infty}(B_1)}$.

For the critical case, we have

Theorem 6.7. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of (6.1). Suppose that

$$|f| = (n-2)\frac{\pi}{2}, \quad f \in C^{k-2,\alpha}(0) \ (k \ge 2).$$

Then $u \in C^{k,\alpha}(0)$ provided $[f]_{C^{\alpha}(0)} \leq \delta$, where $0 < \delta < 1$ depends only on $n, \alpha, \varepsilon_f, \|u\|_{L^{\infty}(B_1)}$ and the modulus of continuity of u.

Remark 6.8. If f is a constant, Theorem 6.6 and Theorem 6.7 have been proved by Warren and Yuan [137] (n = 2), [139] (n = 3) and Wang and Yuan [127] $(n \ge 3)$.

For nonconstant $f \in C^{1,1}$, Bhattacharya [10] proved the supercritical case and Lu [90] proved the critical and supercritical cases. Recently, Zhou [147] extended above results to $f \in C^{0,1}$.

Note that the f is supercritical in the counterexample in Remark 6.5. Hence, the smallness condition can not be removed in Theorem 6.6.

Without the convexity assumption on u or the critical/supercritical assumption on f, Yuan [142] proved $C^{2,\alpha}$ regularity for $C^{1,1}$ viscosity solutions and constant phase in dimension 3. Based on this result, we have the following regularity for a general phase f.

Theorem 6.9. Let $0 < \alpha < 1$ and $u \in C^{1,1}(\overline{B}_1)$ be a viscosity solution of (6.1). Suppose that $f \in C^{k-2,\alpha}(0)$ $(k \ge 2)$. Then $u \in C^{k,\alpha}(0)$.

We first give the

Proof of Theorem 6.3. We only give the proof for the case $[f]_{C^{\alpha}(0)} \leq \delta$ since the case $u \in C^{1,1}(0)$ can be transformed to the former case.

Claim: For any $\delta_1 > 0$, if δ small enough, there exist r > 0 and $P \in \mathcal{P}_2$ such that

(6.3)
$$F(D^2P) = f(0)$$

and

(6.4)
$$||u - P||_{L^{\infty}(B_r)} \le \delta_1 r^2.$$

The proof is similar to the previous (e.g. Lemma 4.12). Suppose not. Then there exist $\varepsilon_0, K > 0$ and sequences of u_m, f_m such that u_m, f_m satisfy (6.1) and

$$||u_m||_{L^{\infty}(B_1)} \le K, \quad [f_m]_{C^{\alpha}(0)} \le 1/m, \quad \varepsilon_{f_m} \ge \varepsilon_0.$$

In addition, for any $P \in \mathcal{P}_2$ with (6.3) holding for f_m , we have

(6.5)
$$||u_m - P||_{L^{\infty}(B_r)} > \delta_1 r^2,$$

where 0 < r < 1 is to be specified later.

Since u_m are convex,

$$||u_m||_{C^{0,1}(\bar{B}_{1/2})} \le C ||u_m||_{L^{\infty}(B_1)} \le K.$$

Hence, u_m are uniformly bounded and equicontinuous. Up to a subsequence (similarly in the following argument), there exists \bar{u} such that

$$u_m \to \bar{u}$$
 in $L^{\infty}(B_{1/2})$.

In addition, since $\varepsilon_{f_m} \geq \varepsilon_0$,

$$f_m(0) \to f_0 \in \left[-n\pi/2 + \varepsilon_0, n\pi/2 - \varepsilon_0\right].$$

By combining with $[f_m]_{C^{\alpha}(0)} \to 0$, we conclude that \bar{u} is a viscosity solution of

$$F(D^2\bar{u}) = f_0$$
 in $B_{1/2}$.

From the regularity for constant phases (see [35]), there exists $\bar{P} \in \mathcal{P}_2$ such that $F(D^2\bar{P}) = f_0$ and

$$|\bar{u}(x) - \bar{P}(x)| \le C|x|^3, \ \forall \ x \in B_{1/2},$$

where C depends only on n and K. Take r small such that $Cr = \delta_1/2$. Then

$$\|\bar{u} - \bar{P}\|_{L^{\infty}(B_r)} \le \frac{1}{2}\delta_1 r^2.$$

In addition, since $f_m(0) \to f_0$, we can choose

$$P_m(x) = \bar{P}(x) + \sum_{i=1}^n c_{i,m} x_i^2$$

such that $c_{i,m} \to 0$ and (6.3) holds for P_m and $f_m(0)$. Thus, (6.5) holds for P_m . By taking the limit in (6.5), we have

$$\|\bar{u}-\bar{P}\|_{L^{\infty}(B_r)} \ge \delta_1 r^2,$$

which is a contradiction. Therefore, the claim holds.

Once the claim is proved, the regularity for u follows as before.

Next, we give the

Proof of Theorem 6.6. We only consider the case

$$f > (n-2)\frac{\pi}{2} + \varepsilon, \quad [f]_{C^{\alpha}(0)} \le \delta.$$

We need the following claim as before:

Claim: For any $\delta_1 > 0$, if δ small enough, there exist r > 0 and $P \in \mathcal{P}_2$ such that

$$F(D^2P) = f(0)$$

and

$$\|u - P\|_{L^{\infty}(B_r)} \le \delta_1 r^2.$$

The proof is similar to the proof of Theorem 6.6. We only need to take care of the compactness of solutions. Indeed, since $f > (n-2)\frac{\pi}{2} + \varepsilon$, the $u + K|x|^2$ is convex where $K = \tan(\pi/2 - \varepsilon)$. This assertion can be proved directly by the definition of viscosity solution. Then

$$||u||_{C^{0,1}(\bar{B}_{1/2})} \le C_1 ||u||_{L^{\infty}(B_1)} + C_2,$$

where C_1 depends only on n and C_2 depends also on ε .

Once the compactness of solutions is built, with the aid of regularity for constant supercritical phase (see [127]), the rest proof is quite similar to that of Theorem 6.6 and we omit it.

Next, we give the

Proof of Theorem 6.7. The difficulty lies in that we can not derive the compactness of solutions and we have to rely on the modulus of continuity of u directly. Hence, the constant δ depends on the modulus of continuity of u.

Then the following claim can be proved as above and we omit it. **Claim:** For any $\delta_1 > 0$, if δ small enough, there exist r > 0 and $P \in \mathcal{P}_2$ such that

$$F(D^2P) = f(0)$$

and

$$\|u - P\|_{L^{\infty}(B_r)} \le \delta_1 r^2.$$

36
Then with the aid of regularity for constant critical phase (see [127]), the regularity of u follows as before.

Next, we give the

Proof of Theorem 6.9. Since $u \in C^{1,1}(\overline{B}_1)$, we have the compactness of solutions a priori. Then with the aid of regularity for constant critical phase (see [142]), the regularity of u follows as before.

Remark 6.10. Note we have assumed $u \in C^{1,1}$. Thus, the equation is in fact uniformly elliptic. In addition, the regularity for constant critical phase has been proved in [142]. Therefore, the $C^{2,\alpha}$ regularity can be proved directly by the theory for uniformly elliptic equations (cf. [27, Chapter 8]). However, for higher order pointwise $C^{k,\alpha}$ regularity, we must apply the regularity theory presented in the introduction.

7. ABP maximum principle, weak Harnack inequality, Hölder regularity

In this section, we develop the basic theory for fully nonlinear locally uniformly elliptic equations. We follows almost exactly the strategy of [16, Chapters 3, 4]. The main difficulty is that we can not make the scaling argument arbitrarily.

First, we introduce some notions.

We also introduce the Pucci's class as follows.

Definition 7.1. We say that $u \in \bar{S}_{\rho}(\lambda, \Lambda, b_0, f)$ if for any $\varphi \in C^2(\Omega)$ with (1.3),

 $\mathcal{M}^{-}(D^{2}\varphi(x_{0}),\lambda,\Lambda) - b_{0}|D\varphi(x_{0})| \leq f(x_{0}).$

Similarly, we denote $u \in \underline{S}_{\rho}(\lambda, \Lambda, b_0, f)$ if for any $\varphi \in C^2(\Omega)$ with

$$\|\varphi\|_{C^{1,1}(\bar{\Omega})} \le \rho, \quad \varphi(x_0) = u(x_0), \quad \varphi \ge u \text{ in } \Omega,$$

we have

$$\mathcal{M}^+(D^2\varphi(x_0),\lambda,\Lambda) + b_0|D\varphi(x_0)| \ge f(x_0)$$

We also define

$$S^*_{\rho}(\lambda, \Lambda, b_0, f) = \underline{S}_{\rho}(\lambda, \Lambda, b_0, -|f|) \cap \overline{S}_{\rho}(\lambda, \Lambda, b_0, |f|).$$

We will denote $\underline{S}_{\rho}(\lambda, \Lambda, b_0, f)$ $(\overline{S}_{\rho}(\lambda, \Lambda, b_0, f), S^*_{\rho}(\lambda, \Lambda, f))$ by $\underline{S}_{\rho}(f)$ $(\overline{S}_{\rho}(f), S^*_{\rho}(f))$ for short if λ, Λ, b_0 are understood well.

Remark 7.2. Note that if $\rho_1 \geq \rho$,

$$\bar{S}_{\rho_1}(\lambda, \Lambda, b_0, f) \subset \bar{S}_{\rho}(\lambda, \Lambda, b_0, f).$$

Hence, for any $\rho > 0$,

$$\bar{S}(\lambda, \Lambda, b_0, f) \subset \bar{S}_{\rho}(\lambda, \Lambda, b_0, f),$$

where \bar{S} denotes the usual Pucci's class.

Remark 7.3. If $u \in \underline{S}_{\rho}(\lambda, \Lambda, b_0, f)$, then $-u \in \overline{S}_{\rho}(\lambda, \Lambda, b_0, f)$. Hence, we only consider the supersolution in the following argument.

As usual, any viscosity solution belongs to the Pucci's class.

Proposition 7.4. Let u be a viscosity supersolution of

$$F(D^2u, Du, u, x) = f \quad in \ \Omega,$$

where F is 5 ρ -uniformly elliptic and $||u||_{L^{\infty}(\Omega)} \leq \rho$. Let $\phi \in C^{2}(\Omega)$ with (1.3). Then (i) if (1.9) holds,

$$u - \phi \in \bar{S}_{4\rho}(\lambda, \Lambda, b_0, f(x) + c_0 | u(x) - \phi(x) | - F(D^2 \phi(x), D\phi(x), \phi(x), x));$$

(ii) if for any $|M|, |p|, |q|, |s| < \rho$ and $x \in B_1$,

(7.1)
$$-b_0|p-q| \le F(M, p, s, x) - F(M, q, s, x) \le b_0|p-q|$$

then

$$u - \phi \in \bar{S}_{4\rho}(\lambda, \Lambda, b_0, f(x) - F(D^2\phi(x), D\phi(x), u(x), x));$$

(iii) if $F(0, p, s, x) \equiv 0$,

$$u \in \bar{S}_{4\rho}(\lambda, \Lambda, 0, f - \mathcal{M}^{-}(D^{2}\phi))$$

Proof. Clearly, $||u - \phi||_{L^{\infty}(\Omega)} \leq 2\rho$. Given $x_0 \in \Omega$, for any $\varphi \in C^2(\Omega)$ with (1.3) (replacing u by $u - \phi$ and ρ by 4ρ there), $\varphi + \phi$ satisfies (1.3) (replacing ρ by 5ρ). By the definition of viscosity solution

$$f(x_0) \ge F(D^2\varphi(x_0) + D^2\phi(x_0), D\varphi(x_0) + D\phi(x_0), \varphi(x_0) + \phi(x_0), x_0).$$

If (1.9) holds,

 $f(x_0) \ge \mathcal{M}^-(D^2\varphi(x_0)) - b_0|D\varphi(x_0)| - c_0|\varphi(x_0)| + F(D^2\phi(x_0), D\phi(x_0), \phi(x_0), x_0),$ which means

$$u - \phi \in \overline{S}_{4\rho}(\lambda, \Lambda, b_0, f(x) + c_0 | u(x) - \phi(x) | - F(D^2 \phi(x), D\phi(x), \phi(x), x)).$$

If (7.1) holds,

$$f(x_0) \ge \mathcal{M}^-(D^2\varphi(x_0)) - b_0 |D\varphi(x_0)| + F(D^2\phi(x_0), D\phi(x_0), \varphi(x_0) + \phi(x_0), x_0).$$

By combining with $u(x_0) = \varphi(x_0) + \phi(x_0),$

$$u - \phi \in \overline{S}_{4\rho}(\lambda, \Lambda, b_0, f(x) - F(D^2\phi(x), D\phi(x), u(x), x))$$

If $F(0, p, s, x) \equiv 0$,

$$f(x_0) \ge \mathcal{M}^-(D^2\varphi(x_0)) + F(D^2\phi(x_0), D\varphi(x_0) + D\phi(x_0), \varphi(x_0) + \phi(x_0), x_0) \\\ge \mathcal{M}^-(D^2\varphi(x_0)) + \mathcal{M}^-(D^2\phi(x_0)).$$

Hence,

$$u \in \bar{S}_{4\rho}(\lambda, \Lambda, 0, f - \mathcal{M}^{-}(D^{2}\phi)).$$

As usual, we have the following maximum principle.

Lemma 7.5. If $u \in C(\overline{\Omega})$ satisfies

$$u \in \bar{S}_{\rho}(\lambda, \Lambda, b_0, 0)$$
 in Ω

and $u \ge 0$ on $\partial \Omega$, then

$$u \geq 0$$
 in Ω .

Proof. Suppose not. Then by choosing α large enough and ε small enough, $\varphi := \varepsilon e^{-\alpha x_1} + c$ will touch u by below at some $x_0 \in \Omega$ and

$$\mathcal{M}^{-}(D^{2}\varphi(x_{0})) - b_{0}|D\varphi(x_{0})| > 0,$$

which is a contradiction.

We also have

Lemma 7.6. Suppose that $\rho \ge \rho_0$,

$$u \in S_{\rho}(\lambda, \Lambda, b_0, f)$$
 in B_1

and

$$u \ge 0 \text{ on } \partial B_1, \quad \|f\|_{L^{\infty}(B_1)} \le 1,$$

where $\rho_0 > 1$ is universal. Then

$$\sup_{B_1} u^- \le C \|f\|_{L^{\infty}(B_1)},$$

where C is universal.

Proof. Take $\alpha > 0$ (universal) large enough such that $v := \|f\|_{L^{\infty}(B_1)} e^{\alpha x_1}$ is a classical solution of

(7.2)
$$\mathcal{M}^{-}(D^{2}v) - b_{0}|Dv| \ge 2||f||_{L^{\infty}(B_{1})}$$

If

$$\sup_{B_1} u^- \ge 2 \|f\|_{L^{\infty}(B_1)} e^{\alpha},$$

v - u has a local maximum at some $x_0 \in B_1$. Then by the definition of viscosity solution,

$$\mathcal{M}^{-}(D^{2}v(x_{0})) - b_{0}|Dv(x_{0})| \le f(x_{0}) \le ||f||_{L^{\infty}(B_{1})},$$

which contradicts to (7.2).

Next, we prove the fundamental Alexandrov-Bakel'man-Pucci maximum principle by the same way as in [27, Chapter 3]. First, we prove a lemma analogous to [27, Lemma 3.3]:

Lemma 7.7. Let $0 < \delta < 1$, $\rho \ge \rho_1$ and

$$\iota \in \overline{S}_{\rho}(\lambda, \Lambda, b_0, f)$$
 in B_{δ}

where $\rho_1 > 1$ is universal. Suppose that

$$\|f\|_{L^{\infty}(B_{\delta})} \le 1.$$

Assume that φ is a convex function in B_{δ} such that $0 \leq \varphi \leq u$ in B_{δ} and $0 = \varphi(0) = u(0)$. Then

$$\varphi(x) \le C(\sup_{B_{\delta}} f^+)|x|^2, \ \forall \ x \in B_{\nu\delta},$$

where $0 < \nu < 1$ and C are universal.

Proof. We prove the lemma in the same way as that of Lemma 3.3 in [27, Chapter 3]. The constant ρ_1 is to be specified later. As in the proof Lemma 3.3 of [27], for $0 < r < \delta/4$, define

$$\bar{C} = \frac{1}{r^2} \sup_{B_r} \varphi.$$

We aim to prove

$$\bar{C} \le \frac{17}{\lambda} \sup_{B_{\delta}} f^+.$$

Suppose not. Let

(7.3)
$$\tilde{C} = \frac{17}{\lambda} \sup_{B_{\delta}} f^+ \le \frac{17}{\lambda}$$

Recall (3.3) in [27], i.e.,

$$\varphi \ge \bar{C}r^2$$
 in $H \cap B_{\delta}$

where H is a hyperplane tangent to B_r at some $x_0 \in \partial B_r$. Since $\tilde{C} < \bar{C}$,

$$\varphi \ge \tilde{C}r^2$$
 in $H \cap B_{\delta}$.

Construct the polynomial P as in the proof of [27, Lemma 3.3], i.e.,

$$P(x) := \frac{\tilde{C}}{8}(x_n + r)^2 - 4\tilde{C}\frac{r^2}{\delta^2}|x'|^2.$$

Then P + c for an appropriate constant c will touch u by below at some point x_0 . Note that

$$||D^2P||_{L^{\infty}} \le \tilde{C}, \quad ||DP||_{L^{\infty}} \le \tilde{C}, \quad ||P+c||_{L^{\infty}} \le \tilde{C} + \rho/2.$$

Choose $\rho_1 \geq 2\tilde{C}$. Then by the definition of viscosity solution,

$$\mathcal{M}^{-}(D^2P) - b_0|DP(x_0)| \le f(x_0) \le \sup_{B_{\delta}} f^+.$$

By choosing

$$\nu = \min\left(\frac{1}{8}\sqrt{\frac{\lambda}{(n-1)\Lambda}}, \frac{\lambda}{12b_0}\right),$$

we have

$$\frac{\lambda \tilde{C}}{16} \le \mathcal{M}^-(D^2 P) - b_0 |DP(x_0)| \le \sup_{B_\delta} f^+.$$

Hence,

$$\tilde{C} \le \frac{16}{\lambda} \sup_{B_{\delta}} f^+,$$

which contradicts to (7.3).

We have another lemma.

Lemma 7.8. Let $\rho \geq \rho_2$,

$$u \in \bar{S}_{\rho}(\lambda, \Lambda, b_0, f)$$
 in B_1

and

$$u \ge 0 \text{ on } \partial B_1, \quad \|u\|_{L^{\infty}(B_1)} \le \frac{\rho}{4}, \quad \|f\|_{L^{\infty}(B_1)} \le 1,$$

where $\rho_2 > \max(\rho_0, \rho_1)$ is universal. Then $\Gamma_u \in C^{1,1}(\bar{B}_1)$ and

 $\mathcal{M}^{-}(D^{2}\Gamma_{u}) - b_{0}|D\Gamma_{u}| \leq f \ a.e. \ in \ \{u = \Gamma_{u}\},\$

where we have extended u by zero outside B_1 and Γ_u is the convex envelop of $-u^-$ in B_2 .

Proof. For any $x_0 \in \{u = \Gamma_u\} \subset B_1$, let L_{x_0} be the supporting affine function of Γ_u at x_0 . Then by Lemma 7.6 and noting $x_0 \in B_1$,

$$\|\Gamma_u\|_{L^{\infty}(B_1)} = \sup_{B_1} u^- \le C_0, \quad |DL_{x_0}| \le C_0, \quad \|L_{x_0}\|_{L^{\infty}(B_1)} \le C_0,$$

where C_0 is universal. Set $v = u - L_{x_0}$. Then $v(x_0) = (\Gamma_u - L_{x_0})(x_0) = 0$ and by choosing ρ_2 large enough,

$$v \in \bar{S}_{3\rho/4}(\lambda, \Lambda, b_0, f + b_0 C_0).$$

Let $w = v/(1 + b_0 C_0)$. Then

$$v \in \bar{S}_{3\rho/(4+4b_0C_0)}(\lambda, \Lambda, b_0, 1).$$

By taking ρ_2 large enough, we can apply Lemma 7.7 for any $0 < \delta < 1$. Then

$$L_{x_0} \le \Gamma_u \le L_{x_0} + C|x - x_0|^2,$$

where C is universal. Therefore, by the same argument in [27] (see the proof of Lemma 3.5), $\Gamma_u \in C^{1,1}(\bar{B}_1)$. Hence, Γ_u is second order differentiable almost everywhere.

Next, take $x_0 \in \{u = \Gamma_u\}$ such that Γ_u is second order differentiable at x_0 . Let P denote the second order polynomial corresponding to Γ_u at x_0 . Then for any $\varepsilon > 0$, $P - \varepsilon |x - x_0|^2$ will touch locally Γ_u and hence u by below at x_0 . Note that $||P|| \leq C$ for some universal C. Then by choosing ρ_2 large enough, $P - \varepsilon |x - x_0|^2$ is an admissible test function. Hence,

$$\mathcal{M}^{-}(D^2P - 2\varepsilon I) - b_0 |DP(x_0) - 2\varepsilon(x - x_0)| \le f(x_0).$$

41

By letting $\varepsilon \to 0$, we arrive at the conclusion.

Based on above lemma, we have the following Alexandrov-Bakel'man-Pucci maximum principle analogous to [27, Theorem 3.2]:

Theorem 7.9 (ABP). Let $\rho \geq \rho_2$,

$$u \in \bar{S}_{\rho}(\lambda, \Lambda, b_0, f)$$
 in B_1

and

$$u \ge 0 \text{ on } \partial B_1, \quad \|u\|_{L^{\infty}(B_1)} \le \frac{\rho}{4}, \quad \|f\|_{L^{\infty}(B_1)} \le 1,$$

where ρ_2 is as in Lemma 7.8. Then

$$\sup_{B_1} u^- \le C \|f^+\|_{L^n(B_1 \cap \{u = \Gamma_u\})},$$

where C is universal.

Proof. The proof is standard and we omit it (see [56, Chapt. 9.1]). \Box

By scaling, we have

Corollary 7.10. Let $\rho \ge \rho_2$, K > 0,

$$u \in \bar{S}_{K\rho}(\lambda, \Lambda, b_0, f)$$
 in B_1

and

$$u \ge 0 \text{ on } \partial B_1, \quad \|u\|_{L^{\infty}(B_1)} \le \frac{K\rho}{4}, \quad \|f\|_{L^{\infty}(B_1)} \le K,$$

where ρ_2 is as in Lemma 7.8. Then

$$\sup_{B_1} u^- \le C \|f^+\|_{L^n(B_1 \cap \{u = \Gamma_u\})}$$

where C is universal.

Proof. Let

$$\tilde{u} = \frac{u}{K}, \quad \tilde{f} = \frac{f}{K}.$$

Then

$$\tilde{u} \in \bar{S}_{\rho}(\lambda, \Lambda, b_0, \tilde{f}) \text{ in } B_1, \quad \|\tilde{u}\|_{L^{\infty}(B_1)} \le \frac{\rho}{4}, \quad \|\tilde{f}\|_{L^{\infty}(B_1)} \le 1$$

By Theorem 7.9,

$$\sup_{B_1} \tilde{u}^- \le C \|\tilde{f}^+\|_{L^n(B_1 \cap \{\tilde{u} = \Gamma_{\tilde{u}}\})}$$

By transforming to u, we obtain the conclusion.

With the aid of the ABP maximum principle, we can prove the following lemma analogous to [27, Lemma 4.5]:

Lemma 7.11. Let $\rho \geq \rho_3$,

$$u \in \bar{S}_{\rho}(\lambda, \Lambda, b_0, |f|)$$
 in $Q_{4\sqrt{n}}$

and

$$u \ge 0 \quad in \ Q_{4\sqrt{n}}, \quad \inf_{Q_3} u \le 1, \quad \|u\|_{L^{\infty}(Q_{4\sqrt{n}})} \le \frac{\rho}{8}, \quad \|f\|_{L^{\infty}(Q_{4\sqrt{n}})} \le \varepsilon_0,$$

where $\rho_3 \ge \rho_2$ is universal. Then

(7.4)
$$|\{u < M\} \cap Q_1| > \mu,$$

where $0 < \varepsilon_0, \mu < 1$ and M > 1 are universal.

Proof. The proof is the same as that of [27, Lemma 4.5]. Construct the auxiliary function φ as in [27, Lemma 4.1] such that the conclusion of [27, Lemma 4.1] holds with

$$\mathcal{M}^+(D^2\varphi,\lambda,\Lambda) \le C\xi$$

replaced by

$$\mathcal{M}^+(D^2\varphi,\lambda,\Lambda) + b_0|D\varphi| \le C\xi.$$

Since

$$\|D^2\varphi\|_{L^{\infty}} \le C_0, \quad \|D\varphi\|_{L^{\infty}} \le C_0, \quad \|D^2\varphi\|_{L^{\infty}} \le C_0,$$

where C_0 is universal. By taking ρ_3 large enough,

$$w := u + \varphi \in \bar{S}_{3\rho/4}(\lambda, \Lambda, b_0, |f| + C\xi), \quad ||w||_{L^{\infty}} \le \frac{3\rho}{16},$$

Now, we can apply the ABP maximum principle Corollary 7.10 to obtain (7.4) as in [27].

By iteration, we have the following lemma analogous to [27, Lemma 4.6]: Lemma 7.12. Let $\rho \ge \rho_3$,

$$u \in \bar{S}_{\rho}(\lambda, \Lambda, b_0, f)$$
 in $Q_{4\sqrt{n}}$

and

$$u \ge 0 \quad in \ Q_{4\sqrt{n}}, \quad \inf_{Q_3} u \le 1, \quad \|u\|_{L^{\infty}(Q_{4\sqrt{n}})} \le \frac{\rho}{8}, \quad \|f\|_{L^{\infty}(Q_{4\sqrt{n}})} \le \varepsilon_0.$$

Then

(7.5)
$$|\{u \ge t\} \cap Q_1| \le Ct^{-\varepsilon}, \ \forall \ t < \frac{\rho}{\rho_3},$$

where C and $0 < \varepsilon < 1$ are universal.

Proof. The proof is the same as in [27, Lemma 4.6]. We only need to prove

(7.6)
$$|\{u > M^k\} \cap Q_1| \le (1-\mu)^k, \ \forall \ 1 \le k \le \frac{1}{\ln M} \ln \frac{\rho}{\rho_3}$$

since (7.5) follows from (7.6) by choosing $\varepsilon > 0$ with

$$1 - \mu = M^{-\varepsilon}.$$

For k = 1, we have just proved in Lemma 7.11. Suppose that (7.6) holds for k - 1. We use the same scaling argument as in [27]:

$$\tilde{u}(y) := \frac{u(x)}{M^{k-1}}.$$

Since $u \in \bar{S}_{\rho}(\lambda, \Lambda, b_0, f)$,

$$\tilde{u} \in \bar{S}_{\rho/M^{k-1}}(\lambda, \Lambda, b_0, \tilde{f}), \quad \tilde{f}(y) = \frac{f(x)}{2^{2i}M^{k-1}}.$$

Note that

$$k \le \frac{1}{\ln M} \ln \frac{\rho}{\rho_3}$$

implies

$$\frac{\rho}{M^{k-1}} \ge \rho_3.$$

Therefore, by Lemma 7.11,

$$|\{\tilde{u} > M\} \cap Q_1| \le 1 - \mu.$$

By transforming to u, we obtain (7.6).

Next, we prove the "weak Harnack inequality" analogous to [27, Theorem 4.8]:

Theorem 7.13. Let $\rho \geq \rho_3$,

$$u \in \bar{S}_{\rho}(\lambda, \Lambda, b_0, f)$$
 in B_1

and

$$0 \le u \le 1$$
 in B_1 , $||u||_{L^{\infty}(B_1)} \le \frac{\rho}{8}$, $||f||_{L^{\infty}(B_1)} \le \varepsilon_0$.

Then

$$||u_{\rho}||_{L^{p_0}(B_{1/2})} \le C\left(\inf_{B_{1/2}} u + ||f||_{L^{\infty}(B_1)}\right),$$

where C and $0 < p_0 < 1$ are universal, and

$$u_{\rho}(x) = \begin{cases} u(x), & \text{if } u(x) \leq \frac{\rho}{\rho_3}; \\ 0, & \text{if } u(x) > \frac{\rho}{\rho_3}. \end{cases}$$

Proof. If

$$\inf_{B_{1/2}} u + \varepsilon_0^{-1} \|f\|_{L^{\infty}(B_1)} \ge 1,$$

44

by Lemma 7.12 and taking $p_0 = \varepsilon/2$,

$$\begin{aligned} \|u_{\rho}\|_{L^{p_{0}}(B_{1/2})} &= p_{0} \int_{0}^{+\infty} t^{p_{0}-1} |\{x \in B_{1/2} : u_{\rho}(x) > t\} | dt \\ &\leq |B_{1/2}| + p_{0} \int_{1}^{\rho/\rho_{3}} t^{p_{0}-1} |\{x \in B_{1/2} : u(x) > t\} | dt \\ &\leq C + C \int_{1}^{\rho/\rho_{3}} t^{p_{0}-1} t^{-\varepsilon} dt \\ &\leq C \\ &\leq C \left(\inf_{B_{1/2}} u + \|f\|_{L^{\infty}(B_{1})}\right). \end{aligned}$$

 \mathbf{If}

$$K^{-1} := \inf_{B_{1/2}} u + \varepsilon_0^{-1} \|f\|_{L^{\infty}(B_1)} < 1,$$

 $\operatorname{consider}$

 $\tilde{u} = Ku.$

Then

$$\tilde{u} \in \bar{S}_{K\rho}(\lambda, \Lambda, b_0, \tilde{f})$$

and

$$\inf_{B_{1/2}} \tilde{u} \le 1, \quad \|\tilde{u}\|_{L^{\infty}(B_1)} \le \frac{\rho}{8}, \quad \|\tilde{f}\|_{L^{\infty}(B_1)} \le \varepsilon_0.$$

Then by Lemma 7.12,

$$\|\tilde{u}_{K\rho}\|_{L^{p_0}(B_{1/2})} \le C.$$

By transferring to u, we obtain the conclusion.

Next, we prove the Hölder regularity.

Theorem 7.14. Let $\rho \geq 2\rho_3$,

$$u \in S^*_{\rho}(\lambda, \Lambda, b_0, f)$$
 in B_1

and

$$||u||_{L^{\infty}(B_1)} \le 1, \quad ||f||_{L^{\infty}(B_1)} \le \varepsilon_1,$$

where $0 < \varepsilon_1 < 1$ is universal. Then

$$\underset{B_r}{\text{osc}} \ u \le Cr^{\alpha}, \ \forall \ \sqrt{\frac{2\rho_3}{\rho}} \le r \le 1,$$

where $0 < \alpha < 1$ and C are universal.

Proof. We only need to prove

(7.7)
$$\underset{B_{1/2^k}}{\operatorname{osc}} u \le 2(1-\mu)^k, \ \forall \ 0 \le k \le \frac{1}{2} \log_2 \frac{\rho}{2\rho_3},$$

45

where $0 < \mu < 1$ is universal. We prove it by induction. For k = 0, (7.7) holds clearly. Suppose that it holds for k. For r > 0, denote

$$M_r = \sup_{B_r} u, \quad m_r = \inf_{B_r} u.$$

Set $r_0 = 1/2^k$. Note that

$$|\{x \in B_{r_0} : u(x) \ge \frac{M_{r_0} + m_{r_0}}{2}\}| \ge \frac{|B_{r_0}|}{2} \text{ or } |\{x \in B_{r_0} : u(x) \le \frac{M_{r_0} + m_{r_0}}{2}\}| \ge \frac{|B_{r_0}|}{2}$$

Without loss of generality, we assume that the former holds. Let

$$y = \frac{x}{r_0}, \quad \tilde{u}(y) = \frac{u(x) - m_{r_0}}{M_{r_0} - m_{r_0}}, \quad \tilde{f}(y) = \frac{r_0^2 f(x)}{M_{r_0} - m_{r_0}}.$$

Without loss of generality, we assume that

$$\underset{B_{r_0}}{\text{osc}} u = M_{r_0} - m_{r_0} \ge r_0^2.$$

Note that $M_{r_0} - m_{r_0} \leq 2$. Then

$$\tilde{u} \in S^*_{r_0^2 \rho/2}(\lambda, \Lambda, b_0, \tilde{f})$$
 in B_1

and

$$0 \le \tilde{u} \le 1$$
 in B_1 , $\|\tilde{f}\|_{L^{\infty}(B_1)} \le \varepsilon_1$.

Moreover,

$$|\{y \in B_1 : \tilde{u}(y) \ge \frac{1}{2}\}| \ge \frac{|B_1|}{2}.$$

Hence, by combining with Theorem 7.13 and noting

$$\frac{r_0^2 \rho}{2} \ge 8, \quad \frac{r_0^2 \rho}{2\rho_3} \ge 1,$$

we have

$$c_0 \le \|\tilde{u}\|_{L^{p_0}(B_{1/2})} = \|\tilde{u}_{r_0^2 \rho/2}\|_{L^{p_0}(B_{1/2})} \le C\left(\inf_{B_{1/2}} \tilde{u} + \|\tilde{f}\|_{L^{\infty}(B_1)}\right) \le C\inf_{B_{1/2}} \tilde{u} + C\varepsilon_1,$$

where $c_0 > 0, C$ are universal constants.

Choose ε_1 small enough such that

$$C\varepsilon_1 \leq \frac{c_0}{2}.$$

Then for some universal constant $\mu > 0$,

$$\inf_{B_{1/2}} \tilde{u} \ge \mu.$$

By transforming back to u,

$$m_{r_0/2} - m_{r_0} \ge \mu \left(M_{r_0} - m_{r_0} \right)$$

Hence,

$$\underset{B_{1/2^{k+1}}}{\operatorname{osc}} u = M_{r_0/2} - m_{r_0/2} \le M_{r_0} - m_{r_0} - \mu \left(M_{r_0} - m_{r_0} \right) = (1 - \mu) \left(M_{r_0} - m_{r_0} \right) \le 2(1 - \mu)^{k+1}.$$

By induction, the proof is completed.

Remark 7.15. Since ρ is finite, we can make the scaling argument only finite times. Hence, we can't obtain the real Hölder regularity. However, it can provide necessary compactness when we use the compactness method to prove higher regularity (see Lemma 8.1 and Lemma 9.1).

By applying above theorem to each $x_0 \in B_{1/2}$ in $B(x_0, 1/2)$, we have

Corollary 7.16. Let $\rho \geq 2\rho_3$,

$$u \in S^*_{\rho}(\lambda, \Lambda, b_0, f)$$
 in B_1

and

$$||u||_{L^{\infty}(B_1)} \le 1, \quad ||f||_{L^{\infty}(B_1)} \le \varepsilon_1,$$

where $0 < \varepsilon_1 < 1$ is universal. Then

$$\underset{B_r(x_0)}{\operatorname{osc}} u \leq \tilde{C}r^{\alpha_0}, \ \forall \ x_0 \in B_{1/2}, \ \forall \ \sqrt{\frac{2\rho_3}{\rho}} \leq r \leq \frac{1}{2},$$

where $0 < \alpha_0 < 1$ and \tilde{C} are universal.

Therefore,

$$|u(x_1) - u(x_2)| \le \tilde{C}|x_1 - x_2|^{\alpha_0}, \ \forall \ x_1, x_2 \in B_{1/2}, \ \sqrt{\frac{2\rho_3}{\rho}} \le |x_1 - x_2| \le \frac{1}{2}.$$

8. Interior $C^{1,\alpha}$ regularity

In this section, we prove the interior pointwise $C^{1,\alpha}$ regularity by the classical technique of perturbation. First, we prove the key step by the compactness method.

Lemma 8.1. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of

$$F(D^{2}u, Du, u, x) = A^{ij}(Du, u, x)u_{ij} + B(Du, u, x) = 0 \text{ in } B_{1},$$

where F is ρ -uniformly elliptic and A is continuous with modulus ω_A . Suppose that (1.7) and (1.8) hold. Let $r \leq r_0$ and assume that for some $P_0 \in \mathcal{P}_1$,

$$||u - P_0||_{L^{\infty}(B_r)} \le r^{1+\alpha}, \quad ||P_0|| \le \bar{C}r_0^{1+\alpha},$$

where \overline{C} depends only on $n, \lambda, \Lambda, \rho, b_0$ and α and $0 < r_0, \delta_0 < 1$ (small) depends also on ω_A .

Then there exists $P \in \mathcal{P}_1$ such that

$$||u - P||_{L^{\infty}(B_{\eta r})} \le (\eta r)^{1+\alpha}, \quad ||P - P_0||_r \le \bar{C}(\eta r)^{1+\alpha},$$

where $0 < \eta < 1/2$ depends only on $n, \lambda, \Lambda, \rho, b_0, c_0$ and α .

Proof. We prove the lemma by contradiction. Suppose that the conclusion is false. Then there exist sequences of A_m, B_m, u_m, P_m, r_m such that $r_m \leq 1/m$,

$$F_m(D^2u_m, Du_m, u_m, x) = A_m^{ij}(Du_m, u_m, x)u_{m,ij} + B_m(Du_m, u_m, x) = 0 \text{ in } B_{r_m}$$

and

$$|u_m - P_m||_{L^{\infty}(B_{r_m})} \le r_m^{1+\alpha}, \quad ||P_m|| \le \frac{\bar{C}}{m},$$

where F_m are ρ -uniformly elliptic and A_m are continuous (with the same modulus ω_A). In addition, (1.7) and (1.8) hold for B_m (with K_B and b_0). Moreover, for any $P \in \mathcal{P}_1$ with

$$\|P - P_m\| \le \bar{C}(\eta r_m)^{1+\alpha},$$

we have

(8.1)
$$||u_m - P||_{L^{\infty}(B_{\eta r_m})} > (\eta r_m)^{1+\alpha},$$

where \bar{C} and $0 < \eta < 1/2$ are to be specified later.

Let

$$\tilde{x} = \frac{x}{r_m}, \quad \tilde{u}_m(\tilde{x}) = \frac{u_m(x) - P_m(x)}{r_m^{1+\alpha}}.$$

Then \tilde{u}_m are viscosity solutions of

(8.2)
$$\tilde{F}_m(D^2\tilde{u}_m, D\tilde{u}_m, \tilde{u}_m, \tilde{x}) = 0 \text{ in } B_1,$$

where

$$\begin{split} \tilde{F}_m(M, p, s, \tilde{x}) = & r_m^{1-\alpha} F_m(r_m^{\alpha-1}M, r_m^{\alpha}p + DP_m(x), r_m^{1+\alpha}s + P_m(x), x), \\ = & A_m^{ij}(r_m^{\alpha}p + DP_m(x), r_m^{1+\alpha}s + P_m(x), x) M_{ij} \\ &+ r_m^{1-\alpha} B_m(r_m^{\alpha}p + DP_m(x), r_m^{1+\alpha}s + P_m(x), x). \end{split}$$

Indeed, since F_m are ρ -uniformly elliptic and $||P_m|| \to 0$, \tilde{F}_m are $\frac{1}{2}r_m^{-\alpha}\rho$ uniformly elliptic for m large enough. Note that

$$\|\tilde{u}_m\|_{L^{\infty}(B_1)} \le 1 \le \frac{1}{4} r_m^{-\alpha} \rho.$$

Then it can be verified directly that \tilde{u}_m are viscosity solutions of (8.2).

Next, by Proposition 7.4, for m large enough,

$$\tilde{u}_m \in S^*_{\frac{1}{2}r_m^{-\alpha}\rho}(\lambda, \Lambda, r_m b_0, \bar{f}_m),$$

where

$$\bar{f}_m(\tilde{x}) = r_m^{1-\alpha} B_m(DP_m(x), r_m^{1+\alpha} \tilde{u}(\tilde{x}) + P_m(x), x)$$

Hence, for any $\varepsilon > 0$, we can take m_0 large enough such that for any $m \ge m_0$,

$$\|\bar{f}_m\|_{L^{\infty}(B_1)} \le \varepsilon_1, \quad \tilde{C}\left(\frac{4\rho_3}{r_m^{-\alpha}\rho}\right)^{\alpha_0/2} \le \frac{\varepsilon}{2},$$

where α_0, C, ρ_3 and ε_1 are as in Corollary 7.16. Set

$$\delta = \left(\frac{4\rho_3}{r_m^{-\alpha}\rho}\right)^{1/2}$$

By Corollary 7.16, for any $x_1, x_2 \in B_{1/2}$ with $|x_1 - x_2| \leq \delta$, by choosing $x_3 \in B_{1/2}$ with $|x_1 - x_3| = |x_2 - x_3| = \delta$, we have for any $m \geq m_0$,

$$|\tilde{u}_m(x_1) - \tilde{u}_m(x_2)| \le |\tilde{u}_m(x_1) - \tilde{u}_m(x_3)| + |\tilde{u}_m(x_2) - \tilde{u}_m(x_3)| \le 2\tilde{C}\delta^{\alpha_0} \le \varepsilon.$$

Thus, \tilde{u}_m are equicontinuous. By Arzelà-Ascoli theorem, there exists $\tilde{u} \in C(\bar{B}_{1/2})$ such that $\tilde{u}_m \to \tilde{u}$ in $L^{\infty}(B_{1/2})$ (up to a subsequence and similarly hereinafter).

Since

$$\lambda I \le A_m(0,0,0) \le \Lambda I, \ \forall \ m \ge 1,$$

there exists a constant symmetric matrix $A = (A^{ij})_{n \times n}$ such that

 $A_m(0,0,0) \to A.$

Now, we show that \tilde{u} is a viscosity solution of

(8.3)
$$A^{ij}\tilde{u}_{ij} = 0 \text{ in } B_{1/2}$$

Given $\tilde{x}_0 \in B_{1/2}$ and $\varphi \in C^2$ touching \tilde{u} strictly by above at \tilde{x}_0 . Then there exist a sequence of $\tilde{x}_m \to \tilde{x}_0$ such that $\varphi + c_m$ touch \tilde{u}_m by above at \tilde{x}_m and $c_m \to 0$. By the definition of viscosity solution, for m large enough (e.g. $r_m^{-\alpha} \rho > 2 \|\varphi\|_{C^2(\bar{B}_{1/2})}$),

$$\tilde{F}_m(D^2\varphi(\tilde{x}_m), D\varphi(\tilde{x}_m), \varphi(\tilde{x}_m) + c_m, \tilde{x}_m) \ge 0.$$

Since

$$\begin{split} F_m(D^2\varphi(\tilde{x}_m), D\varphi(\tilde{x}_m), \varphi(\tilde{x}_m) + c_m, \tilde{x}_m) \\ = & A_m^{ij}(r_m^{\alpha}D\varphi(\tilde{x}_m) + DP_m(x), r_m^{1+\alpha}\varphi(\tilde{x}_m) + P_m(x), x)M_{ij} \\ &+ r_m^{1-\alpha}B_m(r_m^{\alpha}D\varphi(\tilde{x}_m) + DP_m(x), r_m^{1+\alpha}\varphi(\tilde{x}_m) + P_m(x), x), \\ \leq & A_m^{ij}(0, 0, 0)M_{ij} + \omega_A(\frac{C}{m}) \\ &+ r_m^{1-\alpha}B_m(r_m^{\alpha}D\varphi(\tilde{x}_m) + DP_m(x), r_m^{1+\alpha}\varphi(\tilde{x}_m) + P_m(x), x), \end{split}$$

by letting $m \to \infty$, we have

$$A^{ij}\varphi_{ij}(\tilde{x}_0) \ge 0.$$

Hence, \tilde{u} is a subsolution of (8.3). Similarly, we can prove that it is a viscosity supersolution as well. That is, \tilde{u} is a viscosity solution.

Since (8.3) is a linear equation with constant coefficients, $\tilde{u} \in C^{\infty}(B_{1/2})$. Then there exists $\tilde{P} \in \mathcal{P}_1$ such that for any $0 < \eta < 1/4$,

$$\|\tilde{u} - \tilde{P}\|_{L^{\infty}(B_{\eta})} \le C_1 \eta^2 \|\tilde{u}\|_{L^{\infty}(B_{1/2})} \le C_1 \eta^2,$$

and

$$||P|| \le C_2 ||\tilde{u}||_{L^{\infty}(B_{1/2})} \le C_2,$$

where C_1 and C_2 are universal. By taking η small and \overline{C} large such that

$$C\eta^{1-\alpha} \le \frac{1}{2}, \quad C_2 \le \bar{C}\eta^{1+\alpha}.$$

Then

(8.4)
$$\|\tilde{u} - \tilde{P}\|_{L^{\infty}(B_{\eta})} \le \frac{1}{2}\eta^{1+\alpha}, \quad \|\tilde{P}\| \le \bar{C}\eta^{1+\alpha}.$$

Let

$$Q_m(x) = P_m(x) + r_m^{1+\alpha} \tilde{P}(\tilde{x}).$$

Then

$$\|Q_m - P_m\| \le \bar{C}(\eta r_m)^{1+\alpha}.$$

Hence, (8.1) holds for Q_m . That is,

$$||u_m - Q_m||_{L^{\infty}(B_\eta r_m)} > (\eta r_m)^{1+\alpha}.$$

Equivalently,

$$\|\tilde{u}_m - \tilde{P}\|_{L^{\infty}(B_{\eta})} > \eta^{1+\alpha}.$$

Let $m \to \infty$, we have

$$\|\tilde{u} - \tilde{P}\|_{L^{\infty}(B_{\eta})} \ge \eta^{1+\alpha},$$

which contradicts with (8.4).

Now, we give the

Proof of Theorem 1.5. To prove Theorem 1.5, we only to prove the following. There exist a sequence of $P_m \in \mathcal{P}_1 \ (m \ge -1)$ such that for all $m \ge 0$,

(8.5)
$$\|u - P_m\|_{L^{\infty}(B_{\eta^m r_0})} \le (\eta^m r_0)^{1+\alpha},$$

and

(8.6)
$$||P_m - P_{m-1}||_{\eta^m r_0} \le \bar{C}(\eta^m r_0)^{1+\alpha},$$

where η, r_0 and \bar{C} are as in Lemma 8.1.

We prove above by induction. For m = 0, by setting $P_0 \equiv P_{-1} \equiv 0$, (8.5) and (8.6) hold clearly. Suppose that the conclusion holds for $m \leq m_0$. By (8.6),

$$\|P_{m_0}\| \le \sum_{i=1}^{m_0} \|P_i - P_{i-1}\| \le \bar{C}r_0^{1+\alpha} \frac{\eta^{1+\alpha}}{1-\eta^{1+\alpha}} \le \bar{C}r_0^{1+\alpha}.$$

By Lemma 8.1, the conclusion holds for $m = m_0 + 1$. By induction, the proof of Theorem 1.5 is completed.

9. Interior $C^{2,\alpha}$ regularity

In this section, we prove the interior pointwise $C^{2,\alpha}$ regularity. As before, we first prove the key step by the compactness method.

Lemma 9.1. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of

$$F(D^2u, Du, u, x) = f \quad in \ B_1,$$

where F is ρ -uniformly elliptic, $F(0,0,0,x) \equiv 0$ and $D_M F$ is continuous with modulus ω_F . Suppose that (1.9) holds. Let $r \leq r_0$ and assume that for some $P_0 \in \mathcal{P}_2$,

$$|F(M, p, s, x) - F(M, p, s, 0)||_{L^{\infty}(B_r)} \le \delta_0 r^{\alpha}, \ \forall \ |M|, |p|, |s| \le \rho,$$
$$||u - P_0||_{L^{\infty}(B_r)} \le r^{2+\alpha}, \quad ||f - f(0)||_{L^{\infty}(B_r)} \le \delta_0 r^{\alpha}$$

and

$$||P_0|| \le \bar{C}r_0^{2+\alpha}, \quad F(D^2P_0, DP_0(0), P_0(0), 0) = f(0),$$

where \overline{C} depends only on $n, \lambda, \Lambda, \rho, b_0, c_0$ and α and $0 < r_0, \delta_0 < 1$ (small) depends also on ω_F .

Then there exists $P \in \mathcal{P}_2$ such that

$$||u - P||_{L^{\infty}(B_{\eta r})} \le (\eta r)^{2+\alpha}$$

and

$$||P - P_0||_r \le \bar{C}(\eta r)^{2+\alpha}, \quad F(D^2 P, DP(0), P(0), 0) = f(0),$$

where $0 < \eta < 1/2$ depends only on $n, \lambda, \Lambda, \rho, b_0, c_0$ and α .

Proof. We prove the lemma by contradiction. Suppose that the conclusion is false. Then there exist sequences of F_m , u_m , f_m , P_m , r_m such that $r_m \leq 1/m$,

$$F_m(D^2 u_m, Du_m, u_m, x) = f_m \text{ in } B_{r_m},$$
$$\|F_m(M, p, s, x) - F_m(M, p, s, 0)\|_{L^{\infty}(B_{r_m})} \leq \frac{r_m^{\alpha}}{m}, \ \forall \ |M|, |p|, |s| \leq \rho$$
$$\|u_m - P_m\|_{L^{\infty}(B_{r_m})} \leq r_m^{2+\alpha}, \quad \|f_m - f_m(0)\|_{L^{\infty}(B_{r_m})} \leq \frac{r_m^{\alpha}}{m}$$

and

$$F_m(D^2 P_m, DP_m(0), P_m(0), 0) = f_m(0), \quad ||P_m|| \le \frac{C}{m}$$

where F_m are ρ -uniformly elliptic, $F_m(0, 0, 0, x) \equiv 0$ and $D_M F_m$ are continuous (with the same modulus ω_F). In addition, (1.9) holds for F_m with b_0 and c_0 . Moreover, for any $P \in \mathcal{P}_2$ with

$$||P - P_m|| \le \bar{C}(\eta r_m)^{2+\alpha}, \quad F_m(D^2P, DP(0), P(0), 0) = f_m(0),$$

we have

(9.1)
$$||u_m - P||_{L^{\infty}(B_{\eta r_m})} > (\eta r_m)^{2+\alpha},$$

where \bar{C} and $0 < \eta < 1/2$ are to be specified later.

Let

(9.2)
$$\tilde{x} = \frac{x}{r_m}, \quad \tilde{u}_m(\tilde{x}) = \frac{u_m(x) - P_m(x)}{r_m^{2+\alpha}}.$$

Then \tilde{u}_m are viscosity solutions of

(9.3)
$$\tilde{F}_m(D^2\tilde{u}_m, D\tilde{u}_m, \tilde{u}_m, \tilde{x}) = \tilde{f}_m \text{ in } B_1,$$

where

$$\tilde{F}_m(M, p, s, \tilde{x}) = r_m^{-\alpha} F_m(r_m^{\alpha} M + D^2 P_m, r_m^{1+\alpha} p + D P_m(x), r_m^{2+\alpha} s + P_m(x), x),$$

$$\tilde{f}_m(\tilde{x}) = r_m^{-\alpha} f_m(x).$$

Indeed, since F_m are ρ -uniformly elliptic and $||P_m|| \to 0$, \tilde{F}_m are $\frac{1}{2}r_m^{-\alpha}\rho$ uniformly elliptic for m large enough. Note that

$$\|\tilde{u}_m\|_{L^{\infty}(B_1)} \le 1 \le \frac{1}{4} r_m^{-\alpha} \rho.$$

Then it can be verified directly that \tilde{u}_m are viscosity solutions of (9.3).

Next, by Proposition 7.4, for m large enough,

$$\tilde{u}_m \in S^*_{\frac{1}{2}r_m^{-\alpha}\rho}(\lambda, \Lambda, r_m b_0, \bar{f}_m),$$

where

$$\bar{f}_m(\tilde{x}) = |r_m^{-\alpha} \left(f_m(x) - F_m(D^2 P_m, DP_m(x), P_m(x), x) \right)| + r_m^2 c_0 |u_m(x)|.$$

With the aid of $F_m(D^2 P_m, DP_m(0), P_m(0), 0) = f_m(0),$

$$\begin{aligned} |f_m - F_m(D^2 P_m, DP_m, P_m, x)| \\ &= \left| f_m - f_m(0) - \left(F_m(D^2 P_m, DP_m, P_m, x) - F_m(D^2 P_m, DP_m, P_m, 0) \right) \\ &- \left(F_m(D^2 P_m, DP_m, P_m, 0) - F_m(D^2 P_m, DP_m(0), P_m(0), 0) \right) \right| \\ &\leq \frac{r_m^{\alpha}}{m} + \frac{r_m^{\alpha}}{m} + b_0 C r_m + c_0 C r_m. \end{aligned}$$

Hence, for any $\varepsilon > 0$, we can take m_0 large enough such that for any $m \ge m_0$,

$$\|\bar{f}_m\|_{L^{\infty}(B_1)} \le \varepsilon_1, \quad \tilde{C}\left(\frac{4\rho_3}{r_m^{-\alpha}\rho}\right)^{\alpha_0/2} \le \frac{\varepsilon}{2},$$

where $\alpha_0, \tilde{C}, \rho_3$ and ε_1 are as in Corollary 7.16. Set

$$\delta = \left(\frac{4\rho_3}{r_m^{-\alpha}\rho}\right)^{1/2}.$$

By Corollary 7.16, for any $x_1, x_2 \in B_{1/2}$ with $|x_1 - x_2| \leq \delta$, by choosing $x_3 \in B_{1/2}$ with $|x_1 - x_3| = |x_2 - x_3| = \delta$, we have for any $m \geq m_0$,

$$|\tilde{u}_m(x_1) - \tilde{u}_m(x_2)| \le |\tilde{u}_m(x_1) - \tilde{u}_m(x_3)| + |\tilde{u}_m(x_2) - \tilde{u}_m(x_3)| \le 2C\delta^{\alpha_0} \le \varepsilon.$$

Thus, \tilde{u}_m are equicontinuous. By Arzelà-Ascoli theorem, there exists $\tilde{u} \in C(\bar{B}_{1/2})$ such that $\tilde{u}_m \to \tilde{u}$ in $L^{\infty}(B_{1/2})$ (up to a subsequence and similarly hereinafter).

Since F_m are ρ -uniformly elliptic,

$$\lambda I \le D_M F_m(0,0,0,0) \le \Lambda I, \ \forall \ m \ge 1.$$

Hence, there exists a constant symmetric matrix $A = (A^{ij})_{n \times n}$ such that

$$D_M F_m(0,0,0,0) \to A.$$

Now, we show that \tilde{u} is a viscosity solution of

(9.4)
$$A^{ij}\tilde{u}_{ij} = 0 \text{ in } B_{1/2}$$

Given $\tilde{x}_0 \in B_{1/2}$ and $\varphi \in C^2$ touching \tilde{u} strictly by above at \tilde{x}_0 . Then there exist a sequence of $\tilde{x}_m \to \tilde{x}_0$ such that $\varphi + c_m$ touch \tilde{u}_m by above at \tilde{x}_m and $c_m \to 0$. By the definition of viscosity solution, for m large enough (e.g. $r_m^{-\alpha} \rho > 2 \|\varphi\|_{C^2(\bar{B}_{1/2})})$,

$$\tilde{F}_m(D^2\varphi(\tilde{x}_m), D\varphi(\tilde{x}_m), \varphi(\tilde{x}_m) + c_m, \tilde{x}_m) \ge \tilde{f}_m(\tilde{x}_m).$$

We compute

$$\begin{split} F_m(D^2\varphi(\tilde{x}_m), D\varphi(\tilde{x}_m), \varphi(\tilde{x}_m) + c_m, \tilde{x}_m) \\ &= \frac{1}{r_m^{\alpha}} F_m(r_m^{\alpha}D^2\varphi + D^2P_m, r_m^{1+\alpha}D\varphi + DP_m, r_m^{2+\alpha}(\varphi + c_m) + P_m, r_m\tilde{x}_m) \\ &- \frac{1}{r_m^{\alpha}} F_m(r_m^{\alpha}D^2\varphi + D^2P_m, r_m^{1+\alpha}D\varphi + DP_m, r_m^{2+\alpha}(\varphi + c_m) + P_m, 0) \\ &+ \frac{1}{r_m^{\alpha}} F_m(r_m^{\alpha}D^2\varphi + D^2P_m, r_m^{1+\alpha}D\varphi + DP_m, r_m^{2+\alpha}(\varphi + c_m) + P_m, 0) \\ &- \frac{1}{r_m^{\alpha}} F_m(r_m^{\alpha}D^2\varphi + D^2P_m, DP_m(0), P_m(0), 0) \\ &+ \frac{1}{r_m^{\alpha}} F_m(r_m^{\alpha}D^2\varphi + D^2P_m, DP_m(0), P_m(0), 0) \\ &- \frac{1}{r_m^{\alpha}} F_m(D^2P_m, DP_m(0), P_m(0), 0) + \frac{1}{r_m^{\alpha}} f_m(0) \\ &\leq \frac{1}{m} + Cb_0 r_m^{1-\alpha} + Cc_0 r_m^{1-\alpha} \\ &+ F_{m,M_{ij}}(\theta r_m^{\alpha}D^2\varphi(\tilde{x}_m) + D^2P_m, DP_m(0), P_m(0), 0) \varphi_{ij}(\tilde{x}_m) + \frac{1}{r_m^{\alpha}} f_m(0), \end{split}$$

where $0 < \theta < 1$. Note that in above inequality, the variable of φ is \tilde{x}_m and the variable of P_m is $x_m := r_m \tilde{x}_m$. Hence,

$$0 \leq F_m(D^2 \varphi(\tilde{x}_m), D\varphi(\tilde{x}_m), \varphi(\tilde{x}_m) + c_m, \tilde{x}_m) - f_m(\tilde{x}_m)$$

$$\leq \frac{1}{m} + Cb_0 r_m^{1-\alpha} + Cc_0 r_m^{1-\alpha}$$

$$+ F_{m,M_{ij}}(0, 0, 0, 0)\varphi_{ij}(\tilde{x}_m) + \frac{1}{r_m^{\alpha}} |f_m(r_m \tilde{x}_m) - f_m(0)| + \omega_F(\frac{C}{m})$$

Let $m \to \infty$, we have

(9.5)
$$A^{ij}\varphi_{ij}(\tilde{x}_0) \ge 0.$$

Hence, \tilde{u} is a subsolution of (9.4). Similarly, we can prove that it is a viscosity supersolution as well. That is, \tilde{u} is a viscosity solution. bounded and equicontinuous in any compact set of S^n . Then there exists a linear operator \tilde{F} with ellipticity constants λ and Λ such that $\tilde{F}_m \to \tilde{F}$ and $D\tilde{F}_m \to D\tilde{F}$ uniformly in any compact set of S^n .

Since (9.4) is a linear equation with constant coefficients, $\tilde{u} \in C^{\infty}(B_{1/2})$. Then there exists $\tilde{P} \in \mathcal{P}_2$ such that for any $0 < \eta < 1/4$,

$$\|\tilde{u} - \tilde{P}\|_{L^{\infty}(B_{\eta})} \le C_1 \eta^3 \|\tilde{u}\|_{L^{\infty}(B_{1/2})} \le C_1 \eta^3,$$
$$A^{ij} \tilde{P}_{ij} = 0$$

and

$$\|\tilde{P}\| \le C_2 \|\tilde{u}\|_{L^{\infty}(B_{1/2})} \le C_2,$$

where C_1 and C_2 are universal. By taking η small and \overline{C} large such that

$$C\eta^{1-\alpha} \le \frac{1}{2}, \quad C_2 \le (\bar{C}-1)\eta^{2+\alpha}.$$

Then

(9.6)
$$\|\tilde{u} - \tilde{P}\|_{L^{\infty}(B_{\eta})} \le \frac{1}{2}\eta^{2+\alpha}, \quad \|\tilde{P}\| \le (\bar{C} - 1)\eta^{2+\alpha}.$$

By a similar argument to prove (9.5),

$$\tilde{F}_m(D^2\tilde{P}, D\tilde{P}(0), \tilde{P}(0), 0) \to A^{ij}\tilde{P}_{ij} = 0.$$

Note that $\tilde{f}_m(0) \to 0$ as well. Hence, there exist a sequence of constants $t_m \to 0$ such that

$$\tilde{F}_m(D^2\tilde{P} + t_m I, D\tilde{P}(0), \tilde{P}(0), 0) = \tilde{f}_m(0).$$

Let

$$Q_m(x) = P_m(x) + r_m^{2+\alpha} \left(\tilde{P}(\tilde{x}) + \frac{t_m}{2} |\tilde{x}|^2 \right).$$

Then with the aid of (9.6),

$$F_m(D^2Q_m, DQ_m(0), Q_m(0), 0) = f_m(0), \quad ||Q_m - P_m|| \le \bar{C}(\eta r_m)^{2+\alpha}.$$

Hence, (9.1) holds for Q_m . That is,

$$||u_m - Q_m||_{L^{\infty}(B_{\eta}r_m)} > (\eta r_m)^{2+\alpha}.$$

Equivalently,

$$\|\tilde{u}_m - \tilde{P} - \frac{t_m}{2} |\tilde{x}|^2 \|_{L^{\infty}(B_{\eta})} > \eta^{2+\alpha}.$$

Let $m \to \infty$, we have

$$\|\tilde{u} - \tilde{P}\|_{L^{\infty}(B_{\eta})} \ge \eta^{2+\alpha},$$

which contradicts with (9.6).

Now, we give the

Proof of Theorem 1.7. To prove Theorem 1.7, we only to prove the

following. There exist a sequence of $P_m \in \mathcal{P}_2 \ (m \ge -1)$ such that for all $m \ge 0$,

(9.7)
$$\|u - P_m\|_{L^{\infty}(B_{\eta}m_{r_0})} \le (\eta^m r_0)^{2+\alpha},$$

and

(9.8)

$$F(D^2 P_m, DP_m(0), P_m(0), 0) = f(0), \quad ||P_m - P_{m-1}||_{\eta^m r_0} \le \bar{C}(\eta^m r_0)^{2+\alpha},$$

where η, r_0 and \bar{C} are as in Lemma 9.1.

We prove above by induction. Set $P_{-1} \equiv 0$. For m = 0, since F(0, 0, 0, 0) = 0, there exists $t \in \mathbb{R}$ such that

$$F(tI, 0, 0, 0) = f(0), \quad |t| \le \frac{|f(0)|}{n\lambda} \le \frac{\delta}{n\lambda}.$$

Then by choosing $P_0 = t|x|^2/2$ and $\delta = (1 + 1/n\lambda)r_0^{2+\alpha}$,

$$\|u - P_0\|_{L^{\infty}(B_{r_0})} \le \|u\|_{L^{\infty}(B_{r_0})} + \|P_0\|_{L^{\infty}(B_{r_0})} \le \delta + \frac{\delta}{2n\lambda} \le r_0^{2+\alpha}.$$

Hence, (9.7) and (9.8) hold for m = 0. Suppose that the conclusion holds for $m \leq m_0$. By (9.8),

$$\|P_{m_0}\| \le \sum_{i=1}^{m_0} \|P_i - P_{i-1}\| + \|P_0\| \le \bar{C}r_0^{2+\alpha} \frac{\eta^{2+\alpha}}{1-\eta^{2+\alpha}} + r_0^{2+\alpha} \le \bar{C}r_0^{2+\alpha}.$$

By Lemma 9.1, the conclusion holds for $m = m_0 + 1$. By induction, the proof of Theorem 1.7 is completed.

Remark 9.2. Note that we can not prove the pointwise $C^{1,\alpha}$ regularity for a general operator F. The reason is the following. For the $C^{1,\alpha}$ regularity, we will consider for some $P_m \in \mathcal{P}_1$

$$\tilde{x} = \frac{x}{r_m}, \quad \tilde{u}_m(\tilde{x}) = \frac{u_m(x) - P_n(x)}{r_m^{1+\alpha}}$$

instead of (9.2) in the proof of Lemma 9.1. Then \tilde{u}_m are solutions of

$$\tilde{F}_m(D^2\tilde{u}_m, D\tilde{u}_m, \tilde{u}_m, \tilde{x}) := r_m^{1-\alpha}F_m(r_m^{\alpha-1}D^2u_m, r_m^{\alpha}Du_m, r_m^{\alpha+1}u_m, x) = r^{1-\alpha}f_m$$

Note that \tilde{F}_m are only $r_m^{1-\alpha}\rho$ -uniformly elliptic and $r_m^{1-\alpha}\rho \to 0$. Hence, we can not proceed the scaling argument.

YUANYUAN LIAN AND KAI ZHANG

10. Interior $C^{k,\alpha}$ regularity

In this section, we prove the interior pointwise $C^{k,\alpha}$ regularity Theorem 1.9. Since $k \ge 2$, the assumption of Theorem 1.9 is stronger than that of Theorem 1.7. Hence, $u \in C^{2,\alpha}(0)$. Then we only need to prove:

Claim I: If $u \in C^{k-1,\alpha}(0)$ under the assumptions of Theorem 1.9, then $u \in C^{k,\alpha}(0)$.

Indeed, we can prove Theorem 1.9 by induction if **Claim I** has been proved. For k = 3, by Theorem 1.7, $u \in C^{2,\alpha}(0)$. Then **Claim I** implies $u \in C^{3,\alpha}(0)$. Hence, Theorem 1.9 holds for k = 3. We assume that Theorem 1.9 holds for $k = k_0$ and we only need to prove it for $k_0 + 1$. Since Theorem 1.9 holds for k_0 and the assumptions of Theorem 1.9 with $k_0 + 1$ is stronger than that with k_0 , $u \in C^{k_0,\alpha}(0)$. Then **Claim I** implies $u \in C^{k_0+1,\alpha}(0)$. Therefore, Theorem 1.9 holds for $k = k_0 + 1$. By induction, the proof of Theorem 1.9 is completed. Thus, in this section, we only need to prove **Claim I** and assume that $u \in C^{k-1,\alpha}(0)$ throughout this section.

We first prove a special result.

Lemma 10.1. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of

$$F(D^2u, Du, u, x) = f \quad in \ B_1,$$

where F is ρ -uniformly elliptic. Suppose that (1.9) and (1.12) hold (with δ_0) and

 $u(0) = \cdots = |D^{k-1}u(0)| = 0, \quad ||u||_{L^{\infty}(B_1)} \leq \delta_0, \quad |f(x)| \leq \delta_0 |x|^{k-2+\alpha}, \ \forall x \in B_1,$ where $\delta_0 > 0$ depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, \alpha, k$ and K_F . Assume for some $\bar{P} \in \mathcal{HP}_k$,

(10.1)
$$\|\bar{P}\| \le \delta_0$$
, $|F_0(D^2\bar{P}(x), D\bar{P}(x), \bar{P}(x), x)| \le \bar{C}|x|^{k-1}, \ \forall \ x \in B_1$,

where \overline{C} depending only on $n, \lambda, \Lambda, \rho, b_0, c_0, \alpha, k$ and K_F , is to specified in Lemma 10.3.

Then $u \in C^{k,\alpha}(0)$. That is, there exists $P \in \mathcal{HP}_k$ such that

$$|u(x) - P(x)| \le C|x|^{k+\alpha}, \ \forall \ x \in B_1$$

and

$$||P|| \le C\delta_0, \quad |F_0(D^2P(x), DP(x), P(x), x)| \le \bar{C}|x|^{k-1}, \ \forall \ x \in B_1,$$

where C depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, \alpha, k$ and K_F .

Remark 10.2. Since we have assumed $u \in C^{k-1,\alpha}(0)$, $D^i u(0)$ $(1 \le i \le k-1)$ are well defined.

To prove Lemma 10.1, we prove the following key step by the compactness method as before.

Lemma 10.3. Let $0 < \alpha < 1$ and $u \in C(\overline{B}_1)$ be a viscosity solution of

$$F(D^2u, Du, u, x) = f \quad in \ B_1,$$

where F is ρ -uniformly elliptic. Suppose that (1.9) holds. Let $r \leq r_0$ and assume that for some $P_0 \in \mathcal{HP}_k$, (1.12) holds (with δ_1),

$$u(0) = \dots = |D^{k-1}u(0)| = 0, \quad ||u - P_0||_{L^{\infty}(B_r)} \le r^{k+\alpha},$$
$$|f(x)| \le \delta_1 |x|^{k-2+\alpha}, \ \forall \ x \in B_1$$

and

$$||P_0|| \le \bar{C}r_0^{k+\alpha}, \quad |F_0(D^2P_0(x), DP_0(x), P_0(x), x)| \le \bar{C}|x|^{k-1}, \ \forall \ x \in B_1,$$

where \overline{C} depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, \alpha$ and k, and $0 < r_0, \delta_1 < 1$ (small) depend also on K_F .

Then there exists $P \in \mathcal{HP}_k$ such that

$$\|u - P\|_{L^{\infty}(B_{\eta r})} \le (\eta r)^{k+\epsilon}$$

and

$$||P - P_0||_r \le \bar{C}(\eta r)^{2+\alpha}, \quad |F_0(D^2P(x), DP(x), P(x), x)| \le \bar{C}|x|^{k-1},$$

where $0 < \eta < 1/2$ depends only on $n, \lambda, \Lambda, \rho, b_0, c_0, \alpha$ and k.

Proof. We prove the lemma by contradiction. Suppose that the conclusion is false. Then there exist sequences of $F_m, F_{0m}, u_m, f_m, P_m, r_m$ such that $r_m \leq 1/m$,

$$F_m(D^2 u_m, Du_m, u_m, x) = f_m \text{ in } B_{r_m},$$
$$u_m(0) = \dots = |D^{k-1} u_m(0)| = 0, \quad ||u_m - P_m||_{L^{\infty}(B_{r_m})} \le r_m^{k+\alpha},$$
$$|f_m(x)| \le \frac{1}{m} |x|^{k-2+\alpha}, \ \forall \ x \in B_1$$

and

(10.2)

$$||P_m|| \le \frac{C}{m}, \quad |F_{0m}(D^2 P_m(x), DP_m(x), P_m(x), x)| \le \frac{1}{m} |x|^{k-1}, \ \forall \ x \in B_1,$$

where F_m, F_{0m} are ρ -uniformly elliptic and (1.12) (with δ replaced by 1/m) holds for F_m and F_{0m} . Furthermore, $F_{0m} \in C^{k-1}$ (with the same bound K_F). Moreover, for any $P \in \mathcal{HP}_k$ with (10.3)

$$|F_{0m}(D^2P(x), DP(x), P(x), x)| \le \bar{C}|x|^{k-1}, \quad ||P - P_m|| \le \bar{C}(\eta r_m)^{k+\alpha},$$

we have

(10.4)
$$||u_m - P||_{L^{\infty}(B_{\eta r_m})} > (\eta r_m)^{k+\alpha}$$

where \bar{C} and $0 < \eta < 1/2$ are to be specified later.

Let

$$\tilde{x} = \frac{x}{r_m}, \quad \tilde{u}_m(\tilde{x}) = \frac{u_m(x) - P_m(x)}{r_m^{k+\alpha}}.$$

As before, \tilde{u}_m are viscosity solutions of

(10.5)
$$\tilde{F}_m(D^2\tilde{u}_m, D\tilde{u}_m, \tilde{u}_m, \tilde{x}) = \tilde{f}_m \text{ in } B_1.$$

where

$$\tilde{F}_m(M, p, s, \tilde{x})$$

 $=r_m^{-(k-2+\alpha)}F_m(r_m^{k-2+\alpha}M + D^2P_m(x), r_m^{k-1+\alpha}p + DP_m(x), r_m^{k+\alpha}s + P_m(x), x)$
 $-r_m^{-(k-2+\alpha)}F_m(D^2P_m(x), DP_m(x), P_m(x), x),$
 $\tilde{f}_m(\tilde{x}) = r_m^{-(k-2+\alpha)}f_m(x) - r_m^{-(k-2+\alpha)}F_m(D^2P_m(x), DP_m(x), P_m(x), x).$
Next, by Proposition 7.4, for m large anough

Next, by Proposition 7.4, for m large enough,

$$\tilde{u}_m \in S^*_{\frac{1}{2}r_m^{-(k-2+\alpha)}\rho}(\lambda, \Lambda, r_m b_0, \bar{f}_m),$$

where

 $\bar{f}_m(\tilde{x}) = |r_m^{-(k-2+\alpha)} \left(f_m(x) - F_m(D^2 P_m, DP_m(x), P_m(x), x) \right) | + r_m^2 c_0 |u_m(x)|.$ By the assumptions,

$$\|\bar{f}_m\|_{L^{\infty}(B_1)} \le \frac{1}{m} + r_m^{1-\alpha} + c_0 r_m^{k+2+\alpha}.$$

Hence, for any $\varepsilon > 0$, we can take m_0 large enough such that for any $m \ge m_0$,

$$\|\bar{f}_m\|_{L^{\infty}(B_1)} \le \varepsilon_1, \quad \tilde{C}\left(\frac{4\rho_3}{r_m^{-(k-2+\alpha)}\rho}\right)^{\alpha_0/2} \le \frac{\varepsilon}{2},$$

where $\alpha_0, \tilde{C}, \rho_3$ and ε_1 are as in Corollary 7.16. Set

$$\delta = \left(\frac{4\rho_3}{r_m^{-(k-2+\alpha)}\rho}\right)^{1/2}$$

By Corollary 7.16, for any $x_1, x_2 \in B_{1/2}$ with $|x_1 - x_2| \leq \delta$, by choosing $x_3 \in B_{1/2}$ with $|x_1 - x_3| = |x_2 - x_3| = \delta$, we have for any $m \geq m_0$,

$$|\tilde{u}_m(x_1) - \tilde{u}_m(x_2)| \le |\tilde{u}_m(x_1) - \tilde{u}_m(x_3)| + |\tilde{u}_m(x_2) - \tilde{u}_m(x_3)| \le 2\tilde{C}\delta^{\alpha_0} \le \varepsilon.$$

Thus, \tilde{u}_m are equicontinuous. Then there exists $\tilde{u} \in C(\bar{B}_{1/2})$ such that $\tilde{u}_m \to \tilde{u}$ in $L^{\infty}(B_{1/2})$.

As before, there exists a constant symmetric matrix A such that

$$D_M F_{0m}(0,0,0,0) \to A$$

Now, we show that \tilde{u} is a viscosity solution of

(10.6)
$$A^{ij}\tilde{u}_{ij} = 0$$
 in $B_{1/2}$.

Given $\tilde{x}_0 \in B_{1/2}$ and $\varphi \in C^2$ touching \tilde{u} strictly by above at \tilde{x}_0 . Then there exist a sequence of $\tilde{x}_m \to \tilde{x}_0$ such that $\varphi + c_m$ touch \tilde{u}_m by above at \tilde{x}_m and $c_m \to 0$. By the definition of viscosity solution, for *m* large enough,

$$F_m(D^2\varphi(\tilde{x}_m), D\varphi(\tilde{x}_m), \varphi(\tilde{x}_m) + c_m, \tilde{x}_m) \ge f_m(\tilde{x}_m).$$

59

Note that

$$\begin{split} \tilde{f}_{m}(\tilde{x}_{m}) &\leq \tilde{F}_{m}(D^{2}\varphi(\tilde{x}_{m}), D\varphi(\tilde{x}_{m}), \varphi(\tilde{x}_{m}) + c_{m}, \tilde{x}_{m}) \\ &= \frac{1}{r_{m}^{k-2+\alpha}} F_{m}(r_{m}^{k-2+\alpha}D^{2}\varphi + D^{2}P_{m}, r_{m}^{k-1+\alpha}D\varphi + DP_{m}, r_{m}^{k+\alpha}(\varphi + c_{m}) + P_{m}, r_{m}\tilde{x}_{m}) \\ &- \frac{1}{r_{m}^{k-2+\alpha}} F_{0m}(r_{m}^{k-2+\alpha}D^{2}\varphi + D^{2}P_{m}, r_{m}^{k-1+\alpha}D\varphi + DP_{m}, r_{m}^{k+\alpha}(\varphi + c_{m}) + P_{m}, r_{m}\tilde{x}) \\ &+ \frac{1}{r_{m}^{k-2+\alpha}} F_{0m}(r_{m}^{k-2+\alpha}D^{2}\varphi + D^{2}P_{m}, r_{m}^{k-1+\alpha}D\varphi + DP_{m}, r_{m}^{k+\alpha}(\varphi + c_{m}) + P_{m}, r_{m}\tilde{x}) \\ &- \frac{1}{r_{m}^{k-2+\alpha}} F_{0m}(D^{2}P_{m}, DP_{m}, P_{m}, r_{m}\tilde{x}_{m}) \\ &\leq \frac{1}{m} + F_{0m,M_{ij}}(\xi)\varphi_{ij}(\tilde{x}_{m}) + r_{m}F_{0m,p_{i}}(\xi)\varphi_{i}(\tilde{x}_{m}) + r_{m}^{2}F_{0m,s}(\xi)\varphi(\tilde{x}_{m}), \end{split}$$

where

$$\xi = (\theta r_m^{k-2+\alpha} D^2 \varphi + D^2 P_m, \theta r_m^{k-1+\alpha} D \varphi + D P_m, \theta r_m^{k+\alpha} (\varphi + c_m) + P_m, r_m \tilde{x}_m)$$

for some $0 < \theta < 1$. Let $m \to \infty$, we have

for some $0 < \theta < 1$. Let $m \to \infty$, we have

$$A^{ij}\varphi_{ij}(\tilde{x}_0) \ge 0.$$

Hence, \tilde{u} is a subsolution of (10.6). Similarly, we can prove that it is a viscosity supersolution as well. That is, \tilde{u} is a viscosity solution.

Next, we show $\tilde{u}_m \in C^{k-1,\alpha}(0)$ and their norms have a uniform bound for *m* large enough. Take δ small (to be specified later) and set $\hat{u}_m = \delta \tilde{u}_m$. Then \hat{u}_m are viscosity solutions of

(10.7)
$$\hat{F}_m(D^2\hat{u}_m, D\hat{u}_m, \hat{u}_m, \tilde{x}) = \hat{f}_m \text{ in } B_1,$$

where

$$\hat{F}_m(M, p, s, \tilde{x}) = \delta \tilde{F}_m(\delta^{-1}M, \delta^{-1}p, \delta^{-1}s, \tilde{x}), \quad \hat{f}_m(\tilde{x}) = \delta \tilde{f}_m(\tilde{x})$$

and we define \hat{F}_{0m} similarly.

Since $r_m \to 0$ and $||P_m|| \to 0$, for *m* large enough (similarly in the following argument), \hat{F}_m are ρ -uniformly elliptic (although δ is very small) and

$$\|\hat{u}_m\|_{L^{\infty}(B_1)} \le \delta, \quad \|\hat{f}_m\|_{C^{k-2+\alpha}(0)} \le \delta$$

In addition, $\hat{F}_m(0,0,0,\tilde{x}) \equiv 0$ obviously and it can be verified easily that \hat{F}_m satisfies (1.9) with b_0 and c_0 . Next, for any $|M|, |p|, |s| \leq \rho$ and $\tilde{x} \in B_1$,

$$|\hat{F}_m(M, p, s, \tilde{x}) - \hat{F}_{0m}(M, p, s, \tilde{x})| \le \frac{2}{m} |\tilde{x}|^{k-2+\alpha} \le \delta |\tilde{x}|^{k-2+\alpha}.$$

Finally, we show

(10.8)
$$\|\hat{F}_{0m}\|_{C^{k-1}(\bar{\mathbf{B}}_{\rho}\times\bar{B}_1)} \leq \hat{K}_F,$$

where \hat{K}_F depends only on n, ρ and K_F . In fact, note that

$$\hat{F}_{0m}(M, p, s, \tilde{x}) = \hat{F}_{0m}(M, p, s, \tilde{x}) - \hat{F}_{0m}(0, 0, 0, \tilde{x})
= \int_0^1 F_{0m, M_{ij}}(\xi) M_{ij} + r_m F_{0m, p_i}(\xi) p_i + r_m^2 F_{0m, s}(\xi) s dt$$

where

$$\xi = t \left(\delta^{-1} r_m^{k-2+\alpha} M, \delta^{-1} r_m^{k-1+\alpha} p, \delta^{-1} r_m^{k+\alpha} s, x \right) + \left(D^2 P_m(x), D P_m(x), P_m(x), x \right).$$

Hence,

$$\|\hat{F}_{0m}\|_{C^{k-2}(\bar{\mathbf{B}}_{\rho}\times\bar{B}_{1})}\leq\hat{K}_{F}$$

Next, from the definition of \hat{F}_{0m} , if any (k-1)-th derivative of \hat{F}_{0m} involves one derivative with respect to M, p or s, it is bounded; if we take (k-1)-th derivative with respect to \tilde{x} ,

$$D_{\tilde{x}}^{k-1} \hat{F}_{0m}(M, p, s, \tilde{x}) = r_m^{1-\alpha} D_x^{k-1} \Big(F_{0m}(\delta^{-1} r_m^{k-2+\alpha} M + D^2 P_m, \delta^{-1} r_m^{k-1+\alpha} p + D P_m, \delta^{-1} r_m^{k+\alpha} s + P_m, x) \Big) \\ \leq \hat{K}_F, \ \forall \ (M, p, s, x) \in \bar{\mathbf{B}}_{\rho} \times \bar{B}_1.$$

Therefore, (10.8) holds.

Choose δ small enough such that Theorem 1.9 holds for δ and \hat{K}_F . By induction, $\hat{u}_m \in C^{k-1,\alpha}(0)$ and hence $\tilde{u}_m \in C^{k-1,\alpha}(0)$ and their norms have a uniform bound for *m* large enough. Since $u_m(0) = \cdots = D^{k-1}u_m(0) = 0$ and $P_m \in \mathcal{HP}_{k-1}$,

$$|\tilde{u}_m(\tilde{x})| \le C |\tilde{x}|^{k-1+\alpha}, \ \forall \ \tilde{x} \in B_1,$$

where C is a constant independent of m. By taking $m \to \infty$,

(10.9)
$$|\tilde{u}(\tilde{x})| \le C |\tilde{x}|^{k-1+\alpha}, \ \forall \ \tilde{x} \in B_1$$

Since (10.6) is a linear equation with constant coefficients, $\tilde{u} \in C^{\infty}(B_{1/2})$. By noting (10.9), there exists $\tilde{P} \in \mathcal{HP}_k$ such that for any $0 < \eta < 1/4$,

$$\|\tilde{u} - \tilde{P}\|_{L^{\infty}(B_{\eta})} \le C_1 \eta^{k+1} \|\tilde{u}\|_{L^{\infty}(B_{1/2})} \le C_1 \eta^{k+1},$$
$$A^{ij} \tilde{P}_{ij} \equiv 0$$

and

$$\|\tilde{P}\| \le C_2, \quad \|\tilde{u}\|_{L^{\infty}(B_{1/2})} \le C_2,$$

where C_1 and C_2 depend only on n, λ, Λ and k. By taking η small and \overline{C} large such that

$$C\eta^{1-\alpha} \le \frac{1}{2}, \quad C_2 \le (\bar{C}-1)\eta^{k+\alpha}.$$

Then

(10.10)
$$\|\tilde{u} - \tilde{P}\|_{L^{\infty}(B_{\eta})} \leq \frac{1}{2}\eta^{k+\alpha}, \quad \|\tilde{P}\| \leq (\bar{C} - 1)\eta^{k+\alpha}.$$

Now, we try to construct a sequence of $\tilde{P}_m \in \mathcal{HP}_k$ such that $\tilde{P}_m \to \tilde{P}$ as $m \to \infty$ and

(10.11)
$$D^{i}G_{m}(0) = 0, \ \forall \ 1 \le i \le k-2,$$

where

$$G_m(\tilde{x}) = \tilde{F}_{0m}(D^2 \tilde{P}_m(\tilde{x}), D\tilde{P}_m(\tilde{x}), \tilde{P}_m(\tilde{x}), \tilde{x}).$$

The (10.11) implies

$$D^{i}\left(F_{0m}(D^{2}Q(x), DQ(x), Q(x), x)\right)\Big|_{x=0} = 0, \ \forall \ 1 \le i \le k-2,$$

where

$$Q_m(x) = P_m(x) + r_m^{k+\alpha} \tilde{P}_m(\tilde{x}).$$

Then (10.3) holds since $F_{0m} \in C^{k-1}$.

To prove (10.11), let $\hat{P}_m \in \mathcal{HP}_k$ $(m \ge 1)$ with $\|\hat{P}_m\| \le 1$ to be specified later and

$$\tilde{P}_m(\tilde{x}) = \hat{P}_m(\tilde{x}) + \tilde{P}(\tilde{x}).$$

Since $\tilde{P}_m \in \mathcal{HP}_k$ and $\tilde{F}_{0m}(0,0,0,\tilde{x}) \equiv 0$,

$$\begin{aligned} |G_m(\tilde{x})| &= |\tilde{F}_{0m}(D^2 \tilde{P}_m(\tilde{x}), D\tilde{P}_m(\tilde{x}), \tilde{P}_m(\tilde{x}), \tilde{x}) - \tilde{F}_m(0, 0, 0, \tilde{x})| \\ &= |\int_0^1 \tilde{F}_{0m, M_{ij}}(\xi) \tilde{P}_{m, ij} + \tilde{F}_{0m, p_i}(\xi) \tilde{P}_{m, i} + \tilde{F}_{0m, s}(\xi) \tilde{P}_m dt| \\ &\leq C |\tilde{x}|^{k-2}, \end{aligned}$$

where

$$\xi = (tD^2 \tilde{P}_m(x), tD\tilde{P}_m(\tilde{x}), t\tilde{P}_m(\tilde{x}), \tilde{x}).$$

Hence, to verify (10.11) for \tilde{P}_m , we only need to prove $D^{k-2}G_m(0) = 0$. Indeed, since $\tilde{P}_m \in \mathcal{HP}_k$,

$$D^{k-2}G_m(0) = \tilde{F}_{0m,M_{ij}}(0,0,0,0)D^{k-2}\tilde{P}_{m,ij}$$

= $\tilde{F}_{0m,M_{ij}}(0,0,0,0)D^{k-2}\tilde{P}_{ij} + \tilde{F}_{0m,M_{ij}}(0,0,0,0)D^{k-2}\hat{P}_{m,ij},$

Since

$$\tilde{F}_{0m,M_{ij}}(0,0,0,0)D^{k-2}\tilde{P}_{ij} \to \tilde{A}^{ij}D^{k-2}\tilde{P}_{ij} = 0$$

and $\lambda \leq \tilde{F}_{0m,M_{ij}}(0) \leq \Lambda$, we can choose proper \hat{P}_m such that

$$D^{k-2}G_m(0) = 0, \ \forall \ m \ge 1 \ \text{ and } \|\hat{P}_m\| \to 0.$$

Hence, (10.3) holds for Q_m . Then,

$$||u_m - Q_m||_{L^{\infty}(B_\eta r_m)} > (\eta r_m)^{k+\alpha}.$$

Equivalently,

$$\|\tilde{u}_m - \tilde{P} - \hat{P}_m\|_{L^{\infty}(B_{\eta})} > \eta^{k+\alpha}.$$

Let $m \to \infty$, we have

$$\|\tilde{u} - \tilde{P}\|_{L^{\infty}(B_{\eta})} \ge \eta^{k+\alpha},$$

which contradicts with (10.4).

Now, we give the

Proof of Lemma 10.1. To prove Lemma 10.1, we only to prove the following. There exist a sequence of $P_m \in \mathcal{HP}_k$ $(m \ge -1)$ such that for all $m \ge 0$,

(10.12)
$$\|u - P_m\|_{L^{\infty}(B_{\eta^m r_0})} \le (\eta^m r_0)^{k+\alpha},$$

(10.13)
$$|F_0(D^2P_m(x), DP_m(x), P_m(x), x)| \le \hat{C}|x|^{k-1}, \ \forall \ x \in B_1,$$

and

(10.14)
$$||P_m - P_{m-1}||_{\eta^m r_0} \le \bar{C}(\eta^m r_0)^{k+\alpha},$$

where η, r_0 and \bar{C} are as in Lemma 10.3.

We prove above by induction. Set $P_0 = \overline{P}$ and $P_{-1} \equiv 0$. Choose δ_0 such that

$$\delta_0 = \frac{1}{2} r_0^{k+\alpha}.$$

Then by (10.1) and noting

$$\|u - P_0\|_{L^{\infty}(B_{r_0})} \le \|u\|_{L^{\infty}(B_{r_0})} + \|P_0\|_{L^{\infty}(B_{r_0})} \le 2\delta_0 \le r_0^{k+\alpha},$$

(10.12)-(10.14) hold for m = 0. Suppose that the conclusion holds for $m \le m_0$. By (10.14),

$$\|P_{m_0}\| \le \sum_{i=1}^{m_0} \|P_i - P_{i-1}\| + \|P_0\| \le \bar{C}r_0^{k+\alpha} \frac{\eta^{k+\alpha}}{1 - \eta^{k+\alpha}} + \frac{1}{2}r_0^{k+\alpha} \le \bar{C}r_0^{k+\alpha}.$$

By Lemma 10.3, the conclusion holds for $m = m_0 + 1$. By induction, the proof of Lemma 10.1 is completed.

Now, we give the

Proof of Theorem 1.9. As explained at the beginning of this section, we only need to prove **Claim I**. That is, we assume $u \in C^{k-1,\alpha}(0)$ and need to prove $u \in C^{k,\alpha}(0)$.

Let $F_1(M, p, s, x) = F(M, p, s, x) - P_f(x)$ for $(M, p, s, x) \in \mathcal{S}^n \times \mathbb{R}^n \times \mathbb{R} \times \overline{B_1}$. Then u satisfies

$$F_1(D^2u, Du, u, x) = f_1 \text{ in } B_1,$$

where $f_1(x) = f(x) - P_f(x)$. Thus,

$$|f_1(x)| \le [f]_{C^{k-2,\alpha}(0)} |x|^{k-2+\alpha} \le \delta |x|^{k-2+\alpha}, \ \forall \ x \in B_1.$$

Define

$$P_u(x) = \sum_{|\sigma| \le k-1} \frac{1}{\sigma!} D^{\sigma} u(0) x^{\sigma}.$$

Set $u_1 = u - P_u$ and $F_2(M, p, s, x) = F_2(M + D^2 P_u(x), p + DP_u(x), s + P_u(x), x)$. Then u_1 satisfies

(10.15)
$$F_2(D^2u_1, Du_1, u_1, x) = f_1 \text{ in } B_1$$

and

$$u_1(0) = |Du_1(0)| = \dots = |D^{k-1}u_1(0)| = 0.$$

We define F_{01}, F_{02} in a similar way.

Next, we try to construct $\bar{P} \in \mathcal{HP}_k$ such that (10.1) holds. Since we have known the $C^{k-1,\alpha}(0)$ regularity,

 $|F_{02}(0,0,0,x)| = |F_0(D^2 P_u(x), DP_u(x), P_u(x), x) - P_f(x)| \le \bar{C}|x|^{k-2}, \, \forall x \in B_1,$ which implies

$$D^{i}(F_{02}(0,0,0,x))\Big|_{x=0} = 0, \ \forall \ 0 \le i \le k-3.$$

Since $\bar{P} \in \mathcal{HP}_k$,

$$D^{i}\left(F_{2}(D^{2}\bar{P}, D\bar{P}, \bar{P}, x)\right)\Big|_{x=0} = D^{i}\left(F_{2}(0, 0, 0, x)\right)\Big|_{x=0} = 0, \ \forall \ 0 \le i \le k-3.$$

Next, we compute the k-2 order derivatives. Since $F_0(0,0,0,x) \equiv 0$,

$$\begin{aligned} \left| D_x^{k-2} F_0(D^2 P_u(0), DP_u(0), P_u(0), 0) \right| \\ &= \left| D_x^{k-2} F_0(D^2 P_u(0), DP_u(0), P_u(0), 0) - D_x^{k-2} F_0(0, 0, 0, 0) \right| \\ &\leq K_F \|P_u\| \leq C\delta, \end{aligned}$$

which implies

$$\left| D^{k-2} \left(F_2(0,0,0,x) \right) \right|_{x=0} \right| \le C\delta.$$

Note

$$D^{k-2}\left(F_2(D^2\bar{P}, D\bar{P}, \bar{P}, x)\right)\Big|_{x=0} = F_{2,M_{ij}}(0, 0, 0, 0)D^{k-2}\bar{P}_{ij} + D^{k-2}\left(F_2(0, 0, 0, x)\right)\Big|_{x=0}$$

and $\lambda I \leq D_M F_2 \leq \Lambda I$. Then we can choose proper \overline{P} such that

$$D^{k-2}\left(F_2(D^2\bar{P}, D\bar{P}, \bar{P}, x)\right)\Big|_{x=0}, \quad \|\bar{P}\| \le C\delta,$$

where C depends only on $n, \lambda, \Lambda, \rho, k$ and K_F . Therefore, (10.1) holds by taking δ small enough. By Lemma 10.1, u_1 and hence u is $C^{k,\alpha}$ at 0.

Finally, we give the

Proof of Corollary 1.11. By Theorem 1.7, $u \in C^{2,\alpha}(0)$ and there exists $P \in \mathcal{P}_2$ such that

$$|u(x) - P(x)| \le C|x|^{2+\alpha}, \ \forall \ x \in B_1$$

and

(10.16)
$$||P|| \le \bar{C}\delta, \quad F(D^2P, DP(0), P(0), 0) = f(0).$$

For r > 0 to be specified later, let

$$y = \frac{x}{r}, \quad \tilde{u}(y) = \frac{u(x) - P(x)}{r^2}, \quad \tilde{f}(y) = f(x) - F(D^2 P, DP(x), P(x), x).$$

Then \tilde{u} is a solution of

(10.17)
$$\tilde{F}(D^2\tilde{u}, D\tilde{u}, \tilde{u}, x) = \tilde{f} \quad \text{in } B_1,$$

where

$$\tilde{F}(M,p,s,y)=F(M+D^2P,rp+DP(x),r^2s+P(x),x)-F(D^2P,DP(x),P(x),x).$$
 In addition, define

$$\tilde{F}_0(M, p, s, y) = F_0(M + D^2 P, rp + DP(x), r^2 s + P(x), x) - F_0(D^2 P, DP(x), P(x), x).$$

Next, we show that (10.17) satisfies the assumptions of Theorem 1.9. First, $\tilde{F}_0(0,0,0,y) \equiv 0$ clearly. In addition, by (10.16) and taking δ small enough, \tilde{F} and \tilde{F}_0 are $\rho/2$ -uniformly elliptic and

$$K_{\tilde{F}} = \|\bar{F}_0\|_{C^{k-1}(\bar{\mathbf{B}}_{\rho/2} \times \bar{B}_1)} \le CK_F.$$

Moreover, it can be verified easily that for any $|M|, |p|, |s| \le \rho/2, y \in B_1$,

 $|\tilde{F}(M,p,s,y) - \tilde{F}_0(M,p,s,y)| \le C\delta |x|^{k-2+\alpha} \le C\delta r^{k-2+\alpha} |y|^{k-2+\alpha}.$

Finally, by noting $\tilde{f}(0) = 0$,

$$\|\tilde{u}\|_{L^{\infty}(B_1)} \le Cr^{\alpha}, \quad \|f\|_{C^{k-2+\alpha}(0)} \le Cr.$$

Therefore, we can choose r small enough such that the assumptions of Theorem 1.9 are satisfies. Then $\tilde{u} \in C^{k,\alpha}(0)$ and hence $u \in C^{k,\alpha}(0)$. \Box

STATEMENTS AND DECLARATIONS

Competing Interests: All authors declare that they have no conflicts of interest.

References

- [1] A. D. Aleksandrov. Dirichlet's problem for the equation $\text{Det } ||z_{ij}|| = \varphi(z_1, \dots, z_n, z, x_1, \dots, x_n)$. I. Vestnik Leningrad. Univ. Ser. Mat. Meh. Astronom. 13.1 (1958), pp. 5–24.
- [2] A. Alexandroff. Smoothness of the convex surface of bounded Gaussian curvature. C. R. (Doklady) Acad. Sci. URSS (N.S.) 36 (1942), pp. 195–199.
- [3] A. L. Amadori, B. Brandolini, and C. Trombetti. Viscosity solutions of the Monge-Ampère equation with the right hand side in L^p. Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. 18.3 (2007), pp. 221–233. DOI: 10.4171/rlm/491.
- [4] R. Argiolas, F. Charro, and I. Peral. On the Aleksandrov-Bakel'man-Pucci estimate for some elliptic and parabolic nonlinear operators. *Arch. Ration. Mech. Anal.* 202.3 (2011), pp. 875–917. DOI: 10.1007/s00205-011-0434-y.
- S. N. Armstrong and L. Silvestre. Unique continuation for fully nonlinear elliptic equations. *Math. Res. Lett.* 18.5 (2011), pp. 921–926.
 DOI: 10.4310/MRL.2011.v18.n5.a9.
- [6] S. N. Armstrong, L. E. Silvestre, and C. K. Smart. Partial regularity of solutions of fully nonlinear, uniformly elliptic equations. *Comm. Pure Appl. Math.* 65.8 (2012), pp. 1169–1184. DOI: 10.1002/cpa.21394.

- [7] A. Banerjee. A note on the unique continuation property for fully nonlinear elliptic equations. *Commun. Pure Appl. Anal.* 14.2 (2015), pp. 623–626. DOI: 10.3934/cpaa.2015.14.623.
- [8] J. Bao et al. Liouville property and regularity of a Hessian quotient equation. *Amer. J. Math.* 125.2 (2003), pp. 301–316.
- [9] G. Barles. Interior gradient bounds for the mean curvature equation by viscosity solutions methods. *Differential Integral Equations* 4.2 (1991), pp. 263–275.
- [10] A. Bhattacharya. Hessian estimates for Lagrangian mean curvature equation. Calc. Var. Partial Differential Equations 60.6 (2021), Paper No. 224, 23. DOI: 10.1007/s00526-021-02097-0.
- [11] A. Bhattacharya, C. Mooney, and R. Shankar. Gradient estimates for the Lagrangian mean curvature equation with critical and supercritical phase. 2022. arXiv: 2205.13096 [math.AP].
- [12] A. Bhattacharya and R. Shankar. Optimal regularity for Lagrangian mean curvature type equations. 2020. arXiv: 2009.04613 [math.AP].
- [13] A. Bhattacharya and R. Shankar. Regularity for convex viscosity solutions of Lagrangian mean curvature equation. J. Reine Angew. Math. 803 (2023), pp. 219–232. DOI: 10.1515/crelle-2023-0056.
- [14] E. Bombieri, E. De Giorgi, and M. Miranda. Una maggiorazione a priori relativa alle ipersuperfici minimali non parametriche. Arch. Rational Mech. Anal. 32 (1969), pp. 255–267. DOI: 10.1007/BF00281503.
- [15] T. Bourni. $C^{1,\alpha}$ theory for the prescribed mean curvature equation with Dirichlet data. J. Geom. Anal. 21.4 (2011), pp. 982–1035. DOI: 10.1007/s12220-010-9176-6.
- [16] X. Cabré and L. A. Caffarelli. Regularity for viscosity solutions of fully nonlinear equations $F(D^2u) = 0$. Topol. Methods Nonlinear Anal. 6.1 (1995), pp. 31–48. DOI: 10.12775/TMNA.1995.030.
- [17] L. Caffarelli, R. Kohn, and L. Nirenberg. Partial regularity of suitable weak solutions of the Navier-Stokes equations. *Comm. Pure Appl. Math.* 35.6 (1982), pp. 771–831. DOI: 10.1002/cpa.3160350604.
- [18] L. Caffarelli, L. Nirenberg, and J. Spruck. The Dirichlet problem for nonlinear second-order elliptic equations. I. Monge-Ampère equation. *Comm. Pure Appl. Math.* 37.3 (1984), pp. 369–402. DOI: 10.1002/cpa.3160370306.
- [19] L. Caffarelli, L. Nirenberg, and J. Spruck. The Dirichlet problem for nonlinear second-order elliptic equations. III. Functions of the eigenvalues of the Hessian. Acta Math. 155.3-4 (1985), pp. 261–301. DOI: 10.1007/BF02392544.
- [20] L. Caffarelli, L. Nirenberg, and J. Spruck. The Dirichlet problem for the degenerate Monge-Ampère equation. *Rev. Mat. Iberoamericana* 2.1-2 (1986), pp. 19–27. DOI: 10.4171/RMI/23.
- [21] L. A. Caffarelli. A localization property of viscosity solutions to the Monge-Ampère equation and their strict convexity. Ann. of Math. (2) 131.1 (1990), pp. 129–134. DOI: 10.2307/1971509.

- [22] L. A. Caffarelli. Interior $W^{2,p}$ estimates for solutions of the Monge-Ampère equation. Ann. of Math. (2) 131.1 (1990), pp. 135–150. DOI: 10.2307/1971510.
- [23] L. A. Caffarelli. Some regularity properties of solutions of Monge Ampère equation. *Comm. Pure Appl. Math.* 44.8-9 (1991), pp. 965– 969. DOI: 10.1002/cpa.3160440809.
- [24] L. A. Caffarelli. The regularity of mappings with a convex potential. J. Amer. Math. Soc. 5.1 (1992), pp. 99–104. DOI: 10.2307/2152752.
- [25] L. A. Caffarelli. Boundary regularity of maps with convex potentials.
 II. Ann. of Math. (2) 144.3 (1996), pp. 453–496. DOI: 10.2307/2118564.
- [26] L. A. Caffarelli. Monge Ampère equation div-curl theorems in Lagrangian coordinates compression and rotation, NSF-CBMS Lecture. Florida Atlantic University, 1997.
- [27] L. A. Caffarelli and X. Cabré. Fully nonlinear elliptic equations. Vol. 43. American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 1995, pp. vi+104. DOI: 10.1090/coll/043.
- [28] L. A. Caffarelli and Y. Yuan. A priori estimates for solutions of fully nonlinear equations with convex level set. *Indiana Univ. Math. J.* 49.2 (2000), pp. 681–695. DOI: 10.1512/iumj.2000.49.1901.
- [29] L. A. Caffarelli and Y. Yuan. Singular solutions to Monge-Ampère equation. Anal. Theory Appl. 38.2 (2022), pp. 121–127.
- [30] E. Calabi. Improper affine hyperspheres of convex type and a generalization of a theorem by K. Jörgens. *Michigan Math. J.* 5 (1958), pp. 105–126.
- [31] N. Chaudhuri and N. S. Trudinger. An Alexsandrov type theorem for k-convex functions. Bull. Austral. Math. Soc. 71.2 (2005), pp. 305– 314. DOI: 10.1017/S0004972700038260.
- [32] C. Chen. The interior gradient estimate of Hessian quotient equations. J. Differential Equations 259.3 (2015), pp. 1014–1023. DOI: 10.1016/j.jde.2015.02.035.
- [33] C. Chen, W. Dong, and F. Han. Interior Hessian estimates for a class of Hessian type equations. *Calc. Var. Partial Differential Equations* 62.2 (2023), Paper No. 52, 15. DOI: 10.1007/s00526-022-02385-3.
- [34] C. Chen, L. Xu, and D. Zhang. The interior gradient estimate of prescribed Hessian quotient curvature equations. *Manuscripta Math.* 153.1-2 (2017), pp. 159–171. DOI: 10.1007/s00229-016-0877-4.
- [35] J. Chen, R. Shankar, and Y. Yuan. Regularity for convex viscosity solutions of special Lagrangian equation. *Comm. Pure Appl. Math.* 76.12 (2023), pp. 4075–4086. DOI: 10.1002/cpa.22130.
- [36] J. Chen, M. Warren, and Y. Yuan. A priori estimate for convex solutions to special Lagrangian equations and its application. *Comm. Pure Appl. Math.* 62.4 (2009), pp. 583–595. DOI: 10.1002/cpa.20261.

- [37] L. Chen, Q. Tu, and N. Xiang. Pogorelov type estimates for a class of Hessian quotient equations. J. Differential Equations 282 (2021), pp. 272–284. DOI: 10.1016/j.jde.2021.02.030.
- [38] S. Chen, J. Liu, and X.-J. Wang. Global regularity for the Monge-Ampère equation with natural boundary condition. Ann. of Math. (2) 194.3 (2021), pp. 745–793. DOI: 10.4007/annals.2021.194.3.4.
- [39] J. Cheng and Y. Xu. Interior W^{2,p} estimate for small perturbations to the complex Monge-Ampère equation. Calc. Var. Partial Differential Equations 62.8 (2023), Paper No. 231, 46. DOI: 10.1007/s00526-023-02571-x.
- [40] S. Y. Cheng and S. T. Yau. On the regularity of the Monge-Ampère equation $\det(\partial^2 u/\partial x_i \partial s x_j) = F(x, u)$. Comm. Pure Appl. Math. 30.1 (1977), pp. 41–68. DOI: 10.1002/cpa.3160300104.
- [41] S.-K. Chiu and G. Székelyhidi. Higher regularity for singular Kähler-Einstein metrics. *Duke Math. J.* 172.18 (2023), pp. 3521–3558. DOI: 10.1215/00127094-2022-0107.
- [42] K.-S. Chou and X.-J. Wang. A variational theory of the Hessian equation. Comm. Pure Appl. Math. 54.9 (2001), pp. 1029–1064. DOI: 10.1002/cpa.1016.
- [43] J. Chu and S. Dinew. Liouville theorem for a class of Hessian equations. 2023. arXiv: 2306.13825 [math.AP].
- [44] L. Dai, J. Bao, and B. Wang. Interior derivative estimates and Bernstein theorem for Hessian quotient equations. 2023. arXiv: 2305.17831 [math.AP].
- [45] Q. Dai, N. S. Trudinger, and X.-J. Wang. The mean curvature measure. J. Eur. Math. Soc. (JEMS) 14.3 (2012), pp. 779–800. DOI: 10.4171/JEMS/318.
- [46] E. De Giorgi. Frontiere orientate di misura minima. Seminario di Matematica della Scuola Normale Superiore di Pisa, 1960-61. Editrice Tecnico Scientifica, Pisa, 1961, p. 57.
- [47] G. De Philippis, A. Figalli, and O. Savin. A note on interior $W^{2,1+\varepsilon}$ estimates for the Monge-Ampère equation. *Math. Ann.* 357.1 (2013), pp. 11–22. DOI: 10.1007/s00208-012-0895-9.
- [48] G. De Philippis and A. Figalli. W^{2,1} regularity for solutions of the Monge-Ampère equation. Invent. Math. 192.1 (2013), pp. 55–69. DOI: 10.1007/s00222-012-0405-4.
- [49] S.-Z. Du. Necessary and sufficient conditions to Bernstein theorem of a Hessian equation. *Trans. Amer. Math. Soc.* 375.7 (2022), pp. 4873– 4892. DOI: 10.1090/tran/8686.
- [50] L. C. Evans. Classical solutions of fully nonlinear, convex, secondorder elliptic equations. *Comm. Pure Appl. Math.* 35.3 (1982), pp. 333– 363. DOI: 10.1002/cpa.3160350303.
- [51] A. Figalli. The Monge-Ampère equation and its applications. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2017, pp. x+200. DOI: 10.4171/170.
- [52] R. Finn. On equations of minimal surface type. Ann. of Math. (2) 60 (1954), pp. 397–416. DOI: 10.2307/1969841.

- [53] C. Gerhardt. Existence, regularity, and boundary behavior of generalized surfaces of prescribed mean curvature. *Math. Z.* 139 (1974), pp. 173–198. DOI: 10.1007/BF01418314.
- [54] M. Giaquinta. On the Dirichlet problem for surfaces of prescribed mean curvature. *Manuscripta Math.* 12 (1974), pp. 73–86. DOI: 10.1007/BF01166235.
- [55] M. Giaquinta and E. Giusti. Nonlinear elliptic systems with quadratic growth. *Manuscripta Math.* 24.3 (1978), pp. 323–349. DOI: 10.1007/BF01167835.
- [56] D. Gilbarg and N. S. Trudinger. Elliptic partial differential equations of second order. Classics in Mathematics. Reprint of the 1998 edition. Springer-Verlag, Berlin, 2001, pp. xiv+517.
- [57] E. Giusti. On the equation of surfaces of prescribed mean curvature. Existence and uniqueness without boundary conditions. *Invent. Math.* 46.2 (1978), pp. 111–137. DOI: 10.1007/BF01393250.
- [58] E. Giusti. Minimal surfaces and functions of bounded variation. Vol. 80. Monographs in Mathematics. Birkhäuser Verlag, Basel, 1984, pp. xii+240. DOI: 10.1007/978-1-4684-9486-0.
- [59] P. Guan and G. Qiu. Interior C^2 regularity of convex solutions to prescribing scalar curvature equations. *Duke Math. J.* 168.9 (2019), pp. 1641–1663. DOI: 10.1215/00127094–2019–0001.
- [60] P. Guan, N. S. Trudinger, and X.-J. Wang. On the Dirichlet problem for degenerate Monge-Ampère equations. Acta Math. 182.1 (1999), pp. 87–104. DOI: 10.1007/BF02392824.
- [61] C. E. Gutiérrez. The Monge-Ampère equation. Vol. 44. Progress in Nonlinear Differential Equations and their Applications. Birkhäuser Boston, Inc., Boston, MA, 2001, pp. xii+127. DOI: 10.1007/978-1-4612-0195-3.
- [62] E. Heinz. On elliptic Monge-Ampère equations and Weyl's embedding problem. J. Analyse Math. 7 (1959), pp. 1–52. DOI: 10.1007/BF02787679.
- [63] E. Heinz. Über die Differentialungleichung $0 < \alpha \le rt s^2 \le \beta < \infty$. Math. Z. 72 (1959/60), pp. 107–126. DOI: 10.1007/BF01162942.
- [64] N. M. Ivochkina. Classical solvability of the Dirichlet problem for the Monge-Ampère equation. In: vol. 131. Questions in quantum field theory and statistical physics, 4. 1983, pp. 72–79.
- [65] N. M. Ivochkina. Solution of the Dirichlet problem for certain equations of Monge-Ampère type. Mat. Sb. (N.S.) 128(170).3 (1985), pp. 403–415, 447.
- [66] N. Ivochkina, N. Trudinger, and X.-J. Wang. The Dirichlet problem for degenerate Hessian equations. *Comm. Partial Differential Equations* 29.1-2 (2004), pp. 219–235. DOI: 10.1081/PDE-120028851.
- [67] N. M. Ivočkina. A priori estimate of $|u|_{C_2(\overline{\Omega})}$ of convex solutions of the Dirichlet problem for the Monge-Ampère equation. Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 96 (1980), pp. 69–79, 306.

- [68] N. M. Ivočkina. The integral method of barrier functions and the Dirichlet problem for equations with operators of the Monge-Ampère type. Mat. Sb. (N.S.) 112(154).2(6) (1980), pp. 193–206.
- [69] H. Jenkins and J. Serrin. The Dirichlet problem for the minimal surface equation in higher dimensions. J. Reine Angew. Math. 229 (1968), pp. 170–187. DOI: 10.1515/crll.1968.229.170.
- [70] H.-Y. Jian and X.-J. Wang. Continuity estimates for the Monge-Ampère equation. SIAM J. Math. Anal. 39.2 (2007), pp. 608–626.
 DOI: 10.1137/060669036.
- [71] F. John. Extremum problems with inequalities as subsidiary conditions. In: Studies and Essays Presented to R. Courant on his 60th Birthday, January 8, 1948. Interscience Publishers, New York, 1948, pp. 187–204.
- [72] K. Jörgens. Über die Lösungen der Differentialgleichung $rt s^2 = 1$. Math. Ann. 127 (1954), pp. 130–134. DOI: 10.1007/BF01361114.
- [73] N. Korevaar. An easy proof of the interior gradient bound for solutions to the prescribed mean curvature equation. In: Nonlinear functional analysis and its applications, Part 2 (Berkeley, Calif., 1983).
 Vol. 45. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 1986, pp. 81–89. DOI: 10.1090/pspum/045.2/843597.
- [74] N. V. Krylov. Boundedly inhomogeneous elliptic and parabolic equations. Izv. Akad. Nauk SSSR Ser. Mat. 46.3 (1982), pp. 487–523, 670.
- [75] N. V. Krylov. Boundedly inhomogeneous elliptic and parabolic equations in a domain. *Izv. Akad. Nauk SSSR Ser. Mat.* 47.1 (1983), pp. 75–108.
- [76] N. V. Krylov. Degenerate nonlinear elliptic equations. Mat. Sb. (N.S.) 120(162).3 (1983), pp. 311–330, 448.
- [77] N. V. Krylov. Smoothness of the payoff function for a controllable diffusion process in a domain. *Izv. Akad. Nauk SSSR Ser. Mat.* 53.1 (1989), pp. 66–96. DOI: 10.1070/IM1990v034n01ABEH000603.
- [78] N. V. Krylov and M. V. Safonov. An estimate for the probability of a diffusion process hitting a set of positive measure. *Dokl. Akad. Nauk* SSSR 245.1 (1979), pp. 18–20.
- [79] N. V. Krylov and M. V. Safonov. A property of the solutions of parabolic equations with measurable coefficients. *Izv. Akad. Nauk SSSR Ser. Mat.* 44.1 (1980), pp. 161–175, 239.
- [80] O. A. Ladyzhenskaya and N. N. Ural' tseva. Local estimates for gradients of solutions of non-uniformly elliptic and parabolic equations. *Comm. Pure Appl. Math.* 23 (1970), pp. 677–703. DOI: 10.1002/cpa.3160230409.
- [81] G. P. Leonardi and G. E. Comi. The prescribed mean curvature measure equation in non-parametric form. 2023. arXiv: 2302.10592 [math.AP].
- [82] H. Li, X. Li, and S. Zhao. Hessian quotient equations on exterior domains. 2020. arXiv: 2004.06908 [math.AP].
- [83] Y. Li. Strominger-Yau-Zaslow conjecture for Calabi-Yau hypersurfaces in the Fermat family. Acta Math. 229.1 (2022), pp. 1–53.

- [84] Y. Li. Metric SYZ conjecture and non-Archimedean geometry. Duke Math. J. 172.17 (2023), pp. 3227–3255. DOI: 10.1215/00127094-2022-0099.
- [85] Y. Li, L. Nguyen, and J. Xiong. Regularity of viscosity solutions of the σ_k -Loewner-Nirenberg problem. *Proc. Lond. Math. Soc. (3)* 127.1 (2023), pp. 1–34. DOI: 10.1112/plms.12536.
- [86] Y. Lian, L. Wang, and K. Zhang. Pointwise Regularity for Fully Nonlinear Elliptic Equations in General Forms. 2020. arXiv: 2012.00324 [math.AP].
- [87] Y. Lian and K. Zhang. Boundary pointwise C^{1,α} and C^{2,α} regularity for fully nonlinear elliptic equations. J. Differential Equations 269.2 (2020), pp. 1172–1191. DOI: 10.1016/j.jde.2020.01.006.
- [88] Y. Lian et al. Boundary Hölder regularity for elliptic equations. J. Math. Pures Appl. (9) 143 (2020), pp. 311–333. DOI: 10.1016/j.matpur.2020.09.012.
- [89] S. Lu. Interior C^2 estimate for Hessian quotient equation in dimension three. 2023. arXiv: 2311.05835 [math.AP].
- [90] S. Lu. On the Dirichlet problem for Lagrangian phase equation with critical and supercritical phase. *Discrete Contin. Dyn. Syst.* 43.7 (2023), pp. 2561–2575. DOI: 10.3934/dcds.2023020.
- [91] S. Lu. Interior C^2 estimate for Hessian quotient equation in general dimension. 2024. arXiv: 2401.12229 [math.AP].
- [92] M. Miranda. Dirichlet problem with L¹ data for the non-homogeneous minimal surface equation. *Indiana Univ. Math. J.* 24 (1974/75), pp. 227– 241. DOI: 10.1512/iumj.1974.24.24020.
- [93] C. Mooney. Strict 2-convexity of convex solutions to the quadratic Hessian equation. Proc. Amer. Math. Soc. 149.6 (2021), pp. 2473– 2477. DOI: 10.1090/proc/15454.
- [94] C. Mooney and O. Savin. Non C¹ solutions to the special Lagrangian equation. 2023. arXiv: 2303.14282 [math.AP].
- [95] N. Nadirashvili and S. Vlăduţ. Singular solution to special Lagrangian equations. Ann. Inst. H. Poincaré C Anal. Non Linéaire 27.5 (2010), pp. 1179–1188. DOI: 10.1016/j.anihpc.2010.05.001.
- [96] A. V. Pogorelov. The regularity of the generalized solutions of the equation $\det(\partial^2 u/\partial x^i \partial x^j) = \varphi(x^1, x^2, \dots, x^n) > 0$. Dokl. Akad. Nauk SSSR 200 (1971), pp. 534–537.
- [97] A. V. y. Pogorelov. The Minkowski multidimensional problem. Scripta Series in Mathematics. Translated from the Russian by Vladimir Oliker, Introduction by Louis Nirenberg. V. H. Winston & Sons, Washington, DC; Halsted Press [John Wiley & Sons], New York-Toronto-London, 1978, p. 106.
- [98] I. H. Sabitov. Regularity of convex domains with a metric that is regular on Hölder classes. Sibirsk. Mat. Z. 17.4 (1976), pp. 907–915.
- [99] O. Savin. Pointwise $C^{2,\alpha}$ estimates at the boundary for the Monge-Ampère equation. J. Amer. Math. Soc. 26.1 (2013), pp. 63–99. DOI: 10.1090/S0894-0347-2012-00747-4.
- [100] O. Savin. Small perturbation solutions for elliptic equations. Comm. Partial Differential Equations 32.4-6 (2007), pp. 557–578. DOI: 10.1080/03605300500394405.

- [101] O. Savin. Regularity of flat level sets in phase transitions. Ann. of Math. (2) 169.1 (2009), pp. 41–78. DOI: 10.4007/annals.2009.169.41.
- [102] F. Schulz. Über die Differentialgleichung $rt-s^2 = f$ und das Weylsche Einbettungsproblem. *Math. Z.* 179.1 (1982), pp. 1–10. DOI: 10.1007/BF01173911.
- [103] J. Serrin. The problem of Dirichlet for quasilinear elliptic differential equations with many independent variables. *Philos. Trans. Roy. Soc. London Ser. A* 264 (1969), pp. 413–496. DOI: 10.1098/rsta.1969.0033.
- [104] R. Shankar and Y. Yuan. Hessian estimate for semiconvex solutions to the sigma-2 equation. *Calc. Var. Partial Differential Equations* 59.1 (2020), Paper No. 30, 12. DOI: 10.1007/s00526-019-1690-1.
- [105] R. Shankar and Y. Yuan. Regularity for almost convex viscosity solutions of the sigma-2 equation. J. Math. Study 54.2 (2021), pp. 164– 170. DOI: 10.4208/jms.v54n2.21.03.
- [106] R. Shankar and Y. Yuan. Hessian estimates for the sigma-2 equation in dimension four. 2023. arXiv: 2305.12587 [math.AP].
- [107] L. Silvestre and B. Sirakov. Boundary regularity for viscosity solutions of fully nonlinear elliptic equations. *Comm. Partial Differential Equations* 39.9 (2014), pp. 1694–1717. DOI: 10.1080/03605302.2013.842249.
- [108] G. Székelyhidi. Uniqueness of some Calabi-Yau metrics on Cⁿ. Geom. Funct. Anal. 30.4 (2020), pp. 1152–1182. DOI: 10.1007/s00039-020-00543-3.
- [109] G. Tian, Q. Wang, and C.-J. Xu. Local solvability of the k-Hessian equations. Sci. China Math. 59.9 (2016), pp. 1753–1768. DOI: 10.1007/s11425-016-5135-4.
- [110] N. S. Trudinger. On the analyticity of generalized minimal surfaces. Bull. Austral. Math. Soc. 5 (1971), pp. 315–320. DOI: 10.1017/S0004972700047262.
- [111] N. S. Trudinger. A new proof of the interior gradient bound for the minimal surface equation in n dimensions. Proc. Nat. Acad. Sci. U.S.A. 69 (1972), pp. 821–823.
- [112] N. S. Trudinger. Regularity of solutions of fully nonlinear elliptic equations. *Boll. Un. Mat. Ital. A (6)* 3.3 (1984), pp. 421–430.
- [113] N. S. Trudinger. On the Dirichlet problem for Hessian equations. Acta Math. 175.2 (1995), pp. 151–164. DOI: 10.1007/BF02393303.
- [114] N. S. Trudinger. Weak solutions of Hessian equations. Comm. Partial Differential Equations 22.7-8 (1997), pp. 1251–1261. DOI: 10.1080/03605309708821299.
- [115] N. S. Trudinger and J. I. E. Urbas. On second derivative estimates for equations of Monge-Ampère type. Bull. Austral. Math. Soc. 30.3 (1984), pp. 321–334. DOI: 10.1017/S0004972700002069.
- [116] N. S. Trudinger and X.-J. Wang. Hessian measures. I. Topol. Methods Nonlinear Anal. 10.2 (1997). Dedicated to Olga Ladyzhenskaya, pp. 225–239. DOI: 10.12775/TMNA.1997.030.
- [117] N. S. Trudinger and X.-J. Wang. Hessian measures. II. Ann. of Math.
 (2) 150.2 (1999), pp. 579–604. DOI: 10.2307/121089.
- [118] N. S. Trudinger and X.-J. Wang. Hessian measures. III. J. Funct. Anal. 193.1 (2002), pp. 1–23. DOI: 10.1006/jfan.2001.3925.

- [119] N. S. Trudinger and X.-J. Wang. Boundary regularity for the Monge-Ampère and affine maximal surface equations. Ann. of Math. (2) 167.3 (2008), pp. 993–1028. DOI: 10.4007/annals.2008.167.993.
- [120] N. S. Trudinger and X.-J. Wang. The Monge-Ampère equation and its geometric applications. In: *Handbook of geometric analysis. No. 1.* Vol. 7. Adv. Lect. Math. (ALM). Int. Press, Somerville, MA, 2008, pp. 467–524.
- J. Urbas. Some interior regularity results for solutions of Hessian equations. Calc. Var. Partial Differential Equations 11.1 (2000), pp. 1– 31. DOI: 10.1007/s005260050001.
- [122] J. Urbas. An interior second derivative bound for solutions of Hessian equations. *Calc. Var. Partial Differential Equations* 12.4 (2001), pp. 417–431. DOI: 10.1007/PL00009920.
- [123] J. I. E. Urbas. Regularity of generalized solutions of Monge-Ampère equations. *Math. Z.* 197.3 (1988), pp. 365–393. DOI: 10.1007/BF01418336.
- [124] J. I. E. Urbas. On the existence of nonclassical solutions for two classes of fully nonlinear elliptic equations. *Indiana Univ. Math. J.* 39.2 (1990), pp. 355–382. DOI: 10.1512/iumj.1990.39.39020.
- C. Wang and J. Bao. Liouville property and existence of entire solutions of Hessian equations. *Nonlinear Anal.* 223 (2022), Paper No. 113020, 18. DOI: 10.1016/j.na.2022.113020.
- [126] D. Wang and Y. Yuan. Singular solutions to special Lagrangian equations with subcritical phases and minimal surface systems. Amer. J. Math. 135.5 (2013), pp. 1157–1177. DOI: 10.1353/ajm.2013.0043.
- [127] D. Wang and Y. Yuan. Hessian estimates for special Lagrangian equations with critical and supercritical phases in general dimensions. *Amer. J. Math.* 136.2 (2014), pp. 481–499. DOI: 10.1353/ajm.2014.0009.
- [128] L. Wang. On the regularity theory of fully nonlinear parabolic equations. II. Comm. Pure Appl. Math. 45.2 (1992), pp. 141–178. DOI: 10.1002/cpa.3160450202.
- [129] L. Wang. Compactness methods for certain degenerate elliptic equations. J. Differential Equations 107.2 (1994), pp. 341–350. DOI: 10.1006/jdeq.1994.1016.
- [130] X.-J. Wang. Remarks on the regularity of Monge-Ampère equations. In: Proceedings of the International Conference on Nonlinear PDE. Ed. by G. C. Dong and F. H. Lin. 257–263. Beijing: Scientific Press, 1992.
- [131] X. J. Wang. A class of fully nonlinear elliptic equations and related functionals. *Indiana Univ. Math. J.* 43.1 (1994), pp. 25–54. DOI: 10.1512/iumj.1994.43.43002.
- [132] X.-J. Wang. Some counterexamples to the regularity of Monge-Ampère equations. *Proc. Amer. Math. Soc.* 123.3 (1995), pp. 841–845. DOI: 10.2307/2160809.
- [133] X.-J. Wang. Interior gradient estimates for mean curvature equations. Math. Z. 228.1 (1998), pp. 73–81. DOI: 10.1007/PL00004604.
REFERENCES

- [134] X.-J. Wang. The k-Hessian equation. In: Geometric analysis and PDEs. Vol. 1977. Lecture Notes in Math. Springer, Dordrecht, 2009, pp. 177–252. DOI: 10.1007/978-3-642-01674-5_5.
- [135] Y. Wang. Small perturbation solutions for parabolic equations. *Indi*ana Univ. Math. J. 62.2 (2013), pp. 671–697. DOI: 10.1512/iumj.2013.62.4961.
- [136] M. Warren. Special Lagrangian equations. Thesis (Ph.D.)–University of Washington. ProQuest LLC, Ann Arbor, MI, 2008, p. 103.
- M. Warren and Y. Yuan. Explicit gradient estimates for minimal Lagrangian surfaces of dimension two. *Math. Z.* 262.4 (2009), pp. 867– 879. DOI: 10.1007/s00209-008-0403-9.
- M. Warren and Y. Yuan. Hessian estimates for the sigma-2 equation in dimension 3. Comm. Pure Appl. Math. 62.3 (2009), pp. 305–321.
 DOI: 10.1002/cpa.20251.
- [139] M. Warren and Y. Yuan. Hessian and gradient estimates for three dimensional special Lagrangian equations with large phase. Amer. J. Math. 132.3 (2010), pp. 751–770. DOI: 10.1353/ajm.0.0115.
- [140] Y. Xu. Interior estimates for solutions of the quadratic Hessian equations in dimension three. J. Math. Anal. Appl. 489.2 (2020), pp. 124179, 10. DOI: 10.1016/j.jmaa.2020.124179.
- H. Yu. Small perturbation solutions for nonlocal elliptic equations. Comm. Partial Differential Equations 42.1 (2017), pp. 142–158. DOI: 10.1080/03605302.2016.1252395.
- [142] Y. Yuan. A priori estimates for solutions of fully nonlinear special Lagrangian equations. Ann. Inst. H. Poincaré C Anal. Non Linéaire 18.2 (2001), pp. 261–270. DOI: 10.1016/S0294-1449(00)00065-2.
- [143] Y. Yuan. A Bernstein problem for special Lagrangian equations. Invent. Math. 150.1 (2002), pp. 117–125. DOI: 10.1007/s00222-002-0232-0.
- [144] Y. Yuan. Global solutions to special Lagrangian equations. Proc. Amer. Math. Soc. 134.5 (2006), pp. 1355–1358. DOI: 10.1090/S0002-9939-05-08081-0.
- [145] Y. Yuan. Special Lagrangian equations. In: Geometric analysis—in honor of Gang Tian's 60th birthday. Vol. 333. Progr. Math. Birkhäuser/Springer, Cham, [2020] ©2020, pp. 521–536.
- [146] Q. Zhang. Boundary regularity for viscosity solutions of nonlinear singular elliptic equations. *Calc. Var. Partial Differential Equations* 62.3 (2023), Paper No. 87, 21. DOI: 10.1007/s00526-022-02427-w.
- [147] X. Zhou. Hessian estimates for Lagrangian mean curvature equation with sharp Lipschitz phase. 2023. arXiv: 2311.13867 [math.AP].
- [148] X. Zhou. Notes on generalized special Lagrangian equation. 2023. arXiv: 2311.14260 [math.AP].
- [149] W. K. Ziemer. The nonhomogeneous minimal surface equation involving a measure. *Pacific J. Math.* 167.1 (1995), pp. 183–200.
- [150] W. P. Ziemer. Weakly differentiable functions. Vol. 120. Graduate Texts in Mathematics. Sobolev spaces and functions of bounded variation. Springer-Verlag, New York, 1989, pp. xvi+308. DOI: 10.1007/978-1-4612-1015-3.

REFERENCES

Departamento de Análisis Matemático, Instituto de Matemáticas IMAG, Universidad de Granada

Email address: lianyuanyuan.hthk@gmail.com; yuanyuanlian@correo.ugr.es; MR ID:1378049; ORCID:0000-0002-2276-3063

Departamento de Geometría y Topología, Instituto de Matemáticas IMAG, Universidad de Granada

Email address: zhangkaizfz@gmail.com; zhangkai@ugr.es; MR ID:1098004; ORCID:0000-0002-1896-3206