

# ERGODICITY OF SKEW-PRODUCTS OVER TYPICAL IETS

F. ARGENTIERI, P. BERK, F. TRUJILLO

**ABSTRACT.** We prove ergodicity of a class of infinite measure preserving systems, called skew-products. More precisely, we consider systems of the form

$$T_f : \begin{array}{ccc} [0, 1) \times \mathbb{R} & \rightarrow & [0, 1) \times \mathbb{R} \\ (x, t) & \mapsto & (T(x), t + f(x)) \end{array} ,$$

where  $T$  is an interval exchange transformation and  $f$  is a piece-wise constant function with a finite number of discontinuities. We show that such system is ergodic with respect to  $\text{Leb}_{[0,1) \times \mathbb{R}}$  for a typical choice of parameters of  $T$  and  $f$ . This generalizes recent results Chaika and Robertson [4] concerning an exceptional class of interval exchange transformations.

## 1. INTRODUCTION

The interval exchange transformations, or IETs, for short, appear naturally as the Poincaré return maps to a transversal section for many classical surface flows such as translation flows or locally Hamiltonian flows. As such, during the last century, they attracted a lot of attention from a large number of mathematicians. In this article, we study infinite measure preserving systems called *skew-products* built as extensions of said transformations. We focus on ergodic properties of aforementioned systems.

To be more precise, given  $m \geq 1$  and  $M > 0$ , define

$$P_{m,M} = \{(p, q) \in \mathbb{R}^{m+1} \times \mathbb{R}^{m+1} \mid \langle p, q \rangle = 0; |p|_1 = 1; |q|_\infty \leq M\}.$$

We denote by  $C_{m,M}$  the space of mean zero step functions  $f : [0, 1) \rightarrow \mathbb{R}$  of the form

$$f = q_1 \chi_{[0, p_1)} + \dots + q_{m+1} \chi_{[p_1 + \dots + p_{m+1}, 1)},$$

for some  $(p, q) \in P_{m,M}$ . For  $f$  as above, we call

$$\sigma_i(f) := q_{i+1} - q_i \tag{1}$$

the  $i$ -th jump of  $f$ , for any  $1 \leq i \leq m$ , and denote the set of all jumps of  $f$  by  $\sigma(f)$ .

We endow  $C_{m,M}$  with the Lebesgue measure and the  $l_\infty$  metric inherited from  $P_{m,M}$ . Given  $f \in C_{m,M}$  and an IET  $T$ , we associate the *skew-product*  $T_f$  given by

$$T_f : \begin{array}{ccc} [0, 1) \times \mathbb{R} & \rightarrow & [0, 1) \times \mathbb{R} \\ (x, t) & \mapsto & (T(x), t + f(x)) \end{array} . \tag{2}$$

We refer to  $T$  as the *base* of  $T_f$  and to  $f$  as the *cocycle* associated with  $T_f$ .

We also consider the space  $S_0 \times \Lambda^{\mathcal{A}}$  of IETs of  $d \geq 2$  intervals indexed by an alphabet  $\mathcal{A}$  with  $d$  elements. Here  $S_0$  denotes the set of irreducible permutations, that is, permutations that do not divide an interval into two or more subintervals which are being exchanged independently, while  $\Lambda^{\mathcal{A}}$  stands for the unit simplex of real positive vectors indexed by  $\mathcal{A}$ , namely,

$$\Lambda^{\mathcal{A}} = \{v \in \mathbb{R}_+^{\mathcal{A}} \mid |v|_1 = 1\}.$$

On the space  $S_0 \times \Lambda^{\mathcal{A}}$ , we consider the product of counting and Lebesgue measure, and whenever we are speaking about a "typical" or "almost every" IET, we mean almost every with respect to this measure.

The following is the main result of our article.

**Theorem 1.1.** *Let  $m \geq 2$  and  $M > 0$ . For a.e. irreducible IET  $T$  on  $d \geq 2$  intervals the associated cocycle  $T_f$  is ergodic, with respect to  $\text{Leb}_{[0,1) \times \mathbb{R}}$ , for a.e.  $f \in C_{m,M}$ .*

**Remark 1.1.** *The set of IETs considered in Theorem 1.1 are the ones satisfying the conclusions of Theorem 1.2. The set of cocycles for which the result holds depends on  $T$ .*

Before Chaika and Robertson, most of the existing results concerned rotations as the base in the definition of the skew product. Here it is worth to recall that rotations can be seen as IETs of two intervals. The ergodicity of skew-products with rotations in their base was shown for many classes of cocycles, we mention here Oren [13], Hellekalek and Larcher [10], Pask [14], as well as Conze and Piękniewska [6]. The IETs of higher number of intervals were much less explored. Conze and Frączek [5] proved ergodicity of skew products with self-similar IETs in the base (which form a countable set of IETs) and piecewise constant functions, constant over exchanged intervals. Surprisingly, Frączek and Ulcigrai [8], during their study of Ehrenfest wind-tree model, found examples of skew products over IETs with piece-wise constant cocycles, again constant over exchanged intervals, which are not ergodic.

Although much of our arguments seem similar to the ones used in [4], the different approach that we exhibit allows us to obtain a more general result. The biggest similarity is that we also exploit the jumps to obtain ergodicity (although in a slightly different manner, since we do not utilize directly the essential value criterion).

We would like now to highlight major differences. There are typically two obstacles that occur when trying to prove ergodicity of a skew product. First is the required control over close return times (rigid times) of the base transformation. The second is difficulty to get bounded Birkhoff sums of the cocycle along the close return times. In [4], the authors solved both problem by introducing the so-called *quantitative version of Atkinson's Lemma* and greater control over the return times for the bounded type IETs than the for typical IETs. We, on the other hand, utilize the properties of the *Konsevich-Zorich cocycle* (see Subsection 2.3). More precisely we used its ergodicity and that fact that the first Lyapunov exponent of said cocycle is strictly bigger than the second one (it exhibits a spectral gap). Then the Rokhlin towers obtained by this algorithm can be balanced in height and width, while their heights grow much faster than the Birkhoff sums of the cocycle. This forces many of the Birkhoff sums to be bounded. This approach allows us to provide sufficient conditions on the pairs  $(T, f)$  in terms of recurrence properties of  $T$  and the Birkhoff sums  $S_n f$  for Theorem 1.1 to hold (see Properties (1)-(4) below). We then check that for a.e. irreducible IET, there exists a full-measure set of cocycles for which these conditions are satisfied.

**Theorem 1.2.** *For a.e. irreducible IET  $T$  on  $d \geq 2$  intervals there exist  $C, \sigma > 0$  such that the following holds.*

*Given  $m \geq 2$  and  $M > 0$  there exists a full-measure set  $\mathcal{F} \subseteq C_{m,M}$  such that, for any  $f \in C_{m,M}$ , for any positive measure set  $E \subseteq [0, 1)$ , any  $D > mM$  and any  $N > 0$ , there exist  $x \in E$  and  $n > N$  satisfying:*

- (1)  $x, T^n(x) \in E$ .
- (2)  $|S_n f(x)| < D$ .
- (3) *The set  $\{T^i(x)\}_{i=0}^{n-1}$  is  $\frac{C}{n}$ -dense in  $[0, 1)$ .*
- (4) *Either  $[x - \frac{\sigma}{n}, x]$  or  $[x, x + \frac{\sigma}{n}]$  is a continuity interval for  $T^n$ .*

**Remark 1.2.** *It follows from the proof of Theorem 1.2 that the constants  $C, \sigma > 0$  in the statement can be taken uniform for a full-measure set of irreducible IETs on  $d$  intervals.*

Let us point out that, for any IET  $T$  ergodic w.r.t. to  $\text{Leb}_{[0,1]}$ , and any  $f \in C_{m,M}$ , by the recurrence of  $T_f$  it is easy to find times  $n$  where conditions (1) and (2) are verified. Also, by exploiting the ergodicity of the (accelerated) Rauzy-Veech renormalization, is not difficult to find, for a.e.  $T$  and for some constants  $C, \sigma > 0$  not depending on  $T$ , times  $n$  where Conditions (3) and (4) hold. However, these times do not necessarily coincide. The importance of Theorem 1.2 is that it guarantees that these times can be made to coincide, infinitely many times, under generic conditions on  $T$  and  $f$ .

**Remark 1.3.** *The conclusions of Theorem 1.2 are satisfied by the set of linearly recurrent IETs introduced in [4] (as the last two conditions are trivially verified, for any  $n \geq 1$ , for some constants  $C, \sigma$  depending only on  $T$ ). In particular, Theorem 1.1 also applies for this class of IETs.*

Theorem 1.2 is proven in the last section of this work and relies heavily on recurrence properties of the accelerated Rauzy-Veech renormalization as well as in well-known results due to Zorich [20] concerning the deviation of certain ergodic averages over IETs (see Theorems 4.1, 4.2).

Before describing more in detail the proof strategy of Theorem 1.1, let us point out that, as a direct corollary of this theorem, we obtain the following.

**Theorem 1.3.** *Let  $m \geq 2$ ,  $M > 0$  and  $d \geq 2$ . For a.e.  $(T, f) \in S_0 \times \Lambda^A \times C_{m,M}$  the associated skew product  $T_f$  is ergodic with respect to  $\text{Leb}_{[0,1] \times \mathbb{R}}$ .*

*In particular, for a.e.  $f \in C_{m,M}$  there exists a full measure set of IETs for which the associated skew product  $T_f$  is ergodic with respect to  $\text{Leb}_{[0,1] \times \mathbb{R}}$ .*

*Proof.* In view of Theorem 1.1, it is enough to show that the set

$$Z := \{(T, f) \in S_0 \times \Lambda^A \times C_{m,M} \mid T_f \text{ is ergodic}\}$$

is measurable.

For any  $N \in \mathbb{N}$ , let  $\tilde{T}_f^N : [0, 1] \times [-N, N] \rightarrow [0, 1] \times [-N, N]$  denote the first return map of  $T_f$  to  $[0, 1] \times [-N, N]$ . Recall that, due to a classical result by Atkinson [2], if  $T$  is ergodic then the associated skew product  $T_f$  is *recurrent* (see Appendix A for a precise definition). In particular the first return transformation  $\tilde{T}_f^N$  is well-defined on a full-measure subset of  $[0, 1] \times [-N, N]$ .

Note that  $T_f$  is ergodic if and only if  $\tilde{T}_f^N$  is ergodic, for every  $N \in \mathbb{N}$ . Indeed, an induced map of an ergodic transformation is also ergodic. On the other hand, if there was a  $T_f$ -invariant subset of positive measure whose complement has positive measure, then, for  $N \in \mathbb{N}$  sufficiently large, the set  $[0, 1] \times [-N, N]$  would contain a  $\tilde{T}_f^N$ -invariant subset of positive measure whose complement also has positive measure.

Thus,

$$Z = \bigcap_{N=1}^{\infty} Z_N, \quad \text{where } Z_N := \{(T, f) \in S_0 \times \Lambda^A \times C_{m,M} \mid \tilde{T}_f^N \text{ is ergodic}\}.$$

It is not difficult to show that the assignment  $(T, f) \mapsto \tilde{T}_f^N$  is measurable (see Lemma A.4 in Appendix A). Due to the fact that the set of ergodic transformations of any standard probability space is measurable (see [9]), it follows that the set  $Z_N$  is measurable for every  $N \in \mathbb{N}$ . Therefore  $Z$  is measurable, which concludes the proof of the theorem.  $\square$

**1.1. Scheme of the proof of Theorem 1.1.** We suppose  $T$  is an IET ergodic with respect to the Lebesgue measure and satisfying the conclusions of Theorem 1.2.

Suppose, for the sake of contradiction, that for a positive measure set of functions  $f$  in  $C_{m,M}$ , the associated skew-products  $T_f$  are not ergodic. Call this set  $W \subseteq C_{m,M}$ . Since a typical function in  $C_{m,M}$  has at least two rationally independent jumps (see (1)), we can assume WLOG that for any  $f \in W$  the group generated by  $\sigma(f)$  is dense in  $\mathbb{R}$ .

For any  $f$  in an appropriate subset  $V \subseteq W$ , we will define a  $T_f$ -invariant measure  $\mu_f$ , depending measurably on  $f$ , that is not a multiple of  $\text{Leb}_{[0,1] \times \mathbb{R}}$ , nor singular to it.

We then decompose the measure  $\mu_f$  along the fibers  $\{x\} \times \mathbb{R}$  as a family of measures on  $\mathbb{R}$  which we denote  $\{\mu_{f,x}\}_{x \in \mathbb{R}}$ . Since  $\mu_f$  is not a multiple of  $\text{Leb}_{[0,1] \times \mathbb{R}}$ , it is possible to show that for a.e.  $x \in \mathbb{R}$ , the measure  $\mu_{f,x}$  is not a multiple of  $\text{Leb}_{\mathbb{R}}$ . So we may assume that the set  $U$  of  $(f, x) \in V \times [0, 1]$  for which  $\mu_{f,x}$  is not a multiple of  $\text{Leb}_{\mathbb{R}}$  has full measure.

We will derive a contradiction by showing the existence of  $(f, x) \in U$  such that  $\mu_{f,x}$  is a multiple of the Lebesgue measure.

To show this, we first consider cutoffs  $\mu_{f,x,D}$  of the measures  $\mu_{f,x}$  to compact intervals of the form  $[-D, D]$ , and define an appropriate continuity set  $B$  for the family of maps  $(f, x) \mapsto \mu_{f,x,D}$ , for any  $D \in \mathbb{Q}_{>0}$ . We then fix  $(f, x) \in B$  and construct sequences  $(f_{i,k}, x_k^{\pm})$  in  $B$ , converging to  $(f, x)$ , such that for any open interval  $J$  such that  $J, J + \sigma_i(f) \subseteq [-D, D]$ , we have

$$\mu_{f_{i,k}, x_k^+, D}(J) - \mu_{f_{i,k}, x_k^-, D}(J + \sigma_i(f)) \rightarrow 0.$$

Using the continuity of these measures on the set  $B$ , we conclude that  $\mu_{f,x}$  is invariant by the translation by  $\sigma_i(f)$ . Since the set of jumps of  $f$  generates a dense subgroup in  $\mathbb{R}$ , this implies that  $\mu_{f,x}$  is a multiple of the Lebesgue measure, which contradicts our assumptions on the set  $U$ .

**Remark 1.4.** *The proof of the existence of the measures  $\mu_f$  described above follows many well-known techniques but is rather technical. For this reason, we provide a detailed proof of this in Appendix A.*

**1.2. Outline of the article.** First, in Section 2, we introduce the notations used throughout the article as well as the main tool in the proof of Theorem 1.2, namely, Rauzy-Veech renormalization.

In Section 3, assuming Theorem 1.2 which describes the set of pairs of IETs and cocycles to be considered, we follow the scheme described in Section 1.1 to prove our main result, namely, Theorem 1.1.

In Section 4, we prove Theorem 1.2. Finally, in Appendix A, we provide some of the technical details required in the proof of Theorem 1.1.

**Acknowledgments:** We are very grateful to Corinna Ulcigrai for all the fruitful discussions. The first author was supported by the SNSF Grant 188535. The second and third authors acknowledge the support of the Swiss National Science Foundation through Grant 200021\_188617/1. The second author was supported by the NCN Grant 2022/45/B/ST1/00179. The third author was partially supported by the UZH Postdoc Grant, grant no. FK-23-133 and the Spanish State Research Agency, through the Severo Ochoa and María de Maeztu Program for Centers and Units of Excellence in R&D (CEX2020-001084-M).

## 2. NOTATIONS

**2.1. Interval exchange transformations.** We now present main notions and facts used in this article. Let  $I = [0, |I|)$  be an interval of length  $|I|$ , where  $|\cdot|$  denotes the Lebesgue measure of a set. Let also  $\mathcal{A}$  be an alphabet of  $d \geq 2$  elements. We say that  $T$  is an *interval exchange transformation (IET)* of  $d$  elements if there exists a finite partition  $\{I_\alpha\}_{\alpha \in \mathcal{A}}$  of  $I$  into  $d$  subintervals such that  $T : I \rightarrow I$  is a bijection and  $T|_{I_\alpha}$  is a translation for every  $\alpha \in \mathcal{A}$ . In particular,  $T$  preserves Lebesgue measure. Roughly speaking,  $T$  is a permutation of intervals. Since we will be considering  $T$  as a measure-preserving system, we will sometimes, when it is more convenient consider  $T$  as a Lebesgue measure-preserving automorphism of  $[0, 1]$ . Then  $T$  is described by two parameters. First is a permutation  $\pi = (\pi_0, \pi_1)$ , where  $\pi_0, \pi_1 : \mathcal{A} \rightarrow \{1, \dots, d\}$  are bijections and  $\pi_0(\alpha)$  indicates the position of an interval  $I_\alpha$  before the exchange, while  $\pi_1(\alpha)$  the position of  $I_\alpha$  after the exchange. The second parameter is the length vector  $\lambda \in \mathbb{R}_+^{\mathcal{A}}$ , where  $\lambda_\alpha = |I_\alpha|$ . We denote  $T = (\pi, \lambda)$ .

We will often assume some restrictions on  $\pi$  and  $\lambda$ . First, we will only consider  $\pi$  to be irreducible, that is there is no  $k \in \{1, \dots, d-1\}$ , such that  $\pi_1 \circ \pi_0^{-1} \{1, \dots, k\} = \{1, \dots, k\}$ . We denote the set of such permutations as  $S_0^{\mathcal{A}}$ . Moreover, we frequently assume that  $|I| = 1$  and then  $\lambda \in \Lambda^{\mathcal{A}}$ , where  $\Lambda \subseteq \mathbb{R}_+^{\mathcal{A}}$  is a positive unit simplex.

**2.2. Rauzy-Veech induction.** On the space  $S_0^{\mathcal{A}} \times \mathbb{R}_{>0}^{\mathcal{A}}$  we consider an operator  $\tilde{R}$  called *Rauzy-Veech induction*, defined as  $\tilde{R}(\pi, \lambda) = (\pi^{(1)}, \lambda^{(1)})$ , where  $(\pi^{(1)}, \lambda^{(1)})$  is the first return map of  $(\pi, \lambda)$  to the interval  $I^{(1)} := [0, |\lambda| - \min\{\lambda_{\pi_0^{-1}(d)}, \lambda_{\pi_1^{-1}(d)}\})$ , where  $|\lambda| = \sum_{\alpha \in \mathcal{A}} \lambda_\alpha$ . If  $\lambda_{\pi_0^{-1}(d)} > \lambda_{\pi_1^{-1}(d)}$  we say that  $\tilde{R}$  is of "top" type and we say that it is of "bottom" type if  $\lambda_{\pi_0^{-1}(d)} < \lambda_{\pi_1^{-1}(d)}$ .

The map  $\tilde{R}(\pi, \lambda)$  is properly defined as an interval exchange transformation of  $d$  intervals if and only if  $\lambda_{\pi_0^{-1}(d)} \neq \lambda_{\pi_1^{-1}(d)}$ . Keane [11] gave an equivalent condition on  $(\pi, \lambda)$ , for the iterations of Rauzy-Veech induction to be defined indefinitely. More precisely, we say that IET  $T$  satisfies *Keane's condition* if for every two discontinuities  $a$  and  $b$  of  $T$  equality  $T^n(a) = b$  for some  $n \in \mathbb{N}$  implies  $n = 1$ ,  $a = T^{-1}(0)$  and  $b = 0$ . In particular, if the vector  $\lambda$  is rationally independent, that is for every choice of  $c_\alpha \in \mathbb{Z}$ ,  $\alpha \in \mathcal{A}$  we have

$$\sum_{\alpha \in \mathcal{A}} c_\alpha \lambda_\alpha = 0 \Rightarrow c_\alpha = 0 \text{ for every } \alpha \in \mathcal{A},$$

then  $(\pi, \lambda)$  satisfies Keane's condition. When it is well-defined, we denote

$$(\pi^n, \lambda^n) := \tilde{R}^n(\pi, \lambda)$$

for every  $n \in \mathbb{N}$  and we denote its domain by  $I^n$  and the exchanged intervals by  $\{I_\alpha^n\}_{\alpha \in \mathcal{A}}$ .

Note that  $\lambda^1 = A(\pi, \lambda)\lambda$ , where a matrix  $A^{(1)}(\pi, \lambda)$  is defined in the following way

$$A_{\alpha\beta}(\pi, \lambda) = \begin{cases} 1 & \text{if } \alpha = \beta; \\ -1 & \text{if } \alpha = w \text{ and } \beta = l; \\ 0 & \text{otherwise.} \end{cases}$$

Denoting  $A^{(0)}(\pi, \lambda) = \text{Id}$ , we define, inductively,

$$A^{(n)}(\pi, \lambda) = A(\pi^{n-1}, \lambda^{n-1})A^{(n-1)}(\pi, \lambda),$$

for any  $n \geq 1$ . Then

$$\lambda^n = A^{(n)}(\pi, \lambda)\lambda. \quad (3)$$

We will refer to  $A^{(n)}(\pi, \lambda)$  as *Rauzy-Veech matrices*. Note that for every  $n \in \mathbb{N}$ , the matrix  $(A^{(n)}(\pi, \lambda))^{-1}$  is non-negative. One of the many reasons why Rauzy-Veech matrices are important is the fact, that the number  $A_{\alpha\beta}$  indicates the number of visits of interval  $I_\alpha^n$  to  $I_\beta$  before its first return to  $I^n$  via  $T$ . It also gives that for any  $n \in \mathbb{N}$  the interval  $I$  can be represented as a union of  $d = \#\mathcal{A}$  disjoint Rokhlin towers of intervals as

$$I = \bigsqcup_{\alpha \in \mathcal{A}} \bigsqcup_{i=0}^{q_\alpha^n - 1} T^i I_\alpha^n,$$

where we refer to the numbers  $q_\alpha^n$  as *heights* and to intervals  $I_\alpha^n$  as *bases* of the towers. The numbers  $q_\alpha^n$  are the return times of any  $x \in I_\alpha^n$  to  $I^n$  via  $T$ . One can show that if  $[q_\alpha^n]_{\alpha \in \mathcal{A}}$  is an  $n$ -th vector of heights, then

$$(A^{(n)})^*[1, \dots, 1] = [q_\alpha^n]_{\alpha \in \mathcal{A}},$$

where  $(\cdot)^*$  denotes inverse transpose matrix.

Any invariant subset  $\mathfrak{R} \subseteq S_0^{\mathcal{A}}$  under the (projected) action of  $\tilde{R}$  is called a *Rauzy graph*. Note that the value of  $A^{(1)}$  can be derived only from  $\pi$  and  $\pi^{(1)}$ , not the exact value of  $\lambda$ . Any finite sequence of permutations  $[\pi^1, \dots, \pi^k]$  in  $\mathfrak{R}$  such that for every  $2 \leq i \leq k$  the permutation  $\pi^{i+1}$  is attainable as a permutation in  $\tilde{R}(\pi^i, \lambda^i)$  for some  $\lambda^i \in \mathbb{R}_+^{\mathcal{A}}$ , is called a *Rauzy path*. We have the following useful result, which allows to control the heights of towers, while knowing the last path obtained by Rauzy-Veech induction. We denote the inverse Rauzy-Veech matrix (which is non-negative) associated to  $\gamma$  by  $A_\gamma$ . One can show, by considering the set  $A_\gamma(\mathbb{R}_+^{\mathcal{A}})$ , that any Rauzy path is realizable as a sequence of permutations obtained by consecutive action of  $\tilde{R}$  on some  $(\pi, \lambda) \in \mathfrak{R} \times \mathbb{R}_+^{\mathcal{A}}$ .

**2.3. Normalized induction - renormalization process.** The Rauzy-Veech induction allows to construct Rokhlin towers with very useful properties. However, to use it as a renormalization scheme, we need to consider a *normalized Rauzy-Veech induction*  $\mathcal{R} : \mathfrak{R} \times \Lambda^{\mathcal{A}}$ , where  $\mathfrak{R}$  is a Rauzy graph and

$$\mathcal{R}(\pi, \lambda) := \left( \pi^1, \frac{\lambda^1}{|\lambda^1|} \right).$$

In [12] and [16] Masur and Veech independently proved that there exists a measure  $\mu = \mu_{\mathfrak{R}}$  on  $\mathfrak{R} \times \Lambda^{\mathcal{A}}$ , which is equivalent to the product of counting measure on  $\mathfrak{R}$  and Lebesgue measure on  $\Lambda^{\mathcal{A}}$ , such that  $(\mathcal{R}, \mu_{\mathfrak{R}})$  is an ergodic, recurrent, measure-preserving system. However, the measure  $\mu_{\mathfrak{R}}$  is known to be infinite. For this reason, Zorich in [19] introduced an acceleration  $\mathcal{R}_{KZ}$  of  $\mathcal{R}$  on the set of infinitely renormalizable IETs as

$$\mathcal{R}_{KZ}(\pi, \lambda) := \mathcal{R}^{\kappa(\pi, \lambda)}(\pi, \lambda),$$

where  $\kappa(\pi, \lambda) := \max\{n \in \mathbb{N} \mid \mathcal{R}(\pi, \lambda), \dots, \mathcal{R}^n(\pi, \lambda) \text{ are of the same type}\}$ . We sometimes refer to the operator  $\mathcal{R}_{KZ}$  as the *Zorich acceleration*. Similarly, we can define a not normalized version of this acceleration, namely

$$\tilde{\mathcal{R}}_{KZ}(\pi, \lambda) := \tilde{\mathcal{R}}^{\kappa(\pi, \lambda)}(\pi, \lambda).$$

For any  $n \in \mathbb{N}$ , we denote

$$(\pi^{(n)}, \lambda^{(n)}) := \mathcal{R}_{KZ}^n(\pi, \lambda).$$

and denote by  $I^{(n)}$  the domain of definition of  $\tilde{\mathcal{R}}_{KZ}^n(\pi, \lambda)$  and the associated exchanged intervals by  $\{I_\alpha^{(n)}\}_{\alpha \in \mathcal{A}}$ . As before, we can express the interval  $[0, 1)$  as can be represented as a union of disjoint Rokhlin towers with heights  $q^{(n)} = (q_\alpha^{(n)})_{\alpha \in \mathcal{A}}$  and bases  $\{I_\alpha^{(n)}\}_{\alpha \in \mathcal{A}}$

$$I = \bigsqcup_{\alpha \in \mathcal{A}} \bigsqcup_{i=0}^{q_\alpha^{(n)}-1} T^i I_\alpha^{(n)}.$$

We will also consider an associated cocycle of matrices, called *Kontsevich-Zorich cocycle*, defined as  $B(\pi, \lambda) := A^{(n(\pi, \lambda))}$ . We will also use the notation for intermediate matrices

$$B(n, n+1)(\pi, \lambda) := B(\pi^{(n)}, \lambda^{(n)}), \quad (4)$$

and the cocycle notation

$$B^{(0)}(\pi, \lambda) = \text{Id}, \quad B^{(n)}(\pi, \lambda) = B(\pi^{(n-1)}, \lambda^{(n-1)})B^{(n-1)}(\pi, \lambda), \quad n \geq 1. \quad (5)$$

Denoting

$$Q(\pi, \lambda) := B^*(\pi, \lambda),$$

we can express the lengths and heights of these iterates using cocycle notation

$$\lambda^{(n)} = B^{(n)}(\pi, \lambda)\lambda, \quad q^{(n)} = Q^{(n)}(\pi, \lambda)[1, \dots, 1], \quad (6)$$

where  $Q^{(n)}(\pi, \lambda)$  is defined in a manner analogous to (5).

We define the intermediate steps  $Q(n, n+1)(\pi, \lambda)$  in a manner similar to (4). Moreover, if there is no risk of confusion, we will sometimes omit the IET being considered in the notations for intermediate steps of the cocycles  $B$  and  $Q$  and denote, for example,  $Q(n, n+1)(\pi, \lambda)$  simply by  $Q(n, n+1)$ .

The crucial difference between  $\mathcal{R}_{KZ}$  and  $\mathcal{R}$  is that the former preserves a *finite* measure  $\rho = \rho_{\mathfrak{R}}$ , as proven by Zorich in [19]. In the same article, Zorich proved that  $\mathcal{R}_{KZ}$  is actually *integrable*. More precisely, he proved that

$$\int_{\mathfrak{R} \times \Lambda^{\mathcal{A}}} \log \|B(\pi, \lambda)\| d\rho, \int_{\mathfrak{R} \times \Lambda^{\mathcal{A}}} \log \|Q(\pi, \lambda)\| d\rho < \infty. \quad (7)$$

The above fact is one of the crucial tools used in this article since this allow us to consider the Lyapunov exponents associated with these cocycles. Notice that since  $B$  and  $Q$  are *dual* to each other (we refer the interested reader to [20] for a precise definition of dual cocycle) they have the same Lyapunov spectrum when restricted to any ergodic component of  $\mathcal{R}_{KZ}$ .

Let us recall that, as a consequence of several classical works [3, 7, 16, 19], for any Rauzy class  $\mathfrak{R}$ , the Zorich acceleration is ergodic when restricted to  $\mathfrak{R} \times \Lambda^{\mathcal{A}}$  and that, for some  $g \geq 1$  depending only on  $\mathfrak{R}$ , the cocycles  $B$  and  $Q$  restricted to  $\mathfrak{R} \times \Lambda^{\mathcal{A}}$  possess  $2g$  non-zero Lyapunov exponents of the form

$$-\theta_1 < \dots < -\theta_g < 0 < \theta_g < \dots < \theta_1,$$

while the remaining  $d - 2g$  Lyapunov exponents are equal to 0.

For more information on interval exchange transformations and associated renormalizations, we refer the reader to [17] and [18].

**2.4. Path-induced domains.** We prove now a series of simple results concerning the Rauzy-Veech (and Zorich-Kontsevich) cocycle. Their main purpose is to pick a proper domain of the first return of Zorich-Kontsevich cocycle in the proof of Proposition 4.1.

Let  $\mathfrak{R}$  be a Rauzy graph and let  $\gamma$  be a path of length  $L := |\gamma|$  in  $\mathfrak{R}$ . Let

$$\Delta_\gamma := \pi_{\Lambda^A} A_\gamma(\mathbb{R}_+^A) = \{(\pi, \lambda) \in \mathfrak{R} \times \Lambda^A \mid (\pi, \lambda) \text{ follows } \gamma \text{ via } \mathcal{R}\},$$

where  $\pi_{\Lambda^A} : \mathbb{R}_+^A \rightarrow \Lambda^A$  is the projection  $v \mapsto \frac{v}{|v|_1}$  onto the simplex  $\Lambda^A$ . It turns out that by extending the path  $\gamma$  we may guarantee that we do not see path  $\gamma$  as an initial Rauzy path for the first  $L$  iteration of  $\mathcal{R}$ .

**Lemma 2.1.** *There exists a path  $\tilde{\gamma}$  of length  $2L$  such that  $\Delta_{\tilde{\gamma}} \subset \Delta_\gamma$  and*

$$\mathcal{R}^i(\Delta_{\tilde{\gamma}}) \cap \Delta_\gamma = \emptyset \quad \text{for } i = 1, \dots, L-1.$$

*Proof.* WLOG suppose that the last arrow in the path  $\gamma$  was of the "bottom" type, the prove in the opposite case is analogous. Take the path  $\tilde{\gamma} := \gamma * \gamma_L^t$ , where  $\gamma_L^t$  is a path of length  $L$ , starting at the final vertex of  $\gamma$ , such that all its arrows are of "top" type. Since  $\tilde{\gamma}$  starts with  $\gamma$ , we have  $\Delta_{\tilde{\gamma}} \subset \Delta_\gamma$ . Moreover, for any  $(\pi, \lambda) \in \Delta_{\tilde{\gamma}}$  and  $i = 1, \dots, L-1$ , we have that  $\mathcal{R}^{i+L}$  is of the "top" type. Since the last arrow in  $\gamma$  is of "bottom" type, this finishes the proof of the lemma.  $\square$

We will from now on assume that the matrix  $A_\gamma$  is positive. It is a classical fact that as long as  $(\pi, \lambda)$  is infinitely renormalizable (i.e. the infinite orbit of  $(\pi, \lambda)$  via  $\mathcal{R}$  is well defined), then there exists  $m \in \mathbb{N}$  such that the path obtained by following  $m$  steps of  $\mathcal{R}$  on  $(\pi, \lambda)$  is positive (see e.g. Corollary 5.3 in [17]). We have the following well known fact, which we prove for completeness.

**Lemma 2.2.** *There exists a constant  $C_\gamma > 0$  such that for any  $(\pi, \lambda) \in \mathfrak{R} \times \Lambda^A$  and any  $K \geq L \geq 0$  satisfying  $\mathcal{R}^{K-L}(\pi, \lambda) \in \Delta_\gamma$ ,*

$$\max_{\alpha, \beta \in \mathcal{A}} \frac{q_\alpha^K}{q_\beta^K} \leq C_\gamma.$$

*Proof.* Note that

$$[q_\alpha^K]_{\alpha \in \mathcal{A}} = [q_\alpha^{K-L}]_{\alpha \in \mathcal{A}} \cdot A_\gamma$$

and since  $\gamma$  is positive, we also have  $\min_{\alpha \in \mathcal{A}} q_\alpha^K > \max_{\alpha \in \mathcal{A}} q_\alpha^{K-L}$ . Hence

$$\max_{\alpha, \beta \in \mathcal{A}} \frac{q_\alpha^K}{q_\beta^K} \leq \frac{\max_{\alpha \in \mathcal{A}} q_\alpha^{K-L} \cdot \sum_{\alpha, \beta \in \mathcal{A}} (A_\gamma)_{\alpha\beta}}{\max_{\alpha \in \mathcal{A}} q_\alpha^{K-L}}.$$

Thus, to conclude the proof, it is enough to take  $C_\gamma := \sum_{\alpha, \beta \in \mathcal{A}} (A_\gamma)_{\alpha\beta}$ .  $\square$

For convenience, in the proof of Proposition 4.1 we consider a path that is a concatenation  $\gamma * \gamma * \gamma$  of 3 copies of the path  $\gamma$ . It is easy to check that  $\Delta_{\gamma * \gamma * \gamma} = A_\gamma^3(\mathbb{R}_+^A)$ . However, we will require more from our path. In the next lemma we will construct a path  $\gamma$ , which depends on the starting point and is coherent with the Zorich-Kontsevich acceleration.

**Lemma 2.3.** *For a.e.  $(\pi, \lambda) \in \mathfrak{R} \times \Lambda^A$ , there exists an arbitrarily large  $\ell \in \mathbb{N}$  such that path  $\gamma = \gamma(\pi, \lambda)$  of length  $\ell$ , which appears as a path followed by the orbit of  $(\pi, \lambda)$  via  $\mathcal{R}$  satisfies:*

- $A_\gamma$  is a positive matrix,
- $\pi$  is the final vertex of  $\gamma$ ,
- the first and last arrow of  $\gamma$  are of opposite types.

*In particular, if  $(\pi, \lambda') \in \Delta_{\gamma * \gamma * \gamma * \varphi}$ , where  $\varphi$  is a path of length one consisting of the initial arrow of  $\gamma$ , then*

$$\mathcal{R}^{i\ell}(\pi, \lambda') = \mathcal{R}_{KZ}^{iL}(\pi, \lambda') \quad \text{for } i = 1, 2, 3,$$

*where  $L$  is the number of type changes in the path  $\gamma$ .*

*Proof.* Since  $\mathcal{R}$  is ergodic, we may assume that  $(\pi, \lambda)\mathfrak{R} \times \Lambda^A$  is a generic point for  $\mathcal{R}$ . W.l.o.g we may also assume that  $\mathcal{R}(\pi, \lambda)$  is of "top" type, the "bottom" case is symmetric. As in the proof of Lemma 2.1, by iterating  $\mathcal{R}$  long enough, we obtain a path of some length  $\tilde{m} \in \mathbb{N}$ , whose associated matrix is positive. This applies also to any path obtained afterwards, since the product of non-negative matrix with positive diagonal entries with a positive matrix yields a positive matrix.

Consider an arrow which ends at  $\pi$  that is of "bottom" type. Let  $\pi'$  be the initial vertex of this arrow. Then for a positive measure of  $\hat{\lambda}$ , we have that  $\mathcal{R}(\pi', \hat{\lambda})$  is of "bottom" type. Hence by ergodicity of  $\mathcal{R}$ , there exists  $m \geq \tilde{m}$  such that the path  $\gamma$  obtained as a path followed by the orbit of  $(\pi, \lambda)$  of length  $m$  via  $\mathcal{R}$ , which satisfies the desired conditions. If the path  $\gamma$  is not already long enough, then it suffices to consider a long enough concatenation of paths  $\gamma$ .

The last assertion of the Lemma follows from the definition of the Zorich-Kontsevich acceleration and the fact that the first and the last arrow of  $\gamma$  are of opposite types.  $\square$

Finally, we state a result that we use directly later in the proof of Proposition 4.1.

**Proposition 2.1.** *For a.e.  $(\tilde{\pi}, \tilde{\lambda}) \in \mathfrak{R} \times \Lambda^A$  and for any open neighbourhood  $(\tilde{\pi}, \tilde{\lambda}) \in U \subseteq \mathfrak{R} \times \Lambda^A$ , there exists a path  $\gamma$  in  $\mathfrak{R}$  of length  $|\ell|$  such that  $\Delta_\gamma \subset U$  and there exists a path  $\tilde{\gamma}$  with  $\Delta_{\tilde{\gamma}} \subset \Delta_\gamma$  such that for a.e.  $(\pi, \lambda) \in \Delta_{\tilde{\gamma}}$  the following conditions are satisfied*

- $(\pi, \lambda)$  is infinitely renormalizable.
- The next  $3\ell$  Rauzy-Veech renormalizations of  $(\pi, \lambda)$  follow the concatenated path  $\gamma * \gamma * \gamma$ .
- $\mathcal{R}^{i\ell}(\pi, \lambda) = \mathcal{R}_{KZ}^{iL}(\pi, \lambda)$ , for  $i \in \{1, 2, 3\}$ .
- $\mathcal{R}_{KZ}^i(\pi, \lambda) \notin \Delta$ , for  $i \in \{1, \dots, 3L-1\}$ .
- $\Delta \cup \mathcal{R}_{KZ}^L(\Delta) \cup \mathcal{R}_{KZ}^{2L}(\Delta) \subseteq \mathfrak{R} \times \{\lambda \in \Lambda^A \mid |\lambda_\alpha - \lambda_\beta| \leq \nu\}$ . In particular,

$$\max_{\substack{\alpha, \beta \in A, \\ i \in \{0, 1, 2\}}} \frac{\lambda_\alpha^{(iL)}(\pi, \lambda)}{\lambda_\beta^{(iL)}(\pi, \lambda)} < 1 + \nu. \quad (8)$$

•

$$\max_{\substack{\alpha, \beta \in A, \\ i \in \{1, 2, 3\}}} \frac{q_\alpha^{(iL)}(\pi, \lambda)}{q_\beta^{(iL)}(\pi, \lambda)} < C_\gamma. \quad (9)$$

- The entries of  $A_\gamma$  are all  $\geq 2$ .

*Proof.* Recall that  $\gamma$  can be chosen so that all entries of  $A_\gamma$  are positive. It is a well known fact that positive matrices act as contractions on  $\Lambda^A$  (see e.g. Proposition 26.3 in [17]). Hence, by concatenating the path  $\gamma$  multiple times if necessary, we obtain that  $\Delta_\gamma \subset U$ . Also by concatenating if necessary we have that all entries of  $A_\gamma$  are bigger than 2. The remaining conditions follow from Lemmas 2.1, 2.2 applied to the path  $\gamma * \gamma * \gamma$  and Lemma 2.3.  $\square$

**2.5. Nudging of discontinuities.** We now present one of the main tools, which is *nudging* of the cocycles. This notion was first introduced and is directly inspired by the work of Chaika and Robertson [4]. Given  $f \in C_{m,M}$  of the form  $p_1 \chi_{[0, q_1]} + \dots + p_{m+1} \chi_{[q_1 + \dots + q_m, 1]}$  for some  $(q, p) \in \mathbb{R}_+^{m+1} \times \mathbb{R}^{m+1}$ , we can move the location of its  $i$ -th discontinuity by  $\zeta \in \mathbb{R}$  provided that  $|\zeta| < \frac{\Gamma}{2}$ , where  $\Gamma := \min\{q_1, \dots, q_{m+1}\}$ , by considering the function  $\text{nudge}(f, i, \zeta) \in C_{m,M}$  associated to the vector

$$\left( q_1, \dots, q_{i-1}, q_i + \zeta, q_{i+1} - \zeta, q_{i+1}, \dots, q_{m+1}, p_1, \dots, p_i, \frac{p_{i+1}q_{i+1} - \zeta p_i}{q_{i+1} - \zeta}, p_{i+2}, \dots, p_{m+1} \right).$$

Notice that the location of other discontinuities, as well as  $m-1$  of the possible values taken by  $f$ , remain unchanged. Moreover, we have

$$\left| \frac{p_{i+1}q_{i+1} - \zeta p_i}{q_{i+1} - \zeta}, -p_{i+1} \right| = \left| \frac{\zeta p_{i+1} - \zeta p_i}{q_{i+1} - \zeta} \right| \leq \frac{2M|\zeta|}{|q_{i+1} - \zeta|} \leq \frac{4M|\zeta|}{q_{i+1}} \leq \frac{4M|\zeta|}{\Gamma},$$

which implies the following.

**Lemma 2.4** (Lemma 4.3 in [4]). *For any  $f = p_1 \chi_{[0, q_1]} + \dots + p_{m+1} \chi_{[q_1 + \dots + q_m, 1]} \in C_{m,M}$  and any  $\zeta \in \mathbb{R}$  verifying  $|\zeta| < \frac{1}{2} \min\{q_1, \dots, q_{m+1}\}$ ,*

$$|\zeta| \leq |f - \text{nudge}(f, i, \zeta)| \leq |\zeta| \max \left\{ 1, \frac{4M}{\min\{q_1, \dots, q_{m+1}\}} \right\}.$$



### 3. PROOF OF THEOREM 1.1

We will follow the strategy described in Section 1.1. Let  $T$  be an IET ergodic with respect to the Lebesgue measure, satisfying the conclusions of Theorem 1.2, for some  $C, \delta > 0$ . Notice that by Theorem 1.2, this is satisfied by a full-measure set of IETs.

Suppose, for the sake of contradiction, that for a positive measure set  $W$  of functions  $f$  in  $C_{m,M}$ , the associated skew-products  $T_f$  are not ergodic. We assume WLOG that, for any  $f \in W$ , the group generated by  $\sigma(f)$  is dense in  $\mathbb{R}$  and that the conclusions of Theorem 1.2 hold for  $f$ .

**Step 1 - Construction and decomposition of a family of  $T_f$ -invariant measures.** For any  $f$  in an appropriate positive measure subset  $V \subseteq W$ , we will define a measure  $\mu_f$  on  $[0, 1) \times \mathbb{R}$ , depending in a measurable way on  $f$ , such that

- $\mu_f$  is invariant by  $T_f$ ;
- $\mu_f \neq c\text{Leb}_{[0,1) \times \mathbb{R}}$ , for any  $c \in \mathbb{R} \setminus \{0\}$ ;
- $\mu_f \not\perp \text{Leb}_{[0,1) \times \mathbb{R}}$ .

Our construction relies on many well known techniques and schemes, but is rather technical. For the sake of clarity and completeness, we provide a full proof of the existence of the measures described above in Proposition A.1 of Appendix A.

**Step 2 - Decompose the measures  $\mu_f$  on the fibers.** For each  $f \in V$ , we decompose  $\mu_f$  with respect to the partition of  $[0, 1) \times \mathbb{R}$  in vertical lines, and denote this family of measures by  $\{\mu_{f,x}\}_{x \in [0,1]}$ , where  $\mu_{f,x}$  is supported on  $\{x\} \times \mathbb{R}$ . Notice that the map  $(f, x) \mapsto \mu_{f,x}$  is measurable with respect to  $f$  and  $x$ . Indeed, it is formed by gluing the pieces obtained from standard disintegration over the first coordinate on the sets of the form  $[0, 1] \times [N, N+1]$ ,  $N \in \mathbb{Z}$ . Moreover, since  $\mu_f$  is  $T_f$  invariant and by the uniqueness of this decomposition,

$$\mu_{f,T^k(x)} = (T_f^k)_*(\mu_{f,x}), \quad (10)$$

for a.e.  $x$  and any  $k \in \mathbb{Z}$ . As an abuse of notation, in the following, we will treat the measures  $\{\mu_{f,x}\}_{x \in [0,1]}$  as measures on  $\mathbb{R}$ . With this convention, equation (10) becomes

$$\mu_{f,T^k(x)} = (V_{S_k f(x)})_*(\mu_{f,x}), \quad (11)$$

where  $S_k f(x)$  denotes the  $k$ -th Birkhoff sum of  $f$  with respect to  $T$  and

$$\begin{aligned} V_r : \mathbb{R} &\rightarrow \mathbb{R} \\ t &\mapsto t + r \end{aligned} \quad (12)$$

for any  $r \in \mathbb{R}$ .

We will assume WLOG that, for any  $f \in V$ ,

- $\mu_{f,x}$  is not a multiple of  $\text{Leb}_{\mathbb{R}}$ , for a.e.  $x \in [0, 1)$ .

Indeed, if there exists a positive measure set of points in  $[0, 1]$  for which  $\mu_{f,x} = c_x \text{Leb}_{\mathbb{R}}$ , for some constants  $c_x \in \mathbb{R}$ , then, by (10) and the ergodicity of  $T$  with respect to the Lebesgue measure, it follows that a.e.  $x \in [0, 1)$  is of the form  $\mu_{f,x} = c_x \text{Leb}_{\mathbb{R}}$ . Since the function  $x \mapsto c_x$  is  $T$ -invariant, by the ergodicity of  $T$  with respect to the Lebesgue measure, there exists  $c \in \mathbb{R}$  such that  $c_x = c$ , for a.e.  $x \in [0, 1)$ . Therefore,  $\mu_f = c\text{Leb}_{[0,1) \times \mathbb{R}}$ , which contradicts our initial assumption on  $\mu_f$ .

Note that, since the map  $(f, x) \mapsto \mu_{f,x}$  is measurable, the set  $U = \{(f, x) \in V \times [0, 1) \mid \mu_{f,x} \neq c\text{Leb}_{\mathbb{R}}, \forall c \in \mathbb{R}\}$  is measurable as well.

**Step 3 - Consider an appropriate continuity set for the (cutoffs of) the map  $(f, x) \mapsto \mu_{f,x}$ .** For  $L > 0$ , let  $\mu_{f,x,L}$  be the normalized cut-off of the measure of  $\mu_{f,x}$  to  $[-L, L]$ . By Lusin's Theorem, there exists a compact set  $K \subseteq U$  of positive measure such that, for any  $D \in \mathbb{Q} \cap (0, +\infty)$ ,

- $\Phi_D : K \subseteq U \rightarrow \mathcal{M}_L$  is continuous,  
 $(f, x) \mapsto \mu_{f,x,L}$

where  $\mathcal{M}_D$  denotes the space of probability measures on  $[-L, L]$ .

Moreover, there exists a positive measure subset  $B \subseteq K$  so that the following holds.

- The sets  $E_f = \{y \in [0, 1) \mid (f, y) \in B\}$  and  $P_x = \{g \in C_{m,M} \mid (g, x) \in B\}$  have positive Lebesgue measure, for any  $(f, x) \in B$ .

- $x$  (resp.  $f$ ) is a density point of  $E_f$  (resp.  $P_x$ ), for any  $(f, x) \in B$ .
- For any  $0 < \xi < 1$ , there exists  $N \in \mathbb{N}$  such that

$$\frac{\text{Leb}_{C_{m,M}}\{g \in P_x \mid d(f, g) < \frac{1}{n}\}}{\text{Leb}_{C_{m,M}}\{g \in C_{m,M} \mid d(f, g) < \frac{1}{n}\}} > \xi, \quad \frac{\text{Leb}_{[0,1]}\{y \in E_f \mid |x - y| < \frac{1}{n}\}}{\frac{1}{2n}} > \xi, \quad (13)$$

for any  $n > N$  and any  $(f, x) \in B$ .

Indeed, we may define a map  $\Xi : K \times [0, 1] \rightarrow [0, 1]^2$  given by the following formula

$$\Xi((f, x), \epsilon) := \left( \frac{\text{Leb}_{C_{m,M}}\{g \in C_{m,M} \mid (g, x) \in K, d(f, g) < \epsilon\}}{\text{Leb}_{C_{m,M}}\{g \in C_{m,M} \mid d(f, g) < \epsilon\}}, \frac{\text{Leb}_{[0,1]}\{y \in E_f \mid (f, y) \in K, |x - y| < \epsilon\}}{2\epsilon} \right).$$

Note that, if  $(f, x) \in B$  is a density point of  $K$ , then by Lebesgue Density Theorem applied to the set  $K$   $\lim_{\epsilon \rightarrow 0} \Xi((f, x), \epsilon) = (1, 1)$ . Since a.e. point in  $K$  is a density point, we can find a subset  $B' \subseteq K$  of density points of positive Lebesgue measure, such that for any  $0 < \xi < 1$ , there exists  $N_\xi$  such that for any  $(f, x) \in B'$  we have

$$\Xi((f, x), \epsilon) \subseteq (\xi, 1]^2,$$

for any  $\epsilon > 0$  small enough. It is now enough to take  $B$  to be the set of density points of  $B'$ , and notice that for every  $(f, x) \in B$ , vanishing in measure of either of the sets  $E_f$  and  $P_x$  would contradict the definition of a density point.

**Step 4 - Find good returns to the set  $E_f$ .** Fix  $(f, x) \in B$  and  $D > mM$ . We can define sequences  $(x_k)_{k \geq 1} \subseteq E_f$ ,  $(r_k)_{k \geq 1} \subseteq \mathbb{N}$ , with  $r_k \nearrow +\infty$ , such that:

- (1)  $x_k, T^{r_k}(x_k) \in E_f \cap (x - \frac{1}{k}, x + \frac{1}{k})$ ,
- (2)  $|S_{r_k}f(x_k)| < D$ ,
- (3)  $\{T^i(x_k)\}_{i=0}^{r_k-1}$  is  $\frac{C'}{r_k}$ -dense in  $I$ ,
- (4)  $(x_k - \frac{\sigma'}{r_k}, x_k + \frac{\sigma'}{r_k})$  is a continuity interval for  $T^{r_k}$ ,

where  $C' := C + 1$ ,  $\sigma' := \frac{\sigma}{4}$  and  $C, \sigma > 0$  are the constants for which the conclusions of Theorem 1.2 hold.

Indeed, let  $N_0 \geq 1$ . For any  $k \geq 1$ , Theorem 1.2 applied to  $E_k = E_f \cap (x - \frac{1}{2k}, x + \frac{1}{2k})$  with  $N = 2k + N_0$  yields a point  $y_k \in E_f \cap (x - \frac{1}{2k}, x + \frac{1}{2k})$  and a natural number  $r_k > N$  such that  $y_k, T^{r_k}(y_k) \in E_k$ ,  $|S_{r_k}f(y_k)| < D$ ,  $\{T^i(y_k)\}_{i=0}^{r_k-1}$  is  $\frac{C}{r_k}$ -dense in  $I$ , and either  $[y_k - \frac{\sigma}{r_k}, y_k]$  or  $[y_k, y_k + \frac{\sigma}{r_k}]$  is a continuity interval of  $T^{r_k}$ . Let  $J_k$  denote this continuity interval.

By (13), up to consider  $N_0$  sufficiently large, we have

$$|J_k \cap E_f|, |T^{r_k}(J_k) \cap E_f| > \frac{3|J_k|}{4}. \quad (14)$$

Notice that, since  $J_k, T^{r_k}(J_k) \subseteq (x - \frac{1}{k}, x + \frac{1}{k})$ , we have

$$J_k \cap E_f = J_k \cap E_f \cap (x - \frac{1}{k}, x + \frac{1}{k}) \quad T^{r_k}(J_k) \cap E_f = T^{r_k}(J_k) \cap E_f \cap (x - \frac{1}{k}, x + \frac{1}{k}),$$

and thus equation (14) yields

$$|J_k \cap E_f \cap (x - \frac{1}{k}, x + \frac{1}{k}) \cap T^{-r_k}(E_f \cap (x - \frac{1}{k}, x + \frac{1}{k}))| > \frac{|J_k|}{2}.$$

Therefore, there exists  $x_k \in \frac{1}{2}J_k$ , where  $\frac{1}{2}J_k$  denotes the centered interval inside  $J_k$  of length  $\frac{1}{2}|J_k|$ , satisfying the first two assertions above.

Since  $\{T^i(x_k)\}_{i=0}^{r_k-1}$  is simply a translation of  $\{T^i(y_k)\}_{i=0}^{r_k-1}$  by at most  $\frac{\sigma}{r_k}$ ,  $x_k$  verifies the third assertion. Finally, noticing that  $(x_k - \frac{\sigma}{4r_k}, x_k + \frac{\sigma}{4r_k}) \subseteq J_k$ , it follows that  $x_k$  also verifies the fourth assertion.

**Step 5 - Construct appropriate perturbations of  $f$ .** We will define  $\xi_1 > \xi_2 > 0$  and sequences  $(g_{i,k})_{k \geq 1} \subseteq C_{m,M}$  of perturbations of  $f$ , for  $i = 1, \dots, m$ , such that, for any  $k \geq 1$  and any  $i = 1, \dots, m$ ,

- (1)  $|g_{i,k} - f| < \frac{\xi_1}{r_k}$ .

- (2) For any  $g \in C_{m,M}$  verifying  $|g - g_{i,k}| < \frac{\xi_2}{r_k}$ , the  $i$ -th discontinuity of  $g$  belongs to the Rokhlin tower  $\bigsqcup_{j=0}^{r_k-1} T^j(x_k - \frac{\sigma'}{2r_k}, x_k + \frac{\sigma'}{2r_k})$ . Moreover, no other discontinuity of  $h$  belongs to  $\bigsqcup_{j=0}^{r_k-1} T^j(x_k - \frac{3\sigma'}{4r_k}, x_k + \frac{3\sigma'}{4r_k})$ .

We will do it by nudging, as defined in Lemma 2.4. It allows to control a relative position of the discontinuities inside the chosen Rokhlin towers.

Fix  $1 \leq i \leq m$ . By nudging  $f$  using Lemma 2.4, let us construct a function  $g_{i,k} \in C_{m,M}$  such that its  $i$ -th discontinuity belongs to  $\bigsqcup_{j=0}^{r_k-1} T^j(x_k - \frac{\sigma'}{4r_k}, x_k + \frac{\sigma'}{4r_k})$  and no other of its discontinuities belong to  $\bigsqcup_{j=0}^{r_k-1} T^j(x_k - \frac{4\sigma'}{5r_k}, x_k + \frac{4\sigma'}{5r_k})$ .

Recall that any  $f \in C_{m,M}$  is identified with a vector  $(q, p) \in \mathbb{R}^{m+1} \times \mathbb{R}^{m+1}$ . Let us denote  $\Gamma := \min\{q_1, \dots, q_{m+1}\}$ . By nudging  $f$  at most  $m$  times by a distance of at most  $\frac{4\sigma'}{5r_k}$  (which we assume WLOG to be smaller than  $\frac{\Gamma}{4}$ ), we can define a function  $\bar{g}_k \in C_{m,M}$  having no discontinuities in  $\bigsqcup_{j=0}^{r_k-1} T^j(x_k - \frac{4\sigma'}{5r_k}, x_k + \frac{4\sigma'}{5r_k})$  and such that  $|f - \bar{g}_k| \leq \frac{4m\sigma'}{5r_k} \frac{4M+1}{\Gamma}$ . Indeed, we can move the location of any discontinuity of  $f$  in an interval of the form  $T^j(x_k - \frac{4\sigma'}{5r_k}, x_k + \frac{4\sigma'}{5r_k})$  to  $T^j(x_k - \frac{\sigma'}{r_k}, x_k + \frac{\sigma'}{r_k}) \setminus T^j(x_k - \frac{4\sigma'}{5r_k}, x_k + \frac{4\sigma'}{5r_k})$ .

Since  $\{T^i(x_k)\}_{i=0}^{r_k-1}$  is  $\frac{C'}{r_k}$ -dense in  $[0, 1]$ , it follows that  $[0, 1] \setminus \bigsqcup_{j=0}^{r_k-1} T^j(x_k - \frac{\sigma'}{4r_k}, x_k + \frac{\sigma'}{4r_k})$  is a finite union of closed intervals of length at most  $\frac{C'}{r_k}$ . Hence, we can define a function  $g_{i,k} \in C_{m,M}$  whose  $i$ -th discontinuity belongs to  $\bigsqcup_{j=0}^{r_k-1} T^j(x_k - \frac{\sigma'}{4r_k}, x_k + \frac{\sigma'}{4r_k})$  and verifies  $|\bar{g}_k - g_{i,k}| \leq \frac{C'}{r_k} \frac{4M+1}{\Gamma}$  by moving the location of the  $i$ -th discontinuity of  $\bar{g}_k$  by a distance of at most  $\frac{C'}{r_k}$  (which we again assume WLOG to be smaller than  $\frac{\Gamma}{4}$ ) to the closest interval in the Rokhlin tower  $\bigsqcup_{j=0}^{r_k-1} T^j(x_k - \frac{\sigma'}{4r_k}, x_k + \frac{\sigma'}{4r_k})$ . Moreover, we have

$$|f - g_{i,k}| \leq |f - \bar{g}_k| + |\bar{g}_k - g_{i,k}| \leq \frac{\xi_1}{2r_k},$$

where  $\xi_1 = \frac{2(m\sigma' + C)(4M+1)}{\Gamma}$ .

Finally, setting  $\xi_2 = \frac{\sigma'}{20}$ , for any  $h \in C_{m,M}$  verifying  $|h - g_{i,k}| < \frac{\xi_2}{r_k}$  and any  $1 \leq j \leq m$ , the distance between the  $j$ -th discontinuities of  $h$  and  $g_{i,k}$  is at most  $\frac{\sigma'}{20h_k}$ . Hence, for any such  $h$ , only the  $i$ -th discontinuity of  $h$  belongs to the set  $\bigsqcup_{j=0}^{r_k-1} T^j(x_k - \frac{\sigma'}{2r_k}, x_k + \frac{\sigma'}{2r_k})$ .

**Step 6 - Define sequences  $(f_{i,k}, x_k^\pm) \in B$  converging to  $(f, x)$  for which  $S_{r_k} f_{i,k}(x_k^\pm)$  are bounded and differ by  $\sigma'_i(f_{i,k})$ .** We now find two points  $x_k^-$  and  $x_k^+$ , to the left and to the right of  $x_k$ , respectively, such that, roughly speaking, the fiber measures corresponding to  $x_k^+$  and  $x_k^-$  are close to the fiber measure corresponding to  $x_k$  and yet differ via shift given by the jump discontinuity.

By (13), for  $k$  sufficiently large,

$$\min \left\{ \mu \{y \in E_f \mid |x_k - y| < \frac{\sigma'}{r_k}\}, \mu \{y \in E_f \mid |T^{r_k}(x_k) - y| < \frac{\sigma'}{r_k}\} \right\} > \frac{7}{8} \frac{2\sigma'}{r_k}.$$

Hence

$$\begin{aligned} \mu \left\{ y \in (x_k - \frac{3\sigma'}{4r_k}, x_k - \frac{\sigma'}{2r_k}) \mid y \in E_f \text{ and } T^{r_k}(y) \in E_f \right\} &> 0, \\ \mu \left\{ y \in (x_k + \frac{\sigma'}{2r_k}, x_k + \frac{3\sigma'}{4r_k}) \mid y \in E_f \text{ and } T^{r_k}(y) \in E_f \right\} &> 0. \end{aligned}$$

Thus, there exist  $x_k^- \in (x_k - \frac{3\sigma'}{4r_k}, x_k - \frac{\sigma'}{2r_k})$  and  $x_k^+ \in (x_k + \frac{\sigma'}{2r_k}, x_k + \frac{3\sigma'}{4r_k})$  such that

$$(f, x_k^\pm), (f, T^{r_k}(x_k^\pm)) \in B.$$

Similarly, by (13) and for  $k$  sufficiently large,

$$\frac{\min \left\{ \mu \{g \in P_{x_k^\pm} \mid |g - f| < \frac{\xi_1}{r_k}\}, \mu \{g \in P_{T^{r_k}(x_k^\pm)} \mid |g - f| < \frac{\xi_1}{r_k}\} \right\}}{\mu \{g \in C_{m,M} \mid |g - f| < \frac{\xi_1}{r_k}\}} > 1 - \frac{\xi_2^m}{8\sqrt{m+1}}.$$

Hence, noticing that  $\xi_2 < \frac{\xi_1}{2}$  and recalling that  $|f - g_{i,k}| < \frac{\xi_1}{2}$ , there exists  $f_{i,k} \in C_{m,M}$  such that

$$|f_{i,k} - g_{i,k}| < \frac{\xi_2}{r_k}, \quad (f_{i,k}, x_k^\pm), (f_{i,k}, T^{r_k}(x_k^\pm)) \in B.$$

Notice that by the second assertion in Step 5, we have

$$S_{r_k} f_{i,k}(x_k^-) - S_{r_k} f_{i,k}(x_k^+) = \sigma_i(f_{i,k}).$$

Moreover, by the second assertion in Step 4 and since  $|f_{i,k} - f| < \frac{\xi_1}{r_k}$ ,

$$|S_{r_k} f_{i,k}(x_k^\pm)| < D + \xi_1 + M.$$

**Step 7 - Use the measures  $\mu_{f_{i,k}, x_k^\pm}$  to conclude that  $(V_{\sigma_i(f)})_*(\mu_{f,x}) = \mu_{f,x}$ .** In the following, let  $i \in \{1, \dots, m\}$  be fixed. Notice that to prove  $(V_{\sigma_i(f)})_*(\mu_{f,x}) = \mu_{f,x}$ , where  $V_{\sigma_i(f)}$  is as in (12), it is sufficient to show that

$$(V_{\sigma_i(f)})_*(\mu_{f,x})(J) = \mu_{f,x}(J), \quad (15)$$

for any open bounded interval  $J$ .

Recall that by definition of the set  $B$  and by the constructions of the sequences in the previous sections, we have, for any  $L > 0$ ,

$$\mu_{f_{i,k}, x_k^\pm, L}, \mu_{f_{i,k}, T^{r_k}(x_k^\pm), L} \xrightarrow[k \rightarrow \infty]{w} \mu_{f,x,L}. \quad (16)$$

Let  $J = (a, b)$  be an open bounded interval and let  $L > 2 \max\{|a|, |b|\} + D + \xi_1 + M + 1$ . By taking a subsequence, if necessary, we may assume that  $S_{r_k} f_{i,k}(x_k^-)$  converges, as  $k \rightarrow \infty$ . Let us denote its limit by  $t$ . Denote,

$$J_\epsilon = (a - \epsilon, b + \epsilon),$$

for any  $-\frac{(b-a)}{2} < \epsilon < \frac{(b-a)}{2}$ . Notice that for any  $\epsilon$  as before, we have

$$J_\epsilon + t \subseteq [-L, L].$$

By (11), for  $0 < \epsilon < \frac{(b-a)}{2}$  sufficiently small,

$$\begin{aligned} \mu_{f_{i,k}, T^{r_k}(x_k^\pm), L}(\overline{J_{-\epsilon}} - t) &\geq \mu_{f_{i,k}, T^{r_k}(x_k^\pm), L}(J_{-\epsilon} - t) \\ &= \mu_{f_{i,k}, x_k^\pm, L}(J_{-\epsilon} - t + S_{r_k} f_{i,k}(x_k^\pm)) \\ &\geq \mu_{f_{i,k}, x_k^\pm, L}(J_{-2\epsilon} + \frac{\sigma_i(f)}{2} \pm \frac{\sigma_i(f)}{2}). \end{aligned}$$

By Portmanteau's Theorem and (16), taking the limit as  $k \rightarrow \infty$  in the previous expression yields

$$\mu_{f,x,L}(\overline{J_{-\epsilon}} - t) \geq \mu_{f,x,L}(J_{-2\epsilon} + \frac{\sigma_i(f)}{2} \pm \frac{\sigma_i(f)}{2}).$$

By taking the limit as  $\epsilon \searrow 0$ , we obtain

$$\mu_{f,x,L}(J - t) \geq \mu_{f,x,L}(J + \frac{\sigma_i(f)}{2} \pm \frac{\sigma_i(f)}{2}).$$

Similarly, by (11) and for  $0 < \epsilon < \frac{(b-a)}{2}$  sufficiently small,

$$\begin{aligned} \mu_{f_{i,k}, T^{r_k}(x_k^\pm), L}(J_{-2\epsilon} - t) &= \mu_{f_{i,k}, x_k^\pm, L}(J_{-2\epsilon} - t + S_{r_k} f_{i,k}(x_k^\pm)) \\ &\leq \mu_{f_{i,k}, x_k^\pm, L}(\overline{J_{-\epsilon}} + \frac{\sigma_i(f)}{2} \pm \frac{\sigma_i(f)}{2}). \end{aligned}$$

As before, taking limits first as  $k \rightarrow \infty$  and then as  $\epsilon \searrow 0$ , we obtain

$$\mu_{f,x,L}(J - t) \leq \mu_{f,x,L}(J + \frac{\sigma_i(f)}{2} \pm \frac{\sigma_i(f)}{2}).$$

Therefore,

$$\mu_{f,x,L}(J - t) = \mu_{f,x,L}(J + \frac{\sigma_i(f)}{2} \pm \frac{\sigma_i(f)}{2}).$$

In particular,

$$\mu_{f,x}(J + \sigma_i(f)) = \mu_{f,x,L}(J + \sigma_i(f)) = \mu_{f,x,L}(J) = \mu_{f,x}(J),$$

which shows (15).

**Step 8 - Conclude the proof.** Since the set of jumps of  $f$  generates a dense group in  $\mathbb{R}$  and, by the previous step, we have

$$(V_{\sigma_i(f)})_*(\mu_{f,x}) = \mu_{f,x},$$

for any  $i = 1, \dots, m$ , it follows that  $\mu_{f,x}$  is a multiple of the Lebesgue measure on  $\mathbb{R}$ , which contradicts our initial assumption on the measure  $\mu_{f,x}$ .

#### 4. PROOF OF THEOREM 1.2

Throughout this section, we fix  $d, m \geq 2$ ,  $M > 0$ , and a Rauzy class  $\mathfrak{R}$ . We denote by  $\theta_1$  the largest Lyapunov exponent of  $\mathfrak{R}_{KZ}$  when restricted to  $\mathfrak{R} \times \Lambda^{\mathcal{A}}$ . If  $d > 2$ , we denote by  $\theta_2$  the second largest Lyapunov exponent of  $\mathfrak{R}_{KZ}$  when restricted to  $\mathfrak{R} \times \Lambda^{\mathcal{A}}$ . Otherwise, we let  $\theta_2 = 0$ . Notice that in both cases  $0 \leq \theta_2 < \theta_1$ .

First, it will be crucial for us to have control over the maximum possible growth of Birkhoff sums of a given piecewise constant cocycle over a given IET. This will be a direct consequence of the following well-known result due to Zorich [20, Theorem 1].

**Theorem 4.1.** *For a.e. IET  $T = (\pi, \lambda) \in \mathfrak{R} \times \Lambda^{\mathcal{A}}$*

$$\max_{x \in [0,1)} \limsup_{n \rightarrow \infty} \frac{\log |\chi_{\alpha}(x, n) - \lambda_{\alpha} n|}{\log n} = \frac{\theta_2}{\theta_1},$$

where  $\chi_{\alpha}(x, n)$  is the number of visits to  $I_{\alpha}$  of the first  $n$  iterates of  $x$  by  $T$ . Moreover, given  $\varsigma > 0$  there exists  $N(T, \varsigma) \in \mathbb{N}$  such that for any  $n \geq N$ ,

$$\max_{x \in [0,1)} |\chi_{\alpha}(x, n) - \lambda_{\alpha} n| \leq n^{\frac{\theta_2}{\theta_1} + \varsigma}.$$

By considering IETs with  $m$  marked points in the theorem above, one can easily deduce the following.

**Theorem 4.2** (Deviation of ergodic averages). *For a.e. IET  $T = (\pi, \lambda) \in \mathfrak{R} \times \Lambda^{\mathcal{A}}$  there exists a full-measure set  $\mathcal{F} \subseteq C_{m,M}$  such that, for any  $f \in \mathcal{F}$  and any  $\varsigma > 0$ , there exists  $C = C(T, f, \varsigma) > 0$  satisfying*

$$\max_{x \in [0,1)} |S_n^T f(x)| \leq C n^{\frac{\theta_2}{\theta_1} + \varsigma},$$

for any  $n \geq 1$ .

In addition to this, we want appropriate collections of Rokhlin towers. The main requirements to define these Rokhlin towers are the following: First, for any subtower whose height is a fixed proportion of the total height, say  $\frac{1}{\eta}$ , this subtower remains almost as dense in  $[0, 1)$  (up to a constant depending on  $\eta$ ) as the full tower. Second, the heights of the full Rokhlin towers grow at a rate only slightly larger than  $\eta^{1+\epsilon}$ , for some  $\epsilon > 0$  sufficiently small.

**Proposition 4.1.** *Given  $\epsilon > 0$ , there exist  $C, \eta_0, \sigma > 0$  and a positive measure set  $\Delta \subseteq \mathfrak{R} \times \Lambda^{\mathcal{A}}$ , depending only on  $\mathfrak{R}$  and  $\epsilon$ , such that for a.e. IET  $T = (\pi, \lambda) \in \mathfrak{R} \times \Lambda^{\mathcal{A}}$  and any  $\eta > \eta_0$  there exists a sequence of recurrence times  $(n_k)_{k \in \mathbb{N}} \subseteq \mathbb{N}$  (to  $\Delta$  with respect to  $\mathcal{R}$ ) for which the sequence*

$$h_{n_k} := \min_{\alpha \in \mathcal{A}} q_{\alpha}^{(n_k)}, \quad k \geq 1,$$

satisfies the following:

- i) For any  $x \in I$  and any  $k \geq 1$ , either  $\bigsqcup_{i=0}^{h_{n_k}-1} T^i([x - \frac{\sigma}{h_{n_k}}, x])$  or  $\bigsqcup_{i=0}^{h_{n_k}-1} T^i([x, x + \frac{\sigma}{h_{n_k}}])$  consists of  $h_{n_k}$  disjoint intervals.
- ii) For any  $x \in I$  and any  $k \geq 1$ , the set  $\{T^i(x)\}_{i=0}^{\lfloor h_{n_k}/\eta \rfloor - 1}$  is  $\frac{C\eta^{1+\epsilon}}{h_{n_k}}$ -dense in  $I$ .
- iii)  $\limsup_{k \rightarrow \infty} (h_{n_k})^{\frac{1}{k}} \leq C\eta^{1+\epsilon}$ .

As we shall see in Proposition 4.2, the collections of Rokhlin towers given by Proposition 4.1 together with Theorem 4.2 will allow us to combine, for some point  $x$  in a given positive measure subset  $E$ , the recurrence times of  $x$  to  $E$  with the times where the Birkhoff sums  $S_n f(x)$  is uniformly bounded.

*Proof of Proposition 4.1.* Fix  $0 < \epsilon < 1$ . Our construction will depend on two (small) positive constants

$$0 < \nu, \delta < \min\{\epsilon, \frac{1}{10d}\}$$

whose exact value, which depends only on  $\mathfrak{R}$  and  $\epsilon$ , we specify at the end on the proof (see (26)).

Let  $(\pi, \hat{\lambda}) \in \mathfrak{R} \times \Lambda^{\mathcal{A}}$  be an IET, generic w.r.t.  $\mathcal{R}$  such that

- $\max_{\alpha, \beta} |\hat{\lambda}_{\alpha} - \hat{\lambda}_{\beta}| \leq \nu/2$ ,
- $1 \leq i < j \leq d \Rightarrow \hat{\lambda}_{\pi_0^{-1}(i)} < \hat{\lambda}_{\pi_0^{-1}(j)}$ .

Let

$$U := \{\pi\} \times \{\lambda \in \Lambda^{\mathcal{A}} \mid \max_{\alpha, \beta} |\lambda_{\alpha} - \lambda_{\beta}| \leq \nu \text{ and } 1 \leq i < j \leq d \Rightarrow \lambda_{\pi_0^{-1}(i)} < \lambda_{\pi_0^{-1}(j)}\}.$$

Consider the  $\gamma = \gamma(\pi, \lambda)$  of length  $\ell := |\gamma|$  and the path  $\tilde{\gamma}$ , both given by Proposition 2.1 applied to  $(\pi, \hat{\lambda})$  and  $U$ .

Since for any  $(\pi, \lambda) \in \Delta_{\tilde{\gamma}}$  the intervals before exchange are ordered increasingly, for any  $\alpha \in \mathcal{A} \setminus \{\pi_0^{-1}(1)\}$  there exist exactly two distinct symbols  $\alpha_l, \alpha_r \in \mathcal{A}$  such that  $I_{\alpha_l}$  and  $I_{\alpha_r}$  are adjacent intervals (more precisely, the right endpoint of  $I_{\alpha_l}$  and the left endpoint of  $I_{\alpha_r}$  coincide) satisfying

$$T(I_{\alpha}) \cap I_{\alpha_l} \neq \emptyset \neq T(I_{\alpha}) \cap I_{\alpha_r}, \quad T(I_{\alpha}) \subseteq I_{\alpha_l} \cup I_{\alpha_r}.$$

Moreover,

$$T(I_{\pi_0^{-1}(1)}) \subseteq I_{\pi_0^{-1}(1)}.$$

By taking a subset  $\Delta \subset \Delta_{\tilde{\gamma}}$  of positive measure, increasing the constant  $C_{\gamma}$  if necessary, we have that

- For any  $(\pi, \lambda) \in \Delta \cup \mathcal{R}_{KZ}^L(\Delta) \cup \mathcal{R}_{KZ}^{2L}(\Delta)$  with exchanged intervals  $\{I_{\alpha}\}_{\alpha \in \mathcal{A}}$

$$T(I_{\alpha}) \cap I_{\beta} \neq \emptyset \Rightarrow |T(I_{\alpha}) \cap I_{\beta}| \geq C_{\gamma}^{-1}. \quad (17)$$

Indeed, it is enough to consider a finite number of subsets of  $\Delta_{\tilde{\gamma}}$  with connections given by single iterations of  $T$  and remove their small enough neighbourhood. This last property will be useful later in the proof when considering Condition (i).

Let us fix an infinitely renormalizable IET  $(\pi, \lambda)$  returning to  $\Delta$  infinitely often under the action of  $\mathcal{R}$  (recall that a.e. IET verifies this by the ergodicity of  $\mathcal{R}$ ) and denote by  $(m_k)_{k \in \mathbb{N}}$  the associated sequence of first returns. Notice that, by Birkhoff's ergodic theorem,

$$\lim_{k \rightarrow \infty} \frac{m_k}{k} = \frac{1}{\rho(\Delta)},$$

where  $\rho$  denotes the  $\mathcal{RV}_{KZ}$ -invariant probability measure described in Section 2. Moreover, since this measure is equivalent to the Lebesgue measure in any compactly contained subset of  $\mathfrak{R} \times \Lambda^{\mathcal{A}}$  (in particular, in  $\mathfrak{R} \times \{\lambda \in \Lambda^{\mathcal{A}} \mid |\lambda_{\alpha} - \lambda_{\beta}| \leq \frac{1}{10d}\}$  which contains  $\Delta$ ), there exists  $C_d > 0$ , depending only on  $d$ , such that  $\rho(\Delta) < C_d \nu$ . Therefore,

$$\limsup_{k \rightarrow \infty} \frac{k}{m_k} < C_d \nu. \quad (18)$$

The desired sequence of return times  $(n_k)_{k \in \mathbb{N}}$  will be a subsequence of  $(m_k + 2L)_{k \in \mathbb{N}}$ . This choice is made so that, for any  $k \geq 0$  and any  $x \in [0, 1)$ , in addition to the length and height vectors satisfying (8) and (9), the orbit  $\{T^i(x)\}_{i=0}^{h_{n_k}-1}$  goes through all the floors of the every tower in the Rokhlin tower decomposition associated to the renormalization time  $m_k$ . This will be useful only later in the proof of Proposition 4.2 and so, for the sake of clarity, we discuss this (and prove it) in detail in Claim 4.1.

By the integrability of the Zorich cocycle (when induced to  $\Delta$ ) and the dominated convergence theorem, we have

$$\lim_{M \rightarrow \infty} \lim_{N \rightarrow \infty} \left[ \frac{1}{N} \sum_{k=1}^N \log \|Q(m_{k-1}, m_k)\| - \frac{1}{N} \sum_{k=1}^N \chi_{\|Q(m_{k-1}, m_k)\| \leq M} \log \|Q(m_{k-1}, m_k)\| \right] = 0, \quad (19)$$

By (19), there exists  $\eta_0 \geq \exp(\frac{1}{\delta})$ , depending only on  $\Delta$  and  $\delta$ , such that for any  $\eta \geq \eta_0$ ,

$$\frac{1}{N} \sum_{k=1}^N \chi_{\|Q(m_{k-1}, m_k)\| > \eta^\delta} \log \|Q(m_{k-1}, m_k)\| \leq \delta. \quad (20)$$

In the following, we will always assume  $\eta \geq \eta_0$ .

Let  $(m_{r_k})_{k \in \mathbb{N}}$  be the subsequence of return times such that

$$\|Q(m_{r_k-1}, m_{r_k})\| > \eta^\delta. \quad (21)$$

By ergodicity of the Zorich-Kontsevich cocycle, this sequence is infinite for a typical IET. Notice that this subsequence verifies

$$\limsup_{k \rightarrow \infty} \frac{k}{r_k} \leq \frac{\delta}{\log \eta^\delta} < \min \left\{ \delta, \frac{1}{\log \eta} \right\}. \quad (22)$$

Indeed, by (20),

$$\frac{k \log(\eta^\delta)}{r_k} \leq \frac{1}{r_k} \sum_{i=1}^{r_k} \chi_{\|Q(m_{i-1}, m_i)\| > \eta^\delta} \log \|Q(m_{i-1}, m_i)\| \leq \delta.$$

Denote

$$C_\Delta := dC_\gamma^2 \|A_\gamma\|^3$$

and define  $(n_k)_{k \in \mathbb{N}} := (m_{l_k} + 2L)_{k \in \mathbb{N}}$  inductively by setting  $l_0 = 0$  and, for any  $k \geq 0$ ,

$$l_{k+1} := \min \left\{ l > l_k \mid \exists l_k < j < l \text{ s.t. } \begin{cases} \|Q(m_j, m_l)\| \geq \eta C_\Delta, \\ \|Q(m_i, m_{i+1})\| \leq \eta^\delta, \text{ for } j \leq i < l. \end{cases} \right\}. \quad (23)$$

Let us point out that the condition above gives rise to a well-defined infinite sequence  $(l_k)_{k \in \mathbb{N}}$ . Indeed, if only  $K \geq 0$  elements of this sequence can be defined, for any  $N$  such that  $r_N > l_K$  and any  $M > N$ , by (20),

$$\begin{aligned} \|Q(m_{r_N}, m_{r_M})\| &\leq \prod_{i=N+1}^M \|Q(m_{r_i-1}, m_{r_i})\| \prod_{i=N}^M \|Q(m_{r_i}, m_{r_{i+1}-1})\| \\ &< \exp(\delta r_M) (\eta C_\Delta)^{M-N}. \end{aligned}$$

Hence, by (22), if  $M$  is sufficiently large then

$$\begin{aligned} \frac{1}{r_M} \log \|Q(m_{r_N}, m_{r_M})\| &\leq \delta + \frac{M}{r_M} (\log \eta + \log C_\Delta) \\ &< \delta + 1 + \delta \log C_\Delta. \end{aligned}$$

Assuming that  $\delta$  is sufficiently small so that the RHS in the previous equation is smaller than  $\log 2$ , this contradicts the fact that  $\|Q(m_{r_N}, m_{r_M})\| \geq \|A_\gamma\|^{r_M - r_N} \geq 2^{r_M - r_N}$ , for any  $M > N \geq 1$ .

Therefore, using (23), we can define sequences  $(l_k)_{k \in \mathbb{N}}, (s_k)_{k \in \mathbb{N}} \subseteq \mathbb{N}$  such that

$$\eta C_\Delta \leq \|Q(m_{l_k - s_k}, m_{l_k})\| \leq \|Q(m_{l_k - s_k}, m_{l_k - 1})\| \|Q(m_{l_k - 1}, m_{l_k})\| \leq \eta^{1+\delta} C_\Delta, \quad l_k < l_{k+1} - s_{k+1},$$

for any  $k \geq 0$ .

Denote  $(\bar{n}_k)_{k \in \mathbb{N}} := (m_{l_k - s_k})_{k \in \mathbb{N}}$ . The equation above implies that

$$\eta C_\Delta \|A_\gamma\|^{-1} \leq \|Q(\bar{n}_k + L, n_k - L)\| = \|Q(m_{l_k - s_k} + L, m_{l_k} + L)\| \leq \eta^{1+\delta} C_\Delta \|A_\gamma\|. \quad (24)$$

We now check that this sequence satisfies all the properties in the statement of the proposition.

Condition (i): Define  $\sigma := \frac{1}{10dC_\gamma}$ . Fix  $x \in [0, 1)$  and  $k \geq 1$ . Let  $\alpha \in \mathcal{A}$  and  $0 \leq i < q_\alpha^{(n_k)}$  such that  $x \in T^i(I_\alpha^{(n_k)})$ . Let  $\alpha_l, \alpha_r \in \mathcal{A}$  be two symbols such that  $T^{q_\alpha^{(n_k)}}(I_\alpha^{(n_k)}) \subseteq I_{\alpha_l}^{(n_k)} \cup I_{\alpha_r}^{(n_k)}$  and  $I_{\alpha_l}^{(n_k)}, I_{\alpha_r}^{(n_k)}$  are adjacent intervals. Notice that, since  $(\pi^{(n_k)}, \lambda^{(n_k)}) \in \Delta$ , by (8), two such symbols always exist. By (17), either

$$\left[x - \frac{\sigma}{q_\alpha^{(n_k)}}, x\right] \subseteq T^i(I_\alpha^{(n_k)}), \quad \text{and} \quad T^{q_\alpha^{(n_k)} - i} \left([x - \frac{\sigma}{q_\alpha^{(n_k)}}, x]\right) \subseteq I_{\alpha_l}^{(n_k)} \text{ or } T^{q_\alpha^{(n_k)} - i} \left([x - \frac{\sigma}{q_\alpha^{(n_k)}}, x]\right) \subseteq I_{\alpha_r}^{(n_k)},$$

or

$$\left[x, x + \frac{\sigma}{q_\alpha^{(n_k)}}\right] \subseteq T^i(I_\alpha^{(n_k)}), \quad \text{and} \quad T^{q_\alpha^{(n_k)}-i}\left(\left[x, x + \frac{\sigma}{q_\alpha^{(n_k)}}\right]\right) \subseteq I_{\alpha_l}^{(n_k)} \text{ or } T^{q_\alpha^{(n_k)}-i}\left(\left[x, x + \frac{\sigma}{q_\alpha^{(n_k)}}\right]\right) \subseteq I_{\alpha_r}^{(n_k)}.$$

Let us point out that both sets of conditions can hold simultaneously.

Assume without loss of generality that

$$\left[x, x + \frac{\sigma}{q_\alpha^{(n_k)}}\right] \subseteq T^i(I_\alpha^{(n_k)}) \quad \text{and} \quad T^{q_\alpha^{(n_k)}-i}\left(\left[x, x + \frac{\sigma}{q_\alpha^{(n_k)}}\right]\right) \subseteq I_{\alpha_l}^{(n_k)},$$

the other cases being analogous.

Then, since  $h_{n_k} = \min_\alpha q_\alpha^{(n_k)}$ , it follows that  $\bigcup_{i=0}^{h_{n_k}-1} T^i\left(\left[x, x + \frac{\sigma}{q_\alpha^{(n_k)}}\right]\right)$  is a union of disjoint intervals contained in the union of the Rokhlin towers  $\bigcup_{i=0}^{q_\alpha^{(n_k)}-1} T^i(I_\alpha^{(n_k)})$  and  $\bigcup_{i=0}^{q_{\alpha_l}^{(n_k)}-1} T^i(I_{\alpha_l}^{(n_k)})$  associated to the  $n_k$  step of renormalization.

Condition (ii): Notice that for any  $x \in [0, 1)$ , any  $k \geq 0$  and any  $\alpha \in \mathcal{A}$ , the orbit  $\{T^i(x)\}_{i=0}^{q_\alpha^{(m_k+2L)}-1}$  is  $\max_\beta \lambda_\beta^{(m_k+L)}$ -dense in  $[0, 1)$ .

Indeed, since the first  $L$  Zorich renormalizations of  $(\pi^{(m_k+L)}, \lambda^{(m_k+L)})$  follow the path  $\gamma$  and  $A_\gamma$  is a positive matrix, the orbit  $\{T^i(x)\}_{i=0}^{q_\alpha^{(m_k+2L)}-1}$  goes through all Rokhlin towers associated to  $(\pi^{(m_k+L)}, \lambda^{(m_k+L)})$  at least once.

Fix  $x \in [0, 1)$  and  $k \geq 1$ . We have

$$\begin{aligned} h_{n_k} &\geq \frac{1}{dC_\gamma} |q^{(n_k)}| = \frac{1}{dC_\gamma} |Q(\bar{n}_k, n_k) q^{(\bar{n}_k)}| \\ &= \frac{1}{dC_\gamma} |A_\gamma Q(\bar{n}_k + L, n_k - L) A_\gamma q^{(\bar{n}_k)}| \\ &\geq \frac{1}{dC_\gamma} \|Q(\bar{n}_k + L, n_k - L)\| |q^{(\bar{n}_k)}| \\ &\geq \eta |q^{(\bar{n}_k)}|, \end{aligned}$$

where, in the third inequality, we use the fact that the matrix  $A_\gamma$  is positive and the matrix  $Q$  is non-negative, and, in the fourth inequality, we apply (24).

Hence, the orbit  $\{T^i(x)\}_{i=0}^{\lfloor h_{n_k}/\eta \rfloor - 1}$  goes fully along at least one of the Rokhlin towers associated with  $(\pi^{(\bar{n}_k)}, \lambda^{(\bar{n}_k)})$ . It follows from the remark above that this orbit is  $\max_\beta \lambda_\beta^{(\bar{n}_k-L)}$ -dense in  $[0, 1)$ .

Since  $\sum_{\beta \in \mathcal{A}} \lambda_\beta^{(\bar{n}_k-L)} q_\beta^{(\bar{n}_k-L)} = 1$ , by (8) and (9),

$$\max_\beta \lambda_\beta^{(\bar{n}_k-L)} \leq \frac{C_\gamma}{d |q^{(\bar{n}_k-L)}|}.$$

Hence, noticing that

$$h_{n_k} \leq C_\gamma |q^{(n_k)}|, \quad q^{(n_k)} = Q(\bar{n}_k - L, n_k) q^{(\bar{n}_k-L)}, \quad \|Q(\bar{n}_k - L, n_k)\| \leq \eta^{1+\delta} C_\Delta \|A_\gamma\|^2,$$

it follows that

$$\max_\beta \lambda_\beta^{(\bar{n}_k-L)} \leq \frac{C_\gamma^2 \eta^{1+\delta} C_\Delta \|A_\gamma\|^2}{dh_{n_k}}.$$

Therefore, denoting

$$C := \frac{C_\gamma^2 C_\Delta \|A_\gamma\|^2}{d},$$

and since  $0 < \delta < \epsilon$ , it follows that the orbit  $\{T^i(x)\}_{i=0}^{\lfloor h_{n_k}/\eta \rfloor - 1}$  is  $\frac{C\eta^{1+\epsilon}}{h_{n_k}}$ -dense in  $[0, 1)$ .

Condition (iii): Fix  $k \geq 1$ . Notice that

$$\|Q(n_{k-1}, n_k)\| \leq (\eta^{1+\delta} C_\Delta \|A_\gamma\|^2) (\eta C_\Delta)^{p_k} \prod_{\substack{i \in \mathbb{N} \\ n_{k-1} < m_{r_i} \leq n_k}} \|Q(m_{r_{i-1}}, m_{r_i})\|,$$

where  $p_k := \#\{i \in \mathbb{N} \mid n_{k-1} < m_{r_i} \leq n_k\}$ . Indeed, recalling that  $n_k = m_{l_k} + 2L$  (see (23)), the times  $m_{r_i}$  (see (21)) satisfying  $n_{k-1} < m_{r_i} \leq n_k$  divide the return times between  $m_{l_{k-1}} = n_{k-1} - 2L$  and  $m_{l_k} = n_k - 2L$  into at most  $p_k + 1$  blocks so that, on each block, the return times satisfy  $\|Q(m_i, m_{i+1})\| \leq \eta^\delta$ . By (23), the norm of the first  $p_k$  blocks is smaller or equal than  $\eta C_\Delta$  while



the norm of the last block is smaller or equal than  $\eta^{1+\delta}C_\delta$ . More precisely, if we denote these  $p_k$  return times by

$$n_{k-1} < m_{t_1} < m_{t_2} < \dots m_{t_{p_k}} < \bar{n}_k < n_k,$$

then,

$$\begin{cases} \|Q(m_{t_i-1}, m_{t_i})\| \geq \eta^\delta, & \text{for } i = 1, \dots, p_k - 1, \\ \|Q(m_{t_i}, m_{t_{i+1}-1})\| \leq \eta C_\Delta, & \text{for } i = 1, \dots, p_k - 1, \\ \|Q(n_{k-1}, m_{t_1-1})\| \leq \|m_{l_{k-1}}, m_{t_1-1}\| \leq \eta C_\Delta, \\ \|Q(m_{t_{p_k}}, n_k)\| \leq \|Q(m_{t_{p_k}}, m_{l_k-1})\| \|Q(m_{l_k-1}, m_{l_k})\| \|Q(m_{l_k}, m_{l_k+2L})\| \leq \eta^{1+\delta} C_\Delta \|A_\gamma\|^2. \end{cases}$$

Hence,

$$\|Q(n_0, n_k)\| \leq (\eta^{1+\delta} C_\Delta \|A_\gamma\|^2)^k (\eta C_\Delta)^{P_k} \prod_{m_0 < m_{r_i} \leq m_{l_k}} \|Q(m_{r_i-1}, m_{r_i})\|, \quad (25)$$

where  $P_k := \sum_{j=1}^k p_j$ . Notice that by (20),  $P_k \leq \frac{l_k}{\log \eta}$ .

By (18), for  $k$  sufficiently large,  $\frac{l_k}{n_k} < C_d \nu$ . Hence, using (20) to bound the product along the return times  $m_{r_i}$  in (25), for  $k$  sufficiently large, we obtain

$$\begin{aligned} \frac{1}{k} \log \|Q(n_0, n_k)\| &\leq \log(\eta^{1+\delta} C_\Delta \|A_\gamma\|^2) + \frac{l_k}{k \log \eta} \log(\eta C_\Delta) + \delta \frac{l_k}{k} \\ &\leq \log(\eta^{1+\delta} C_\Delta \|A_\gamma\|^2) + \frac{l_k}{k} \left(1 + \frac{\log(C_\Delta)}{\log \eta} + \delta\right) \\ &\leq \log(\eta^{1+\delta} C_\Delta \|A_\gamma\|^2) + \frac{n_k}{k} C_d \nu (1 + 2\delta), \end{aligned}$$

assuming  $\eta$  is sufficiently large so that  $\frac{\log(C_\Delta)}{\log \eta} < \delta$ .

Since  $\lim_{k \rightarrow \infty} \frac{\log \|Q(0, N)\|}{N} = \theta_1$  then, for  $k$  sufficiently large, the previous equation yields,

$$\begin{aligned} \frac{n_k}{k} &= \frac{n_k}{\log \|Q(n_0, n_k)\|} \frac{\log \|Q(n_0, n_k)\|}{k} \\ &\leq \frac{1}{\theta_1 - \delta} \left( \log(\eta^{1+\delta} C_\Delta \|A_\gamma\|^2) + \frac{n_k}{k} C_d \nu (1 + 2\delta) \right). \end{aligned}$$

Thus

$$\frac{n_k}{k} \leq \frac{\log(\eta^{1+\delta} C_\Delta \|A_\gamma\|^2)}{\theta_1 - \delta} \left(1 - \frac{C_d \nu (1 + 2\delta)}{\theta_1 - \delta}\right)^{-1}.$$

Finally, for  $k$  sufficiently large,

$$\begin{aligned} (h_{n_k})^{1/k} &\leq \|Q(0, n_k)[1, \dots, 1]\|^{1/k} \\ &\leq \exp\left((\theta_1 + \delta) \frac{n_k}{k}\right) \\ &\leq (\eta^{1+\delta} C_\Delta \|A_\gamma\|^2)^{\frac{\theta_1 + \delta}{\theta_1 - 2\delta}} \left(1 - \frac{C_d \nu (1 + 2\delta)}{\theta_1 - \delta}\right)^{-1} \\ &\leq (\eta C_\Delta \|A_\gamma\|^2)^{1+\epsilon}, \end{aligned}$$

where we assume that  $\nu$  and  $\delta$  are sufficiently small, so that

$$(1 + \delta) \frac{\theta_1 + \delta}{\theta_1 - 2\delta} \left(1 - \frac{C_d \nu (1 + 2\delta)}{\theta_1 - \delta}\right)^{-1} \leq 1 + \epsilon, \quad (26)$$

where  $C_d$  is the constant in (18), which depends only on  $d$ .  $\square$

Using the previous ‘balanced times’ given by Proposition 4.1 we can prove the following.

**Proposition 4.2.** *Fix  $\epsilon > 0$  satisfying*

$$(1 + \epsilon) \frac{\theta_2}{\theta_1} < 1, \quad (27)$$

and let  $\Delta, \gamma, L, C, C_\gamma, \nu, \eta_0, \sigma$  as in Proposition 4.1.

There exists  $\eta \geq \eta_0$ , depending only on  $\mathfrak{R}$  and  $\epsilon$ , such that for any IET  $T = (\pi, \lambda) \in \mathfrak{R} \times \Lambda^A$  as in Proposition 4.1 and Theorem 4.2, and for a.e  $f \in C_{m,M}$ , the following holds.

Let  $(n_k)_{k \geq 1}, (h_{n_k})_{k \geq 1} \subseteq \mathbb{N}$  be the sequences given by Proposition 4.1 associated to  $T$  and  $\eta$ . Let  $D > mM$  and  $E \subseteq [0, 1)$  with positive Lebesgue measure. Then, for any  $P \in \mathbb{N}$  there exist  $y \in E$ , and natural numbers  $p \geq P$  and  $\frac{h_{n_p}}{\eta} \leq n \leq h_{n_p}$ , such that:

- (1)  $T^n(y) \in E$ ,
- (2)  $|S_n f(y)| < D$ .

*Proof of Proposition 4.2.* Fix  $\varsigma > 0$  satisfying

$$(1 + \epsilon) \left( \frac{\theta_2}{\theta_1} + \varsigma \right) < 1 - \varsigma. \quad (28)$$

Notice that such a value exists since, by assumption,  $\epsilon$  verifies (27). Fix  $\eta \in 2\mathbb{N}$  satisfying

$$\eta > \max \left\{ \eta_0, (4C)^{1/\varsigma} \right\}, \quad (29)$$

and let  $(n_k)_{k \geq 1}, (h_{n_k})_{k \geq 1} \subseteq \mathbb{N}$  be the sequences given by Proposition 4.1 associated to  $T$  and  $\eta$ .

Assume, for the sake of contradiction, that the conclusions of Proposition 4.2 do not hold for some  $D, P > 0$  and some  $E \subseteq [0, 1)$  with  $\bar{\delta} := |E| > 0$ .

Up to considering a positive measure subset of  $E$ , we may assume, without loss of generality,

$$\inf_{x \in E} \left( \inf \{ i \geq 1 \mid T^i(x) \in E \} \right) \geq h_{n_P}. \quad (30)$$

Fix  $\vartheta > 0$  satisfying

$$\vartheta < \frac{1}{8\sigma \|A_\gamma\|^2 C_\gamma (1+\nu)}, \quad (31)$$

and let  $x \in E$  be a density point of  $E$ . Fix  $k > P$  and define

$$L_0(k) := \left\{ 0 \leq i < h_{n_k} \mid |F_k(T^i(x)) \cap E| > (1 - \vartheta) |F_k(T^i(x))| \right\},$$

where  $F_k(T^i(x))$  denotes the unique floor containing  $T^i(x)$  in the Rokhlin towers decomposition associated to the renormalization time  $n_k - 2L$ .

**Claim 4.1.** *There exists  $k_0 > 0$  and  $0 < \delta < \bar{\delta}$  such that  $|L_0(k)| \geq \delta h_{n_k}$ , for any  $k \geq k_0$ .*

*Proof of Claim 4.1.* Notice that by the choice of  $\Delta$  in Proposition 4.1, the orbit  $\{x, \dots, T^{h_{n_k}-1}(x)\}$  intersects all floors in the Rokhlin towers decomposition associated to the renormalization time  $n_k - 2L$ . In particular,

$$[0, 1) = \bigcup_{i=0}^{h_{n_k}-1} F_k(T^i(x)). \quad (32)$$

Indeed, as  $q^{(n_k)} = A_\gamma q^{(n_k-L)}$  and all the entries of  $A_\gamma$  are larger or equal to 2, it follows that  $h_{n_k} \geq 2|q^{(n_k-L)}|$ . In particular, the orbit  $\{x, \dots, T^{h_{n_k}-1}(x)\}$  goes through at least one full Rokhlin tower associated with the renormalization time  $n_k - L$ . Moreover, since  $q^{(n_k-L)} = A_\gamma q^{(n_k-2L)}$  and  $A_\gamma$  is positive, this orbit goes through all the Rokhlin towers associated with the renormalization time  $n_k - 2L$  at least once.

Since a.e.  $x \in E$  is a density point of  $E$ , there exists  $t_0 > 0$  such that the set

$$\overline{E} := \left\{ x \in E \mid \forall 0 < t < t_0, \quad \frac{\text{Leb}_{[0,1]} \{ y \in E \mid |x - y| < t \}}{2t} > 1 - \frac{\vartheta}{2} \right\}$$

satisfies

$$|\overline{E}| > \frac{\bar{\delta}}{2}.$$

Let  $k_0 > 0$  sufficiently large so that, for any  $k \geq k_0$ ,

$$\max_{\alpha \in \mathcal{A}} \lambda_\alpha^{(n_k-2L)} < t_0.$$

Fix  $k \geq k_0$  and let us denote

$$\overline{L} = \{ 0 \leq i < h_{n_k} \mid \overline{E} \cap F_k(T^i(x)) \neq \emptyset \}.$$

Notice that, by definition of  $\overline{E}$ ,

$$\overline{L} \subseteq L_0(k).$$

By (32) we have  $\overline{E} \subseteq \bigcup_{i=0}^{h_{n_k}-1} F_k(T^i(x))$ . Thus,

$$\frac{\overline{\delta}}{2} < |\overline{E}| \leq \bigcup_{i \in \overline{L}} F_k(T^i(x)) \leq |\overline{L}| \max_{\alpha \in \mathcal{A}} \lambda_{\alpha}^{(n_k-2L)} \leq |L_0(k)| \max_{\alpha \in \mathcal{A}} \lambda_{\alpha}^{(n_k)} \leq |L_0(k)| \frac{d(1+\nu)C_{\gamma}}{h_{n_k}},$$

where in the last inequality we use  $\sum_{\alpha \in \mathcal{A}} \lambda_{\alpha}^{(n_k)} q_{\alpha}^{(n_k)} = 1$ , together with (8) and (9).

The result holds by setting  $\delta := \frac{1}{\overline{\delta} \frac{1}{2d(1+\nu)C_{\gamma}}}$ .  $\square$

In the following, we assume that  $k \geq \max\{P, k_0\}$ , where  $k_0$  is given by the previous claim.

**Claim 4.2.** *There exists  $L(k) \subseteq L_0(k)$  such that  $|L(k)| \geq \delta \left(\frac{\eta}{4}\right)^{k-P}$  and, for any  $a, b \in L(k)$ , if  $a \neq b$  then there exists  $P \leq p \leq k$  such that  $\frac{h_{n_p}}{\eta} \leq |a - b| \leq h_{n_p}$ .*

*Proof of the Claim.* Starting from  $L_0(k)$ , we define a nested sequence of sets

$$L(k) := L_{k-P}(k) \subseteq L_{k-P-1}(k) \subseteq \dots \subseteq L_0(k) \subseteq \{0, \dots, h_{n_k}\},$$

as follows. First, we split  $\{0, \dots, h_{n_k}\}$  in  $\eta$  disjoint pieces with at most  $\left\lfloor \frac{h_{n_k}}{\eta} \right\rfloor$  consecutive elements by setting

$$J_{0,i}(k) = \left\{ i \left\lfloor \frac{h_{n_k}}{\eta} \right\rfloor, \dots, i \left\lfloor \frac{h_{n_k}}{\eta} \right\rfloor - 1 \right\}, \quad i = 0, \dots, \eta - 1,$$

and

$$J_{0,\eta}(k) = \left\{ \eta \left\lfloor \frac{h_{n_k}}{\eta} \right\rfloor, \dots, h_{n_k} \right\}.$$

Denote  $L_{0,i}(k) := L_0(k) \cap J_{0,i}(k)$ , for  $i = 0, \dots, \eta$ . It follows from Claim 4.1 that

$$\left| \bigsqcup_{i \in 2\mathbb{N} + \varphi(k)} L_{0,i}(k) \right| = \max \left\{ \left| \bigsqcup_{i \text{ odd}} L_{0,i}(k) \right|, \left| \bigsqcup_{i \text{ even}} L_{0,i}(k) \right| \right\} \geq \frac{|L_0(k)|}{2} \geq \delta \frac{h_{n_k}}{2}, \quad (33)$$

where  $\varphi(k) = 1$  if the maximum in the equation above is attained for  $i$  odd, and  $\varphi(k) = 0$  otherwise.

Since by assumption  $h_{n_{k-1}} \leq \left\lfloor \frac{h_{n_k}}{\eta} \right\rfloor$ , for each  $1 \leq i \leq \eta$  there exists a subset  $\overline{J}_{0,i} \subseteq J_{0,i}(k)$  of  $h_{n_{k-1}}$  consecutive elements such that  $\overline{L}_{0,i}(k) := L_0(k) \cap \overline{J}_{0,i}(k)$  verifies

$$|\overline{L}_{0,i}(k)| \geq \frac{\eta h_{n_{k-1}}}{2 h_{n_k}} |L_0(k) \cap J_{0,i}(k)|.$$

We define

$$L_1(k) := \bigsqcup_{i \in 2\mathbb{N} + \varphi(k)} \overline{L}_{0,i}(k),$$

which by (33) verifies

$$|L_1(k)| \geq \frac{\eta h_{n_{k-1}}}{2 h_{n_k}} \frac{|L_0(k)|}{2} \geq \delta h_{n_{k-1}} \frac{\eta}{4}.$$

Notice that by construction,  $L_1(k)$  is the disjoint union of  $\frac{\eta}{2}$  disjoint sets, and each of these sets is contained in a subset of  $h_{n_{k-1}}$  consecutive natural numbers. Moreover, if  $a$  and  $b$  belong to different pieces of this union, then  $|b - a| > h_{n_k}$ .

Repeating the process described above to each of these sets in the disjoint union we can define, recursively, sets  $L_{k-P}(k) \subseteq L_{k-P-1}(k) \subseteq \dots \subseteq L_0(k) \subseteq \{0, \dots, h_{n_k}\}$ , such that, for any  $0 \leq i \leq k - P$ , the set  $L_i(k)$  verifies  $|L_i(k)| \geq \delta h_{n_{k-i}} \left(\frac{\eta}{4}\right)^i$  and it is the disjoint union of (possibly empty)  $\left(\frac{\eta}{2}\right)^i$  sets, each of these contained in a subset of  $h_{n_{k-i}}$  consecutive natural numbers. Moreover, if  $a$  and  $b$  belong to different pieces of this union, there exists  $k - i \leq p \leq k$  such that  $\frac{h_{n_p}}{\eta} \leq |a - b| \leq h_{n_p}$ .

Finally, the claim follows by noticing that by (30) the  $\left(\frac{\eta}{2}\right)^{k-P}$  disjoint sets defining  $L(k) = L_{k-P}(k)$  consist of at most one element. Indeed, if this was not the case, there exist  $a, b \in L(k) \subseteq L_0(k)$  such that  $0 < b - a < h_{n_p}$ . In particular,

$$|F_k(T^a(x)) \cap E| > (1 - \vartheta) |F_k(T^a(x))|, \quad |F_k(T^b(x)) \cap E| > (1 - \vartheta) |F_k(T^b(x))|. \quad (34)$$

If  $T^a(x)$  and  $T^b(x)$  belong to the same tower in the decomposition associated with the renormalization time  $n_k - 2L$ , then, by (31) and (34),  $T^{b-a}(E) \cap E \neq \emptyset$ , which contradicts (30).

Otherwise, if  $T^a(x)$  and  $T^b(x)$  belong to a different tower in this decomposition, by (8) and (17),

$$|T^{b-a}(F_k(T^a(x)) \cap F_k(T^b(x)))| > \frac{1}{C_\gamma(1+\nu)} |F_k(T^b(x))|,$$

and thus, by (31) and (34),  $T^{b-a}(E) \cap E \neq \emptyset$  which again contradicts (30).  $\square$

Recall that either  $[x - \frac{\sigma}{h_{n_k}}, x]$  or  $[x, x + \frac{\sigma}{h_{n_k}}]$  is a continuity interval of  $T^{h_{n_k}}$ . Let us denote by  $J_x$  this continuity interval. Notice that, since  $f \in C_{m,M}$  is piece-wise constant and  $\bigsqcup_{i=0}^{h_{n_k}-1} T^i(J_x)$  is a disjoint union of intervals, we have

$$|S_i f(y) - S_i f(y')| < mM, \quad \text{for any } y, y' \in J_x \text{ and } 0 \leq i < h_{n_k}. \quad (35)$$

Moreover, for any  $i, j \in L(k)$  with  $i < j$ , there exists  $y \in T^i(J_x) \cap E$  such that  $T^{j-i}(y) \in E$ . Indeed, by the definition of  $L(k)$ , we have

$$\begin{aligned} |T^{j-i}(T^i(J_x \cap E)) \cap E| &= |T^i(J_x \cap E)| - |T^{j-i}(T^i(J_x \cap E)) \setminus E| \\ &\geq |T^i(J_x \cap E)| - |F_k(T^j(x)) \setminus E| \\ &\geq |T^i(J_x \cap E)| - \vartheta |F_k(T^j(x))| \\ &\geq |T^i(J_x)| - \vartheta |F_k(T^i(x))| - \vartheta |F_k(T^j(x))| \\ &\geq \frac{\sigma}{h_{n_k}} - 2\vartheta \max_{\beta \in \mathcal{A}} \lambda_\beta^{(n_k-2L)}. \end{aligned}$$

Since  $\sum_{\beta \in \mathcal{A}} \lambda_\beta^{(n_k-2L)} q_\beta^{(n_k-2L)} = 1$  and  $q^{(n_k)} = A_\gamma^2 q^{(n_k-2L)}$ , by (8),

$$\max_{\beta} \lambda_\beta^{(n_k-2L)} \leq \frac{(1+\nu)\|A_\gamma\|^2}{h_{n_k}}.$$

Hence, by (31),

$$|T^{j-i}(T^i(J_x \cap E)) \cap E| \geq \frac{\sigma - 2\vartheta(1+\nu)\|A_\gamma\|^2}{h_{n_k}} \geq \frac{3\sigma}{4h_{n_k}}.$$

and thus there exists  $y \in T^i(J_x) \cap E \cap T^{i-j}(E)$ .

Therefore, if we assume that the conclusions of Proposition 4.2 are false, it follows that

$$|S_{j-i} f(y)| > D,$$

for any  $i, j \in L(k)$  with  $i < j$  and any  $y \in T^i(J_x) \cap E \cap T^{i-j}(E)$ . In view of (35), this implies that

$$|S_{j-i} f(y)| > D - mM,$$

for any  $i, j \in L(k)$  with  $i < j$  and any  $y \in T^i(J_x)$ . In particular, letting  $i_k := \min(L(k))$ ,

$$|S_{j-i_k} f(T^{i_k}(z)) - S_{i-i_k} f(T^{i_k}(z))| = |S_{j-i} f(T^i(z))| > D - mM.$$

for any  $i, j \in L(k)$  with  $i_k < i < j$  and any  $z \in J_x$ .

Denote  $D' := D - mM$ . The previous equation implies

$$\left| \bigsqcup_{i \in L(k) \setminus \{i_k\}} \left( S_{i-i_k} f(T^{i_k}(x)) - \frac{D'}{2}, S_{i-i_k} f(T^{i_k}(x)) + \frac{D'}{2} \right) \right| \geq D'(|L(k)| - 1).$$

In particular, there exists  $i \in L \setminus \{i_k\}$  such that

$$|S_{i-i_k} f(T^{i_k}(x))| \geq \frac{D'}{2}(|L(k)| - 1) \geq \frac{D'}{2} \left( \delta \left( \frac{n}{4} \right)^{k-P} - 1 \right).$$

Hence

$$\max \{|S_{i_k} f(x)|, |S_i f(x)|\} \geq \frac{D'}{4} \left( \delta \left( \frac{n}{4} \right)^{k-P} - 1 \right).$$

On the other hand, it follows from Theorem 4.2 that

$$\max \{|S_{i_k} f(x)|, |S_i f(x)|\} \leq C_0 h_{n_k}^{\frac{\theta_2}{\theta_1} + \varsigma}.$$

where  $C_0$  is a positive constant depending only on  $T, f, \varsigma$ . The last two equations imply that

$$C_0 h_{n_k}^{\frac{\theta_2}{\theta_1} + \varsigma} \geq \frac{D'}{2} \left( \delta \left( \frac{\eta}{4} \right)^{k-P} - 1 \right).$$

Since the construction above can be done for any  $k > P$ , taking  $k$ -th root and making  $k$  go to infinity, Proposition 4.1 yields

$$(C\eta)^{(1+\epsilon)\left(\frac{\theta_2}{\theta_1} + \varsigma\right)} \geq \frac{\eta}{4}.$$

Thus, by (28),

$$C\eta^{1-\varsigma} \geq \frac{\eta}{4},$$

which contradicts (29).  $\square$

Theorem 1.2 now follows easily from Propositions 4.1 and 4.2.

## APPENDIX A.

This section gives a detailed proof of the first step in the proof of Theorem 1.1, namely, the following.

**Proposition A.1.** *Let  $M > 0$ ,  $m \in \mathbb{N}$  and  $d \geq 2$  be fixed. Given a minimal IET  $T$  on  $d$  intervals and a positive measure set  $W \subseteq C_{m,M}$  such that for any  $f \in W$  the associated skew-product  $T_f$  given by (2) is not ergodic w.r.t. to the Lebesgue measure, there exists a positive measure subset  $V \subseteq W$  and a measurable function  $f \mapsto \mu_f$  from  $V$  to the space of Radon measures on  $[0, 1) \times \mathbb{R}$  such that*

- $\mu_f$  is invariant by  $T_f$ ;
- $\mu_f \neq c \text{Leb}_{[0,1) \times \mathbb{R}}$ , for any  $c \in \mathbb{R} \setminus \{0\}$ ;
- $\mu_f \not\perp \text{Leb}_{[0,1) \times \mathbb{R}}$ .

Given  $T$  and  $f$  as above we will construct the measure  $\mu_f$  by using the ergodic decomposition of the measure  $\text{Leb}_{[0,1] \times [-1/2, 1/2]}$  with respect to the first return map of  $T_f$  to the set  $[0, 1) \times [-1/2, 1/2]$ . Let us denote the first return map by  $\tilde{T}_f$ . Recall first, that due to Atkinson [2] we know that if  $T$  is ergodic, almost every point is *recurrent*, that is, for every  $\epsilon > 0$ , there exists arbitrarily large  $N > 0$  such that  $|S_N f(x)| < \epsilon$ . In particular, the map  $\tilde{T}_f : [0, 1] \times [-1/2, 1/2] \rightarrow [0, 1] \times [-1/2, 1/2]$  is well-defined and preserves  $\text{Leb}_{[0,1] \times [-1/2, 1/2]}$ .

For any  $f \in V$ , let us consider the ergodic decomposition of  $\text{Leb}_{[0,1] \times [-1/2, 1/2]}$  with respect to  $\tilde{T}_f$ :

$$\text{Leb}_{[0,1] \times [-1/2, 1/2]} = \int_{[0,1] \times [-1/2, 1/2] \setminus \text{Inv}(\tilde{T}_f)} p_{\bar{x}} d\rho_f(\bar{x}), \quad (36)$$

where  $\rho_f$  is a probability measure on the space of ergodic components  $[-1/2, 1/2] \times \mathbb{R} \setminus \text{Inv}(\tilde{T}_f)$  and  $p_{\bar{x}}$  is an ergodic measure with respect to  $\tilde{T}_f$  (see [15]).

If  $\tilde{T}_f$  is ergodic then the space of invariant components is trivial and hence it is one point, while if it is not ergodic, then the space is expected to be uncountable. Thus, to get a decomposition of  $\text{Leb}_{[0,1] \times [-1/2, 1/2]}$  that will allow us to measurably assign a measure to a point and a cocycle, we need to consider a different decomposition.

Since  $(\tilde{T}_f, p_{\bar{x}})$  is ergodic for almost every  $\bar{x} \in [-1/2, 1/2] \times \mathbb{R} \setminus \text{Inv}(\tilde{T}_f)$ , in view of Birkhoff Ergodic Theorem,  $p_{\bar{x}}$ -almost every point is generic, that is the set of points  $x \in [-1/2, 1/2] \times \mathbb{R}$  such that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \delta_{\tilde{T}_f^i x} = p_{\bar{x}}$$

in the weak-\* topology. In particular, almost every point with respect to  $\text{Leb}_{[0,1] \times [-1/2, 1/2]}$  is generic for the measure  $p_{\bar{x}}$  for some unique  $\bar{x} \in [0, 1] \times [-1/2, 1/2] \setminus \text{Inv}(\tilde{T}_f)$ . Let us denote this measure by  $p_{x,f}$ . We have the following result.

**Lemma A.1.** *We have the following decomposition:*

$$\text{Leb}_{[0,1] \times [-1/2, 1/2]} = \int_{[0,1] \times [-1/2, 1/2]} p_{x,f} dx.$$

*Proof.* By (36) we have

$$\begin{aligned} \text{Leb}_{[0,1] \times [-1/2, 1/2]} &= \int_{[0,1] \times [-1/2, 1/2] \setminus \text{Inv}(\tilde{T}_f)} p_{\bar{x}} d\rho_f(\bar{x}) \\ &= \int_{[0,1] \times [-1/2, 1/2] \setminus \text{Inv}(\tilde{T}_f)} \int_{[0,1] \times [-1/2, 1/2]} p_{\bar{x}} dp_{\bar{x}} d\rho_f(\bar{x}) \\ &= \int_{[0,1] \times [-1/2, 1/2] \setminus \text{Inv}(\tilde{T}_f)} \int_{[x|p_{x,f}=p_{\bar{x}}]} p_{\bar{x}} dp_{\bar{x}} d\rho_f(\bar{x}) \end{aligned}$$

Since the set  $[x|p_{x,f}=p_{\bar{x}}]$  coincides with the invariant component  $\bar{x}$  (up to the measure  $p_{\bar{x}}$ ), we get

$$\begin{aligned} \text{Leb}_{[0,1] \times [-1/2, 1/2]} &= \int_{[0,1] \times [-1/2, 1/2] \setminus \text{Inv}(\tilde{T}_f)} \int_{\bar{x}} p_{x,f} dp_{\bar{x}}(x) d\rho_f(\bar{x}) \\ &= \int_{[0,1] \times [-1/2, 1/2]} p_{x,f} d\text{Leb}(x). \end{aligned}$$

□

**Remark A.1.** *By the Birkhoff Ergodic Theorem, the decomposition in the above lemma is trivial, that is every measure in the decomposition is Lebesgue, if and only if  $\tilde{T}_f$  is ergodic, which, by recurrence, is equivalent to the ergodicity of  $T_f$ .*

We need the following, simple lemma on the properties of the set of point convergence.

**Lemma A.2.** *Let  $\{f_n\}_{n \in \mathbb{N}}$  be a sequence of continuous functions on a topological space  $X$  to the complete metric space  $(Y, d)$ . Then the set*

$$C := \{x \in X \mid \lim_{n \rightarrow \infty} f_n(x) \text{ exists}\}$$

*is Borel.*

*Proof.* Since the space  $(Y, d)$  is complete, we have

$$C = \{x \in X \mid \{f_n(x)\}_{n \in \mathbb{N}} \text{ is a Cauchy sequence}\}.$$

Then

$$C = \bigcap_{m=1}^{\infty} \bigcup_{N=1}^{\infty} \bigcap_{k=N}^{\infty} \bigcap_{\ell=N}^{\infty} \{x \in X \mid |f_k(x) - f_{\ell}(x)| < 1/m\},$$

which, together with the assumption on the continuity, finishes the proof.

□

We now turn into one of the main measurability results, which is going to be used to construct proper measure assignments. First we introduce some auxilliary sets. Take

$$D := \{(x, f) \in ([0, 1] \times [-1/2, 1/2]) \times W \mid S_n f(x) \neq \pm 1/2 \text{ for } n \in \mathbb{Z}\}$$

and

$$E := \left\{ (x, f) \in ([0, 1] \times [-1/2, 1/2]) \times W \mid \begin{array}{l} T_f^k(x) \text{ does not belong to the vertical line given} \\ \text{by the discontinuity of } f \text{ or } T \text{ for any } k \in \mathbb{Z} \end{array} \right\}.$$

Note that the complements of these sets are of 0 measure. Indeed, the points that do not belong to one of those sets, satisfy one of countably many linear equations. Note also, that  $E$  and  $F$  sets are defined topologically, not just up zero-measure sets.

**Proposition A.2.** *The assignment  $(x, f) \mapsto p_{x,f}$ , as in Lemma A.1, is measurable with respect to the product topology in the domain and the weak-\* topology in the image.*

*Proof.* Consider the function  $G_n : ([0, 1] \times [-1/2, 1/2]) \times W \rightarrow \mathcal{M}$ , where  $\mathcal{M}$  is the space of probability measures on  $[0, 1] \times [-1/2, 1/2]$ , given by the formula

$$G_n(x, f) := \frac{1}{n} \sum_{i=0}^{n-1} \delta_{\tilde{T}_f^i x}.$$

On  $([0, 1] \times [-1/2, 1/2]) \times W$  we consider the product of Lebesgue measures. We will show that  $G_n$  is continuous on a full measure set.

Take  $(x, f) \in D \cap E$  and let  $(x_m, f_m) \rightarrow (x, f)$  and let  $p_1, \dots, p_m$  be discontinuities of  $f$ . Moreover, let  $s_1, \dots, s_d$  be the discontinuities of  $T$ . We will prove that

$$\lim_{m \rightarrow \infty} G_n(x_m, f_m) = G_n(x, f). \quad (37)$$

Pick any  $\epsilon > 0$  such that

$$\epsilon < \frac{1}{2} \min \left\{ \min_{k=0, \dots, N(x)} (|S_k f(x) - 1/2|), \min_{k=0, \dots, N(x)} (|S_k f(x) + 1/2|), \right. \\ \left. \min_{k=0, \dots, N(x); \ell=1, \dots, m} |T^k(x) - p_\ell|, \min_{k=0, \dots, N(x); j=1, \dots, d} |T^k(x) - s_j| \right\}, \quad (38)$$

where  $N(x)$  is  $n-1$ -th return time of  $x$  to  $[0, 1] \times [-1/2, 1/2]$  via  $T_f$ . Note that by definition of  $D$  and  $E$ , such  $\epsilon$  exists. Take  $M \in \mathbb{N}$  such that for every  $m \geq M$  we have

$$\max_{k=0, \dots, N(x)} \{|S_k f(x) - S_k f_m(x_m)|, |T^k(x) - T^k(x_m)|\} < \epsilon.$$

Such  $M$  exists due to (38). In particular, the return times up to  $N(x)$  of iterations of  $T_f$  to  $[0, 1] \times [-1/2, 1/2]$  are identical as those of  $x$ . Hence, the distance of  $\delta_{\tilde{T}_f^i x}$  and  $\delta_{\tilde{T}_f^i x_m}$  in Levy-Prokhorov metric is less than  $\epsilon$ , for every  $m > M$ . This finishes the proof of continuity of  $G_n$  on  $D \cap E$ . It follows, by Lemma A.2, that the set  $\Omega$  of points  $(x, f) \in D \cap E$  such that  $x$  is generic for some  $T_f$ -invariant measure is measurable. Since for any  $f \in W$ , the set of points  $x$  which are generic for  $T_f$  is of full measure, the set  $\Omega$  is of full product measure.

To conclude the proof of the proposition, it suffices to notice that  $\lim_{n \rightarrow \infty} G_n(x, f) = p_{x,f}$  for every  $(x, f) \in \Omega$ . In other words  $(x, f) \mapsto p_{x,f}$  on  $\Omega$  is a point-wise limit of continuous maps, thus it is well-defined and measurable on a full measure set.  $\square$

The decomposition obtained in the above proposition is not done on the whole space but rather on a subset of  $[0, 1] \times \mathbb{R}$ . We now show how to obtain a measure which is absolutely continuous with respect to the Lebesgue measure, using the aforementioned decomposition.

For every  $f \in W$ , consider a dynamical system  $T_f^N : [0, 1] \times [-N, N]$ , which is the first return map of  $T_f$  to  $[0, 1] \times [-N, N]$ . Note that the map  $\tilde{T}_f : [0, 1] \times [-1/2, 1/2]$  can be viewed as the first return map of  $\tilde{T}_f^N$  to  $[0, 1] \times [-1/2, 1/2]$ . Denote by  $\mathcal{M}^N$  the set of finite Radon measures on  $[0, 1] \times [-N, N]$ . We have the following corollary of Proposition A.2.

**Lemma A.3.** *There exists a measurable assignment*

$$([0, 1] \times [-N, N]) \times W \ni (x, f) \rightarrow p_{x,f}^N \in \mathcal{M}^N,$$

where

$$\mu_N^f := \int_{[0,1] \times [-N,N]} p_{x,f}^N dx \ll \text{Leb}_{[0,1] \times [-N,N]}$$

the integrated measures are invariant and ergodic. In particular, for every  $N \geq 1$  we have

$$p_{x,f}^N|_{[0,1] \times [-1/2, 1/2]} = p_{x,f}$$

and if  $N_1 < N_2$ , then

$$p_{x,f}^{N_2}|_{[0,1] \times [-N_1, N_1]} = p_{x,f}^{N_1}.$$

*Proof.* Let  $N \in \mathbb{N}$  and for every  $m \in \mathbb{N}$  let

$$U_m := \{x \in [0, 1] \times [-1/2, 1/2] \mid m \text{ is the first return time of } x \text{ via } T_f^N \text{ to } [0, 1] \times [-1/2, 1/2]\}.$$

Note that the first return map to  $[0, 1] \times [-1/2, 1/2]$  via  $T_f^N$  is equal  $\tilde{T}_f$ . Consider the measure on  $[0, 1] \times [-1/2, 1/2]$  given by

$$\mu_f^N := \sum_{m=1}^{\infty} \sum_{j=0}^{m-1} (T_f^N)_* (\text{Leb}_{[0,1] \times [-1/2, 1/2]})|_{U_m}.$$

Since each of the summands is absolutely continuous w.r.t. Lebesgue measure, then so is  $\mu_f^N$ . By Lemma A.1 we have

$$\mu_f^N = \int_{[0,1] \times [-1/2, 1/2]} \sum_{m=1}^{\infty} \sum_{j=0}^{m-1} (T_f^N)_*^j p_{x,f}|_{U_m} dx$$

Take

$$p_{x,f}^N := \sum_{m=1}^{\infty} \sum_{j=0}^{m-1} (T_f^N)_*^j p_{x,f}|_{U_m}.$$

Since the restriction of measures and taking images of measures are measurable operations, by Proposition A.2 we obtain the measurability of the assignment  $(x, f) \mapsto p_{x,f}^N$ . The invariance and ergodicity of  $p_{x,f}^N$  follows directly from invariance and ergodicity of  $p_{x,f}$ , while the last two equations follow from the construction.  $\square$

**Remark A.2.** *With the notation of the above proof, we actually have that  $\mu_f^N$  is a restriction of the Lebesgue measure on the strip to the disjoint union  $\bigcup_{m=1}^{\infty} \bigcup_{j=0}^{m-1} (T_f^N)^j(U_m)$ . This does **not** have to be the whole measure  $\text{Leb}_{[0,1] \times [-N, N]}$ .*

In the following result we pass from the decomposition on the bounded subsets to the decomposition on the whole strip.

**Proposition A.3.** *There exists a measurable assignment*

$$([0, 1] \times [-1/2, 1/2]) \times W \ni (x, f) \mapsto \tilde{p}_{x,f} \in \mathcal{M}_R,$$

where  $\mathcal{M}_R$  is the space of Radon measures on  $[0, 1] \times \mathbb{R}$  such that

- for a.e every  $(x, f) \in ([0, 1] \times [-1/2, 1/2]) \times W$  with respect to the product Lebesgue measure, the measure  $\tilde{p}_{x,f}$  is  $T_f$ -invariant and  $\tilde{p}_{x,f}|_{[0,1] \times [-1/2, 1/2]} = p_{x,f}$ ,
- $\hat{\mu}_f := \int_{[0,1] \times [-1/2, 1/2]} \tilde{p}_{x,f} dx$  is well defined and  $\hat{\mu}_f \ll \text{Leb}_{[0,1] \times \mathbb{R}}$ .

*Proof.* For every  $x \in [0, 1] \times [-1/2, 1/2]$  and  $f \in W$  consider the sequence of measures  $\{p_{x,f}^N\}$  given by the Lemma A.3, here seen as measures on  $[0, 1] \times \mathbb{R}$ . Define the measure  $\tilde{p}_{x,f}$  by putting for every compact subset  $K \subseteq [0, 1] \times \mathbb{R}$

$$\tilde{p}_{x,f}(K) = p_{x,f}^{N_K}(K), \text{ where } N_K := \min\{N \in \mathbb{N} \mid K \subseteq [0, 1] \times [-N, N]\}.$$

Note that in view of Lemma A.3, we can replace in the above definition  $N_K$  by any  $N$  bigger than  $N_K$ . Hence the above equation really defines a measure.

We claim that

$$\lim_{N \rightarrow \infty} p_{x,f}^N = \tilde{p}_{x,f} \tag{39}$$

exists. Indeed, if  $g$  is a continuous function on  $[0, 1] \times \mathbb{R}$  with a compact support  $K_g$ , then there exists  $N_g \in \mathbb{N}$ , such that  $K_g \subseteq [0, 1] \times [-N_g, N_g]$ . Then, by Lemma A.3 we get

$$\lim_{N \rightarrow \infty} \int_{[0,1] \times \mathbb{R}} g dp_{x,f}^N = \int_{[0,1] \times [-N_g, N_g]} g dp_{x,f}^{N_g} = \int_{[0,1] \times [-N_g, N_g]} g d\tilde{p}_{x,f} = \int_{[0,1] \times \mathbb{R}} g d\tilde{p}_{x,f}.$$

Note that the assignment  $P(x, f) := \tilde{p}_{x,f}$  is measurable. Indeed, recall that the topology of  $\mathcal{M}_R$  is generated by the sets of the form  $C(\nu, f_1, \dots, f_n, \epsilon_1, \dots, \epsilon_n) = \{\rho \in \mathcal{M}_R \mid |\int f_i d\rho - \int f_i d\nu| < \epsilon_i\}$ ,



where  $n \in \mathbb{N}$ ,  $\nu \in \mathcal{M}_R$ ,  $\epsilon_1, \dots, \epsilon_n > 0$  and  $f_1, \dots, f_n \in C_c([0, 1] \times \mathbb{R})$ . Then, there exists  $N \in \mathbb{N}$  such that the supports of functions  $f_1, \dots, f_n$  are included in  $[0, 1] \times [-N, N]$ . Thus

$$\begin{aligned} & \{(x, f) \in ([0, 1] \times [-1/2, 1/2]) \times W \mid \left| \int f_i d\tilde{p}_{x,f} - \int f_i d\nu \right| < \epsilon_i\} = \\ & \{(x, f) \in ([0, 1] \times [-1/2, 1/2]) \times W \mid \left| \int f_i dp_{x,f}^N - \int f_i d\nu \right| < \epsilon_i\}. \end{aligned}$$

Since the assignment  $(x, f) \mapsto p_{x,f}^N$  is measurable, so is the above set. By the fact that all parameters  $\nu$ ,  $f_j$  and  $\epsilon_j$  are arbitrary, it follows that  $P$  is measurable.

In a similar fashion, if  $A \subseteq [0, 1] \times \mathbb{R}$  is a bounded set, then since  $f$  is bounded, there exists  $N \in \mathbb{N}$  such that  $A, T_f A \subseteq [0, 1] \times [-N, N]$ . Since  $p_{x,f}^N$  is  $T_f^N$ -invariant, then  $\tilde{p}_{x,f}(A) = \tilde{p}_{x,f}(T_f(A))$ . Hence the measure  $\tilde{p}_{x,f}$  is  $T_f$  invariant.

Finally, it remains to notice that by Lemma A.3, we have for every  $N \in \mathbb{N}$  that

$$\mu_f^N = \int_{[0,1] \times [-1/2, 1/2]} p_{x,f}^N dx \ll \int_{[0,1] \times [-N, N]} p_{x,f}^N dx = \text{Leb}_{[0,1] \times [-N, N]}. \quad (40)$$

Then by (39) we get that  $\hat{\mu}_f := \lim_{N \rightarrow \infty} \mu_f^N$  is well defined,  $\hat{\mu}_f|_{[0,1] \times [-N, N]} = \mu_f^N$  and by (40), we have that  $\hat{\mu}_f \ll \text{Leb}_{[0,1] \times \mathbb{R}}$ .  $\square$

**Remark A.3.** *It is worth to mention that the measure  $\hat{\mu}_f$  obtained in the above proposition does not need to be the whole Lebesgue measure on the strip  $[0, 1] \times \mathbb{R}$ . Indeed if the function  $f$  is a bounded coboundary then the support of  $\hat{\mu}_f$  is contained in the set  $[0, 1] \times [-\|f\|_\infty - 1, \|f\|_\infty + 1]$  (see also Remark A.2).*

One can observe that until now we did not use the assumption on non-ergodicity of considered transformations. Indeed, this assumption is only used in the following proof of the Proposition A.1.

*Proof of Proposition A.1.* Consider the assignment  $(x, f) \mapsto \tilde{p}_{x,f}$  given by Proposition A.3 and denote it by  $P : ([0, 1] \times [-1/2, 1/2]) \times W \rightarrow \mathcal{M}_R$ . Since it is a measurable map, by Lusin's Theorem, there exists a compact subset  $K \subseteq ([0, 1] \times [-1/2, 1/2]) \times W$  of positive measure such that  $P|_K$  is continuous. Fix  $(x_0, f_0) \in K$ , a point of density. Since  $f_0 \in W$ , we have that  $\tilde{p}_{x_0, f_0}|_{[0,1] \times [-1/2, 1/2]} \neq \text{Leb}_{[0,1] \times [-1/2, 1/2]}$ . In particular, if  $d_{LP}$  denotes the Levy-Prokhorov metric on  $\mathcal{M}$  (not  $\mathcal{M}_R$ !), we have that there exists  $\epsilon > 0$  such that  $\text{Leb}_{[0,1] \times [-1/2, 1/2]} \notin B_{LP}(\tilde{p}_{x_0, f_0}|_{[0,1] \times [-1/2, 1/2]}, \epsilon)$ . Since  $P|_K$  is continuous, there exists  $\delta > 0$  such that for every  $(x, f) \in K \cap D((x_0, f_0), \delta) =: \bar{K}$ , we have  $d_{LP}(\tilde{p}_{x_0, f_0}|_{[0,1] \times [-1/2, 1/2]}, \tilde{p}_{x,f}|_{[0,1] \times [-1/2, 1/2]}) < \epsilon/2$ . Since  $(x_0, f_0)$  is a density point of  $K$ , we have  $\text{Leb}_{[0,1] \times [-1/2, 1/2]}(\bar{K}) > 0$ .

Let  $V \subseteq W$  be the set of those elements  $f$  for which sets  $K_f := \{x \in [0, 1] \times [-1/2, 1/2] \mid (x, f) \in \bar{K}\}$  have positive measure. Since  $\bar{K}$  is of positive measure, then so is  $V$ . Consider the map

$$V \ni f \mapsto \bar{\mu}_f := \int_{K_f} \tilde{p}_{x,f} dx.$$

Note that it is actually the integration of  $P|_{\bar{K}}$ , a continuous function, with respect to the  $x$ -coordinate. In particular, it is continuous. Let  $\tilde{\mu}_f := \frac{1}{\text{Leb}(K_f)} \bar{\mu}_f$ . Note that  $\text{Leb}(K_f) = \bar{\mu}_f([0, 1] \times [-1/2, 1/2])$ . Since the map  $\mathcal{M} \ni \nu \mapsto \nu([0, 1] \times [-1/2, 1/2])$  is measurable, then so is  $V \ni f \mapsto \tilde{\mu}_f \in \mathcal{M}$ .

Moreover, since  $\bar{K} \subseteq D((x_0, f_0), \delta)$ , by the choice of  $\delta$  we have that measure  $\tilde{\mu}_f$  is a convex combination of measures from the set  $B_{LP}(p_{x_0, f_0}, \epsilon/2)$  and as such,  $\mu_f \in B_{LP}(p_{x_0, f_0}, \epsilon/2)$ . In particular

$$\tilde{\mu}_f|_{[0,1] \times [-1/2, 1/2]} \neq \text{Leb}_{[0,1] \times [-1/2, 1/2]},$$

hence

$$\tilde{\mu}_f \neq \text{Leb}_{[0,1] \times \mathbb{R}}.$$

Moreover,

$$\tilde{\mu}_f = \frac{1}{\text{Leb}(K_f)} \int_{K_f} \tilde{p}_{x,f} dx \ll \int_{[0,1] \times [-1/2, 1/2]} \tilde{p}_{x,f} dx = \hat{\mu}_f \ll \text{Leb}_{[0,1] \times \mathbb{R}},$$

where  $\hat{\mu}_f$  is given by Proposition A.3. This finishes the proof.  $\square$

We will now show Lemma A.4 which in turn was used earlier to show that Theorem 1.3 follows from Theorem 1.1. First, we recall some classical facts from the general ergodic theory. Let  $(X, \mathcal{B}, \nu)$  be a standard probability space. We denote by  $\text{Aut}(X)$  the space of all  $\nu$ -measure preserving automorphisms on  $X$ . Let  $\{B_k\}_{k=1}^\infty$  be a countable family of subsets generating  $\mathcal{B}$ . It is a classical fact (see, e.g., [1]), that this space endowed with the metric

$$d_{\text{Aut}(X)}(S_1, S_2) = \sum_{k=1}^{\infty} 2^{-k} (\nu(S_1(B_k) \triangle S_2(B_k)) + \nu(S_1^{-1}(B_k) \triangle S_2^{-1}(B_k))) \quad (41)$$

is a Polish space. Let also  $\text{Erg}(X) \subseteq \text{Aut}(X)$  be a subset of ergodic automorphism. It is a classical fact that it is measurable (see ex. [9]).

We now show that by twitching both the IET and the cocycle, the first return maps to the cylinder in  $[0, 1) \times \mathbb{R}$  obtained in this way are close in an appropriate space of automorphisms.

**Lemma A.4.** *With the notation as in Lemma A.3, the map*

$$((S_0 \times \Lambda^A) \cap \text{Erg}([0, 1))) \times C_{m,M} \ni (T, f) \mapsto \tilde{T}_f^N \in \text{Aut}([0, 1) \times [-N, N], \mathcal{B}, \text{Leb}_{[0,1] \times [-N,N]})$$

*is continuous for every  $N \in \mathbb{R}_{>0}$ .*

*Proof.* We will prove the lemma for  $N = 1/2$ , for other cases the proof differs only by proper rescaling of measures. Let  $(B_k)_{k=1}^\infty$  be a family of rectangles generating the Borel  $\sigma$ -algebra on  $[0, 1) \times [-N, N]$ . We consider the metric  $d := d_{\text{Aut}([0,1] \times [-N,N])}$  as in (41).

Fix  $\pi \in S_0$  and let  $(T_n, f_n)_{n=1}^\infty$  be a sequence converging to  $(T, f)$  in  $\{\pi\} \times \Lambda^A \times C_{m,M}$  with respect to the product metric (on  $\Lambda^A$  we consider standard Euclidean metric), with  $T_n, T$  being ergodic. Denote by  $T_{n,f_n}$  the skew product given by  $(T_n, f_n)$ . Note that due to ergodicity assumption, the first return transformations under consideration are all well defined due to recurrence. Let  $\varepsilon > 0$  and let  $K$  be such that

$$\sum_{i=K+1}^{\infty} 2^{-i+1} \text{Leb}_{[0,1] \times [-1/2, 1/2]}(B_i) < \varepsilon/2.$$

It is enough to show that there exist  $L \in \mathbb{N}$  such that for every  $n \geq L$  we have

$$\sum_{i=1}^K 2^{-i} \left( \text{Leb}_{[0,1] \times [-1/2, 1/2]}(\tilde{T}_{n,f_n}(B_i) \triangle \tilde{T}_f(B_i)) + \text{Leb}_{[0,1] \times [-1/2, 1/2]}(\tilde{T}_{n,f_n}^{-1}(B_i) \triangle \tilde{T}_f^{-1}(B_i)) \right) < \varepsilon/2.$$

On the other hand, to show the above inequality, it is enough to find  $L_i$  for every  $i \in \{1, \dots, K\}$  such that for every  $n \geq L_i$

$$\text{Leb}_{[0,1] \times [-1/2, 1/2]} \left( \tilde{T}_{n,f_n}(B_i) \triangle \tilde{T}_f(B_i) \right) + \text{Leb}_{[0,1] \times [-1/2, 1/2]} \left( \tilde{T}_{n,f_n}^{-1}(B_i) \triangle \tilde{T}_f^{-1}(B_i) \right) < \varepsilon/2K$$

and take  $L := \max_{i=1, \dots, K} L_i$ .

Fix  $i \in \{1, \dots, K\}$  and let  $B := B_i \cap [0, 1) \times [-1/2 + \varepsilon/16K, 1/2 - \varepsilon/16K]$ . By picking sufficiently large  $L_i$ , we obtain a subset  $\hat{B} \subseteq B$  with  $\text{Leb}(B \setminus \hat{B}) < \varepsilon/16K$ , consisting of finitely many rectangles, such that for every  $x \in \hat{B}$  the first return times to  $[0, 1) \times [-1/2, 1/2]$  via  $T_{n,f_n}$  and  $T_f$  as well as by  $T_{n,f_n}^{-1}$  and  $T_f^{-1}$  are identical, for any  $n \geq L_i$ . By increasing  $L_i$  further if necessary, we get that these rectangles are shifted via  $T_{n,f_n}^{\pm 1}$  for every  $n \geq L_i$ , and via  $T_f^{\pm 1}$  by an arbitrary small amount. In particular, for  $L_i$  large enough, we get

$$\text{Leb}_{[0,1] \times [-1/2, 1/2]} \left( \tilde{T}_{n,f_n}(\hat{B}) \triangle \tilde{T}_f(\hat{B}) \right) + \text{Leb}_{[0,1] \times [-1/2, 1/2]} \left( \tilde{T}_{n,f_n}^{-1}(\hat{B}) \triangle \tilde{T}_f^{-1}(\hat{B}) \right) < \varepsilon/8K.$$

Hence

$$\begin{aligned} & \text{Leb}_{[0,1] \times [-1/2, 1/2]} \left( \tilde{T}_{n,f_n}(B_i) \triangle \tilde{T}_f(B_i) \right) + \text{Leb}_{[0,1] \times [-1/2, 1/2]} \left( \tilde{T}_{n,f_n}^{-1}(B_i) \triangle \tilde{T}_f^{-1}(B_i) \right) \\ & \leq \text{Leb}_{[0,1] \times [-1/2, 1/2]} \left( \tilde{T}_{n,f_n}(B) \triangle \tilde{T}_f(B) \right) + \text{Leb}_{[0,1] \times [-1/2, 1/2]} \left( \tilde{T}_{n,f_n}^{-1}(B) \triangle \tilde{T}_f^{-1}(B) \right) + \varepsilon/4K \\ & \leq \left( \tilde{T}_{n,f_n}(\hat{B}) \triangle \tilde{T}_f(\hat{B}) \right) + \text{Leb}_{[0,1] \times [-1/2, 1/2]} \left( \tilde{T}_{n,f_n}^{-1}(\hat{B}) \triangle \tilde{T}_f^{-1}(\hat{B}) \right) + 3\varepsilon/8K \leq \varepsilon/2K, \end{aligned}$$

which finishes the proof.  $\square$

## REFERENCES

- [1] AARONSON, J. *An introduction to infinite ergodic theory*, vol. 50 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [2] ATKINSON, G. Recurrence of Co-Cycles and Random Walks. *Journal of the London Mathematical Society* *s2-13*, 3 (1976), 486–488. \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1112/jlms/s2-13.3.486>.
- [3] AVILA, A., AND VIANA, M. Simplicity of Lyapunov spectra: proof of the Zorich-Kontsevich conjecture. *Acta Mathematica* *198*, 1 (2007), 1–56. Publisher: Institut Mittag-Leffler.
- [4] CHAIKA, J., AND ROBERTSON, D. Ergodicity of skew products over linearly recurrent IETs. *Journal of the London Mathematical Society* *100*, 1 (2019), 223–248. \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1112/jlms.12210>.
- [5] CONZE, J.-P., AND FRĄCZEK, K. Cocycles over interval exchange transformations and multivalued Hamiltonian flows. *Advances in Mathematics* *226*, 5 (Mar. 2011), 4373–4428.
- [6] CONZE, J.-P., AND PIĘKNIEWSKA, A. On multiple ergodicity of affine cocycles over irrational rotations. *Israel Journal of Mathematics* *201*, 2 (Apr. 2014), 543–584.
- [7] FORNI, G. Deviation of Ergodic Averages for Area-Preserving Flows on Surfaces of Higher Genus. *Annals of Mathematics* *155*, 1 (2002), 1–103. Publisher: Annals of Mathematics.
- [8] FRĄCZEK, K., AND ULCIGRAI, C. Non-ergodic  $\mathbb{Z}$ -periodic billiards and infinite translation surfaces. *Inventiones mathematicae* *197*, 2 (Aug. 2014), 241–298.
- [9] HALMOS, P. R. *Lectures on ergodic theory*. Chelsea Publishing Co., New York, 1960.
- [10] HELLEKALEK, P., AND LARCHER, G. On the ergodicity of a class of skew products. *Israel Journal of Mathematics* *54*, 3 (Oct. 1986), 301–306.
- [11] KEANE, M. Interval exchange transformations. *Mathematische Zeitschrift* *141*, 1 (Feb. 1975), 25–31.
- [12] MASUR, H. Interval Exchange Transformations and Measured Foliations. *Annals of Mathematics* *115*, 1 (1982), 169–200. Publisher: Annals of Mathematics.
- [13] OREN, I. Ergodicity of cylinder flows arising from irregularities of distribution. *Israel Journal of Mathematics* *44*, 2 (June 1983), 127–138.
- [14] PASK, D. A. Skew products over the irrational rotation. *Israel Journal of Mathematics* *69*, 1 (Feb. 1990), 65–74.
- [15] ROHLIN, V. A “general” measure-preserving transformation is not mixing. *Doklady Akad. Nauk SSSR (N.S.)* *60* (1948), 349–351.
- [16] VEECH, W. A. Gauss Measures for Transformations on the Space of Interval Exchange Maps. *Annals of Mathematics* *115*, 2 (1982), 201–242. Publisher: Annals of Mathematics.
- [17] VIANA, M. Ergodic Theory of Interval Exchange Maps. *Revista Matemática Complutense* *19*, 1 (Apr. 2006), 7–100. Number: 1.
- [18] YOCOZ, J.-C. Interval exchange maps and translation surfaces. In *Homogeneous flows, moduli spaces and arithmetic*, vol. 10 of *Clay Math. Proc.* Amer. Math. Soc., Providence, RI, 2010, pp. 1–69.
- [19] ZORICH, A. Finite Gauss measure on the space of interval exchange transformations. Lyapunov exponents. *Annales de l’Institut Fourier* *46*, 2 (1996), 325–370.
- [20] ZORICH, A. Deviation for interval exchange transformations. *Ergodic Theory and Dynamical Systems* *17*, 6 (Dec. 1997), 1477–1499. Publisher: Cambridge University Press.