Healthy Horndeski gravities with torsion

S. Mironov^{1, 2, 3, *} and M. Valencia-Villegas^{2, †}

¹Institute for Nuclear Research of the Russian Academy of Sciences,

60th October Anniversary Prospect, 7a, 117312 Moscow, Russia

²Institute for Theoretical and Mathematical Physics, MSU, 119991 Moscow, Russia

³NRC, "Kurchatov Institute", 123182, Moscow, Russia

We show that the full Horndeski theory with both curvature and torsion can support nonsingular, stable and subluminal cosmological solutions at *all times*. Thus, with torsion, the usual No-Go theorem that holds in a curved spacetime is avoided. In particular, it is essential to include the nonminimal derivative couplings of the \mathcal{L}_5 part of the Horndeski action $(G^{\mu\nu} \nabla_{\mu} \nabla_{\nu} \phi, \text{ and } (\nabla^2 \phi)^3)$. Without the latter a No-Go already impedes the eternal subluminality of nonsingular, stable cosmologies.

I. INTRODUCTION

Modifications of General Relativity (GR) with scalar fields have been widely investigated mainly motivated by the need to address the singularity issues of GR, the Dark Energy and Dark Matter problems, and the recent phenomenological advances which promise guidelines and constraints to the vast theoretical possibilities [1].

In this work we consider the most general modification of GR with second derivatives of a scalar in the action, but with second order equations of motion, which is known as Horndeski theory [2–6]. It reunites under one class a wide variety of models, ranging from the cosmological constant, to k-essence and more generally, minimally coupled and non-minimally coupled scalars, such as Brans-Dicke (See for instance [7] for a review).

The main attractive feature of Horndeski theory -later rediscovered as Galileons [3–6]- is that it permits to violate without obvious pathologies the Null Energy Condition (NEC)¹, upon which the singularity theorems of Penrose and Hawking hold [9, 10]. Indeed it has been proven possible to build *locally* stable and subluminal *bouncing* cosmologies, which however, generally have an issue with *global* stability [11–18]. In other words, even if the solutions are *locally* healthy around the bounce - or the most physically relevant phase - an instability will certainly happen at some time earlier or later in the evolution of the Universe [19]. The certainty of instabilities at some time is what we call *global* stability issues, which were established as No-Go theorems in [13, 14] for the case without torsion. In the case with torsion, the certainty of instabilities *or* an eventual superluminality of the graviton was proven in [20] for a restricted class of theories. We refer to the latter also as a "global stability" issue, because having "stability" would force an arguably unphysical and *potentially* unacceptable period of superluminality [21–23].

All in all, it has been argued that the global stability issues can be postponed and cured with other types of matter that could become relevant at other phases [19]. However, there are generalizations of the No-Go theorems that hold for even more general modifications of GR [17], and furthermore, some analysis have suggested that violating the NEC potentially brings issues of some sort such as superluminality, even if singularities are successfully avoided [22]. Thus, the global stability seems indeed a pressing issue that has to be solved concretely. Among the possible options, one is to consider altogether other theories such as Beyond Horndeski [24, 25]. In this case the equations of motion are of higher than second order, but there is no Ostrogradsky ghost by construction. Even for Horndeski theory there are very particular solutions to the global stability issue [14, 18, 26, 27], but one is restricted to one of the following three options: either the model propagates no scalar perturbation about a nonsingular Friedmann-Lemaître-Robertson-Walker (FLRW) background - which may be unsatisfactory because we do expect small deviations from FLRW on cosmological scales - or the scalar perturbation propagates about Minkowski spacetime [26], or one is forced to consider non conventional asymptotics, such as gravity being the strongest force in the past [14, 27]. Finally, substantially reconsidering Horndeski theory, now on a flat spacetime and with extra terms, fully *exchanging* curvature for torsion through the teleparallel connection, the usual No-Go theorems break [28].

We suggest that another concrete solution - on the familiar curved spacetime - may be to simply lift some of the assumptions that are usually taken for Horndeski theory. Indeed, we relax the mathematically and physically unjustified² assumption of considering a spacetime *only* with curvature, as research in Horndeski theory historically

^{*} sa.mironov_1@physics.msu.ru

[†] mvalenciavillegas@itmp.msu.ru

¹ See for instance [8] for a review.

² Indeed, there are motivations - beyond those considered in this paper - that suggest the relevance of torsion [29]. See also the discussions in [30]. In fact, torsion had already been explored as a potential cure for the singularities of GR, but without addressing potential issues with global stability [29, 31–33].

developed. Because, as we show below, assuming vanishing torsion only helps to simplicity, but at the expense of "artificially" enabling the global stability issues. Indeed, it is remarkable that for such a general theory as Horndeski, a No-Go theorem for the linearization holds at all. It may be that in fact that the simplifying assumptions of the spacetime on which Horndeski theory was formulated are too restrictive given that so many criteria - 1) nonsingular, 2) stable, 3) sub/ luminal cosmology- must be satisfied. We take 1) to 3) as a definition of *"healthiness"* for what follows.

This work builds on the results for **up to quartic** Horndeski theory with both curvature and torsion -namely, with up to $(\nabla^2 \phi)^2$ terms in the action- [20], where a No-Go theorem for healthy cosmological evolution was formulated ³. In this paper, it is shown that once we include the more complex \mathcal{L}_5 part of the action -namely, with up to $(\nabla^2 \phi)^3$ terms-, a No-Go theorem cannot hold. We first argue that the pressing issues with the simpler forms of Horndeski theory -including the torsionless [13, 14] and the "up to quartic torsionful theory" in [20]- can be seen as simply accidental to the assumptions: namely, in the familiar Horndeski theory without torsion many of the coefficients of the quadratic action for the graviton are tightly related to the coefficients of the quadratic action for the scalar mode, and this is nothing more than the consequence of having taken *ad-hoc* simplifications. This link between coefficients ends up in the well-known contradictory requirements for the healthiness of the graviton and the scalar mode (No-Go theorems). A simple pragmatical solution is to lift the assumptions: Turning on torsion modifies the action for the graviton and the scalar mode in increasingly *divergent* ways, as more terms of the full Horndeski action are considered. These modifications come from nontrivial tensor and scalar perturbations of torsion, which mix with the metric perturbations. The more mixing, the more the link between the graviton and scalar actions is broken. For instance, in the "up to quartic torsionful theory" in [20] the mixing of more perturbations is enough to break the standard No-Go theorem [13, 14], thus ending up in a weaker form of this No-Go, where one can achieve stability and nonsingular cosmology, but with an arbitrarily short superluminality of the graviton [20]. In this paper we show that if we lift further the "simplicity assumptions" and consider the complete form of Horndeski theory -including the \mathcal{L}_5 part - more torsion perturbations do not decouple, but rather they mix more with the metric perturbations. This ends up untangling the equations that must be imposed for the healthiness of both sectors. They become two sets of very different combinations of the Lagrangian functions \mathcal{L}_2 to \mathcal{L}_5 . Thus, it is easier to independently satisfy these equations not only locally but also without meeting any contradictions at some time in the evolution. Namely, reaching global stability. We show in this work an explicit toy model with globally healthy cosmological solutions.

Furthermore, the gravitational waves in all potentially healthy Horndeski gravities with torsion follow a characteristic dispersion relation that is not common to any other simpler form of Horndeski theory about the spatially flat FLRW background. Their dispersion relation depends non-trivially on the wavelength. This follows because of the more torsion perturbations that do not decouple, which is necessary to avoid the No-Go's.

We proceed as follows: In section II we define the model. In section III A we summarize the gravitational waves' characteristic features in all potentially healthy models.

In section III B we compare the situation regarding the global stability between three forms of Horndeski theory: (i) without torsion, (ii) on a spacetime with torsion but without \mathcal{L}_5 (namely "up to quartic Horndeski-Cartan theory"), and (iii) on a spacetime with torsion with the complete Horndeski action (which we denote as the full Horndeski-Cartan action). We explain how the No-Go holds for the simpler forms of the theory (i),(ii), and how the usual analysis breaks for (iii), thus first suggesting that with the full Horndeski-Cartan action (iii) healthy solutions can be built.

Finally, in section III C we show that a toy model exists, taken from within the general Horndeski-Cartan theories, which has a classical solution that is nonsingular, stable and sub/luminal *at all times*, thus showing that a "No-Go theorem" cannot exist. We finalize with the conclusions in section IV.

In the Appendix VA we report the construction details of the stable toy model, in VB we present details of the notation for torsion, and in VC, VD, VE we show all necessary details of the quadratic actions for the tensor and scalar modes.

II. THE MODEL: HORNDESKI WITH CURVATURE AND TORSION

Horndeski theory is built on top of GR with four general functions G_2 , G_3 , G_4 , G_5 that depend on a real scalar field ϕ and its first derivatives $X = -\frac{1}{2}\partial_{\mu}\phi \partial^{\mu}\phi$. These general functions appear in specific Lorentz invariant combinations of the second derivative term $(\tilde{\nabla}_{\mu}\tilde{\nabla}_{\nu}\phi)^p$ with $p \leq 3$, such that the equations of motion of all the fields are at most of second order, hence avoiding the Ostrogradsky ghost.

³ We consider torsion in the second order, metric formalism. Namely, we assume from the start that the action can be written with a connection that can be expressed in terms of the Christoffel connection plus torsion [34, 35]. See [28, 36–48] for other formalisms and modifications of Horndeski.

If we consider a spacetime with curvature and torsion, the basic block to build the second-order part of the Horndeski action $(\tilde{\nabla}_{\mu}\tilde{\nabla}_{\nu}\phi)$ is not symmetric under the exchange of indices $\mu \leftrightarrow \nu$. Thus, there are more ways to build Lorentz invariant combinations of the type $(\tilde{\nabla}_{\mu}\tilde{\nabla}_{\nu}\phi)^p$ and still keep the second order equations of motion. These multiple choices of Horndeski Lagrangians on a spacetime with torsion may be named by a handful of free parameters⁴, which for different choices may lead to fundamentally different dynamics within the family of Horndeski-Cartan theories [34, 35].

Currently, it is known that the most physically compelling type⁵ of Horndeski-Cartan theories with only G_2 , G_3 , G_4 functions ("up to quartic") eventually suffers of a pathology in the classical solutions at some time in the evolution of the universe, or the graviton needs to become superluminal for at least an arbitrarily short time [20]. More precisely, in up to quartic theories that can be written as the $\mathcal{L}_2 + \mathcal{L}_3 + \mathcal{L}_4$ part of the action (1), namely, with up to $(\tilde{\nabla}_{\mu}\tilde{\nabla}_{\nu}\phi)^2$ Lorentz invariant combinations, it is not possible to obtain a FLRW cosmology that is always subluminal, nonsingular and stable. Denoting torsionful⁶ quantities with a "tilde" (~), and $\partial G_A / \partial X =: G_{A,X}$,

$$S = \int d^4x \left(\mathcal{L}_2 + \mathcal{L}_3 + \mathcal{L}_4 + \mathcal{L}_5 \right), \qquad (1)$$

$$\mathcal{L}_2 = G_2, \tag{2}$$

$$\mathcal{L}_3 = -G_3 \,\nabla_\mu \nabla^\mu \phi \,, \tag{3}$$

$$\mathcal{L}_{4} = G_{4}\tilde{R} + G_{4,X}\left(\left(\tilde{\nabla}_{\mu}\tilde{\nabla}^{\mu}\phi\right)^{2} - \left(\tilde{\nabla}_{\mu}\tilde{\nabla}_{\nu}\phi\right)\tilde{\nabla}^{\nu}\tilde{\nabla}^{\mu}\phi\right).$$

$$\tag{4}$$

In this letter we show that the situation changes drastically if we also consider the remaining structure of Horndeski theory, namely with the G_5 part which contains up to $(\tilde{\nabla}_{\mu}\tilde{\nabla}_{\nu}\phi)^3$ Lorentz invariant combinations.

Specifically, let us take the complete action (1) and, among the many possible terms that a spacetime with curvature and torsion allows, we consider only the following Lorentz contractions of $(\tilde{\nabla}_{\mu}\tilde{\nabla}_{\nu}\phi)^3$ terms

$$\mathcal{L}_{5} = G_{5} \tilde{G}^{\mu\nu} \tilde{\nabla}_{\mu} \tilde{\nabla}_{\nu} \phi - \frac{1}{6} G_{5, X} \left((\tilde{\nabla}_{\mu} \tilde{\nabla}^{\mu} \phi)^{3} + (\tilde{\nabla}_{\nu} \tilde{\nabla}_{\rho} \phi) \left(2 (\tilde{\nabla}^{\mu} \tilde{\nabla}^{\nu} \phi) \tilde{\nabla}^{\rho} \tilde{\nabla}_{\mu} \phi - 3 (\tilde{\nabla}_{\mu} \tilde{\nabla}^{\mu} \phi) \tilde{\nabla}^{\rho} \tilde{\nabla}^{\nu} \phi \right) \right).$$
(5)

Any other choice of contractions of Lorentz indices in both \mathcal{L}_4 and \mathcal{L}_5 leads to fundamentally different dynamics⁷, even if the equations of motion are kept of second order.

The specific subclass (1) - (5) which we have chosen within the family of all Horndeski-Cartan theories will be enough to show that there exist at least some Horndeski-Cartan theory that can support an all time stable, nonsingular and subluminal cosmology.

III. STABLE COSMOLOGIES WITH A DISTINCTIVE GRAVITON

We examine the stability of the perturbations against a spatially flat FLRW background. As in GR there is no dynamical vector perturbation. And similar as in Horndeski without torsion, there are the usual two polarizations of the graviton and a scalar mode. However, their speed and stability are markedly different for the full theory with torsion:

A. Modified graviton

For all actions of the type (1), containing the G_5 structure on a torsionful spacetime, the dispersion relation of gravitational waves (ω) on the FLRW background is such that the speed generally depends on the wavelength⁸,

$$\omega^2 = \frac{f_0 + \vec{p}^2 f_1}{f_2 + \vec{p}^2 f_3} \vec{p}^2 , \qquad (6)$$

⁴ Namely, for Horndeski with torsion with up to \mathcal{L}_4 there are two free parameters [35]. Including \mathcal{L}_5 it is clear that many more free parameters are possible, but it has not been classified.

⁵ Namely, the theory that besides a graviton also propagates a scalar mode with a relativistic dispersion relation. This corresponds to a particular choice of parameters of a two-parameter family of (up to quartic G_4) Horndeski-Cartan theories [34, 35].

 $^{^6}$ In particular, note that the Einstein tensor $\tilde{G}^{\mu\nu}$ is not symmetric due to torsion.

⁷ See for instance [34] for the discussion in \mathcal{L}_4 . In general they may have less constraints, making them algebraically more difficult to analyze. Their analysis is left for future work.

⁸ This is not that surprising given that the cosmological background spontaneously breaks Lorentz invariance. Indeed, this is common in some derivatively coupled scalars, such as in ghost condensates, which usually have a non relativistic scalar dispersion relation [49].

where we have written all the dependance on momentum (\vec{p}) explicitly, and f_0 , f_1 , f_2 , f_3 depend on the Lagrangian functions. They are given in the Appendix VD.

For our analysis it is appropriate to take the short wavelength approximation and consider the speed of gravitational waves as $\vec{p}^2 \to \infty$,

$$c_g^2 = \frac{f_1}{f_3}.$$
 (7)

This peculiar dispersion relation for the graviton prevails for all Horndeski theories with torsion that contain G_5 , irrespectful of the order of contraction of Lorentz indices in any term of the action (1). Indeed, it is useful to notice in Eq. (6), $f_1 \propto f_3 \propto G_5^2$. Thus, we can already conclude that all potentially healthy models of Horndeski on a space-time with Torsion will be characterized by the dispersion relation (6), because they must contain the G_5 part. Indeed, in [20] it was proven that a No-Go theorem holds for the simpler form of Horndeski with torsion without G_5 (See the analysis below).

Let us now see how this new graviton with the distinctive Eq. (6) helps to the stability of Horndeski models in comparison to simpler forms of the theory:

B. Dodging the No-Go

Let us briefly see how the dispersion relation in Eq. (6) comes to be markedly different in comparison to simpler forms of Horndeski theory. It is obtained as follows: Defining torsion as the difference between connections $T^{\rho}_{\mu\nu} = \tilde{\Gamma}^{\rho}_{\mu\nu} - \tilde{\Gamma}^{\rho}_{\nu\mu}$, we note that by its antisymmetry in lower indices it has 24 independent components, of which we count two twocomponent tensor perturbations $T^{(1)}_{ij}$, $T^{(2)}_{ij}$ about the spatially flat FLRW background⁹. Hence, denoting with h_{ij} the tensor perturbation of the metric, the action (1) for all tensor perturbations about the FLRW background (in conformal time), in momentum space takes the form

$$S_{\tau} = \int d\eta d^{3}p \left(b_{1} (\dot{h}_{ij})^{2} + b_{2} \vec{p}^{2} (h_{ij})^{2} + b_{3} (h_{ij})^{2} + \left(c_{1} \vec{p}^{2} (T_{ij}^{(2)})^{2} + c_{2} h_{ij} T_{ij}^{(1)} + c_{3} \dot{h}_{ij} T_{ij}^{(1)} + c_{4} (T_{ij}^{(1)})^{2} \right) + \vec{p}^{2} \left(d_{1} T_{ij}^{(1)} + d_{2} \dot{h}_{ij} + d_{3} h_{ij} \right) T_{ij}^{(2)} + d_{4} \vec{p}^{2} h_{ij} T_{ij}^{(1)} \right),$$
(8)

where b_A , c_A , d_A depend on three background fields: the scale factor of the FLRW metric $a(\eta)$, the Horndeski scalar, which in the context of linearized expressions we also denote as $\phi(\eta)$ and a non-trivial torsion background $x(\eta)$, which, however, can be solved in terms of $a(\eta)$, $\phi(\eta)$ (See Appendices V B 1 and V C).

The situation for the gravitational waves compares as follows between different forms of Horndeski theory:

(i) without torsion,

(ii) on a spacetime with torsion with the action (1) but without \mathcal{L}_5 (which we refer as "up to quartic Horndeski-Cartan theory"), and

(iii) on a spacetime with torsion with the action (1) including \mathcal{L}_5 (which we denote as the full Horndeski-Cartan action¹⁰).

- (i) On a spacetime without torsion only the terms b_A in the first line in (8) contribute. In this case one recognizes that the graviton is massless ($b_3 = 0$) after using the equations for the background fields.
- (ii) On a spacetime with torsion but without \mathcal{L}_5 , the first and second lines in (8) do contribute, but the third does not (namely, $d_A = 0$). Again the graviton turns out to be massless, but its speed is modified due to its coupling c_2 , c_3 to one of the torsion perturbations, $T_{ij}^{(1)}$. The essential aspect is that in this case $T_{ij}^{(2)}$ fully decouples, because $d_1 = d_2 = d_3 = 0$ (namely, its equation gives $c_1 T_{ij}^{(2)} \equiv 0$), and as a consequence the modification to the graviton of the up to quartic Horndeski-Cartan theory is not as marked as in (6), as reported in [20, 34, 35].

⁹ See Appendix VB for more details on our notation for torsion and Appendix VB1 for a detailed decomposition of perturbations and background fields. We follow the same notation as in [34].

¹⁰ Let us however note that one could still consider additional contractions of Lorentz indices.

(iii) On the other hand, for the full Horndeski-Cartan action (1), the third line in (8) also contributes and now $T_{ij}^{(2)}$ no longer decouples. This is the critical difference. Its equation of motion solves torsion as

$$T_{ij}^{(2)} = -\frac{1}{2c_1} \left(d_1 T_{ij}^{(1)} + d_2 \dot{h}_{ij} + d_3 h_{ij} \right).$$
(9)

With this solution back in (8) we now obtain a $-\frac{d_1^2}{4c_1}\vec{p}^2$ momentum contribution to $(T_{ij}^{(1)})^2$, besides the c_4 term. Indeed, now there is a term of the form $-\frac{1}{4c_1}(-4c_1c_4 + \vec{p}^2 d_1^2) (T_{ij}^{(1)})^2$. This lies at the core of the peculiar dispersion relation of the graviton, because $T_{ij}^{(1)}$ also couples to \dot{h}_{ij} and h_{ij} , and, if $d_1 \propto G_5$ is not identically zero, then there will be a $\frac{1}{\mathcal{O}(\vec{p}^2)}$ contribution to the kinetic and gradient terms for the graviton.

Indeed, skipping unimportant details for this discussion, the torsion perturbation $T_{ij}^{(1)}$ can be finally solved in (1) in terms of h_{ij} , \dot{h}_{ij} , and critically, with terms of order $\frac{1}{\mathcal{O}(\vec{p}^2)}$, provided c_1 , c_2 , c_3 are not identically zero,

$$T_{ij}^{(1)} = \frac{1}{f_2 + \vec{p}^2 f_3} \left(\left(2 c_1 c_3 - \vec{p}^2 d_1 d_2 \right) \dot{h}_{ij} + \left(2 c_1 c_2 - \vec{p}^2 (d_1 d_3 - 2 c_1 d_4) \right) h_{ij} \right), \tag{10}$$

with $f_2 = -4 c_1 c_4$, $f_3 = d_1^2$. Finally, using the solutions Eq. (10), (9) we can write the action (8) for the usual graviton with two polarizations, as:

$$S_{\tau} = \int d\eta \, d^3 p \, a^4 \, \left[\frac{1}{2 \, a^2} \, \frac{1}{f_2 + \vec{p}^2 \, f_3} \, \left(\bar{\mathcal{G}}_{\tau} \, (\dot{h}_{ij})^2 - \, \vec{p}^2 \, \bar{F}_{\tau} \, (h_{ij})^2 \right) \right] \tag{11}$$

with,

$$\bar{\mathcal{G}}_{\tau} = \frac{2}{a^2} c_1 \left(c_3^2 - 4 \, b_1 \, c_4 \right), \ \bar{\mathcal{F}}_{\tau}(\vec{p}^{\,2}) = \frac{\bar{f}_0 + \vec{p}^{\,2} \, \bar{f}_1}{f_2 + \vec{p}^{\,2} \, f_3} \tag{12}$$

from which we identify the dispersion relation (6) with $f_0 = \bar{f}_0/\bar{\mathcal{G}}_{\tau}$, $f_1 = \bar{f}_1/\bar{\mathcal{G}}_{\tau}$. It is convenient to also define¹¹

$$\mathcal{G}_{\tau} = \frac{\bar{\mathcal{G}}_{\tau}}{f_2 + \vec{p}^2 f_3}, \ \mathcal{F}_{\tau} = \frac{\bar{\mathcal{F}}_{\tau}}{f_2 + \vec{p}^2 f_3}.$$
 (13)

Now, in most forms of Horndeski theory the stability or no-ghost requirements for both the tensor and scalar sectors are contradictory. The issue is that the coefficients in the action for the tensor and scalar sectors are tightly related. In up to quartic Horndeski-Cartan also the subluminality of the graviton is part of the inconsistent assumptions. Let us see how the tight relation between tensor and scalar actions is broken for the full Horndeski action with torsion (1).

The scalar sector for the action (1) can be brought to a form that is typical of Horndeski theories in the unitary gauge, after integrating out all scalar perturbations of torsion, because they are non dynamical (We show this in detail in the Appendix VE). Namely:

$$\mathcal{S}_{s} = \int \mathrm{d}\eta \,\mathrm{d}^{3}x \,a^{4} \left(-3 \,\frac{\bar{\mathcal{G}}_{\mathcal{S}}}{a^{2}} \,\dot{\psi}^{2} + \frac{\bar{\mathcal{F}}_{\mathcal{S}}}{a^{2}} \,(\partial_{i}\psi)^{2} + 6 \,\frac{\Theta}{a} \,\alpha \,\dot{\psi} + 2 \,\frac{T}{a^{2}} \,\partial_{i}\alpha \,\partial_{i}\psi + 2 \frac{\partial_{i}\partial_{i}B}{a^{2}} \left(a \,\Theta \,\alpha - \,\bar{\mathcal{G}}_{\mathcal{S}} \,\dot{\psi} \right) + \Sigma \,\alpha^{2} \right) \,, \quad (14)$$

where ψ , α and B are scalar perturbations for the metric, and the coefficients $\bar{\mathcal{G}}_{\mathcal{S}}$, $\bar{\mathcal{F}}_{\mathcal{S}}$, Θ , T, Σ , given in the Appendix V E, depend on the backgrounds $a(\eta)$, $\phi(\eta)$, but they do not depend on spatial momentum. Or, using the constraint equation $\alpha = \frac{\bar{\mathcal{G}}_{\mathcal{S}}}{a \Theta} \dot{\psi}$ imposed by the Lagrange multiplier B in Eq. (14), the action for a single dynamical scalar finally reads

$$\mathcal{S}_s = \int \mathrm{d}\eta \,\mathrm{d}^3 x \,a^4 \,\left(\frac{1}{a^2} \,\mathcal{G}_{\mathcal{S}} \,\dot{\psi}^2 - \frac{1}{a^2} \,\mathcal{F}_{\mathcal{S}} \,(\partial_i \psi)^2\right)\,,\tag{15}$$

¹¹ The global factor in the part of the action for the tensor modes, of order $\frac{1}{\mathcal{O}(\vec{p}^{2})}$, may indicate nothing more than that at very high momenta the theory is strongly coupled, which we already expect from this effective theory. However, it is clear that the speed of gravitational waves is well defined in any case. Now, whether this presents a danger in the perturbative expansion falls beyond the reach of the first approximation in this paper, restricted to linear order.



6



(a) N monotonous growing around any zero η_z .

(b) N in between any two zeros η_z . N is continuous in that interval and hence vanishes at some point.



where,

$$\mathcal{G}_{\mathcal{S}} = 3\,\bar{\mathcal{G}}_{\mathcal{S}} + \frac{\bar{\mathcal{G}}_{\mathcal{S}}^2\,\Sigma}{\Theta^2}\,,\ \mathcal{F}_{\mathcal{S}} = \frac{1}{a^2}\frac{\mathrm{d}N}{\mathrm{d}\eta} - \bar{\mathcal{F}}_{\mathcal{S}}\,,\tag{16}$$

with $c_{\mathcal{S}}^2 = \mathcal{F}_{\mathcal{S}}/\mathcal{G}_{\mathcal{S}}$ the speed squared of the scalar mode, and where we have defined

$$N =: \frac{a\,\bar{\mathcal{G}}_{\mathcal{S}}\,T}{\Theta}\,,\tag{17}$$

which is a quantity of major importance for the discussions below.

Now, let us see how the argument on the stability compares between the different forms of Horndeski theory (i) to (iii) defined before: assuming "normal" asymptotics -more precisely, assuming no strong gravity in the asymptotic past or future $\mathcal{F}_{\tau}(\eta) > b_2 > 0$ as $\eta \to \pm \infty$ - the following compelling assumptions (A), (B) are already mutually inconsistent in the case (i) of Horndeski theory without torsion¹²:

- A) nonsingular cosmological solution (a lower bound on the scale factor $a(\eta) > b_1 > 0$),
- B) the graviton and the scalar mode are not ghosts and they suffer no gradient instabilities, $\mathcal{G}_{\tau} > 0$, $\mathcal{F}_{\tau} > 0$, $\mathcal{F}_{\mathcal{S}} > 0$, $\mathcal{G}_{\mathcal{S}} > 0$.

In the case (ii) of torsionful, up to quartic Horndeski theory one finds an inconsistency if together with the assumptions (A, B) we also assume that

C) the graviton is always sub/ luminal $(c_a^2 \leq 1)$.

Indeed, following the argument initially shown in [12–14] and then extended to the case with torsion in [20], we can see the inconsistency in both cases (i), (ii), as follows:

On one hand N must not vanish provided the assumption of a nonsingular cosmology together with the no-ghost condition and sub/luminality of the graviton: namely, in (i) $\bar{\mathcal{G}}_{\mathcal{S}} = T = \mathcal{G}_{\tau}$ and $\bar{\mathcal{F}}_{\mathcal{S}} = \mathcal{F}_{\tau}$, hence $N \propto a \mathcal{G}_{\tau}^2 \neq 0$. And in (ii), $\bar{\mathcal{G}}_{\mathcal{S}} = \mathcal{G}_{\tau} > 0$, $\bar{\mathcal{F}}_{\mathcal{S}} = \mathcal{F}_{\tau}$ and $T = \mathcal{F}_{\tau} (c_g^2 - 2) < 0$ hence $N \propto a \mathcal{G}_{\tau} T \neq 0$.

On the other hand, the stability requirement for the scalar mode $\mathcal{F}_{\mathcal{S}} > 0$ in (B) tells, firstable, that the function N is monotonous increasing $\frac{\mathrm{d}_N}{\mathrm{d}_\eta} > a^2 \mathcal{F}_{\tau} > 0$. This reveals the behavior of N around any isolated zeros of Θ , which we denote as η_z , as in Figure 1a. In particular it implies that in between any two zeros of Θ , N must vanish as in Figure 1b. Secondly, $\mathcal{F}_{\mathcal{S}} > 0$ also tells that the slope of N is bounded from below asymptotically: $\frac{\mathrm{d}_N}{\mathrm{d}_\eta} > b_1^2 b_2 > 0$

¹² In this discussion we remain within the framework of general Horndeski theories not defined by the particular equation $\Theta \equiv 0$, which solves the issues in the torsionless Horndeski theory at the expense of loss of generality, having only two options: either the model propagates no scalar perturbation about a nonsingular FLRW background, or the scalar perturbation propagates about Minkowski spacetime [26].



FIG. 2: N in a semi-infinite interval $(-\infty, \eta_z)$. Similarly N vanishes at some time in an interval (η_z, ∞) or if there are no zeros η_z at any time of the evolution, because N cannot have horizontal asymptotes $\frac{dN}{d\eta} > b_1^2 b_2 > 0$ with b_1 and b_2 positive constants, as $\eta \to \pm \infty$.

as $\eta \to \pm \infty$. This implies that N will also vanish in any infinite interval - even if Θ never vanishes - because N cannot have a horizontal asymptote. See for instance Figure 2 in the case of the left-most zero of Θ . This contradicts the last paragraph, hence, the physically compelling assumptions in each case (i), (ii) are mutually inconsistent. A detailed proof is given in [13, 14, 16, 20].

Finally, in the case (iii) the situation is markedly different. Considering the expression (17) for N, let us see that it can vanish in many ways: first, notice that in general the condition in (B) $\mathcal{G}_{\tau} > 0$ does not constrain the sign of $\bar{\mathcal{G}}_{\mathcal{S}}$, because $\bar{\mathcal{G}}_{\mathcal{S}} \neq \mathcal{G}_{\tau}(\vec{p}^2)^{-13}$. This is obvious because while the latter depends on momentum, the former does not. Moreover, close inspection of the expression for T in Eq. (85) in the Appendix VE, shows that the latter is not related in a simple way to the speed of the graviton -defined only in the short wavelength *approximation* (7)-, which is the key aspect that would usually relate the stability to superluminality in up to quartic Horndeski-Cartan theory (ii). This is not surprising, because the dispersion relation in Eq. (6) depends in a complicated way on momentum, while the coefficient T in the scalar sector is independent of momentum.

Therefore, the usual assumptions to have a classically healthy solution (A)-(C) at some momenta -in particular $\mathcal{G}_{\tau}(\vec{p}^{\,2}) > 0$ and $c_g^2 \leq 1$ - do not restrict $\bar{\mathcal{G}}_S$ nor T, and one could potentially design a theory where (A)-(C) hold, and $\bar{\mathcal{G}}_S$ or T vanish at some time, and hence N vanishes too. All in all, if there exists a No-Go argument it must be a momentum dependent statement. A general statement fails because the action for the tensor and scalar modes are modified in substantially diverging ways by torsion.

Now, it is important to mention a potentially special case. For that, let us first notice the general identity

$$\bar{\mathcal{G}}_{\tau} = f_2 \, \bar{\mathcal{G}}_{\mathcal{S}} \,, \tag{18}$$

which is still not a strong link between the tensor and scalar modes at all momenta - even if $\mathcal{G}_{\tau} > 0$ and $\bar{\mathcal{G}}_{\tau} > 0$ by (B) - because the sign of $f_2(\eta)$ is in principle not constrained. However, by the relation (18) and the form of \mathcal{G}_{τ} in Eqn. (13) it is clear that at low momentum a closer relation between the tensor and scalar modes arises¹⁴: namely as $\vec{p}^2 \to 0$ we find

$$\mathcal{G}_{\tau}(\vec{p}^2 = 0) = \bar{\mathcal{G}}_{\mathcal{S}}, \ \mathcal{F}_{\tau}(\vec{p}^2 = 0) = \bar{\mathcal{F}}_{\mathcal{S}}.$$
 (19)

Thus, according to the inequalities (B) the no-ghost and stability of the tensor modes in the low momentum constrain the sign of $\overline{\mathcal{G}}_{\mathcal{S}}$ and $\overline{\mathcal{F}}_{\mathcal{S}}$ of the scalar mode. Evidently this reduces the ways in which N can vanish at low momentum, but let us recall that T is momentum independent and as we show in the Appendix VE it may vanish in many different ways, thus making unplausible an exact No-Go even in this low momentum case. Indeed, we show an explicit counterexample to a No-Go in the next section.

¹³ Although the notation for $\bar{\mathcal{G}}_{\mathcal{S}}$ and $\mathcal{G}_{\mathcal{S}}$ is similar, they are very different functions (See Eqn. (16), and note that Σ is a complicated function of the Lagrangian functions.). So, the condition in (B) $\mathcal{G}_{\mathcal{S}} > 0$ does not constrain the sign of $\bar{\mathcal{G}}_{\mathcal{S}}$.

¹⁴ This is not that surprising given that in Eq. (6), the low momentum limit $\vec{p}^2 \to 0$ is equivalent to the limit $G_5 \to 0$. Namely, the low momentum limit recovers a dispersion relation of the graviton similar to the form of "up to quartic" theories (let us recall that $f_1 \propto G_5^2$ and $f_3 \propto G_5^2$).



8



(a) Speed of gravitational waves and of the scalar mode. The Hubble showing a short bounce phase centered at $\eta = 0$ and of width $\tau = 1$. The isotropic torsion background, $x(\eta)$.

(b) Detail of the speed of gravitational waves close to the bounce phase.

FIG. 3: Physically relevant quantities.

C. A counter example to a "No-Go": an all time stable, nonsingular and subluminal toy model

We support the previous discussion showing that a model exists, taken from within the general Horndeski-Cartan theories (with G_5) (1), which has a classical solution that is nonsingular, stable and sub/luminal *at all times*, thus showing that a "No-Go theorem" cannot exist.

Our sole intention is to show that there are such potentially interesting models. Thus, our criteria is limited to the latter and whether further features can be achieved - such as physically interesting asymptotics - is left as an open question.

The strategy to build the model is to take an Ansatz for the scalar potentials G_2 to G_5 , as an expansion in powers of X, with coefficients that are general functions of ϕ . Then, assuming a set of physically compelling solutions for a bouncing cosmology, H and ϕ (Eqns. (20), (21)), we work backwards to solve the free functions in the Ansatz such that the equations of motion of the background fields are satisfied. In this section we summarize the key features of the model. The details are relegated to the Appendix VA.

In our model the background fields are

$$a = (\tau^2 + \eta^2)^{\frac{1}{6}}, \ H = \frac{\dot{a}}{a^2} = \frac{\eta}{3(\tau^2 + \eta^2)^{\frac{7}{6}}}, \ \phi = \eta, \ x = -\frac{1}{3(1 + \eta^2)^{\frac{1}{6}}},$$
(20)

and to satisfy the conditions (A)-(C) for the tensor perturbation at all momenta, the following inequalities hold

$$\bar{\mathcal{G}}_{\tau} > 0, \ \bar{f}_0 > 0, \ \bar{f}_1 > 0, \ f_2 > 0, \ f_3 = d_1^2 > 0, \ c_g^2 = \frac{f_1}{f_3} \le 1,$$
(21)

which we plot in Figures 4a and 3. The fourth inequality in (21) guarantees that in the global factor of the tensor sector the following holds for all \vec{p} : $(f_2 + \vec{p}^2 d_1^2) > 0$. Hence, the stability and non-ghosty conditions remain the same for all momenta. Similarly, the second inequality is required for the stability of the graviton at all momenta.

For the (momentum independent) scalar sector the inequalities that must hold are simply those in (B) in the last section, which we plot for our model in Figure 4b.

Now, because N must vanish and simultaneously we have no ghosty graviton at low momentum $\mathcal{G}_{\tau}(\vec{p}^2 = 0) = \bar{\mathcal{G}}_S > 0$, we have chosen a model where T vanishes, as in the Figure 5, which was the only option left to avoid any pathologies. Importantly, all Euler-Lagrange equations for the background fields $a(\eta)$, ϕ , x (20) are satisfied and the Lagrangian functions are everywhere regular. Indeed, the Ansatz for the Lagrangian functions G_2 to G_5 is finally solved as in Figures 6, 7. In Figure 3 we also show the Hubble parameter, the speed of the tensor modes and the scalar, and the torsion background.

IV. CONCLUSIONS

We showed that with the most general Horndeski theories with curvature and torsion it is possible to build classically "healthy" models, namely, with nonsingular, all time stable and subluminal cosmological solutions, at least at linear order in a perturbative expansion.

We first argued how the latter compares to simpler forms of the theory, namely: the historically common form of Horndeski without torsion, and a simplified version of the theory with torsion (namely, without G_5). We explained where the usual No-Go arguments break when generalizing from the latter two to the former full Horndeski theory with curvature and torsion.

In brief, we noted that the mathematically and physically unjustified assumption of a torsionless spacetime (or the restriction to up to quartic torsionful theory) leads to accidental relations at linear order, which restrict the healthiness of the solutions in these simpler variations of the theory, in the form of the "No-Go theorems".

Torsion solves the issues by mere "force", simply because the *tensor sector* and the *scalar sector* couple *in different* ways to torsion perturbations, thus breaking the usual links between the two sectors, which would otherwise lead to the "No-Go". Namely, despite the fact that the torsion perturbations are non dynamical, these couplings of the metric perturbation and the Horndeski scalar to the Torsion field, greatly modify *in diverging ways* the coefficients in the action for the final graviton and the scalar mode. Thus, the stability, sub/ luminality and no ghost criteria restrict very different combinations of the Lagrangian functions for both sectors, allowing to independently satisfy them without meeting any contradictions at some time in the evolution.

We also showed that all the healthy models must have a peculiar dispersion relation for the gravitational waves. It opens the question whether there are phenomenological imprints left by this distinctive graviton.

We showed a toy model where all the criteria of healthiness can be satisfied. However, the usual construction methods used in the literature (e.g. [19] or [24]) prove hard to implement, because the usual criteria for healthiness splits into many conditions that usually cannot be solved algebraically (See Appendix VA). We presume that obtaining models with physically relevant asymptotics besides the criteria of healthiness is a computational challenge rather than a fundamental obstruction.

ACKNOWLEDGEMENTS

The work of S.M. is partly supported by the grant of the Foundation for the Advancement of Theoretical Physics and Mathematics "BASIS" and by RFBR grant 21-51-46010.





mode.



FIG. 4: Stability and no-ghost condition for tensor and scalar modes at all momenta.

V. APPENDICES

A. Construction of the healthy toy model

In this appendix we show the details of how to construct the model given in section III C.

A reasonable assumption is that taking the all-time nonsingular and stable bouncing solution with a short period of superluminality that was built in [20], one can cure the superluminality issue with a contribution from the new G_5 part of the action (1). This is our approach below ¹⁵.

1. Lagrangian functions G_3, G_4

As a first step to a "Healthy model" including G_5 , let us first build a model with a bouncing solution, but with up to G_4 . By [20] we know that without G_5 the model will suffer at least instabilities or superluminal periods. We will refer to the latter as the "Unhealthy part" of the model. The strategy is then to introduce G_5 to remedy these problems.

For the "Unhealthy part" we will closely follow the example built in [20]. Therefore, for this subsection let us first consider the action (1) without \mathcal{L}_5 . In that case we can also reach the same form of the actions (11), (14) and we use the same notation for these simplified functions that, let us stress again, *only* in this section do not contain G_5 nor its derivatives.

Using in our advantage the generality of Horndeski theory, let us choose the following Ansatz for the general Lagrangian functions G_3 and G_4 :

$$G_3(\phi, X) = g_{30}(\phi) + g_{31}(\phi) \left(X - X_0\right), \tag{22}$$

$$G_4(\phi, X) = g_{40}(\phi) + g_{41}(\phi) \left(X - X_0 \right), \qquad (23)$$

where we choose X_0 as the function X valued on the choice of background fields

$$a = (\tau^2 + \eta^2)^{\frac{1}{6}}, \quad \phi = \eta.$$
(24)

This choice must be such that the Euler-Lagrange equations for the background fields of the full Horndeski-Cartan theory (including G_5) are solved. We guarantee it in the sections below by suitably fixing the g_{30} , G_2 Lagrangian functions.



FIG. 5: Avoiding the No-Go

¹⁵ This approach is taken by mere simplicity. Namely, close inspection shows that building such a model including G_5 with the "reconstruction" method used in [20, 24] is computationally challenging. The main issue is that simple Ansätze usually lead to differential equations of the Lagrangian functions G_A , which are difficult to solve.

The parameter $\tau > 0$ fixes the maximum of H and the width of the bounce phase (the domain where $\dot{H}(\eta) > 0$ around $\eta = 0$), as in Figure 3. With the solutions (24) we have the following function of ϕ for the Ansatz: $X_0 = 1/(2(\tau^2 + \phi^2)^{\frac{1}{3}})$ and we choose a short bounce $\tau = 1$.

Now, let us start with a similar model as the one built in [20]. Namely, let us solve for g_{31} , g_{40} , g_{41} algebraically from the following three equations,

$$\mathcal{F}_{\tau}(g_{40}, g_{41}) = \frac{23}{25}, \qquad (25)$$

$$\frac{T(g_{40}, g_{41})}{\mathcal{F}_{\tau}} = -\frac{7}{5} + \frac{1}{4}\operatorname{sech}\left(\frac{29}{50}\frac{\eta}{\tau}\right)^2 + \frac{1}{2}\operatorname{sech}\left(\frac{2}{5}\left(\frac{\eta}{\tau} + \frac{3}{2}\right)\right)$$
(26)

$$\Theta(g_{30}, g_{31}, g_{40}, g_{41}) = -H, \tag{27}$$

where only for these equations we plug-in the Ansatz

$$x =: -\frac{8a^3 H X G_{4,X} + a^2 \dot{\phi} (G_3 - 2G_{4,\phi})}{4(G_4 + 2X G_{4,X})}.$$
(28)

Let us stress that the equation (28) is only physically meaningful within the context of up to quartic Horndeski-Cartan theory (See [20] for more details). Namely, the expression (28) is devoid of meaning for the full Horndeski-Cartan theory with G_5 and in this note it is simply an Ansatz for the function $x(\eta)$ that appears in the functions $\mathcal{F}_{\tau}(g_{40}, g_{41}), T(g_{40}, g_{41}), \Theta(g_{30}, g_{31}, g_{40}, g_{41})$ in this section. Here, we take this definition for the mere practical purpose of building the "Unhealthy part of the model" in the same way as in [20]. Up to this point **this amounts to nothing more than a choice of a part of the free Lagrangian functions** G_3, G_4 .

Now, the key issue of solving the Euler-Lagrange equation derived from the full Horndeski-Cartan theory (with G_5) will be resolved later by suitably fixing the free Lagrangian functions g_{30} , G_2 .

2. Lagrangian functions G_2, G_5

Let us take this "Unhealthy part of the model" as a basis. Namely, keeping our choice of g_{31} , g_{40} , g_{41} which are functions of time and g_{30} , but are independent of G_5 and G_2 , let us now consider the full Horndeski-Cartan action (1) with G_5 , and let us take the Ansatz

$$G_2(\phi, X) = g_{20}(\phi) + g_{21}(\phi) (X - X_0) + g_{22}(\phi) (X - X_0)^2, \qquad (29)$$

$$G_5(\phi, X) = g_{50}(\phi) + g_{51}(\phi) \left(X - X_0 \right) + g_{52}(\phi) \left(X - X_0 \right)^2.$$
(30)

With this particular Ansatz all equations to satisfy the stability, subluminality and no-ghost criteria become differential equations of the free Lagrangian functions g_{50} , g_{51} . To avoid these issues, we take a practical simple approach and choosing an Ansatz for the functions g_{50} , g_{51} we show that a model exists with the desired features. Indeed, careful choice of parameters with a general Ansatz (and potentially fine tuning¹⁶) proves that the following choice is enough for our purpose:

$$g_{50} = \frac{1}{100} \left(-94 - 26 \operatorname{sech} \left(\frac{1}{2} - \eta \right) + 49 \operatorname{tanh} \left(\frac{3}{2} - \eta \right) - 90 \operatorname{sech} (\eta) \operatorname{tanh}(\eta) - 80 \operatorname{sech} \left(\frac{7}{10} (\eta + 1) \right) \operatorname{tanh} \left(\frac{7}{10} (\eta + 1) \right)^2 \right),$$
(31)

$$g_{51} = \frac{1}{20} \left(95 + 97 \operatorname{sech} \left(\frac{37}{25} - \eta \right) + 20 \operatorname{tanh} (\eta + 1) + 150 \operatorname{sech} \left(\frac{3}{5} \left(\eta - \frac{7}{4} \right) \right) \operatorname{tanh} \left(\frac{3}{5} \left(\eta - \frac{7}{4} \right) \right) + 20 \operatorname{tanh} \left(\frac{7}{10} \left(\eta + \frac{11}{20} \right) \right)^2 \right).$$
(32)

¹⁶ This would most probably be a problem of the Ansatz and our approach. Whether it can be avoided is an open question that we do not address in this note.



(a) Plot of g_{20} , g_{21} , g_{22} which we chose to satisfy two of the equations of motion for the background fields, and the luminality of the scalar.

(b) Plot of g_{30} , g_{31} which we chose to satisfy the third independent equation of motion of background fields, and to fix the function Θ

FIG. 6: Everywhere regular Lagrangian functions G_2, G_3

Now, plugging-in these Lagrangian functions g_{50} , g_{51} and g_{31} , g_{40} , g_{41} of the last section, we can solve for g_{30} , g_{52} algebraically from the following equations

$$\Theta(g_{30}, g_{52}) = -H + \frac{1}{100} \left(7 \operatorname{sech} \left(3 \left(\eta + \frac{27}{20} \right) \right) - 10 \operatorname{sech} \left(2 \left(\eta - 1 \right) \right) \right),$$
(33)

$$\mathcal{E}_{{}^{0}K_{0ii}}(g_{30}) = 0, \qquad (34)$$

where $\mathcal{E}_f = 0$, which denotes the Euler-Lagrange equation for the background field f, and $\Theta(g_{30}, g_{52})$ (which corresponds to the function Θ obtained from the action (14)) are computed from the full Horndeski-Cartan lagrangian (1) with G_5 . It is important to note that $\Theta(g_{30}, g_{52})$ in (33) is totally different from the function (27) $\Theta(g_{30}, g_{31}, g_{40}, g_{41})$ of the last section, which was computed without G_5 only to choose the Lagrangian functions g_{31}, g_{40}, g_{41} .

Furthermore, for simplicity -and in order to mimic the $x(\eta)$ profile in [20]- when solving (33), (34), we have chosen a model within the full Horndeski-Cartan theories (1) whose torsion background is

$$x(\eta) = -\frac{1}{3\left(1+\eta^2\right)^{\frac{1}{6}}}.$$
(35)

In other words, by solving the Euler-Lagrange equation (34) we have suitably found the Lagrangian function g_{30} that leads to our desired solution (35). Thus, the torsion background (35) has actual physical meaning because it is used to solve the equations of motion (34) and (36) below, in contrast to the function (28) that we took as Ansatz for mere practical purposes in three *ad-hoc* equations in the last section just to select the Lagrangian functions g_{31} , g_{40} , g_{41} .

The approach in this section of fixing $x(\eta)$ and finding g_{30} from the Euler-Lagrange equation is simpler because $\mathcal{E}_{0_{K_{0ij}}}(g_{30})$ is quadratic in the former and linear in the latter.

Finally, to make our choices of background fields $a(\eta)$, $\phi(\eta)$, $x(\eta)$ (24) and (35) consistent, we must satisfy the remaining Euler-Lagrange equations. We do this by choosing a model within the general action (1) (with G_5) with g_{20} , g_{21} , g_{22} solving algebraically the three independent equations

$$\mathcal{G}_{\mathcal{S}} = \mathcal{F}_{\mathcal{S}}, \ \mathcal{E}_{g_{00}} = 0, \ \mathcal{E}_{g_{11}} = \mathcal{E}_{g_{22}} = \mathcal{E}_{g_{33}} = 0,$$
 (36)

of which the first equation implies a model with a scalar mode that propagates exactly at the speed of light.

The solution for the Lagrangian functions are nonsingular everywhere. They are plotted in Figures 6 and 7.

B. Notation for torsion and its decomposition into irreducible components

We consider torsion in the second order (metric formalism). We closely follow the notation in [20, 34]. We give the necessary details below for completeness. As before, let us denote the torsion tensor as the difference between the



FIG. 7: Everywhere regular Lagrangian functions G_4 , G_5 . Let us notice that $f_3 \propto d_1 \propto G_5 = g_{50}$ on-shell, which does not vanish. Hence, the speed of gravitational waves $\propto f_3^{-1}$ in Eq. (7) is well defined.

non-symmetric connection:

$$T^{\rho}{}_{\mu\nu} = \tilde{\Gamma}^{\rho}{}_{\mu\nu} - \tilde{\Gamma}^{\rho}{}_{\nu\mu} \,, \tag{37}$$

and for convenience we also introduce the contortion tensor:

$$K^{\rho}{}_{\mu\nu} = -\frac{1}{2} \left(T^{\rho}{}_{\nu}{}^{\rho}{}_{\mu} + T^{\rho}{}_{\mu}{}_{\nu} + T^{\rho}{}_{\mu\nu} \right) , \qquad (38)$$

and let us notice the antisymmetry $T^{\rho}_{\mu\nu} = -T^{\rho}_{\nu\mu}$, $K_{\mu\nu\sigma} = -K_{\sigma\nu\mu}$. It is easy to see that we can express torsionful quantities in terms of torsionless covariant derivatives (∇) associated with the Christoffel connection $(\Gamma^{\rho}_{\mu\nu})$, plus contortion, as follows:

$$\tilde{\Gamma}^{\rho}_{\mu\nu} = \Gamma^{\rho}_{\mu\nu} - K^{\rho}_{\ \mu\nu} \,, \tag{39}$$

$$\tilde{\nabla}_{\mu}V^{\nu} = \partial_{\mu}V^{\nu} + \tilde{\Gamma}^{\nu}_{\mu\lambda}V^{\lambda} \tag{40}$$

$$\tilde{\nabla}_{\mu}V^{\nu} = \nabla_{\mu}V^{\nu} - K^{\nu}{}_{\mu\lambda}V^{\lambda} \,, \tag{41}$$

where we draw attention to our convention to sum over the second lower index of the torsionful connection in Eq. (40). Thus, we can explicitly write the action (1) in terms of contortion, the metric and the Horndeski scalar using Eq. (41), such as in [34]. For instance, we can rewrite the Ricci tensor computed with the nonsymmetric connection, \tilde{R} , in terms of the Ricci tensor computed with the Christoffel connection, R, as

$$\tilde{R}_{\mu\nu} = R_{\mu\nu} + \nabla_{\nu} K^{\rho}{}_{\rho\mu} - \nabla_{\rho} K^{\rho}{}_{\nu\mu} + K^{\rho}{}_{\rho\sigma} K^{\sigma}{}_{\nu\mu} - K^{\rho}{}_{\nu\sigma} K^{\sigma}{}_{\rho\mu} , \qquad (42)$$

and similarly for the Einstein tensor. Thus, the latter are not symmetric [29] and they introduce far from trivial $\mathcal{O}(K^2, K^3)$ contributions in the first term of the action (5), $G_5 \tilde{G}^{\mu\nu} \tilde{\nabla}_{\mu} \tilde{\nabla}_{\nu} \phi$, which, as discussed before, is responsible for the momentum dependent dispersion relation of the graviton (6). More precisely, the $p^2 T_{ij}^{(1)} T_{ij}^{(2)}$, which is essential to the stark modification of the graviton in comparison to other forms of Horndeski theory, stems from the third term of the Ricci tensor in Eq. (42) plugging in the first term of the action (5): $G_5 (\nabla_{\rho} K^{\rho\nu\mu}) \tilde{\nabla}_{\mu} \tilde{\nabla}_{\nu} \phi$. This turns out to be essential to allow globally healthy Horndeski gravities.

1. Linearization

For the perturbative expansion at linear order about a spatially flat FLRW background let us decompose the perturbations into irreducible components under small rotation group. We show all details below for completeness. The perturbed metric is denoted as

$$\mathrm{d}s^2 = (\eta_{\mu\nu} + \delta g_{\mu\nu}) \,\mathrm{d}x^\mu \,\mathrm{d}x^\nu \tag{43}$$

where

$$\eta_{\mu\nu} \mathrm{d}x^{\mu} \,\mathrm{d}x^{\nu} = a^2(\eta) \left(-\mathrm{d}\eta^2 + \delta_{ij} \,\mathrm{d}x^i \,\mathrm{d}x^j\right) \tag{44}$$

and η is conformal time. The metric perturbation is

$$\delta g_{\mu\nu} \,\mathrm{d}x^{\mu} \,\mathrm{d}x^{\nu} = a^2(\eta) \left(-2\,\alpha\,\mathrm{d}\eta^2 + 2\left(\partial_i B + S_i\right)\mathrm{d}\eta\,\mathrm{d}x^i + \left(-2\,\psi\,\delta_{ij} + 2\,\partial_i\partial_j E + \partial_i F_j + \partial_j F_i + 2\,h_{ij}\right)\mathrm{d}x^i\,\mathrm{d}x^j\right)\,,\tag{45}$$

with α , B, ψ , E scalar perturbations, S_i , F_i transverse vector perturbations, and h_{ij} , a symmetric, traceless and transverse tensor perturbation.

For contortion, antisymmetric in the first and third indices, there are 24 independent components. They are written as eight scalars denoted as $C^{(n)}$ with n = 1, ..., 8, six (two-component) transverse vectors denoted as $V_i^{(m)}$ with m = 1, ..., 6 and two (two-component) traceless, symmetric, transverse tensors $T_{ij}^{(1)}, T_{ij}^{(2)}$.

Explicitly, the decomposition of contortion perturbation reads, for the scalar sector

$$\delta K_{i00}^{\text{scalar}} = \partial_i C^{(1)}$$

$$\delta K_{ij0}^{\text{scalar}} = \partial_i \partial_j C^{(2)} + \delta_{ij} C^{(3)} + \epsilon_{ijk} \partial_k C^{(4)}$$

$$\delta K_{i0k}^{\text{scalar}} = \epsilon_{ikj} \partial_j C^{(5)}$$

$$\delta K_{iik}^{\text{scalar}} = (\delta_{ij} \partial_k - \delta_{kj} \partial_i) C^{(6)} + \epsilon_{ikl} \partial_l \partial_j C^{(7)} + (\epsilon_{ijl} \partial_l \partial_k - \epsilon_{kjl} \partial_l \partial_i) C^{(8)}, \qquad (46)$$

for the vector sector

$$\begin{split} \delta K_{i00}^{\text{vector}} &= V_i^{(1)} \\ \delta K_{ij0}^{\text{vector}} &= \partial_i V_j^{(2)} + \partial_j V_i^{(3)} \\ \delta K_{i0k}^{\text{vector}} &= \partial_i V_k^{(4)} - \partial_k V_i^{(4)} \\ \delta K_{ijk}^{\text{vector}} &= \delta_{ij} V_k^{(5)} - \delta_{kj} V_i^{(5)} + \partial_j \partial_i V_k^{(6)} - \partial_j \partial_k V_i^{(6)}, \end{split}$$

$$(47)$$

and for the tensor sector

$$\delta K_{ij0}^{\text{tensor}} = T_{ij}^{(1)}$$

$$\delta K_{ijk}^{\text{tensor}} = \partial_i T_{jk}^{(2)} - \partial_k T_{ji}^{(2)}, \qquad (48)$$

Thus, the components of contortion perturbation are

$$\delta K_{i\mu\nu} = \delta K_{i\mu\nu}^{\text{scalar}} + \delta K_{i\mu\nu}^{\text{vector}} + \delta K_{i\mu\nu}^{\text{tensor}} \,. \tag{49}$$

The non-vanishing components of the background contortion tensor on a homogeneous and isotropic background spacetime are

$${}^{0}K_{0jk} = x(\eta)\delta_{jk}$$

$${}^{0}K_{ijk} = y(\eta)\epsilon_{ijk}, \qquad (50)$$

thus the contortion tensor with all indices down is written as:

$$K_{\mu\nu\sigma} = {}^{0}K_{\mu\nu\sigma} + \delta K_{\mu\nu\sigma} \tag{51}$$

Finally, the scalar field is written in terms of a time dependent part $\phi(\eta)$ plus a spacetime dependent perturbation Π . The distinction between the background field $\phi(\eta)$ and the spacetime dependent field $\phi(x)$ will be clear from the context.

C. Equations of motion for the background fields

From (1) we can compute the action for the background fields a, ϕ, x, y . The Euler-Lagrange equations $\mathcal{E}_f = \frac{\partial \mathcal{L}}{\partial f} = 0$ for f one of the following components $g_{00}, g_{ij}, K_{ij0}, K_{ijk}, \phi$ are of up to second order in the background fields. In particular,

$$\mathcal{E}_{K_{ijk}} = -y \, \frac{2 \, a^4 \, G_4 + \, G_5 \left(a^2 \, \ddot{\phi} + \, x \, \dot{\phi} \right)}{a^{10}} \,, \tag{52}$$

leads to two branches of background solutions: namely, either the torsion background y vanishes or not. Throughout this note we have assumed the simpler, former branch

$$y(\eta) \equiv 0. \tag{53}$$

Let us notice that if there is no G_5 in the action (up to quartic Horndeski-Cartan), Eq. (53) would be the only possible solution to Eq. (52). Thus, the branch (53) is a natural continuation to previous results in up to quartic Horndeski-Cartan theory in [20, 34, 35].

With (53) the remaining equations can be written with

$$\begin{aligned} \mathcal{E}_{g_{00}} &= \frac{1}{2 a^{12}} \Big(a^{10} \big(-G_2 - 6 \, G_4 \, H^2 + \big(2 \, G_{2,X} - 2 \, G_{3,\phi} + \big(24 \, G_{4,X} - 18 \, G_{5,\phi} \big) \, H^2 \big) \, X + \big(24 \, G_{4,XX} - 12 \, G_{5,\phi X} \big) \, H^2 \, X^2 \big) \\ &+ a^7 \big(\, H(72 \, G_{4,X} \, x - 36 \, G_{5,\phi} \, x) \, X + \, H(48 \, G_{4,XX} \, x - 24 \, G_{5,\phi X} \, x) \, X^2 \big) \\ &+ a^4 \big(6 \, G_4 \, x^2 + \big(48 \, G_{4,X} \, x^2 - 18 \, G_{5,\phi} \, x^2 \big) \, X + \big(24 \, G_{4,XX} \, x^2 - 12 \, G_{5,\phi X} \, x^2 \big) \, X^2 \big) \\ &+ a^9 \big(-6 \, G_{4,\phi} \, H + \big(\big(6 \, G_{3,X} - 12 \, G_{4,\phi X} \big) \, H + 10 \, G_{5,X} \, H^3 \big) \, X + 4 \, G_{5,XX} \, H^3 \, X^2 \big) \, \dot{\phi} \\ &+ a^6 \big(3 \, G_3 \, x - 6 \, G_{4,\phi} \, x + 9 \, G_5 \, H^2 \, x + \big(6 \, G_{3,X} \, x - 12 \, G_{4,\phi X} \, x + 36 \, G_{5,X} \, H^2 \, x \big) \, X + 12 \, G_{5,XX} \, H^2 \, x \, X^2 \big) \, \dot{\phi} \\ &+ a^3 \big(18 \, G_5 \, H \, x^2 + 42 \, G_{5,X} \, H \, x^2 \, X + 12 \, G_{5,XX} \, H \, x^2 \, X^2 \big) \, \dot{\phi} + \big(9 \, G_5 \, x^3 + 16 \, G_{5,X} \, x^3 \, X + 4 \, G_{5,XX} \, x^3 \, X^2 \big) \, \dot{\phi} \Big) \,, \end{aligned}$$

$$\begin{split} \mathcal{E}_{g_{ij}} &= \frac{\delta_{ij}}{2 a^{12}} \left(a^{10} (G_2 + 6 G_4 H^2 + (-2 G_{3,\phi} + 4 G_{4,\phi\phi}) X) + (-3 G_5 x^3 - 2 G_{5,X} x^3 X) \dot{\phi} \right. \\ &+ a^6 (G_3 x - 2 G_{4,\phi} x + 3 G_5 H^2 x + (-8 G_{4,\phi\chi} x + 4 G_{5,\phi\phi} x + 18 G_{5,X} H^2 x) X + 8 G_{5,XX} H^2 x X^2) \dot{\phi} \\ &+ a^3 (6 G_5 H x^2 + 14 G_{5,X} H x^2 X + 4 G_{5,XX} H x^2 X^2) \dot{\phi} \\ &+ a^9 (4 G_4 \dot{H} + (2 G_{4,\phi} H + ((2 G_{3,X} - 12 G_{4,\phi\chi} + 4 G_{5,\phi\phi}) H + 2 G_{5,X} H^3) X + 4 G_{5,XX} H^3 X^2) \dot{\phi}) \\ &+ a^8 (2 G_{4,\phi} \ddot{\phi} - 4 G_{5,XX} H^2 X^2 \ddot{\phi} + X (-2 G_{3,X} \ddot{\phi} + 4 G_{4,\phi\chi} \ddot{\phi} - 6 G_{5,X} H^2 \ddot{\phi}) - 4 G_{5,X} H X \dot{H} \dot{\phi} \\ &+ ((-2 G_{4,X} - G_{5,\phi}) H^2 + (8 G_{4,XX} - 6 G_{5,\phi\chi}) H^2 X) \dot{\phi}^2) \\ &+ a^7 ((H (-4 G_{4,X} \ddot{\phi} + 4 G_{5,\phi} \ddot{\phi}) + H X (-8 G_{4,XX} \ddot{\phi} + 4 G_{5,\phi\chi} \ddot{\phi})) \dot{\phi} + (-4 G_{4,X} + 2 G_{5,\phi}) \dot{H} \dot{\phi}^2) \\ &+ a^4 (-2 G_4 x^2 + (-8 G_{4,X} x \ddot{\phi} + 4 G_{5,\phi} x \ddot{\phi} + X (-8 G_{4,XX} x \ddot{\phi} + 4 G_{5,\phi\chi} x \ddot{\phi})) \dot{\phi} + \dot{\phi}^2 (-4 G_{4,X} \dot{x} + 2 G_{5,\phi} \dot{x})) \\ &+ a^2 (-2 G_5 x^2 \ddot{\phi} - 10 G_{5,X} x^2 X \ddot{\phi} - 4 G_{5,XX} x^2 X^2 \ddot{\phi} + (-2 G_{4,X} x^2 - G_{5,\phi} x^2 - 2 G_{5,\phi\chi} x^2 X) \dot{\phi}^2 \\ &+ \dot{\phi} (-4 G_5 x \dot{x} - 4 G_{5,X} x X \dot{x})) + a^5 (-2 G_5 H x \ddot{\phi} - 16 G_{5,X} H x X \ddot{\phi} - 8 G_{5,XX} H x X^2 \ddot{\phi} \\ &+ (H (12 G_{4,X} x - 8 G_{5,\phi} x) + H (8 G_{4,XX} x - 8 G_{5,\phi\chi} x) X) \dot{\phi}^2 \\ &+ \dot{\phi} (-2 G_5 x \dot{H} - 2 G_5 H \dot{x} + X (-4 G_{5,X} x \dot{H} - 4 G_{5,X} H \dot{x}))) \Big), \end{split}$$

$$\mathcal{E}_{K_{ij0}} = \frac{\delta_{ij}}{2 a^{10}} \Big(a^7 (-8 G_{4,X} + 4 G_{5,\phi}) H X + a^4 (-4 G_4 x + (-8 G_{4,X} x + 4 G_{5,\phi} x) X) \\ + a^6 (-G_3 + 2 G_{4,\phi} - G_5 H^2 - 2 G_{5,X} H^2 X) \dot{\phi} + a^3 (-4 G_5 H x - 4 G_{5,X} H x X) \dot{\phi} + (-3 G_5 x^2 - 2 G_{5,X} x^2 X) \dot{\phi} \Big),$$
(56)

and, due to gauge redundancy, the equation for ϕ is trivially satisfied by the latter and their time derivatives. Indeed,

$$\mathcal{E}_{\phi} = a^2 \left(\dot{\mathcal{E}}_{g_{00}} + a \left(5 \,\mathcal{E}_{g_{00}} + 3 \,\mathcal{E}_{g_{ii}} \right) \, H \right) + 3 \, x \left(\dot{\mathcal{E}}_{K_{ii0}} + 4 \, a \, H \, \mathcal{E}_{K_{ii0}} \right) \,, \tag{57}$$

where repeated spatial indices are not summed in (57).

D. Quadratic action for the tensor sector

From (1) we can write the quadratic action for the tensor modes in the form (8). The coefficients b_A , c_A , d_A are functions of time *only*. They are given by the following expressions:

$$b_1 = a^2 (G_4 + (-2G_{4,X} + G_{5,\phi})X) - aG_{5,X} H X \dot{\phi} - \frac{x(G_5 + 2G_{5,X}X)\dot{\phi}}{2a^2},$$
(58)

$$b_2 = -\frac{2a^4(G_4 - G_{5,\phi}X) + aG_{5,X}H\dot{\phi}^3 - \dot{\phi}(3G_5x + G_{5,X}\ddot{\phi}\dot{\phi})}{2a^2},$$
(59)

$$b_{3} = -\frac{2}{a^{6}} \left(x^{3} (3G_{5} + 2G_{5,X}X) \dot{\phi} - 4a^{3} H x^{2} (G_{5} + X(3G_{5,X} + G_{5,XX}X)) \dot{\phi} - a^{6} x(2(-2G_{4,\phi X} + G_{5,\phi\phi})X + H^{2}(G_{5} + 4X(2G_{5,X} + G_{5,XX}X))) \dot{\phi} + 2a^{2} x(x(G_{5} + X(5G_{5,X} + 2G_{5,XX}X))) \ddot{\phi} + x(2G_{4,X} + G_{5,\phi X}X) \dot{\phi}^{2} + 2(G_{5} + G_{5,X}X) \dot{\phi} \dot{x}) - a^{5} (x(-H(G_{5} + 4X(2G_{5,X} + G_{5,XX}X))) \ddot{\phi} - (G_{5} + 2G_{5,X}X) \dot{H} \dot{\phi} + H(4G_{4,X} - 3G_{5,\phi} + 4G_{4,XX}X - 4G_{5,\phi X}X) \dot{\phi}^{2}) - H(G_{5} + 2G_{5,X}X) \dot{\phi} \dot{x}) - a^{4} (2x(-2G_{4}x + (-2G_{4,X} + G_{5,\phi} - 2G_{4,XX}X + G_{5,\phi X}X) \ddot{\phi} \dot{\phi}) + (-2G_{4,X} + G_{5,\phi}) \dot{\phi}^{2} \dot{x}) \right),$$

$$(60)$$

$$c_1 = \frac{2 a^4 G_4 + a^2 G_5 \ddot{\phi} + G_5 x \dot{\phi}}{2 a^6}, \qquad (61)$$

$$c_{2} = -\frac{2x(2a^{4}(G_{4} + 2G_{4,X}X - G_{5,\phi}X) + 2a^{3}H(G_{5} + G_{5,X}X)\dot{\phi} + x(3G_{5} + 2G_{5,X}X)\dot{\phi})}{a^{6}}, \qquad (62)$$

$$c_3 = \frac{\dot{\phi}(2x(G_5 + G_{5,X}X) + a^3H(G_5 + 2G_{5,X}X) + a^2(2G_{4,X} - G_{5,\phi})\dot{\phi})}{a^4},$$
(63)

$$c_4 = \frac{1}{4x} c_2 \,, \tag{64}$$

$$d_1 = \frac{G_5 \dot{\phi}}{a^4} = -\frac{1}{a^2} d_2 = \frac{1}{2x} d_3 = \frac{1}{a^2} d_4, \qquad (65)$$

Let us highlight that one can always get rid of second derivatives of the background fields by using their equations of motion. In this note, we have used the equations of motion for the background fields to express G_2 , $G_{4,X}$, $G_{4,XX}$, $G_{3,\phi}$, $G_{2,\phi X}$ (non-vanishing by assumption) in terms of other derivatives of the Lagrangian functions in order to obtain shorter expressions.

1. Details for the speed of the graviton

We wrote the dispersion relation of the gravitational waves as,

$$\omega^2 = \frac{f_0 + \vec{p}^2 f_1}{f_2 + \vec{p}^2 f_3} \vec{p}^2 ,$$

and the speed $c_g^2 = f_1/f_3$, where $f_0 = \bar{f}_0/\bar{\mathcal{G}}_{\tau}$, $f_1 = \bar{f}_1/\bar{\mathcal{G}}_{\tau}$, $f_2 = -4c_1c_4$, $f_3 = d_1^2$ and $\bar{\mathcal{G}}_{\tau} = c_1(c_3^2 - 4b_1c_4)$. The latter are fixed by the coefficients b_A , c_A , d_A given in the first part of this appendix, and

$$\begin{split} \bar{f}_{1} &= -\frac{2}{a^{20}} \Big(G_{5}^{\ 2} X(-8 \, a^{10} \, G_{4}(\, G_{4}^{\ 2} + 2 \, G_{4,X} \, G_{4} \, X - G_{5}(\, G_{3} + 2 \, G_{5} \, H^{2}) \, X) \\ &\quad + 2 \, a^{9} \, G_{5} \, H(4 \, G_{4}^{\ 2} + 8 \, G_{4}(\, G_{4,X} - G_{5,\phi}) \, X + G_{5} \, X(\, G_{3} + \, H^{2}(\, G_{5} + 2 \, G_{5,X} \, X))) \, \dot{\phi} \\ &\quad + 2 \, G_{5}^{\ 2} \, x^{2} \, \dot{\phi}(\, x \, X(3 \, G_{5} + 2 \, G_{5,X} \, X) - G_{5} \, \ddot{\phi} \, \dot{\phi}) + a^{3} \, G_{5}^{\ 2} \, H \, x \, \dot{\phi}(2 \, x \, X(5 \, G_{5} + 6 \, G_{5,X} \, X) - G_{5} \, \ddot{\phi} \, \dot{\phi}) \\ &\quad + 2 \, a^{7} \, G_{5} \, \dot{\phi}(- \, H((-4 \, G_{4,X} + G_{5,\phi}) \, G_{5} \, X + 2 \, G_{4}(\, G_{5} + 2 \, G_{5,X} \, X)) \, \ddot{\phi} + G_{4} \, G_{5} \, \dot{H} \, \dot{\phi}) \\ &\quad + 2 \, a^{8}((4 \, G_{4} \, G_{5,\phi} \, G_{5} \, X + G_{3} \, G_{5}^{\ 2} \, X - 2 \, G_{4}^{\ 2}(3 \, G_{5} + 2 \, G_{5,X} \, X)) \, \ddot{\phi} \\ &\quad + G_{5} \, H^{2}(2(\, G_{4,X} - G_{5,\phi}) \, G_{5} \, X + G_{4}(\, G_{5} + 4 \, G_{5,X} \, X)) \, \dot{\phi}^{2}) \\ &\quad + 2 \, a^{6}(-2 \, G_{4}^{\ 2} \, x(\, G_{5} + 2 \, G_{5,X} \, X) \, \dot{\phi} + G_{4} \, G_{5}(-2 \, G_{5} \, \ddot{\phi}^{\ 2} + 8 \, G_{4,X} \, x \, X \, \dot{\phi} - 4 \, G_{5,\phi} \, x \, X \, \dot{\phi} + G_{5} \, \ddot{\phi} \, \dot{\phi}) \\ &\quad + 2 \, a^{6}(-2 \, G_{4}^{\ 2} \, x(\, G_{5} + 2 \, G_{5,X} \, X) \, \dot{\phi} + G_{4} \, G_{5}(-2 \, G_{5} \, \ddot{\phi}^{\ 2} + 8 \, G_{4,X} \, x \, X \, \dot{\phi} - 4 \, G_{5,\phi} \, x \, X \, \dot{\phi} + G_{5} \, \ddot{\phi} \, \dot{\phi}))) \\ &\quad + 2 \, a^{4} \, G_{5}(x(6 \, G_{4} \, G_{5} \, x \, X + ((4 \, G_{4,X} + G_{5,\phi}) \, G_{5} \, X - 2 \, G_{4}(3 \, G_{5} + 2 \, G_{5,X} \, X)) \, \ddot{\phi} \, \dot{\phi}) + 2 \, G_{4} \, G_{5} \, \dot{\phi} \, \dot{\phi$$

 \overline{f}_0 is of little significance for the speed of the graviton and its form is cumbersome. Thus, we do not show it in this note. However, it can be easily obtained from the action (8).

E. Quadratic action for the scalar mode

1. Initial form of the action

From (1), a direct computation gives the quadratic action for the scalar mode. It can be written in the unitary gauge (namely, with $\Pi = E = 0$), as follows:

$$S_{\tau} = \frac{1}{2} \int d\eta \, d^3 p \, \left(M_1 \, C^{(3)} \, \dot{\psi} + M_2 \, B \, \dot{\psi} + M_3 \, C^{(2)} \, C^{(3)} + M_4 \, B \, C^{(3)} + M_5 \, C^{(2)} \, \dot{\psi} + M_6 \, C^{(2)} \, \psi + M_7 \, B \, \psi + M_8 \, C^{(3)} \, \psi \right. \\ \left. + M_9 \, \alpha \, \dot{\psi} + M_{10} \, C^{(3)} \, \alpha + M_{11} \, B \, \alpha + M_{12} \, C^{(2)} \, \alpha + M_{13} \, \alpha \, \psi + M_{14} \, \dot{\psi}^2 + M_{15} \, \left(C^{(3)} \right)^2 + M_{16} \, \alpha^2 + M_{17} \, \psi^2 \right),$$

$$(67)$$

Eq. (67) is trivially obtained after using the constraint equations for the Lagrange multipliers $C^{(1)}$, $C^{(5)}$, and $C^{(7)}$, which impose $C^{(4)} = C^{(6)} = C^{(8)} = 0$. The coefficients M_A depend on conformal time and momentum, and can be written as:

$$M_{1} = \frac{a^{4}(48\,G_{4,X} - 24\,G_{5,\phi})\,X + a^{3}(12\,G_{5}\,H + 24\,G_{5,X}\,H\,X)\,\phi + (24\,G_{5}\,x + 24\,G_{5,X}\,x\,X)\,\phi}{a^{4}}\,,\tag{68}$$

$$M_2 = -\frac{p^2 a^2}{3} M_1 + \frac{4(2 a^4 G_4 + G_5(a^3 H + x) \dot{\phi}) p^2}{a^2}, \qquad (69)$$

$$M_3 = \frac{1}{a^4} M_2 - \frac{8(2 a^4 G_4 + G_5(a^3 H + x) \dot{\phi}) p^2}{a^6}, \qquad (70)$$

$$M_4 = M_5 = -\frac{p^2}{3} M_1 \,, \tag{71}$$

$$M_6 = -2 x M_3 , (72)$$

$$M_7 = \frac{2\,p^2\,x}{3}\,M_1\,,\tag{73}$$

$$p^2 M_8 = 6 x M_3 + \frac{4}{a^2} p^4 G_5 \dot{\phi}, \qquad (74)$$

$$M_{9} = \frac{1}{a^{4}} \left(a^{7} (-24 G_{4} H + (48 G_{4,X} - 48 G_{5,\phi}) H X + (96 G_{4,XX} - 48 G_{5,\phiX}) H X^{2}) + a^{4} (-24 G_{4} x + (96 G_{4,XX} - 48 G_{5,\phiX}) X + (96 G_{4,XX} x - 48 G_{5,\phiX} x) X^{2}) + a^{6} (-6 G_{3} - 6 G_{5} H^{2} + (12 G_{3,X} - 24 G_{4,\phiX} + 48 G_{5,X} H^{2}) X + 24 G_{5,XX} H^{2} X^{2}) \dot{\phi} + a^{3} (12 G_{5} H x + 120 G_{5,X} H x X + 48 G_{5,XX} H x X^{2}) \dot{\phi} + (18 G_{5} x^{2} + 72 G_{5,X} x^{2} X + 24 G_{5,XX} x^{2} X^{2}) \dot{\phi} \right),$$

$$(75)$$

$$M_{10} = \frac{1}{a^6} \left(a^7 ((96 G_{4,X} - 48 G_{5,\phi}) H X + (96 G_{4,XX} - 48 G_{5,\phiX}) H X^2) + a^4 ((144 G_{4,X} x - 48 G_{5,\phiX} x) X + (96 G_{4,XX} x - 48 G_{5,\phiX} x) X^2) + a^6 (12 G_5 H^2 + (12 G_{3,X} - 24 G_{4,\phiX} + 60 G_{5,X} H^2) X + 24 G_{5,XX} H^2 X^2) \dot{\phi} + a^3 (48 G_5 H x + 144 G_{5,X} H x X + 48 G_{5,XX} H x X^2) \dot{\phi} + (36 G_5 x^2 + 84 G_{5,X} x^2 X + 24 G_{5,XX} x^2 X^2) \dot{\phi} \right),$$

$$(76)$$

$$M_{11} = -\frac{1}{3} p^2 M_9 \,, \tag{77}$$

$$M_{12} = -\frac{1}{3} p^2 M_{10} , \qquad (78)$$

$$M_{13} = \frac{1}{a^6} \left(a^7 (H(-192 G_{4,X} x + 96 G_{5,\phi} x) X + H(-192 G_{4,XX} x + 96 G_{5,\phi X} x) X^2) \right. \\ \left. + a^4 ((-288 G_{4,X} x^2 + 96 G_{5,\phi} x^2) X + (-192 G_{4,XX} x^2 + 96 G_{5,\phi X} x^2) X^2) \right. \\ \left. + a^6 (-24 G_5 H^2 x + (-24 G_{3,X} x + 48 G_{4,\phi X} x - 120 G_{5,X} H^2 x) X - 48 G_{5,XX} H^2 x X^2) \dot{\phi} \right. \\ \left. + a^3 (-96 G_5 H x^2 - 288 G_{5,X} H x^2 X - 96 G_{5,XX} H x^2 X^2) \dot{\phi} + (-72 G_5 x^3 - 168 G_{5,X} x^3 X - 48 G_{5,XX} x^3 X^2) \dot{\phi} \right. \\ \left. + (a^8 (-8 G_4 + (16 G_{4,X} - 8 G_{5,\phi}) X) + 8 a^7 G_{5,X} H X \dot{\phi} + a^4 (4 G_5 x + 8 G_{5,X} x X) \dot{\phi}) p^2 \right),$$

$$(79)$$

$$p^2 M_{14} = -\frac{3}{2} M_2 \,, \tag{80}$$

$$p^2 M_{15} = -\frac{3}{2} M_3 \,, \tag{81}$$

$$\begin{split} M_{16} &= \frac{1}{a^{6}} \left(a^{10} (-12\,G_{4}\,H^{2} + (-2\,G_{2,X} - 36\,G_{5,\phi}\,H^{2})\,X + (4\,G_{2,XX} - 4\,G_{3,\phi X} + (144\,G_{4,XX} - 84\,G_{5,\phi X})\,H^{2})\,X^{2} \\ &+ (48\,G_{4,XXX} - 24\,G_{5,\phi XX})\,H^{2}\,X^{3}) \right. \\ &+ a^{3} (46\,G_{5}\,H\,x^{2} + 208\,G_{5,X}\,H\,x^{2}\,X + 156\,G_{5,XX}\,H\,x^{2}\,X^{2} + 24\,G_{5,XXX}\,H\,x^{2}\,X^{3})\,\dot{\phi} \\ &+ (18\,G_{5}\,x^{3} + 82\,G_{5,X}\,x^{3}\,X + 56\,G_{5,XX}\,x^{3}\,X^{2} + 8\,G_{5,XXX}\,x^{3}\,X^{3})\,\dot{\phi} \\ &+ a^{9} (-8\,G_{4}\,\dot{H} + (-2\,G_{3}\,H - 2\,G_{5}\,H^{3} + ((12\,G_{3,X} - 36\,G_{4,\phi X})\,H + 44\,G_{5,X}\,H^{3})\,X \\ &+ ((12\,G_{3,XX} - 24\,G_{4,\phi XX})\,H + 44\,G_{5,XX}\,H^{3})\,X^{2} + 8\,G_{5,XXX}\,H^{3}\,X^{3})\,\dot{\phi}) + a^{7} (8\,G_{4}\,H\,x - 72\,G_{5,\phi}\,H\,x\,X \\ &+ H(336\,G_{4,XX}\,x - 168\,G_{5,\phi X}\,x)\,X^{2} + H(96\,G_{4,XXX}\,x - 48\,G_{5,\phi XX}\,x)\,X^{3} - 8\,G_{4,X}\,H\,\ddot{\phi}\,\dot{\phi}) \\ &+ a^{4} (-4\,G_{4}\,x^{2} - 36\,G_{5,\phi}\,x^{2}\,X + (192\,G_{4,XX}\,x^{2} - 84\,G_{5,\phi X}\,x^{2})\,X^{2} + (48\,G_{4,XXX}\,x^{2} - 24\,G_{5,\phi XX}\,x^{2})\,X^{3} \\ &- 8\,G_{4,X}\,x\,\ddot{\phi}\,\dot{\phi}\,) + a^{8} (-2\,G_{3}\,\ddot{\phi} - 2\,G_{5}\,H^{2}\,\ddot{\phi} - 4\,G_{5,X}\,H^{2}\,X\,\ddot{\phi} - 4\,G_{5}\,H\,\dot{H}\,\dot{\phi}\,+ 22\,G_{4,X}\,H^{2}\,\dot{\phi}^{2}) \\ &+ a^{6} ((-4\,G_{3}\,x + 26\,G_{5}\,H^{2}\,x + (18\,G_{3,X}\,x - 36\,G_{4,\phi X}\,x + 170\,G_{5,X}\,H^{2}\,x)\,X \\ &+ (12\,G_{3,XX}\,x - 24\,G_{4,\phi XX}\,x + 144\,G_{5,XX}\,H^{2}\,x)\,X^{2} + 24\,G_{5,XXX}\,H^{2}\,x\,X^{3})\,\dot{\phi} - 8\,G_{4}\,\dot{x}) \\ &+ a^{2} (-2\,G_{5}\,x^{2}\,\ddot{\phi} - 4\,G_{5,X}\,x^{2}\,X\,\ddot{\phi}\,+ 50\,G_{4,X}\,x^{2}\,\dot{\phi}^{2} - 4\,G_{5}\,x\,\dot{\phi}\,\dot{x}) \\ &+ a^{5} (-4\,G_{5}\,H\,x\,\ddot{\phi} - 8\,G_{5,X}\,H\,x\,X\,\ddot{\phi}\,+ 72\,G_{4,X}\,H\,x\,\dot{\phi}^{2}\,+ \dot{\phi}(-4\,G_{5}\,x\,\dot{H}\,- 4\,G_{5}\,H\,\dot{x}))) \Big), \end{split} \tag{82}$$

$$\begin{split} M_{17} &= \frac{1}{a^6} \left(\left(72\,G_5\,x^3 + 48\,G_{5,X}\,x^3\,X \right) \dot{\phi} \right. \\ &+ a^6 (-12\,G_5\,H^2\,x + (48\,G_{4,\phi X}\,x - 24\,G_{5,\phi\phi}\,x - 96\,G_{5,X}\,H^2\,x)\,X - 48\,G_{5,XX}\,H^2\,x\,X^2) \dot{\phi} \\ &+ a^3 (-48\,G_5\,H\,x^2 - 144\,G_{5,X}\,H\,x^2\,X - 48\,G_{5,XX}\,H\,x^2\,X^2) \dot{\phi} \\ &+ a^4 (48\,G_4\,x^2 + (48\,G_{4,X}\,x\,\ddot{\phi} - 24\,G_{5,\phi}\,x\,\ddot{\phi} + X(48\,G_{4,XX}\,x\,\ddot{\phi} - 24\,G_{5,\phi X}\,x\,\ddot{\phi})) \dot{\phi} + \dot{\phi}^2 (24\,G_{4,X}\,\dot{x} - 12\,G_{5,\phi}\,\dot{x})) \\ &+ a^2 (24\,G_5\,x^2\,\ddot{\phi} + 120\,G_{5,X}\,x^2\,X\,\ddot{\phi} + 48\,G_{5,XX}\,x^2\,X^2\,\ddot{\phi} + (48\,G_{4,X}\,x^2 + 24\,G_{5,\phi X}\,x^2\,X) \dot{\phi}^2 \\ &+ \dot{\phi} (48\,G_5\,x\,\dot{x} + 48\,G_{5,X}\,x\,X\,\dot{x})) + a^5 (12\,G_5\,H\,x\,\ddot{\phi} + 96\,G_{5,X}\,H\,x\,X\,\ddot{\phi} + 48\,G_{5,XX}\,H\,x\,X^2\,\ddot{\phi} \\ &+ (H(-48\,G_{4,X}\,x + 36\,G_{5,\phi}\,x)) + H(-48\,G_{4,XX}\,x + 48\,G_{5,\phi X}\,x)\,X) \dot{\phi}^2 \\ &+ \dot{\phi} (12\,G_5\,x\,\dot{H} + 12\,G_5\,H\,\dot{x} + X(24\,G_{5,X}\,x\,\dot{H} + 24\,G_{5,X}\,H\,\dot{x}))) \\ &+ (a^8 (4\,G_4 - 4\,G_{5,\phi}\,X) - 4\,a^6\,G_{5,X}\,X\,\ddot{\phi} - 6\,a^4\,G_5\,x\,\dot{\phi} + 4\,a^7\,G_{5,X}\,H\,X\,\dot{\phi})\,p^2 \Big) \,. \end{split}$$

$$\tag{83}$$

Again, as in the Appendix VD, we use the equations of motion for the background fields to express G_2 , $G_{4,X}$, $G_{4,XX}$, $G_{3,\phi}$, $G_{2,\phi X}$ (non-vanishing by assumption) in terms of other derivatives of the Lagrangian functions in order to obtain shorter expressions.

2. Obtaining the quadratic action for the scalar mode in its final form

Now, from the action (67) we can obtain the form (14) as follows: the Euler-Langrange equation for the Torsion scalar $C^{(2)}$ -which is a Lagrange multiplier- is a constraint that can be used to express $C^{(3)}$ in terms of α , ψ , $\dot{\psi}$. Plugging back $C^{(3)}(\alpha, \psi, \dot{\psi})$ in (67) gives (14). In particular, as explained before, the functions of time $\bar{\mathcal{G}}_{\mathcal{S}}$ and T in Eq. (14) are essential to avoid the No-Go theorems. Let us explicitly write them:

$$\bar{\mathcal{G}}_{\mathcal{S}} = \frac{(2a^4 G_4 + G_5(a^3 H + x)\phi)^2}{2a^8 G_4 + 2a^7 H(G_5 + G_{5,X}X)\dot{\phi} + a^4 x(3G_5 + 2G_{5,X}X)\dot{\phi} + a^6(2G_{4,X} - G_{5,\phi})\dot{\phi}^2},$$
(84)

$$T = \frac{t_1}{t_2},\tag{85}$$

with

$$t_{1} = \left(2\left(x^{2} X^{2} (G_{5,X} G_{5} + 4 G_{5,X}^{2} X - 2 G_{5,XX} G_{5} X\right) - 2 a^{3} H x X (G_{5}^{2} - 4 G_{5,X}^{2} X^{2} + 2 G_{5,XX} G_{5} X^{2}) + a^{6} (-2 G_{4}^{2} + X (-G_{5}^{2} H^{2} + 2 X ((-2 G_{4,X} + G_{5,\phi})^{2} + 2 G_{5,X}^{2} H^{2} X) - G_{5} X (G_{3,X} - 2 G_{4,\phi X} + H^{2} (G_{5,X} + 2 G_{5,XX} X)))) - 2 a^{5} H (G_{4} G_{5} + (-4 G_{4,X} G_{5,X} + 2 G_{5,\phi} G_{5,X} + 2 G_{4,XX} G_{5} - G_{5,\phi X} G_{5}) X^{2}) \dot{\phi} - 2 a^{2} x (G_{4} G_{5} + (-G_{4,X} + G_{5,\phi}) G_{5} X + (-4 G_{4,X} G_{5,X} + 2 G_{5,\phi} G_{5,X} + 2 G_{4,XX} G_{5} - G_{5,\phi X} G_{5}) X^{2}) \dot{\phi})\right),$$

$$(86)$$

$$t_2 = \left(2a^6(G_4 + 2G_{4,X}X - G_{5,\phi}X) + 2a^5H(G_5 + G_{5,X}X)\dot{\phi} + a^2x(3G_5 + 2G_{5,X}X)\dot{\phi}\right).$$
(87)

Thus, by inspection it is clear that even if Eq. (19) obligues us to satisfy $\mathcal{G}_{\mathcal{S}} > 0$ in order to have a stable graviton at low momentum, t_1 and hence T can still vanish in many ways with a careful choice of Lagrangian functions (it is worth to note at this point that in general $T \neq \Theta$).

That we can choose Lagrangian functions that allow T to vanish at some point -while still satisfying the stability of all modes at all momenta, and subluminality- is the key tool to build theories without suffering the usual No-Go theorems.

The technical detail that leads to these new possibilities boils down to a new G_5 term in the action (67) in comparison to up to quartic Horndeski-Cartan theories: namely, the p^2 part in the term $M_8 C^{(3)} \psi$. Indeed, the constraint imposed by $C^{(2)}$ is of the form $C^{(3)} = -\frac{M_{12}}{M_3} \alpha + \ldots$, which then gives a new $\mathcal{O}(p^2 G_5)$ term in $-\frac{M_{12}}{M_3} M_8 \alpha \psi$ which gives a totally new T in comparison to simpler forms of the theory.

VI. REFERENCES

- S. Arai, K. Aoki, Y. Chinone, R. Kimura, T. Kobayashi, H. Miyatake, D. Yamauchi, S. Yokoyama, K. Akitsu, T. Hiramatsu et al., "Cosmological gravity probes: Connecting recent theoretical developments to forthcoming observations," Progress of Theoretical and Experimental Physics, vol. 2023, no. 7, p. 072E01, 2023.
- [2] G. W. Horndeski, "Second-order scalar-tensor field equations in a four-dimensional space," International Journal of Theoretical Physics, vol. 10, no. 6, pp. 363–384, 1974.
- [3] A. Nicolis, R. Rattazzi, and E. Trincherini, "Galileon as a local modification of gravity," *Physical Review D*, vol. 79, no. 6, p. 064036, 2009.
- [4] C. Deffayet, X. Gao, D. A. Steer, and G. Zahariade, "From k-essence to generalised Galileons," Phys. Rev. D, vol. 84, p. 064039, 2011.
- [5] T. Kobayashi, M. Yamaguchi, and J. Yokoyama, "Generalized G-inflation: Inflation with the most general second-order field equations," Prog. Theor. Phys., vol. 126, pp. 511–529, 2011.
- [6] C. Deffayet, G. Esposito-Farese, and A. Vikman, "Covariant galileon," Physical Review D, vol. 79, no. 8, p. 084003, 2009.
- [7] T. Kobayashi, "Horndeski theory and beyond: a review," Reports on Progress in Physics, vol. 82, no. 8, p. 086901, 2019.
- [8] V. A. Rubakov, "The null energy condition and its violation," *Physics-Uspekhi*, vol. 57, no. 2, p. 128, 2014.
- [9] R. Penrose, "Gravitational collapse and space-time singularities," Phys. Rev. Lett., vol. 14, pp. 57–59, 1965.
- [10] S. Hawking, "Occurrence of singularities in open universes," Phys. Rev. Lett., vol. 15, pp. 689–690, 1965.
- [11] J. Evslin and T. Qiu, "Closed Timelike Curves in the Galileon Model," JHEP, vol. 11, p. 032, 2011.
- [12] V. A. Rubakov, "More about wormholes in generalized Galileon theories," Theor. Math. Phys., vol. 188, no. 2, pp. 1253– 1258, 2016.
- [13] M. Libanov, S. Mironov, and V. Rubakov, "Generalized Galileons: instabilities of bouncing and Genesis cosmologies and modified Genesis," JCAP, vol. 08, p. 037, 2016.
- [14] T. Kobayashi, "Generic instabilities of nonsingular cosmologies in Horndeski theory: A no-go theorem," Phys. Rev. D, vol. 94, no. 4, p. 043511, 2016.
- [15] R. Kolevatov and S. Mironov, "Cosmological bounces and Lorentzian wormholes in Galileon theories with an extra scalar field," Phys. Rev. D, vol. 94, no. 12, p. 123516, 2016.
- [16] S. Mironov, "Mathematical Formulation of the No-Go Theorem in Horndeski Theory," Universe, vol. 5, no. 2, p. 52, 2019.
- [17] S. Akama and T. Kobayashi, "Generalized multi-Galileons, covariantized new terms, and the no-go theorem for nonsingular cosmologies," *Phys. Rev. D*, vol. 95, no. 6, p. 064011, 2017.
- [18] P. Creminelli, D. Pirtskhalava, L. Santoni, and E. Trincherini, "Stability of Geodesically Complete Cosmologies," JCAP, vol. 11, p. 047, 2016.
- [19] A. Ijjas and P. J. Steinhardt, "Classically stable nonsingular cosmological bounces," Phys. Rev. Lett., vol. 117, no. 12, p. 121304, 2016.
- [20] S. Mironov and M. Valencia-Villegas, "Stability of nonsingular cosmologies in galileon models with torsion: A no-go theorem for eternal subluminality," *Phys. Rev. D*, vol. 109, p. 044073, Feb 2024.
- [21] A. Adams, N. Arkani-Hamed, S. Dubovsky, A. Nicolis, and R. Rattazzi, "Causality, analyticity and an IR obstruction to UV completion," JHEP, vol. 10, p. 014, 2006.
- [22] S. Dubovsky, T. Gregoire, A. Nicolis, and R. Rattazzi, "Null energy condition and superluminal propagation," JHEP, vol. 03, p. 025, 2006.
- [23] P. Creminelli, O. Janssen, and L. Senatore, "Positivity bounds on effective field theories with spontaneously broken Lorentz invariance," JHEP, vol. 09, p. 201, 2022.
- [24] S. Mironov, V. Rubakov, and V. Volkova, "Bounce beyond Horndeski with GR asymptotics and γ-crossing," JCAP, vol. 10, p. 050, 2018.
- [25] S. Mironov, V. Rubakov, and V. Volkova, "Subluminal cosmological bounce beyond Horndeski," JCAP, vol. 05, p. 024, 2020.
- [26] S. Mironov and A. Shtennikova, "Stable cosmological solutions in Horndeski theory," JCAP, vol. 06, p. 037, 2023.
- [27] Y. Ageeva, P. Petrov, and V. Rubakov, "Nonsingular cosmological models with strong gravity in the past," *Phys. Rev. D*, vol. 104, no. 6, p. 063530, 2021.
- [28] B. Ahmedov, K. F. Dialektopoulos, J. Levi Said, A. Nosirov, Z. Oikonomopoulou, and O. Yunusov, "Stable bouncing solutions in Teleparallel Horndeski gravity: violations of the no-go theorem," 11 2023.
- [29] F. W. Hehl, P. Von Der Heyde, G. D. Kerlick, and J. M. Nester, "General Relativity with Spin and Torsion: Foundations and Prospects," *Rev. Mod. Phys.*, vol. 48, pp. 393–416, 1976.
- [30] S. M. Carroll and G. B. Field, "Consequences of propagating torsion in connection dynamic theories of gravity," Phys. Rev. D, vol. 50, pp. 3867–3873, 1994.
- [31] W. Kopczyński, "A non-singular universe with torsion," Physics Letters A, vol. 39, no. 3, pp. 219–220, 1972.
- [32] J. Tafel, "A non-singular homogeneous universe with torsion," Phys. Lett. A, vol. 45, no. 4, pp. 341–342, 1973.
- [33] A. Trautman, "On the structure of the einstein-cartan equations," in Symp. Math., vol. 12, 1973, pp. 139–162.
- [34] S. Mironov and M. Valencia-Villegas, "Quartic Horndeski-Cartan theories in a FLRW universe," Phys. Rev. D, vol. 108,

no. 2, p. 024057, 2023.

- [35] M. Valencia-Villegas, "Quartic Horndeski-Cartan theories: a review on the stability of nonsingular cosmologies," PoS, vol. ICPPCRubakov2023, p. 033, 2024.
- [36] B. Ahmedov, K. F. Dialektopoulos, J. Levi Said, A. Nosirov, Z. Oikonomopoulou, and O. Yunusov, "Cosmological perturbations in the teleparallel analog of Horndeski gravity," JCAP, vol. 08, p. 074, 2023.
- [37] T. Helpin and M. S. Volkov, "Varying the horndeski lagrangian within the palatini approach," Journal of Cosmology and Astroparticle Physics, vol. 2020, no. 01, p. 044, 2020.
- [38] T. Helpin and M. S. Volkov, "A metric-affine version of the horndeski theory," International Journal of Modern Physics A, vol. 35, no. 02n03, p. 2040010, 2020.
- [39] Y.-Q. Dong, Y.-Q. Liu, and Y.-X. Liu, "Constraining Palatini-Horndeski theory with gravitational waves after GW170817," Eur. Phys. J. C, vol. 83, no. 8, p. 702, 2023.
- [40] E. Davydov, "Comparing metric and palatini approaches to vector horndeski theory," International Journal of Modern Physics D, vol. 27, no. 04, p. 1850038, 2018.
- [41] Y.-Q. Dong and Y.-X. Liu, "Polarization modes of gravitational waves in palatini-horndeski theory," *Physical Review D*, vol. 105, no. 6, p. 064035, 2022.
- [42] S. Capozziello, M. Caruana, J. Levi Said, and J. Sultana, "Ghost and Laplacian instabilities in teleparallel Horndeski gravity," JCAP, vol. 03, p. 060, 2023.
- [43] S. Bahamonde, G. Trenkler, L. G. Trombetta, and M. Yamaguchi, "Symmetric teleparallel Horndeski gravity," Phys. Rev. D, vol. 107, no. 10, p. 104024, 2023.
- [44] K. F. Dialektopoulos, J. L. Said, and Z. Oikonomopoulou, "Classification of teleparallel horndeski cosmology via noether symmetries," The European Physical Journal C, vol. 82, no. 3, pp. 1–15, 2022.
- [45] R. C. Bernardo, J. L. Said, M. Caruana, and S. Appleby, "Well-tempered minkowski solutions in teleparallel horndeski theory," *Classical and Quantum Gravity*, vol. 39, no. 1, p. 015013, 2021.
- [46] S. Bahamonde, K. F. Dialektopoulos, M. Hohmann, and J. L. Said, "Post-newtonian limit of teleparallel horndeski gravity," *Classical and Quantum Gravity*, vol. 38, no. 2, p. 025006, 2020.
- [47] S. Bahamonde, K. F. Dialektopoulos, V. Gakis, and J. L. Said, "Reviving horndeski theory using teleparallel gravity after gw170817," *Physical Review D*, vol. 101, no. 8, p. 084060, 2020.
- [48] S. Bahamonde, K. F. Dialektopoulos, and J. L. Said, "Can horndeski theory be recast using teleparallel gravity?" *Physical Review D*, vol. 100, no. 6, p. 064018, 2019.
- [49] N. Arkani-Hamed, H.-C. Cheng, M. A. Luty, and S. Mukohyama, "Ghost condensation and a consistent infrared modification of gravity," *Journal of High Energy Physics*, vol. 2004, no. 05, p. 074, 2004.