

Total coloring of graphs via efficient dominating sets

Italo J. Dejter
 University of Puerto Rico
 Rio Piedras, PR 00936-8377
italo.dejter@gmail.com

Abstract

A totally efficient coloring of a regular graph with color set formed by one unit more than its degree is a total coloring in which each vertex color class is an efficient dominating set, that is a perfect code. We find that the 3-cube graph has a totally efficient coloring and conjecture that this is the only existing case of a totally effective coloring. We also deal with a related problem exemplified by a star transposition graph and related graphs, leading to edge-girth colorings on their prism graphs.

1 A problem in total coloring and the 3-cube

A *total coloring* of a graph G is an assignment of colors to the vertices and edges of G such that no two incident or adjacent elements (vertices, edges) are assigned the same color [4]. A total coloring of G such that the vertices adjacent to each $v \in V(G)$ together with v itself are assigned pairwise different colors will be said to be an *efficient coloring*. This will be said to be *totally efficient* if: **(a)** G is r -regular; **(b)** the color set is $[r] = \{0, 1, \dots, r-1\}$; and **(c)** each $v \in V(G)$ together with its neighbors are assigned all the colors in $[r]$ via a bijection between $N[v] = N(v) \cup \{v\}$ and $[r] = \{1, 2, \dots, r\}$, where $N[v]$ and $N(v)$ are the *closed neighborhood* and *open neighborhood* of v , respectively; see [2].

Let G be a graph of regular degree $\kappa \geq 3$ and girth and chromatic index equal to $r = \kappa + 1$. Does G possess a totally efficient coloring with r colors? This is answered in the affirmative in the case of the binary 3-cube Q_3 , but at the moment, that is the only answer we are able to offer. We conjecture this is the only existing totally efficient coloring.

Theorem 1. *The binary 3-cube graph Q_3 admits a totally efficient coloring via color set $[\kappa]$, for $\kappa = 3$.*

Proof. The claimed coloring is depicted on the upper-left of Figure 1. Observe that each 4-cycle of Q_3 (a girth cycle) has its edges in bijective correspondence with color set $[r] = [4]$, and its vertices also in bijective correspondence with color set $[r] = [4]$, where color 0 is hazel, color 1 is red, color 2 is blue, color 3 is green. \square

It is not difficult to see that a totally efficient coloring of a graph $G = (V, E)$ partitions the vertex set V into *efficient dominating sets*, or *perfect codes*, that is vertex subsets S such that every vertex of V is at distance no larger than 1 from S . In our case, such partition is composed by the colored subsets




Figure 1: Totally effective coloring of Q_3 and associated proper edge coloring of $K_{4,4}$

$$C_0 = \{000, 111\}, C_1 = \{100, 011\}, C_2 = \{010, 101\}, C_3 = \{001, 110\}, \quad (1)$$

that is the pairs of opposite, antipodal, or complementary, vertices in colors hazel, red, blue and green, respectively, and/or numbers 0,1,2,3, respectively, in the figure. In addition, a 1-factor exists in each of the 6-cycles forming the complements $Q_3 \setminus C_i$, ($i = 0, 1, 2, 3$), of the pairs C_i in Q_3 , as depicted in the figure with the color assignment adopted for the vertices

of Q_3 extended to the edges of Q_3 . Such color assignment takes care of all the edges of Q_3 , as follows:

$$\begin{aligned}
\{(010, 110), (100, 101), (001, 011)\} &\subset E(Q_3) \setminus C_0 \\
\{(000, 010), (011, 111), (100, 110)\} &\subset E(Q_3) \setminus C_1 \\
\{(000, 001), (011, 111), (001, 011)\} &\subset E(Q_3) \setminus C_2 \\
\{(000, 100), (010, 011), (101, 111)\} &\subset E(Q_3) \setminus C_3
\end{aligned} \tag{2}$$

Let Γ be a finite connected κ -regular simple graph with chromatic index $\chi'(\Gamma) = \kappa$. Let $g = g(\Gamma)$ be the girth of Γ . We say that Γ is a *g -tight graph* if $g = g(\Gamma) = \kappa = \chi'(\Gamma)$. In each g -tight graph Γ , it makes sense to look for a proper edge-coloring via κ colors, each girth cycle colored via a bijection between the cycle edges and the colors they are assigned, precisely κ colors. We will say that such a coloring is an *edge-girth coloring* of Γ and, in such a case, that Γ is *edge-girth chromatic*, or *egc* for short.

In [3], focus was on g -tight κ -regular graphs Γ that are *girth-regular* [5], adapted as follows. Let Γ be one such graph. Let $\{e_1, \dots, e_\kappa\}$ be the set of edges incident in Γ to a vertex v . Let (e_i) be the number of κ -cycles containing an edge e_i , for $1 \leq i \leq \kappa$. Assume $(e_1) \geq (e_2) \geq \dots \geq (e_\kappa)$. Let the *signature* of v be the κ -tuple $((e_1), (e_2), \dots, (e_\kappa))$. The graph Γ is said to be *girth-regular* if all its vertices have a common signature. In such a case, the signature of any vertex of Γ is said to be the *signature* of Γ .

Girth-regular graphs that are g -tight are said to be *girth- g -regular graphs* as well as $((e_1), (e_2), \dots, (e_g))$ -graphs, or $(e_1)(e_2) \cdots (e_g)$ -graphs, if no confusion arises, where $g = \kappa$. In this notation, a prefix

$$a_1^{(1)} a_1^{(2)} \cdots a_1^{(m_1)} a_2^{(1)} a_2^{(2)} \cdots a_2^{(m_2)} \cdots a_t^{(1)} a_t^{(2)} \cdots a_t^{(m_t)} \tag{3}$$

with $a_i^{(j)} = a_{i'}^{(j')}$ iff $i = i'$ may be abbreviated as $a_1^{m_1} a_2^{m_2} \cdots a_t^{m_t}$, where superscripts equal to 1 may be omitted (e.g., 3221 abbreviates to 32^21).

Edge-girth colorings of $(e_1)(e_2) \cdots (e_g)$ -graphs are equivalent to 1-factorizations [6] such that the cardinality of the intersection of each 1-factor with each girth cycle is 1. These factorizations are said to be *tight*, and the resulting colored girth cycles, are said to be *tightly colored*. Notice that a Γ as above with a tight factorization is egc.

Corollary 2. *The complete bipartite graph $K_{4,4}$ is a girth-4-regular graph with signature 3^4 obtained from Q_3 by adding the antipodal edges*

$$e_0 = (000, 111), e_1 = (100, 011), e_2 = (010, 101), e_3 = (001, 110), \tag{4}$$

associated to the perfect codes C_0, C_1, C_2, C_3 in display (1), respectively. The end-vertices of e_i have common color i in the totally effective coloring of Q_3 obtained in Theorem 1, for $i = 0, 1, 2, 3$. By assigning color i to edge e_i , for $i = 0, 1, 2, 3$, an edge-girth coloring of $K_{4,4}$ is obtained, which is egc.

Proof. On the upper right of Figure 1, a copy of $K_{4,4}$ is obtained by adding the edges e_i , ($i = 0, 1, 2, 3$) to the copy of Q_3 depicted to its left in the figure. Such addition was done carrying the shown totally efficient coloring; by recognizing that the two vertices of each C_i have a common color r_i , then, the corresponding e_i also is assigned the color r_i , yielding the claimed edge-girth coloring of $K_{4,4}$. \square

From the upper right of Figure 1, three mutually orthogonal Latin squares of order 4, or MOLS(4), [1] are obtained, namely

	1	2	4	7	
0	132	213	321	000	
3	220	101	033	312	(5)
5	303	022	110	231	
6	011	330	202	123	

where the rows are headed by the even-weight vertices x_{ew} , namely (in both decimal and 3-binary-digit notation)

$$0 = 000, 3 = 011, 5 = 101, 6 = 110,$$

and the columns are headed by the odd-weight vertices x_{ow} , namely

$$1 = 001, 2 = 010, 4 = 100, 7 = 111,$$

and where each triple in (5) corresponding to a row-and-column position contains: **(i)** the directional color number of the edge (x_{ew}, x_{ow}) (namely: 1=horizontal, 2=depth, 3=vertical and 0=antipodal); **(ii)** the actual color number of such edge; and **(iii)** a complementary color that contributes in the edge-girth coloring of the binary 4-cube in Corollary 3, represented in the lower portion of Figure 1.

Corollary 3. *The binary 4-cube Q_4 is a girth-regular graph with signature 3^4 and an edge-girth coloring obtained from that of Q_3 in Theorem 1 as an extension. Moreover, let $\psi : Q_4 \rightarrow K_{4,4}$ be the graph map that sends antipodal elements (i.e., vertices, edges) of Q_4 into corresponding elements of $K_{4,4}$ considered as antipodal classes mod \mathbb{Z}_2 of such elements. Then, the edge-girth coloring of Q_4 is obtained via the inverse image ψ^{-1} of the edge-girth coloring of $K_{4,4}$.*

Proof. Let the vertices of the binary 4-cube or binary tesseract Q_4 being denoted as follows:

$$\begin{aligned} 0 &= 0000, 1 = 1000, 2 = 0100, 3 = 1100, 4 = 0010, 5 = 1010, 6 = 0110, 7 = 1110, \\ \underline{0} &= 0001, \underline{1} = 1001, \underline{2} = 0101, \underline{3} = 1101, \underline{4} = 0011, \underline{5} = 1011, \underline{6} = 0111, \underline{7} = 1111. \end{aligned} \quad (6)$$

On the lower part of Figure 1, the totally effective colored copy H of Q_3 in the upper left of the figure is reproduced as an inner 3-cube $\phi(H)$ in a copy G of Q_4 , where $\phi : H \rightarrow G$ is inclusion graph map, with radial edges from each vertex x of $\phi(H)$ to the corresponding vertex \underline{x} of $G \setminus \phi(H)$. An edge-girth coloring of G is obtained by extending the image via ϕ of the totally effective coloring of $Q_3 = H$ in Theorem 1 by assigning to each radial edge the color of its incident vertex in $\phi(H)$. Then, there is a unique color compatible with the already assigned edge colors for each edge of $G \setminus \phi(H)$, allowing to close the claimed edge-girth coloring of $G = Q_4$. In particular, any two edges of Q_4 that are antipodal to each other bear a common color. Now, $\phi\psi$ is the inclusion graph map of $H = Q_3$ into $K_{4,4}$, where the edges of $K_{4,4} \setminus Q_3$ receive the colors of the their endvertices, which is the same color for each such edge. This allows to establish the last assertion of the statement. \square

2 A related problem via a star transposition graph

Let G be a graph of regular degree $\kappa \geq 3$, girth 2κ and chromatic index $r = \kappa + 1$. Does G possess a totally efficient coloring with r colors? This is answered in the affirmative in the case of the star transposition graph ST_4 [2] represented as a toroidal cutout both on the left and right of Figure 2 (to be used in the proof of Theorem 4), with the edges traced thick black and the uniformly colored bright areas (green, blue, yellow) having each a central 6-cycle with darker interior (denoted a3, b3, c3, respectively). The straight borders between the colored bright areas are indicated as dashed-traced segments, with their extremes being the members of the efficient dominating set given by $N(a3) = N(b3) = N(c3) = \{3120, 3102, 3201, 3210, 3012, 3021\}$. In fact, each 6-cycle xj indicates the value $j \in \{0, 1, 2, 3\}$ assumed by position $x \in \{a, b, c\}$ in the six existing vertices of the cycle, where $x \in \{a, b, c\}$ indicates respectively the second, third and fourth entries of 4-permutations $y_0y_ay_by_c$ such that $\{y_0, y_a, y_b, y_c\} = \{0, 1, 2, 3\}$.




Figure 2: Pair of vertex-fixed edge-orthogonal totally efficient colorings of ST_4

Theorem 4. *The star transposition graph ST_4 admits a totally efficient coloring via the color set $[\kappa]$, for $\kappa = 3$, in two vertex-fixed edge-orthogonally instances that can be combined to produce an edge-girth coloring of the prism graph of ST_4 .*

Proof. Throughout, each vertex $y_0y_ay_by_c$ of ST_4 is assigned color y_0 , which insures that the vertex set of ST_4 has a partition into efficient dominating sets [2]. On the other hand, each edge of ST_4 separates two faces of adjacent 6-cycles. Also, the edges of each 6-cycle determine two 2-factors of three edges each. An edge coloring of ST_4 is obtained by selecting in each 6-cycle xj one of its 1-factors, say F , and assigning color j to the three edges of F in such a way that all edges of ST_4 get a well-defined color. More specifically, the 6-cycle xj has the

edges of its remaining 1-factor F' (so $F \cup F' = E(xj)$), with the colors j_1, j_3, j_5 of the three alternate adjacent 6-cycles xj_1, xj_3, xj_5 of F' , where the associated edge color sequence of xj is $(j_0, j_1, j_2, j_3, j_4, j_5)$, with j_0, j_2, j_4 being the colors of the edges of F . This produces a totally efficient coloring C of ST_4 while determining which 2-factors of the 6-cycles xj of ST_4 take the roles of F and of F' . Now, by exchanging the roles of the F 's and F' 's in the argument above, a second totally efficient coloring of ST_4 is obtained, that behaves as a similar vertex coloring as that of C but that is edge-orthogonal to that of C in the sense that the roles of the 2-factors F and F' are exchanged. The resulting pair of vertex-fixed edge-orthogonal totally efficient colorings of ST_4 is shown in Figure 2, where the red edge color numbers behave in our sense of orthogonality while the red vertex color numbers remain fixed. This edge-orthogonality allows to consider the prism $ST_4 \square K_2$, assign the vertex colors of ST_4 to the corresponding edges of the copies $\{v\} \square K_2$ of K_2 , ($v \in V(ST_4)$), where the endvertices of each such edge have a common color, and take the orthogonality to justify the claimed edge-girth coloring. \square




Figure 3: Another view of the previous figure

Figure 3 shows a modification of Figure 2 to stress the difference between the two vertex-fixed edge-orthogonal colorings by means of colored triangles whose vertices are the edge color numbers, so on the left of the figure these triangles are pointing down while on the right they point up. The colors of these triangles are; light-gray in a_0, b_0, c_0 ; light-green in a_1, b_1, c_1 ; light-blue in a_2, b_2, c_2 ; yellow in a_3, b_3, c_3 .

Corollary 5. *The result of Theorem 4 is extended periodically to any larger toroidal cutout in the plane formed by continuation of the toroidal cutout in Theorem 4, up to the whole euclidean plane.*

Proof. A continuation of the two orthogonal totally effective colorings of ST_4 to any case as in the statement is sufficient to complete its proof. \square

Theorem 1 and Corollaries 2-3 can also be combined to look formally like Theorem 4, as follows.

Theorem 6. *The binary 3-cube graph Q_3 admits a totally efficient coloring via the color set $[\kappa]$, for $\kappa = 3$, in two vertex-fixed edge-orthogonally instances that can be combined to produce an edge-girth coloring of the binary 4-cube Q_4 .*

Proof. Both in Figure 1 and display (5), the four 6-cycles of Q_3 avoiding a perfect code C_i are seen showing their 2-factors. We were using their fixed-vertex edge-orthogonality properties in a way similar to that of Theorem 4. In this context, $K_{4,4}$ replaces the prism of Theorem 5 in the context of Section 1. \square

References

- [1] C. J. Colbourn, J. H. Dinitz, Handbook of Combinatorial Designs (2nd ed.), Boca Raton: Chapman & Hall/ CRC, (2007).
- [2] I. J. Dejter and O. Serra, *Efficient dominating sets in Cayley graphs*, Discrete Appl. Math., **129** (2003), 319–328.
- [3] I. J. Dejter, *Tight factorizations of girth- g -regular graphs*, Arxiv 2102.06956, (2024).
- [4] J. Geetha, N. Narayanan and K. Somasundaram, Total coloring-a survey, AKCE int. Jour. of Graphs and Combin., **20**, (2023), issue 3. 339-351.
- [5] P. Potočník, J. Vidali, *Girth-regular graphs*, Ars Mathematica Contemporanea, **17** (2019), 349–368.
- [6] W. D. Wallis, One-Factorizations (Math. and its Appl.), Springer, NY, 1997.