A NOTE ON CONVERGENCE OF DENSITIES TO FREE EXTREME VALUE DISTRIBUTIONS

YAMATO KINDAICHI AND YUKI UEDA

ABSTRACT. The concept of free extreme value distributions as universal limit laws for the spectral maximum of free noncommutative real random variables was discovered by Ben Arous and Voiculescu in 2006. This paper contributes to study the convergence of densities towards free extreme value distributions under the von Mises condition for sample distributions.

1. Introduction

Extreme value theory plays a pivotal role in analyzing maximum or minimum within a large dataset of samples. Its most notable feature is its universality, as established by the Fisher-Tippett-Gnedenko theorem [7, 8] (see also [19]). This theorem asserts that the limit laws of the maximum for a large number of samples are classified into only three types: Fréchet, Weibull and Gumbel distributions. The study of convergence towards extreme value distributions is one of significant topics in pure mathematics of extreme value theory. Smith [20] and Omey [18] explored the rate of uniform convergence in the Fréchet case. Subsequently, Bartholmé and Swan [2] also examined similar outcomes using Stein's method. Hall [13] has already studied the rate of uniform convergence between extreme values in normal i.i.d. random variables and the Gumbel distributions. More comprehensively, Kusumoto and Takeuchi [15] employed Stein's method to analyze the rate of uniform convergence in the Gumbel case. Generally, convergence in distributions does not necessarily imply the convergence of their corresponding densities (even if densities exist). Haan and Resnick [12] proved that the normalize maximum converges to the densities of extreme value distributions under the so-called von Mises conditions. Additionally, Omey [18] investigated the rate of convergence for the density of normalized maximum towards extreme values under certain regularly varying conditions for a sample distribution.

In the framework of Voiculescu's free probability theory (see [16, 25] for details), Ben Arous and Voiculescu [4] developed a theory to analyze the maximum of a large number of non-commutative real random variables on some W^* -probability space (i.e. selfadjoint operators affiliated with some von Neumann algebra). Let (\mathcal{M}, τ) be a W^* -probability space. According to [4], for any free non-commutative real random variables X, Y on (\mathcal{M}, τ) , the spectral distribution function $F_{X \vee Y}(x) = \tau(\mathbf{1}_{(-\infty,x]}(X \vee Y))$ of the maximum $X \vee Y$ (with respect to spectral order [1, 17]) is given by

$$F_{X\vee Y}(x) = \max\{F_X(x) + F_Y(x) - 1, 0\} =: (F_X \boxtimes F_Y)(x),$$

²⁰¹⁰ Mathematics Subject Classification. Primary: 46L54; Secondary: 60B10, 62G32.

Key words and phrases. free extreme value distribution, convergence of densities and von Mises condition.

where $F_X(x) = \tau(\mathbf{1}_{(-\infty,x]}(X))$ and F_Y is similarly defined. In general, for any distribution functions F and G on \mathbb{R} , we define

$$F \boxtimes G := \max\{F + G - 1, 0\}$$
 and $F^{\boxtimes n} := \underbrace{F \boxtimes \cdots \boxtimes F}_{n \text{ times}}.$

The operation \square is called the *free max-convolution*. Similarly as classical extreme value theory, the limiting distributions of the normalized maximum $W_n = ((X_1 \vee \cdots \vee X_n) - b_n)/a_n$ (for some $a_n > 0$ and $b_n \in \mathbb{R}$) of a large number of freely independent, identically distributed non-commutative real random variables X_1, \ldots, X_n on some W^* -probability space, are also characterized by just three types:

$$\Phi_{\alpha}^{\text{free}}(x) := \begin{cases} (1-x^{-\alpha})\mathbf{1}_{[1,\infty)}(x), & \alpha > 0 \quad \text{(free Fr\'echet distribution);} \\ \{1-(-x)^{-\alpha}\}\mathbf{1}_{[-1,0]}(x)+\mathbf{1}_{(0,\infty)}(x), & \alpha < 0 \quad \text{(free Weibull distribution);} \\ (1-e^{-x})\mathbf{1}_{[0,\infty)}(x) & \alpha = 0 \quad \text{(free Gumbel distribution).} \end{cases}$$

The distributions $\Phi_{\alpha}^{\text{free}}$ are called the *free extreme value distributions*. Rigorously, for a distribution function F on \mathbb{R} , there exist $a_n > 0$, $b_n \in \mathbb{R}$ and a non-degenerate distribution function G such that

$$F^{\boxtimes n}(a_n x + b_n) \to G(x)$$
, as $n \to \infty$, $x \in \mathbb{R}$,

then G is the free extreme value distribution.

Since [4], free extreme value theory has continued to advance from various perspectives. Ben Arous and Kargin [3] clarified the relationships between the convergence of free point processes and the max-domains of attraction of free extreme value distributions. Benaych-Georges and Cabanal-Duvillard [5] established a certain random matrix model such that the empirical eigenvalue distributions of random matrices converge to the free extreme value distribution as the matrix size tends to infinity. Grela and Nowak [9] provided explicit calculations for several extreme random matrix ensembles. Also they presented a formula, showing the equivalence of free extreme distributions to the POT (Peak-Over-Threshold) method in classical probability. Recently, the author [23] has discovered a close relationship between limit theorems for classical and free max-convolutions. This results are the max-analogue presented in Bercovici and Pata's seminal work [6]. In [24] and [14] (with Hasebe), the author further found interesting relations between additive convolution semigroups and max-convolution semigroups in classical, free and boolean cases via Tucci, Haagerup and Möller limit theorem [21, 10].

In accordance with the author's work [22], the uniform convergence of $F^{\boxtimes n}(a_n \cdot +b_n)$ to the free extreme value distribution G was proven under certain analytic assumptions regarding the density of the normalized maximum W_n , and the rate of convergence has been provided via Stein's method. We prompt a natural question here. Can we assert the convergence of densities of W_n towards the density of free extreme value distribution? To be more precise, if the density function w_n of the normalized maximum W_n exists for almost all $n \in \mathbb{N}$, our inquiry pertains to whether $w_n(x)$ converges to the density of free extreme value distributions as characterized by

$$\varphi_{\alpha}^{\text{free}}(x) := \begin{cases} \alpha x^{-\alpha-1} \mathbf{1}_{(1,\infty)}(x), & \alpha > 0 \quad \text{(free Fr\'echet density);} \\ -\alpha (-x)^{-\alpha-1} \mathbf{1}_{(-1,0)}(x), & \alpha < 0 \quad \text{(free Weibull density);} \\ e^{-x} \mathbf{1}_{(0,\infty)}(x), & \alpha = 0 \quad \text{(free Gumbel density).} \end{cases}$$

In this paper, we establish the convergence of densities towards the free extreme values distributions, under the von Mises condition and a few analytic assumptions regarding sample distributions.

Theorem 1.1. Consider a distribution function $F = \exp\{-e^{-\phi}\}$ where ϕ is a function with some analytic assumptions. Let us define $h_{\alpha}(x)$ by

$$h_{\alpha}(x) := \begin{cases} x\phi'(x) - \alpha, & \alpha > 0, \ \omega_F = \infty; \\ (\omega_F - x)\phi'(x) + \alpha, & \alpha < 0, \ \omega_F < \infty; \\ (1/\phi'(x))', & \alpha = 0, \end{cases}$$

where $\omega_F := \sup\{x \in \mathbb{R} : F(x) < 1\}$ and we understand $\omega_F = \infty$ when F(x) < 1 for all $x \in \mathbb{R}$. Assume that there exists a non-increasing function g such that $g(x) \to 0$ as $x \to \omega_F$ and $|h_{\alpha}| \leq g$ (von Mises condition). Then we obtain the followings.

• The case $\alpha > 0$ and $\omega_F = \infty$. If $a_n > 0$ satisfies $F(a_n) = e^{-\frac{1}{n}}$, then

$$\sup_{x>1} |w_n(x) - \varphi_\alpha^{\text{free}}(x)| \le O(n^{-1} \vee g(a_n)).$$

• The case $\alpha < 0$ and $\omega_F < \infty$. Let us consider $\alpha < -1$. If $a_n > 0$ satisfies $F(\omega_F - a_n) = e^{-\frac{1}{n}}$ and $b_n = \omega_F$, then

$$\sup_{-1 \le x \le 0} |w_n(x) - \varphi_\alpha^{\text{free}}(x)| \le O(n^{-1} \lor g(\omega_F - a_n)).$$

When $\alpha < 0$ in general, it is true when the supremum runs over any compact sets in (-1,0).

• The case $\alpha = 0$. If $b_n \in \mathbb{R}$ satisfies $F(b_n) = e^{-\frac{1}{n}}$ and $a_n = \frac{F(b_n)}{nF'(b_n)}$, then

$$\sup_{x>0} |w_n(x) - \varphi_0^{\text{free}}(x)| \le O(n^{-1} \vee g(b_n)).$$

For details, see Theorem 2.5 for $\alpha > 0$, Theorem 3.4 and Corollary 3.5 for $\alpha < 0$ and Theorem 4.5 for $\alpha = 0$.

2. The case
$$\alpha > 0$$
 (Fréchet type)

In this section, we consider a distribution function F defined as $F = \exp\left\{-e^{-\phi}\right\}$, where $\omega_F = \infty$ and ϕ is a C^1 -function with $\phi' > 0$ in a neighborhood of ∞ . Choose $a_n = F^{\leftarrow}(e^{-\frac{1}{n}})$, where $F^{\leftarrow}(y) := \inf\{x \in \mathbb{R} : F(x) \geq y\}$. Since F is continuous, we get $F(a_n) = e^{-\frac{1}{n}}$, equivalently, $\phi(a_n) = \log n$. Since ϕ is differentiable, the function $x \mapsto F^{\boxtimes n}(a_n x)$ has the density function w_n given by

$$w_n(x) = na_n F'(a_n x) = na_n \phi'(a_n x)(-\log F(a_n x))F(a_n x),$$

for all $x > A_n := a_n^{-1} F^{\leftarrow}(1 - n^{-1}) \in (-\infty, 1)$. Since $a_n \sim F^{\leftarrow}(1 - n^{-1})$, we get $A_n \to 1$ as $n \to \infty$. We further assume the von Mises condition for sample distributions F.

Assumption 2.1. For $\alpha > 0$, we define

$$h_{\alpha}(x) := x\phi'(x) - \alpha = \frac{xF'(x)}{F(x)(-\log F(x))} - \alpha, \qquad x > 0.$$

Assume $h_{\alpha}(x) \to 0$ as $x \to \infty$ (see [19, (2.42)]).

Under Assumption 2.1, we get $F^n(a_n \cdot) \xrightarrow{w} \Phi_{\alpha}$ by [19, Pages 107–108], where $\Phi_{\alpha}(x) = \exp(-x^{-\alpha})\mathbf{1}_{(0,\infty)}(x)$ is the Fréchet distribution. Due to [4, Theorem 6.12], one can also see that $F^{\boxtimes n}(a_n \cdot) \xrightarrow{w} \Phi_{\alpha}^{\text{free}}$. We summarize the above discussion as follows.

Lemma 2.2. Under Assumption 2.1, we have $F^{\boxtimes n}(a_n \cdot) \xrightarrow{w} \Phi_{\alpha}^{\text{free}}$.

According to the above discussion, we are interested in the following class of sample distributions.

Definition 2.3. Let us consider $\alpha > 0$. Denote by \mathcal{F}_{α} the set of all distribution functions F such that

- (F-1) $F = \exp\{-e^{-\phi}\}$, where $\omega_F = \infty$ and ϕ is a C^1 -function with $\phi' > 0$ in a neighborhood of ∞ ;
- (F-2) there exists a nonincreasing continuous function g on $(0, \infty)$ such that $g(x) \to 0$ as $x \to \infty$ and $|h_{\alpha}(x)| \le g(x)$ for all x > 0.

Obviously, Assumption 2.1 achieves from the condition (F-2). Therefore if $F \in \mathcal{F}_{\alpha}$ then $F^{\boxtimes n}(a_n \cdot) \xrightarrow{w} \Phi_{\alpha}^{\text{free}}$ by Lemma 2.2.

For any $F \in \mathcal{F}_{\alpha}$, we prove the uniform convergence of the density w_n of $x \mapsto F^{\boxtimes n}(a_n x)$ to the free Fréchet density $\varphi_{\alpha}^{\text{free}}$ on $(1, \infty)$ that is a domain in which $\varphi_{\alpha}^{\text{free}} > 0$. To show this, we prepare the following inequality for the free Fréchet distributions.

Lemma 2.4. For any
$$\alpha_1, \alpha_2 > 0$$
, we have $\sup_{x \in \mathbb{R}} |\Phi_{\alpha_1}^{\text{free}}(x) - \Phi_{\alpha_2}^{\text{free}}(x)| \le e^{-1} \frac{|\alpha_2 - \alpha_1|}{\alpha_1 \vee \alpha_2}$.

Proof. Without loss of generality, we may assume that $\alpha_1 < \alpha_2$. Note that $|\Phi_{\alpha_1}^{\text{free}}(x) - \Phi_{\alpha_2}^{\text{free}}(x)| = 0$ for all $x \leq 1$. Since $\frac{\partial}{\partial \beta} x^{-\beta} = -x^{-\beta} \log x$ for $\beta > 0$ and x > 1, we get

$$\begin{split} \sup_{x>1} |\Phi_{\alpha_1}^{\text{free}}(x) - \Phi_{\alpha_2}^{\text{free}}(x)| &= \sup_{x>1} |x^{-\alpha_1} - x^{-\alpha_2}| \\ &\leq \sup_{x>1} \sup_{\beta \in [\alpha_1, \alpha_2]} |-x^{-\beta} \log x| |\alpha_2 - \alpha_1| \\ &= (\alpha_2 - \alpha_1) \sup_{x>1} x^{-\alpha_2} \log x = e^{-1} \ \frac{\alpha_2 - \alpha_1}{\alpha_2}. \end{split}$$

Theorem 2.5. Let $F \in \mathcal{F}_{\alpha}$, a_n and g be defined as above. Then w_n converges uniformly to the free Fréchet density $\varphi_{\alpha}^{\text{free}}$ on $(1, \infty)$. More precisely, we get

$$\sup_{x>1} |w_n(x) - \varphi_\alpha^{\text{free}}(x)| \le O(n^{-1} \vee g(a_n)),$$

for sufficiently large n.

Proof. For all x > 1 and for sufficiently large $n \in \mathbb{N}$ (such that infinitely many a_n are in a neighborhood of ∞), we have

$$|w_{n}(x) - \varphi_{\alpha}^{\text{free}}(x)| \leq |na_{n}\phi'(a_{n}x)(-\log F(a_{n}x)) - \alpha x^{-\alpha-1}| + \alpha x^{-\alpha-1}(1 - F(a_{n}x))$$

$$\leq \underbrace{|na_{n}\phi'(a_{n}x)(-\log F(a_{n}x)) - \alpha x^{-\alpha-1}|}_{=:I} + \alpha(1 - e^{-\frac{1}{n}}).$$

On the other hands, we obtain

$$I \leq \left| \frac{n(-\log F(a_n x))}{x} (a_n x \phi'(a_n x) - \alpha) \right| + \left| \frac{(n(-\log F(a_n x)))}{x} \alpha - \alpha x^{-\alpha - 1} \right|$$

$$\leq g(a_n x) \frac{n \exp(-\phi(a_n x))}{x} + |n(-\log F(a_n x)) - x^{-\alpha}| \alpha x^{-1} \qquad \text{(by condition (F-2))}$$

$$\leq g(a_n) + \alpha \underbrace{|n(-\log F(a_n x)) - x^{-\alpha}|}_{=:J},$$

where the last inequality holds since $\phi' > 0$ in a neighborhood of ∞ and

$$\frac{n \exp(-\phi(a_n x))}{x} \le n \exp(-\phi(a_n)) = 1.$$

According to [19, Page 108], for $x \ge 1$, we have

$$(\alpha - g(a_n)) \log x \le \phi(a_n x) - \phi(a_n) \le (\alpha + g(a_n)) \log x,$$

which implies that $\Phi_{\alpha-g(a_n)}^{\text{free}}(x) \leq 1 + n \log F(a_n x) \leq \Phi_{\alpha+g(a_n)}^{\text{free}}(x)$. Thus,

$$\Phi_{\alpha}^{\text{free}}(x) - \Phi_{\alpha+g(a_n)}^{\text{free}}(x) \le n(-\log F(a_n x)) - x^{-\alpha} \le \Phi_{\alpha}^{\text{free}}(x) - \Phi_{\alpha-g(a_n)}^{\text{free}}(x)$$

for all $x \geq 1$, and therefore

$$J \le e^{-1} \frac{2g(a_n)}{\alpha + g(a_n)} \le \frac{2}{\alpha e} g(a_n)$$

by Lemma 2.4. Hence, the desired result holds true.

Example 2.6. As below, since all F are differentiable, we take $a_n > 0$ such that $F(a_n) = e^{-\frac{1}{n}}$. Let w_n be the density of the function $x \mapsto F^{\boxtimes n}(a_n x)$.

(1) (Fréchet distribution) Consider $F = \Phi_{\alpha}$ for $\alpha > 0$. The choice of a_n leads to $a_n = n^{\frac{1}{\alpha}}$ for $n \in \mathbb{N}$. As per the definition of h_{α} , we have $h_{\alpha}(x) = 0$. Hence,

$$\sup_{x>1} |w_n(x) - \varphi_\alpha^{\text{free}}(x)| \le O(n^{-1})$$

by Theorem 2.5. Note that the uniform convergence of w_n to $\varphi_{\alpha}^{\text{free}}$ on $(-\infty, 1]$ is not true. Actually, since $A_n = (-n \log(1 - n^{-1}))^{-\frac{1}{\alpha}}$, we get

$$w_n(A_n+) = \alpha \left\{ -n \log \left(1 - n^{-1} \right) \right\}^{1 + \frac{1}{\alpha}} \left(1 - n^{-1} \right) \to \alpha$$

as $n \to \infty$. Thus, we conclude

$$\lim_{n \to \infty} \sup_{A_n < x \le 1} |w_n(x) - \varphi_\alpha^{\text{free}}(x)| = \lim_{n \to \infty} \sup_{A_n < x \le 1} |w_n(x)| \ne 0.$$

(2) (Log-logistic distribution) Consider $\alpha > 0$ and $F(x) = \{1 - (1 + x^{\alpha})^{-1}\}\mathbf{1}_{(0,\infty)}(x)$. The choice of $a_n > 0$ leads to $a_n = (e^{\frac{1}{n}} - 1)^{-\frac{1}{\alpha}}$ for $n \in \mathbb{N}$. By definition of h_{α} , one can see

$$|h_{\alpha}(x)| = \alpha - \frac{\alpha}{(1+x^{\alpha})(-\log(1-(1+x^{\alpha})^{-1}))} \le \frac{\alpha}{1+x^{\alpha}} =: g(x), \quad x > 0,$$

and $g(a_n) = \alpha(1 - e^{-\frac{1}{n}})$. According to Theorem 2.5, we also obtain

$$\sup_{x>1} |w_n(x) - \varphi_\alpha^{\text{free}}(x)| \le O(n^{-1})$$

(3) (Cauchy distribution) Consider $F(x) = \frac{1}{2} + \frac{1}{\pi} \operatorname{Tan}^{-1} x$ for $x \in \mathbb{R}$. The choice of a_n leads to $a_n = \tan(\pi e^{-\frac{1}{n}} - \frac{\pi}{2})$ for $n \in \mathbb{N}$. One can see that $F^{\boxtimes n}(a_n \cdot) \xrightarrow{w} \Phi_1^{\text{free}}$. By definition of h_{α} , we observe

$$|h_1(x)| = \left| \frac{x}{\pi(1+x^2)\left(\frac{1}{2} + \frac{1}{\pi}\operatorname{Tan}^{-1}x\right)\left(-\log\left(\frac{1}{2} + \frac{1}{\pi}\operatorname{Tan}^{-1}x\right)\right)} - 1 \right|$$

$$\leq 1 - \frac{x}{\pi(1+x^2)\left(-\log\left(\frac{1}{2} + \frac{1}{\pi}\operatorname{Tan}^{-1}x\right)\right)} =: g(x)$$

for any x > 0. Furthermore, $g(x) \to 0$ as $x \to \infty$. This establishes

$$g(a_n) = 1 + \frac{n}{2\pi} \sin(2\pi e^{-\frac{1}{n}})$$

$$= 1 + \frac{n}{2\pi} \sin\left(2\pi - \frac{2\pi}{n} + \frac{\pi}{n^2} + O(n^{-3})\right)$$

$$= 1 + \frac{n}{2\pi} \sin\left(-\frac{2\pi}{n} + \frac{\pi}{n^2} + O(n^{-3})\right)$$

$$= 1 + \frac{n}{2\pi} \left\{ \left(-\frac{2\pi}{n} + \frac{\pi}{n^2} + O(n^{-3})\right) + O(n^{-3}) \right\}$$

$$= \frac{1}{2n} + O(n^{-2}).$$

Therefore we get $\sup_{x>1} |w_n(x) - \varphi_1^{\text{free}}(x)| \leq O(n^{-1})$ by Theorem 2.5.

3. The case
$$\alpha < 0$$
 (Weibull Type)

In this section, we consider a distribution function F defined as $F = \exp\left\{-e^{-\phi}\right\}$, where $\omega_F < \infty$ and ϕ is a C^1 -function with $\phi' > 0$ in a left neighborhood of ω_F . Define $a'_n = F^{\leftarrow}(e^{-\frac{1}{n}})$. Thus we have $F(a'_n) = e^{-\frac{1}{n}}$, equivalently, $\phi(a'_n) = \log n$. Let us set $a_n = \omega_F - a'_n$ and $b_n = \omega_F$ for all $n \in \mathbb{N}$. Since ϕ is differentiable, the density w_n of the function $x \mapsto F^{\boxtimes n}(a_n x + b_n)$ exists and is given by

$$w_n(x) = na_n \phi'(a_n x + b_n)(-\log F(a_n x + b_n))F(a_n x + b_n),$$

for all $A_n < x < 0$, where

$$A_n = \frac{F^{\leftarrow}(1 - n^{-1}) - b_n}{a_n}$$
 and $A_n \to -1$ as $n \to \infty$.

Similarly as in Fréchet case, we assume the von Mises condition in Weibull case.

Assumption 3.1. For $\alpha < 0$, we define

$$h_{\alpha}(x) := (\omega_F - x)\phi'(x) + \alpha = \frac{(\omega_F - x)F'(x)}{F(x)(-\log F(x))} + \alpha.$$

Assume $h_{\alpha}(x) \to 0$ as $x \to \omega_F - 0$.

Under Assumption 3.1 and [19, Pages 59-60], we get $F^n(a_n \cdot +b_n) \xrightarrow{w} \Phi_{\alpha}$ as $n \to \infty$, where $\Phi_{\alpha}(x) := \exp(-(-x)^{-\alpha})\mathbf{1}_{(-\infty,0)}(x) + \mathbf{1}_{[0,\infty)}(x)$ is the Weibull distribution. Due to [4], it is equivalent to $F^{\boxtimes n}(a_n \cdot +b_n) \xrightarrow{w} \Phi_{\alpha}^{\text{free}}$ as $n \to \infty$.

Lemma 3.2. Under Assumption 3.1, we have $F^{\boxtimes n}(a_n \cdot +b_n) \xrightarrow{w} \Phi_{\alpha}^{\text{free}}$ as $n \to \infty$.

According to the above discussion, we set the collection of certain sample distributions.

Definition 3.3. Consider $\alpha < 0$. Denote by \mathcal{F}_{α} the set of all distribution functions F such that

- (W-1) $F = \exp\{-e^{-\phi}\}\$, where ϕ is a C^1 -function with $\phi' > 0$ in a left neighborhood of ω_F .
- (W-2) there exists a nonincreasing function g such that $g(x) \to 0$ as $x \to \omega_F$ and $|h_{\alpha}(x)| \le g(x)$ for all $x < \omega_F$.

It follows from Lemma 3.2 that $F \in \mathcal{F}_{\alpha}$ implies that $F^{\boxtimes n}(a_n \cdot +b_n) \xrightarrow{w} \Phi_{\alpha}^{\text{free}}$. The same strategy as in the Fréchet case can be used to prove density convergence in the Weibull case.

Theorem 3.4. Consider $\alpha < -1$. Let $F \in \mathcal{F}_{\alpha}$, a_n, b_n and g be defined as above. Then we have

$$\sup_{-1 < x < 0} |w_n(x) - \varphi_\alpha^{\text{free}}(x)| \le O(n^{-1} \lor g(\omega_F - a_n)),$$

for sufficiently large n.

Proof. Consider $\beta = -\alpha > 1$. Let us put $F_*(x) := F(\omega_F - x^{-1})$ and $\phi_*(x) := \phi(\omega_F - x^{-1})$. Then $F_* = \exp\{-e^{-\phi_*}\}$ and ϕ_* is a C^1 -function with $\phi'_* > 0$ in a neighborhood of ∞ . Moreover, we define the function $h^*_{\beta}(x) := x\phi'_*(x) - \beta$. Then we get

$$h_{\beta}^*(x) = h_{\alpha}(\omega_F - x^{-1}) \to 0$$
, as $x \to \infty$,

by Assumption 3.1. Thus h_{β}^* satisfies the von Mises condition in Fréchet case (see Assumption 2.1). And also, $|h_{\beta}^*(x)| \leq g(\omega_F - x^{-1})$ by (W-2). If we put $a_n^* := a_n^{-1}$, then $a_n^* = F_*^{\leftarrow}(e^{-\frac{1}{n}})$ (and hence $F_*(a_n^*) = e^{-\frac{1}{n}}$) and $F_*^{\boxtimes n}(a_n^* \cdot) \xrightarrow{w} \Phi_{\beta}^{\text{free}}$. Let w_n^* be the density of $x \mapsto F_*^{\boxtimes n}(a_n^*x)$. Since one can see that $w_n^*(x) = x^{-2}w_n(-x^{-1})$ and $\varphi_{\beta}^{\text{free}}(x) = x^{-2}\varphi_{\alpha}^{\text{free}}(-x^{-1})$, we have

(3.1)
$$\sup_{\substack{-1 < x < 0 \\ x > 1}} |w_n(x) - \varphi_\alpha^{\text{free}}(x)| = \sup_{\substack{x > 1 \\ x > 1}} |w_n(-x^{-1}) - \varphi_\alpha^{\text{free}}(-x^{-1})| = \sup_{\substack{x > 1 \\ x > 1}} x^2 |w_n^*(x) - \varphi_\beta^{\text{free}}(x)|.$$

For any x > 1, we get

$$\begin{split} x^{2}|w_{n}^{*}(x) - \varphi_{\beta}^{\text{free}}(x)| &= x^{2}|na_{n}^{*}\phi_{*}'(a_{n}^{*}x)(-\log F_{*}(a_{n}^{*}x))F_{*}(a_{n}^{*}x) - \beta x^{-\beta-1}| \\ &= x|(h_{\beta}^{*}(a_{n}^{*}x) + \beta)n(-\log F_{*}(a_{n}^{*}x))F_{*}(a_{n}^{*}x) - \beta x^{-\beta}| \\ &\leq xg(\omega_{F} - (a_{n}^{*}x)^{-1})n(-\log F_{*}(a_{n}^{*}x)) + \beta \underbrace{\lfloor nx(-\log F_{*}(a_{n}^{*}x)) - x^{-\beta+1} \rfloor}_{=:I} \\ &+ \beta x^{-\beta+1}(1 - F_{*}(a_{n}^{*}x)) \\ &\leq g(\omega_{F} - a_{n})nx \exp(-\phi_{*}(a_{n}^{*}x)) + \beta I + \beta(1 - e^{-\frac{1}{n}}). \end{split}$$

For sufficiently large n and for any x > 1, we get $\{x \exp(-\phi_*(a_n^*x))\}' = \exp(-\phi_*(a_n^*x))(1 - \beta - h_\alpha(a_n^*x)) < 0$, and therefore

$$nx \exp(-\phi_*(a_n^*x)) \le n \exp(-\phi_*(a_n^*)) = 1.$$

A proof similar to that of Lemma 2.4 yields that for $\beta_1, \beta_2 > 1$,

(3.2)
$$\sup_{x>1} |x\Phi_{\beta_1}(x) - x\Phi_{\beta_2}(x)| \le e^{-1} \frac{|\beta_2 - \beta_1|}{(\beta_1 - 1) \vee (\beta_2 - 1)}.$$

Since the inequality (2.1) implies that for any x > 1,

$$x\Phi_{\beta}^{\text{free}}(x) - x\Phi_{\beta+g(\omega_F-a_n)}^{\text{free}} \leq nx(-\log F_*(a_n^*x)) - x^{-\beta+1} \leq x\Phi_{\beta}^{\text{free}}(x) - x\Phi_{\beta-g(\omega_F-a_n)}^{\text{free}},$$

we get $I \leq e^{-1} \frac{2g(\omega_F - a_n)}{\beta + g(\omega_F - a_n) - 1} \leq \frac{2}{e(\beta - 1)} g(\omega_F - a_n)$ by (3.2). Finally, the equation (3.1) implies that

$$\sup_{-1 < x < 0} |w_n(x) - \varphi_{\alpha}^{\text{free}}(x)| \le \left(1 + \frac{2\beta}{e(\beta - 1)}\right) g(\omega_F - a_n) + \beta(1 - e^{-\frac{1}{n}}),$$

as desired. \Box

According to the above proof and Theorem 2.5, it is easy to show the locally uniformly convergence.

Corollary 3.5. Consider $\alpha < 0$. Under assumptions in Theorem 3.4, we get

$$\sup_{x \in K} |w_n(x) - \varphi_\alpha^{\text{free}}(x)| \le O(n^{-1} \vee g(\omega_F - a_n))$$

for any compact sets K in (-1,0).

Proof. Put $\beta = -\alpha > 0$ and use the symbols in the proof of Theorem 3.4. By Theorem 2.5, we have

$$\sup_{-1 < x < 0} x^2 |w_n(x) - \varphi_\alpha^{\text{free}}(x)| = \sup_{x > 1} |w_n^*(x) - \varphi_\beta^{\text{free}}(x)| \le O(n^{-1} \vee g(\omega_F - a_n)).$$

For any compact set $K = [a, b] \subset (-1, 0)$ and for any $x \in K$, we have

$$|w_n(x) - \varphi_\alpha^{\text{free}}(x)| \le \frac{1}{b^2} O(n^{-1} \vee g(\omega_F - a_n)),$$

as desired. \Box

Example 3.6. As below, since all F are differentiable, we take $a'_n \in \mathbb{R}$ such that $F(a'_n) = e^{-\frac{1}{n}}$ and put $a_n = \omega_F - a'_n$ and $b_n = \omega_F$. Let w_n be the density of $x \mapsto F^{\boxtimes n}(a_n x + b_n)$.

(1) (Weibull distribution) Consider $F = \Phi_{\alpha}$ for $\alpha < -1$. Then $\omega_F = 0$. The choice of a'_n leads to $a'_n = -n^{\frac{1}{\alpha}}$ for $n \in \mathbb{N}$. By definition of h_{α} , we get $h_{\alpha}(x) = 0$. By Theorem 3.4,

$$\sup_{-1 < x < 0} |w_n(x) - \varphi_{\alpha}^{\text{free}}(x)| \le O(n^{-1}).$$

(2) Let F be a distribution function such that $1 - F(x) = K(\omega_F - x)^{-\alpha}$ for some K > 0 and $\alpha < -1$. Since $F(a'_n) = e^{-\frac{1}{n}}$, we have $a'_n = \omega_F - K^{\frac{1}{\alpha}}(1 - e^{-\frac{1}{n}})^{-\frac{1}{\alpha}}$ (i.e. $a_n = K^{\frac{1}{\alpha}}(1 - e^{-\frac{1}{n}})^{-\frac{1}{\alpha}}$). By definition of h_{α} , we get

$$h_{\alpha}(x) = \frac{-\alpha K(\omega_F - x)^{-\alpha}}{(1 - K(\omega_F - x)^{-\alpha})(-\log(1 - K(\omega_F - x)^{-\alpha}))} + \alpha,$$

for all $\omega_F - K^{-\frac{1}{\alpha}} < x < \omega_F$. Then we define $g(x) = |h_{\alpha}(x)|$ and $g(\omega_F - a_n) = -\alpha n e^{\frac{1}{n}} (1 - e^{-\frac{1}{n}}) = O(n^{-1})$. Hence,

$$\sup_{-1 < x < 0} |w_n(x) - \varphi_\alpha^{\text{free}}(x)| \le O(n^{-1})$$

by Theorem 3.4.

(3) (Uniform distribution) We define $F(x) = x\mathbf{1}_{[0,1]}(x) + \mathbf{1}_{(1,\infty)}(x)$. Then $\omega_F = 1$. The choice of a'_n leads to $a'_n = e^{-\frac{1}{n}}$. If we take $a_n = 1 - e^{-\frac{1}{n}}$ and $b_n = 1$, then $F^{\boxtimes n}(a_n \cdot + b_n) \xrightarrow{w} \Phi_{-1}^{\text{free}}$. By definition of h_{α} , we get

$$h_{-1}(x) = \frac{1-x}{x(-\log x)} - 1 > 0, \quad 0 < x < 1,$$

and define $g(x) = h_{-1}(x)$. Then $g(\omega_F - a_n) = g(e^{-\frac{1}{n}}) = ne^{\frac{1}{n}}(1 - e^{-\frac{1}{n}}) - 1 = O(n^{-1})$, and therefore

$$\sup_{x \in K} |w_n(x) - \varphi_{-1}^{\text{free}}(x)| \le O(n^{-1})$$

for any compact sets K in (-1,0) by Corollary 3.5.

Remark 3.7. More strongly, for w_n in Example 3.6 (1), a numerical calculation tells us that the supremum of the absolute difference between $w_n(x)$ and $\varphi_{\alpha}^{\text{free}}(x)$ over the interval (-1,0) is bounded by $O(n^{-1})$ for any $\alpha < 0$. We expect that the uniformly convergence of w_n to $\varphi_{\alpha}^{\text{free}}$ on (-1,0) holds true for any $\alpha < 0$.

4. The case
$$\alpha = 0$$
 (Gumbel Type)

In this section, we suppose a distribution function F expressed by $F = \exp\{-e^{-\phi}\}$, where ϕ is a twice differentiable function with $\phi' > 0$ in a (left) neighborhood of $\omega_F \leq \infty$. We define

$$f(x) = \frac{1}{\phi'(x)}.$$

The function f is called an auxiliary function of F. Define $b_n = F^{\leftarrow}(e^{-\frac{1}{n}})$, leading to $F(b_n) = e^{-\frac{1}{n}}$, equivalently $\phi(b_n) = \log n$. Put $a_n := f(b_n) = \frac{F(b_n)}{nF'(b_n)}$. Since ϕ is differentiable, the function $x \mapsto F^{\boxtimes n}(a_n x + b_n)$ has the density w_n defined by

$$w_n(x) = na_n \phi'(a_n x + b_n)(-\log F(a_n x + b_n))F(a_n x + b_n),$$

for all $A_n < x < B_n$, where

$$A_n := \frac{F^{\leftarrow}(1 - n^{-1}) - b_n}{a_n} \quad \text{and} \quad A_n \to 0 \text{ as } n \to \infty,$$

$$B_n := \frac{\omega_F - b_n}{a_n}, \quad \text{and} \quad B_n \to \infty \text{ as } n \to \infty,$$

where we understand $B_n = \infty$ if $\omega_F = \infty$.

We now assume the von Mises condition for sample distributions F.

Assumption 4.1. We define

$$h_0(x) := f'(x) = -\log F(x) - \left\{ \frac{F(x)F''(x)(-\log F(x))}{(F'(x))^2} + 1 \right\}.$$

Assume $h_0(x) \to 0$ as $x \to \omega_F$ (see [19, (2.52)]).

According to [19, Pages 114], under Assumption 4.1, we have $F^n(a_n \cdot +b_n) \xrightarrow{w} \Phi_0$ as $n \to \infty$, where $\Phi_0(x) := \exp(-e^{-x})\mathbf{1}_{\mathbb{R}}(x)$ is the Gumbel distribution. Due to [4, Theorem 6.11], one can see that $F^{\boxtimes n}(a_n \cdot +b_n) \xrightarrow{w} \Phi_0^{\text{free}}$.

Lemma 4.2. Under Assumption 4.1, we have $F^{\boxtimes n}(a_n \cdot +b_n) \xrightarrow{w} \Phi_0^{\text{free}}$.

Here, we focus on the following class of sample distributions.

Definition 4.3. Denote by \mathcal{F}_0 the set of all distribution functions F such that

(G-1) $F = \exp\{-e^{-\phi}\}\$, where ϕ is a twice differentiable function with $\phi' > 0$ in a neighborhood of ω_F

(G-2) there exists a nonincreasing continuous function g such that $g(x) \to 0$ as $x \to \omega_F$ and $|h(x)| \le g(x)$ for all $x < \omega_F$.

By Lemma 4.2, it is easy to see that if $F \in \mathcal{F}_0$, then $F^{\boxtimes n}(a_n \cdot +b_n) \xrightarrow{w} \Phi_0^{\text{free}}$. In this case, we investigate the convergence of densities towards the free Gumbel distribution as follows.

For a positive real number a, we define the following distribution functions:

$$U_{+}(a,x) := \{1 - (1+ax)^{-a^{-1}}\} \mathbf{1}_{(-a^{-1},\infty)}(x),$$

$$U_{-}(a,x) := \{1 - (1-ax)^{a^{-1}}\} \mathbf{1}_{(-\infty,a^{-1})}(x) + \mathbf{1}_{[a^{-1},\infty)}(x).$$

Lemma 4.4. For 0 < a < 1, we get $\sup_{x \in \mathbb{R}} |U_{\pm}(a,x) - \Phi_0^{\text{free}}(x)| \le e^{-1}a$.

Proof. We provide only the bound on $U_{-}(a,x) - \Phi_{0}^{\text{free}}(x)$. Note that

$$\sup_{x \in \mathbb{R}} |U_{-}(a, x) - \Phi_{0}^{\text{free}}(x)| = \sup_{x \le 0} |U_{-} - \Phi_{0}^{\text{free}}| \vee \sup_{0 < x < a^{-1}} |U_{-} - \Phi_{0}^{\text{free}}| \vee \sup_{a^{-1} \le x} |U_{-} - \Phi_{0}^{\text{free}}|$$
$$=: I_{1} \vee I_{2} \vee I_{3}.$$

For $x \ge a^{-1}$, one can see $|U_{-}(a,x) - \Phi_{0}^{\text{free}}(x)| = e^{-x} \le e^{-a^{-1}} \le e^{-1}a$, and therefore $I_{3} \le e^{-1}a$. Next, for $x \le 0$, it is easy to see $|U_{-}(a,x) - \Phi_{0}^{\text{free}}(x)| = 1 - (1-ax)^{a^{-1}} \le e^{-1}a$, and hence $I_{1} \le e^{-1}a$. Finally, for $0 < x < a^{-1}$, we have

$$|U_{-}(a,x) - \Phi_0^{\text{free}}(x)| = e^{-x} - (1-ax)^{a^{-1}} =: u(x) > 0.$$

It is easy to verify that there is a unique point $0 < x_0 < a^{-1}$ such that $u'(x_0) = 0$ and $u(x) \le u(x_0)$ by the intermediate theorem. Therefore u(x) takes the supremum of on $(0, a^{-1})$ if and only if $u'(x_0) = 0$, that is, $(1 - ax_0)^{a^{-1} - 1} = e^{-x_0}$. Hence

$$I_2 \le \sup_{0 < x < a^{-1}} (e^{-x} - e^{-x}(1 - ax)) = a \sup_{0 < x < a^{-1}} xe^{-x} \le aa^{-1}e^{-a^{-1}} \le e^{-1}a.$$

Thus, the desired result is obtained.

Theorem 4.5. Let $F \in \mathcal{F}_0$, a_n , b_n , g, f be defined as above. Then w_n converges uniformly to the free Gumbel density φ_0^{free} on $(0, \infty)$. More strictly, we obtain

$$\sup_{x>0} |w_n(x) - \varphi_0^{\text{free}}(x)| \le O(n^{-1} \vee g(b_n)),$$

for sufficiently large n.

Proof. For all x > 0 and for sufficiently large $n \in \mathbb{N}$ (such that infinitely many $b_n > 0$), we have

$$|w_{n}(x) - \varphi_{0}^{\text{free}}(x)| \leq |na_{n}\phi'(a_{n}x + b_{n})(-\log F(a_{n}x + b_{n}))F(a_{n}x + b_{n}) - e^{-x}F(a_{n}x + b_{n})| + |e^{-x}F(a_{n}x + b_{n}) - e^{-x}| \leq |na_{n}\phi'(a_{n}x + b_{n})(-\log F(a_{n}x + b_{n})) - e^{-x}| + (1 - F(b_{n})) = \underbrace{\left|\frac{f(b_{n})}{f(a_{n}x + b_{n})}(-n\log F(a_{n}x + b_{n})) - e^{-x}\right|}_{=:I} + (1 - e^{-\frac{1}{n}}).$$

By [19, (2.54)], for x > 0, we have

$$U_{+}(g(b_n), x) \le 1 + n \log F(a_n x + b_n) \le U_{-}(g(b_n), x).$$

According to [12, Lemma 2], for any $\epsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that for x > 0 and $n \ge n_0$, we get

$$\frac{1}{1+\epsilon} \left(\frac{-\log F(b_n)}{-\log F(a_n x + b_n)} \right)^{-\epsilon} \le \frac{f(b_n)}{f(a_n x + b_n)} \le \frac{1}{1-\epsilon} \left(\frac{-\log F(b_n)}{-\log F(a_n x + b_n)} \right)^{\epsilon}.$$

Hence, for any x > 0, $n \ge n_0$,

$$\frac{1}{1+\epsilon}(1-U_{-}(g(b_n),x))^{1+\epsilon} \le \frac{f(b_n)}{f(a_nx+b_n)}(-n\log F(a_nx+b_n)) \le \frac{1}{1-\epsilon}(1-U_{+}(g(b_n),x))^{1-\epsilon}.$$

Finally, we get

$$\Phi_0^{\text{free}}(x) - U_-(g(b_n), x) \le \frac{f(b_n)}{f(a_n x + b_n)} (-n \log F(a_n x + b_n)) - e^{-x} \le \Phi_0^{\text{free}}(x) - U_+(g(b_n), x),$$

for sufficiently large n. By Lemma 4.4, we get $I \leq e^{-1}g(b_n)$, as desired.

Example 4.6. Since all F are differentiable as below, we take b_n such that $F(b_n) = e^{-\frac{1}{n}}$ and $a_n = f(b_n)$. Let w_n be the density of the function $x \mapsto F^{\boxtimes n}(a_n x + b_n)$ as below.

(1) (Classical Gumbel distribution) Suppose that $F = \Phi_0$. By definition of the functions ϕ , f and h_0 , we get $\phi(x) = x$, f(x) = 1 and $h_0(x) = f'(x) = 0$. The choices of a_n and b_n imply that $a_n = 1$ and $b_n = \log n$ for $n \in \mathbb{N}$, respectively. By Theorem 4.5,

$$\sup_{x>0} |w_n(x) - \varphi_0^{\text{free}}(x)| \le O(n^{-1}).$$

(2) Define $F(x) = \exp\{-e^{-x^{\alpha}}\}$ for $\alpha > 0$ with $\alpha \neq 1$. Then we get $\phi(x) = x^{\alpha}$, $f(x) = \alpha^{-1}x^{-\alpha+1}$ and $h_0(x) = (-1 + \alpha^{-1})x^{-\alpha}$. The choices of a_n and b_n lead to $a_n = \alpha^{-1}(\log n)^{-1 + \frac{1}{\alpha}}$ and $b_n = (\log n)^{\frac{1}{\alpha}}$ for $n \in \mathbb{N}$, respectively. Define $g(x) = |h_0(x)| = |-1 + \alpha^{-1}|x^{-\alpha}$. Then we get

$$\sup_{x>0} |w_n(x) - \varphi_0^{\text{free}}(x)| \le O\left(\frac{1}{n} \vee \left| -1 + \frac{1}{\alpha} \right| \frac{1}{\log n}\right) = O\left(\frac{1}{\log n}\right).$$

(3) (Normal distribution) Let F be the standard normal distribution function and let us set $p(t) := \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2}{2}}$ as the standard normal density. We define b_n such that $F(b_n) = e^{-\frac{1}{n}}$ and $a_n = \frac{F(b_n)}{nF'(b_n)}$. It is easy to see that F'(x) = p(x) and F''(x) = -xp(x), and therefore $h_0(x) = -\log F(x)(1 + \frac{xF(x)}{p(x)}) - 1$. Here we take $g(x) = 1 + \log F(x)(1 + \frac{x}{p(x)})$. Hence

$$g(b_n) = 1 - \frac{1}{n} \left(1 + \frac{b_n}{p(b_n)} \right) = 1 - \frac{1}{n} - \frac{\sqrt{2\pi}b_n}{n} \exp\left(\frac{b_n^2}{2}\right).$$

Since it must be $\frac{\sqrt{2\pi}b_n}{n} \exp\left(\frac{b_n^2}{2}\right) \to 1$ as $n \to \infty$, we get

$$b_n = \sqrt{2\log n} - \frac{\log\log n + \log 4\pi}{2\sqrt{2\log n}},$$

see e.g. [13]. Finally, we have $g(b_n) = O((\log n)^{-\frac{1}{2}})$, and therefore

$$\sup_{x>0} |w_n(x) - \varphi_0^{\text{free}}(x)| \le O\left(\frac{1}{n} \vee O\left(\frac{1}{\sqrt{\log n}}\right)\right) = O\left(\frac{1}{\sqrt{\log n}}\right),$$

by Theorem 4.5.

ACKNOWLEDGEMENTS

This work was started when Y.K. was affiliated with Department of Mathematics, Hokkaido University. Furthermore, Y.U. was supported by JSPS Grant-in-Aid for Young Scientists 22K13925.

References

- [1] T. Ando, Majorization, doubly stochastic matrices and comparison of eigenvalues, Linear Algebra Appl. 18 (1989) 163–248.
- [2] C. Bartholmé and Y. Swan, Rates of convergence towards the Fréchet distribution, preprint, arXiv:1311.3896.
- [3] G. Ben Arous and V. Kargin, Free point processes and free extreme values, Probab. Theory Relat. Fields 147 (2010) 161–183.
- [4] G. Ben Arous and D. Voiculescu, Free extreme values, Ann. Probab. 34 (2006) 2037–2059.
- [5] F. Benaych-Georges and T. Cabanal-Duvillard, A matrix interpolation between classical and free max operations.
 I. The univariate case, J. Theoret. Probab. 23 (2010) 447–465.
- [6] H. Bercovici and V. Pata, Stable laws and domains of attraction in free probability theory (with an appendix by P. Biane), Ann. Math. (2) 149 (1999) 1023–1060.
- [7] R. A. Fisher and L.H.C. Tippett, Limiting forms of the frequency distribution of the largest or smallest member of a sample. Proc. Camb. Philos. Soc. 24 180. (1928).
- [8] B. V. Gnedenko, Sur la distribution limite du terme maximum d'une série aléatoire, Ann. Math. 44 (1943) 423–453 (Translated and reprinted in: Breakthroughs in Statistics, Vol. I, eds. S. Kotz and N. L. Johnson (Springer-Verlag, 1992), pp. 195–225)
- [9] J. Grela and M. A. Nowak, Extreme matrices or how an exponential map links classical and free extreme laws, Phys. Rev. E 102 (2020) 022109.
- [10] U. Haagerup and S. Möller, The law of large numbers for the free multiplicative convolution, in Operator Algebra and Dynamics, Springer Proceedings in Mathematics and Statistics, Vol. 58 (2013), pp. 157–186.
- [11] L. de Haan, On regular variation and its applications to the weak convergence of sample extremes, Math. Centre Tracts 32 (1970).
- [12] L. de Haan and S. I. Resnick, Local limit theorems for sample extremes, Ann. Probab., 10, no.2, 396–413 (1982).
- [13] P. Hall, On the Rate of Convergence of Normal Extremes, J. Appl. Probab. 16, no.2, 433–439 (1979).
- [14] T. Hasebe and Y. Ueda, Homomorphisms relative to additive convolutions and max-convolutions: Free, boolean and classical cases, Proc. Amer. Math. Soc. 149 (2021) 4799–4814.
- [15] H. Kusumoto and A. Takeuchi, Remark on rates of convergence to extreme value distributions via the Stein equations, Extremes 23 (2020) 411–423.
- [16] A. Nica and R. Speicher, Lectures on the Combinatorics of Free Probability. London Mathematical Society Lecture Note Series 335. Cambridge: Cambridge Univ. Press. (2006).
- [17] M. P. Olson, The selfadjoint operators of a von Neumann algebra form a conditionally complete lattice, Proc. Amer. Math. Soc. 28 (1971) 537–544.
- [18] E. Omey, Rates of convergence for densities in extreme value theory, Ann. Probab. 16 (1988) 479-486.
- [19] S. I. Resnick, Extreme Values, Regular Variation and Point Processes, Springer Series in Operations Research and Financial Engineering (1987).
- [20] R. L. Smith, Uniform rates of convergence in extreme-value theory, Adv. Appl. Probab. 14 (1982) 600–622.
- [21] G. H. Tucci, Limits laws for geometric means of free random variables, Indiana Univ. Math. J. 59 (2010) 1–13.
- [22] Y. Ueda, Rates of convergence for laws of the spectral maximum of free random variables. Infin. Dimens. Anal. Quantum Probab. Relat. Top. 25, No. 3, 2250019 (2022).
- [23] Y. Ueda, Limit theorems for classical, freely and Boolean max-infinitely divisible distributions. J. Theoret. Probab. 35 (2022), no.1, 89–114.
- [24] Y. Ueda, Max-convolution semigroups and extreme values in limit theorems for the free multiplicative convolution, Bernoulli 27 (2021) 502–531.
- [25] D. Voiculescu, Addition of certain noncommuting random variables. J. Funct. Anal. 66 323–346. (1986).

YAMATO KINDAICHI: SAPPORO, HOKKAIDO, JAPAN Email address: e.ifswb2mussw2@gmail.com

Yuki Ueda: Department of Mathematics, Hokkaido University of Education, 9 Hokumon-cho, Asahikawa, Hokkaido 070-8621, Japan

Email address: ueda.yuki@a.hokkyodai.ac.jp