## CONNECTING DISCLINATIONS BY RIDGES

PETER GLADBACH AND HEINER OLBERMANN

ABSTRACT. We consider a thin elastic sheet with a finite number of disclinations in a variational framework in the Föppl–von Kármán approximation. Under the non-physical assumption that the out-of-plane displacement is a convex function, we prove that minimizers display ridges between the disclinations. We prove the associated energy scaling law with upper and lower bounds that match up to logarithmic factors in the thickness of the sheet. One of the key estimates in the proof that we consider of independent interest is a generalization of the monotonicity property of the Monge-Ampère measure.

## 1. INTRODUCTION

The present paper draws its motivation from the mathematical analysis of strongly compressed thin elastic structures, corresponding to the *crumpling* of paper. This phenomenon has received a lot of attention in the physics literature over the last decades, see e.g. [LW05, Wit07, DW01, LGL<sup>+</sup>95, WL93, CM98, CM05, PDK<sup>+</sup>08]. Viewing it as an instance of the minimization of (free) energy, one expects from the everyday experience of crumpling paper that the structure of minimizers should be characterized by a complex network of ridges that meet in vertices, with the elastic energy being focused in this network. Closer inspection shows that elastic energy should be focused in the ridges only, the formation of vertices being energetically less costly. As the thickness of the sheet h is sent to 0 with all other parameters in the setting being fixed, one expects that a finite number of ridges should carry the elastic energy, which scales with  $h^{5/3}$ . This conjecture has been supported by the lower bounds proved by Venkataramani in [Ven04], where a single ridge is considered as the result of clamping an elastic sheet along the lateral boundaries of a bent frame. Venkataramini's lower bound holds in the Föppl-von Kármán approximation, and was later upgraded by Conti and Maggi [CM08] to a lower bound with the same scaling in the geometrically fully nonlinear setting. The latter authors also provide an upper bound with the same scaling  $h^{5/3}$  for a much more general situation, in the form of an approximation result for piecewise affine isometric maps. The boundary conditions considered for a single fold in [CM08, Ven04] are such that any deviation from the sharp fold satisfying these conditions involves stretching of the sheet, and is hence very costly. This fact is at the heart of the respective proofs. In the sequel, we will call boundary conditions with this feature *tensile*.

From a mathematical point of view, the next step in the asymptotic analysis would be to prove the *emergence* of ridges without enforcing them by tensile boundary conditions, and an associated lower bound with the expected scaling. In sufficient generality (e.g., just assuming that the elastic sheet is confined by a container whose diameter is much smaller than that of the sheet, a setting that we will refer to as the "crumpling problem" from now on), this is a very difficult and up to now entirely open problem. One way of understanding

Date: May 17, 2024.

the difficulty is the following: If boundary conditions and other constraints that define the problem allow for a large class of *short* maps (i.e., deformations  $y : \Omega \to \mathbb{R}^3$  that satisfy  $\nabla y^T \nabla y \leq \mathrm{Id}_{2\times 2}$  in the sense of positive definite matrices), then by the famous Nash-Kuiper theorem [Nas54, Kui55] there exist very many competitors that are arbitrarily close to isometric immersions. Since the distance of the induced metric from that of an isometric immersion is the leading order term in the typical elastic energy

$$I_h(y) = \int_{\Omega} |\nabla y^T \nabla y - \mathrm{Id}_{2 \times 2}|^2 + |\nabla^2 y|^2 \mathrm{d}x, \qquad (1)$$

one needs quantitative lower bounds for the bending energy for this large set of competitors.

A setting that at first sight seems well adapted to investigate the emergence of ridges would be an elastic sheet featuring two *disclinations* (metric defects). The case of one disclination has been investigated in [Olb16, Olb17, Olb18], where, aside from an energy scaling law, it has been shown that minimizers converge to a conical configuration as the thickness of the sheet approaches zero. In the case of two disclinations, one expects approximately conical shapes in the neighborhood of each of them, and the formation of a ridge between them.<sup>1</sup> Since one single ridge is expected at a known position, it seems that this setting should be much more tractable than the full crumpling problem as described above. Alas, also in this setting the derivation of a lower bound with the conjectured  $h^{5/3}$ scaling seems to be prohibitively difficult. To the best of the authors' knowledge, this remains an entirely open problem.

In the present article we will solve the problem we have just described under an additional non-physical assumption: Working in the Föppl–von Kármán approximation, we assume that the out-of-plane deformation is *convex*. Under this assumption, we are able to show the emergence of a ridge between two disclinations, and an associated energy scaling law with upper and lower bounds that match up to logarithmic factors in h.

More precisely, we work with a finite number of disclinations, and show that up to logarithmic factors, the energy of the ridges between them scales with  $h^{4/3}$ . As is to be expected, the energy under the convexity assumption is larger than without. In Section 3.2, we show that an energy bounded by  $Ch^{5/3}$  can be achieved in the absence of the constraint of convexity of the out-of-plane component. This means that our results constitute a rigorous proof that for h small enough, minimizers have non-convex out-of-plane component (see Corollary 3.4).

1.1. Statement of the main result. Let  $\Omega \subseteq \mathbb{R}^2$  be convex, open, bounded with  $C^{1,1}$  boundary,  $\{a_1, \ldots, a_N\} \subseteq \Omega$ ,  $\sigma_i > 0$  for  $i = 1, \ldots, N$ , and  $\bar{\mu} := \sum_{i=1}^N \sigma_i \delta_{a_i}$ . For a convex function  $u : \Omega \to \mathbb{R}$ , let  $\partial^- u(x)$  denote the subdifferential of u at x,

$$\partial^{-}u(x) = \{ p \in \mathbb{R}^{n} : p \cdot (y - x) + u(x) \le u(y) \text{ for all } y \in \Omega \}$$

and let  $\mu_u$  denote its Monge-Ampère measure defined for a Borel set  $E \subseteq \Omega$  as

$$\mu_u(E) = \mathcal{L}^2\left(\partial^- u(E)\right) \,,$$

<sup>&</sup>lt;sup>1</sup>This comes with the caveat that the formation of a ridge in such a situation can be avoided, unless further constraints or boundary conditions are introduced into the problem, see Section A in the appendix. In the statement of Theorem 1.1, this additional assumption is given by the Dirichlet boundary values v = 0 on  $\partial\Omega$ .

where  $\mathcal{L}^2$  denotes the Lebesgue measure on  $\mathbb{R}^2$ . By [Fig17, Theorem 2.13], there exists a unique convex solution  $v_0$  of the Monge-Ampère equation

$$\begin{cases} \mu_{v_0} = \bar{\mu} \text{ in } \Omega\\ v_0 = 0 \text{ on } \partial\Omega. \end{cases}$$
(2)

By Lemma 2.11 below,  $v_0 \notin W^{2,2}(\Omega)$ . We define the Föppl–von Kármán energy for deformations  $(\mathbf{u}, v) \in W^{1,2}(\Omega; \mathbb{R}^2) \times W^{2,2}(\Omega)$ , with the reference configuration given by the "singular" configuration  $v_0$ ,

$$E_{h}(\mathbf{u}, v) = \int_{\Omega} \left| \nabla \mathbf{u} + \nabla \mathbf{u}^{T} + \nabla v \otimes \nabla v - \nabla v_{0} \otimes \nabla v_{0} \right|^{2} + h^{2} |\nabla v|^{2} \mathrm{d}x.$$
(3)

This functional can be viewed as a partial linearization of the geometrically nonlinear functional (1), with metric defects. A rigorous relation between these energies can be established in the sense of  $\Gamma$ -convergence, see [FJM06]. In the definition of the energy (3), we have chosen to absorb the in-plane deformation  $\mathbf{u}_0$  that would yield vanishing stress in the reference configuration via  $\nabla \mathbf{u}_0 + \nabla \mathbf{u}_0^T + \nabla v_0 \otimes \nabla v_0 = 0$ , into  $\mathbf{u}$ . Let

$$\mathcal{A} := \left\{ (\mathbf{u}, v) \in W^{1,2}(\Omega; \mathbb{R}^2) \times W^{2,2}(\Omega) : v \text{ convex}, v = 0 \text{ on } \partial\Omega \right\}$$

We will prove:

**Theorem 1.1.** There exist  $h_0, C > 0$  that only depend on  $\Omega$  and  $\overline{\mu}$  such that

$$\frac{1}{C}h^{4/3} \left(\log \frac{1}{h}\right)^{-2/3} \le \inf_{(\mathbf{u},v)\in\mathcal{A}} E_h(\mathbf{u},v) \le Ch^{4/3} \left(\log \frac{1}{h}\right)^{1/3}$$

for all  $h < h_0$ .

- **Remark 1.2.** (i) The sources of the logarithmic factors in the upper and lower bounds are discussed in Remarks 3.2 and 4.1 below, respectively. For a sketch of the idea of the lower bound, see the beginning of Section 4.
  - (ii) The boundary conditions v = 0 on  $\partial\Omega$  are necessary in order to "force" the emergence of a ridge between the vertices defined by  $\bar{\mu}$ . If these boundary conditions are omitted, then it is possible to achieve an energy that scales with  $h^2 \log \frac{1}{h}$ , which is much smaller than  $h^{4/3}$ . For a proof of this fact, see Section A of the appendix.

1.2. The method of proof: Gauss curvature and convexity. As should be clear from the discussion so far, of the two inequalities in Theorem 1.1, the lower bound is the more interesting and difficult one to prove. Its proof is based on a principle that has already been used in [Olb16, Olb17, Olb18, Olb19, GO24]. In these papers lower bounds for the elastic energy are derived starting from a certain control over the (linearized) Gauss curvature of the deformed surface, which is obtained either directly from the constraints in the problem or from energetic considerations. The cited papers do this in settings that produce approximately conical configurations, which are characterized by an elastic energy that scales with  $h^2 \log \frac{1}{h}$ .

The idea is that the leading order term in the elastic energy (distance of the induced metric from that of an isometric immersion) controls the Gauss curvature. Knowledge about the Gauss curvature can be translated into information about the image of the deformation gradient. This information can in turn be exploited to obtain lower bounds for the second gradient of the deformation.

If one assumes convexity of the out-of-plane deformation v and hence non-negativity of the linearized Gauss curvature (in other words: the Monge-Ampère measure  $\mu_v$ ), this chain of arguments can be made more powerful by noting that knowledge about the Monge-Ampère measure directly gives some information about the deformation itself via the monotonicity of the subdifferential of convex functions and the Alexandrov maximum principle. In fact, our proof uses as an essential tool the following generalization of the monotonicity property of the Monge-Ampère measure, which we consider of interest in its own right (see Proposition 2.6 below): If  $\Omega \subseteq \mathbb{R}^n$  is a convex bounded open domain, and  $\phi \in C^0(\overline{\Omega})$  a non-negative concave function, then whenever  $u, v \in C^0(\overline{\Omega})$  are convex with u = v on  $\partial\Omega$  and  $u \geq v$ , we have

$$\int_{\Omega} \phi \mathrm{d}\mu_u \le \int_{\Omega} \phi \mathrm{d}\mu_v \,. \tag{4}$$

The classical monotonicity property is the special case  $\phi = 1$ . The proof will be given in Section 2.1 below.

The combination of the Föppl–von Kármán approximation with the assumption of convexity of the out-of-plane component has been studied in [GO24], where a partially clamped elastic sheet in the shape of the sector of a disk has been considered. The method of proof used in that paper is adapted to the particular setting considered there. The tools we will present here are different and more versatile.

1.3. Scientific context. Crumpling is only one of many possible *patterns* that thin elastic sheets may form under the influence of external forces. Explaining these patterns as the result of energy minimization has proved to be a fruitful principle within the physics and mathematics community, see e.g. the overview article by Witten [Wit07] or the book by Audoly and Pomeau [AP10]. In the calculus of variations, two types of results have emerged: Firstly, the derivation of reduced models of three-dimensional nonlinear elasticity in the limit of small film thickness by means of  $\Gamma$ -convergence. Notable successes include the derivation of a membrane model by Le Dret and Raoult [LDR95] as well as the derivation of nonlinear plate models by Friesecke, James and Müller [FJM02, FJM06]. The  $\Gamma$ -convergence analysis has also been extended to settings involving shells or curved reference configurations, see [LMP11, Lew20, Tob21].

Second, there is the investigation of the qualitative properties of low-energy states in the variational formulation of elasticity. An early instance of this approach in nonlinear elasticity is due to Ball and James [BJ87]. A popular way of carrying out such an analysis is by the derivation of energy scaling laws (and is, of course, the approach followed in the current paper), such as as in the work by Kohn and Müller [KM92, KM94]. Since then there have been many successful applications to phenomena in materials science. In the case of thin elastic sheets, there have been notable results for sheets wrinkling under tension [BK14], under compression [BCDM02], or when attached to a substrate [KN13, BK17].

1.4. Notation. For  $A \subseteq \mathbb{R}^n$ , we will denote the convex envelope of A as convA, and the interior of A by  $A^\circ$ . The characteristic function of A is denoted by  $\mathbb{1}_A$ . For a finite number of points  $a_1, \ldots, a_k \in \mathbb{R}^n$ , we will write  $[a_1a_2 \ldots a_k] = \operatorname{conv}\{a_1, \ldots, a_k\}$ . For  $x = (x_1, x_2) \in \mathbb{R}^2$  we write  $x^{\perp} = (-x_2, x_1)$ .

The k-dimensional Hausdorff measure is denoted by  $\mathcal{H}^k$ , the n-dimensional Lebesgue measure by  $\mathcal{L}^n$ . The set of signed Radon measures on  $\Omega \subset \mathbb{R}^n$  is denoted by  $\mathcal{M}(\Omega)$ . The subset of non-negative elements of  $\mathcal{M}(\Omega)$  is denoted by  $\mathcal{M}_+(\Omega)$ . For every  $\mu \in \mathcal{M}(\Omega)$ there exists a decomposition  $\mu = \mu_+ - \mu_-$  with  $\mu_{\pm} \in \mathcal{M}_+(\Omega)$ . We write  $|\mu| = \mu_+ + \mu_-$ . For a Radon measure  $\mu$  and a  $|\mu|$ -integrable function f, the signed Radon measure  $\mu \sqcup f$ is defined by  $\mu \sqcup f(A) = \int_A f d\mu$  for  $A \subseteq \mathbb{R}^n$   $\mu$ -measurable. For sequences  $\mu_k \in \mathcal{M}(\Omega)$  and  $\mu \in \mathcal{M}(\Omega)$  such that

$$\lim_{k \to \infty} \int_{\Omega} \varphi d\mu_k \to \int_{\Omega} \varphi d\mu \quad \text{for all } \varphi \in C_c^0(\Omega) , \qquad (5)$$

we say that  $\mu_k$  converges weakly-\* to  $\mu$  and write  $\mu_k \stackrel{*}{\rightharpoonup} \mu$ . We note that (5) is equivalent to

$$\mu(U) \leq \liminf_{k \to \infty} \mu_k(U) \quad \text{for all open sets } U \subseteq \Omega$$
  
and  $\mu(K) \geq \limsup_{k \to \infty} \mu_k(K) \quad \text{for all compact sets } K \subseteq \Omega.$ 

For matrices  $A, B \in \mathbb{R}^{n \times n}$ , we define the dot product  $A : B = \text{Tr}(A^T B)$ , where Tr denotes the trace. The matrix  $\operatorname{cof} A \in \mathbb{R}^{n \times n}$  denotes the matrix of cofactors of A,  $(\operatorname{cof} A)_{ij} = (-1)^{i+j} \det \hat{A}_{ij}$ , where  $\hat{A}_{ij}$  is the  $(n-1) \times (n-1)$  matrix obtained from A by deleting the *i*-th row and the *j*-th column.

The symbol "C" will be used as follows: An inequality such as  $f \leq Cg$  is always to be understood as the statement "there exists a constant C > 0 such that  $f \leq Cg$ "; sometimes we will put the full statement for clarity, in other places we just write the inequality in an attempt to increase readability. When we write C(a, b, ...), then C may depend on a, b, ... Dependence of C may be omitted from the notation when it is clear from the context. Instead of  $f \leq Cg$ , we also write  $f \leq g$ . In chains of inequalities, the value of C may change at each time that the symbol is used.

1.5. Plan of the paper. In Section 2, we derive some results in convex analysis; in particular we prove the generalized monotonicity property of the Monge-Ampère measure and we will study the structure of the solution  $v_0$  of (2). Using this structure we may then construct the upper bound in Theorem 1.1 in Section 3.1. In Section 3.2 we provide the upper bound construction for the situation when the convexity constraint for the out-of-plane component is discarded, achieving a lower energy. Section 4 contains the proof of the lower bound. It starts off with a rough overview of the proof, followed up by the details in Subsections 4.1-4.5. In the appendix, we show that when no boundary conditions are imposed on v, an energy that scales with  $h^2 \log \frac{1}{h}$  can be achieved (much lower than what we found in Theorem 1.1), with test functions that do not display any ridges.

## 2. Some convex analysis

Let  $\Omega \subseteq \mathbb{R}^n$  be open, convex and bounded. For a convex function  $u : \overline{\Omega} \to \mathbb{R}$ , the notions of subgradient  $\partial^- u$  and Monge-Ampère measure that we have introduced above for the special case n = 2 have their obvious analogue in the general case. We recall that  $\mu_u$  is a Radon measure, and the following elementary fact concerning subgradients, that we will repeatedly use in our proofs below without further comment:

$$p \in \partial^- u(x) \quad \Leftrightarrow \quad p \cdot x - u(x) \ge p \cdot y - u(y) \quad \text{ for all } y \in \overline{\Omega} \,.$$

As is well known, any convex function  $u: \overline{\Omega} \to \mathbb{R}$  can be written as the supremum of affine functions that are smaller than u,

 $u(x) = \sup\{a(x) : a \text{ affine }, a \le u\}.$ 

For a possibly non-convex function  $u: \overline{\Omega} \to \mathbb{R}$  we define the convex envelope as

 $u^{c}(x) = \sup\{v(x) : v \text{ convex }, v \le u\}.$ 

Combining with the previous identity, we have

$$u^c(x) = \sup\{v(x): a \text{ affine }, \, a \leq u\}$$

For  $E \subseteq A$  and  $u: A \to \mathbb{R}$ , we define the restriction  $R_E u: A \to \mathbb{R} \cup \{+\infty\}$ ,

$$R_E u(x) = \begin{cases} u(x) & \text{if } x \in E \\ +\infty & \text{else.} \end{cases}$$

We define the lifting of a convex function on a bounded set:

**Definition 2.1.** Let  $K \subseteq \mathbb{R}^n$  be compact,  $\Omega := (\operatorname{conv} K)^\circ$ , and  $u \in C^0(\overline{\Omega})$ . We define the lifting of u on K as  $L_K u \in C^0(\overline{\Omega})$ ,

$$L_K u := (R_K u)^c$$

Here it is understood that the supremum in the definition of  $(R_K u)^c$  is taken over convex functions  $v: \overline{\Omega} \to \mathbb{R}$  that satisfy  $v \leq R_K u$ .

We note that  $L_K u$  is a well-defined convex function  $\overline{\Omega} \to \mathbb{R}$ . If u itself is convex, then  $L_K u \ge u$  in convK, and  $L_K u = u$  on K. As a consequence of the above identities,

$$L_K u(x) = \sup\{a(x) : a \text{ affine }, a \le u \text{ on } K\}.$$
(6)

The lifting  $L_{K}u$  has the property that its Monge-Ampère measure is concentrated on the set K, as we will show in the upcoming lemma.

**Lemma 2.2.** Let  $K \subseteq \mathbb{R}^n$  be compact,  $\Omega = (\operatorname{conv} K)^\circ$ ,  $u \in C^0(\overline{\Omega})$  convex,  $w = L_K u$ , and  $p \in \partial^- w(\Omega)$ . Then all extreme points of  $(\partial^- w)^{-1}(p)$  are contained in K. In addition, we have

$$\mu_w(\Omega \setminus K) = 0.$$

*Proof.* Assume there are  $x \in \Omega$ ,  $p \in \partial^- w(x)$  and an extreme point y of  $(\partial^- w)^{-1}(p)$  with  $y \notin K$ . Then  $a(z) = u(x) + p \cdot (z - x)$  defines the affine function satisfying  $(\partial^- w)^{-1}(p) = \{z : a(z) = w(z)\}$ . Since y is an extreme point of the convex set  $(\partial^- w)^{-1}(p)$  and y has positive distance from the compact set K, there exists an affine function  $b : \mathbb{R}^n \to \mathbb{R}$  and  $\delta > 0$  such that b(y) > 0 and

$$b(x') \le 0$$
 if  $x' \in K$  and  $\operatorname{dist}(x', (\partial^- w)^{-1}(p)) < \delta$ .

We observe that by choice of the affine function a,

 $m := \max\{a(x') - w(x') : \operatorname{dist}(x', (\partial^{-}w)^{-1}(p)) \ge \delta\} < 0,$ 

and hence by choosing

$$0 < \varepsilon < \frac{|m|}{2 \max_{\operatorname{conv} K} |b|}$$

and recalling w = u on K, we obtain that

$$\begin{array}{c} a + \varepsilon b \leq u \text{ on } K \, . \\ 6 \end{array}$$

Using (6), it follows  $w(y) \ge a(y) + \varepsilon b(y) > a(y) = w(y)$ , a contradiction. This shows  $y \in K$ .

To see that  $\mu_w(\Omega \setminus K) = 0$ , we define the set

$$N = \{ p \in \mathbb{R}^n : \#(\partial^- w)^{-1}(p) > 1 \}.$$

Denoting by  $w^*$  the Legendre transform of w, we have that  $(\partial^- w)^{-1} = \partial^- w^*$  (see [Roc70, Theorem 23.5]), and hence  $N = \{p \in \mathbb{R}^n : \#\partial^- w^*(p) > 1\}$ . Since  $w^*$  is locally Lipschitz, and  $\#\partial^- w^*(p) > 1$  implies non-differentiability of  $w^*$  in p, we have that  $\mathcal{L}^n(N) = 0$  by Rademacher's Theorem. By what we have already shown, if  $x \in (\partial^- w)^{-1}(p) \setminus K$ , then x is not an extreme point of  $(\partial^- w)^{-1}(p)$ , which of course implies  $\#(\partial^- w)^{-1}(p) > 1$ , i.e.,  $p \in N$ . Thus we have  $\partial^- w(\Omega \setminus K) \subseteq N$  which in turn yields  $\mu_w(\Omega \setminus K) = \mathcal{L}^n(\partial^- w(\Omega \setminus K)) \leq \mathcal{L}^n(N) = 0$ .

The following approximation lemma will be repeatedly useful in our proofs below.

**Lemma 2.3.** Let  $u \in W^{1,\infty}(\mathbb{R}^n)$  be a convex function,  $\varepsilon > 0$ , and

$$P_{\varepsilon}u(x) := \inf \left\{ q(x) : q \in C^{2}(\Omega), \nabla^{2}q \ge \varepsilon^{-1} \mathrm{Id}, q \ge u \right\} \quad \text{for } x \in \mathbb{R}^{n} \,.$$
(7)

Then  $P_{\varepsilon}u$  is convex, with  $\|\nabla^2 P_{\varepsilon}u\|_{L^{\infty}(\mathbb{R}^n)} \leq \frac{1}{\varepsilon}$ , and writing  $c = 4\|\nabla u\|_{L^{\infty}(\mathbb{R}^n)}$  we have that  $u = P_{\varepsilon}u$  on the set

$$\left\{x \in \mathbb{R}^n : u \in W^{2,\infty}(B(x,c\varepsilon)), \|\nabla^2 u\|_{L^{\infty}(B(x,c\varepsilon))} \leq \frac{1}{\varepsilon}\right\}.$$

*Proof.* First we note that the right hand side in (7) can be slightly changed, replacing the inequality  $\nabla^2 q \leq \varepsilon^{-1}$ Id by an equality, and that the infimum is achieved:

$$P_{\varepsilon}u(x) = \min\left\{q(x) : q \in C^2(\mathbb{R}^n), \nabla^2 q = \varepsilon^{-1} \mathrm{Id}, q \ge u\right\}.$$
(8)

Here the existence of the minimum for every x follows from a straightforward compactness argument.

Now we show that  $P_{\varepsilon}u$  is convex. Let  $x_0, x_1 \in \mathbb{R}^n$  and  $x = (1 - \lambda)x_0 + \lambda x_1$  for some  $\lambda \in [0, 1]$ . For i = 0, 1 let  $Q_i$  be quadratic functions with  $\nabla^2 Q_i = \frac{\text{Id}}{\varepsilon}, Q_i \geq u$  on  $\Omega$ , and  $Q_i(x_i) = u(x_i)$ .

Define

$$\tilde{Q} := (\min(Q_0, Q_1))^c \,.$$

Since u is convex, we have  $\tilde{Q} \geq u$ . We note that  $\tilde{Q} \in W^{2,\infty}(\mathbb{R}^n)$  with  $\nabla^2 \tilde{Q} \leq \frac{\mathrm{Id}}{\varepsilon}$  almost everywhere. This may either be deduced by computing  $\tilde{Q}$  explicitly, or from the general regularity results of convex envelopes from [KK01, GR90].

Define the quadratic function  $Q(y) := \tilde{Q}(x) + \nabla \tilde{Q}(x) \cdot (y-x) + \frac{|y-x|^2}{2\varepsilon}$ . Then  $Q \ge \tilde{Q} \ge u$  on  $\Omega$  and

$$P_{\varepsilon}u(x) \le Q(x) = \tilde{Q}(x) \le (1-\lambda)Q_0(x_0) + \lambda Q_1(x_1) \le (1-\lambda)P_{\varepsilon}u(x_0) + \lambda P_{\varepsilon}u(x_1).$$

This proves the convexity of  $P_{\varepsilon}u$ .

To see that  $\nabla^2 P_{\varepsilon} u \leq \frac{\mathrm{Id}}{\varepsilon}$ , we use that for every  $x \in \mathbb{R}^n$ , there is a quadratic function Q with  $\nabla^2 Q = \frac{\mathrm{Id}}{\varepsilon}$ ,  $Q \geq P_{\varepsilon} u$  and  $Q(x) = P_{\varepsilon} u(x)$  as we have noted in (8). Since  $P_{\varepsilon} u$  is also convex, this implies that  $P_{\varepsilon} u$  is everywhere differentiable and that  $\nabla P_{\varepsilon} u$  is Lipschitz, with  $0 \leq \nabla^2 P_{\varepsilon} u \leq \frac{\mathrm{Id}}{\varepsilon}$  almost everywhere.

Finally, assume that  $x \in \mathbb{R}^n$  with  $\|\nabla^2 u\|_{L^{\infty}(B(x,4\|\nabla u\|_{\infty}\varepsilon))} \leq \frac{1}{\varepsilon}$ . Define the function  $Q(y) = u(x) + \nabla u(x) \cdot (y-x) + \frac{|y-x|^2}{2\varepsilon}$ . By our choice of x, we have  $u \leq Q$  in  $\Omega \cap B(x,4\|\nabla u\|_{\infty}\varepsilon)$ . For  $y \in \Omega \setminus B(x,4\|\nabla u\|_{\infty}\varepsilon)$ , we have by Lipschitz continuity  $u(y) \leq u(x) + \|\nabla u\|_{\infty}|y-x|$ , whereas

$$Q(y) \ge u(x) - \|\nabla u\|_{\infty} |y - x| + \frac{|y - x|^2}{2\varepsilon}$$
  
$$\ge u(x) - \|\nabla u\|_{\infty} |y - x| + \frac{|y - x|^2}{2\varepsilon}$$
  
$$\ge u(y).$$

Hence  $Q \ge u$  on  $\Omega$  and u(x) = Q(x), which implies  $P_{\varepsilon}u(x) = u(x)$ , completing the proof.

2.1. Monotonicity properties for convex functions. We now state several invariances of Monge-Ampère measures:

**Lemma 2.4.** Let  $\Omega \subseteq \mathbb{R}^n$  be convex, open, and bounded. Let  $u, v \in C^0(\mathbb{R}^n)$  be convex functions with u = v in  $\mathbb{R}^n \setminus \Omega$ . Then

(i)  $\partial^{-}u(\bar{\Omega}) = \partial^{-}v(\bar{\Omega})$ . In particular,  $\mu_{u}(\bar{\Omega}) = \mu_{v}(\bar{\Omega})$ . (ii)  $\int_{\bar{\Omega}} x d\mu_{u} = \int_{\bar{\Omega}} x d\mu_{v}$ .

*Proof.* To prove (i), let  $p \in \partial^- v(\overline{\Omega})$ . Then the concave function  $x \mapsto p \cdot x - v(x)$  achieves its maximum on  $\overline{\Omega}$ , and hence the maximum of that function on  $\mathbb{R}^n \setminus \Omega$  is achieved on  $\partial\Omega$ ,

$$\max_{x \in \mathbb{R}^n \setminus \Omega} p \cdot x - v(x) = \max_{x \in \partial \Omega} p \cdot x - v(x).$$

Since u = v on  $\mathbb{R}^n \setminus \Omega$ , this identity reads

$$\max_{x \in \mathbb{R}^n \setminus \Omega} p \cdot x - u(x) = \max_{x \in \partial \Omega} p \cdot x - u(x) \,,$$

which implies  $p \in \partial^- u(\bar{\Omega})$ . Reversing the roles of u, v yields the claim.

To prove (ii), assume first that  $u, v \in W^{2,\infty}(\mathbb{R}^n)$ . Let  $i \in \{1, \ldots, n\}$ . Writing w = v - u, we have

$$\frac{\mathrm{d}}{\mathrm{d}t}\Big|_{t=0} \int_{\Omega} x_i \mathrm{d}\mu_{u+t(v-u)} = \left.\frac{\mathrm{d}}{\mathrm{d}t}\right|_{t=0} \int_{\Omega} x_i \det \nabla^2(u+tw) \mathrm{d}x$$
$$= \int_{\Omega} x_i \mathrm{cof} \,\nabla^2 u(x) : \nabla^2 w(x) \,\mathrm{d}x \,.$$

We will show that the right hand side vanishes, which is sufficient to prove our claim. By the distributional Piola identity div cof  $\nabla^2 u = 0$ , we get that

div 
$$(x_i \operatorname{cof} \nabla^2 u \nabla w)$$
, div  $(w \operatorname{cof} \nabla^2 u e_i) \in L^2(\Omega)$ ,

and hence

$$x_i \operatorname{cof} \nabla^2 u \nabla w, \ w \operatorname{cof} \nabla^2 u e_i \in H(\operatorname{div}; \Omega) := \{ f \in L^2(\Omega; \mathbb{R}^n) : \operatorname{div} f \in L^2(\Omega) \}$$

It is well known that  $C^{\infty}(\overline{\Omega})$  is dense in  $H(\operatorname{div};\Omega)$  and that a continuous normal trace operator  $H(\operatorname{div};\Omega) \to H^{1/2}(\partial\Omega)$  exists (see e.g. [Tar07, Chapter 20]). In particular, one may integrate by parts as follows,

r

$$\int_{\Omega} x_i \operatorname{cof} \nabla^2 u(x) : \nabla^2 w(x) \, \mathrm{d}x$$
  
=  $-\int_{\Omega} \operatorname{cof} \nabla^2 u(x) : \nabla w(x) \otimes e_i \, \mathrm{d}x + \int_{\partial\Omega} x_i \operatorname{cof} \nabla^2 u(x) : \underbrace{\nabla w(x)}_{=0} \otimes n(x) b \, \mathrm{d}\mathcal{H}^{n-1}$   
=  $-\int_{\partial\Omega} \underbrace{w(x)}_{=0} \operatorname{cof} \nabla^2 u(x) : e_i \otimes n(x) \, \mathrm{d}\mathcal{H}^{n-1}$   
= 0.

Here we have used the fact that  $w = \nabla w = 0$  on  $\partial \Omega$ . This completes the proof for the case  $u, v \in W^{2,\infty}_{\text{loc.}}(\mathbb{R}^n)$ . We note in passing that for this case, the statement and proof hold true also for non-convex u, v.

If u, v are only continuous, we approximate both u and v using Lemma 2.3, yielding  $u_{\varepsilon}, v_{\varepsilon} \in W^{2,\infty}_{\text{loc.}}(\mathbb{R}^n)$  with  $u_{\varepsilon} = v_{\varepsilon}$  outside of  $B(\Omega, \rho(\varepsilon))$ , with  $\rho(\varepsilon) \to 0$  as  $\varepsilon \to 0$ , and  $u_{\varepsilon} \to u, v_{\varepsilon} \to v$  locally uniformly. From this convergence follows the weak-\* convergence of the Monge-Ampère measures  $\mu_{u_{\varepsilon}}, \mu_{v_{\varepsilon}}$  to  $\mu_u, \mu_v$  respectively (see e.g. [Fig17, Proposition 2.6]). Let us fix  $i \in \{1, \ldots, n\}$ , and choose a > 0 such that  $\phi(x) := a + x_i \ge 0$  on  $\overline{\Omega}$ . Multiplying the convergent sequences by  $\phi$  we get

$$\mu_{v_{\varepsilon}} \sqcup \phi \stackrel{*}{\rightharpoonup} \mu_{v} \sqcup \phi, \quad \mu_{u_{\varepsilon}} \sqcup \phi \stackrel{*}{\rightharpoonup} \mu_{u} \sqcup \phi.$$

We choose a non-increasing sequence of open convex bounded sets  $\Omega_{\delta}$  with  $\Omega \subset \Omega_{\delta}$  and  $\bigcap_{\delta>0}\Omega_{\delta} = \overline{\Omega}$ . Now using the approximation properties of Radon measures (see e.g. [EG92, Chapter 1]), we obtain the following chain of inequalities,

$$\mu_{u} \sqcup \phi(\overline{\Omega}) = \inf_{\delta > 0} \mu_{u} \sqcup \phi(\Omega_{\delta})$$

$$\leq \inf_{\delta > 0} \liminf_{\varepsilon \to 0} \mu_{u_{\varepsilon}} \sqcup \phi(\Omega_{\delta})$$

$$= \inf_{\delta > 0} \liminf_{\varepsilon \to 0} \mu_{v_{\varepsilon}} \sqcup \phi(\Omega_{\delta})$$

$$\leq \inf_{\delta > 0} \limsup_{\varepsilon \to 0} \mu_{v_{\varepsilon}} \sqcup \phi(\overline{\Omega_{\delta}})$$

$$\leq \inf_{\delta > 0} \mu_{v} \sqcup \phi(\overline{\Omega_{\delta}})$$

$$= \mu_{v} \sqcup \phi(\overline{\Omega}) .$$

Reversing the roles of u and v yields the equality  $\mu_u \sqcup \phi(\overline{\Omega}) = \mu_v \sqcup \phi(\overline{\Omega})$ , and hence, after subtracting  $a\mu_u(\overline{\Omega}) = a\mu_v(\overline{\Omega})$  from this equality, we obtain our claim.  $\Box$ 

Remark 2.5. The equality

$$\frac{\mathrm{d}}{\mathrm{d}t}\bigg|_{t=0}\int_{\Omega}x_{i}\det\nabla^{2}(u+tw)\mathrm{d}x=0\quad\text{ for all }w\in W^{2,\infty}_{0}(\Omega)\,,$$

which we have proved above, means of course that  $x_i \det \nabla^2 u$  is a null Lagrangian. To the best of the authors' knowledge, this fact has gone unnoticed in the literature so far.

As a consequence of the previous lemma, we obtain the following local monotonicity properties:

**Proposition 2.6.** Let  $\Omega \subseteq \mathbb{R}^n$  be convex, open, and bounded,  $u, v \in W^{1,\infty}(\Omega)$  convex with u = v on  $\partial\Omega$ ,  $u \ge v$  in  $\Omega$ , and  $\phi \in C^0(\overline{\Omega})$  concave and non-negative. Then

$$\int_{\Omega} \phi \mathrm{d}\mu_u \le \int_{\Omega} \phi \mathrm{d}\mu_v \,. \tag{9}$$

*Proof. Step 1.* First we suppose that for an arbitrary set of points  $\{a_1, \ldots, a_M\} \subseteq \Omega, u, v$  are of the form  $u = L_K u, v = L_K v$ , where  $K = \partial \Omega \cup \{a_1, \ldots, a_M\}$ , and  $u(a_j) = v(a_j)$  for  $j = 2, \ldots, M$ . We extend u and v to convex functions  $U, V \in W^{1,\infty}(\mathbb{R}^n)$  respectively by

$$U(x) = \min_{y \in \bar{\Omega}} u(y) + L|x - y|, \quad V(x) = \min_{y \in \bar{\Omega}} v(y) + L|x - y|,$$

where  $L \ge \max(\|\nabla u\|_{\infty}, \|\nabla v\|_{\infty})$  is an upper bound for the Lipschitz constants of u, v.

Recalling that  $p \in \partial^- u(a_1)$  if and only if the function  $x \mapsto p \cdot x - u(x)$  attains its maximum over K in  $a_1$ , and  $v(a_1) \leq u(a_1)$ , v = u on  $K \setminus \{a_1\}$ , we have that

$$\partial^{-}U(a_1) = \partial^{-}u(a_1) \subseteq \partial^{-}v(a_1) = \partial^{-}V(a_1).$$

Furthermore, since we have  $\partial^- U(\bar{\Omega}) = \partial^- V(\bar{\Omega})$  by Lemma 2.4 (i),

$$\partial^{-}U(y) = \partial^{-}V(y) \cup (\partial^{-}U(y) \cap \partial^{-}v(a_{1})) \text{ for all } y \in K \setminus \{a_{1}\}.$$

In other words, only the excess subgradient  $\partial^- v(a_1) \setminus \partial^- u(a_1)$  is redistributed to  $K \setminus \{a_1\}$ , and

$$0 \le \mu_V(\{a_1\}) - \mu_U(\{a_1\})$$
  
$$0 \ge \mu_V - \mu_U \quad \text{on } K \setminus \{a_1\}.$$

In particular, the latter implies  $\mu_V \leq \mu_U$  on  $\partial\Omega$ . Now let  $b: \Omega \to \mathbb{R}$  be an affine function with  $b(a_1) = \phi(a_1)$  and  $b \geq \phi$ . Then

$$\begin{split} \int_{\Omega} \phi \mathrm{d}(\mu_{v} - \mu_{u}) &= \int_{\Omega} \phi \mathrm{d}(\mu_{V} - \mu_{U}) \\ &\geq \int_{\overline{\Omega}} \phi \mathrm{d}(\mu_{V} - \mu_{U}) \\ &= (\mu_{V} - \mu_{U}) \sqcup \phi(\{a_{1}\}) + (\mu_{V} - \mu_{U}) \sqcup \phi(K \setminus \{a_{1}\}) \\ &\geq (\mu_{V} - \mu_{U}) \sqcup b(\{a_{1}\}) + (\mu_{V} - \mu_{U}) \sqcup b(K \setminus \{a_{1}\}) \\ &= \int_{\overline{\Omega}} b \, \mathrm{d}(\mu_{V} - \mu_{U}) \\ &= 0, \end{split}$$

where we used Lemma 2.4 (ii) in the last equality.

Step 2. Now let us suppose u, v are of the form  $u = L_K u, v = L_K v$ , with  $K = \partial \Omega \cup \{a_1, \ldots, a_M\}$ . The inequality (9) follows by applying Step 1 M times.

Step 3. Now let  $u, v \in W^{1,\infty}(\overline{\Omega})$  be as in the statement of the current proposition. Let  $\{a_k : k \in \mathbb{N}\} \subseteq \Omega$  be a dense subset of  $\Omega$ . For  $k \in \mathbb{N}$ , set  $u_k = L_{\partial \Omega \cup \{a_1, \dots, a_k\}} v$ ,  $v_k = L_{\partial \Omega \cup \{a_1, \dots, a_k\}} v$ .

We have that  $u_k \to u, v_k \to v$  uniformly, which implies

$$\begin{array}{cccc}
\mu_{v_k} \stackrel{*}{\rightharpoonup} \mu_v, & \mu_{u_k} \stackrel{*}{\rightharpoonup} \mu_u, \\
\mu_{v_k} \sqcup \phi \stackrel{*}{\rightharpoonup} \mu_v \sqcup \phi, & \mu_{u_k} \sqcup \phi \stackrel{*}{\rightharpoonup} \mu_u \sqcup \phi. \\
\end{array} \tag{10}$$

By the classical monotonicity of the subdifferential (see [Fig17, Lemma 2.7]),

$$\mu_{v_k}(\Omega) \le \mu_v(\Omega), \quad \mu_{u_k}(\Omega) \le \mu_u(\Omega) \quad \text{for all } k \in \mathbb{N}.$$

By (10),

$$\liminf_{k \to \infty} \mu_{v_k}(\Omega) \ge \mu_v(\Omega), \quad \liminf_{k \to \infty} \mu_{u_k}(\Omega) \ge \mu_u(\Omega).$$

Thus

$$\lim_{k \to \infty} \mu_{v_k}(\Omega) = \mu_v(\Omega) \,, \quad \lim_{k \to \infty} \mu_{u_k}(\Omega) = \mu_u(\Omega) \,,$$

which implies

$$\lim_{k \to \infty} \mu_{v_k} \, \lfloor \, \phi(\Omega) = \mu_v \, \lfloor \, \phi(\Omega) \,, \quad \lim_{k \to \infty} \mu_{u_k} \, \lfloor \, \phi(\Omega) = \mu_u \, \lfloor \, \phi(\Omega) \,,$$

and the claim of the proposition follows from the previous step.

In order to substantiate our claim from the introduction that Proposition 2.6 is of independent interest, we note in passing that as a direct byproduct, we obtain the following generalization of the well-known Alexandrov-Bakel'man-Pucci maximum principle (see e.g. [CC95] for the classical statement).

**Corollary 2.7.** Let  $\Omega \subseteq \mathbb{R}^n$  be convex, open and bounded, let  $u \in C^2(\Omega) \cap C^0(\overline{\Omega})$  satisfy the linear differential inequality

$$Lu = \sum_{i=1}^{n} a^{ij} \partial_{x_i} \partial_{x_j} u \le g \text{ in } \Omega$$

where  $a^{ij}: \Omega \to \mathbb{R}, i, j = 1, ..., n, A = (a^{ij})_{i,j=1,...,n} > 0$  in the sense of positive definite matrices,  $g \in L^n(\Omega)$ , and suppose that  $\phi \in C^0(\overline{\Omega})$  is a non-negative concave function. Then

$$u(x) \ge \min_{\partial \Omega} u - \frac{\operatorname{diam}\left(\Omega\right)}{n(\omega_n \phi(x))^{1/n}} \left\| \left(\frac{\phi}{\det A}\right)^{1/n} g \right\|_{L^n(\Gamma^-)} \quad \text{for } x \in \Omega \,,$$

where  $\Gamma^{-}$  is the set where u agrees with its convex envelope.

*Proof.* Let  $u^c$  denote the convex envelope of u, and  $\hat{u} := L_{\{x\}\cup\partial\Omega}u$ . We note

$$\omega_n \left(\frac{|u(x) - \min_{\partial \Omega} u|}{\operatorname{diam}\left(\Omega\right)}\right)^n \le \mu_{\hat{u}}(\{x\}).$$
(11)

By Proposition 2.6, we have

$$\phi(x)\mu_{\hat{u}}(\{x\}) = \int \phi \mathrm{d}\mu_{\hat{u}} \le \int \phi \mathrm{d}\mu_{u^c} = \int_{\Gamma^-} \phi \det \nabla^2 u \mathrm{d}x \,. \tag{12}$$

Furthermore,

$$\det A \det \nabla^2 u \le \left(\frac{\operatorname{Tr} A \nabla^2 u}{n}\right)^n . \tag{13}$$
11)-(13) and reordering yields the claim.

Combining the inequalities (11)-(13) and reordering yields the claim.

The following lemma is a slightly generalized version of the Alexandrov maximum principle [Fig17, Theorem 2.8]:

**Lemma 2.8.** Let  $\Omega \subseteq \mathbb{R}^n$  be open, convex, bounded,  $U \subseteq \Omega$  open and  $u \in C^0(\overline{\Omega})$  convex. Then for every  $x \in U$ 

$$\frac{L_{\partial U}u(x) - u(x)}{\operatorname{diam}(U)} \le C(n)\mu_u(U)^{1/n}.$$
(14)

*Proof.* We define the auxiliary function  $v := L_{\partial U \cup \{x\}} u$ . We will show that

$$\mathcal{L}^{n}(\partial^{-}v(x)) \ge C(n) \frac{|L_{\partial U}u(x) - u(x)|^{n}}{\operatorname{diam}(U)^{n}}.$$
(15)

By the monotonicity of the subdifferential (see [Fig17, Lemma 2.7]),

$$\mu_u(U) \ge \mu_v(U) = \mathcal{L}^n(\partial^- v(x)) \ge C(n) \frac{|L_{\partial U}u(x) - u(x)|^n}{\operatorname{diam}(U)^n},$$

which after rearranging yields (14).

Now to show (15):

Let 
$$p \in \partial^{-}L_{\partial U}u(x)$$
. If  $q \in B\left(p, \frac{L_{\partial U}u(x)}{\operatorname{diam}(U)}\right)$ , then  
 $q \cdot x - u(x) = p \cdot x - L_{\partial U}u(x) + (q - p) \cdot x + L_{\partial U}u(x) - u(x)$   
 $\geq \max_{z \in \partial U} p \cdot z - L_{\partial U}u(z) + (q - p) \cdot x + L_{\partial U}u(x) - u(x)$   
 $\geq \max_{z \in \partial U} q \cdot z - u(z) - |q - p||x - z| + L_{\partial U}u(x) - u(x)$   
 $\geq \max_{z \in \partial U} q \cdot z - u(z).$ 
This shows that  $\overline{B\left(p, \frac{L_{\partial U}u(x)}{U}\right)} \subseteq \partial^{-}v(x)$ .

s shows that L  $\left(p, \frac{1}{\operatorname{diam}(U)}\right) \subseteq 0 \quad v(x)$ 

As a consequence we get the following identity of the solution of (2).

**Lemma 2.9.** Let  $v_0$  be the solution of (2), and  $K := \partial \Omega \cup \{a_1, \ldots, a_N\}$ . Then  $v_0 = L_K v_0$ .

*Proof.* The inequality  $v_0 \leq L_K v_0$  follows directly from the definition of  $L_K v_0$ . Assume  $v_0(x) < L_K v(x)$  for some  $x \in \Omega \setminus \{a_1, \ldots, a_N\}$ . By Lemma 2.8 with  $U = \Omega \setminus \{a_1, \ldots, a_N\}$ , we obtain  $\mu_{L_K v_0}(U) > 0$ , and hence a contradiction.

2.2. Analysis of the singular Monge-Ampère equation (2). We will now treat the special case  $\Omega \subseteq \mathbb{R}^2$  (still supposing it to be open, convex and bounded). Let  $\bar{\mu} =$  $\sum_{i=1}^{N} \sigma_i \delta_{a_i} \in \mathcal{M}_+(\Omega)$ , with  $\sigma_i > 0, \{a_1, \ldots, a_N\} \subseteq \Omega$  and  $N \ge 1$ . We characterize the solution  $v_0$  to the Monge-Ampère equation with right hand side  $\bar{\mu}$ , see (2). In particular, Lemma 2.11 below will show that that  $\nabla v_0 \in BV(\Omega; \mathbb{R}^2)$ , and that its jump set  $J_{\nabla v_0}$  is contained in the set of line segments connecting the vertices  $\{a_1, \ldots, a_N\}$ . First we state and prove an auxiliary lemma:

**Lemma 2.10.** Let  $v_0 \in C^0(\overline{\Omega})$  be the solution to (2). Suppose that  $p \in \partial^- v_0(\Omega)$ ,  $[a_1a_2] \subseteq$  $(\partial^{-}v_{0})^{-1}(p), and$ 

$$(\partial^{-}v_{0})^{-1}(p) \cap \{x : (x - a_{1}) \cdot e_{12}^{\perp} > 0\} = \emptyset,$$
(16)

where  $e_{12} = \frac{a_2-a_1}{|a_2-a_1|}$ . Then there exists  $\varepsilon_0 > 0$  such that  $p + \varepsilon e_{12}^{\perp} \in \partial^- v_0(x)$  for every  $x \in [a_1a_2]$  and every  $\varepsilon \in (0, \varepsilon_0)$ .

*Proof.* Let  $x \in [a_1a_2]$ . The supporting function

v

$$A(z) = v_0(x) + p \cdot (z - x) = v_0(a_1) + p \cdot (z - a_1)$$

satisfies  $\{A = v_0\} = (\partial^- v_0)^{-1}(p)$ . There exists r > 0 such that

$$a_0(a) - A(a) > r \quad \text{for all } a \in \{a_1, \dots, a_N\} \setminus (\partial^- v_0)^{-1}(p)$$

Writing  $L := (\partial^{-} v_0)^{-1}(p) \cap \partial \Omega$ , the assumption (16) implies that there exists s > 0 with

$$e_{12}^{\perp} \cdot (z - a_1) < -s \quad \text{for } z \in L$$

Furthermore for every subset S of  $\partial \Omega$  with positive distance from L, there exists c(S) > 0 such that

$$v - A \ge c(S)$$
 on  $S$ .

Combining the above estimates, there exists  $\varepsilon_0 > 0$  depending on r and s such that for  $\varepsilon < \varepsilon_0$ , the affine function

$$A_{\varepsilon}(z) := A(z) + \varepsilon e_{12}^{\perp} \cdot (z - a_1)$$

satisfies  $A_{\varepsilon} = v_0$  on  $[a_1a_2]$  and

$$A_{\varepsilon} \leq v_0$$
 on  $\{a_1, \ldots, a_N\} \cup \partial \Omega$ .

By Lemma 2.9,  $v_0 = L_K v_0$  with  $K = \{a_1, \ldots, a_N\} \cup \partial \Omega$ , and hence the above implies that  $A_{\varepsilon}$  is a supporting function for  $v_0$ , and in turn that  $p + \varepsilon e_{12}^{\perp} \in \partial^- v_0(x)$ .

We will use the following notation to denote non-ordered pairs of labels: For  $(i, j) \in \{1, \ldots, N\}^2$ , we define the equivalence relation "~" by requiring  $(i, j) \sim (j, i)$  for all  $i, j \in \{1, \ldots, N\}$ . The set of non-ordered pairs is  $\{(i, j) \in \{1, \ldots, N\}^2 : i \neq j\} / \sim$ .

**Lemma 2.11.** Let  $\Omega \subseteq \mathbb{R}^2$  be open, convex, bounded, with  $C^{1,1}$  boundary. Let  $v_0 \in C^0(\overline{\Omega})$  be the solution to (2). Define the singular set

$$S_{v_0} := \{x \in \Omega : v_0 \text{ is not differentiable at } x\}.$$

Then

(i) There exists  $\mathcal{R}_{v_0} \subseteq \{(i,j) \in \{1,\ldots,N\}^2 : i \neq j\} / \sim$  such that

$$S_{v_0} = igcup_{(i,j)\in\mathcal{R}_{v_0}}[a_i a_j]$$
 .

(ii) There exists C > 0 only depending on  $\Omega, \bar{\mu}$  such that

$$|\nabla^2 v_0(x)| \le \frac{C}{\operatorname{dist}(x, \{a_1, \dots, a_N\})}$$

for almost every  $x \in \Omega \setminus S_{v_0}$ . In particular,  $v_0 \in W^{2,\infty}_{\text{loc}}(\Omega \setminus S_{v_0})$ .

- (iii) For each  $(i, j) \in \mathcal{R}_{v_0}$ , there exist  $b_{ij}^+, b_{ij}^- \in \Omega$  satisfying the following property:  $[a_i b_{ij}^+ a_j b_{ij}^-]$  is a non-degenerate rhombus with diagonal  $[a_i a_j]$ , and  $v_0$  is affine on  $[a_i b_{ij}^+ a_j]$  and on  $[a_i b_{ij}^- a_j]$ , with two different values of the gradient  $\nabla v_0$  on these sets.
- (iv) If  $N \ge 2$ , then for every  $i \in \{1, ..., N\}$  there exists  $j \in \{1, ..., N\}$  such that  $(i, j) \in \mathcal{R}_{v_0}$ .

**Remark 2.12.** Whenever  $(i, j) \in \mathcal{R}_{v_0}$ , we will say that  $[a_i a_j]$  is a *ridge*.

Proof. Let  $x_0 \in \Omega \setminus \{a_1, \ldots, a_N\}$  and  $p \in \partial^- v_0(x_0)$ . By Lemma 2.2, the extreme points of  $(\partial^- v_0)^{-1}(p)$  are contained in  $\partial \Omega \cup \{a_1, \ldots, a_N\}$ . At least one of the following holds: (a)  $(\partial^- v_0)^{-1}(p) \cap \partial \Omega \neq \emptyset$ , or (b)  $\#\partial^- v_0(x_0) > 1$ , or (c)  $(\partial^- v_0)^{-1}(p) \cap \partial \Omega = \emptyset$  and  $\#\partial^- v_0(x_0) = 1$ . (a)  $(\partial^- v_0)^{-1}(p) \cap \partial\Omega \neq \emptyset$ . In this case, since  $v_0(z) = 0 \ge v_0(x_0) + p \cdot (z - x_0)$  for all  $z \in \partial\Omega$ , p must be a positive multiple of the outer normal  $n(z_0)$  for any  $z_0 \in (\partial^- v_0)^{-1}(p) \cap \partial\Omega$ . In addition, there must be at least one vertex  $a \in (\partial^- v_0)^{-1}(p) \cap \{a_1, \ldots, a_N\}$ , since otherwise  $v_0 = 0$  on  $(\partial^- v_0)^{-1}(p)$ , which would imply  $v_0 = 0$  in  $\overline{\Omega}$ . This gives us the exact form

$$p = \frac{v_0(a)}{n(z_0) \cdot (a - z_0)} n(z_0) \,. \tag{17}$$

(b)  $\#\partial^- v_0(x_0) > 1$ . Since  $0 = \overline{\mu}(\{x_0\}) = \mathcal{L}^2(\partial^- v_0(x_0))$ , the subdifferential must be a line segment  $\partial^- v_0(x_0) = [p^- p^+]$  with  $p^+ \neq p^-$ . For subgradients  $p \in [p^- p^+] \setminus \{p^-, p^+\}$ , we have

$$v_0(x_0) + p \cdot (x - x_0) \le \max(v_0(x_0) + p^- \cdot (x - x_0), v_0(x_0) + p^+ \cdot (x - x_0)) \quad \text{for } x \in \Omega,$$

with equality precisely on  $x_0 + (p^+ - p^-)^{\perp}$ . This shows that  $(\partial^- v_0)^{-1}(p)$  is a line segment orthogonal to  $p^+ - p^-$  for such p.

By (a), an extreme point of such a line segment cannot be a boundary point, so that  $(\partial^- v_0)^{-1}(p) = [a_i a_j]$  for some  $i, j \in \{1, \ldots, N\}$ . In addition,

$$[a_i a_j] \subsetneq (\partial^- v_0)^{-1} (p^{\pm}), \qquad (18)$$

since the equality  $[a_i a_j] = (\partial^- v_0)^{-1} (p^{\pm})$  would imply the existence of  $\varepsilon_0 > 0$  such that

$$p^{\pm} + t e_{ij}^{\perp} \in \partial^- v_0(x_0) \quad \text{ for } |t| < \varepsilon_0$$

by Lemma 2.10, where  $e_{ij} = (a_j - a_i)/|a_j - a_i|$ . This would be a contradiction to the choice of  $p^{\pm}$  as extreme points of  $\partial v^-(x_0)$ . Now (18) implies that  $\nabla v_0 = p^-$ ,  $\nabla v_0 = p^+$  respectively on at least two non-degenerate triangles bordering  $[a_i a_j]$ , which we may denote by  $[a_i b_{ij}^+ a_j]$ ,  $[a_i b_{ij}^+ a_j]$ , with a choice of  $b_{ij}^{\pm}$  that makes  $[a_i b_{ij}^+ a_j b_{ij}^-]$  a rhombus.

(c)  $(\partial^{-}v_{0})^{-1}(p) \cap \partial\Omega = \emptyset$  and  $\#\partial^{-}v_{0}(x_{0}) = 1$ . In this case all extreme points of

$$(\partial^{-}v_{0})^{-1}(p) = (\partial^{-}v_{0})^{-1}(\nabla v_{0}(x_{0}))$$

are vertices, and we may write  $(\partial^- v_0)^{-1}(p) = [a_{i_1} \dots a_{i_M}]$ . Hence, if  $x_0 \in \partial(\partial^- v_0)^{-1}(p)$ , there exist  $i, j \in \{1, \dots, N\}$  such that  $x_0 \in [a_i a_j]$  and

$$(\partial^{-}v_{0})^{-1}(p) \cap \{x : (x - x_{0}) \cdot e_{ij}^{\perp} > 0\} = 0$$

which produces the contradiction  $\#\partial^- v_0(x_0) > 1$  by Lemma 2.10. Hence  $x_0$  must be an interior point of  $(\partial^- v_0)^{-1}(p)$ , i.e.  $v_0$  is affine in a neighborhood of  $x_0$ .

Proof of (i), (ii), (iii):

In particular we may draw the following from the case distinction (a)-(c) above: If  $x \in \Omega \setminus \bigcup_{(i,j) \in \mathcal{R}_{v_0}} [a_i a_j]$ , then either  $v_0$  is affine in a neighborhood of x, or by (17),

$$\nabla v_0(x) = p = \frac{v_0(a)}{n(z) \cdot (a-z)} n(z), \tag{19}$$

for some vertex  $a \in \{a_1, \ldots, a_N\}$  and some boundary point  $z \in \partial \Omega$ . If  $x \in [az]$ , the boundary point z = z(x, a) is defined through the locally Lipschitz implicit function

$$\begin{cases} z(x,a) \in \partial \Omega\\ x \in [az(x,a)] \end{cases}$$

A direct calculation shows that  $|\nabla_x z(x,a)| \leq \frac{C(\Omega,a)}{|x-a|}$ . By the chain rule,  $|\nabla^2 v_0(x)| \leq \frac{C}{|x-a|}$ , where C only depends on  $\Omega$  and  $\bar{\mu}$ . This proves (i) and (ii), letting  $\mathcal{R}_{v_0}$  be the set of pairs (i, j) that appear in (b). Point (iii) follows immediately from the constructions in (b).

Proof of (iv): Assuming that N > 1, consider a vertex  $a \in \{a_1, \ldots, a_N\}$ . We claim that there exists  $p \in \partial^- v_0(a)$  such that  $(\partial^- v_0)^{-1}(p)$  contains at least one more vertex  $a' \neq a$ . Indeed, if this were not so, then by convexity of  $\Omega$  we would have

$$\Omega \subseteq \bigcup_{p \in \partial^- v_0(a)} (\partial^- v_0)^{-1}(p) \,,$$

where for each of the sets  $(\partial^- v_0)^{-1}(p)$  on the right hand side, all extreme points except a are contained in  $\partial\Omega$ . But  $p \in \partial^- v_0(a)$  and  $x \in (\partial^- v_0)^{-1}(p) \setminus \{a\}$  implies that x is an element but not an extreme point of  $(\partial^- v_0)^{-1}(p)$ , which in turn implies  $\mu_{v_0}(\{x\}) = 0$ . Hence  $\operatorname{supp} \bar{\mu} = \operatorname{supp} \mu_{v_0} = \{a\}$ , which is a contradiction to the assumption that the support of  $\bar{\mu} = \mu_{v_0}$  contains at least two points. This proves the claim that there exists  $a' \in \{a_1,\ldots,a_N\} \setminus \{a| \text{ such that } [aa'] \subseteq (\partial^- v_0)^{-1}(p)$ . We note that a' is an extreme point of  $(\partial^- v_0)^{-1}(p)$ , since otherwise we would again obtain by an analogous argument to the one that we have just used, that  $\mu_{v_0}(\{a'\}) = 0$ .

So let  $p \in \partial^{-}v_{0}(a) \cap \partial^{-}v_{0}(a')$  with  $a, a' \in \{a_{1}, \ldots, a_{N}\}, a \neq a', a$  and a' being extreme points of  $(\partial^{-}v_{0})^{-1}(p)$ . Since  $v_{0} = 0$  on  $\partial\Omega$  and  $v_{0} < 0$  on  $\Omega$ , we have that  $(\partial^{-}v_{0})^{-1}(p) \cap$  $\partial\Omega$  is either empty, consists of a single point, or of a line segment. Since all extreme points of  $(\partial^{-}v_{0})^{-1}(p)$  lie either on  $\partial\Omega$  or in  $\{a_{1}, \ldots, a_{N}\}$ , there must be a vertex  $a'' \in$  $\partial(\partial^{-}v_{0})^{-1}(p) \cap \{a_{1}, \ldots, a_{N}\} \setminus \{a\}$  (possibly, but not necessarily identical to a') such that  $[a a''] \subseteq \partial(\partial^{-}v_{0})^{-1}(p)$ . Without loss of generality, we may assume that the conditions of Lemma 2.10 are fulfilled for  $a_{1} \equiv a, a_{2} \equiv a''$ . Hence, for  $x \in [aa''] \setminus \{a, a''\} = [a_{1}a_{2}] \setminus$  $\{a_{1}, a_{2}\}, \#\partial^{-}v_{0}(x) > 1$ , which yields  $(1, 2) \in \mathcal{R}_{v_{0}}$ .

## 3. PROOF OF THE UPPER BOUND

## 3.1. Proof of the upper bound for convex out-of-plane component.

**Lemma 3.1.** Assume that  $[abc] \subseteq \mathbb{R}^2$  is a non-degenerate triangle whose angles are bounded away from 0 and  $\pi$ ,  $w_0 \in \mathbb{R}^2$ , and  $C_1 > 0$ . Then there exists  $\mathbf{w} \in W^{1,2}_{\text{loc}}([abc]^\circ; \mathbb{R}^2)$ and  $\varepsilon_0 > 0$  such that for every  $\varepsilon \in (0, \varepsilon_0)$ , the following properties are fulfilled:

$$\mathbf{w} = w_0 \qquad on \ [ab]$$
$$\mathbf{w} = 0 \qquad on \ (\partial [abc]) \setminus [ab]$$
$$|\mathbf{w}| \lesssim |w_0| \qquad in \ [abc]$$
$$|\nabla \mathbf{w}(x)| \lesssim |w_0| \operatorname{dist}(x, \{a, b\})^{-1} \ in \ [abc]$$
$$\int_{[abc] \setminus B_{\varepsilon}} |\nabla \mathbf{w}|^2 \mathrm{d}x \lesssim |w_0|^2 \log \frac{1}{\varepsilon}$$
15

where  $B_{\varepsilon} = B(a, C_1 \varepsilon) \cup B(b, C_1 \varepsilon) \cup B(c, C_1 \varepsilon)$ . The implicit constants in the inequalities above may depend on [abc] and  $C_1$ , but not on  $w_0$  or  $\varepsilon$ .

*Proof.* We may assume that a = (-L, 0), b = (L, 0) and  $c \cdot e_2 > 0$ . By assumption there exists  $d \in (0, 1)$  bounded away from 0 such that

$$\left|\frac{(b-a)^{\perp}}{|b-a|} \cdot \frac{c-a}{|c-a|}\right| > d.$$

Choose  $f \in C^2([0, L]; [0, \infty))$  such that f(x) = dx for  $x \in [0, L/2]$ ,  $f' \ge 0$ ,  $f'' \le 0$  and f'(L) = 0. Choose  $\varphi \in C^{\infty}(\mathbb{R})$  such that  $\varphi(x) = 1$  for  $x \le 0$ ,  $\varphi(x) = 0$  for  $x \ge 1$  and  $0 \le \varphi \le 1$ . Now for  $x \in [abc]$  set

$$\mathbf{w}(x) = \begin{cases} w_0 \varphi\left(\frac{x_2}{f(L+x_1)}\right) & \text{if } x_1 \le 0\\ w_0 \varphi\left(\frac{x_2}{f(L-x_1)}\right) & \text{if } x_1 \ge 0 \,. \end{cases}$$
(20)

From this definition it follows

$$\nabla \mathbf{w}(x)|^{2} = |w_{0}|^{2} \left( \left| \frac{x_{2}f'}{f^{2}} \varphi' \right|^{2} + \frac{|\varphi'|^{2}}{f^{2}} \right)$$
$$\lesssim |w_{0}|^{2} \max(|x-a|^{-2}, |x-b|^{-2}),$$

and hence

$$\int_{[abc]\setminus B_{\varepsilon}} |\nabla \mathbf{w}|^2 \mathrm{d}x \lesssim |w_0|^2 \log \frac{1}{\varepsilon}$$

as claimed. The other inequalities also follow straightforwardly from the explicit formulae for  $\mathbf{w}$  and  $\nabla \mathbf{w}$ .

Proof of the upper bound in Theorem 1.1. For  $(i, j) \in \mathcal{R}_{v_0}$  (see Lemma 2.11 for the notation), we will write

$$e_{ij} = \frac{a_j - a_i}{|a_j - a_i|} \,.$$

According to Lemma 2.11, for each  $(i, j) \in \mathcal{R}_{v_0}$ , we may choose two points  $b_{ij}^+, b_{ij}^-$  such that  $\nabla v_0 = p^+$  on  $[a_i b_{ij}^+ a_j]$  and  $\nabla v_0 = p^-$  on  $[a_i b_{ij}^- a_j]$ , for some  $p^+, p^- \in \mathbb{R}^2$  with  $p^+ \neq p^-$ . We may define the convex one-homogeneous function  $W_{ij} : \mathbb{R} \to \mathbb{R}$  by requiring

$$v_0(x) = v_0(a_i + e_{ij} \cdot (x - a_i)e_{ij}) + W_{ij}((x - a_i) \cdot e_{ij}^{\perp}) \quad \text{for } x \in R_{ij} = [a_i b_{ij}^+ a_j b_{ij}^-].$$

Now choose a convex  $C^2$  function  $w_{ij} : \mathbb{R} \to \mathbb{R}$  and  $M_{ij} > 0$  such that  $w_{ij} \ge W_{ij}$ ,  $w_{ij}(x) = W_{ij}(x)$  for  $|x| > M_{ij}$  and

$$\int_{-M_{ij}}^{M_{ij}} (W'_{ij}(t))^2 - (w'_{ij}(t))^2 dt = W'_{ij}(M_{ij}) - W'_{ij}(-M_{ij})$$

Such a choice of  $w_{ij}, M_{ij}$  is always possible (note that the right hand side above does not depend on  $M_{ij}$ ) and depends only on the two values of  $W'_{ij}$  (and hence on  $v_0$ ). For  $(i, j) \in \mathcal{R}_{v_0}$  let  $S^{\varepsilon}_{ij}$  denote the strip of width  $2\varepsilon M_{ij}$  around  $[a_i a_j]$ ,

$$S_{ij}^{\varepsilon} = \{x : (x - a_i) \cdot e_{ij} \in [0, |a_i - a_j|], \varepsilon^{-1}(x - a_i) \cdot e_{ij}^{\perp} \in [-M_{ij}, M_{ij}]\}.$$

For  $\varepsilon > 0$  we may define the convex function  $w_{ij,\varepsilon} : \Omega \to \mathbb{R}$ ,

$$w_{ij,\varepsilon}(x) := v_0(a_i + e_{ij} \cdot (x - a_i)e_{ij}) + \varepsilon w_{ij}(\varepsilon^{-1}(x - a_i) \cdot e_{ij}^{\perp})$$
16

We note that there exists  $C_1 > 0$  that only depends on  $v_0$  (and hence on  $\Omega, \bar{\mu}$ ) such that

 $v_0(x) \ge w_{ij,\varepsilon}(x)$  for  $x \in \Omega \setminus \left(S_{ij}^{\varepsilon} \cup B(a_i, C_1 \varepsilon) \cup B(a_j, C_1 \varepsilon)\right)$  for all  $(i, j) \in \mathcal{R}_{v_0}$ ,

and

$$\|\nabla^2 w_{ij,\varepsilon}\|_{L^{\infty}} \lesssim \varepsilon^{-1}$$

Now we define the convex function

$$\tilde{v}_{\varepsilon}(x) = \max\{v_0(x), w_{ij,\varepsilon}(x) : (i,j) \in \mathcal{R}_{v_0}\} \text{ for } x \in \Omega$$

Writing

$$B_{\varepsilon} = \bigcup_{i=1}^{N} B(a_i, C_1 \varepsilon)$$

our construction guarantees that

 $\tilde{v}_{\varepsilon} \in C^2(\Omega \setminus B_{\varepsilon})$ ,

with

$$\|\nabla^2 \tilde{v}_{\varepsilon}\|_{L^{\infty}(\Omega \setminus B_{\varepsilon})} \le C\varepsilon^{-1}.$$
(21)

We further modify  $\tilde{v}_{\varepsilon}$  in order to obtain an analogous estimate on all of  $\Omega$  by using Lemma 2.3. More precisely, we extend  $\tilde{v}_{\varepsilon}$  to a globally Lipschitz convex function on  $\mathbb{R}^2$ , and choose some convex open set  $\tilde{\Omega} \subset \Omega$  with  $\{a_1, \ldots, a_N\} \subseteq \tilde{\Omega}$ . We choose c > 0 and  $\varepsilon_0$  that only depend on  $v_0, C_1, \tilde{\Omega}$  such that for every  $\varepsilon \in (0, \varepsilon_0)$  the following are fulfilled:

- The function  $\hat{v}_{\varepsilon} := P_{c\varepsilon} \tilde{v}_{\varepsilon}$  satisfies  $\hat{v}_{\varepsilon} = \tilde{v}_{\varepsilon}$  on  $\tilde{\Omega} \setminus B_{\varepsilon}$ ,  $\|\nabla^2 v_0\|_{L^{\infty}(\Omega \setminus \tilde{\Omega})} \leq C \varepsilon^{-1}$ .

Then we set

$$v_{\varepsilon}(x) = \begin{cases} v_0(x) & \text{if } x \in \Omega \setminus \tilde{\Omega} \\ \hat{v}_{\varepsilon}(x) & \text{if } x \in \tilde{\Omega} \,. \end{cases}$$

By construction this choice satisfies

$$\begin{split} \|\nabla^2 v_{\varepsilon}\|_{L^{\infty}(\Omega)} &\leq C\varepsilon^{-1} \\ v_{\varepsilon} &= w_{ij,\varepsilon} \quad \text{in } S^{\varepsilon}_{ij} \setminus B_{\varepsilon} \\ v_{\varepsilon} &= 0 \quad \text{on } \partial\Omega \,. \end{split}$$

Next we define the in-plane deformation  $\mathbf{u}_{\varepsilon}$ . For  $(i, j) \in \mathcal{R}_{v_0}$ , set

$$S_{ij}^{\varepsilon*} := S_{ij}^{\varepsilon} \setminus B_{\varepsilon}$$

By possibly increasing  $C_1$  we get that for each  $(i, j) \in \mathcal{R}_{v_0}$ ,

$$S_{ij}^{\varepsilon*} \subseteq R_{ij} = \left[a_i b_{ij}^+ a_j b_{ij}^-\right],$$

and for  $(i, j) \neq (k, l) \in \mathcal{R}_{v_0}$ ,

$$S_{ij}^{\varepsilon*} \cap S_{kl}^{\varepsilon*} = \emptyset.$$

For  $(i, j) \in \mathcal{R}_{v_0}$  and  $|t| < M_{ij}$ , set

$$f_{ij}(t) = W'_{ij}(-M_{ij}) + \int_{-M_{ij}}^{t} (W'_{ij}(s))^2 - (w'_{ij}(s))^2 ds$$
$$g_{ij}(t) = 2 \frac{v_0(a_j) - v_0(a_i)}{|a_j - a_i|} \int_{-M_{ij}}^{t} W'_{ij}(s) - w'_{ij}(s) ds$$

and for  $x \in S_{ij}^{\varepsilon}$ ,

$$\mathbf{u}_{ij}^{\varepsilon}(x) := \frac{\varepsilon}{2} \left( f_{ij} \left( \varepsilon^{-1} (x - a_i) \cdot e_{ij}^{\perp} \right) e_{ij}^{\perp} + g_{ij} \left( \varepsilon^{-1} (x - a_i) \cdot e_{ij}^{\perp} \right) e_{ij} \right) \,.$$

We note that this definition implies

$$\mathbf{u}_{ij}^{\varepsilon}(x) = \frac{\varepsilon}{2} W_{ij}'(\pm M_{ij}) e_{ij}^{\perp} \quad \text{for } \varepsilon^{-1}(x - a_i) \cdot e_{ij}^{\perp} = \pm M_{ij} , \qquad (22)$$

and for  $x \in S_{ij}^{\varepsilon} \setminus [a_i a_j]$ ,

$$\nabla \mathbf{u}_{ij}^{\varepsilon}(x) + (\nabla \mathbf{u}_{ij}^{\varepsilon})^{T}(x) 
= f_{ij}'\left(\varepsilon^{-1}(x-a_{i})\cdot e_{ij}^{\perp}\right)e_{ij}^{\perp}\otimes e_{ij}^{\perp} + g_{ij}'\left(\varepsilon^{-1}(x-a_{i})\cdot e_{ij}^{\perp}\right)\left(e_{ij}\otimes e_{ij}^{\perp} + e_{ij}^{\perp}\otimes e_{ij}\right) 
= \left(\left(W_{ij}'\left(\varepsilon^{-1}(x-a_{i})\cdot e_{ij}^{\perp}\right)\right)^{2} - \left(w_{ij}'\left(\varepsilon^{-1}(x-a_{i})\cdot e_{ij}^{\perp}\right)\right)^{2}\right)e_{ij}^{\perp}\otimes e_{ij}^{\perp} 
+ \left(W_{ij}'\left(\varepsilon^{-1}(x-a_{i})\cdot e_{ij}^{\perp}\right) - w_{ij}'\left(\varepsilon^{-1}(x-a_{i})\cdot e_{ij}^{\perp}\right)\right) 
\times \frac{v(a_{j}) - v(a_{i})}{|a_{j} - a_{i}|}\left(e_{ij}\otimes e_{ij}^{\perp} + e_{ij}^{\perp}\otimes e_{ij}\right) 
= \nabla v_{0}(x)\otimes\nabla v_{0}(x) - \nabla v_{\varepsilon}(x)\otimes\nabla v_{\varepsilon}(x).$$
(23)

Consider the part of  $\partial S_{ij}^{\varepsilon*}$  that is parallel to  $[a_i a_j]$ . It consists of two line segments, that we denote by  $l_{ij}^{\pm\varepsilon}$ . Let  $T_{ij}^{\pm\varepsilon}$  denote the isosceles triangles with base  $l_{ij}^{\pm\varepsilon}$  and base angle  $\alpha$ , oriented such that the interior of  $T_{ij}^{\pm\varepsilon}$  has no intersection with  $S_{ij}^{\varepsilon}$ . We may choose  $\alpha > 0$  independently of  $\varepsilon$  such that the intersection for any two triangles in the set  $\{T_{ij}^{\pm\varepsilon}, (i, j) \in \mathcal{R}_{v_0}\}$  is at most a single point. Now we apply Lemma 3.1 to the triangles  $T_{ij}^{\pm\varepsilon}$ with the constant boundary value on  $l_{ij}^{\pm\varepsilon}$  given by the value of  $\mathbf{u}_{ij}^{\varepsilon}$  on that set, obtaining  $\mathbf{w}_{ij}^{\pm} \in W_{\text{loc}}^{1,2}((T_{ij}^{\pm\varepsilon})^{\circ}; \mathbb{R}^2)$  with the properties

$$\begin{split} \mathbf{w}_{ij}^{\pm} &= \frac{\varepsilon}{2} W_{ij}'(\pm M_{ij}) e_{ij}^{\pm} \text{ on } l_{ij}^{\pm\varepsilon} \\ \mathbf{w}_{ij}^{\pm} &= 0 & \text{ on } \partial T_{ij}^{\pm\varepsilon} \setminus l_{ij}^{\pm\varepsilon} \\ |\mathbf{w}_{ij}^{\pm}| &\lesssim \varepsilon & \text{ on } \partial B_{2\varepsilon} \cap T_{ij}^{\pm\varepsilon} \\ |\nabla \mathbf{w}_{ij}^{\pm}| &\lesssim 1 & \text{ on } \partial B_{2\varepsilon} \cap T_{ij}^{\pm\varepsilon} \\ \int_{T_{ij}^{\pm\varepsilon} \setminus B_{2\varepsilon}} |\nabla \mathbf{w}_{ij}^{\pm}|^2 \mathrm{d}x \lesssim \varepsilon^2 \log \frac{1}{\varepsilon} \end{split}$$

where

$$B_{2\varepsilon} = \bigcup_{i=1}^{N} B(a_i, 2C_1\varepsilon) \,.$$

On  $\Omega \setminus B_{2\varepsilon}$ , we may now define  $\mathbf{u}_{\varepsilon}$  by

$$\mathbf{u}_{\varepsilon}(x) = \begin{cases} \mathbf{u}_{ij}^{\varepsilon}(x) & \text{if } x \in S_{ij}^{\varepsilon*} \setminus B_{2\varepsilon} \\ \mathbf{w}_{ij}^{\pm}(x) & \text{if } x \in T_{ij}^{\pm\varepsilon} \\ 0 & \text{else.} \\ 18 \end{cases}$$

This definition satisfies  $\mathbf{u}_{\varepsilon} \in W^{1,2}(\Omega \setminus B_{2\varepsilon})$ . We note

$$\sum_{(i,j)\in\mathcal{R}_{v_0}} \int_{S_{ij}^{\varepsilon*} \setminus B_{2\varepsilon}} |\nabla \mathbf{u}_{\varepsilon} + \nabla \mathbf{u}_{\varepsilon}^T + \nabla v \otimes \nabla v - \nabla v_0 \otimes \nabla v_0|^2 \mathrm{d}x = 0$$

and

$$\sum_{(i,j)\in\mathcal{R}_{v_0}} \int_{T_{ij}^{\pm\varepsilon}\setminus B_{2\varepsilon}} |\nabla \mathbf{u}_{\varepsilon} + \nabla \mathbf{u}_{\varepsilon}^T + \nabla v \otimes \nabla v - \nabla v_0 \otimes \nabla v_0|^2 \mathrm{d}x$$

$$\leq 2 \sum_{(i,j)\in\mathcal{R}_{v_0}} \int_{T_{ij}^{\pm\varepsilon}\setminus B_{2\varepsilon}} |\nabla \mathbf{u}_{\varepsilon}|^2 \mathrm{d}x$$

$$\lesssim \varepsilon^2 \log \frac{1}{\varepsilon}.$$
(24)

By the estimates on  $\mathbf{u}_{ij}^{\varepsilon}$ ,  $\nabla \mathbf{u}_{ij}^{\varepsilon}$ ,  $\mathbf{w}_{ij}^{\pm}$ ,  $\nabla \mathbf{w}_{ij}^{\pm}$  on  $\partial B_{2\varepsilon}$ , we may extend  $\mathbf{u}_{\varepsilon}$  to a map in  $W^{1,2}(\Omega; \mathbb{R}^2)$  satisfying

$$\int_{B_{2\varepsilon}} |\nabla \mathbf{u}_{\varepsilon}|^2 \mathrm{d}x \lesssim \varepsilon^2 \,,$$

and hence

$$\int_{B_{2\varepsilon}} |\nabla \mathbf{u}_{\varepsilon} + \nabla \mathbf{u}_{\varepsilon}^{T} + \nabla v \otimes \nabla v - \nabla v_{0} \otimes \nabla v_{0}|^{2} \mathrm{d}x \lesssim \varepsilon^{2} \,.$$

Combining these estimates, we get

$$\int_{\Omega} |\nabla \mathbf{u}_{\varepsilon} + \nabla \mathbf{u}_{\varepsilon}^{T} + \nabla v_{\varepsilon} \otimes \nabla v_{\varepsilon} - \nabla v_{0} \otimes \nabla v_{0}|^{2} \mathrm{d}x \lesssim \varepsilon^{2} \log \frac{1}{\varepsilon}$$

Furthermore,

$$\begin{split} \int_{\Omega} |\nabla^2 v_{\varepsilon}|^2 \mathrm{d}x &\lesssim \int_{B_{\varepsilon}} |\nabla^2 v_{\varepsilon}|^2 \mathrm{d}x + \int_{\Omega \setminus B_{\varepsilon} \setminus U_2^{\varepsilon}} |\nabla^2 v_{\varepsilon}|^2 \mathrm{d}x + \sum_{(i,j) \in \mathcal{R}_{v_0}} \int_{S_{ij}^{\varepsilon*}} |\nabla^2 v_{\varepsilon}|^2 \mathrm{d}x \\ &\lesssim 1 + \log \frac{1}{\varepsilon} + \varepsilon^{-1} \,. \end{split}$$

Choosing  $\varepsilon := h^{2/3} \left( \log \frac{1}{h} \right)^{-1/3}$ , we obtain

$$E_h(\mathbf{u}_{\varepsilon}, v_{\varepsilon}) \lesssim h^{4/3} \left(\log \frac{1}{h}\right)^{1/3}.$$

**Remark 3.2.** The logarithmic factor in the upper bound comes from the membrane energy density away from the vertices, see (24). Since  $\nabla v_{\varepsilon}$  necessarily needs to transition between values that differ by O(1) when crossing a ridge, convexity forces the integral of the in-plane contribution to the strain  $\nabla \mathbf{u} + \nabla \mathbf{u}^T$  across the ridge to be of the same order as the transition layer. The authors believe that this cannot be counterbalanced by a different choice of  $v_{\varepsilon}$  in a way that would make the logarithmic factors disappear, since any deviation from  $v_0$  far away from the vertices is energetically costly. 3.2. Upper bound for non-convex out-of-plane component. If we allow for nonconvex out-of-plane deformations v, then we may achieve the scaling  $h^{5/3}$ . In the statement below,  $\Omega \subseteq \mathbb{R}^2$  and  $\bar{\mu} = \sum_{i=1}^N \sigma_i \delta_{a_i}$  are as in Theorem 1.1.

**Proposition 3.3.** There exists C > 0 that only depends on  $\Omega, \overline{\mu}$  with the following property: For h > 0 there exists  $(\mathbf{u}_h, v_h) \in W^{1,2}(\Omega; \mathbb{R}^2) \times W^{2,\infty}(\Omega)$  such that

$$E_h(\mathbf{u}_h, v_h) \le Ch^{5/3}$$

Before proving Proposition 3.3, we note that in combination with Theorem 1.1 it implies that minimizers of  $E_h$  have non-convex out-of-plane component:

**Corollary 3.4.** There exists  $h_1 < h_0$  such that the following holds true for every  $0 < h < h_1$ : If  $(\mathbf{u}_h, v_h) \in \arg \min E_h$ , then  $v_h$  is non-convex.

The proof of Proposition 3.3 is based on Lemma 16 from [COT17], which itself is an adaptation of the construction for the single ridge by Conti and Maggi [CM08] to the Föppl–von Kármán approximation. Below we cite the Föppl–von Kármán version in full for the reader's convenience. In the lemma we use the notation

$$\bar{E}_h[\mathbf{u}, v, W] = \int_W |\nabla \mathbf{u} + \nabla \mathbf{u}^T + \nabla v \otimes \nabla v|^2 + h^2 |\nabla^2 v|^2 \mathrm{d}x$$

**Lemma 3.5.** [COT17, Lemma 16] Let  $[abcd] \subseteq \mathbb{R}^2$  be a non-degenerate quadrilateral with diagonals [ac] and [bd], contained in the square with diagonal [ac]. Furthermore, let  $(\mathbf{u}_0, w_0) : [abcd] \to \mathbb{R}^2 \times \mathbb{R}$  be continuous functions, affine on [abc] and [adc], with  $\nabla \mathbf{u}_0 + \nabla \mathbf{u}_0^T + \nabla w_0 \otimes \nabla w_0 = 0$ . Then for all  $h \in (0, l/8)$  there are  $(\mathbf{u}_h, w_h) \in C^2([abcd] \setminus \{a, c\}; \mathbb{R}^2 \times \mathbb{R})$  such that  $\mathbf{u}_0 = \mathbf{u}_h, \nabla \mathbf{u}_0 = \nabla \mathbf{u}_h, w_0 = w_h$  on  $\partial [abcd] \setminus \{a, c\}, |\nabla \mathbf{u}_h| + |\nabla w_h| \leq C(\mathbf{u}_0, w_0)$  and

$$\bar{E}_{h}[\mathbf{u}_{h}, w_{h}, [abcd] \setminus (B(a, h) \cup B(c, h))] \le Ch^{5/3} l^{1/3},$$
(25)

where l = |a - c|, and the constant C may depend on the angles of [abcd], and on  $\mathbf{u}_0$  and  $w_0$ , but not on h.

Proof of Proposition 3.3. Using the notations of Lemmata 2.11 and 3.5, for every pair  $(i, j) \in \mathcal{R}_{v_0}$ , we may write  $R_{ij} = [abcd]$  with  $a_i = a$ ,  $a_j = c$ , set  $w_0^{ij} = v_0$  and define  $\mathbf{u}_0^{ij}$  by requiring it to be continuous on [abcd] and affine on both [abc] and [adc], with

$$\nabla \mathbf{u}_0^{ij} + \left(\nabla \mathbf{u}_0^{ij}\right)^T = -\nabla v_0 \otimes \nabla v_0 \,, \tag{26}$$

where  $v_0$  is the unique solution of (2). The equation (26) indeed possesses a solution: Note that after a suitable rotation of the domain, we may assume that  $[ac] \subseteq \mathbb{R} \times \{0\}$ , and

$$v_0(x) = \begin{cases} A_1 x_1 + A_2 x_2 & \text{if } x_2 \ge 0\\ A_1 x_1 + A_3 x_2 & \text{if } x_2 < 0 \end{cases}$$

Then a solution of (26) is given by

$$\mathbf{u}_{0}^{ij}(x_{1}, x_{2}) = \begin{cases} (A_{4}x_{1} + A_{5}x_{2}, A_{6}x_{2}) & \text{if } x_{2} \ge 0\\ (A_{4}x_{1} + A_{7}x_{2}, A_{8}x_{2}) & \text{if } x_{2} < 0 \end{cases}$$

where  $A_4, \ldots, A_8$  are given by

$$-2A_4 = A_1^2, \quad -A_5 = A_1A_2, \quad -2A_6 = A_2^2, \quad -A_7 = A_1A_3, \quad -2A_8 = A_3^2.$$

Applying Lemma 3.5 we obtain  $(\mathbf{u}_h^{ij}, w_h^{ij})$  in  $C^2([abcd] \setminus \{a, c\}; \mathbb{R}^2 \times \mathbb{R})$  that satisfy the bound (25), and  $\mathbf{u}_h^{ij} = \mathbf{u}_0^{ij}, w_h^{ij} = v_0$  on  $\partial[abcd]$ . This finishes the construction on  $R_{ij}$ . Now we set

$$\mathbf{u}_{h} = \begin{cases} \mathbf{u}_{h}^{ij} - \mathbf{u}_{0}^{ij} & \text{in } R_{ij} \text{ for } (i,j) \in \mathcal{R}_{v_{0}} \\ 0 & \text{in } \Omega \setminus \bigcup_{(i,j) \in \mathcal{R}_{v_{0}}} R_{ij} \end{cases}$$

and

$$v_h = \begin{cases} w_h^{ij} & \text{in } R_{ij} \text{ for } (i,j) \in \mathcal{R}_{v_0} \\ v_0 & \text{in } \Omega \setminus \bigcup_{(i,j) \in \mathcal{R}_{v_0}} R_{ij} . \end{cases}$$

The energy satisfies

$$\int_{\Omega \setminus \bigcup_{i=1}^{N} B(a_i,h)} |\nabla \mathbf{u}_h + \nabla \mathbf{u}_h^T + \nabla v_h \otimes \nabla v_h - \nabla v_0 \otimes \nabla v_0|^2 + h^2 |\nabla^2 v_h|^2 dx$$
$$= \sum_{(i,j) \in \mathcal{R}_{v_0}} \bar{E}_h \left[ \mathbf{u}_h^{ij}, w_h^{ij}, R_{ij} \setminus (B(a_i,h) \cup B(a_j,h)) \right]$$
$$+ h^2 \int_{\Omega \setminus (\cup_{(i,j) \in \mathcal{R}_{v_0}} R_{ij} \cup (\cup_{i=1}^{N} B(a_i,h)))} |\nabla^2 v_0|^2 dx$$
$$< Ch^{5/3}.$$

It remains to smoothen our construction on  $\bigcup_{i=1}^{N} B(a_i, h)$ . For notational simplicity we discuss only the treatment of one vertex placed in the origin,  $a_1 = (0, 0)$ . Let  $\phi_h \in C_c^{\infty}(B(0, h))$  be such that  $\phi = 1$  on B(0, h/2). Let  $\eta \in C_c^{\infty}(B(0, 1))$  be a standard mollifier, and let  $\eta_h(\cdot) = h^{-2}\eta(\cdot/h)$ . We set  $\tilde{v}_h = \phi_h(v_h*\eta_h) + (1-\phi_h)v_h$ . Then  $\|\nabla^2 \tilde{v}_h\|_{L^2(B(0,h))}^2 \leq C$ . We note that  $\nabla v_h$  is uniformly bounded, therefore  $\|\nabla v_h \otimes \nabla v_h - \nabla \tilde{v}_h \otimes \nabla \tilde{v}_h\|_{L^2(B(0,h))}^2 \leq Ch^2$ . We conclude that

$$E_h(\mathbf{u}_h, \tilde{v}_h) = \int_{\Omega} |\nabla \mathbf{u}_h + \nabla \mathbf{u}_h^T + \nabla \tilde{v}_h \otimes \nabla \tilde{v}_h - \nabla v_0 \otimes \nabla v_0|^2 + h^2 |\nabla^2 v_h|^2 \mathrm{d}x$$
  
$$\leq C h^{5/3}.$$

This proves the proposition.

## 4. PROOF OF THE LOWER BOUND

In the present section, we will prove the lower bound in Theorem 1.1. This will proceed in several steps. As before,  $v_0$  denotes the unique solution of the Monge-Ampère equation (2).

In Subsection 4.1 we recall the lower bound estimate

$$\|\nabla \mathbf{u} + \nabla \mathbf{u}^T + \nabla v \otimes \nabla v - \nabla v_0 \otimes \nabla v_0\|_{L^2(\Omega)} \gtrsim \|\mu_v - \bar{\mu}\|_{W^{-2,2}(\Omega)}$$

that has already been used in [Olb19, GO24]. This justifies our approach to look for lower bounds of the functional

$$I_h(v) = \|\mu_v - \bar{\mu}\|_{W^{-2,2}(\Omega)}^2 + h^2 \|\nabla^2 v\|_{L^2(\Omega)}^2$$

in the sequel.

We will then go on to consider sequences  $v_h$  that satisfy the upper bound in Theorem 1.1. By the results of Subsection 4.1, this implies  $I_h(v_h) \lesssim h^{4/3} \left(\log \frac{1}{h}\right)^{1/3}$ . For such a

sequence, we will first prove "suboptimal" estimates for  $||v_h - v_0||_{L^{\infty}}$ . More precisely, we will prove in Subsection 4.2 that  $||v_h - v_0||_{L^{\infty}(\Omega)} \leq h^{\gamma}$  for some fixed positive  $\gamma$ . We call these estimates suboptimal since we will prove much sharper ones later on.

In Subsection 4.3, we consider the behavior of  $v_h$  satisfying the upper bound near any one of the vertices  $a_i$ , i = 1, ..., N. By the suboptimal estimates obtained in Subsection 4.2, we know that for a suitably chosen affine function  $\tilde{\ell}^i$ ,  $v_h$  is close to the "conical" function  $\hat{v} := L_{\{v_h = \tilde{\ell}^i\} \cup \{a_i\}} v$  on  $\{v_h \leq \tilde{\ell}^i\}$ , see Figures 1 and 2. We prove a statement that allows us to upgrade the suboptimal estimates to *optimal* estimates for  $|v_h(x) - \hat{v}(x)|$ on  $\{v_h \leq \tilde{\ell}^i\}$ , see Proposition 4.8. Here and in the following, by "optimal" estimates we mean that one can construct examples in which the estimate is attained up to at most a logarithmic factor in the small parameter h.



FIGURE 1. The graphs of  $v_h$  and  $\tilde{\ell}^i$ , with  $v_h(a_i) < \tilde{\ell}^i(a_i)$ ,  $v_h(a_l) > \tilde{\ell}^k(a_l)$ for  $l \in \{1, \ldots, N\} \setminus \{i\}$ . The map  $\tilde{\ell}^i$  will be constructed in Proposition 4.11 as a slight perturbation of some other affine map  $\ell^i$ , introduced in Lemma 4.10.

In Subsection 4.4, we consider a ridge  $(i, j) \in \mathcal{R}_{v_0}$  (where we are using the notation introduced in Lemma 2.11), and the rhombus  $R_{ij} = [a_i b_{ij}^+ a_j b_{ij}^-]$ , on which the gradient of  $v_0$  has exactly two values. We use the results from the previous subsections as follows: By the suboptimal estimates from Subsection 4.2, v is close to  $v_0$  on the corners of  $R_{ij}$ . Consider the continuous function  $w_h$  that is affine on  $[a_i b_{ij}^+ a_j]$  and on  $[a_i b_{ij}^- a_j]$ , with the values of  $w_h$  at  $a_i, a_j, b_{ij}^{\pm}$  given by those of  $v_h$ . By the results of Subsection 4.3, we have that  $v_h$  is close to  $w_h$  on  $\partial R_{ij}$ , with optimal estimates, see Figures 2 and 3. Using the generalized monotonicity property of the Monge-Ampère measure (Proposition 2.6), we are able to transfer these estimates from the boundary to the bulk of  $R_{ij}$ . This crucial step is contained in Proposition 4.11.

The latter allows us to prove the lower bound in Subsection 4.5: The optimal bounds in the interior imply that there is a small transition layer parallel to the ridge of size  $\approx h^{-2/3}$  (up to logarithms in h), across which the gradient  $\nabla v_h$  has a jump of order one. This is enough to establish the lower bound.

**Remark 4.1.** The logarithmic factors in the lower bound have the following source: When estimating the  $W^{-2,2}$  norm of  $\mu_v - \bar{\mu}$  from below (in the "optimal" estimates Propositions 4.8 and 4.11), we have to work with test functions that behave approximately



FIGURE 2. The set  $E^i = \{x : v_h(x) \leq \tilde{\ell}^i(x)\}$ . The function  $\hat{v}$  is defined by requiring to be affine on each of the line segments  $[a_i x]$  with  $x \in \partial E^i$ , agreeing with  $v_h$  on the endpoints of these line segments. By Proposition 4.8,  $\|v_h - \hat{v}\|_{L^{\infty}}$  is controlled by the energy. This estimate is optimal in the sense explained in the outline of the proof at the beginning of Section 4.



FIGURE 3. Since  $b_{ij}^{\pm} \in \partial E^i \cap \partial E^j$  and  $R_{ij} = [a_i b_{ij}^{+} a_j b_{ij}^{-}]$ , the optimal estimates for  $v_h$  hold on all of  $\partial R_{ij}$ . These estimates control the distance of  $v_h$  from a continuous function on  $\partial R_{ij}$  that is affine on  $[a_i b_{ij}^{+} a_j]$  and on  $[a_i b_{ij}^{-} a_j]$ .

like  $x \mapsto |x - a_i|$ , which fail to be elements of  $W^{2,2}$ , producing logarithmic singularities after smoothing on the appropriate scale. It seems that this cannot be improved within our method of proof, since we need to work with non-negative test functions that vanish at vertices (where  $\bar{\mu}$  is concentrated) and are positive everywhere else in order to detect the deviations of v from  $v_0$ .

4.1. Lower bounds via the  $W^{-2,2}$  norm. For the reader's convenience, we state and prove a lemma that is basically identical to [Olb19, Lemma 4.4]:

**Lemma 4.2.** Let  $(\mathbf{u}, v) \in W^{1,2}(\Omega; \mathbb{R}^2) \times W^{2,2}(\Omega)$ , and as above  $\bar{\mu} = \mu_{v_0} = \sum_{i=1}^N \sigma_i \delta_{a_i}$ . Then

$$\|\mu_{v} - \bar{\mu}\|_{W^{-2,2}(\Omega)} \leq \frac{1}{2} \|\nabla \mathbf{u} + \nabla \mathbf{u}^{T} + \nabla v \otimes \nabla v - \nabla v_{0} \otimes \nabla v_{0}\|_{L^{2}(\Omega)}.$$
23

*Proof.* The Hessian determinant of a smooth function w in its very weak form is given by

$$\det \nabla^2 w = \partial_1 \partial_2 (\partial_1 w \partial_2 w) - \frac{1}{2} \partial_2^2 (\partial_1 w^2) - \frac{1}{2} \partial_1^2 (\partial_2 w^2) = -\frac{1}{2} \operatorname{curl} \operatorname{curl} \nabla w \otimes \nabla w$$

Here, we have used the notation  $\operatorname{curl}(w_1, w_2) = \partial_2 w_1 - \partial_1 w_2$ . In the equation above,  $\operatorname{curl}$ is first applied in each row of the matrix  $\nabla w \otimes \nabla w$ , and then on the components of the resulting column vector. When replacing w by  $v \in W^{2,2}(\Omega)$  the equality between the left hand side and the right hand side holds in  $W^{-2,2}(\Omega)$ . Since curl curl  $(\nabla \mathbf{w}^T + \nabla \mathbf{w}) = 0$  for every  $\mathbf{w} \in W^{1,2}(\Omega; \mathbb{R}^2)$ , we obtain

$$\det \nabla^2 v = -\frac{1}{2} \operatorname{curl} \operatorname{curl} \left( \nabla \mathbf{u} + \nabla \mathbf{u}^T + \nabla v \otimes \nabla v \right) \,.$$

again in  $W^{-2,2}(\Omega)$ . An analogous identity for  $v_0$  holds in  $W^{-2,2}(\Omega)$ ,

$$\mu_{v_0} = -\frac{1}{2} \operatorname{curl} \operatorname{curl} \left( \nabla v_0 \otimes \nabla v_0 \right) \,,$$

Thus for every  $\varphi \in W_0^{2,2}(\Omega)$ , we obtain by two integrations by parts, and the Cauchy-Schwarz inequality,

$$\int_{\Omega} \varphi d(\mu_{v} - \bar{\mu}) = -\frac{1}{2} \int_{\Omega} \left( \nabla \mathbf{u} + \nabla \mathbf{u}^{T} + \nabla v \otimes \nabla v - \nabla v_{0} \otimes \nabla v_{0} \right) : \operatorname{cof} \nabla^{2} \varphi dx$$
$$\leq \frac{1}{2} \| \nabla \mathbf{u} + \nabla \mathbf{u}^{T} + \nabla v \otimes \nabla v - \nabla v_{0} \otimes \nabla v_{0} \|_{L^{2}(\Omega)} \| \varphi \|_{W^{2,2}_{0}(\Omega)}.$$

This proves our claim.

### 4.2. Suboptimal global estimates.

**Lemma 4.3.** Let  $\Omega \subseteq \mathbb{R}^n$  be open, bounded, convex. Let  $\mu = \sum_{i=1}^N \sigma_i \delta_{a_i}, \nu = \sum_{i=1}^N \tau_i \delta_{a_i} = 0$  $\mathcal{M}_+(\Omega)$  be two non-negative discrete measures with the same support. Let  $u, v \in C^0(\overline{\Omega})$ be the respective solutions to the Dirichlet problems

$$\begin{cases} \mu_u = \mu, \mu_v = \nu & \text{ in } \Omega\\ u = v = 0 & \text{ on } \partial\Omega. \end{cases}$$
(27)

Then

$$|u - v||_{L^{\infty}} \le C(\Omega, N) \sum_{i=1}^{N} |\sigma_i^{1/n} - \tau_i^{1/n}|.$$
(28)

*Proof.* Let  $\delta = ||u - v||_{L^{\infty}} > 0$ . Without loss of generality, there is  $x \in \Omega$  with  $\delta =$ u(x) - v(x). Let  $A := (\partial^- v)^{-1} (\partial^- v(x)) \subseteq \overline{\Omega}$  be the maximal simplex containing x where v is affine. Since u - v is convex on A, its maximum is attained on the extreme points of A. By Lemmata 2.2 and 2.9, the extreme points of A are contained in  $\partial \Omega \cup \{a_1, \ldots, a_N\}$ . Hence the maximum must be attained at some vertex, so that without loss of generality  $x = a_1$ . For  $k = 0, \ldots, N$  let  $A_k = \{a_j : u(a_j) - v(a_j) \ge \frac{k\delta}{N}\}$ . Since  $A_k \supseteq A_{k+1}$  and  $A_N \supseteq \{a_1\}$ , there is  $k_0 \in \{0, \ldots, N-1\}$  with  $A_{k_0} = A_{k_0+1}$ . We will show that

$$\partial^{-}v(A_{k_0}) \supseteq B\left(\partial^{-}u(A_{k_0}), \frac{\delta}{N\operatorname{diam}\left(\Omega\right)}\right),$$
(29)

so that by the Brunn-Minkowski inequality

$$\mu_v(A_{k_0})^{1/n} \ge \mu_u(A_{k_0})^{1/n} + c(n) \frac{\delta}{N \text{diam}\,(\Omega)},$$

which after rearranging yields

$$\|u - v\|_{L^{\infty}} = \delta \le C(\Omega, N)(\mu_v(A_{k_0})^{1/n} - \mu_u(A_{k_0})^{1/n}) \le C(\Omega, N)\sum_{i=1}^N |\sigma_i^{1/n} - \tau_i^{1/n}|.$$

To show (29), let  $p \in \partial^- u(a_j)$  for some  $a_j \in A_{k_0} = A_{k_0+1}$  and  $|q| < \frac{\delta}{N \operatorname{diam}(\Omega)}$ . We estimate

$$\begin{split} (p+q) \cdot a_j - v(a_j) &\geq p \cdot a_j - u(a_j) + \frac{(k_0+1)\delta}{N} + q \cdot a_j \\ &\geq \max_{y \in \partial \Omega \cup \{a_1, \dots, a_N\} \setminus A_{k_0}} p \cdot y - u(y) + \frac{(k_0+1)\delta}{N} + q \cdot y - |q||y - a_j| \\ &\geq \max_{y \in \partial \Omega \cup \{a_1, \dots, a_N\} \setminus A_{k_0}} (p+q) \cdot y - v(y) + \frac{\delta}{N} - \frac{\delta}{N}. \end{split}$$

By Lemma 2.2,  $\max_{y \in \partial \Omega \cup \{a_1, \dots, a_N\}} (p+q) \cdot y - v(y) = \max_{y \in \Omega} (p+q) \cdot y - v(y)$ , and hence  $p+q \in \partial^- v(A_{k_0})$ , completing the proof.

In the upcoming lemma, we will use the notation

$$U_{\rho} = \{ x \in U : \operatorname{dist}(x, \partial U) > \rho \}.$$

for  $U \subseteq \mathbb{R}^2$ . The assumptions that will be made on U are chosen such that they accommodate the case  $U = \Omega \setminus \{a_1, \ldots, a_N\}$ , where  $\Omega$  and  $\{a_1, \ldots, a_N\}$  are as in Theorem 1.1.

**Lemma 4.4.** Let  $U \subseteq \mathbb{R}^2$  be open, bounded, such that  $\overline{U}$  is convex and  $U_{\rho}$  has Lipschitz boundary, with

$$\mathcal{H}^{n-1}(\partial U_{\rho}) \leq C(U)$$
  
Furthermore let  $\alpha \in (0,1]$ , and  $u \in C^{0,\alpha}(\overline{U})$  convex. Then
$$\|u - L_{\partial U}u\|_{L^{\infty}(U)} \leq C(U,\alpha)\|\mu_{u}\|_{W^{-2,2}(U)}^{\frac{4\alpha}{n(4\alpha+3)}}\|u\|_{C^{0,\alpha}}^{\frac{3}{4\alpha+3}}.$$
(30)

Proof. Let  $\rho > 0$  to be chosen later. Pick a test function  $\phi_{\rho} \in C_c^{\infty}(U)$  with  $\phi_{\rho} = 1$  in  $U_{\rho}$  and  $\|\phi_{\rho}\|_{W^{2,2}} \leq C(U)\rho^{-3/2}$ . This can be achieved by choosing  $\phi_{\rho} = \eta_{\rho/3} * \mathbb{1}_{U_{\rho}}$ , where  $\eta_{\varepsilon} = \varepsilon^{-2}\eta(\cdot/\varepsilon)$  and  $\eta \in C_c^{\infty}(\mathbb{R}^n)$  is a standard symmetric mollifier with  $\operatorname{supp} \eta \subseteq B(0,1)$ . Then on the one hand,

$$\mu_u(U_\rho) \le \int \phi_\rho \mathrm{d}\mu_u \le \|\mu_u\|_{W^{-2,2}(U)} \|\phi_\rho\|_{W^{2,2}(U)} \le C(U) \|\mu_u\|_{W^{-2,2}(U)} \rho^{-3/2}.$$

By the Alexandrov maximum principle, Lemma 2.8, we have

$$\|u - L_{\partial U_{\rho}} u\|_{L^{\infty}(U_{\rho})} \le C(U)(\mu_u(U_{\rho}))^{1/n} \le C(U) \|\mu_u\|_{W^{-2,2}}^{1/n} \rho^{-3/4}$$

On the other hand, we note that  $L_{\partial U \cap \partial U_{\rho}} u = L_{\partial U_{\rho}}$  on  $U_{\rho}$ , and

 $\|L_{\partial U}u - L_{\partial U \cap \partial U_{\rho}}u\|_{L^{\infty}(U)} = \|L_{\partial U}u - u\|_{L^{\infty}(\partial U_{\rho})} \le C(U, \alpha)\rho^{\alpha}\|u\|_{C^{0,\alpha}}.$ 

Similarly,

for all  $\rho > 0$ .

$$\|u - L_{\partial U \cap \partial U_{\rho}} u\|_{L^{\infty}(U \setminus U_{\rho})} \le C(U, \alpha) \rho^{\alpha} \|u\|_{C^{0, \alpha}}$$

Putting everything together via the triangle inequality

$$\begin{aligned} \|u - L_{\partial U}u\|_{L^{\infty}(U)} &\leq \|u - L_{\partial U\rho}u\|_{L^{\infty}(U\rho)} + \|u - L_{\partial U\cap\partial U\rho}u\|_{L^{\infty}(U\setminus U\rho)} \\ &+ \|L_{\partial U}u - L_{\partial U\cap\partial U\rho}u\|_{L^{\infty}(U)} \\ &25 \end{aligned}$$

and choosing the optimal  $\rho$ ,

$$\rho = \left(\frac{\|\mu_u\|_{W^{-2,2}}^{1/n}}{\|u\|_{C^{0,\alpha}}}\right)^{\frac{1}{3/4+\alpha}}$$

yields (30).

From now on we assume that  $v_h \in W^{2,2}(\Omega)$  satisfies the upper bound in Theorem 1.1, in particular (using Lemma 4.2)

$$I_h(v_h) \le C_1 h^{4/3} \left( \log \frac{1}{h} \right)^{1/3}$$
 (31)

**Lemma 4.5.** Let  $v_0 \in C^0(\overline{\Omega})$  be a solution of (2), and  $\beta \in (\frac{3}{4}, 1)$ . Then there exists  $C(\Omega, \beta, C_1, a_1, \ldots, a_N) > 0$  and  $h_0(\Omega, \beta, C_1, a_1, \ldots, a_N) > 0$  such that for all  $h < h_0$ , and all  $v_h \in W^{2,2}(\Omega)$  with  $v_h = 0$  on  $\partial\Omega$  satisfying (31), we have that

$$\|v_h - L_{\partial \Omega \cup \{a_1, \dots, a_N\}} v_h\|_{L^{\infty}} \le Ch^{\gamma}$$

where

$$\gamma = \frac{4\beta - 3}{3(4\beta + 3)} > 0$$

Proof. By the continuous embedding  $W^{2,2}(\Omega) \hookrightarrow C^{0,\alpha}(\Omega)$ , we have that  $||v_h||_{C^{0,\alpha}(\Omega)} \leq Ch^{-1/3} \left(\log \frac{1}{h}\right)^{1/6}$ . Thus we may apply Lemma 4.4 with  $U = \Omega \setminus \{a_1, \ldots, a_N\}$  and (for example)  $\alpha = \frac{1+\beta}{2}$ , to obtain

$$\begin{aligned} \|v_h - L_{\Omega \cup \{a_1, \dots, a_N\}} v_h\|_{L^{\infty}} \\ &\leq C(\Omega, a_1, \dots, a_N) \left( h^{2/3} (\log h^{-1})^{1/6} \right)^{\frac{2\alpha}{4\alpha + 3}} \left( h^{-1/3} (\log h^{-1})^{1/6} \right)^{\frac{3}{4\alpha + 3}} \\ &\leq C h^{\frac{4\alpha - 3}{3(4\alpha + 3)}} (\log h^{-1})^{\frac{2\alpha + 3}{6(4\alpha + 3)}} . \end{aligned}$$

The statement of the present lemma follows from choosing  $h_0$  and C appropriately. **Lemma 4.6.** Assume that  $v_h \in W^{2,2}(\Omega)$  with  $v_h = 0$  on  $\partial\Omega$  satisfying (31). Then for i = 1, ..., N,

$$|v_h(a_i)| \leq C$$
.

*Proof.* For  $\rho > 0$ , let  $\Omega_{\rho} := \{x \in \Omega : \operatorname{dist}(x, \partial \Omega) > \rho\}$ . We estimate  $|v_h(a_i)| \le |L_{\partial \Omega_{\rho}} v_h(a_i)| + |L_{\partial \Omega_{\rho}} v_h(a_i) - v_h(a_i)|$ 

$$|L_{\partial\Omega_{\rho}}v_{h}(a_{i})| \leq |L_{\partial\Omega_{\rho}}v_{h}(a_{i})| + |L_{\partial\Omega_{\rho}}v_{h}(a_{i}) - v_{h}(a_{i})|$$
$$\leq \sup_{x \in \partial\Omega_{\rho}} |v_{h}| + \sup_{x \in \Omega_{\rho}} |L_{\partial\Omega_{\rho}}v_{h}(x) - v_{h}(x)|$$

Using again the fact that  $\|v_h\|_{C^{0,\alpha}(\Omega)} \leq C(\alpha)h^{-1/3} \left(\log \frac{1}{h}\right)^{1/6}$  for every  $\alpha \in (0,1)$  by the continuous embedding  $W^{2,2}(\Omega) \hookrightarrow C^{0,\alpha}(\Omega)$ , we obtain

$$\sup_{x \in \partial \Omega_{\rho}} |v_h| \le C \rho^{\alpha} h^{-1/3} \left( \log \frac{1}{h} \right)^{1/6}.$$

By Lemma 2.8,

$$\sup_{x \in \Omega_{\rho}} \left| L_{\partial \Omega_{\rho}} v_h(x) - v_h(x) \right| \le C \mu_{v_h}(\Omega_{\rho})^{1/2}$$

Let  $\phi_{\rho} \in C_c^{\infty}(\Omega)$  such that  $\phi_{\rho} = 1$  on  $\Omega_{\rho}$  and such that  $\|\phi_{\rho}\|_{W_0^{2,2}(\Omega)} \leq C\rho^{-3/2}$ . We have that

$$\begin{aligned} \mu_{v_h}(\Omega_{\rho}) &\leq \int \phi_{\rho} d\mu_{v_h} \\ &\leq \int \phi_{\rho} d\bar{\mu} + \|\mu_{v_h} - \bar{\mu}\|_{W^{-2,2}(\Omega)} \|\phi_{\rho}\|_{W^{2,2}_{0}(\Omega)} \\ &\leq \bar{\mu}(\Omega) + Ch^{2/3} \left(\log \frac{1}{h}\right)^{1/6} \rho^{-3/2} \end{aligned}$$

and inserting this last estimates in the previous ones, we obtain

$$|v_h(a_i)| \le C(\bar{\mu}, \Omega) + C(\alpha)\rho^{\alpha}h^{-1/3} \left(\log\frac{1}{h}\right)^{1/6} + Ch^{2/3} \left(\log\frac{1}{h}\right)^{1/6}\rho^{-3/2}.$$

Choosing  $\alpha = 1 - \varepsilon$  for some small  $\varepsilon > 0$ , and  $\rho = h^{1/3+\varepsilon}$ , we obtain the claim of the present lemma.

**Proposition 4.7.** Let  $\Omega \subseteq \mathbb{R}^2$  and  $\bar{\mu} = \sum_{i=1}^N \sigma_i \delta_{a_i}$  be as in the statement of Theorem 1.1,  $v_h$  a sequence satisfying the upper bound in Theorem 1.1,  $\beta \in (\frac{3}{4}, 1)$  and  $h_0, C, \gamma$  chosen as in Lemma 4.5. Then

$$\|v_h - v_0\|_{L^{\infty}} \le Ch^{\gamma} \quad \text{for } h < h_0.$$

*Proof.* Let  $r = \frac{1}{4} \min_{i,j} \min(|a_i - a_j|, \operatorname{dist}(a_i, \partial \Omega)) > 0$ . Find for each  $i = 1, \ldots, N$  two test functions  $\phi_i, \Phi_i \in C_c^{\infty}(\Omega)$  with  $\phi_i(a_j) = \Phi_i(a_j) = \delta_{ij}$  and  $0 \le \phi_i \le \mathbb{1}_{B(a_i,r)} \le \Phi_i$ . Then denoting by  $\langle \cdot, \cdot \rangle$  the dual pairing  $W^{-2,2} \times W_0^{2,2} \to \mathbb{R}$ ,  $|\langle \mu_{v_h} - \bar{\mu}, \phi_i \rangle| + |\langle \mu_{v_h} - \bar{\mu}, \Phi_i \rangle| \lesssim h^{2/3} \log \frac{1}{h}$  by the assumed estimates on  $I_h(v_h)$ , and hence

$$\sigma_{i} - Ch^{2/3} \left( \log \frac{1}{h} \right)^{1/6} \leq \langle \mu_{v_{h}}, \phi_{i} \rangle$$

$$\leq \mu_{v_{h}} (B(a_{i}, r))$$

$$\leq \langle \mu_{v_{h}}, \Phi_{i} \rangle$$

$$\leq \sigma_{i} + Ch^{2/3} \left( \log \frac{1}{h} \right)^{1/6}.$$
(32)

Let us write

$$v_h := L_{\partial \Omega \cup \{a_1, \dots, a_N\}} v_h \,.$$

For every  $i \in \{1, ..., N\}$ ,  $\partial^- w_h(a_i)$  is convex. Furthermore  $|v_h(a_i)| \leq C$  by Lemma 4.6 and hence

$$\operatorname{diam} \partial^- w_h(a_i) \le C \tag{33}$$

by the definition of  $w_h$ . Combining (33) with the convexity of  $\partial^- w_h(a_i)$ , we get that there exists  $\varepsilon_0$  such that

$$\mathcal{L}^{2}\left(\left\{p \in \mathbb{R}^{2} : \operatorname{dist}(p, \partial(\partial^{-}w_{h}(a_{i})) < \varepsilon\right\}\right)\right) \leq C\varepsilon$$
(34)

for every  $\varepsilon < \varepsilon_0$ .

Now let  $p \in \partial^- w_h(a_i)$  with  $\operatorname{dist}(p, \partial(\partial^- w_h(a_i))) \ge \frac{\|v_h - w_h\|_{L^{\infty}}}{r}$ . Then for any  $x \in \overline{\Omega}$ with  $|x - a_i| > r$  we may choose  $p' \in \partial B\left(p, \frac{\|v_h - w_h\|_{L^{\infty}}}{r}\right)$  such that  $(p - p') \cdot (x - a_i) = \frac{27}{27}$ 

$$\begin{aligned} -|p - p'||x - a_i|. \text{ Thus} \\ p \cdot x - v_h(x) &\leq p \cdot x - w_h(x) + \|v_h - w_h\|_{L^{\infty}} \\ &= p' \cdot x - w_h(x) + (p - p') \cdot x + \|v_h - w_h\|_{L^{\infty}} \\ &\leq p' \cdot a_i - w_h(a_i) + (p - p') \cdot x + \|v_h - w_h\|_{L^{\infty}} \\ &\leq p \cdot a_i - w_h(a_i) - |p - p'||x - a_i| + \|v_h - w_h\|_{L^{\infty}} \\ &$$

which implies that

$$p \notin \partial^- v_h(x)$$
,

and hence

$$\left\{p \in \partial^- w_h(a_i) : \operatorname{dist}(p, \partial(\partial^- w_h(a_i))) > \frac{\|v_h - w_h\|_{L^{\infty}}}{r}\right\} \subseteq \partial^- v_h(B(a_i, r)).$$

From this inclusion we now obtain using (32), (34) and Lemma 4.5,

$$\mu_{w_h}(\{a_i\}) = \mathcal{L}^2(\partial^- w_h(a_i))$$
  

$$\leq \mathcal{L}^2(\partial^- v_h(B(a_i, r)) + C \|v_h - w_h\|_{L^{\infty}}$$
  

$$\leq \sigma_i + Ch^{2/3} \left(\log \frac{1}{h}\right)^{1/6} + Ch^{\gamma}.$$

Next we prove

$$\partial^{-}v_{h}(B(a_{i},r)) \subseteq \left\{ p : \operatorname{dist}(p,\partial^{-}w_{h}(a_{i})) < \frac{4\|v_{h} - w_{h}\|_{L^{\infty}}}{r} \right\}.$$
(35)

To show (35), observe that  $p \in \partial^- w_h(a_i)$  if and only if there exists an affine function A with  $\nabla A = p$ ,  $w_h \ge A$  on  $B(a_i, r)$  and  $w_h = A$  on at least one point in  $B(a_i, r)$ . For any  $x_0 \in B(a, r_i)$  and  $p \in \partial^- v_h(x_0)$ , we have

$$v_h(x_0) + p \cdot (x - x_0) \le v_h(x) \le w_h(x) \quad \text{for } x \in B(a_i, 2r)$$

Assume  $v_h(x_0) + p \cdot (a_i - x_0) < w_h(a_i) - 4 \|v_h - w_h\|_{L^{\infty}}$ . Then with  $x_1 = 2x_0 - a_i$  and the fact that  $w_h$  is affine on the line segment  $[a_1x_1]$ , we get  $v_h(x_1) \ge v_h(x_0) + p \cdot (x_1 - x_0) \ge w_h(x_1) + 2 \|v_h - w_h\|_{L^{\infty}}$ , which is a contradiction. Hence

$$w_h(a_i) + p \cdot (x - a_i) \le w_h(x) + 4 ||v_h - w_h||_{L^{\infty}}$$
 for  $x \in B(a_i, r)$ .

Thus one may find  $p' \in B\left(p, \frac{4\|v_h - w_h\|_{L^{\infty}}}{r}\right)$  that satisfies

$$w_h(a_i) + p' \cdot (x - a_i) \le w_h(x)$$
 for  $x \in B(a_i, r)$ ,

which implies  $p' \in \partial^- w_h(a_i)$  and hence (35). From (35) and (32), we obtain

$$\mu_{w_h}(\{a_i\}) = \mathcal{L}^2(\partial^- w_h(a_i))$$
  

$$\geq \mathcal{L}^2(\partial^- v_h(B(a_i, r)) - C \|v_h - w_h\|_{L^{\infty}}$$
  

$$\geq \sigma_i - Ch^{2/3} \left(\log\frac{1}{h}\right)^{1/6} - Ch^{\gamma}.$$

Hence we have proved

$$|\mu_{w_h}(\{a_i\}) - \sigma_i| \le Ch^{\gamma} \quad \text{for } i = 1, \dots, N.$$
28

By Lemma 4.3, we obtain  $||w_h - v_0||_{L^{\infty}} \leq Ch^{\gamma}$ . Thus

$$||v_h - v_0||_{L^{\infty}} \le ||w_h - v_0||_{L^{\infty}} + ||w_h - v_h||_{L^{\infty}} \le Ch^{\gamma}.$$

4.3. Estimates for nearly conical configurations. In this subsection we consider a closed convex set E with  $0 \in E$ , and  $|x| \simeq 1$  for  $x \in \partial E$ , and a convex function  $v \in C^0(\bar{E})$  with v = 0 on  $\partial E$ . We are going to define a weighted integral of the Hessian determinant that can serve as a replacement for the energy  $\|\mu_v - \delta_0\|_{W^{-2,2}}$ ,

$$F(v) = \int_{E} |x| \mathrm{d}\mu_{v} \,. \tag{36}$$

**Proposition 4.8.** Let  $E \subseteq \mathbb{R}^n$  be a convex open set with  $0 \in E$ , with  $1/R \leq |x| \leq R$  for  $x \in \partial E$ , and  $U \subseteq \mathbb{R}^n$  open with  $E \subset U$ . Let  $v \in C^0(U)$  be convex with v = 0 on  $\partial E$  and  $\mu_v(\partial E) = 0$ , and  $\hat{v} : E \to \mathbb{R}$  defined by

$$\hat{v} = L_{\partial E \cup \{0\}} v \,.$$

Then

$$\hat{v}(x) - v(x) \le C(R, n) \min(|x|^{(n-1)/n} F(v)^{1/n}, |v(0)||x|) \quad \text{for all } x \in E.$$
(37)

*Proof.* First we assume that the restriction of v to E is in  $W^{2,\infty}$ ,  $v \in W^{2,\infty}(E)$ . We may write by a change of variables

$$F(v) = \int_{E} |x| \det \nabla^{2} v dx$$
  
= 
$$\int_{\partial^{-} v(E)} |(\nabla v)^{-1}(p)| dp$$
  
$$\geq \int_{\partial^{-} \hat{v}(0)} |(\nabla v)^{-1}(p)| dp.$$
 (38)

In the last step we have used the monotonicity of the subdifferential (see [Fig17, Lemma 2.7] or Proposition 2.6 with  $\phi = 1$ ). The point  $(\nabla v)^{-1}(p) \in E$  can be written for almost every  $p \in \partial^{-}v(E)$  as

$$(\nabla v)^{-1}(p) = \operatorname*{arg\,max}_{x \in E} p \cdot x - v(x). \tag{39}$$

Let

$$x_0 \in \operatorname*{arg\,max}_{x \in E \setminus \{0\}} \frac{\hat{v}(x) - v(x)}{|x|^{(n-1)/n}} \tag{40}$$

and  $y_0 := \hat{v}(x_0) - v(x_0)$ . (A maximizer exists since  $v, \hat{v}$  are Lipschitz at 0 and coincide on  $\partial E$ .) Let  $p_0 \in \partial^- \hat{v}(x_0)$ . We claim that

$$|(\nabla v)^{-1}(p)| \ge \frac{1}{2}|x_0| \quad \text{for all } p \in \partial^- \hat{v}(0) \cap B\left(p_0, \frac{y_0}{2|x_0|}\right).$$
(41)

Since  $\mathcal{L}^n(\partial^- \hat{v}(0) \cap B(p_0, r)) \ge cr^n$  for all  $r \le |v(0)|$ , and  $r = \frac{y_0}{2|x_0|} \le C|v(0)|$  by the Lipschitz continuity of  $v, \hat{v}$ , we obtain

$$\begin{split} F(v) &\geq \int_{\partial^{-}\hat{v}(0)\cap B(p_{0},\frac{y_{0}}{2|x_{0}|})} |(\nabla v)^{-1}(p)| \,\mathrm{d}p \\ &\geq \frac{|x_{0}|}{2} c \left(\frac{y_{0}}{|x_{0}|}\right)^{n} \\ &= c \frac{y_{0}^{n}}{|x_{0}|^{n-1}} \\ &= c \max_{x \in E} \frac{(\hat{v}(x) - v(x))^{n}}{|x|^{n-1}}. \end{split}$$

Reordering yields

$$\hat{v}(x) - v(x) \le C|x|^{(n-1)/n} F(v)^{1/n}$$
 for all  $x \in E$ .

The inequality  $\hat{v}(x) - v(x) \leq C|v(0)||x|$  follows from the uniform Lipschitz bounds near 0. Now to show (41):

Let  $p \in \partial^- \hat{v}(0)$ . Then  $\hat{v}(x) \ge p \cdot x$  for all  $x \in E$  and for  $|x| < \frac{|x_0|}{4}$  we have

$$p \cdot x - v(x) \le \hat{v}(x) - v(x) \le \frac{|x|^{1/2}}{|x_0|^{1/2}} y_0 < \frac{y_0}{2}.$$
 (42)

On the other hand for  $|p - p_0| \le \frac{y_0}{2|x_0|}$  we have since  $p_0 \cdot x_0 = \hat{v}(x_0)$ 

$$p \cdot x_0 - v(x_0) = p_0 \cdot x_0 - v(x_0) + (p - p_0) \cdot x_0 \ge y_0 - |p - p_0| |x_0| \ge \frac{y_0}{2}.$$
 (43)

In particular  $|(\nabla v)^{-1}(p)| \ge \frac{|x_0|}{4}$ . This completes the proof for  $v \in W^{2,\infty}(E)$ .

In the general case, consider the approximations  $v_{\varepsilon} := P_{\varepsilon}v|_U \in W^{2,\infty}(U)$  of v given obtained by applying the smoothing operation  $P_{\varepsilon}$  from Lemma 2.3. Let  $E_{\varepsilon} := \{x \in U : v_{\varepsilon} < 0\}$ . Then  $\mu_{v_{\varepsilon}} \stackrel{*}{\rightharpoonup} \mu_{v}$  weakly-\* as measures on  $\overline{E}$  and  $\mathbb{1}_{E_{\varepsilon}} \to \mathbb{1}_{E}$  in  $L^{1}(U)$ . Using the fact  $\mu_{v}(\partial E) = 0$ , we obtain

$$\int_{E_{\varepsilon}} |x| \mathrm{d}\mu_{v_{\varepsilon}} \to \int_{E} |x| \mathrm{d}\mu_{v}$$
$$\hat{v}_{\varepsilon}(x) - v_{\varepsilon}(x) \to \hat{v}(x) - v(x) \quad \text{for every } x \in E.$$

This finishes the proof.

**Remark 4.9.** The scaling exponent  $\frac{1}{n}$  with which the energy enters the right hand side in (37) is optimal in the sense that one may construct a family of examples that satisfies this scaling. To alleviate the notation, we assume n = 2, but the general case can be treated in analogous fashion.

Consider 
$$E = [-1, 1]^2$$
, and for  $0 < R < H < 1$ , define the piecewise affine function

$$v_{R,H}(x) = \{\sup w(x) : w \text{ convex }, w|_{\partial E} = 0, w(0) = -1, w(Re_1) = R - H\}.$$

We observe that with  $\hat{v}_{R,H} = L_{\partial E \cup \{0\}} v_{R,H}$  defined as in the proposition, we have that  $\hat{v}_{R,H} = v_{0,0} = \max(|x_1|, |x_2|)$ . We have that

$$F(v_{R,H}) = \int |x| \mathrm{d}\mu_{v_{R,H}}(x) = R\mathcal{L}^2(\partial^- v_{R,H}(Re_1)),$$
  
30

and calculate

$$\partial^{-}v_{R,H}(Re_{1}) = \operatorname{conv}\left(\left(1 - \frac{H}{R}, \frac{H}{R}\right), \left(1 - \frac{H}{R}, -\frac{H}{R}\right), \left(\frac{1 + H}{1 - R}, 0\right)\right),$$
$$F(v_{R,H}) \ge R\left(\frac{H}{R}\right)^{2} = \frac{H^{2}}{R},$$

and of course

so that

$$\hat{v}_{R,H}(Re_1) - v_{R,H}(Re_1) = H \ge \sqrt{F(v_{R,H})R}$$

This proves our claim.

4.4. Quantitative behavior near ridges. Before demonstrating Proposition 4.11 below, which is pivotal in obtaining control over  $v_h$  in the vicinity of the ridges, we establish some helpful notation in the forthcoming lemma. This lemma introduces affine maps  $\ell^k$  $(\ell_{ij}^k$  in the statement of the lemma). The maps  $\tilde{\ell}^k$  mentioned at the outset of the present section are perturbations of the maps  $\ell^k$ , see (47) below. The maps  $\tilde{\ell}^k$  will act as auxiliary tools facilitating the utilization of Proposition 4.8 in proximity to the vertices.

**Lemma 4.10.** Let  $v_0$  be the solution of (2). Then the choice of points  $b_{ij}^+, b_{ij}^- \in \Omega$  for  $(i, j) \in \mathcal{R}_{v_0}$  from Lemma 2.11 may be modified in a way such that, additionally to the properties claimed in that lemma, there exist affine functions  $\ell_{ij}^i, \ell_{ij}^j$  for  $(i, j) \in \mathcal{R}_{v_0}$  with the following properties (see Figure 1 for an illustration):

(i) For every  $(i, j) \in \mathcal{R}_{v_0}$ ,

$$v_{0}(a_{k}) < \ell_{ij}^{k}(a_{k}) \quad for \ k \in \{i, j\}$$
  
$$v_{0}(a_{l}) > \ell_{ij}^{k}(a_{l}) \quad for \ k \in \{i, j\}, l \in \{1, \dots, N\} \setminus \{k\}$$
  
$$\ell_{ij}^{k}(b_{ij}^{l}) = v_{0}(x) \quad for \ k \in \{i, j\}, l \in \{\pm\}.$$

(ii) For every  $(i, j) \in \mathcal{R}_{v_0}$ , with

$$K_{ij} := \operatorname{conv}\left(R_{ij} \cup \{x : v_0(x) \le \ell_{ij}^i\} \cup \{x : v_0(x) \le \ell_{ij}^j\}\right)$$
(44)

we have that

$$a_l \notin K_{ij}$$
 for  $l \notin \{i, j\}$ .

*Proof.* Fix  $(i, j) \in \mathcal{R}_{v_0}$ . We initially assume that  $b_{ij}^{\pm}$  are as in the statement of Lemma 2.11. By an affine change of variables, we may assume  $[a_i a_j] = [-L, L] \times \{0\}, \ b_{ij}^{\pm} = (0, \pm \beta)$  for some  $L, \beta > 0$  and

$$\nabla v_0 = p^+ \text{ on } [a_i b_{ij}^+ a_j]$$
  
$$\nabla v_0 = p^- \text{ on } [a_i b_{ij}^- a_j]$$

with  $p^+ \neq p^-$ . Clearly  $\bar{p} := \frac{1}{2}(p^+ + p^-) \in \partial^- v_0(a_i) \cap \partial^- v_0(a_j)$ . Furthermore, the affine function

$$\ell_{\bar{p},d}(x) = v(a_i) + d + (x - a_i) \cdot \bar{p} = v(a_j) + (x - a_j) \cdot \bar{p} + d$$

satisfies, for d > 0 small enough,

$$\begin{aligned} \ell_{\bar{p},d}(a_l) &= v(a_l) & \text{ for } l \in \{i,j\}, \\ \ell_{\bar{p},d}(a_l) &< v(a_l) & \text{ for } l \in \{1,\dots,N\} \setminus \{i,j\}. \end{aligned}$$

Without loss of generality, for  $\alpha > 0$  small enough, we may assume that

 $p_i(\alpha) := \bar{p} + \alpha (p^+ - p^-)^\perp \in \left(\partial^- v_0(a_i)\right)^\circ,$ and  $p_i(\alpha) := \bar{p} - \alpha (p^+ - p^-)^\perp \in \left(\partial^- v_0(a_j)\right)^\circ.$ 

Now define  $d(\alpha) > 0$  by requiring that the affine functions

$$\ell_{p_i(\alpha),d(\alpha)} := v_0(a_i) + d(\alpha) + p_i(\alpha) \cdot (x - a_i)$$
  
$$\ell_{p_j(\alpha),d(\alpha)} := v_0(a_j) + d(\alpha) + p_j(\alpha) \cdot (x - a_j)$$

satisfy

$$\ell_{p_i(\alpha),d(\alpha)} \left( (+L/2,0) \right) = v_0 \left( (+L/2,0) \right) \\ \ell_{p_i(\alpha),d(\alpha)} \left( (-L/2,0) \right) = v_0 \left( (-L/2,0) \right) \,.$$

(Note that  $(L/2, 0) = \frac{3}{4}a_j + \frac{1}{4}a_i$ ,  $(-L/2, 0) = \frac{3}{4}a_j + \frac{1}{4}a_i$ .) For  $\alpha$  small enough and  $k \in \{i, j\}$ , this implies

$$\ell_{p_k(\alpha),d(\alpha)}(a_k) > v_0(a_k),$$
  

$$\ell_{p_k(\alpha),d(\alpha)}(a_l) < v_0(a_l) \quad \text{for } l \in \{1,\dots,N\} \setminus \{k\}.$$

Moreover, it implies that there exists  $b(\alpha) > 0$ , b continuous with  $\lim_{\alpha \to 0} b(\alpha) = 0$ , such that

$$\ell_{p_i(\alpha),d(\alpha)}\left((0,\pm b(\alpha))\right) = v_0\left((0,\pm b(\alpha))\right) = \ell_{p_j(\alpha),d(\alpha)}\left((0,\pm b(\alpha))\right) \,.$$

Choosing some small positive  $\alpha$  and replacing  $b_{ij}^{\pm}$  by  $(0, \pm b(\alpha))$  completes the construction, setting  $\ell_{ij}^k = \ell_{p_k(\alpha), d(\alpha)}$  for  $k \in \{i, j\}$ .

In the statement of the following proposition, we assume that  $\Omega \subseteq \mathbb{R}^2$  and  $\bar{\mu} = \sum_{i=1}^{N} \sigma_i \delta_{a_i}$  are as in Theorem 1.1,  $v_0$  is the solution of (2), and  $(i, j) \in \mathcal{R}_{v_0}$ . Define  $U_{ij} \subseteq \Omega$  to be an open convex neighborhood of  $K_{ij}$  defined in (44) such that

$$a_l \notin \overline{U_{ij}}$$
 for  $l \in \{1, \ldots, N\} \setminus \{i, j\}$ .

**Proposition 4.11.** There exist constants  $\varepsilon_0, C > 0$  such that for every convex  $v \in C^0(U_{ij})$ with  $\|v - v_0\|_{L^{\infty}(U_{ij})} \leq \varepsilon_0$ ,

$$|v(x) - L_{\{a_j, b_{ij}^+, a_i, b_{ij}^-\}} v(x)| \le C\sqrt{F(v)} \sqrt{F(v)} + \operatorname{dist}(x, [a_i a_j]) \quad \text{for } x \in R_{ij},$$
(45)

where  $F(v) = \int_{U_{ij}} \operatorname{dist}(\cdot, \{a_i, a_j\}) d\mu_v$ .

*Proof.* Without loss of generality we assume that  $a_j = -a_i = Le_1$ , L > 0. Let  $\ell_{ij}^i, \ell_{ij}^j$  be the affine functions introduced in Lemma 4.10. We will write  $b^{\pm} \equiv b_{ij}^{\pm}, U \equiv U_{ij}, R \equiv R_{ij}, \ell_{ij}^k \equiv \ell_{ij}^k$ .

Step 1: Conical structure of  $v_0$ . For  $k \in \{i, j\}$ , let

$$E_0^k = \{x : v_0(x) \le \ell^k(x)\}.$$

By the properties of  $\ell^i, \ell^j$  stated in Lemma 4.10 we have that

$$a_k \in E_0^k, \quad \operatorname{dist}(a_k, \partial E_0^k) \gtrsim 1, \quad \operatorname{dist}(a_k, E_0^l) \gtrsim 1$$
  
$$\operatorname{dist}(E_0^k, \partial U) \gtrsim 1, \quad \operatorname{dist}(\partial E_0^i \cap \partial E_0^j, [a_i a_j]) \gtrsim 1, \quad \mathcal{L}^2(E_0^i, E_0^j) \gtrsim 1.$$
(46)

The implicit constants in the above inequalities depend only on  $\Omega, U$  and  $\bar{\mu}$ .

Step 2: Near-conical structure of v.

For  $k \in \{i, j\}$ , define the affine functions  $\tilde{\ell}^i, \tilde{\ell}^j$  by requiring

$$\tilde{\ell}^{k}(b^{\pm}) = v(b^{\pm})$$

$$\tilde{\ell}^{k}(a_{k}) = \ell^{k}(a_{k})$$
(47)

for  $k \in \{i, j\}$ . Note that this choice implies  $\|\tilde{\ell}^k - \ell^k\|_{L^{\infty}} \leq C\varepsilon_0$ . Hence, defining convex sets  $E^k \subseteq \Omega$  through

$$E^k := \left\{ x \in \Omega \, : \, v(x) \le \tilde{\ell}^k(x) \right\},\,$$

we obtain that (46) also holds with  $E_0^k$  replaced by  $E^k$  provided that  $\varepsilon_0$  is small enough, the choice depending only on  $v_0$ , and hence on  $\Omega, \bar{\mu}$ . We now apply Proposition 4.8 to  $v - \tilde{\ell}^k$  in the two sets  $E^k$ ,  $k \in \{i, j\}$ , where for the application of that proposition we translate  $a_k$  into the origin. This yields two convex functions  $\hat{g}_k : E^k \to \mathbb{R}$  with  $\hat{g}_k = 0$  on  $\partial E^k$  and  $\hat{g}_k(a_k) = v(a_k) - \tilde{\ell}^k(a_k), k \in \{i, j\}$ . We set

$$\hat{v}_k = \hat{g}_k + \ell^k \,.$$

By Proposition 4.8 we have

$$|v(x) - \hat{v}_k(x)| = |(v - \tilde{\ell}^k) - \hat{g}_k|$$
  
$$\leq C \sqrt{\int_{E^k} |x' - a_k| \mathrm{d}\mu_v(x')} \sqrt{|x - a_k|} \quad \text{for } x \in E^k.$$

Using  $|\cdot -a_k| \leq C \operatorname{dist}(\cdot, \{a_i, a_j\})$  on  $E^k$ , we deduce

$$|v(x) - \hat{v}_k(x)| \le C\sqrt{F(v)}\sqrt{|x - a_k|} \quad \text{for } x \in E^k.$$
(48)

We define the function

$$v := L_{\{a_j, b_+, a_i, b_-\}} v$$

which is affine on  $[a_i b_+ a_i]$  and on  $[a_i b_- a_i]$ . For this choice of w, (48) implies

$$|v(x) - w(x)| \le C\sqrt{F(v)}\sqrt{|x_2|} \quad \text{for all } x \in \partial R,$$
(49)

since  $w = \hat{v}_i$  and  $|x - a_i| \leq |x_2|$  on  $[a_i b_{ij}^+] \cup [a_i b_{ij}^-]$  with analogous estimates with the roles of i and j reversed.

Step 3: Boundary comparison function. We will extend a weaker version of (49) to the interior of R, losing some control near the ridge  $[a_i a_j] \subseteq \{x_2 = 0\}$ . We define the comparison function

$$w_0(x) := w(x) - C\sqrt{F(v)}\sqrt{\alpha F(v) + |x_2|},$$
(50)

where  $\alpha > 0$  is chosen large enough so that  $w_0 : \mathbb{R}^2 \to \mathbb{R}$  is convex with

$$|[\partial_2 w_0]| \ge |[\partial_2 w]|/2$$
 on  $\{x_2 = 0\}$ ,

where  $[\partial_2 w_0]$  denotes the jump of  $\partial_2 w_0$  at  $\{x : x_2 = 0\}$ . We note that such a choice of  $\alpha$  is possible and depends only on  $v_0$ . We also note that by (49),

$$w_0 \le v \le w \text{ on } \partial R.$$
  
33

Of course,  $w_0(x_1, x_2)$  only depends on  $x_2$ . For later reference, we note the following inequality for arguments  $x_2, y_2$  with opposite signs, that follows from the convexity of  $w_0$ ,  $w_0(t, w_2) = \lim_{t \to 0} (w_0(t, w_2) + \partial_x w_2(t, w_2)) = w_0(t, w_2) = w_0(t, w_2)$ 

$$w_{0}(t, y_{2}) - \lim_{x_{2} \downarrow 0} (w_{0}(t, x_{2}) + \partial_{2}w_{0}(t, x_{2})(y_{2} - x_{2})) = w_{0}(t, y_{2}) - w_{0}(t, 0) - y_{2}\partial_{2}w_{0}(t, 0+)$$

$$\geq |[\partial_{2}w_{0}]||y_{2}| \quad \text{for } y_{2} < 0,$$
(51)

with an obvious analogue for  $y_2 > 0$  and the limit  $x_2 \uparrow 0$ .

Step 4: Upper bound for  $w_0 - v$ . We now show that for all  $x = (x_1, x_2) \in R$ , we have

$$v(x) \ge w_0(x) - C\sqrt{F(v)}\sqrt{F(v) + |x_2|}.$$
 (52)

To do so, consider the non-negative concave quadratic test function

$$\phi(x) := ((x - a_i) \cdot e_1)((a_j - x) \cdot e_1) \ge 0 \quad \text{for } x \in R,$$

which is constant on all vertical lines and satisfies

$$\frac{1}{C}\operatorname{dist}(x, \{a_i, a_j\}) \le \phi(x) \le C \operatorname{dist}(x, \{a_i, a_j\}) \text{ on } R.$$
(53)

Next we will apply the monotonicity formula in Proposition 2.6 to the functions  $v \leq L_{\partial R \cup \{x\}} v$  which coincide on  $\partial R$ . The application of the proposition yields

$$\phi(x)\mathcal{L}^2(\partial^- L_{\partial R\cup\{x\}}v(x)) = \int \phi \mathrm{d}\mu_{L_{\partial R\cup\{x\}}v} \leq \int \phi \mathrm{d}\mu_v \lesssim F(v).$$
(54)

If  $v(x) \ge w_0(x)$ , (52) holds. Otherwise, we claim that  $\partial^- L_{\partial R \cup \{x\}} v(x)$  has area bounded from below by

$$\mathcal{L}^{2}(\partial^{-}L_{\partial R\cup\{x\}}v(x)) \gtrsim \min\left(\frac{(w_{0}(x) - v(x))^{2}}{\operatorname{dist}(x,\{a_{i},a_{j}\})|x_{2}|}, \frac{w_{0}(x) - v(x)}{\operatorname{dist}(x,\{a_{i},a_{j}\})}\right).$$
(55)

 $\langle \rangle$ 

To prove (55), take  $p_0 \in \partial^- w_0(x)$ . Write

$$(-L,0) = a_i = a_-, \quad (+L,0) = a_j = a_+$$

and set

$$p_1 = p_0 - \operatorname{sgn}(x_1) \frac{w_0(x) - v(x)}{|a_{\operatorname{sgn}(x_1)} - x|} e_1.$$

We note

$$-\operatorname{sgn}(x_1)\frac{e_1 \cdot (y-x)}{|a_{\operatorname{sgn}(x_1)} - x|} \le 1 \quad \text{for all } y \in \partial R$$

Thanks to this observation we obtain for every  $y \in \partial R$  the following chain of estimates:

$$p_{1} \cdot y - v(y) = p_{0} \cdot y - w_{0}(y) + (p_{1} - p_{0}) \cdot y + w_{0}(y) - v(y)$$

$$\leq p_{0} \cdot x - w_{0}(x) + (p_{1} - p_{0}) \cdot y$$

$$\leq p_{1} \cdot x - w_{0}(x) + (p_{1} - p_{0}) \cdot (y - x)$$

$$= p_{1} \cdot x - w_{0}(x) - (w_{0}(x) - v(x)) \frac{(y - x) \cdot e_{1} \operatorname{sgn}(x_{1})}{|a_{\operatorname{sgn}(x_{1})} - x|}$$

$$\leq p_{1} \cdot x - v(x) ,$$
(56)

where we have used  $p_0 \in \partial^- w_0(x)$  and  $w_0(y) \leq v(y)$  for all  $y \in \partial R$  to obtain the first inequality. The estimate (56) implies  $p_1 \in \partial^- L_{\partial R \cup \{x\}} v(x)$ .

In a similar way, we set

$$p_2 = p_0 - \operatorname{sgn}(x_2) \min\left(\frac{w_0(x) - v(x)}{|x_2|}, \frac{1}{2} |[\partial_2 w_0]|\right) e_2$$

We estimate assuming  $y \in \partial R$ , this time keeping the non-positive term  $r(x_2, y_2) := -w_0(y) + w_0(x) + (y - x) \cdot p_2$ , whose analogue we discarded in the previous calculation:

$$p_{2} \cdot y - v(y) \leq p_{2} \cdot x - w_{0}(x) + (p_{2} - p_{0}) \cdot (y - x) + r(x_{2}, y_{2})$$

$$\leq p_{2} \cdot x - w_{0}(x) - \operatorname{sgn}(x_{2}) \min\left(\frac{w_{0}(x) - v(x)}{|x_{2}|}, \frac{1}{2} |[\partial_{2}w_{0}]|\right) e_{2} \cdot (y - x) \quad (57)$$

$$+ r(x_{2}, y_{2}).$$

In order to derive from (57) the wished for estimate

$$p_2 \cdot y - v(y) \le p_2 \cdot x - v(x), \qquad (58)$$

we will assume that  $x_2 \ge 0$ , the proof for  $x_2 \le 0$  working out in an analogous fashion. We make a case distinction: If  $y_2 \ge x_2$ , then of course

$$-\min\left(\frac{w_0(x) - v(x)}{|x_2|}, \frac{1}{2}|[\partial_2 w_0]|\right)e_2 \cdot (y - x) \le 0,$$

and (58) follows from (57) and  $r(x_2, y_2) \leq 0$ . If  $0 \leq y_2 \leq x_2$  then

$$-\min\left(\frac{w_0(x)-v(x)}{|x_2|},\frac{1}{2}|[\partial_2 w_0]|\right)e_2\cdot(y-x)\le w_0(x)-v(x)\,,$$

and again (58) follows from (57) and  $r(x_2, y_2) \leq 0$ . Finally, for the case  $y_2 < 0$  we note that by the convexity of  $w_0$  and (51),

$$r(x_2, y_2) \le \lim_{s \downarrow 0} r(s, y_2) \equiv r(0+, y_2) \le -|[\partial_2 w_0]||y_2|.$$

Hence

$$-\min\left(\frac{w_0(x) - v(x)}{|x_2|}, \frac{1}{2}|[\partial_2 w_0]|\right) e_2 \cdot (y - x) \le w_0(x) - v(x) + \frac{1}{2}|[\partial_2 w_0]||y_2| \le w_0(x) - v(x) - \frac{1}{2}r(x_2, y_2),$$

and again (58) follows from (57). Hence we have proved

$$p_2 \in \partial^- L_{\partial R \cup \{x\}} v(x) \,.$$

Summarizing, we have that  $p_0, p_1, p_2 \in \partial^- L_{\partial R \cup \{x\}} v(x)$  and hence, by the convexity of the subdifferential at a single point, also

$$\partial^{-}L_{\partial R\cup\{x\}}v(x) \supseteq \left[p_0p_1p_2\right],$$

which implies

$$\mathcal{L}^{2}(\partial^{-}L_{\partial R\cup\{x\}}v(x)) \geq \frac{w_{0}(x) - v(x)}{|a_{\operatorname{sgn}(x_{1})} - x|} \min\left(\frac{w_{0}(x) - v(x)}{|x_{2}|}, \frac{1}{2}|[\partial_{2}w_{0}]|\right)$$
$$\gtrsim \min\left(\frac{(w_{0}(x) - v(x))^{2}}{\operatorname{dist}(x, \{a_{i}, a_{j}\})|x_{2}|}, \frac{w_{0} - v(x)}{\operatorname{dist}(x, \{a_{i}, a_{j}\})}\right),$$

i.e. (55). Combining (53), (54) and (55) yields our claim (52). Combining the latter with (50) yields the statement of the proposition.  $\Box$ 

## 4.5. Proof of the lower bound in Theorem 1.1.

*Proof of the lower bound in Theorem 1.1.* By the upper bound in Theorem 1.1 and Lemma 4.2 we may assume that

$$\|\mu_{v_h} - \bar{\mu}\|_{W^{-2,2}(\Omega)}^2 + h^2 \|\nabla^2 v_h\| \le Ch^{4/3} \left(\log \frac{1}{h}\right)^{1/3}$$

where the constant on the right hand side only depends on  $\Omega, \bar{\mu}$ . Choose  $(i, j) \in \mathcal{R}_{v_0}$ , and recall the definition of the rhombus  $R_{ij}$  from Lemmata 2.11 and 4.10. By Lemma 4.10,

 $a_l \notin R_{ij}$  for  $l \in \{1, \ldots, N\} \setminus \{i, j\}$ .

We may choose two open convex sets  $U_{ij}^1, U_{ij}^2$  such that  $R_{ij} \subset U_{ij}^1 \subset U_{ij}^2$  and

$$a_l \notin \overline{U_{ij}^2}$$
 for  $l \in \{1, \dots, N\} \setminus \{i, j\}$ .

We will write  $U^1 \equiv U^1_{ij}, U^2 \equiv U^2_{ij}$ . We claim that

$$F(v_h) := \int_{U^1} \operatorname{dist}(x, \{a_1, \dots, a_N\}) \mathrm{d}\mu_{v_h}(x)$$

satisfies

$$F(v_h) \le C(\Omega, \bar{\mu}) \left(h \log \frac{1}{h}\right)^{2/3}.$$
(59)

To see that (59) holds, choose  $r_i, r_j > 0$  independently of h such that

$$B(a_i, r_i), B(a_j, r_j) \subseteq U^1$$
  
$$B(a_i, r_i) \cap B(a_j, r_j) = \emptyset,$$

for  $k \in \{i, j\}$  choose  $\psi_k \in C_c^{\infty}(B(a_k, r_k))$  such that  $0 \leq \psi_k \leq 1$  and  $\psi_k = 1$  on  $B(a_k, r_k/2)$ , and complete the set  $\{\psi_i, \psi_j\}$  to a partition of unity  $\{\psi_i, \psi_j, \varphi\}$  of  $U^1$ , where  $\varphi \in C_c^{\infty}(U^2)$ . Let  $\eta \in C_c^{\infty}(\mathbb{R}^2)$  be a standard mollifier, and  $\eta_{\varepsilon}(\cdot) = \frac{1}{\varepsilon^2}\eta(\cdot/\varepsilon)$  for  $\varepsilon > 0$ . Now the *h*-dependent choice of test function we make to show (59) is the following: We write  $\varepsilon = h^{2/3}$  and set

$$\phi_h = (|\cdot - a_i| * \eta_{\varepsilon}) \psi_i + (|\cdot - a_j| * \eta_{\varepsilon}) \psi_j + \varphi.$$

A calculation yields

$$\begin{cases} \phi_h(x) \simeq \operatorname{dist}(x, \{a_i, a_j\}) + h^{2/3} & \text{in } U^1 \\ |\nabla \phi_h(x)| \le C & \text{in } U^2 \\ |\nabla^2 \phi_h(x)| \le \frac{C}{\operatorname{dist}(x, \{a_i, a_j\}) + h^{2/3}} & \text{in } U^2 \,, \end{cases}$$
(60)

where all constants on the right hand side only depend on  $U^1, U^2, \psi_i, \psi_j, \varphi$  (and may hence be said to depend on  $\Omega, \bar{\mu}$ ). Thus

$$\|\phi_h\|_{W^{2,2}_0(\Omega)} \le C \left(\log \frac{1}{h}\right)^{1/2}$$
36

and we obtain

$$F(v_h) \leq C \int \phi_h d\mu_{v_h}$$
  

$$\leq \|\phi_h\|_{W_0^{2,2}(\Omega)} \|\mu_{v_h} - \bar{\mu}\|_{W^{-2,2}(\Omega)} + \int \phi_h d\bar{\mu}$$
  

$$\leq Ch^{2/3} \log(1/h)^{1/2+1/6} + (\sigma_i + \sigma_j)h^{2/3}$$
  

$$\leq Ch^{2/3} \log(1/h)^{2/3},$$
(61)

proving (59).

We recall the notation  $R_{ij} = [a_i b_{ij}^+ a_j b_{ij}^-]$ , and assume without loss of generality that  $a_i = -a_j = Le_1$ . We set

$$w_h := L_{\{a_i, a_j, b_{ij}^+, b_{ij}^-\}} v_h$$

By Lemma 2.11,  $v_0$  is affine on  $[a_i b_{ij}^+ a_j]$  and on  $[a_i b_{ij}^- a_j]$ , with

$$\nabla v_0 = p^+ \text{ on } [a_i b_{ij}^+ a_j]$$
  
$$\nabla v_0 = p^- \text{ on } [a_i b_{ij}^- a_j]$$

and  $p^+ \neq p^-$ . By definition, there exist  $\tilde{p}_h^+, \tilde{p}_h^- \in \mathbb{R}^2$  such that

$$\nabla w_h = \tilde{p}_h^+ \text{ on } [a_i b_{ij}^+ a_j]$$
  
$$\nabla w_h = \tilde{p}_h^- \text{ on } [a_i b_{ij}^- a_j]$$

and by Proposition 4.7, we have

$$|p^{+} - \tilde{p}_{h}^{+}| + |p^{-} - \tilde{p}_{h}^{-}| \le Ch^{\gamma}$$
(62)

for  $h < h_0$ . Now we may apply Proposition 4.11 with  $U \equiv U^2$ ,  $\varepsilon_0 \equiv Ch_0^{\gamma}$ . We note that the hypothesis  $||v - v_0||_{L^{\infty}} \equiv ||v_h - v_0||_{L^{\infty}} \leq Ch^{\gamma} \leq \varepsilon_0$  is met by Proposition 4.7, provided that  $h_0$  is chosen small enough. Thus we obtain the existence of a constant  $C_1 > 0$  (that only depends on  $\Omega, \overline{\mu}$ ) such that

$$|v_h(x) - w_h(x)| \le C_1 \left( \left( h \log \frac{1}{h} \right)^{2/3} + \sqrt{\left( h \log \frac{1}{h} \right)^{2/3} |x_2|} \right) \quad \text{for } x \in R_{ij}.$$
(63)

Here we may choose  $C_1$  such that it additionally satisfies  $C_1 \ge \max(|p^+ - p^-|, 1)$ . In particular, setting

$$l(h) := \frac{16C_1^2 \left(h \log \frac{1}{h}\right)^{2/3}}{|\tilde{p}_h^+ - \tilde{p}_h^-|^2},$$

we obtain,

$$v_h(x_1, \pm l(h)) \ge w_h(x_1, \pm l(h)) - C_1 \left(1 + \frac{4C_1}{|\tilde{p}_h^+ - \tilde{p}_h^-|}\right) \left(h \log \frac{1}{h}\right)^{2/3},$$
 (64)

where we have assumed that  $(x_1, \pm l(h)) \in R_{ij}$ , which is the case for h small enough and  $x_1 \in [-L/2, L/2]$ . Moreover by convexity

$$v_h(x_1,0) \le w_h(x_1,0).$$
 (65)  
37

Combining (64) with (65) and using the convexity of  $v_h$ , we obtain

$$\begin{split} \int_{-l(h)}^{l(h)} \partial_{2}^{2} v_{h}(x_{1}, x_{2}) dx_{2} \\ &= \partial_{2} v_{h}(x_{1}, l(h)) - \partial_{2} v_{h}(x_{1}, -l(h)) \\ &\geq \frac{v_{h}(x_{1}, l(h)) - v_{h}(x_{1}, 0)}{l(h)} + \frac{v_{h}(x_{1}, -l(h)) - v_{h}(x_{1}, 0)}{l(h)} \\ &\geq \frac{w_{h}(x_{1}, l(h)) - w_{h}(x_{1}, 0)}{l(h)} + \frac{w_{h}(x_{1}, -l(h)) - w_{h}(x_{1}, 0)}{l(h)} \\ &- \frac{C_{1} \left(1 + \frac{4C_{1}}{|\tilde{p}_{h}^{+} - \tilde{p}_{h}^{-}|\right)}{8C_{1}^{2}} \\ &\geq |\tilde{p}_{h}^{+} - \tilde{p}_{h}^{-}| - \frac{|\tilde{p}_{h}^{+} - \tilde{p}_{h}^{-}|^{2}}{8C_{1}} - \frac{|\tilde{p}_{h}^{+} - \tilde{p}_{h}^{-}|}{2} \\ &\geq \frac{1}{4} |\tilde{p}_{h}^{+} - \tilde{p}_{h}^{-}| \,. \end{split}$$
(66)

Using the Cauchy-Schwarz inequality, we deduce

$$h^{2} \int_{-L/2}^{L/2} \int_{-l(h)}^{l(h)} |\partial_{2}^{2} v_{h}(x_{1}, x_{2})|^{2} dx_{2} dx_{1} \ge h^{2} L \frac{|\tilde{p}_{h}^{+} - \tilde{p}_{h}^{-}|^{2}}{8l(h)} \ge C h^{4/3} \log(1/h)^{-2/3},$$
(67)

with C > 0 depending only on  $\Omega$  and  $\bar{\mu}$ , where in the last inequality, we have used (62).  $\Box$ 

# APPENDIX A. SCALING WITHOUT BOUNDARY CONDITIONS: $h^2 \log \frac{1}{h}$

We observe that in fact the scaling for M disclinations without any further constraints such as boundary conditions will *not* be  $h^{4/3}$  but  $h^2 \log \frac{1}{h}$ , like a single disclination. We claim the following:

**Remark A.1.** Let  $\Omega \subseteq \mathbb{R}^2$  and  $\bar{\mu} = \sum_{i=1}^N \sigma_i \delta_{a_i}$  be as in Theorem 1.1. Then there exist constants  $C, h_0 > 0$  that only depend on  $\Omega$  and  $\bar{\mu}$  such that for every  $h < h_0$ , there exist  $(\mathbf{u}_h, v_h) \in W^{1,2}(\Omega; \mathbb{R}^2) \times W^{2,2}(\Omega)$  with

$$E_h(\mathbf{u}_h, v_h) \le Ch^2 \log \frac{1}{h}.$$

In the proof of the remark, we are going to use the following lemma:

**Lemma A.2.** Let  $\varepsilon \in (0,1)$  and  $u_{\varepsilon} : \mathbb{R}^2 \to \mathbb{R}$  the function

$$u_{\varepsilon}(x_1, x_2) := \begin{cases} |x_2| & \text{if } |x_2| \ge \varepsilon |x_1| \\ \frac{\varepsilon |x_1|}{2} + \frac{x_2^2}{2\varepsilon |x_1|} & \text{if } |x_2| < \varepsilon |x_1| \end{cases}$$

Then  $u_{\varepsilon} \in W^{2,\infty}_{\text{loc}}(\mathbb{R}^2 \setminus \{0\})$  is convex and one-homogeneous, with  $\mu_{u_{\varepsilon}} = \frac{4\varepsilon}{3}\delta_0$ . In addition,  $u_{\varepsilon}$  is linear in the two cones  $\{x \in \mathbb{R}^2 : x_2 > \varepsilon | x_1 |\}$  and  $\{x \in \mathbb{R}^2 : x_2 < \varepsilon | x_1 |\}$ 

In addition,  $u_{\varepsilon}$  is linear in the two cones  $\{x \in \mathbb{R}^2 : x_2 > \varepsilon | x_1 |\}$  and  $\{x \in \mathbb{R}^2 : x_2 < -\varepsilon | x_1 |\}$ .

*Proof.* We check that for  $x \neq 0$ 

$$\nabla u(x) = \begin{cases} e_2 & \text{if } x_2 > \varepsilon |x_1| \\ \begin{pmatrix} \frac{\varepsilon}{2} - \frac{x_2^2}{2\varepsilon x_1^2} \\ \frac{x_2}{\varepsilon x_1} \end{pmatrix} & \text{if } |x_2| \le \varepsilon |x_1| \\ -e_2 & \text{if } x_2 < \varepsilon |x_1| \end{cases}$$

In particular,  $\nabla u$  is locally Lipschitz in  $\mathbb{R}^2 \setminus \{0\}$ . A direct calculation shows that  $u(x) = \max_{y \in \mathbb{R}^2 \setminus \{0\}} x \cdot \nabla u(y)$ , showing that u is convex.

Since  $\nabla u$  is 0-homogeneous,  $\mu_u = \sigma \delta_0$  for some  $\sigma \ge 0$ . We calculate

$$\sigma = \mathcal{L}^2(\operatorname{conv}\{\nabla u(y) : y \in \mathbb{R}^2 \setminus \{0\}\}) = 2\int_{-1}^1 \frac{\varepsilon}{2}(1-t^2) \, dt = \frac{4\varepsilon}{3}.$$

Proof of Remark A.1. First we construct a solution  $v_1$  of  $\mu_{v_1} = \bar{\mu}$  that is in  $W^{2,\infty}_{\text{loc}}(\Omega \setminus \{a_1,\ldots,a_N\})$ . Define for  $\varepsilon > 0$  the horizontal double cone  $K_{\varepsilon} := \{x \in \mathbb{R}^2 : |x_2| \le \varepsilon |x_1|\}$ . Choose  $e \in S^1$  and  $\varepsilon > 0$  depending on  $\Omega$  and  $a_1,\ldots,a_N$  such that the sets

$$K_i := \Omega \cap (a_i + R_e K_{\varepsilon}), \quad i = 1, \dots, N,$$

have mutually positive distance from each other, where  $R_e \in SO(2)$  is the rotation mapping  $e_1$  to e. See Figure 4 for an illustration. Then define

$$v_1(x) := \sum_{i=1}^N \sqrt{\frac{3\sigma_i}{4\varepsilon}} u_{\varepsilon}(R_e^T(x-a_i)),$$

where  $u_{\varepsilon} : \mathbb{R}^2 \to \mathbb{R}$  is the function from Lemma A.2. By our choice of  $e, \varepsilon$ , it follows that  $\mu_{v_1} = \bar{\mu}$ . Note here that additivity of the Monge-Ampère measure holds only since  $\nabla^2 u_{\varepsilon} = 0$  outside of  $K_{\varepsilon}$ . Hence curl curl  $(\nabla v_1 \otimes \nabla v_1 - \nabla v_0 \otimes \nabla v_0) = 0$  in  $W^{-2,2}(\Omega)$ . By [ACGK06, Theorem 7.2]<sup>2</sup> this implies the existence of

$$\mathbf{u}_0 \in W^{1,2}(\Omega; \mathbb{R}^2)$$

such that

$$\nabla v_1 \otimes \nabla v_1 - \nabla v_0 \otimes \nabla v_0 = -(\nabla \mathbf{u}_0 + \nabla \mathbf{u}_0^T)$$

Now we define  $v_h$  on  $B(a_i, h)$  by requiring it to be convex, in  $W^{2,\infty}(\Omega)$  with  $|\nabla^2 v_h| \leq Ch^{-1}$ and

$$v_h = v_1, \quad \nabla v_h = \nabla v_1 \quad \text{ on } \partial B(a_i, h).$$

Setting  $v_h = v_1$  on  $\Omega \setminus \bigcup_{i=1}^N B(a_i, h)$  and  $\mathbf{u}_h = \mathbf{u}_0$  completes our construction. The estimate  $E_h(\mathbf{u}_h, v_h) \leq Ch^2 \log \frac{1}{h}$  is the result of a straightforward calculation.  $\Box$ 

<sup>&</sup>lt;sup>2</sup>In [ACGK06] the statement is made for  $\mathbb{R}^3$  valued functions, but the relevant proofs in this paper work out verbatim for  $\mathbb{R}^2$  valued functions.



FIGURE 4. Construction of the convex function  $v_1$  in the proof of Remark A.1. The double cones  $K_1, \ldots, K_5$  are shown in blue. Their orientation and width are chosen so as to make them pairwise disjoint, thus ensuring that  $\mu_{v_1} = \bar{\mu}$ .

#### References

- [ACGK06] C. Amrouche, P. G. Ciarlet, L. Gratie, and S. Kesavan. On the characterizations of matrix fields as linearized strain tensor fields. *Journal de mathématiques pures et appliquées*, 86(2):116–132, 2006.
- [AP10] B. Audoly and Y. Pomeau. Elasticity and geometry: from hair curls to the non-linear response of shells. Oxford University Press, 2010.
- [BCDM02] H. Ben Belgacem, S. Conti, A. DeSimone, and S. Müller. Energy scaling of compressed elastic films—three-dimensional elasticity and reduced theories. Arch. Ration. Mech. Anal., 164(1):1– 37, 2002.
- [BJ87] J. M. Ball and R. D. James. Fine phase mixtures as minimizers of energy. Archive for Rational Mechanics and Analysis, 100:13–52, 1987.
- [BK14] P. Bella and R.V. Kohn. Wrinkles as the result of compressive stresses in an annular thin film. Comm. Pure Appl. Math., 67(5):693–747, 2014.
- [BK17] P. Bella and R. V. Kohn. Wrinkling of a thin circular sheet bounded to a spherical substrate. *Philos. Trans. Roy. Soc. A*, 375(2093):20160157, 20, 2017.
- [CC95] L. A. Caffarelli and X. Cabré. Fully nonlinear elliptic equations, volume 43. American Mathematical Soc., 1995.
- [CM98] E. Cerda and L. Mahadevan. Conical surfaces and crescent singularities in crumpled sheets. *Phys. Rev. Lett.*, 80:2358–2361, Mar 1998.
- [CM05] E. Cerda and L. Mahadevan. Confined developable elastic surfaces: cylinders, cones and the elastica. Proc. Roy. Soc. London Ser. A, 461(2055):671–700, 2005.
- [CM08] S. Conti and F. Maggi. Confining thin elastic sheets and folding paper. Arch. Ration. Mech. Anal., 187(1):1–48, 2008.
- [COT17] S. Conti, H. Olbermann, and I. Tobasco. Symmetry breaking in indented elastic cones. Math. Mod. Meth. Appl. S., 27:291–321, 2017.
- [DW01] B.A. DiDonna and T.A. Witten. Anomalous strength of membranes with elastic ridges. Phys. Rev. Lett., 87:206105, Oct 2001.
- [EG92] L.C. Evans and R.F. Gariepy. Measure Theory and Fine Properties of Functions. CRC Press, 1992.
- [Fig17] A. Figalli. The Monge–Ampère equation and its applications. EMS, 2017.
- [FJM02] G. Friesecke, R.D. James, and S. Müller. A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity. *Comm. Pure Appl. Math.*, 55(11):1461–1506, 2002.

- [FJM06] G. Friesecke, R.D. James, and S. Müller. A hierarchy of plate models derived from nonlinear elasticity by Gamma-convergence. Arch. Ration. Mech. Anal., 180(2):183–236, 2006.
- [GO24] P. Gladbach and H. Olbermann. Variational competition between the full Hessian and its determinant for convex functions. *Nonlinear Analysis*, 242:113498, 2024.
- [GR90] A. Griewank and P.J. Rabier. On the smoothness of convex envelopes. Transactions of the American Mathematical Society, 322(2):691–709, 1990.
- [KK01] B. Kirchheim and J. Kristensen. Differentiability of convex envelopes. C. R. Acad. Sci. Paris Sér. I Math., 333(8):725–728, 2001.
- [KM92] R. V. Kohn and S. Müller. Relaxation and regularization of nonconvex variational problems. In Proceedings of the Second International Conference on Partial Differential Equations (Italian) (Milan, 1992), volume 62, pages 89–113 (1994), 1992.
- [KM94] R. V. Kohn and S. Müller. Surface energy and microstructure in coherent phase transitions. Comm. Pure Appl. Math., 47(4):405–435, 1994.
- [KN13] R.V. Kohn and H.-M. Nguyen. Analysis of a compressed thin film bonded to a compliant substrate: the energy scaling law. J. Nonlinear Sci., 23(3):343–362, 2013.
- [Kui55] N.H. Kuiper. On C<sup>1</sup>-isometric imbeddings. I, II. Nederl. Akad. Wetensch. Proc. Ser. A. 58, Indag. Math., 17:545–556, 683–689, 1955.
- [LDR95] H. Le Dret and A. Raoult. The nonlinear membrane model as variational limit of nonlinear three-dimensional elasticity. *Journal de mathématiques pures et appliquées*, 74(6):549–578, 1995.
- [Lew20] M. Lewicka. Quantitative immersability of Riemann metrics and the infinite hierarchy of prestrained shell models. Arch. Ration. Mech. Anal., 236(3):1677–1707, 2020.
- [LGL<sup>+</sup>95] A. Lobkovsky, S. Gentges, H. Li, D. Morse, and T.A. Witten. Scaling properties of stretching ridges in a crumpled elastic sheet. *Science*, 270(5241):1482–1485, 1995.
- [LMP11] M. Lewicka, M. G. Mora, and M. Reza Pakzad. The matching property of infinitesimal isometries on elliptic surfaces and elasticity of thin shells. Arch. Ration. Mech. Anal., 200(3):1023– 1050, 2011.
- [LW05] T. Liang and T.A. Witten. Crescent singularities in crumpled sheets. Phys. Rev. E, 71:016612, Jan 2005.
- [Nas54] J. Nash.  $C^1$  isometric imbeddings. Ann. Math. (2), 60:383–396, 1954.
- [Olb16] H. Olbermann. Energy scaling law for the regular cone. J. Nonlinear Sci., 26:287–314, 2016.
- [Olb17] H. Olbermann. Energy scaling law for a single disclination in a thin elastic sheet. Arch. Ration. Mech. Anal., 224:985–1019, 2017.
- [Olb18] H. Olbermann. The shape of low energy configurations of a thin elastic sheet with a single disclination. Analysis & PDE, 11(5):1285–1302, 2018.
- [Olb19] H. Olbermann. On a boundary value problem for conically deformed thin elastic sheets. Analysis & PDE, 12:245–258, 2019.
- [PDK<sup>+</sup>08] L. Pocivavsek, R. Dellsy, A. Kern, S. Johnson, B. Lin, K. Y. C. Lee, and E. Cerda. Stress and fold localization in thin elastic membranes. *Science*, 320(5878):912–916, 2008.
- [Roc70] R. T. Rockafellar. Convex analysis, volume No. 28 of Princeton Mathematical Series. Princeton University Press, Princeton, NJ, 1970.
- [Tar07] Luc Tartar. An introduction to Sobolev spaces and interpolation spaces, volume 3. Springer Science & Business Media, 2007.
- [Tob21] I. Tobasco. Curvature-driven wrinkling of thin elastic shells. Archive for Rational Mechanics and Analysis, 239(3):1211–1325, 2021.
- [Ven04] S.C. Venkataramani. Lower bounds for the energy in a crumpled elastic sheet—a minimal ridge. Nonlinearity, 17(1):301–312, 2004.
- [Wit07] T.A. Witten. Stress focusing in elastic sheets. Rev. Mod. Phys., 79:643–675, Apr 2007.
- [WL93] T.A. Witten and H. Li. Asymptotic shape of a fullerene ball. *Europhys. Lett.*, 23(1):51, 1993.

(Peter Gladbach) INSTITUT FÜR ANGEWANDTE MATHEMATIK, UNIVERSITÄT BONN, 53115 BONN, GER-MANY (Heiner Olbermann) INSTITUT DE RECHERCHE EN MATHÉMATIQUE ET PHYSIQUE, UCLOUVAIN, 1348 LOUVAIN-LA-NEUVE, BELGIUM

Email address, Peter Gladbach: gladbach@iam.uni-bonn.de

Email address, Heiner Olbermann: heiner.olbermann@uclouvain.be