# Double transverse spin asymmetry in the $p^{\uparrow}\bar{p}^{\uparrow}$ Drell-Yan process from Sivers functions

Zhun Lu,<sup>1</sup> Bo-Qiang Ma,<sup>2, \*</sup> and Ivan Schmidt<sup>1,†</sup>

<sup>1</sup>Center of subatomic studies and Departamento de Física, Universidad

Técnica Federico Santa María, Casilla 110-V, Valparaíso, Chile

<sup>2</sup>School of Physics and MOE Key Laboratory of Heavy Ion Physics, Peking University, Beijing 100871, China

We show that the transverse double spin asymmetry (DSA) in the Drell-Yan process contributed only from the Sivers functions can be picked out by the weighting function  $\frac{Q_T}{M^2}(\cos(\phi - \phi_{S_1})\cos(\phi - \phi_{S_2}) + 3\sin(\phi - \phi_{S_1})\sin(\phi - \phi_{S_2}))$ . The asymmetry is proportional to the product of two Sivers functions from each hadron  $f_{1T}^{\perp(1)} \times f_{1T}^{\perp(1)}$ . Using two sets of Sivers functions extracted from the semi-inclusive deeply elastic scattering data at HERMES, we estimate this asymmetry in the  $p^{\uparrow}\bar{p}^{\uparrow}$ Drell-Yan process which is possible to be performed in HESR at GSI. The prediction of DSA in the Drell-Yan process contributed by the function  $g_{1T}(x, \mathbf{k}_T^2)$ , which can be extracted by the weighting function  $\frac{Q_T}{M^2}(3\cos(\phi - \phi_{S_1})\cos(\phi - \phi_{S_2}) + \sin(\phi - \phi_{S_1})\sin(\phi - \phi_{S_2}))$ , is also given at GSI.

PACS numbers: 13.88.+e, 13.85.Qk

#### I. INTRODUCTION

The Sivers effect [1] was proposed originally to explain the large single spin asymmetries (SSA) observed in inclusive pion hadro-production  $(p^{\uparrow}p \rightarrow \pi X)$  at FNAL [2]. The effect can be quantitatively described by a  $\mathbf{k}_T$ dependent distribution named as Sivers function [3, 4]  $f_{1T}^{\perp}(x, \mathbf{k}_T^2)$ , which is the distribution of unpolarized partons in a transversely polarized proton. It arises from a non-trivial correlation between the nucleon transverse spin and the intrinsic transverse momenta in the nucleon. Despite its (naively) *T*-odd property [5], Sivers function has been proven to be non-vanishing [6] due to its special gauge-link property [7, 8, 9].

Recently the SSA measured in semi-inclusive deeply inelastic scattering (SIDIS) processes with transversely polarized targets at HERMES [10, 11, 12] and COM-PASS [13, 14], has been shown to be interpreted by the Sivers effect. The asymmetry is identified by the angular dependence  $\sin(\phi - \phi_S)$ , where  $\phi$  and  $\phi_S$  denote respectively the azimuthal angles of the produced hadron and of the nucleon spin polarization, with respect to the lepton scattering plane. The coexistent Collins asymmetry [5], with a angular dependence  $\sin(\phi + \phi_S)$ , has also been measured in those experiments. The data on the Sivers SSA has been utilized by different groups [15, 16, 17, 18, 19] to extract the Sivers functions of the proton, especially those for the u and d quarks, on the basis of the generalized factorization [20, 21]. Those sets of parametrization of the Sivers functions are qualitatively in agreement [22] among themselves, and were applied to predict the Sivers SSA in various processes in the established or planed facilities, such as the SIDIS at JLab, and the Drell-Yan process at COMPASS, RHIC

and GSI.

In this paper, we will investigate the role of the Sivers function on the transverse double spin asymmetry (DSA) in the Drell-Yan process. The transverse DSA has been investigated [23] for many years, and is believed to have advantage to unravel the transverse spin property of the nucleon [24], especially the transversity distribution  $h_1(x)$  [25]. Various azimuthal asymmetries contributed by different  $\mathbf{k}_T$ -dependent distribution functions have been analyzed and given in Refs. [26] and [27]. As shown in Ref. [27], the Sivers function contributes to the DSA in the Drell-Yan process through the product  $f_{1T}^{\perp} \times f_{1T}^{\perp}$ . However, this DSA is mixed with the contribution from another  $\mathbf{k}_T$ -dependent distribution function  $g_{1T}(x, \mathbf{k}_T^2)$ . We will show that through the appropriate weighting function  $\frac{Q_T^2}{M^2}(\cos(\phi - \phi_{S_1})\cos(\phi - \phi_{S_2}) + 3\sin(\phi - \phi_{S_1})\sin(\phi - \phi_{S_2}))$ , the asymmetry from the Sivers function can be isolated without mixing with the contribution from other functions. Using two sets of parameterizations [16, 18] of the Sivers functions we calculate the double spin asymmetry from the Sivers functions in the  $p^{\uparrow}\bar{p}^{\uparrow}$  Drell-Yan process at GSI. An asymmetry around 1 % is predicted. The asymmetries estimated from these two sets of Sivers functions are quantitatively different. Therefore measuring the DSA in the Drell-Yan process can provide new information on Sivers functions, especially their sizes. The transverse DSA contributed by  $g_{1T}(x, \mathbf{k}_T^2)$  through the product  $g_{1T} \times g_{1T}$  can also be picked out by another weighting function. We estimate this asymmetry by adopting a  $g_{1T}$  coming from the combination of a Lorentz invariance relation presented in Refs. [28, 29] and the Wandura-Wilzeck approximation [30].

## II. EXTRACTING DSA CONTRIBUTED BY THE SIVERS FUNCTIONS

The importance of the transverse-momentum distributions of quarks for a full understanding of the struc-

<sup>\*</sup>Corresponding author. Electronic address: mabq@phy.pku.edu.cn

<sup>&</sup>lt;sup>†</sup>Corresponding author. Electronic address: ivan.schmidt@usm.cl

ture of hadrons has been widely recognized in the last decade [4, 29, 31, 32]. A comprehensive leading-twist tree level analysis of the (spin dependent) Drell-Yan process in terms of  $\mathbf{k}_T$ -dependent distributions has been given in Ref. [26]. The role of the *T*-odd  $\mathbf{k}_T$ -dependent distributions in this process has been presented in Ref. [27]. In the Collins-Soper frame [33] the leading order unpolarized differential cross-section for the Drell-Yan process  $h_1(P_1) + h_2(P_2) \rightarrow \gamma^*(q) + X \rightarrow l^+(l_1) + l^-(l_2) + X$  has the form [27]

$$\frac{d\sigma^{(0)}(h_1h_2 \to l\bar{l}X)}{d\Omega dx_1 dx_2 d^2 \mathbf{q}_T} = \frac{\alpha_{em}^2}{3Q^2} \sum_q e_q^2 \left\{ A(y)\mathcal{F}[f_1^q f_1^{\bar{q}}] + B(y)\cos 2\phi \mathcal{F}\left[ (2\hat{\mathbf{h}} \cdot \mathbf{p}_T \hat{\mathbf{h}} \cdot \mathbf{k}_T - \mathbf{p}_T \cdot \mathbf{k}_T) \frac{h_1^{\perp q} h_1^{\perp \bar{q}}}{M_1 M_2} \right] \right\}$$
(1)

where q denotes the quark flavors, the notation

$$\mathcal{F}[f_1 f_1] = \int d^2 \mathbf{p}_\perp d^2 \mathbf{k}_\perp \delta^2 (\mathbf{p}_T + \mathbf{k}_T - \mathbf{q}_T) \\ \times f_1(x_1, \mathbf{p}_T^2) f_1(x_2, \mathbf{k}_T^2)$$
(2)

shows the convolution of transverse momenta,  $Q^2 = q^2$ is the invariance mass of the lepton pair,  $q_T$  is the transverse momentum of the lepton pair,  $\hat{\mathbf{h}} = \mathbf{q}_T/Q_T$ ,  $\phi$  is the angle between the hadron plane and the lepton plane, and

$$A(y) = \left(\frac{1}{2} - y + y^2\right) = \frac{1}{4}(1 + \cos^2\theta), \qquad (3)$$

$$B(y) = y(1-y) = \frac{1}{4}\sin^2\theta,$$
 (4)

in the c.m. frame of the lepton pair.

The function  $h_1^{\perp}$  in the second line of (1) is the Boer-Mulders function [4], the chiral-odd partner of the Sivers function. This function has attracted a lot of interest [34, 35, 36] recently because it can account for the anomalous  $\cos 2\phi$  asymmetries [37, 38] observed in the unpolarized Drell-Yan process, given by the second term of Eq. (1) has shown.

The leading order differential cross-section for the double transversely polarized Drell-Yan process is [27]

$$\frac{d\sigma^{(2)}(h_1^{\uparrow}h_2^{\uparrow} \to l\bar{l}X)}{d\Omega dx_1 dx_2 d^2 \mathbf{q}_T} = \frac{\alpha_{em}^2}{3Q^2} \sum_q \left\{ \cdots + \frac{A_1(y)}{2} |\mathbf{S}_{1T}| |\mathbf{S}_{2T}| \cos(2\phi - \phi_{S_1} - \phi_{S_2}) \right. \\ \times \mathcal{F}\left[ \hat{\mathbf{h}} \cdot \mathbf{p}_T \ \hat{\mathbf{h}} \cdot \mathbf{k}_T \frac{f_{1T}^{\perp q} f_{1T}^{\perp \bar{q}} - g_{1T}^q g_{1T}^{\bar{q}}}{M_1 M_2} \right] \\ \left. - \frac{A_1(y)}{2} |\mathbf{S}_{1T}| |\mathbf{S}_{2T}| \cos(\phi - \phi_{S_1}) \cos(\phi - \phi_{S_2}) \right. \\ \times \mathcal{F}\left[ \mathbf{p}_T \cdot \mathbf{k}_T \frac{f_{1T}^{\perp q} f_{1T}^{\perp \bar{q}}}{M_1 M_2} \right]$$

$$-\frac{A_{1}(y)}{2} |\mathbf{S}_{1T}| |\mathbf{S}_{2T}| \sin(\phi - \phi_{S_{1}}) \sin(\phi - \phi_{S_{2}}) \times \mathcal{F}\left[\mathbf{p}_{T} \cdot \mathbf{k}_{T} \frac{g_{1T}^{q} g_{1T}^{\bar{q}}}{M_{1} M_{2}}\right] \bigg\},$$
(5)

The ... indicates the terms which will not contribute in our analysis below,  $\phi_{S_1}$  and  $\phi_{S_2}$  are the angles between  $S_{1T}$ ,  $S_{2T}$  and the lepton plane, respectively.

As shown in (5), the Sivers function can contribute to the transverse DSA through the product  $f_{1T}^{\perp} \times f_{1T}^{\perp}$ . However this asymmetry is mixed with the asymmetry to which it contributes another  $\mathbf{k}_T$ -dependent distribution  $g_{1T}(x, \mathbf{k}_T^2)$ . The main goal of this paper is to isolate the asymmetry contributed by the Sivers function. The starting point is the method introduced in Ref. [39], by which one integrates the differential cross section with a proper weighting function  $W(Q_T, \phi, \phi_{S_1}, \phi_{S_2})$ , as follows:

$$\langle W(Q_T, \phi, \phi_{S_1}, \phi_{S_2}) \rangle$$

$$= \int d\phi d\phi_{S_1} d\mathbf{q}_T^2 \frac{d\sigma(h_1 h_2 \to l\bar{l}X)}{d\Omega dx_1 dx_2 d^2 \mathbf{q}_T}$$

$$\times W(Q_T, \phi, \phi_{S_1}, \phi_{S_2}). \tag{6}$$

With the above weighting procedure, one can pick up the terms in which one is interested. Besides this, one can de-convolute the transverse momentum integration in a model independent way.

The unpolarized angular independent cross section can be picked out by using the weighting function 1, from Eq. (1):

$$\left(\frac{A(y)\alpha_{em}^{2}}{3Q^{2}}\right)^{-1} \cdot \langle 1 \rangle_{UU} \\ = 4\pi^{2} \sum_{q} e_{q}^{2} f_{1}^{q}(x_{1}) f_{1}^{\bar{q}}(x_{2})$$
(7)

We denote  $W_C = \cos(\phi - \phi_{S_1})\cos(\phi - \phi_{S_2})$  and  $W_S = \sin(\phi - \phi_{S_1})\sin(\phi - \phi_{S_2})$ . Given the weighting function  $\frac{Q_T^2}{M^2}W_C$  (assuming  $M_1 = M_2 = M$ , i.e. the colliding two hadrons are nucleons), we can obtain the following term from (5):

$$\left(\frac{A(y)\alpha_{em}^{2}}{3Q^{2}}\right)^{-1} \cdot \left\langle \frac{Q_{T}^{2}}{M_{p}^{2}}W_{C} \right\rangle_{TT} \\
= \pi^{2} \sum_{q} e_{q}^{2} \left\{ 3 \left[ f_{1T}^{\perp(1)q}(x_{1}) f_{1T}^{\perp(1)\bar{q}}(x_{2}) - g_{1T}^{(1)q}(x_{1}) g_{1T}^{(1)\bar{q}}(x_{2}) \right] - 2 f_{1T}^{\perp(1)q}(x_{1}) f_{1T}^{\perp(1)\bar{q}}(x_{2}) \right\} \\
= \pi^{2} \sum_{a} e_{a}^{2} \left[ f_{1T}^{\perp(1)q}(x_{1}) f_{1T}^{\perp(1)\bar{q}}(x_{2}) - 3 g_{1T}^{(1)q}(x_{1}) g_{1T}^{(1)\bar{q}}(x_{2}) \right], \quad (8)$$

where  $f_{1T}^{\perp(1)}(x)$  and  $g_{1T}^{(1)}(x)$  are the first  $\mathbf{k}_T^2$ -moments, defined as:

$$f_{1T}^{\perp(1)}(x) = \int d^2 \mathbf{k}_T \frac{\mathbf{k}_T^2}{2M^2} f_{1T}^{\perp}(x, \mathbf{k}_T^2), \qquad (9)$$

$$g_{1T}^{(1)}(x) = \int d^2 \mathbf{k}_T \frac{\mathbf{k}_T^2}{2M^2} g_{1T}(x, \mathbf{k}_T^2).$$
(10)

The factor  $Q_T^2$  introduced in the weighting function ensures that the transverse momentum integration in (8) can be de-convoluted (for details, refer to the Appendix). Again, applying the weighting function  $\frac{Q_T^2}{M^2}W_S$  on (5), we arrive at

$$\left(\frac{A(y)\alpha_{em}^2}{3Q^2}\right)^{-1} \cdot \left\langle \frac{Q_T^2}{M_p^2} W_S \right\rangle_{TT}$$

$$= -\pi^2 \sum_q e_q^2 \left\{ 3 \left[ f_{1T}^{\perp(1)q}(x_1) f_{1T}^{\perp(1)\bar{q}}(x_2) - g_{1T}^{(1)q}(x_1) g_{1T}^{(1)\bar{q}}(x_2) \right] - 2g_{1T}^{(1)q}(x_1) g_{1T}^{(1)\bar{q}}(x_2) \right\}$$

$$= \pi^2 \sum_a e_a^2 \left[ -3f_{1T}^{\perp(1)q}(x_1) f_{1T}^{\perp(1)\bar{q}}(x_2) + g_{1T}^{(1)q}(x_1) g_{1T}^{(1)\bar{q}}(x_2) \right],$$

$$(11)$$

Therefore, combining (8) and (11), we can extract the term contributing to the transverse DSA and coming only from the Sivers functions:

$$\left(\frac{A(y)\alpha_{em}^2}{3Q^2}\right)^{-1} \cdot \left\langle \frac{Q_T^2}{M_p^2} \left(W_C + 3W_S\right) \right\rangle_{TT} = -8\pi^2 \sum_q e_q^2 f_{1T}^{\perp(1)q}(x_1) f_{1T}^{\perp(1)\bar{q}}(x_2), \quad (12)$$

with the weighting function  $\frac{Q_T^2}{M^2}(W_C + 3W_S)$ . By taking the ratio between (13) and (7), we define

By taking the ratio between (13) and (7), we define the weighted double spin asymmetry as follows

$$A_{TT}^{f} = \frac{\left\langle \frac{Q_{T}^{2}}{M^{2}} (W_{C} + 3W_{S}) \right\rangle_{TT}}{\langle 1 \rangle_{UU}} \\ = -\frac{2\sum_{q} e_{q}^{2} f_{1T}^{\perp(1)q}(x_{1}) f_{1T}^{\perp(1)\bar{q}}(x_{2})}{\sum_{q} e_{q}^{2} f_{1}^{q}(x_{1}) f_{1}^{\bar{q}}(x_{2})}.$$
 (13)

The above equation thus provides a possibility to study the Sivers function by measuring the transverse DSA in the Drell-Yan process.

Also, from (8), (11) and (7) we can get another type of DSA:

$$A_{TT}^{g} = \frac{\left\langle \frac{Q_{T}^{2}}{M^{2}} (3W_{C} + W_{S}) \right\rangle_{TT}}{\langle 1 \rangle_{UU}} \\ = -2 \frac{\sum_{q} e_{q}^{2} g_{1T}^{(1)q}(x_{1}) g_{1T}^{(1)\bar{q}}(x_{2})}{\sum_{q} e_{q}^{2} f_{1}^{q}(x_{1}) f_{1}^{\bar{q}}(x_{2})}, \qquad (14)$$

which is contributed only by  $g_{1T}$ .

### III. NUMERICAL RESULTS

In this section we will give numerical results on the DSA from the Sivers functions. We consider the transversely polarized proton antiproton Drell-Yan process,



FIG. 1: The DSA in the proton antiproton Drell-Yan process at GSI coming only from the Sivers functions, and calculated from Eq. (13). The kinematics is  $s = 45 \text{ GeV}^2$  and  $Q^2 =$  $2.5 \text{ GeV}^2$ . The solid and dashed curve use the Sivers functions in Ref. [16] and in Ref. [18], respectively.

where the valence Sivers functions are involved, so that a larger asymmetry should be measured compared to the  $p^{\uparrow}p^{\uparrow}$  Drell-Yan process. The  $p^{\uparrow}\bar{p}^{\uparrow}$  Drell-Yan process is possible to be performed in the planned high energy storage ring (HESR) [40] at GSI. We study the transverse DSA at GSI from the Sivers functions, based on Eq. (13). For this end we need to know the input for the Sivers functions. Several groups [16, 18, 19] have parameterized the Sivers functions based on the data of SIDIS at HERMES [11, 12], and partially based on COM-PASS data [13]. The kinematics in GSI can be chosen as the c.m. energy  $s = 45 \text{ GeV}^2$ . For the invariance mass square of the lepton pair we choose  $Q^2 = 2.5 \text{ GeV}^2$ , which is close to the scale at HERMES. Therefore these sets of Sivers functions extracted from the data of HERMES can be applied to predict the asymmetries at GSI in the kinematics regime we give above. We will adopt two sets of Sivers functions, which are the sets in Refs. [16] and [18], respectively. The Sivers functions in Ref. [19] can not be applied here since in that paper  $f_{1T}^{\perp(1/2)}(x)$  is given while we use  $f_{1T}^{\perp(1)}(x)$  in our calculation.

To use these Sivers functions one should notice that T-odd distribution functions in the DIS and in the Drell-Yan process have a minus sign difference [7]. However in the  $p^{\uparrow}\bar{p}^{\uparrow}$  Drell-Yan process two Sivers functions appear in the product, therefore the sign difference doesn't matter here and the functions can be used directly.

In Ref. [16] the Sivers functions are parameterized as

$$-\frac{\mathbf{k}_T}{M}f_{1T}^{\perp,q}(x,\mathbf{k}_T^2) = N_q(x)f_1^q(x)g(\mathbf{k}_T^2)h(\mathbf{k}_T^2), \quad (15)$$



FIG. 2: The DSA in the proton antiproton Drell-Yan process at GSI coming from the function  $g_{1T}$ , defined in Eq. (14). The kinematics is  $s = 45 \text{ GeV}^2$  and  $Q^2 = 2.5 \text{ GeV}^2$ .

with

$$N_q(x) = N_q x^{a_q} (1-x)^{b_q} \frac{(a_q + b_q)^{(a_q + b_q)}}{a_q^{a_q} b_q^{b_q}}, \quad (16)$$

$$g(\mathbf{k}_T^2) = \frac{e^{-\mathbf{k}_T^2/\langle k_T^2 \rangle}}{\pi \langle k_T^2 \rangle},\tag{17}$$

for q = u, d. For the function  $h(\mathbf{k}_T^2)$  two options are considered:

(a) 
$$h(\mathbf{k}_T^2) = \frac{2k_T M_0}{k_T^2 + M_0^2},$$
 (b)  $\sqrt{2e} \frac{p_T}{M'} e^{-k_T^2/M'}.$  (18)

In our calculation we will adopt option (b) in Eq.(18), and the central values of their fit. This parametrization has taken advantage of the more precise data [12] at HERMES.

In Ref. [18] the authors give the set of Sivers functions for the u and d quark as,

$$x f_{1T}^{\perp(1),u}(x) = -x f_{1T}^{\perp(1),d} = -0.17 x^{0.66} (1-x)^5,$$
 (19)

extracted from the published HERMES data [11], and whose form is based on the limit of a large number of colours  $N_c$ .

For the unpolarized distribution we use the MRST2001(LO set) parametrization [41]. In Fig. 1 we present the DSA from Sivers functions at GSI, as a function of  $x_1$ . A sizable asymmetry is predicted. The asymmetry (solid line) based on the Sivers functions from Ref. [16] is much larger than the asymmetry (dashed line) based on the Sivers functions from Ref. [18]. As explained in Ref. [42], taking into account the more precise data [12] of HERMES, larger Sivers functions can be

dashed curve in Fig. 1. Thus the difference between the asymmetries from the two sets of Sivers functions may be reduced. Depending on the accuracy of the experimental measurements on the transverse DSA at GSI, useful constraints on the Sivers functions could be obtained, but it might be hard to distinguish between different parameterizations without high precision measurements.

Finally, we will predict the DSA contributed by the function  $g_{1T}(x, \mathbf{k}_T^2)$  at GSI. This function, describing longitudinal polarization of quarks in the transversely polarized target, also plays role in the double polarized (longitudinal-transverse) SIDIS process [28, 43]. A treatment on  $g_{1T}(x, \mathbf{k}_T^2)$  is the so-called Lorentz invariance relation that connect the first  $\mathbf{k}_T^2$  moment of  $g_{1T}(x, \mathbf{k}_T^2)$  with the twist-three distribution function  $g_2(x)$ :

$$g_2^q(x) = \frac{d}{dx} g_{1T}^{q(1)}(x).$$
(20)

Using the Wandzura and Wilczek approximation for  $g_2^q$ 

$$g_2^q(x) \approx -g_1^a(x) + \int_x^1 dy \frac{g_1^q(x)}{y},$$
 (21)

the following relation was derived in Ref. [28]

$$g_{1T}^{(1)q}(x) \approx x \int_{x}^{1} dy \frac{g_{1}^{q}(x)}{y}.$$
 (22)

For the polarized parton distribution we apply the GRSV2001 (standard scenario) parametrization [44], and for the unpolarized distribution we use GRV98 LO parametrization [45], following the choice in Ref. [43]. In Fig. 2 we show the DSA contributed by  $g_{1T}(x, \mathbf{k}_T^2)$  in the  $p^{\uparrow}\bar{p}^{\uparrow}$  Drell-Yan process at GSI with  $s = 45 \text{ GeV}^2$  and  $Q^2 = 2.5 \text{ GeV}^2$ . An asymmetry of 1% is predicted.

We end this section with some comment. In our calculation, especially in the case of Siver DSA, we choose  $Q^2 = 2.5 \text{ GeV}^2$ . This value is consistent with the averaged scale  $\langle Q^2 \rangle$  in the HERMES experiment, from which the Sivers functions were extracted. Therefore, the parameterizations for Sivers functions in Refs. [16, 18] can be applied here without further assumptions. Experimental measurements at GSI can also cover the continuous Drell-Yan masses 2-5 GeV which corresponds to  $Q^2$  in the range 4 - 25 GeV<sup>2</sup>. To estimate the asymmetries in this region one should use the fitted functions evolved to the relevant scale, which is not trivial for the  $\mathbf{k}_T$ -dependent distributions [46]. Therefore we assume that the ratios in Eqs.(13) and (14) scale with  $Q^2$ . In this region, The result is similar to the one which can be obtained at the fixed value of  $Q^2 = 2.5 \text{ GeV}^2$ . Also there is the subtlety that the next to leading order correction of the hard process could lead the substantial K-factor on the transversely polarized cross-section. Since we calculate an asymmetry, which is essentially a ratio where the  $Q^2$  dependences in the numerator and denominator tend

to cancel each other, the effect of both the  $Q^2$  dependence and K-factors do not introduce a strong influence on the resulting prediction coming from Eqs. (13) and (14).

### IV. SUMMARY

We have performed an analysis of the transverse DSA in the Drell-Yan process contributed by the Sivers functions through the term  $f_{1T}^{\perp} \times f_{1T}^{\perp}$ . The asymmetry can be isolated through the appropriate weighting function  $\frac{Q_T^2}{M^2}(\cos(\phi-\phi_{S_1})\cos(\phi-\phi_{S_2})+3\sin(\phi-\phi_{S_1})\sin(\phi-\phi_{S_2})),$  without mixing with the contribution from other distribution functions. Using two sets of Sivers functions parameterizing the SSA data in the SIDIS process, we calculate the double spin asymmetry in the  $p^{\uparrow}\bar{p}^{\uparrow}$  Drell-Yan process from the Sivers functions at GSI. An asymmetry around to 1% is predicted. The asymmetries estimated from these two sets of Sivers functions are quantitatively different. Therefore measurements of the DSA in Drell-Yan process can provide new information on the Sivers functions, especially their sizes. The transverse DSA contributed by  $g_{1T}(x, \mathbf{k}_T^2)$  through the product  $g_{1T} \times g_{1T}$  in the Drell-Yan process can also be picked out by a weighting function. We estimate this asymmetry at GSI by adopting  $g_{1T}$  from the combination of the Lorentz invariance relation and the Wandura-Wilzeck approximation. The investigation on the double transversely polarized Drell-Yan process thus can shed light on the knowledge of  $\mathbf{k}_{T}$ -dependent distribution functions, including the Sivers functions.

#### Acknowledgments

This work is partially supported by National Natural Science Foundation of China (Nos. 10421503, 10575003, 10505011, 10528510), by the Key Grant Project of Chinese Ministry of Education (No. 305001), by the Research (China), by Fondecyt (Chile) under Project No. 3050047.

Fund for the Doctoral Program of Higher Education

# **APPENDIX: MOMENTS**

To derive (8) and (11) we have used the following transverse momentum integrations:

$$\int d^{2}\mathbf{k}_{T} d^{2}\mathbf{p}_{T} \delta^{2}(\mathbf{q}_{T} - \mathbf{k}_{T} - \mathbf{p}_{T}) \frac{Q_{T}^{2}}{M^{2}}(\mathbf{k}_{T} \cdot \mathbf{p}_{T})$$

$$\times f(x_{1}, \mathbf{k}_{T}^{2}) f(x_{2}, \mathbf{p}_{T}^{2})$$

$$= \frac{1}{M^{2}} \int d^{2}\mathbf{k}_{T} d^{2}\mathbf{p}_{T}(\mathbf{k}_{T} + \mathbf{p}_{T})^{2}\mathbf{k}_{T} \cdot \mathbf{p}_{T}$$

$$\times f(x_{1}, \mathbf{k}_{T}^{2}) f(x_{2}, \mathbf{p}_{T}^{2})$$

$$= \frac{2}{M^{2}} \int d^{2}\mathbf{k}_{T} d^{2}\mathbf{p}_{T}(\mathbf{k}_{T} \cdot \mathbf{p}_{T})^{2} f(x_{1}, \mathbf{k}_{T}^{2}) f(x_{2}, \mathbf{p}_{T}^{2})$$

$$= \frac{2}{M^{2}} \int d^{2}\mathbf{k}_{T} d^{2}\mathbf{p}_{T}(\mathbf{k}_{T}^{2} \mathbf{p}_{T}^{12} + \mathbf{k}_{T}^{2} \mathbf{p}_{T}^{2})$$

$$\times f(x_{1}, \mathbf{k}_{T}^{2}) f(x_{2}, \mathbf{p}_{T}^{2})$$

$$\times f(x_{1}, \mathbf{k}_{T}^{2}) f(x_{2}, \mathbf{p}_{T}^{2})$$

$$= 4M^{2} f^{(1)}(x_{1}) f^{(1)}(x_{2}). \qquad (A.1)$$

$$\int d^{2}\mathbf{k}_{T}d^{2}\mathbf{p}_{T}\delta^{2}(\mathbf{q}_{T}-\mathbf{k}_{T}-\mathbf{p}_{T})\frac{Q_{T}^{2}}{4M^{2}}\hat{\mathbf{h}}\cdot\mathbf{k}_{T}\hat{\mathbf{h}}\cdot\mathbf{p}_{T}$$

$$\times f(x_{1},\mathbf{k}_{T}^{2})f(x_{2},\mathbf{p}_{T}^{2})$$

$$=\frac{1}{M^{2}}\int d^{2}\mathbf{k}_{T}d^{2}\mathbf{p}_{T}(\mathbf{k}_{T}+\mathbf{p}_{T})\cdot\mathbf{k}_{T}(\mathbf{k}_{T}+\mathbf{p}_{T})\cdot\mathbf{p}_{T}$$

$$\times f(x_{1},\mathbf{k}_{T}^{2})f(x_{2},\mathbf{p}_{T}^{2})$$

$$=\frac{1}{M^{2}}\int d^{2}\mathbf{k}_{T}d^{2}\mathbf{p}_{T}(\mathbf{p}_{T}^{2}\mathbf{k}_{T}^{2}+(\mathbf{k}_{T}\cdot\mathbf{p}_{T})^{2})$$

$$\times f(x_{1},\mathbf{k}_{T}^{2})f(x_{2},\mathbf{p}_{T}^{2})$$

$$= 6M^{2}f^{(1)}(x_{1})f^{(1)}(x_{2}) \qquad (A.2)$$

In the above integrals, the terms containing odd numbers of  $\mathbf{k}_T^i$  or  $\mathbf{p}_T^i$  vanish after being integrated over  $\mathbf{k}_T$  or  $\mathbf{p}_T$ .

- [1] D. Sivers, Phys. Rev. D 41, 83 (1990); 43, 261 (1991).
- [2] D.L.Adams *et.al*, Phys. Lett. B **261**, 201 (1991); **264**, 462 (1991); Z. Phys. C **56**, 181 (1992).
- [3] M. Anselmino, M. Boglione, and F. Murgia, Phys. Lett. B 362, 164 (1995).
- [4] D. Boer and P.J. Mulders, Phys. Rev. D 57, 5780 (1998).
- [5] J.C. Collins, Nucl. Phys. **B396**, 161 (1993).
- [6] S.J. Brodsky, D.S. Hwang, and I. Schmidt, Phys. Lett. B 530, 99 (2002); Nucl. Phys. B642, 344 (2002).
- [7] J.C. Collins, Phys. Lett. B 536, 43 (2002).
- [8] X. Ji and F. Yuan, Phys. Lett. B 543, 66 (2002); A.V. Belitsky, X. Ji, and F. Yuan, Nucl. Phys. B656, 165 (2003).
- [9] D. Boer, P.J. Mulders, and F. Pijlman, Nucl. Phys. B667, 201 (2003).
- [10] N. C. Makins (HERMES Collaboration), Transversity Workshop, Athens, Greece; R. Seidl (HERMES Col-

laboration), Proc. of DIS'2004, Štrbské Pleso, Slovakia, 2004; I.M. Gregor (HERMES Collaboration), Acta Phys. Polon. B **36**, 209 (2005).

- [11] A. Airapetian *et al.* (HERMES Collaboration), Phys. Rev. Lett. **94**, 012002 (2005).
- [12] M. Diefenthaler (HERMES Collaboration), in Proceedings of DIS 2005, Madison, Wisconsin (USA), hep-ex/0507013.
- [13] V.Yu. Alexakhin *et al.* (COMPASS Collaboration), Phys. Rev. Lett. **94** 202002 (2005).
- [14] E.S. Ageev et al. (COMPASS Collaboration), hep-ex/0610068.
- [15] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, and A. Prokudin, Phys. Rev. D 71, 074006 (2005).
- [16] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian,

F. Murgia, and A. Prokudin, Phys. Rev. D **72**, 094007 (2005); Erratum-ibid. D **72**, 099903 (2005).

- [17] A.V. Efremov, K. Goeke, S. Menzel, A. Metz and P. Schweitzer, Phys. Lett. B 612, 233 (2005).
- [18] J.C. Collins, A.V. Efremov, K. Goeke, S. Menzel, A. Metz and P. Schweitzer, Phys. Rev. D 73, 014021 (2006).
- [19] W. Vogelsang and F. Yuan, Phys. Rev. D 72, (2005) 054028.
- [20] X. Ji, J.P. Ma, and F. Yuan, Phys. Rev. D 71, 034005 (2004); Phys. Lett. B 597 (2004) 299.
- [21] J.C. Collins and A. Metz, Phys. Rev. Lett. 93, 252001 (2004).
- [22] For the comparison for these extrations, see M. Anselmino arXiv:hep-ph/0511017.
- [23] R.L. Jaffe and X.-D. Ji, Nucl. Phys. B375, 527 (1992);
  X.-D. Ji, Phys. Lett. B284, 137 (1992); W. Vogelsang and A. Weber, Phys. Rev. D 48, 2073 (1993); V. Barone, T. Calarco, and A. Drago, Phys. Rev. D 56, 527 (1997);
  O. Martin, A. Schafer, M. Stratmann, and W. Vogelsang, Phys. Rev. D 60, 117502 (1999); J. Soffer, M. Stratmann, and W. Vogelsang, Phys. Rev. D 65, 114024 (2002); V. Barone et. al., Phys. Lett. B 639, 483 (2006).
- [24] For a review on tranverse polarization phenomena, see V. Barone, A. Drago, and P.G. Ratcliffe, Phys. Rep. 359, 1 (2002).
- [25] J.P. Ralston and D.E. Soper, Nucl. Phys. B152 ,109 (1979).
- [26] R.D. Tangerman and P.J. Mulders, Phys. Rev. D 51, 3357 (1995); hep-ph/9408305.
- [27] D. Boer, Phys. Rev. D 60, 014012 (1999).
- [28] A. Kotzinian and P.J. Mulders, Phys. Rev. D 54, 1229 (1996).
- [29] P.J. Mulders and R.D. Tangerman, Nucl. Phys. B461, 197 (1996); B484, 538(E) (1997).
- [30] S. Wandzura and F. Wilczek, Phys. Lett. B 72, 195 (1977).
- [31] J. Levelt and P.J. Mulders, Phys. Rev. D 49, 96 (1994).
- [32] A. Kotzinian, Nucl. Phys. B441, 234 (1995).

- [33] J.C. Collins and D.E. Soper, Phys. Rev. D 16, 2219 (1977).
- [34] G. R. Goldstein, L. Gamberg, Talk given at 31st International Conference on High Energy Physics (ICHEP 2002), Amsterdam, The Netherlands, hep-ph/0209085; hep-ph/0506127.
- [35] Z. Lu and B.-Q. Ma, Phys. Rev. D 70, (2004) 094044;
   Phys. Lett. B 615, (2005) 200; Z. Lu, B.-Q. Ma, and I. Schmidt, Phys. Lett. B 639, 494 (2006).
- [36] D. Boer, S.J. Brodsky, and D.S. Hwang, Phys. Rev. D 67, 054003 (2003); D. Boer, A. Brandenburg, O. Nachtmann, and A. Utermann, Eur. Phys. J. C 40, 55 (2005).
- [37] S. Falciano *et al.* (NA10 Collaboration), Z. Phys. C **31**, 513 (1986); M. Guanziroli *et al.* (NA10 Collaboration), Z. Phys. C **37**, 545 (1988).
- [38] J.S. Conway *et al.* (E615 Collaboration), Phys. Rev. D 39, 92 (1989).
- [39] A. Kotzinian and P.J. Mulders, Phys. Lett. B 406, 373 (1997).
- [40] PAX Collaboration, V. Barone et al., hep-ex/0505054.
- [41] A. D. Martin, R.G. Roberts, W.J. Stirling, and R. S. Thorne, Phys. Lett. B 531, 216 (2002).
- [42] J. C. Collins, A. V. Efremov, K. Goeke, M. Grosse Perdekamp, S. Menzel, B. Meredith, A. Metz, and P. Schweitzer, Phys. Rev. D 73, 094023 (2006); hep-ph/0510342.
- [43] A. Kotzinian, B. Parsamyan, and A. Prokudin, Phys. Rev. D 73, 114017 (2006).
- [44] M. Gluck, E. Reya, M. Stratmann, and W. Vogelsang, Phys. Rev. D 63, 094005 (2001).
- [45] M. Gluck, E. Reya, and A. Vogt, Eur. Phys. J. C 5, 461 (1998).
- [46] There is some work on the  $Q^2$  evolution of the  $\mathbf{k}_T$ moment of  $\mathbf{k}_T$ -dependent distributions held in the Large  $N_c$  limit, see: A.A. Henneman, D. Boer, and P.J. Mulders, Nucl. Phys. **B620**, 331 (2002).