

Effective gravity from a quantum gauge theory in Euclidean space-time

R. F. Sobreiro* , V. J. Vasquez Otoya†

*CBPF, Centro Brasileiro de Pesquisas Físicas,
Rua Xavier Sigaud 150, 22290-180 Urca, Rio de Janeiro, Brasil*

Abstract

We consider a $SO(d)$ gauge theory in an Euclidean d -dimensional space-time, which is known to be renormalizable to all orders in perturbation theory for $2 \leq d \leq 4$. Then, with the help of a space-time representation of the gauge group, the gauge theory is mapped into a curved space-time with linear connection. Further, in that mapping the gauge field plays the role of the linear connection of the curved space-time and an effective metric tensor arises naturally from the mapping. The obtained action, being quadratic in the Riemann-Christoffel tensor, at a first sight, spoils a gravity interpretation of the model. Thus, we provide a sketch of a mechanism that breaks the $SO(d)$ color invariance and generates the Einstein-Hilbert term, as well as a cosmological constant term, allowing an interpretation of the model as a modified gravity in the Palatini formalism. In that sense, gravity can be visualized as an effective classical theory, originated from a well defined quantum gauge theory. We also show that, in the four dimensional case, two possibilities for particular solutions of the field equations are the de Sitter and Anti de Sitter space-times.

1 Introduction

Since the beginning of the last century many efforts to give gravity a quantum description have been developed. However, a complete consistent quantum theory of gravity still lacks. In particular, attempts to quantize general relativity have failed since it is a nonrenormalizable theory where higher order derivative counterterms already appear at one-loop quantum corrections [1, 2, 3]. In order to solve this problem gravity has been described by several alternative approaches. Some of them are: The higher derivative gravities [4]; The loop quantum gravity (LQG) [5, 6]; The Einstein-Cartan gauge formalism [7, 8]; The interpretation that gravity would be an effective theory originated from more general theories such as superstring theories [9, 10]. It is important to highlight though that it is not our intention to give a complete description on those issues, for that we refer to the cited bibliography and references therein.

To discuss quantum aspects of gravity, let us take under consideration the pure quantum mechanical point of view. It is reasonably unanimous that Quantum Physics is the fundamental

*sobreiro@cbpf.br

†vjose@cbpf.br

theory while Classical Physics is a particular case of the quantum theory. From this point of view, quantization methods employed to achieve a quantum model from a classical model are just *ansätze* or algorithms that might not always work. A renowned example is a general dissipative system. This is the main quantum mechanics point, implying that the word *quantize* should not be taken as a fundamental principle [11, 12]. In quantum field theory, the picture is not different. The quantization methods are just *ansätze* successfully utilized to obtain a quantum theory from a classical model. The starting point would be the fundamental quantum theory from where, in some suitable limits, one recovers the classical equivalent theory. This might be the reason why gravity resists to be quantized by the usual quantization methods, requiring then a more fundamental understanding of quantum theory.

Thus, based on the fundamental character of the quantum theory and on the success of gauge theories when utilized to describe the fundamental interactions, we opt, in this work, for the idea that gravity can be visualized as an effective theory originated from a simple, non-Abelian, four-dimensional Euclidean quantum gauge field theory. To be more specific, we start our investigation on the proposed idea by considering a $SO(d)$ gauge theory in an Euclidean d -dimensional space-time*. Under this prescription the graviton is simply defined as the gauge field, A_i^a . We recall that an essential ingredient of this description is that there is no curved space in the quantum sector of the theory. In fact, the Euclidean space-time is a motivation to define the path integral allowing a useful consistent scenario to perform quantum computations.

Once we have established the properties of the theory at quantum level, we can relate it with a curved space-time through a mapping from the Euclidean gauge theory to a dynamical space-time with $GL(d, \mathbb{R})$ symmetry. For that, one can use a representation of the $SO(d)$ gauge group borrowing the Euclidean space-time indices, then the group index a can be represented by a pair of antisymmetric space-time indices, $A_i^{jk} = -A_i^{kj}$. In this mapping the Euclidean space is identified with a curved space through transformation matrices e_i^μ , which are related to an effective metric tensor g . The gauge field is mapped into a linear connection of the space-time $A_{ij}^k \mapsto \Gamma_{\mu\nu}^\alpha$ which is independent of the metric tensor. Thus, the Euclidean gauge theory is mapped into a dynamical space-time in the Palatini formalism. The original action, $\int \mathbf{F}^2$, is mapped into an action quadratic in the Riemann tensor $\int \sqrt{g} \mathbf{R} \cdot \mathbf{R}$.

In order to relate the dynamical curved space with gravity, it would be useful to generate the Einstein-Hilbert (EH) term in the action. However, the term $\mathbf{R} \cdot \mathbf{R}$ still carries the global gauge symmetry of the original gauge theory while the EH term, $R = R_{\mu\nu}^{\mu\nu} = \text{Tr } \mathbf{R}$, breaks color invariance. Thus, we provide a sketch of a color symmetry breaking mechanism which generates the EH term. The first feature of the color symmetry breaking is the appearance of physical spin-2 excitations originated from the antisymmetry structure of the color indices. Before the breaking, those excitations were hidden behind the color symmetry and thus were unphysical. It is important to know that this mechanism is employed in the quantum sector of the gauge theory, *i.e.*, the mapping is only performed after the mechanism is worked out. This detail ensures the existence of a framework for well defined quantum computations, as an Euclidean QFT is. An extra feature of the mechanism is that a cosmological constant (CC) appears naturally in the resulting action. We remark at this point, and will be pointed further, that this mechanism is merely a possible sketch. In fact, further investigation on the formalization of the present

*Here we are considering a d -dimensional space just for generality purposes, the main focus being the case $d = 4$.

mechanism (or other) is needed and are currently under investigation [13].

Finally, once the previously mechanism is worked out, the mapping can be performed. The obtained action for the dynamical space-time is now constituted by a quadratic term in the Riemann tensor, the EH term and the CC term. This new action can now be related with gravity, in the Palatini formalism. In the particular case of four dimensions, the equations of motion provide the de Sitter space-time (dS_4) or the Anti de Sitter (AdS_4) as the fundamental space-time.

This work is organized as follows: In section 2, we will show how a $SO(d)$ gauge model, in d -dimensional Euclidean space-time, can be mapped into a dynamical curved space. The resulting dynamical space-time enjoys $GL(d, \mathbb{R})$ symmetry and $SO(d)$ gauge symmetry (color invariance). In section 3, we will argue that, to obtain an effective gravity theory from the previous method, a color symmetry breaking mechanism is required. In section 4, a color symmetry breaking mechanism will be sketched and, due to this mechanism, the EH term is generated together with the CC. Still in this section, we will discuss the resulting classical field equations. Finally, the conclusions will be displayed in section 5.

2 From $SO(d)$ gauge theory to dynamical curved space-time

We start this section with a small review of the definitions, notations and properties of the $SO(d)$ gauge theory.

2.1 Generalities of the $SO(d)$ gauge theory

Let us consider the following principal bundle

$$\mathcal{P} \equiv \{SO(d) ; \mathbb{R}^d ; \mathcal{A}\} , \quad (1)$$

where $SO(d)$ is the group of orthogonal matrices characterizing the fiber and the structure group of the fiber. There are $D = d(d - 1)/2$ generators of this group which we label by λ^a , $a \in \{1, 2, \dots, D\}$. The algebra is a typical Lie algebra,

$$[\lambda^a, \lambda^b] = f^{abc} \lambda^c , \quad (2)$$

where f^{abc} represents the structure constants of the group. Also, we denote the elements of the group by $u = e^{\omega^a \lambda^a}$. The total space of the principal bundle (1) is chosen to be an Euclidean d -dimensional space, \mathbb{R}^d , which enjoys a global $O(d)$ symmetry. The set \mathcal{A} is the space of the algebra-valued connections, the base space[†],

$$A_i = A_i^a \lambda^a . \quad (3)$$

The homeomorphisms of the principal bundle are characterized by the gauge transformations

$$A_i \rightarrow A_i + u^\dagger D_i u , \quad (4)$$

[†]The first part of the Latin alphabet, $\{a, b, c, \dots, h\}$, labels the group indices while the second part, $\{i, j, k, \dots, z\}$, will be used to label the space-time coordinates.

where the covariant derivative is just

$$D_i \cdot = \partial_i \cdot + [A_i, \cdot] . \quad (5)$$

Infinitesimal gauge variations are then defined as

$$\delta A_i = D_i \omega . \quad (6)$$

The gauge invariant action is

$$S_{\text{YM}} = \frac{1}{4\kappa^2} \text{Tr} \int d^d x F_{ij} F^{ij} , \quad (7)$$

where the field strength (fiber curvature) is given by

$$F_{ij} = F_{ij}^a \lambda^a = \partial_i A_j - \partial_j A_i + [A_i, A_j] . \quad (8)$$

The quantity κ is the coupling constant.

To quantize the model a gauge fixing is usually needed. Here we choose the simplest covariant gauge, the Landau gauge $\partial^i A_i = 0$. With that purpose we add to the gauge invariant action (7) a gauge fixing term

$$S = S_{\text{YM}} + S_{\text{gf}} , \quad (9)$$

where

$$S_{\text{gf}} = \text{Tr} \int d^d x (b \partial^i A_i + \bar{c} \partial^i D_i c) . \quad (10)$$

The fields $c = c^a \lambda^a$ and $\bar{c} = \bar{c}^a \lambda^a$ are, respectively, the Faddeev-Popov ghost and antighost fields while $b = b^a \lambda^a$ is the Lautrup-Nakanishi field which plays the role of a Lagrange multiplier enforcing the Landau gauge condition. The action (9) enjoys renormalizability, *i.e.*, the model is consistent at quantum level. In fact, as discussed in Ap.A, the renormalizability of action (9) can be established to all orders in perturbation theory for $2 \leq d \leq 4$.

After gauge fixing, a global $SO(d)$ gauge symmetry survives, the so called color invariance. This invariance is characterized by the non-observational character of the group indices. In terms of a Ward identity, color invariance is described by

$$\int d^d x \left(\left[A_i, \frac{\delta S}{\delta A_i} \right] + \left[c, \frac{\delta S}{\delta c} \right] + \left[\bar{c}, \frac{\delta S}{\delta \bar{c}} \right] + \left[b, \frac{\delta S}{\delta b} \right] \right) = 0 . \quad (11)$$

2.2 Space-time representation

Unless the contrary is said, we will not consider the gauge fixing term for the rest of the article. Thus, from now on, we are dealing with the gauge invariant action (7) and global color invariance is now described by

$$\int d^d x \left[A_i, \frac{\delta S_{\text{YM}}}{\delta A_i} \right] = 0 . \quad (12)$$

First, we will explore the fact that the $SO(d)$ group, in an Euclidean space-time, can borrow the space-time structure to fix a representation of the group. For that, we observe that the

dimension of the total space \mathbb{R}^d coincides with the Casimir d of the group $SO(d)$, which means that the generators can be represented as a set of $(d \times d)$ -matrices. Also, the group dimension, D , coincides with the number of independent elements of an antisymmetric $(d \times d)$ -matrix. Those properties allow a unique representation of the gauge group where the generators are in the form of matrices labeled by space-time indices,

$$\begin{aligned}\lambda^a &\mapsto \lambda^{ij}, \\ f^{abc} &\mapsto f^{ijklmn},\end{aligned}\tag{13}$$

where

$$\begin{aligned}\lambda^{ij} &= -\lambda^{ji}, \\ f^{ijklmn} &= -\frac{1}{2} \left[\left(\delta^{il} \delta^{mj} - \delta^{jl} \delta^{mi} \right) \delta^{kn} + \left(\delta^{jk} \delta^{im} - \delta^{ik} \delta^{mj} \right) \delta^{ln} \right].\end{aligned}\tag{14}$$

Thus

$$\begin{aligned}A_i^a \lambda^a &= A_{ij}^k \lambda_k^j, \\ F_{ij}^a \lambda^a &= F_{ijk}^l \lambda_l^k.\end{aligned}\tag{15}$$

In the above described representation the action (7) reads

$$S_{\text{YM}} = \frac{1}{4\kappa^2} \int d^d x F_{ijk}^l F^{ijk}_l,\tag{16}$$

where

$$F_{ijk}^l = \partial_i A_{jk}^l - \partial_j A_{ik}^l + A_{im}^l A_{jk}^m - A_{jm}^l A_{ik}^m.\tag{17}$$

Thus, we are dealing with a gauge theory, in an Euclidean space-time with a representation where the color indices are represented by space-time indices.

As previously pointed, the color invariance has a fundamental role in this article, therefore, for further use, let us write the color symmetry Ward identity in the space-time representation

$$\int d^d x \left(A_{ki}^n \delta_{mj} - A_{kj}^n \delta_{mi} \right) \frac{\delta S_{\text{YM}}}{\delta A_{km}^n} = 0.\tag{18}$$

2.3 Pseudo-vielbein and curved space-time

Looking at the field strength expression in (17), one can recognize an extreme similarity with a Riemann-Christoffel tensor of a curved space-time. Notwithstanding, one can think in absorb the gauge field as a geometrical structure of the space-time and end up in an effective geometrical theory, completely equivalent to a gauge theory. In fact, we can perform a mapping from the Euclidean space to a curved space[‡],

$$\{x_i\} \mapsto \{x_\mu, x^\mu\},\tag{19}$$

[‡]The Greek indices label the coordinates of the curved space time.

by introducing the pseudo-vielbein and its inverse

$$\begin{aligned} e^i_\mu &= \frac{\partial x^i}{\partial x^\mu}, \\ \bar{e}^\mu_i &= \frac{\partial x^\mu}{\partial x^i}. \end{aligned} \quad (20)$$

Thus

$$\begin{aligned} dx^i &= e^i_\mu dx^\mu, \\ dx^\mu &= \bar{e}^\mu_i dx^i. \end{aligned} \quad (21)$$

The name *pseudo-vielbein*[§] is here employed due to the fact that e is not the vielbein of the EC theory of gravity [7, 8], as it will become evident in what follows. In that sense, the flat space is completely independent of the curved space. Differently of the EC formalism the p-vielbein does not define a reference frame which travels tangently on the curved space. In here, the p-vielbein is a point-to-point identification of the two spaces which have no direct geometrical relation. We will return to this issue soon.

The requirement of metric preservation in both spaces implies on the existence of a symmetric metric tensor $\{g_{\mu\nu}, g^{\mu\nu}\}$, in fact the following relations hold,

$$\begin{aligned} e^i_\mu e_{i\nu} &= g_{\mu\nu}, \\ \bar{e}^\mu_i \bar{e}^j_\mu &= \delta_i^j, \\ e^\mu_i \bar{e}^j_\mu &= \delta_i^j, \\ e^\mu_i \bar{e}^i_\nu &= \delta^\mu_\nu, \end{aligned} \quad (22)$$

where the first two are related to the metric invariance in both spaces. In that sense the p-vielbein is equivalent to the usual vielbein, which justifies the name. However, here, e is essentially a mapping field.

For the ordinary derivative and for the gauge field the mapping occurs according to

$$\begin{aligned} \partial_i &= \bar{e}^\mu_i \partial_\mu, \\ A_{ij}{}^k &= \bar{e}^\mu_i \bar{e}^\nu_j e^k_\alpha \Gamma_{\mu\nu}{}^\alpha - \bar{e}^\mu_i e^k_\nu \partial_\mu \bar{e}^\nu_j. \end{aligned} \quad (23)$$

The quantity Γ is identified with the connection of the curved space.

Now we return to the geometrical properties of the p-vielbein. In the EC formalism the vielbein connects the curved space with a tangent space. This association implies that the partial derivatives do not commute in flat space while the derivatives in curved space do commute. In our formalism, the original flat space-time is not a tangent space of the curved space. They are essentially independent spaces, modulo the mapping through e^μ_i . Thus, the partial derivative commutes in Euclidean space as well as in curved space,

$$\begin{aligned} [\partial_i, \partial_j] &= 0, \\ [\partial_\mu, \partial_\nu] &= 0. \end{aligned} \quad (24)$$

[§]For economical purposes we will call the *pseudo-vielbein* simply by *p-vielbein*.

Those commutation rules imply that the non-holonomicity coefficients of both spaces vanish. Consequently, the p-vielbein behaves exactly as a coordinate matrix transformation.

Finally, we wish to let clear that there is no *a priori* proof of the existence of consistent (invertible) transformations e_i^μ . For now, we naively assume that e_i^μ exists and is invertible. Nevertheless, this assumption will be confirmed *a posteriori* when we solve the field equations in Sec.4.4. Essentially, a kind of proof of the existence of an invertible e_i^μ would rely on solving the field equations in the curved space-time and find an invertible metric tensor as a consistent solution.

2.4 Action for the dynamical curved space

A problem to be faced at this point is that we must generate a scalar action on the curved space, which means that F , in the curved space, should be a tensor. Substituting (23) in (17) and making use of the relations (24) we find

$$F_{ijk}{}^l = \bar{e}_i^\mu \bar{e}_j^\nu \bar{e}_k^\alpha e_l^\beta R_{\mu\nu\alpha}{}^\beta, \quad (25)$$

where $R_{\mu\nu\alpha}{}^\beta$ is a four rank tensor in the form

$$R_{\mu\nu\alpha}{}^\beta = \partial_\mu \Gamma_{\nu\alpha}{}^\beta - \partial_\nu \Gamma_{\mu\alpha}{}^\beta + \Gamma_{\mu\gamma}{}^\beta \Gamma_{\nu\alpha}{}^\gamma - \Gamma_{\nu\gamma}{}^\beta \Gamma_{\mu\alpha}{}^\gamma, \quad (26)$$

recognized as the Riemann-Christoffel tensor. At the action (16), this transformation results in

$$S_{YM}^e = \frac{1}{4\kappa^2} \int d^d x e R_{\mu\nu\alpha}{}^\beta R^{\mu\nu\alpha}{}_\beta, \quad (27)$$

where $e = \det e_\mu^i = \sqrt{-\det g_{\mu\nu}}$ comes from the Jacobian of the transformation (21). The action (27) displays color invariance under the $SO(d)$ group characterized by the following functional identity

$$\int d^d x (\Gamma_{\mu\nu}{}^\alpha \delta_{\beta\gamma} - \Gamma_{\mu\gamma}{}^\alpha \delta_{\beta\nu}) \frac{\delta S_{YM}}{\delta \Gamma_{\mu\beta}{}^\alpha} = 0. \quad (28)$$

Further, it is clear that (27) enjoys $GL(d, \mathbb{R})$ symmetry, not at the gauge sector, but in the curved space-time itself. What does it mean? It means that we started from a renormalizable gauge theory. Due to the possibility of the space-time representation the classical theory is then mapped into a curved space with linear connection. The gauge field turns into the linear connection and an effective metric tensor arises from the p-vielbein.

2.5 On-shell field equations

The action (27) describes the dynamics of a curved space-time. The fundamental field being the connection Γ . The explicit form of Γ , which dictates the class of geometry we are dealing with, is unknown till now. Thus, at classical level, the field equations of the connection are just

$$- \mathcal{D}_\mu R^{\mu\nu\alpha}{}_\beta - 2 \left[\frac{d}{2} Q_\mu + T_{\mu\kappa}{}^\kappa \right] R^{\mu\nu\alpha}{}_\beta + T_{\mu\kappa}{}^\nu R^{\mu\kappa\alpha}{}_\beta = 0, \quad (29)$$

where

$$Q_\mu = \frac{1}{d} g^{\alpha\beta} \mathcal{D}_\mu g_{\alpha\beta}, \quad (30)$$

is the Weyl convector and

$$T_{\mu\nu}{}^\alpha = \frac{1}{2} (\Gamma_{\mu\nu}{}^\alpha - \Gamma_{\nu\mu}{}^\alpha) , \quad (31)$$

is the torsion tensor. The covariant derivative \mathcal{D} is the usual covariant derivative of a general curved space-time with linear connection Γ , which is different of the definition of the covariant derivative (5).

Evidently, the p-vielbein constitutes a field and it will have its respective field equation. However, it is important to keep in mind that the fundamental field here is the connection which has well defined quantum properties. The p-vielbein is a pure classical field. The classical field equations of the connection are the usual classical limit selecting the main contribution to the path integral. On the other hand, the field equations for the p-vielbein are interpreted as a minimization principle related to the vacuum energy stability. In fact, there are several possible solutions to e . Nevertheless, through the field equations, we select the p-vielbein which makes the vacuum energy stable. In principle, it is just a matter of minimal principles. Thus, the variation of the action (27) with respect to the metric tensor provides[¶]

$$\frac{1}{2} g^{\mu\nu} R^{\sigma\rho\alpha\beta} R_{\sigma\rho\alpha\beta} - R^\mu{}_{\alpha\beta\gamma} R^{\nu\alpha\beta\gamma} - R_{\alpha\beta\gamma}{}^\mu R^{\alpha\beta\gamma\nu} = 0 . \quad (32)$$

Now, contracting the equation for the p-vielbein with $g_{\mu\nu}$ one finds

$$\frac{(d-4)}{2} R^{\sigma\rho\alpha\beta} R_{\sigma\rho\alpha\beta} = 0 . \quad (33)$$

We recall that the geometry involved is determined by the field equations (29) and (32). The solution of those equations will provide the geometrical properties of the curved space. *The curved space is simply a space with single linear connection*, which generalizes several geometries such as the Riemannian geometry and the Einstein-Cartan geometry. From that point of view, we are dealing with the Palatini formalism, the so called metric-affine formulation. The difference is that action (27) no longer describes gravity. In the next section, we start the discussion whether the presented formalism can be related to a gravity theory or not.

Exploiting, the field equations (29), (32) and (33) for $d \neq 4$, we observe that equation (33) implies that $R^{\sigma\rho\alpha\beta} = 0$, which is also a solution of (29) and (32). In other words, the case $d \neq 4$ is trivial when matter is not being considered.

For the case $d = 4$ equation (33) is satisfied for any non-vanishing curvature. In that case, the nontrivial solutions of (29) are allowed and nontrivial curved spaces show up. Obviously, this phenomenon occurs due to the nonlinear character of the action (27). Physically, this character is associated to the self interaction of Γ .

3 Intermediate discussion

We wish to relate the above described theory with gravity through the following point of view:

[¶]The same equation can be achieved by varying the action (27) with respect to the p-vielbein.

Quantum gravity would be described by the action (7) in Euclidean space-time, which is unitary and renormalizable. In that hypothesis, the gauge field (3) is interpreted as the graviton. Thus, at very high energies gravity is simply a renormalizable quantum gauge theory in Euclidean space-time, where a spin-1 gauge field plays the role of the graviton.

On the other hand, at low energies, the theory can be described by a deformed space-time where the gauge field is visualized as a linear connection. In that regime an effective metric tensor arises due to the presence of the p -vielbein. Thus, action (27) can be regarded as a kind of effective theory of gravity. In that sense, classical gravity is no longer a fundamental theory. The deformation of the space-time occurs from quantum effects of a gauge theory as well as the covariance principle of general relativity.

We remark that the present approach to treat gravity is essentially different of the Einstein-Cartan formalism [7, 8]. In the EC formalism there is an initial assumption of the existence of a curved space as the fundamental space. The tangent flat space is used in order to define the spin connection which allows the introduction of fermion fields with the help of the vielbein. However, the formalism still lacks renormalizability once the EH term is taken as the fundamental action for gravity. In our prescription the fundamental theory is a consistent Euclidean quantum gauge theory of spin-1 fields which, for a certain limit, is equivalent to a pure dynamical curved space-time where no reference to a tangent space neither spin connection is made. Thus, the curved space is a direct consequence of the dynamics of the gauge theory and not the contrary. Furthermore, the original flat space is not a tangent space of the curved space, as it is evident from (24).

This idea turns on essentially three issues to be faced. First of all: *Where is the EH term $\propto R = R^{\mu\nu}{}_{\nu\mu}$, which ensures the relation of a theory with gravity?* It is easy to see that this question hits directly the color symmetry of action (27). In fact, the EH term would require contraction between group indices and space-time indices, which breaks color invariance.

The second point concerns the spin of the physical excitations. At quantum level, the physical excitations carries spin-1, as any gauge theory of vector connection. Curiously, in the space-time representation, spin-2 excitations also show up, due to the identification of the group indices with the space-time indices. However, due to color symmetry, those are nonphysical excitations, which will never be observed. Thus, the question of the spin-2 physical excitations persists: *Would color invariance forbid the existence of spin-2 excitations at the physical sector of the theory?*

Finally, the equivalence between actions (7) and (27), for now, due to color invariance, seems to be just a point of view, *i.e.*, is just another way to look at the nature. It happens because color invariance forbids the observational character of color indices. For example, in $R_{\mu\nu\alpha\beta}$ the last two indices are related to color invariance. Then, if one wishes to measure, directly, the curvature of the space time, it would never been successful, since the last two indices are non-observable. Therefore, the third question is: *Which physical effect pushes the theory to the action (27), characterizing the low energy regime of (7) in terms of an observable curved space-time?*

Those questions are then closely related to the color symmetry of action (27). Further, a color symmetry breaking mechanism seems to be necessary if one wishes to describe gravity with the above program. With that purpose, in the next section, we will provide a sketch of

a possible color symmetry breaking mechanism. We will show that this mechanism can generate the physical spin-2 excitations as well as the EH term. Further, the same mechanism would dictate the vacuum of the theory, allowing a physical motivation for the mappings (21) and (23).

We would like to recall, at this point, that the above statements and what follows next only work for $2 \leq d \leq 4$, as is required by the renormalizability condition of the theory.

4 Color symmetry breaking and gravity

For the beginning of this section we forget the dynamical curved space-time equivalent theory and turn back at the Euclidean theory (1).

4.1 A background field

Let us consider, again, the action (16). We suppose the existence of a background field Υ which generates the following field strength

$$F_{ijk}{}^l(\Upsilon) = m^2(\delta_i^l \delta_{jk} - \delta_{ik} \delta_j^l), \quad (34)$$

where m has dimension of a mass. It is evident that the field strength renormalizes according to the renormalization of the gauge field and the coupling constant. Thus, a field strength for a background would require that m renormalizes nontrivially according to the renormalization of the gauge field and the coupling constant. In fact, according to Ap.B, the renormalization factor of m^2 is not independent and is given by

$$Z_{m^2} = Z_A Z_\kappa^2. \quad (35)$$

The condition (34) implies that Υ is a solution of the classical field equations if one requires the following extra condition

$$\Upsilon_i{}^{li} \delta_{jk} - \Upsilon_{ik}{}^i \delta_j^l + \Upsilon_{kj}{}^l - \Upsilon_k{}^l{}_j = 0. \quad (36)$$

It is not difficult to see that equation (34) suggests that Υ is a singular configuration. For that, we write Υ as a pure gauge, which is just a gauge equivalent of the vacuum configuration $\Upsilon' = 0$,

$$\Upsilon_{ik}{}^l = \partial_i \theta_k^l. \quad (37)$$

And then, considering the linear approximation of equation (34), one finds

$$(\partial_i \partial_j - \partial_j \partial_i) \theta_k^l = m^2(\delta_i^l \delta_{jk} - \delta_{ik} \delta_j^l). \quad (38)$$

This equation establishes the singular nature of Υ since θ is clearly singular. An explicit example can be found in [15], where Υ is associated with flux tubes in an Abelian superconductor.

4.2 Color symmetry breaking

From (34) we see that Υ explicitly breaks color invariance since it mixes color and space-time indices. As a consequence it will turn the color indices into observable space-time indices. On the other hand, one might argue on the physical consequences of the background in the sense that the field strength, in the form (34), is an exclusive solution of the space-time representation. This means that this effect can be described exclusively in the space-time representation. To answer this question, we can take a look on the Noether current for global color symmetry,

$$j_{pij}(A) = \frac{\delta S_{YM}}{\delta(\partial_p A_{km}^n)} \delta_{(ij)} A_{km}^n, \quad (39)$$

where, from (18),

$$\delta_{(ij)} A_{km}^n = \delta_{mj} A_{ki}^n - \delta_{mi} A_{kj}^n. \quad (40)$$

Thus

$$j_{pij}(A) = A_{ki}^n F_{pkj}^n - A_{kj}^n F_{pki}^n. \quad (41)$$

For the background we have then

$$j_{pij}(\Upsilon) = m^2 \left(\Upsilon_{ki}^k \delta_{jp} - \Upsilon_{kj}^k \delta_{ip} + \Upsilon_{ijp} - \Upsilon_{jip} \right). \quad (42)$$

Making use of condition (36) we achieve, identically,

$$j_{pij}(\Upsilon) = 0. \quad (43)$$

Thus, the background configuration lies at the nonphysical sector of the theory. This means that the massless Goldstones states, associated with the color symmetry breaking, should decouple from the physical spectrum of the theory.

Obviously, considering the gauge fixing term, (10), our conclusion remains the same since the difference is just a BRST exact term

$$j_p^{ij}(\Upsilon) = s[\mathcal{F}(\Upsilon, A, \bar{c}, c, b)], \quad (44)$$

where \mathcal{F} is a functional of the fields and the background. In (44) the BRST variation of the background is defined according to

$$s\Upsilon_k^a = f^{abc} \Upsilon_k^b c^c, \quad (45)$$

or, in the space-time representation,

$$s\Upsilon_{ki}^j = \Upsilon_{ki}^m c_m^j - \Upsilon_{km}^j c_i^m. \quad (46)$$

For the other fields the BRST transformations are given in (68).

4.3 Effective action for the background

Now, we write the gauge field as a perturbation around the background

$$A_{ij}^k \rightarrow \Upsilon_{ij}^k + A_{ij}^k. \quad (47)$$

Thus,

$$F_{ijk}{}^l(A) \rightarrow F_{ijk}{}^l(A) + m^2(\delta_{ik}\delta_j^l - \delta_i^l\delta_{jk}) + \Upsilon_i{}^m{}_k A_{jm}{}^l - \Upsilon_{im}{}^l A_j{}^m{}_k - \Upsilon_j{}^m{}_k A_{im}{}^l + \Upsilon_{jm}{}^l A_i{}^m{}_k. \quad (48)$$

To avoid terms depending on the background in (48) and, consequently, in the resulting action, we write Υ as^{||}

$$\Upsilon_{ij}{}^k = \mathcal{F}_{ij}{}^k - D_{ijm}{}^{kl} h_l^m, \quad (49)$$

in such a way that Υ is fixed in order to maintain the relation (34) while \mathcal{F} and h are arbitrary and the covariant derivative is taken with respect the background itself, $D = D(\Upsilon)$. Thus, we can get rid of the Υ dependent term in (48) by making the smooth limit of the arbitrary functions $\mathcal{F} \leftrightarrow D \cdot h$ while $F(\Upsilon)$ remains fixed and Υ becomes small. Thus, we are taken the limit $\Upsilon \rightarrow 0$ while keeping the singular character of Υ . This trick might be interpreted as follows: A fixed background breaks the gauge invariance of the background field method [16]. However, in order to control this breaking while keeping some gauge freedom on the background, the fields \mathcal{F} and h were introduced in (49). As a consequence, the functions \mathcal{F} and h allow Υ to vary through a class of backgrounds that generates (34). Then, by making Υ as small as possible, expression (48) reads

$$F_{ijk}{}^l(A) \rightarrow F_{ijk}{}^l(A) + m^2(\delta_{ik}\delta_j^l - \delta_i^l\delta_{jk}). \quad (50)$$

And the action (16) now reads

$$S_{\text{eff}} = \frac{1}{4\kappa^2} \int d^d x \left[F_{ijk}{}^l F^{ijk}{}_l - 4m^2 F + 2d(d-1)m^4 \right]. \quad (51)$$

We have introduced a new parameter, m , as a free parameter. However, this parameter is not present in the starting action. Thus, it might exist a condition to fix this parameter to a physical consistent value. For instance, the mass parameter would be such that it characterizes the vacuum nature of the background. To do so, m would be fixed, in a self consistent way, by requiring minimal dependence in m of the vacuum energy,

$$\frac{\partial \mathcal{W}}{\partial m^2} = 0, \quad (52)$$

where the quantum action is defined as

$$e^{-\mathcal{W}} = \int DADbD\bar{c}Dc e^{-S_{\text{eff}} - S_{gf}}. \quad (53)$$

The gap equation (52) in fact reads

$$\langle F \rangle = d(d-1)m^2, \quad (54)$$

where $\langle F \rangle$ is the expectation value of F related to the functional (53). The gap equation (54) fixes m to a physical value m_* which stabilizes the vacuum. We also remark that this kind of gap equation (52) have been frequently used in Physics, in particular, in QCD where the gap equation (52) is imposed to find an optimal value for the mass gap in QCD [17]. Also in QCD, a similar gap equation is used to fix the so called Gribov parameter, associated with the improvement of the quantization of Yang-Mills theories [18, 19, 20].

^{||}Notice that the covariant derivative possess two color indices. Thus, together with the ordinary space-time index, there will be five indices in the space-time representation.

Once m_* is determined, the existence of the background turns out to be completely characterized by m_* and by the presence of the the color breaking term F in the action. Thus, after the computation of m_*^2 ,

$$S_{\text{eff}} = \frac{1}{4\kappa^2} \int d^d x \left[F_{ijk}{}^l F^{ijk}{}_l - 4m_*^2 F + 2d(d-1)m_*^4 \right]. \quad (55)$$

The theory is then described by the effective action (55). We notice that in the action (55) the parameters κ and m_* are assumed to be already fixed from usual loop quantum computations [21].

Notice that, due to color symmetry breaking mechanism, the group indices are transformed into observable space-time indices. Thus $A_{ij}{}^k$ would describe physical spin-1 excitations as well as spin-2 physical excitations.

4.4 Gravity

Since we have now an effective theory, where a kind of non-perturbative vacuum is consistently defined, we can perform the mapping (21) and (23), providing now

$$S_{\text{eff}}^e = \frac{1}{4\kappa^2} \int d^d x e \left[R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\beta} - 4m_*^2 R + 2d(d-1)m_*^4 \right], \quad (56)$$

where $R = R_{\mu\nu}{}^{\mu\nu}$.

The action (56) is $GL(d, \mathbb{R})$ invariant in the space-time sector, and, due to the presence of the EH term, we can associate it with gravity. In that sense, the mass parameter m_*^2 is related to both, Newton constant, G , and the cosmological constant, Λ , through

$$\begin{aligned} G &= \frac{\kappa^2}{16\pi m_*^2}, \\ \Lambda &= \frac{d(d-1)}{2} m_*^2. \end{aligned} \quad (57)$$

Thus,

$$S_{\text{eff}}^e = \frac{1}{16\pi G} \int d^d x e \left[\frac{d(d-1)}{8\Lambda} R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\beta} - R + \Lambda \right], \quad (58)$$

which is recognized as a generalization of the EH action in the Palatini formalism.

Let us derive the field equations of action (58). For the connection Γ , the field equations read

$$\begin{aligned} & - \mathcal{D}_\mu R^{\mu\nu\alpha}{}_\beta - 2 \left[\frac{d}{2} Q_\mu + T_{\mu\kappa}{}^\kappa \right] R^{\mu\nu\alpha}{}_\beta + T_{\mu\kappa}{}^\nu R^{\mu\kappa\alpha}{}_\beta \\ & + \frac{2\Lambda}{d(d-1)} \left[Q_\beta{}^{\nu\alpha} - \delta_\beta^\nu \bar{Q}^\alpha + 2T_\beta{}^{\alpha\nu} + U^{\mu\nu\alpha}{}_\beta \left(\frac{d}{4} Q_\mu + T_{\mu\kappa}{}^\kappa \right) \right] = 0, \end{aligned} \quad (59)$$

where

$$U_{\mu\nu\alpha\beta} = g_{\mu\beta} g_{\nu\alpha} - g_{\mu\alpha} g_{\nu\beta}, \quad (60)$$

again, Q_μ is the Weyl convector (30), which can be obtained from the non-metricity tensor $Q_{\mu\nu\alpha}$, defined as

$$Q_{\mu\alpha\beta} = \mathcal{D}_\mu g_{\alpha\beta} . \quad (61)$$

The quantity \bar{Q}_μ is also a vector constructed from the non-metricity,

$$\bar{Q}_\mu = Q_{\nu\mu}{}^\nu . \quad (62)$$

Finally, $T_{\mu\nu\alpha}$ is the torsion tensor (31).

The variation with respect to the metric tensor provides

$$\frac{1}{2}g^{\mu\nu} R^{\sigma\rho\alpha\beta} R_{\sigma\rho\alpha\beta} - R^\mu{}_{\alpha\beta\gamma} R^{\nu\alpha\beta\gamma} - R_{\alpha\beta\gamma}{}^\mu R^{\alpha\beta\gamma\nu} - \frac{8\Lambda}{d(d-1)} \left[\frac{1}{2}g^{\mu\nu} (R - \Lambda) - R^{\mu\nu} \right] = 0 . \quad (63)$$

Contraction of this equation, (63), with $g_{\mu\nu}$ results on the trace equation

$$\frac{(d-4)}{4} R^{\sigma\rho\alpha\beta} R_{\sigma\rho\alpha\beta} + \frac{2\Lambda}{d(d-1)} (2-d) R + \frac{2\Lambda^2}{(d-1)} = 0 . \quad (64)$$

Let us focus on the more interesting case $d = 4$. The field equations (59) and (63) now read

$$\begin{aligned} & -\mathcal{D}_\mu R^{\mu\nu\alpha}{}_\beta - 2 [2Q_\mu + T_{\mu\kappa}{}^\kappa] R^{\mu\nu\alpha}{}_\beta + T_{\mu\kappa}{}^\nu R^{\mu\kappa\alpha}{}_\beta \\ & + \frac{\Lambda}{6} \left[Q_\beta{}^{\nu\alpha} - \delta_\beta^\nu \bar{Q}^\alpha + 2T_\beta{}^\alpha{}_\nu + U^{\mu\nu\alpha}{}_\beta (Q_\mu + T_{\mu\kappa}{}^\kappa) \right] = 0 , \\ & \frac{1}{2}g^{\mu\nu} R^{\sigma\rho\alpha\beta} R_{\sigma\rho\alpha\beta} - R^\mu{}_{\alpha\beta\gamma} R^{\nu\alpha\beta\gamma} - R_{\alpha\beta\gamma}{}^\mu R^{\alpha\beta\gamma\nu} - \frac{2\Lambda}{3} \left[\frac{1}{2}g^{\mu\nu} (R - \Lambda) - R^{\mu\nu} \right] = 0 , \end{aligned} \quad (65)$$

while the trace equation (64) now provides

$$R = 2\Lambda . \quad (66)$$

In the usual gravitation theory with cosmological constant, *i.e.*, the action (58) with no quadratic terms in the curvature, the equation (66) is also obtained and the solution is the dS_4 space-time where the curvature is given by

$$R_{\mu\nu\alpha\beta} = \frac{\Lambda}{6} U_{\mu\nu\alpha\beta} . \quad (67)$$

Remarkably, this expression is also an exact solution of (65). To see this one has to use the fact that the geometry of the dS_4 space-time is Riemannian, $Q = T = 0$.

It is worth mention that one would generate a gravity theory with the AdS_4 as a solution if, in (52), a negative value for m_*^2 shows up. In that case $\Lambda < 0$ and we are in fact facing the AdS_4 instead of dS_4 .

5 Discussions and conclusions

In this work we have established a mapping from an Euclidean $SO(d)$ gauge theory to a dynamical space-time with linear connection and independent metric tensor. Moreover, in order

to relate the mapping with a model for gravity, we have developed a sketch for a color symmetry breaking mechanism which generates spin-2 excitations in Euclidean space-time. Thus, the mapping from the Euclidean space with a gauge field to a curved space with linear connection provides a modified gravity model with cosmological constant in the Palatini formalism.

We noticed that that is exactly the mappings (21) and (23) which leads action (7) into action (27). However, we strongly remark again that the presented color symmetry breaking mechanism is just a sketch. Realistic quantum computations are needed in order to give estimates for κ and m and, consequently, to G and Λ . By contrast, in principle, we have a vast freedom to find suitable solutions for κ and m within the renormalization group equations. This happens because, being renormalizable to all orders in perturbation theory, the theory is scale invariant as well as renormalization scheme invariant. Furthermore, the value the m is related to the value of Λ which receives several contributions from the non-perturbative vacuum energy of the rest of fundamental interactions [22]. For instance, at the electroweak sector there is a contribution coming from the Higgs vacuum while from the QCD sector there would be a contribution from the Gribov ambiguities and from condensates [20]. Obviously, even if consistent solutions might not show up analytically, numerical computations would be applied as well. Another point is that the gap equation (52) is an *ansatz* justified by the minimization principle of the vacuum energy. It might be possible to exist another way to compute m at the quantum level. We emphasize that this kind of computation (for κ and m within the renormalization group) is beyond the scope of this article and must be analyzed [13]. From now on, for the rest of the conclusion we will assume that the computation is possible and gives reasonable values for m and κ .

Concerning the space-time nature, we can conclude from the present proposal that, for very high energies, the fundamental space-time would be a continuous Euclidean space. Now, once we start to decrease the energy, the background starts to show its presence, breaking the color symmetry. Thus, spin-2 physical excitations show up and the theory can be visualized as a dynamical curved space-time. Under this scope, the fundamental space-time configurations would be the dS_4 space (Or AdS_4 if $\Lambda < 0$).

Another point to be explored in the future would be the inclusion of matter fields (non-gravitational fields) in the model. In principle, we can include the standard model in this formalism since we are dealing with an Euclidean space-time. However, it should result, at the gravitational low energy regime, in a consistent quantum standard model in a classical curved space.

An evident interpretation of the method is that there is no incompatibility between GR and QM. In fact, the quantum gravity theory is described in an Euclidean space-time, thus, there are no problems to be faced in the space and time definitions. At the quantum level, the space and time are the usual quantum mechanical parameters. At the classical level, the space and time are translated into the GR space-time. However, quantum effects are no longer associated with them. Thus, we can say that QG and GR would be different sides of the same coin.

Finally, we recall attention to the fact that the approach presented in this article might be generalized to other groups but the $SO(d)$, as well as for other classes of gauge theories. For instance, one may start with a $GL(d, \mathbb{R})$ gauge theory and see if it can be casted into a dynamical space-time. Obviously, in that particular case, the space-time representation exists as well.

Acknowledgments

The authors would like to acknowledge A. Accioly, H. J. M. Cuesta, S. A. Dias, E. L. Rodrigues and J. F. Villas da Rocha for fruitful and interesting discussions. Also J. A. Helayël-Neto is gratefully acknowledge for fruitful and clarifying discussions, suggestions and for reviewing the article. Still, we would like to kindly thank T. R. B. M. Rodrigues for suggestions in the correction of the text. Finally, the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq-Brazil) is acknowledged for financial support.

A Renormalizability of the model

To study renormalizability of the action (9) we will use the algebraic renormalization technique [14]. A basic ingredient of this method relies on the so called BRST symmetry. For the action (9) the BRST transformations are

$$\begin{aligned}
 sA_i &= -D_i c , \\
 sc &= \frac{1}{2}[c, c] , \\
 s\bar{c} &= b , \\
 sb &= 0 ,
 \end{aligned}
 \tag{68}$$

where s , the BRST operator, is nilpotent $s^2 = 0$.

In order to have a consistent quantum description of the model we introduce a set of external BRST invariant sources coupled to the nonlinear BRST variations,

$$\begin{aligned}
 S_{ext} &= s\text{Tr} \int d^d x (-\Omega_i A_i + Lc) \\
 &= \text{Tr} \int d^d x \left(-\Omega_i D_i c + \frac{1}{2}L[c, c] \right) ,
 \end{aligned}
 \tag{69}$$

where

$$s\Omega = sL = 0 . \tag{70}$$

Then, in this appendix we will work with the general BRST invariant action

$$\Sigma = \frac{1}{4}\text{Tr} \int d^d x F_{ij} F_{ij} + \text{Tr} \int d^d x (b \partial_i A_i + \bar{c} \partial_i D_i c) + \text{Tr} \int d^d x \left(-\Omega_i D_i c + \frac{1}{2}L[c, c] \right) . \tag{71}$$

Let us take a look at the power counting renormalizability of (9). For that, the UV dimensions of the fields and sources are displayed in table 1. Observing the dimension of the coupling constant we see that renormalizability would be a feature of the restricted cases where $2 \leq d \leq 4$.

A.1 Symmetries and Ward identities

The symmetries of the model are described by the following Ward identities:

fields	∂	s	A	c	\bar{c}	b	κ
UV dimension	1	0	1	0	$d-2$	$d-2$	$(4-d)/2$
Ghost number	0	1	0	1	-1	0	0

Table 1: Quantum numbers of the relevant quantities.

- The Slavnov-Taylor identity

$$\mathcal{S}(\Sigma) = \text{Tr} \int d^d x \left(\frac{\delta \Sigma}{\delta \Omega_i} \frac{\delta \Sigma}{\delta A_i} + \frac{\delta \Sigma}{\delta L} \frac{\delta \Sigma}{\delta c} + b \frac{\delta \Sigma}{\delta \bar{c}} \right) = 0. \quad (72)$$

- The ghost equation

$$\int d^d x \left(\frac{\delta \Sigma}{\delta c} + \left[\bar{c}, \frac{\delta \Sigma}{\delta b} \right] \right) = \int d^d x ([\Omega_i, A_i] + [L, c]). \quad (73)$$

- The gauge condition and the antighost equation

$$\begin{aligned} \frac{\delta \Sigma}{\delta b} &= \partial_i A_i, \\ \frac{\delta \Sigma}{\delta \bar{c}} + \partial_i \frac{\delta \Sigma}{\delta \Omega_i} &= 0. \end{aligned} \quad (74)$$

- The $SL(2, \mathbb{R})$ symmetry

$$\text{Tr} \int d^d x \left(c \frac{\delta \Sigma}{\delta \bar{c}} + \frac{\delta \Sigma}{\delta L} \frac{\delta \Sigma}{\delta b} \right) = 0. \quad (75)$$

- Global color symmetry

$$\int d^d x \left(\left[A_i, \frac{\delta \Sigma}{\delta A_i} \right] + \left[c, \frac{\delta \Sigma}{\delta c} \right] + \left[\bar{c}, \frac{\delta \Sigma}{\delta \bar{c}} \right] + \left[b, \frac{\delta \Sigma}{\delta b} \right] + \left[\Omega_i, \frac{\delta \Sigma}{\delta \Omega_i} \right] + \left[L, \frac{\delta \Sigma}{\delta L} \right] \right) = 0. \quad (76)$$

A.2 Characterization of the most general counterterm

Now, to show the renormalizability of the action (71) we will make use of the algebraic renormalization technique [14]. For that, one adds to the classical action a perturbation term,

$$\Sigma^{(1)} = \Sigma + \epsilon \Sigma^c, \quad (77)$$

where the counterterm Σ^c is an integrated polynomial on the fields and sources with UV dimension up-bounded by four and vanishing ghost number. The enforcement that the action (77) obeys the Ward identities (72-76) implies on the following constraints for the counterterm,

$$\begin{aligned} \mathcal{S}_\Sigma \Sigma^c &= 0, \\ \int d^d x \frac{\delta \Sigma^c}{\delta c} &= 0, \\ \frac{\delta \Sigma^c}{\delta b} &= 0, \\ \frac{\delta \Sigma^c}{\delta \bar{c}} + \partial_i \frac{\delta \Sigma^c}{\delta \Omega_i} &= 0, \\ \text{Tr} \int d^d x c \frac{\delta \Sigma^c}{\delta \bar{c}} &= 0, \\ \int d^d x \left(\left[A_i, \frac{\delta \Sigma^c}{\delta A_i} \right] + \left[c, \frac{\delta \Sigma^c}{\delta c} \right] + \left[\bar{c}, \frac{\delta \Sigma^c}{\delta \bar{c}} \right] + \left[\Omega_i, \frac{\delta \Sigma^c}{\delta \Omega_i} \right] + \left[L, \frac{\delta \Sigma^c}{\delta L} \right] \right) &= 0. \end{aligned} \quad (78)$$

where \mathcal{S}_Σ stands for the linearized Slavnov-Taylor operator,

$$\mathcal{S}_\Sigma = \text{Tr} \int d^d x \left(\frac{\delta \Sigma}{\delta \Omega_i} \frac{\delta}{\delta A_i} + \frac{\delta \Sigma}{\delta A_i} \frac{\delta}{\delta \Omega_i} + \frac{\delta \Sigma}{\delta L} \frac{\delta}{\delta c} + \frac{\delta \Sigma}{\delta c} \frac{\delta}{\delta L} + b \frac{\delta}{\delta \bar{c}} \right). \quad (79)$$

The first of (78) defines a cohomological problem whose general solution is given by

$$\Sigma^c = \Delta_0 + \mathcal{S}_\Sigma \Delta^{-1}, \quad (80)$$

where Δ_0 is the nontrivial part of the cohomology,

$$\Delta_0 \neq \mathcal{S}_\Sigma \cdot (\text{something}), \quad (81)$$

and $\mathcal{S}_\Sigma \Delta^{-1}$ is the trivial part. The nontrivial part is an integrated polynomial on the fields and sources with UV dimension up-bounded by four and vanishing ghost number while Δ^{-1} is an integrated polynomial with UV dimension up-bounded by four and ghost number given by -1 . In fact, it is straightforward to show that the most general counterterm (80) is determined by [14]

$$\Delta_0 = a_0 S_{\text{YM}}, \quad (82)$$

and

$$\Delta^{-1} = a_1 \text{Tr} \int d^d x (\Omega_i + \partial_i \bar{c}) A_i. \quad (83)$$

where a_0 and a_1 are independent renormalization parameters.

A.3 Quantum stability

The last step to prove renormalizability is to show that the counterterm (80) can be reabsorbed in the classical action (71) by means of multiplicative redefinition of the fields, sources and parameters according to

$$\Sigma(\phi_0, J_0, \kappa_0) = \Sigma(\phi, J, \kappa) + \epsilon \Sigma^c(\phi, J, \kappa), \quad (84)$$

where

$$\begin{aligned} \phi_0 &= Z_\phi^{1/2} \phi, \\ J_0 &= Z_J J, \\ \kappa_0 &= Z_\kappa \kappa, \end{aligned} \quad (85)$$

and

$$\begin{aligned} \phi &\in \{A, b, c, \bar{c}\}, \\ J &\in \{\Omega, L\}. \end{aligned} \quad (86)$$

In fact it is not difficult to show that the action (71) is multiplicatively renormalizable, where the independent renormalization factors are given by

$$\begin{aligned} Z_\kappa &= 1 - \epsilon \frac{a_0}{2}, \\ Z_A^{1/2} &= 1 + \epsilon \left(\frac{a_0}{2} + a_1 \right), \end{aligned} \quad (87)$$

and the nonindependent renormalization factors read

$$\begin{aligned} Z_c^{1/2} = Z_{\bar{c}}^{1/2} = Z_\Omega^{-1/2} &= Z_\kappa^{-1/2} Z_A^{-1/4}, \\ Z_b^{1/2} = Z_L^{-2} &= Z_A^{-1}. \end{aligned} \tag{88}$$

This ends the proof of the renormalizability of pure $SO(d)$ gauge theory in a $2 \leq d \leq 4$ -dimensional Euclidean space-time.

B Renormalization of m_*^2

In this appendix we provide the program for derivation of the renormalization rule of the background parameter m_*^2 . We omit the details since they are somewhat simple to be performed. For that we consider as a starting point the fact that the background is a particular configuration of the gauge field A . Thus

$$\Upsilon_0 = Z_A^{1/2} \Upsilon. \tag{89}$$

Now, perturbing the equation (84) according to (47) we will find an expression where the mass parameter m_*^2 is present. Thus, performing the usual multiplicative renormalization (85-86) together with

$$m_{*0}^2 = Z_{m^2} m_*^2, \tag{90}$$

we find expression (35).

References

- [1] G. 't Hooft and M. J. G. Veltman, *Annales Poincare Phys. Theor. A* **20**, 69 (1974).
- [2] S. Deser and P. van Nieuwenhuizen, *Phys. Rev. D* **10**, 401 (1974).
- [3] S. Deser and P. van Nieuwenhuizen, *Phys. Rev. D* **10**, 411 (1974).
- [4] K. S. Stelle, *Phys. Rev. D* **16**, 953 (1977).
- [5] C. Rovelli, “*Quantum Gravity*,” Cambridge, UK: Univ. Pr. (2004) 455 p.
- [6] L. Smolin, arXiv:hep-th/0408048.
- [7] R. Utiyama, *Phys. Rev.* **101**, 1597 (1956).
- [8] T. W. B. Kibble, *J. Math. Phys.* **2**, 212 (1961).
- [9] S. Ferrara, *Phys. Scripta* **T15**, 132 (1987).
- [10] S. Deser, *Phys. Scripta* **T15**, 138 (1987).
- [11] D. M. Gitman and I. V. Tyutin, “*Quantization of fields with constraints*,” Springer (1990) 291 p. (Springer series in nuclear and particle physics).

- [12] A. O. Bolivar, “*Quantum-Classical correspondence*,” Springer (2004) 186 p. (The Frontiers Collection).
- [13] *Work in progress*.
- [14] O. Piguet and S. P. Sorella, “*Algebraic renormalization: Perturbative renormalization, symmetries and anomalies*,” Lect. Notes Phys. **M28**, 1 (1995).
- [15] N. C. Ribeiro, R. F. Sobreiro and S. P. Sorella, Phys. Lett. A **317**, 6 (2003) [arXiv:hep-th/0302202].
- [16] G. 't Hooft, In **Karpacz 1975, Proceedings, Acta Universitatis Wratislaviensis No.368, Vol.1**, Wroclaw 1976, 345-369.
- [17] S. P. Sorella, Annals Phys. **321**, 1747 (2006).
- [18] V. N. Gribov, Nucl. Phys. B **139**, 1 (1978).
- [19] D. Zwanziger, Nucl. Phys. B **399**, 477 (1993).
- [20] D. Dudal, R. F. Sobreiro, S. P. Sorella and H. Verschelde, Phys. Rev. D **72**, 014016 (2005) [arXiv:hep-th/0502183].
- [21] C. Itzykson and J. B. Zuber, “*Quantum Field Theory*,” New York, USA: McGraw-Hill (1980) 705 P.(International Series In Pure and Applied Physics).
- [22] I. L. Shapiro and J. Sola, J. Phys. A **40**, 6583 (2007) [arXiv:gr-qc/0611055].