Boundary State for Dissipative Quantum Mechanics and Thirring Model

Taejin Lee∗

Department of Physics, Kangwon National University, Chuncheon 200-701, Korea and

Pacific Institute for Theoretical Physics Department of Physics and Astronomy,

University of British Columbia, 6224 Agricultural Road, Vancouver, British Columbia V6T 1Z1, Canada

(Dated: October 30, 2018)

The dissipative quantum system is studied using the Thirring model with a boundary mass. At the critical point where the Thirring coupling vanishes, the theory reduces to a free fermion theory with a boundary mass. We construct boundary states for the dissipative quantum systems in one dimension, which describes the system off the critical points as well as at the critical points.

PACS numbers: 03.65.-w, 05.40.Jc, 11.10.-z, 11.25.-w Keywords:

I. INTRODUCTION

The two space-time dimensional quantum field theory of massless bosons with a periodic boundary potential has recently come into the spotlight again. This boundary conformal field theory has received constant attention along the years, since it describes the various condensed physics systems such as the dissipative quantum mechanics of a particle in a one-dimensional periodic potential [1–4], Josephson junction arrays [5–7] and the dissipative Hofstadter problem [8, 9]. The application of the theory also includes the Kondo problem [10, 11], the one-dimensional conductors [12], tunnelling between Hall edge states [13], and junctions of quantum wires [14]. The recent revival of interest in the theory is mainly due to the string theory application. The string with its ends on a unstable D-brane develops a marginal periodic boundary interaction and the tachyon field of the open string condensates [15–20]. This process called "rolling tachyon" [21, 22] is believed to be responsible for the decay of the unstable D-brane.

Recently the theory with a periodic boundary potential is discussed in detail by applying the fermionization technique [23, 24]. The advantage of the fermion formulation is that one can explicitly construct the boundary state for the rolling tachyon, since the periodic boundary potential becomes a boundary fermion mass term, which is quadratic in fermion field. The fermion formulation of the theory, however, has been performed only for the theory at the critical point. In this paper we shall generalize the fermion formulation to the theory off the critical point and develop a perturbative theory around the critical point by using the Thirring model.

The massless Thirring model [25], which is the first example of an exactly solvable relativistic interacting field theory, has served as an excellent laboratory for the study of various aspects of the quantum field theory in two dimensions. Since the seminal work of Thirring, extensive studies of the model have been carried out by numerous authors [26–29]; notably the complete solution of the theory was obtained by Klaiber [29]. Along this line Schwinger [30] obtained an exact solution of quantum electrodynamics in 1+1 dimensions and Coleman [31] proved the equivalence between the sine-Gordon model and the massive Thirring model.

In this Letter we will widen the range of applications of the Thirring model by describing the dissipative quantum system in terms of the Thirring model. We find that the Thirring model with a boundary mass is the most suitable framework to discuss the dissipative quantum system: At the critical point the Thirring coupling, which is directly related to the friction constant of the system, vanishes and the theory reduces to a free fermion theory with a boundary mass. The boundary state takes a simple form at the critical point when it is expressed in fermion variables as discussed in refs. [23, 24]. Thus, the Thirring model provides a perturbative theory expanded in the Thirring coupling near the critical point. The previous analysis of the dissipative quantum system is based on this Coulomb gas expansion; the duality symmetries and the phase diagram of the system have been studied in this framework. However, the perturbative theory based on the Coulomb gas expansion has some limitation to study the system near or off the critical point, being one dimensional theory with a non-local interaction. In the followings we formulate the dissipative system of a particle moving in one dimension subject to a periodic potential, known as the Schmid model, near the critical point in terms of the Thirring model and the boundary state formulation.

[∗]Electronic address: taejin@kangwon.ac.kr

II. THE SCHMID MODEL AND THE THIRRING MODEL

Caldeira and Leggett [33, 34] discussed first the quantum mechanical description of dissipation by coupling a bath or environment, which consists of an infinite number of harmonic oscillators, to the system. In the quantum theory the interaction with the bath produces a non-local effective interaction. Subsequenlty Schmid [1] studied the dissipative system in the presence of a periodic potential. The one dimensional dissipative model with a periodic potential, called Schmid model, is described by the following action

$$
S_{SM} = \frac{\eta}{4\pi\hbar} \int_{-T/2}^{T/2} dt dt' \frac{(X(t) - X(t'))^2}{(t - t')^2} - \frac{V_0}{\hbar} \int_{-T/2}^{T/2} dt \cos\frac{2\pi X}{a}.
$$
 (1)

The first non-local term is responsible for the dissipation, and the second term denotes the periodic potential respectively. An interesting feature of the model is that it exhibits a phase transition, unlike one dimensional quantum mechanical systems with local interactions only. Depending on the value of the friction constant η , the system has two phases; the localized phase and the delocalized one.

We can map the Schmid model to the string theory on a disk by identifying the time as the boundary parameter σ in string theory and scaling the field variable X:

$$
t = \frac{T}{2\pi}\tau, \quad X \to \frac{a}{2\pi}X. \tag{2}
$$

Then, the action for the Schmid model reads as

$$
S_{SM} = \frac{\eta}{4\pi\hbar} \left(\frac{a}{2\pi}\right)^2 \int d\tau d\tau' \frac{\left(X(\tau) - X(\tau')\right)^2}{(\tau - \tau')^2} - \frac{V_0}{\hbar} \frac{T}{2\pi} \int d\tau \frac{1}{2} \left(e^{iX} + e^{-iX}\right). \tag{3}
$$

This action can be interpreted as the boundary effective action for the open bosonic string subject to a boundary periodic potential on a disk with a boundary condition; on the boundary ∂M , $X(\sigma, \tau) = X(\tau)$,

$$
e^{-iS_{SM}} = \int D[X] \exp\bigg[-i\left(\frac{1}{4\pi\alpha'}\int_M d\tau d\sigma \partial_\alpha X \partial^\alpha X - \frac{m}{2}\int_{\partial M} d\tau \left(e^{iX} + e^{-iX}\right)\right)\bigg].\tag{4}
$$

Here, we identify the physical parameters of the two theories as

$$
\frac{\eta}{4\pi\hbar} \left(\frac{a}{2\pi}\right)^2 = \frac{1}{8\pi^2\alpha'}, \quad -\frac{V_0}{\hbar} \frac{T}{2\pi} = m.
$$
\n⁽⁵⁾

In string theory the periodic potential describes the interaction between the open string the unstable D-brane.

The open string dynamics is often described more efficiently in its equivalent closed string picture by the boundary state formulation. The corresponding closed string action is obtained from its open string action by simply taking $\sigma \to \tau, \, \tau \to \sigma,$

$$
S = \frac{1}{4\pi\alpha'} \int d\tau d\sigma \, \partial_{\alpha} X \partial^{\alpha} X - \frac{m}{2} \int d\sigma \left(e^{iX} + e^{-iX} \right) \tag{6}
$$

The open string dynamics can be encoded completely by the boundary state.

When $\alpha = 1/\alpha' = 1$, the system becomes critical. This can be easily understood if we introduce an auxiliary boson field Y and fermionize the system [23, 24, 35]. Introducing an auxiliary boson field Y which satisfies the Dirichlet condition $Y|_{\partial M} = 0$ at the boundary, and defining the boson fields, $\phi_1 = \frac{X+Y}{\sqrt{2}}$ $\frac{Y}{2}$, $\phi_2 = \frac{X-Y}{\sqrt{2}}$, we may rewrite the action as

$$
S = \frac{\alpha}{4\pi} \int_M d\tau d\sigma \sum_i^2 \partial \phi_i \partial \phi_i - \frac{m}{4} \int_{\partial M} d\tau \sum_i^2 \left(e^{i\sqrt{2}\phi_i} + e^{-i\sqrt{2}\phi_i} \right). \tag{7}
$$

Since Y is a free boson field and it vanishes at the boundary, Y is completely decoupled from the physical degrees of freedom. We see that if $\alpha = 1$, $e^{\pm i\sqrt{2}\phi_i}$ are marginal boundary operator with the scaling dimension 1. An explicit calculation of the current correlation function or the mobility shows that

$$
\langle 0|\partial_{\sigma}X(\sigma)\partial_{\sigma}X(\sigma')|B\rangle = -\frac{1}{2}(1-\pi^2m^2)\sin^{-2}\frac{(\sigma-\sigma')}{2}
$$
\n(8)

the theory becomes indeed critical where $\alpha = 1$.

An explicit evaluation of the boundary state at the critical point is possible if we fermionize the model; the boson fields are mapped to the fermion fields as

$$
\psi_{1L}(z) = \zeta_{1L} : e^{-\sqrt{2}i\phi_{1L}(z)} : , \quad \psi_{2L}(z) = \zeta_{2L} : e^{\sqrt{2}i\phi_{2L}(z)} : \tag{9a}
$$

$$
\psi_{1R}(\bar{z}) = \zeta_{1R} : e^{\sqrt{2}i\phi_{1R}(\bar{z})} : , \quad \psi_{2R}(\bar{z}) = \zeta_{2R} : e^{-\sqrt{2}i\phi_{2R}(\bar{z})} : \tag{9b}
$$

where $\zeta_{iL/R}$ are co-cycles, ensuring the anti-commutation relations between the fermion operators. Since the boundary interaction term can be written as a boundary fermion mass term, which is only quadratic in fermion field, the model is exactly solvable

$$
S = \int \frac{d\tau d\sigma}{2\pi} \left(\bar{\psi}_1 \gamma^\mu \partial_\mu \psi_1 + \bar{\psi}_2 \gamma^\mu \partial_\mu \psi_2 \right) + m \int \frac{d\sigma}{2\pi} \left(\bar{\psi}_1 \psi_1 + \bar{\psi}_2 \psi_2 \right) \tag{10}
$$

where $\psi_i = (\psi_{iL}, \psi_{iR})^t$, and

$$
\gamma^0 = \sigma_1, \quad \gamma^1 = \sigma_2, \quad \gamma^5 = \sigma_3 = -i\gamma^0\gamma^1. \tag{11}
$$

The boundary state is given formally as

$$
|B\rangle =: \exp\left[m \int \frac{d\sigma}{2\pi} \left(\bar{\psi}_1 \psi_1 + \bar{\psi}_2 \psi_2\right)\right] : |N, D\rangle \tag{12}
$$

where $|N, D\rangle$ is a simple boundary state satisfying

$$
\left(\psi_R(0,\sigma) + i\sigma^2\psi_L(0,\sigma)\right)|N,D\rangle = 0, \quad \left(\psi_R^{\dagger}(0,\sigma) + i\psi_L^{\dagger}(0,\sigma)\sigma^2\right)|N,D\rangle = 0.
$$
\n(13)

We refer the reader to ref. [24] for the explicit expression of the boundary state.

Now let us discuss the dissipative system off the critical points. When $\alpha \neq 1$, we may write the action as

$$
S = \frac{1}{4\pi} \int_{M} d\tau d\sigma \sum_{i}^{2} \partial \phi_{i} \partial \phi_{i} + \frac{1}{4\pi} (\alpha - 1) \int_{M} d\tau d\sigma \sum_{i}^{2} \partial \phi_{i} \partial \phi_{i}
$$

$$
- \frac{m}{4} \int_{\partial M} d\tau \sum_{i}^{2} \left(e^{i\sqrt{2}\phi_{i}} + e^{-i\sqrt{2}\phi_{i}} \right). \tag{14}
$$

and treat the second term as an interaction. In terms of the fermion fields the second term can be written as the Thirring interaction term. Hence, the fermionized action is given by

$$
S = \frac{1}{2\pi} \int_M d\tau d\sigma \sum_i^2 \left(\bar{\psi}_i \gamma^\mu \partial_\mu \psi_i + \frac{g}{4\pi} j_i^\mu j_{i\mu} \right) + \frac{m}{2} \int_{\partial M} d\sigma \sum_i^2 \bar{\psi}_i \psi_i \tag{15}
$$

where $g = \pi(\alpha - 1)$. This is the Thirring model with a boundary mass. At the critical point where $g = 0$ ($\alpha = 1$), the action reduces to the free fermion theory with a boundary mass. Near the critical point, we can use this Thirring action to develop perturbative theory for the dissipative quantum system.

III. BOUNDARY STATE NEAR THE CRITICAL POINT

In order to apply the boundary state formulation to the dissipative system near the critical point, the bulk action should be free. We may transmute the bulk Thirring interaction into a boundary one by introducing Abelian gauge fields

$$
S = \frac{1}{2\pi} \int_M d\tau d\sigma \sum_i^2 \left[\bar{\psi}_i \gamma^\mu \left(\partial_\mu + i A_{i\mu} \right) \psi_i + \frac{\pi}{g} A_{i\mu} A_i{}^\mu \right] + \frac{m}{2} \int_{\partial M} d\sigma \sum_i^2 \bar{\psi}_i \psi_i. \tag{16}
$$

Since in general the Abelian gauge vector fields in $1 + 1$ dimensions may be decomposed as

$$
A_i^{\mu} = \epsilon^{\mu\nu} \partial_{\nu} \theta_i + \partial^{\mu} \chi_i, \quad i = 1, 2,
$$
\n(17)

the interaction between the gauge fields and the fermion fields may be removed by a gauge transformation

$$
\psi_i = e^{-\gamma_5 \theta_i - i\chi_i} \psi_i_0, \quad \bar{\psi}_i = \bar{\psi}_i_0 e^{-\gamma_5 \theta_i + i\chi_i}.
$$
\n
$$
(18)
$$

Then the bulk action becomes a free field one

$$
S_{bulk} = \frac{1}{2\pi} \int_M d\tau d\sigma \sum_{i=1}^2 \left[\bar{\psi}_i{}_0 \gamma^\mu \partial_\mu \psi_i{}_0 + \left(\frac{\pi}{g} + 1 \right) (\partial \theta_i)^2 + \frac{\pi}{g} (\partial \chi_i)^2 \right]. \tag{19}
$$

The additional kinetic action for θ_i is a manifestation of the $U(1)$ chiral anomaly:

$$
D[\psi]D[\bar{\psi}] = D[\psi_0]D[\bar{\psi}_0] \exp\left[\frac{1}{4\pi} \int d\tau d\sigma \sum_i (\partial \theta_i)^2\right].
$$
\n(20)

Since the boundary mass term is not invariant under the $U(1)$ chiral gauge transformation, it transforms as

$$
\sum_{i} \bar{\psi}_i \psi_i = \sum_{i} \bar{\psi}_{i0} e^{-2\gamma_5 \theta_i} \psi_{i0}.
$$
\n(21)

Note that scalar fields χ_i are free in the bulk and do not appear in the boundary action. Since the physical operators, being $U(1)_V$ gauge invariant, do not depend on χ_i , we may drop them. For the sake of convenience, we scale the scalar fields

$$
\theta_i \to \kappa \theta_i, \quad \kappa = \sqrt{\frac{g}{2(\pi + g)}}.
$$
\n(22)

It brings us to

$$
S = \frac{1}{2\pi} \int_M d\tau d\sigma \sum_{i=1}^2 \left[\bar{\psi}_{i0} \gamma^\mu \partial_\mu \psi_{i0} + \frac{1}{2} (\partial \theta_i)^2 \right] + m \int_{\partial M} d\sigma \sum_i \bar{\psi}_{i0} e^{-2\gamma_5 \kappa \theta_i} \psi_{i0}.
$$
 (23)

It is clear that in the limit of the critical point, $\kappa \to 0$, the interaction between the scalar fields θ_i and ψ_0 vanishes. Thus, θ_i , becoming free fields, can be dropped and the action reduces to one at the critical point.

The next step to construct the perturbative boundary state formulation is to find appropriate boundary conditions for ψ_0^i and θ_i . We note that the fermion fields ψ_0^i should be understood as those at the critical point. So the boundary conditions for them are the same as those in Eq.(13) The boundary conditions for the fermion fields would remain intact, since the boundary conditions for the boson fields ϕ_i would not change if the Thirring interaction term is turned on. The boundary conditions for the fermion fields can be formally kept unchanged, if we require the boson fields satisfy the following conditions

$$
\theta_1|B_0\rangle = -\theta_2|B_0\rangle, \quad \chi_1|B_0\rangle = \chi_2|B_0\rangle. \tag{24}
$$

As we already noticed that χ_i are completely decoupled from the physical degrees of freedom, we are not concerned with the boundary conditions for χ_i . Eq.(24) only fixes the boundary condition for $\frac{1}{\sqrt{2}}$ $\frac{1}{2}(\theta_1 + \theta_2)$. For the sake of completness, we may choose a Neumann conditions for $\frac{1}{\sqrt{2}}$ $\frac{1}{2}(\theta_1 - \theta_2)$. Once, the simple boundary state $|B_0\rangle$ is constructed, the boundary state for the dissipated system $|B(m, \kappa)\rangle$ may be given as

$$
|B(m,\kappa)\rangle = \exp\left[m \int_{\partial M} d\sigma \sum_{i} \bar{\psi}_{i} e^{-2\gamma_{5}\kappa \theta_{i}} \psi_{i}\right] |B_{0}\rangle
$$
\n(25)

where we drop the subscript " $_0$ " of the fermion fields for notational convenience.

It is interesting to see that the scalar fields θ_i appear only through the boundary interaction and the physical operators such as currents do not depend upon them. Near the critical point where $|\kappa| < 1$, we may expand the boundary state $|B(m, \kappa)\rangle$ in κ as follows

$$
|B(m,\kappa)\rangle = \exp\left[m\int d\sigma \sum_{i} \bar{\psi}_{i}\psi_{i} - 2m\kappa \int d\sigma \sum_{i} \bar{\psi}_{i}\gamma^{5}\psi_{i}\theta_{i}\right]|B(0,0)\rangle
$$

$$
= \sum_{n} \frac{(-2m\kappa)^{n}}{n!} \left[\int d\sigma \sum_{i} \bar{\psi}_{i}\gamma^{5}\psi_{i}\theta_{i}\right]^{n} |B(m,0)\rangle
$$
(26)

where $|B(m,0)\rangle$ corresponds to the boundary state at the critical point, of which explicit expression can be found in [24].

As we expand $|B(m,\kappa)\rangle$ in κ , we encounter a divergent term, proportional to the boundary mass term with a coefficient

$$
2m^{2}\kappa^{2}\int d\sigma_{1}\int d\sigma_{2}\sum_{i}\langle\theta_{i}(\sigma_{1})\theta_{i}(\sigma_{2})\rangle \Big(\langle\bar{\psi}\gamma^{5}\psi(\sigma_{1})\bar{\psi}\gamma^{5}\psi(\sigma_{2})\rangle
$$

$$
=2m^{2}\kappa^{2}\int d\sigma_{1}\int d\sigma_{2}\ln\left|1-\frac{z_{1}}{z_{2}}\right|\left(\frac{1}{z_{1}-z_{2}}\frac{1}{\bar{z}_{1}-\bar{z}_{2}}\right)
$$
(27)

where $z_i = e^{i\sigma_i}$ and $\bar{z}_i = e^{-i\sigma_i}$. In order to regularize it we may introduce an infinitesimal parameter $|\epsilon| \ll 1$ as follows

$$
z_1 = e^{i\sigma_1}, \quad \bar{z}_1 = \frac{1}{z_1}, \quad z_2 = e^{-\epsilon}e^{i\sigma_2}, \quad \bar{z}_2 = \frac{e^{-2\epsilon}}{z_2}.
$$
 (28)

Then we find,

$$
\int d\sigma_1 \int d\sigma_2 \ln \left| 1 - \frac{z_1}{z_2} \right| \left(\frac{1}{z_1 - z_2} \frac{1}{\bar{z}_1 - \bar{z}_2} \right) = \frac{\pi^2}{2\epsilon}.
$$
\n(29)

This divergence can be taken care of by renormalization of the boundary mass. It yields

$$
m = m_0 \left[1 + \frac{\alpha - 1}{2\alpha} \ln \frac{\Lambda^2}{\mu^2} \right] = m_0 \left(\frac{\Lambda^2}{\mu^2} \right)^{\frac{(\alpha - 1)}{2\alpha}}
$$
\n(30)

where $\ln \frac{\Lambda^2}{\mu^2} = \frac{m_0^2 \pi^2}{\epsilon}$ $\frac{p^{\pi}}{\epsilon}$. The result is in complete agreement with the previous work ref.[3]: If $\alpha > 1$, m tends to grow and if α < 1, it scales to zero. The corection to m vanishes when $\alpha = 1$. In fact, an explicit construction of the boundary state showes that perturbative corrections to m vanish at all orders if $\alpha = 1$.

A few remarks are in order to conclude this Letter. We discuss the dissipative quantum system in one dimension, employing the boundary state formulation of string theory and the Thirring model in two dimensions. The framework presented here has some advantages over the previous one based on the Coulomb gas expansion which deals with the non-local interaction: We need to deal with a local boundary interaction only and can take an advantage of the string theory techniques to explore various aspects of the system. Since the system is described in terms of the (bulk) massless Thirring model, all the physical quantities are exactly calculable at any given order.

Acknowledgement: The author thanks Gordon Semenoff, Philip Stamp for informative discussions.

- [1] A. Schmid, Phys. Rev. Lett. **51**, 1506 (1983).
- [2] F. Guinea, V. Hakim and A. Muramatsu, Phys. Rev. Lett. 54, 263 (1985).
- [3] M.P.A. Fisher and W. Zwerger, Phys. Rev. B32, 6190 (1985).
- [4] C. G. Callan and L. Thorlacius, Nucl. Phys. B **329**, 117 (1990).
- [5] L.I. Glazman and A.I. Larkin, Phys.Rev.Lett. 79, 3736 (1997).
- [6] R. Fazio and H. van der Zant, Phys. Rep. 355, 235 (2001)
- [7] D. Giuliano and P. Sodano, Nucl. Phys. B711, 480-504, (2005).
- [8] C. G. Callan and D. Freed, Nucl. Phys. B 374, 543 (1992).
- [9] C. G. Callan, A. G. Felce and D. E. Freed, Nucl. Phys. B 392, 551 (1993).
- [10] I. Affleck and A. W. W. Ludwig, Nucl. Phys. B 360, 641 (1991).
- [11] I. Affleck and A. W. W. Ludwig, Nucl. Phys. B 352, 849 (1991).
- [12] C. L. Kane and M. P. A. Fisher, Phys. Rev. **B46**, 15233 (1992).
- [13] C.L. Kane and M.P.A. Fisher, [arXiv:cond-mat/9506116].
- [14] C. Chamon, M. Oshikawa and I. Affleck, Phys. Rev. Lett. 91, 206403 (2003).
- [15] E. Witten, Phys. Rev. D **46**, 5467 (1992).
- [16] S. L. Shatashvili, Phys. Lett. B 311, 83 (1993).
- [17] S. L. Shatashvili, Alg. Anal. 6, 215 (1994).
- [18] T. Lee, Phys. Rev. D 64, 106004 (2001).
- [19] T. Lee, Phys. Lett. B **520**, 385 (2001).
- [20] T. Lee, J. Korean Phys. Soc. 42, 34 (2003).
- [21] A. Sen, JHEP 0204, 048 (2002).
- [22] T. Lee, JHEP 0611, 056 (2006).
- [23] T. Lee and G. W. Semenoff, JHEP 0505, 072 (2005).
- [24] M. Hasselfield, T. Lee, G. W. Semenoff and P. C. .E. Stamp, Ann. Phys. 321, 2849 (2006).
- [25] W. Thirring, Ann. Phys. 3, 91 (1958).
- [26] V. Glaser, Nuovo Cim. 9, 990 (1958).
- [27] K. Johnson, Nuovo Cim. 20, 773 (1961).
- [28] C. R. Hagen, Nuovo Cim. 51B, 169 (1967).
- [29] B. Klaiber, in "Lectures in Theoretical Physics," edited by A. Barut and W. Brittin, Gordon and Breach, New York, 1968, p. 141-176.
- [30] J. S. Schwinger, Phys. Rev. **125**, 397 (1962).
- [31] S. Coleman, Phys. Rev. D 11, 2088 (1975).
- [32] S. Ji, J.-Y. Koo and T. Lee, to be published in J. Korean Phys. Soc. (2007).
- [33] A. O. Caldeira and A. J. Leggett, Ann. Phys. **149**, 374 (1983).
- [34] A. O. Caldeira and A. J. Leggett, Physica 121A, 587 (1983).
- [35] J. Polchinski and L. Thorlacius, Phys. Rev. D **50**, R622 (1994).