

# Bispinor Formulation of Spin 3/2 Field Theory

Haryanto M. Siahaan\*

## Abstract

In this paper, we investigate an alternative formulation for spin 3/2 field equation. First we will review equation of motion of Dirac and Maxwell, and then construct the equation for spin 3/2 in the similar fashion. Our method actually a generalization of relativistic equation of motion based on spin operator and Hamiltonian similar to Dirac and Maxwell equation in ref. [1] pp. 50. Furthermore, we could not bring the equation to the Klein Gordon limit for every wave function component in spin 3/2 formalism.

## 1 Introduction

The famous Dirac equation can be read as

$$(i\gamma^\mu\partial_\mu - \beta m) = 0 \quad (1)$$

which also can be written as

$$i\frac{\partial}{\partial t}\psi = \left(\frac{1}{i}\vec{\alpha} \cdot \nabla - \beta m\right)\psi \quad (2)$$

Then  $\left(\frac{1}{i}\vec{\alpha} \cdot \nabla - \beta m\right)$  is recognized as Dirac Hamiltonian. Each matrix in equation (2) is

$$\vec{\alpha} = \begin{pmatrix} 0 & \vec{\sigma} \\ \vec{\sigma} & 0 \end{pmatrix}; \beta = \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \quad (3)$$

where  $\vec{\sigma}$  are Pauli matrices. In bispinor context, equation (2) can be written as a coupled equation which relate the particle wave function and it's antiparticle.

$$i\frac{\partial}{\partial t}\varphi = m\varphi + \frac{2}{i}\vec{S} \cdot \nabla\chi \quad (4)$$

---

\*E-mail : hm\_siahaan@gmail.com

$$i\frac{\partial}{\partial t}\chi = -m\chi + \frac{2}{i}\vec{S} \cdot \nabla\varphi \quad (5)$$

where  $\vec{S} = \frac{\vec{\sigma}}{2}$  as our spin operator for every two component spinor in Dirac equation. This spin operator satisfy Lie algebra for  $SU(2)$  group,  $[S^i, S^j] = i\epsilon^{ijk}S^k$ . The square of this operator also satisfy  $\vec{S} \cdot \vec{S} = s(s+1) = \frac{3}{4}$  for  $s = \frac{1}{2}$ .

To get the plane wave with  $E^2 = \vec{p}^2 + m^2$  as a solution, equation (2) should satisfy the Klein Gordon equation. This can be done by square each operator at left hand side and right hand side. The properties of  $\{\vec{\alpha}, \beta\} = 0$  and  $\{\alpha_i, \alpha_j\} = 0$  which guarantee that we can get the Klein Gordon expression for each  $\psi$ .

A similar method can be applied for Maxwell equation. In the case of no source, Maxwell equation can be read as

$$\nabla \times \vec{E} + \frac{\partial \vec{B}}{\partial t} = 0, \quad (6)$$

$$\nabla \times \vec{B} - \frac{\partial \vec{E}}{\partial t} = 0. \quad (7)$$

Now, by introducing such a spinor for electromagnetic fields that is  $\psi_{EM} = (\vec{E}, i\vec{B})$ , so equation (6) and (7) can be rewritten as

$$i\frac{\partial}{\partial t}\psi_{EM} = \frac{1}{i}\vec{\alpha}_{EM} \cdot \nabla\psi_{EM} \quad (8)$$

where

$$\vec{\alpha}_{EM} = \begin{pmatrix} 0 & \vec{S}_{EM} \\ \vec{S}_{EM} & 0 \end{pmatrix} \quad (9)$$

and  $\vec{S}_{EM}$  is generator of rotation in 3 dimensional Euclidean space. In this way, we can say that  $(\frac{1}{i}\vec{\alpha}_{EM} \cdot \nabla)$  is the "Maxwell Hamiltonian". The  $\vec{S}_{EM}$  can be recognized as for spin operator for  $s = 1$ , because it satisfy  $[S^i, S^j] = i\epsilon^{ijk}S^k$  and  $\vec{S} \cdot \vec{S} = s(s+1) = 2$ .

$$S_{EM}^1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, S_{EM}^2 = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix}, S_{EM}^3 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & 0 & 0 \end{pmatrix} \quad (10)$$

Furthermore, we can check that equation (8) does not satisfy the Klein Gordon equation. In other words, each component of  $\psi_{EM}$  does not have

plane wave solution with  $E^2 = \vec{p}^2 + m^2$  as the eigenvalue of energy. It is clear because at elementary level we know that the dynamical variable is  $A^\mu(x) = (A^0(x), \vec{A}(x))$  which has relation with  $\vec{E}(x)$  and  $\vec{B}(x)$  by  $\vec{E}(x) = -\left(\frac{\partial \vec{A}(x)}{\partial t} - \nabla A^0(x)\right)$  and  $\vec{B}(x) = \nabla \times \vec{A}(x)$ . By imposing this relation and the Lorentz gauge constraint<sup>1</sup>, we can get the massless Klein Gordon expression for every  $A^\mu$ 's component. At this point, we can learn that the Klein Gordon equation, or in other words relativistic mass-energy-momentum relation, could not be directly satisfied when we construct a higher spin formulation in "bispinor" fashion.

## 2 Construction of Spin 3/2 Field Equation

Learning from the pattern that we get in Dirac and Maxwell formalism, we can make a generalization for arbitrary spin by denoting that Hamiltonian operator could have the form  $(\frac{a}{i}\vec{\alpha}_s \cdot \nabla - bm)$  with undetermined complex matrices a and b.  $\vec{\alpha}_s$  is a left diagonal matrix which contains spin operator and s denote the value of spin. So in general we could have an equation which can be read as

$$i\frac{\partial}{\partial t}\Psi_s(x) = \left(\frac{a}{i}\vec{\alpha}_s \cdot \nabla + bm\right)\Psi_s(x) \quad (11)$$

where  $\Psi_s(x)$  is the bispinor for arbitrary spin field and has  $2s(s+1)$  components.

So in spin 3/2 formalism, we would get an equation of motion that written as

$$i\frac{\partial}{\partial t}\Psi_{3/2}(x) = \left(\frac{a}{i}\vec{\alpha}_{3/2} \cdot \nabla + bm\right)\Psi_{3/2}(x) \quad (12)$$

where

$$\vec{\alpha}_{3/2} = \begin{pmatrix} 0 & \vec{S}_{3/2} \\ \vec{S}_{3/2} & 0 \end{pmatrix} \quad (13)$$

and  $\vec{S}_{3/2} = (S_{3/2}^1, S_{3/2}^2, S_{3/2}^3)$ .

$$S_{3/2}^1 = \frac{1}{2} \begin{pmatrix} 0 & \sqrt{3} & 0 & 0 \\ \sqrt{3} & 0 & 2 & 0 \\ 0 & 2 & 0 & \sqrt{3} \\ 0 & 0 & \sqrt{3} & 0 \end{pmatrix}, \quad (14)$$

---

<sup>1</sup>This constraint must be included

$$S_{3/2}^2 = \frac{i}{2} \begin{pmatrix} 0 & -\sqrt{3} & 0 & 0 \\ \sqrt{3} & 0 & -2 & 0 \\ 0 & 2 & 0 & -\sqrt{3} \\ 0 & 0 & \sqrt{3} & 0 \end{pmatrix}, \quad (15)$$

$$S_{3/2}^3 = \begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -3 \end{pmatrix}. \quad (16)$$

Matrices of spin operator are produced by usual angular momentum relations,

$$S_{\pm} = S^1 \pm iS^2, \quad (17)$$

$$S^3 |s, m\rangle = m |s, m\rangle, \quad (18)$$

$$S_{\pm} |s, m\rangle = \sqrt{(s \mp m)(s \pm m + 1)} |s, m \pm 1\rangle. \quad (19)$$

For simplification, we try to evaluate equation (12) in massless case. In an explicit form, we can write our equation as

$$i \frac{\partial}{\partial t} \begin{pmatrix} \Phi(x) \\ \Omega(x) \end{pmatrix} = \frac{a}{i} \begin{pmatrix} 0 & \vec{S}_{3/2} \cdot \nabla \\ \vec{S}_{3/2} \cdot \nabla & 0 \end{pmatrix} \begin{pmatrix} \Phi(x) \\ \Omega(x) \end{pmatrix}. \quad (20)$$

Before further investigation, we should recognize that our spin operator, or more precisely our  $\vec{\alpha}_{3/2}$  do not anticommute for each different components<sup>2</sup> and  $\alpha_{3/2}^i \alpha_{3/2}^i \neq \frac{1}{a} I_{4 \times 4}$  which means equation (20) cannot simply have Klein Gordon expression. Our formalism cannot directly fulfill this requirement to be such a relativistic equation. Probably this problem is similar to Maxwell formalism when we use the "spinor"  $\psi_{EM} = (\vec{E}, i\vec{B})$  as the dynamical variable. Perhaps we need another constraints to satisfy this relativistic wave equation requirement or finding another "dynamical variable" that can be related to our bispinor for describing spin 3/2 fields just like in Maxwell formalism case before.

### 3 Conclusion

Trying to get such a consistent theory of higher spin fields (higher than spin 1) is a complicated manner. Beside, there still no single particle with higher

---

<sup>2</sup>Compared to Dirac's  $\vec{\alpha}$

spin that needs some explanation ever exist in laboratory. But effort to get a good equation is an interesting thing. Furthermore, finding the expression (11) for standard Rarita Schwinger formulation can be proved imposible.

\*\*\*

I am grateful to Prof. Triyanta for interesting discussion and his encouragement.

## 4 References

1. C. Itzykson, and J. Zuber, "Quantum Field Theory," McGraw-Hill (1980). pp. 48-50.
2. W.Rarita, and J. Schwinger, Phys. Rev. 60 (1941) 61.