SYMMETRIES OF SPATIAL GRAPHS AND SIMON INVARIANTS

RYO NIKKUNI AND KOUKI TANIYAMA

ABSTRACT. An ordered and oriented 2-component link L in the 3-sphere is said to be achiral if it is ambient isotopic to its mirror image ignoring the orientation and ordering of the components. Kirk-Livingston showed that if Lis achiral then the linking number of L is not congruent to 2 modulo 4. In this paper we study orientation-preserving or reversing symmetries of 2-component links, spatial complete graphs on 5 vertices and spatial complete bipartite graphs on 3 + 3 vertices in detail, and determine the necessary conditions on linking numbers and Simon invariants for such links and spatial graphs to be symmetric.

1. INTRODUCTION

Throughout this paper we work in the piecewise linear category. Let $L = J_1 \cup J_2$ be an ordered and oriented 2-component link in the unit 3-sphere \mathbb{S}^3 . Unless otherwise stated, the links in this paper will be ordered and oriented. A link L is said to be component preserving achiral (CPA) if there exists an orientation-reversing self-homeomorphism φ of \mathbb{S}^3 such that $\varphi(J_1) = J_1$ and $\varphi(J_2) = J_2$, and component switching achiral (CSA) if there exists an orientation-reversing self-homeomorphism φ of \mathbb{S}^3 such that $\varphi(J_2) = J_1$ [3]. If L is either CPA or CSA, then L is said to be achiral. Note that L may be both CPA and CSA (a trivial link, for example). The following was shown by Kirk-Livingston.

Theorem 1.1. ([4, 6.1 COROLLARY]) If L is achiral then lk(L) is not congruent to 2 modulo 4, where lk denotes the linking number.

See also [7, Theorem 5.1] for an elementary proof of Theorem 1.1. Note that for any odd integer n there exists a 2-component link of linking number n which is both CPA and CSA; see Fig. 1.1 (cf. [7, §5]). In [7], Livingston gave an example of a CSA link with linking number 4 (a cabling of the Hopf link) and stated open problems: to find an achiral link of linking number 4m for any integer m, and to find a CPA link of linking number 4m for any integer m". For the latter, Kidwell showed the following.

Theorem 1.2. ([3, Theorem 4]) A 2-component link of nonzero even linking number cannot be CPA.

²⁰⁰⁰ Mathematics Subject Classification. Primary 57M15; Secondary 57M25.

Key words and phrases. Symmetric spatial graph, achiral link, Simon invariant, linking number.

The first author was partially supported by Grant-in-Aid for Young Scientists (B) (No. 18740030), Japan Society for the Promotion of Science.

The second author was partially supported by Grant-in-Aid for Scientific Research (C) (No. 18540101), Japan Society for the Promotion of Science.





Kidwell also gave an example of a CSA link of linking number 4m for any odd integer m [3, §3]. But as far as the authors know, a CSA link of linking number 4m for any nonzero even integer m has not been exhibited yet.

To give a complete answer to Livingston's problem, we will present a new family of achiral links. For an integer m and $\varepsilon_1, \varepsilon_2 \in \{-1, 1\}$, let $L(m, \varepsilon_1, \varepsilon_2)$ be a 2component link of linking number $4m + (\varepsilon_1 + \varepsilon_2)/2$ as illustrated in Fig. 1.2. Note that $L(0, \varepsilon, -\varepsilon)$ is trivial for $\varepsilon = \pm 1$. Therefore $L(0, \varepsilon, -\varepsilon)$ is both a CPA and CSA link of linking number 0. Moreover we have the following.



+1 = -1 = -1

FIGURE 1.2.

Theorem 1.3. (1) For any integer m and $\varepsilon_1, \varepsilon_2 \in \{-1, 1\}$, $L(m, \varepsilon_1, \varepsilon_2)$ is CSA. (2) For any integer m and $\varepsilon = \pm 1$, $L(m, \varepsilon, \varepsilon)$ is CPA.

Proof. In Fig. 1.2, we may suppose that a regular diagram of $L(m, \varepsilon_1, \varepsilon_2)$ on the standard 2-sphere \mathbb{S}^2 in \mathbb{S}^3 is such that the crossing of sign ε_1 is on the north pole and that of sign ε_2 is on the south pole.

(1) A $\pi/2$ rotation of \mathbb{S}^2 around the earth's axis maps $L(m, \varepsilon_1, \varepsilon_2)$ onto its mirror image. Thus we have the assertion.

(2) The π rotation of \mathbb{S}^2 around an axis through the equator interchanges the components of $L(m, \varepsilon, \varepsilon)$. Then by composing it with a $\pi/2$ rotation of \mathbb{S}^2 around the earth's axis, we see that $L(m, \varepsilon, \varepsilon)$ is mapped onto its mirror image preserving the components. Thus we have the result.

The following corollary shows that Theorems 1.1 and 1.2 give the best possible necessary conditions on the linking number for a 2-component link to be CPA or to be CSA.

Corollary 1.4. (1) For any integer m and $\varepsilon = \pm 1$, $L(m, \varepsilon, \varepsilon)$ is both a CPA and CSA link of linking number $4m + \varepsilon$.

(2) For any integer m and $\varepsilon = \pm 1$, $L(m, \varepsilon, -\varepsilon)$ is a CSA link of linking number 4m.

Remark 1.5. A further step in the investigation of the achirality of 2-component links is to take the invertibility of each component into account, as follows. For a 2-component link $L = J_1 \cup J_2$, assume that there exists a self-homeomorphism φ of \mathbb{S}^3 such that $\varphi(L) = L$. Then the candidates of $(\varphi(J_1), \varphi(J_2))$ are (1) (J_1, J_2) , (2) (J_2, J_1) , (3) $(-J_1, -J_2)$, (4) $(-J_2, -J_1)$, (5) $(-J_1, J_2)$, (6) $(J_1, -J_2)$, (7) $(-J_2, J_1)$, (8) $(J_2, -J_1)$. When $lk(L) \neq 0$, by observing the change of the sign of lk(L) it can be seen that the first four cases may occur only if φ is orientation-preserving, and the last four only if φ is orientation-reversing, where L is CPA in case of (5) or (6), and CSA in case of (7) or (8). The first four cases can be realized by a (2, 2n)-torus link simultaneously for any integer n, so these symmetries do not depend on the linking number. On the other hand, since $L(m, \varepsilon_1, \varepsilon_2)$ is invertible, only Theorems 1.1 and 1.2 are restrictions of the linking number for the latter four cases.

Next, let us consider finite graphs which are embedded in \mathbb{S}^3 . An embedding f of a finite graph G into \mathbb{S}^3 is called a *spatial embedding of* G or simply a *spatial graph*. Two spatial embeddings f and g of G are said to be *ambient isotopic* if there exists an orientation-preserving self-homeomorphism φ of \mathbb{S}^3 such that $\varphi \circ f = g$. In [21], the second author introduced the notion of (*spatial graph-)homology* as a fundamental equivalence relation on spatial graphs which is weaker than ambient isotopy. It is known that a homology is generated by *Delta moves* and ambient isotopies [9, Theorem 1.3], where a Delta move is a local deformation of a spatial graph as illustrated in Fig. 1.3. Linking numbers of constituent 2-component links are typical homological invariants of spatial graphs. In particular, two k-component links are homologous if and only if they have the same pairwise linking numbers [10, Theorem 1.1].

On the other hand, let K_5 and $K_{3,3}$ be a *complete graph* on five vertices and a *complete bipartite graph* on 3 + 3 vertices respectively, known as obstructions in Kuratowski's graph planarity criterion [6]. For a spatial embedding f of K_5 or $K_{3,3}$, the *Simon invariant* $\mathcal{L}(f)$ is defined [21, §4], which is an odd integer



FIGURE 1.3.

valued homological invariant calculated from the regular diagram of f, like the linking number. We give a precise definition of $\mathcal{L}(f)$ in the next section. It is known that two spatial embeddings of a graph are homologous if and only if they have the same *Wu invariant* [22], or equivalently, their corresponding constituent 2-component links have the same linking number and their corresponding spatial subgraphs which are homeomorphic to K_5 or $K_{3,3}$ have the same Simon invariant [17]. Thus, linking numbers and Simon invariants play a primary role in classifying spatial graphs up to homology. We remark here that both the linking number and the Simon invariant come from the Wu invariant as special cases, see [22, §2] for details.

Our main purpose in this paper is to reveal the relationship between orientationpreserving or reversing symmetries of spatial embeddings of K_5 and $K_{3,3}$ and their Simon invariants in addition to the case of 2-component links and their linking numbers. A study of symmetries of spatial graphs is not only a fundamental problem in low dimensional topology as a generalization of achirality of knots and links but also an important research theme from a standpoint of application to macromolecular chemistry, which is called molecular topology [23], [16], [1], [2]. Actually, both K_5 and $K_{3,3}$ have been realized chemically as underlying molecule graphs for some chemical compounds [14], [12], [24], [5]. We refer the reader to [15] for a pioneer work on symmetries of spatial embeddings of K_5 and $K_{3,3}$.

From now on we assume that G is K_5 or $K_{3,3}$. Let $\operatorname{Aut}(G)$ be the *automorphism* group of G. We regard each automorphism of G as a self-homeomorphism of G. For a spatial embedding f of G and an automorphism σ on G, we say that f is σ symmetric (resp. rigidly σ -symmetric) if there exists a self-homeomorphism (resp. periodic self-homeomorphism) φ of \mathbb{S}^3 such that $f \circ \sigma = \varphi \circ f$. In particular, if φ is orientation-reversing then we say that f is σ -achiral (resp. rigidly σ -achiral). In the first author's preliminary report [11], it was shown that for any odd integer nthere exist an automorphism σ on G and a σ -achiral spatial embedding f of G such that $\mathcal{L}(f) = n$. Then it is natural to ask the following question.

Question 1.6. For an automorphism σ on G and an odd integer n, does there exist a σ -symmetric spatial embedding f of G such that $\mathcal{L}(f) = n$?

For an automorphism σ of G and an odd integer n, we say that the pair (σ, n) is *realizable* if there exists a σ -symmetric spatial embedding f of G such that $\mathcal{L}(f) = n$. Then the following proposition holds.

Proposition 1.7. Let σ and τ be two automorphisms of G which are conjugate in Aut(G), and n an odd integer. Then (σ, n) is realizable if and only if (τ, n) is realizable.

Therefore it is sufficient to consider only conjugacy classes in $\operatorname{Aut}(G)$. Note that we may identify $\operatorname{Aut}(G)$ with a subgroup of the symmetric group \mathfrak{S}_m of degree m, where m is the number of vertices of G. If G is $K_{3,3}$, we assume that the vertices corresponding to 1, 2 and 3 are adjacent to the vertices corresponding to 4, 5 and 6. Then we have the following.

Proposition 1.8. (1) Representatives for all conjugacy classes in $Aut(K_5)$ are

id, (1 2), (1 2 3), (1 2 3 4), (1 2 3 4 5), (1 2)(3 4), (1 2)(3 4 5).

(2) Representatives for all conjugacy classes in $Aut(K_{3,3})$ are

id, $(1 \ 2)$, $(1 \ 2 \ 3)$, $(1 \ 4 \ 2 \ 5 \ 3 \ 6)$, $(1 \ 2)(4 \ 5)$, $(1 \ 2)(4 \ 5 \ 6)$, $(1 \ 2 \ 3)(4 \ 5 \ 6)$, $(1 \ 4 \ 2 \ 5)(3 \ 6)$, $(1 \ 4)(2 \ 5)(3 \ 6)$.

Now we state our main theorems.

Theorem 1.9. Let G be K_5 and n an odd integer. Then we have the following. (1) ((1 2 3), n) is realizable if and only if n is congruent to ± 1 modulo 6.

- (2) $((1 \ 2 \ 3 \ 4 \ 5), n)$ is realizable.
- $(3) ((1 \ 2)(3 \ 4), n)$ is realizable.
- $(4) ((1 \ 2 \ 3 \ 4), n)$ is realizable.
- (5) ((1 2), n) is realizable if and only if $n = \pm 1$.
- (6) $((1\ 2)(3\ 4\ 5), n)$ is realizable if and only if $n = \pm 1$.

Theorem 1.10. Let G be $K_{3,3}$ and n an odd integer. Then we have the following. (1) ((1 2 3), n) is realizable if and only if n is congruent to ± 1 modulo 6.

- (2) $((1 \ 2)(4 \ 5), n)$ is realizable.
- (3) $((1 \ 2 \ 3)(4 \ 5 \ 6), n)$ is realizable.
- (4) $((1 \ 4)(2 \ 5)(3 \ 6), n)$ is realizable.
- (5) $((1 \ 4 \ 2 \ 5 \ 3 \ 6), n)$ is realizable.
- (6) $((1 \ 4 \ 2 \ 5)(3 \ 6), n)$ is realizable.
- (7) ((1 2), n) is realizable if and only if $n = \pm 1$.
- (8) ((1 2)(4 5 6), n) is realizable if and only if $n = \pm 1$.

Furthermore, each realizable pair (σ, n) may be realized by a rigidly σ -symmetric spatial embedding of G. By combining Theorems 1.9 and 1.10 with Propositions 1.7 and 1.8, we can give a complete answer to Question 1.6. We can also determine when (σ, n) is realized by a σ -achiral spatial embedding of G as follows.

Theorem 1.11. (1) Let σ be an automorphism of K_5 and n an odd integer. Suppose that (σ, n) is realizable. Then (σ, n) is realized by a σ -achiral spatial embedding of K_5 if and only if σ is conjugate to $(1 \ 2 \ 3 \ 4)$, $(1 \ 2)$ or $(1 \ 2)(3 \ 4 \ 5)$.

(2) Let σ be an automorphism of $K_{3,3}$ and n an odd integer. Suppose that (σ, n) is realizable. Then (σ, n) is realized by a σ -achiral spatial embedding of $K_{3,3}$ if and only if σ is conjugate to $(1 \ 4 \ 2 \ 5)(3 \ 6)$, $(1 \ 2)$ or $(1 \ 2)(4 \ 5 \ 6)$.

Actually we will show that a realizable pair (σ, n) in Theorem 1.9 (1), (2), (3) and Theorem 1.10 (1), (2), (3), (4), (5) can only be realized by an orientation-preserving self-homeomorphism of \mathbb{S}^3 , and a realizable pair (σ, n) in Theorem 1.9 (4), (5), (6) and Theorem 1.10 (6), (7), (8) can only be realized by an orientation-reversing self-homeomorphism of \mathbb{S}^3 .

In the next section we give a precise definition of the Simon invariant. In section 3, we prove some propositions including Propositions 1.7 and 1.8 that are needed later. The proofs of Theorems 1.9, 1.10 and 1.11 are given in section 4.

2. Simon invariant

Let G be K_5 or $K_{3,3}$ where the vertices have a fixed numbering. In this section we review the Simon invariant of spatial embeddings of G. We give an orientation to each edge of G as illustrated in Fig. 2.1, according to the numbering of the vertices. For an unordered pair x, y of disjoint edges K_5 , we define the sign $\varepsilon(x, y)$ by $\varepsilon(e_i, e_j) = 1$, $\varepsilon(d_k, d_l) = -1$ and $\varepsilon(e_i, d_k) = -1$. For an unordered pair x, y of disjoint edges of $K_{3,3}$, we also define the sign $\varepsilon(x, y)$ by $\varepsilon(c_i, c_j) = 1$, $\varepsilon(b_k, b_l) = 1$ and $\varepsilon(c_i, b_k) = 1$ if c_i and b_k are parallel in Fig. 2.1, and -1 if they are anti-parallel. For a spatial embedding f of G, we fix a regular diagram of f and denote the sum of the signs of the crossing points between f(x) and f(y) by l(f(x), f(y)), where x, y is an unordered pair of disjoint edges of G. Now we define an integer $\mathcal{L}(f)$ by

$$\mathcal{L}(f) = \sum_{(x,y)} \varepsilon(x,y) l(f(x), f(y)),$$

where the summation is taken over all unordered pairs of disjoint edges of G. This integer $\mathcal{L}(f)$ is called the *Simon invariant* of f. Actually, this is an odd integer valued ambient isotopy invariant [21, THEOREM 4.1] up to the numbering of the vertices. In paricular, for a different numbering of the vertices, the value of the Simon invariant may be different. The Simon invariant can also be described from a cohomological viewpoint as follows. See [22, Example 2,4, Example 2.5] for details. Let $C_2(X)$ be the *configuration space* of ordered two points of a topological space X, namely

$$C_2(X) = \{(x, y) \in X \times X \mid x \neq y\}.$$

Let ι be an involution on $C_2(X)$ defined by $\iota(x, y) = (y, x)$. Then we call the integral cohomology group of Ker $(1 + \iota_{\sharp})$ the *skew-symmetric integral cohomology group* of the pair $(C_2(X), \iota)$ and denote it by $H^*(C_2(X), \iota)$. It is known that $H^2(C_2(\mathbb{R}^3), \iota) \cong$ \mathbb{Z} [25] and $H^2(C_2(G), \iota) \cong \mathbb{Z}$. We denote a generator of $H^2(C_2(\mathbb{R}^3), \iota)$ by Σ . Let $f : G \to \mathbb{S}^3 \setminus \{(0, 0, 0, 1)\}$ be a spatial embedding of G, regarded as an embedding into \mathbb{R}^3 . This embedding f naturally induces an equivariant embedding $f \times f : C_2(G) \to C_2(\mathbb{R}^3)$ with respect to the action ι and therefore induces a homomorphism

$$(f \times f)^* : H^2(C_2(\mathbb{R}^3), \iota) \longrightarrow H^2(C_2(G), \iota).$$

Then the Simon invariant $\mathcal{L}(f)$ coincides with $(f \times f)^*(\Sigma)$ up to sign. Thus the absolute value $|\mathcal{L}(f)|$ is an ambient isotopy invariant independent of the numbering of the vertices.

The Simon invariant $\mathcal{L}(f)$ is also closely related to the constituent knots of f. Let $\Gamma(G)$ be the set of all cycles of G, where a *cycle* is a subgraph of G which is homeomorphic to a circle. We say that a cycle is a *k*-cycle if it contains exactly kedges. Let $\omega : \Gamma(G) \to \mathbb{Z}$ be the map defined by

$$\omega(\gamma) = \begin{cases} 1 & \text{(if } \gamma \text{ is a 5-cycle)} \\ -1 & \text{(if } \gamma \text{ is a 4-cycle)} \\ 0 & \text{(if } \gamma \text{ is a 3-cycle)} \end{cases}$$

for $G = K_5$, and

$$\omega(\gamma) = \begin{cases} 1 & \text{(if } \gamma \text{ is a 6-cycle)} \\ -1 & \text{(if } \gamma \text{ is a 4-cycle)} \end{cases}$$



FIGURE 2.1.

for $G = K_{3,3}$. For a spatial embedding f of G, we define an integer $\alpha_{\omega}(f)$ by

$$\alpha_{\omega}(f) = \sum_{\gamma \in \Gamma(G)} \omega(\gamma) a_2(f(\gamma)),$$

where $a_2(J)$ is the second coefficient of the Conway polynomial of a knot J. This integer $\alpha_{\omega}(f)$ is called the α -invariant of f [20]. In [9], Motohashi and the second author showed that if $\mathcal{L}(f) = 2j - 1$ then

$$\alpha_{\omega}(f) = \frac{j(j-1)}{2}.$$

This implies that

(2.1)
$$\alpha_{\omega}(f) = \frac{\mathcal{L}(f)^2 - 1}{8}$$

for a spatial embedding f of G. Then we have the following.

Lemma 2.1. Let $G = K_5$ or $K_{3,3}$, and let f be a spatial embedding of G. Then $\alpha_{\omega}(f)$ is a multiple of 3 if and only if $\mathcal{L}(f) \equiv \pm 1 \pmod{6}$.

Proof. It is easy to check that $m \equiv \pm 1 \pmod{6}$ if and only if $m^2 \equiv \pm 1 \pmod{6}$. Therefore $\mathcal{L}(f) \equiv \pm 1 \pmod{6}$ if and only if $\mathcal{L}(f)^2 - 1$ is a multiple of 6. By (2.1) we have $\mathcal{L}(f)^2 - 1 = 8\alpha_{\omega}(f)$. Since $8\alpha_{\omega}(f)$ is a multiple of 6 if and only if $\alpha_{\omega}(f)$ is a multiple of 3 we have the desired conclusion.

3. Conjugacy classes in Aut(G) and Simon invariants

In this section we discuss the relationship between conjugacy classes in Aut(G) and Simon invariants.

Lemma 3.1. Let $G = K_5$ or $K_{3,3}$. For an automorphism σ of G and an odd integer n, if (σ, n) is realizable then $(\sigma, -n)$ is realizable.

Proof. Let ρ be an orientation-reversing self-homeomorphism of \mathbb{S}^3 defined by $\rho(x_1, x_2, x_3, x_4) = (x_1, x_2, x_3, -x_4)$, and f a spatial embedding of G. We call $\rho \circ f$ the mirror image embedding of f and denote it by f!. If (σ, n) is realized by f, there exists a self-homeomorphism φ of \mathbb{S}^3 such that $f \circ \sigma = \varphi \circ f$. Then the following diagram is

commutative.



Thus, f! is σ -symmetric. By the definition of the Simon invariant, we can see that $\mathcal{L}(f!) = -\mathcal{L}(f) = -n$. Therefore $(\sigma, -n)$ is realizable.

Lemma 3.2. Let $G = K_5$ or $K_{3,3}$, and f a spatial embedding of G. Then $\mathcal{L}(f \circ \xi) = \pm \mathcal{L}(f)$ for any automorphism ξ of G.

Proof. The automorphism ξ naturally induces an equivariant homeomorphism $\xi \times \xi : C_2(G) \to C_2(G)$ with respect to the action ι and therefore induces an isomorphism

$$(\xi \times \xi)^* : H^2(C_2(G), \iota) \xrightarrow{\cong} H^2(C_2(G), \iota).$$

Then we have

$$\mathcal{L}(f \circ \xi) = ((f \circ \xi) \times (f \circ \xi))^*(\Sigma) = (\xi \times \xi)^*((f \times f)^*(\Sigma)) = (\xi \times \xi)^*(\mathcal{L}(f))$$

where Σ is a suitable generator of $H^2(C_2(\mathbb{R}^3), \iota)$; see the following commutative diagram:

Since $H^2(C_2(G), \iota) \cong \mathbb{Z}$, we have the desired conclusion.

Proof of Proposition 1.7. Assume that there exist a spatial embedding f of G, a self-homeomorphism φ of \mathbb{S}^3 and an automorphism ξ of G such that $f \circ \sigma = \varphi \circ f$ and $\tau = \xi^{-1}\sigma\xi$. Then the following diagram is commutative:

$$\begin{array}{ccc} G & \stackrel{\xi}{\longrightarrow} G & \stackrel{f}{\longrightarrow} \mathbb{S}^{3} \\ \tau & & \sigma & & \downarrow \varphi \\ G & \stackrel{\xi}{\longrightarrow} G & \stackrel{f}{\longrightarrow} \mathbb{S}^{3} \end{array}$$

Hence, $f \circ \xi$ is τ -symmetric. Thus, by Lemma 3.2, $\mathcal{L}(f \circ \xi) = \pm \mathcal{L}(f) = \pm n$. If $\mathcal{L}(f \circ \xi) = n$, then (τ, n) is realizable. If $\mathcal{L}(f \circ \xi) = -n$, then (τ, n) is also realizable by Lemma 3.1.

Proof of Proposition 1.8. (1) We may identify $\operatorname{Aut}(K_5)$ with the symmetric group of degree 5. It is not hard to see that all conjugacy classes in $\operatorname{Aut}(K_5)$ are classified as follows:

(i) id,

- (ii) $(i \ j)$ for $\{i, j\} \subset \{1, 2, 3, 4, 5\}$,
- (iii) $(i \ j \ k)$ for $\{i, j, k\} \subset \{1, 2, 3, 4, 5\}$,
- (iv) $(i \ j \ k \ l)$ for $\{i, j, k, l\} \subset \{1, 2, 3, 4, 5\}$,
- (v) (i j k l m) for $\{i, j, k, l, m\} = \{1, 2, 3, 4, 5\},\$
- (vi) $(i \ j)(k \ l)$ for $\{i, j\} \subset \{1, 2, 3, 4, 5\}$ and $\{k, l\} \subset \{1, 2, 3, 4, 5\} \setminus \{i, j\}$,

8

(vii) $(i \ j)(k \ l \ m)$ for $\{i, j\} \subset \{1, 2, 3, 4, 5\}$ and $\{k, l, m\} = \{1, 2, 3, 4, 5\} \setminus \{i, j\}$. Hence we have the desired representatives. We omit the details. (2) In a similar way, we may identify $\operatorname{Aut}(K_{3,3})$ with a subgroup of \mathfrak{S}_6 of order 10 (Note that $\operatorname{Aut}(K_{3,3})$ has a structure of the *wreath product* $\mathfrak{S}_2[\mathfrak{S}_3]$). It is not hard to see that all conjugacy classes in $\operatorname{Aut}(K_{3,3})$ are classified as follows:

(i) id.

(ii) $(i \ j)$ for $\{i, j\} \subset \{1, 2, 3\}$ or $\{i, j\} \subset \{4, 5, 6\}$,

(iii) $(i \ j \ k)$ for $\{i, j, k\} = \{1, 2, 3\}$ or $\{4, 5, 6\}$,

(iv) $(i \ l \ j \ m \ k \ n)$ for $\{i, j, k\} = \{1, 2, 3\}$ and $\{l, m, n\} = \{4, 5, 6\}$,

(v) $(i \ j)(l \ m)$ for $\{i, j\} \subset \{1, 2, 3\}$ and $\{l, m\} \subset \{4, 5, 6\}$,

(vi) $(i \ j)(l \ m \ n)$ for $\{i, j\} \subset \{1, 2, 3\}$ and $\{l, m, n\} = \{4, 5, 6\}$, or $\{i, j\} \subset \{4, 5, 6\}$ and $\{l, m, n\} = \{1, 2, 3\}$,

(vii) $(i \ j \ k)(l \ m \ n)$ for $\{i, j, k\} = \{1, 2, 3\}$ and $\{l, m, n\} = \{4, 5, 6\}$,

(viii) $(i \ l \ j \ m)(k \ n)$ for $\{i, j\} \subset \{1, 2, 3\}, \{l, m\} \subset \{4, 5, 6\}$ and $k \in \{1, 2, 3\} \setminus \{i, j\}, n \in \{4, 5, 6\} \setminus \{l, m\},$

(ix) $(i \ l)(j \ m)(k \ n)$ for $\{i, j, k\} = \{1, 2, 3\}$ and $\{l, m, n\} = \{4, 5, 6\}$.

Hence we have the desired representatives. We also omit the details.

Remark 3.3. We can decide the sign of $\mathcal{L}(f)$ in Lemma 3.2 as follows. Let $D_2(G)$ be the union of $s \times t$ where (s,t) varies over all pairs of disjoint edges of G. It is known that $D_2(G)$ is homotopy equivalent to $C_2(G)$, equivariantly with respect to the action ι [22, Proposition 1.4]. Moreover, $D_2(K_5)$ and $D_2(K_{3,3})$ are homeomorphic to the closed connected orientable surfaces of genus 6 and 4, respectively [13]. Then, for an automorphism σ of G, $\mathcal{L}(f \circ \sigma) = \mathcal{L}(f)$ (resp. $\mathcal{L}(f \circ \sigma) = -\mathcal{L}(f)$ if and only if $(\sigma \times \sigma)|_{D_2(G)}$ is orientation-preserving (resp. orientation-reversing). Note that if two automorphisms σ and τ are conjugate and $(\sigma \times \sigma)|_{D_2(G)}$ is orientation-preserving (resp. orientation-reversing), then $(\tau \times$ τ)|_{D₂(G)} is also orientation-preserving (resp. orientation-reversing). As a consequence, $(\sigma \times \sigma)|_{D_2(K_5)}$ is orientation-reversing if and only if σ is conjugate to $(1 \ 2 \ 3 \ 4), (1 \ 2)$ or $(1 \ 2)(3 \ 4 \ 5), and (\sigma \times \sigma)|_{D_2(K_{3,3})}$ is orientation-reversing if and only if σ is conjugate to $(1 \ 4 \ 2 \ 5)(3 \ 6)$, $(1 \ 2)$ or $(1 \ 2)(4 \ 5 \ 6)$ by a direct observation how the oriented 2-cells of $D_2(G)$ are mapped (or we may check that by a direct calculation of the Simon invariants under two different numberings of the vertices that differ by an automorphism of G). In only these cases, it may happen that $\mathcal{L}(f \circ \sigma) = -\mathcal{L}(f)$.

4. Proofs of main theorems

For the proofs of Theorems 1.9 and 1.10, we need two results which have been proved as consequences of 3-manifold topology. The following is a direct corollary of [19, Theorem 2].

Theorem 4.1. Let $G = K_5$ or $K_{3,3}$, σ an automorphism of G, and f a σ -symmetric spatial embedding of G. Then there exist a spatial embedding g of G which is homologous to f and a periodic self-homeomorphism φ of \mathbb{S}^3 such that $g \circ \sigma = \varphi \circ g$.

The following is well known as Smith's theorem.

Theorem 4.2. ([18]) Let φ be an orientation-reversing periodic self-homeomorphism of \mathbb{S}^3 . Then the fixed point set $\operatorname{Fix}(\varphi)$ is homeomorphic to \mathbb{S}^0 or \mathbb{S}^2 .

In the following proofs we denote the image of the vertex with label i under the spatial embedding also by i, so long as no confusion arises.

Proof of Theorem 1.9. (3) and (4) Let $f_{m,\pm}$ be the spatial embeddings of K_5 as in Fig. 4.1. By a calculation we have $\mathcal{L}(f_{m,\pm}) = 4m \pm 1$, which may be any odd number for a suitable choice of m and ± 1 . A $\pi/2$ rotation around the axis through the vertex 5, the middle point of the edge $\overline{13}$ and the middle point of the edge $\overline{24}$ maps $f_{m,\pm}(K_5)$ onto its mirror image. Therefore $f_{m,\pm}$ is (1 2 3 4)-symmetric. Thus ((1 2 3 4), n) is realizable for any odd integer n. A π rotation maps $f_{m,\pm}(K_5)$ onto itself and hence $f_{m,\pm}$ is (1 3)(2 4)-symmetric. Therefore ((1 3)(2 4), n) is realizable for any odd integer n. Since (1 2)(3 4) is conjugate to (1 3)(2 4), Proposition 1.7 implies ((1 2)(3 4), n) is realizable for any odd integer n.



FIGURE 4.1.

(2) Let f_m be the spatial embedding of K_5 as in Fig. 4.2. Note that $\mathcal{L}(f_m) = 2m + 1$, which may be any odd number. Observe that $f_m(K_5)$ is contained in an unknotted Möbius strip in \mathbb{S}^3 containing m full twists. Then by considering a suitable ambient isotopy of \mathbb{S}^3 that preserves the Möbius strip setwise we conclude that f_m is $(1 \ 2 \ 3 \ 4 \ 5)$ -symmetric.



FIGURE 4.2.

(1) ("if" part) Let $f_{m,\pm}$ be the spatial embedding of K_5 as in Fig. 4.3. Note that $\mathcal{L}(f_{m,\pm}) = 6m \pm 1$. A $2\pi/3$ rotation around the axis through the vertices 4 and 5 maps $f_{m,\pm}(K_5)$ onto itself and so $f_{m,\pm}$ is (1 2 3)-symmetric.



FIGURE 4.3.

(5) ("if" part) and (6) ("if" part) Let f_{\pm} be the spatial embedding of K_5 as in Fig. 4.4. Note that $\mathcal{L}(f_{\pm}) = \pm 1$. By considering the reflection of \mathbb{S}^3 with respect to the 2-sphere containing the cycle [3 4 5], we see that f_{\pm} is (1 2)-symmetric. Then by composing this reflection and a $2\pi/3$ rotation around the axis through the vertices 1 and 2, we find that f_{\pm} is (1 2)(3 4 5)-symmetric.



FIGURE 4.4.

(1) ("only if" part) Suppose that f is a (1 2 3)-symmetric spatial embedding of K_5 . We will show that $\alpha_{\omega}(f)$ is a multiple of 3. Then by Lemma 2.1 we have the result. There exist twelve 5-cycles and fifteen 4-cycles in K_5 . By the permutation

 $(1\ 2\ 3)$ they are divided into the following nine orbits:

Note that each orbit contains exactly three cycles and these cycles are mapped onto the same knot under f. Therefore they have the same a_2 and we have the desired conclusion.

(5) ("only if" part) Suppose that f is a (1 2)-symmetric spatial embedding of K_5 . Then by Theorem 4.1 there exist a (1 2)-symmetric spatial embedding g of K_5 which is homologous to f and a periodic self-homeomorphism φ of \mathbb{S}^3 such that $g \circ (1 2) = \varphi \circ g$. As pointed out in Remark 3.3, it follows that $\mathcal{L}(g \circ (1 2)) = -\mathcal{L}(g)$. Thus $\mathcal{L}(\varphi \circ g) = \mathcal{L}(g \circ (1 2)) = -\mathcal{L}(f)$, so φ is orientation-reversing. Then by Theorem 4.2, Fix(φ) is homeomorphic to \mathbb{S}^0 or \mathbb{S}^2 . Since Fix(φ) contains at least three points g(3), g(4) and g(5), it is homeomorphic to \mathbb{S}^2 . Then Fix(φ) $\cap g(K_5)$ is the union of the 3-cycle g([3 4 5]) and the middle point of the edge $g(\overline{1 2})$. By considering the regular projection which is almost perpendicular to Fix(φ) and observing that the signs of crossings by the dotted lines in Fig. 4.5 are not counted in the Simon invariant we deduce that $\mathcal{L}(g) = \pm 1$. Since f and g are homologous it follows that $\mathcal{L}(f) = \mathcal{L}(g) = \pm 1$.



FIGURE 4.5.

(6) ("only if" part) Note that the cube of $(1 \ 2)(3 \ 4 \ 5)$ equals $(1 \ 2)$. Therefore $(1 \ 2)(3 \ 4 \ 5)$ -symmetric spatial embeddings of K_5 are $(1 \ 2)$ -symmetric. Then we have the result by (5) ("only if" part).

Proof of Theorem 1.10. (2) and (6) Let $f_{m,\pm}$ be the spatial embeddings of $K_{3,3}$ as in Fig. 4.6. Note that $\mathcal{L}(f_{m,\pm}) = 4m \pm 1$, which may be any odd number. A $\pi/2$ rotation maps $f_{m,\pm}(K_{3,3})$ onto its mirror image. Therefore $f_{m,\pm}$ is $(1 \ 4 \ 2 \ 5)(3 \ 6)$ symmetric. Also, a π rotation maps $f_{m,\pm}(K_{3,3})$ onto itself and hence $f_{m,\pm}$ is $(1 \ 2)(4 \ 5)$ -symmetric.



FIGURE 4.6.

(3) (4) (5) Let f_m be the spatial embedding of $K_{3,3}$ as in Fig. 4.7. Note that $\mathcal{L}(f_m) = 2m + 1$. Observe that $f_m(K_{3,3})$ is contained in an unknotted Möbius strip in \mathbb{S}^3 containing 2m + 1 half twists. Then by considering a suitable ambient isotopy of \mathbb{S}^3 that preserves the Möbius strip setwise we see that f_m is $(1 \ 4 \ 2 \ 5 \ 3 \ 6)$ -symmetric. Note that the square of $(1 \ 4 \ 2 \ 5 \ 3 \ 6)$ is $(1 \ 2 \ 3)(4 \ 5 \ 6)$. Therefore f_m is $(1 \ 2 \ 3)(4 \ 5 \ 6)$ -symmetric. It is also easy to see that f_m is $(1 \ 5)(2 \ 6)(3 \ 4)$ -symmetric. Since $(1 \ 5)(2 \ 6)(3 \ 4)$ is conjugate to $(1 \ 4)(2 \ 5)(3 \ 6)$ we have the result.



FIGURE 4.7.

(1) ("if" part) Let $f_{m,\pm}$ be the spatial embedding of $K_{3,3}$ as in Fig. 4.8. Note that $\mathcal{L}(f_{m,\pm}) = 6m \pm 1$. A $2\pi/3$ rotation around the axis through the vertices 5, 4 and 6 maps $f_{m,\pm}(K_{3,3})$ onto itself, so $f_{m,\pm}$ is (1 2 3)-symmetric.

(7) ("if" part) and (8) ("if" part) Let f_{\pm} be the spatial embedding of $K_{3,3}$ as in Fig. 4.9. Note that $\mathcal{L}(f_{\pm}) = \pm 1$. By considering the reflection of \mathbb{S}^3 with respect to the 2-sphere containing the vertices 3, 4, 5 and 6, we find that f_{\pm} is (1 2)-symmetric. Then by composing this reflection and a $2\pi/3$ rotation around the axis through the vertices 1, 2 and 3, we conclude that f_{\pm} is (1 2)(4 5 6)-symmetric.

(1) ("only if" part) Suppose that f is a (1 2 3)-symmetric spatial embedding of $K_{3,3}$. We will show that $\alpha_{\omega}(f)$ is a multiple of 3. Then by Lemma 2.1 we have



FIGURE 4.8.



FIGURE 4.9.

the result. There exist six 6-cycles and nine 4-cycles in $K_{3,3}$. By the permutation (1 2 3) they are divided into the following five orbits:

$$\begin{array}{c} [1 \ 4 \ 2 \ 5 \ 3 \ 6] \mapsto [2 \ 4 \ 3 \ 5 \ 1 \ 6] \mapsto [3 \ 4 \ 1 \ 5 \ 2 \ 6] \mapsto [1 \ 4 \ 2 \ 5 \ 3 \ 6] \\ [1 \ 6 \ 2 \ 5 \ 3 \ 4] \mapsto [2 \ 6 \ 3 \ 5 \ 1 \ 4] \mapsto [3 \ 6 \ 1 \ 5 \ 2 \ 4] \mapsto [1 \ 6 \ 2 \ 5 \ 3 \ 4] \\ [1 \ 4 \ 2 \ 5] \mapsto [2 \ 4 \ 3 \ 5] \mapsto [3 \ 4 \ 1 \ 5] \mapsto [1 \ 4 \ 2 \ 5] \\ [1 \ 4 \ 2 \ 5] \mapsto [2 \ 4 \ 3 \ 5] \mapsto [3 \ 4 \ 1 \ 5] \mapsto [1 \ 4 \ 2 \ 5] \\ [1 \ 5 \ 2 \ 6] \mapsto [2 \ 5 \ 3 \ 6] \mapsto [3 \ 5 \ 1 \ 6] \mapsto [1 \ 4 \ 2 \ 5] \\ [1 \ 5 \ 2 \ 6] \mapsto [2 \ 5 \ 3 \ 6] \mapsto [3 \ 5 \ 1 \ 6] \mapsto [1 \ 5 \ 2 \ 6] \\ [1 \ 6 \ 2 \ 4] \mapsto [2 \ 6 \ 3 \ 4] \mapsto [3 \ 6 \ 1 \ 4] \mapsto [1 \ 6 \ 2 \ 4] \end{array}$$

Note that each orbit contains exactly three cycles and they are mapped onto the same knot under f. Therefore they have the same a_2 and we have the desired conclusion.

(7) ("only if" part) Suppose that f is a (1 2)-symmetric spatial embedding of $K_{3,3}$. Then by Theorem 4.1 there exist a (1 2)-symmetric spatial embedding g of $K_{3,3}$ which is homologous to f and a periodic self-homeomorphism φ of \mathbb{S}^3 such that $g \circ (1 2) = \varphi \circ g$. By the same reason as in the proof of Theorem 1.9 (5) ("only if" part), φ is orientation-reversing. Then by Theorem 4.2, Fix(φ) is homeomorphic to \mathbb{S}^0 or \mathbb{S}^2 . Since Fix(φ) contains at least four points g(3), g(4), g(5) and g(6), it is homeomorphic to \mathbb{S}^2 . Then Fix(φ) $\cap g(K_{3,3})$ is the induced subgraph of the vertices g(3), g(4), g(5) and g(6). By considering the regular projection which is almost perpendicular to Fix(φ) and by the same reason as in the proof of Theorem 1.9 (5) ("only if" part), we deduce that $\mathcal{L}(g) = \pm 1$ (see Fig. 4.10). Since f and gare homologous it follows that $\mathcal{L}(f) = \mathcal{L}(g) = \pm 1$.



FIGURE 4.10.

(8) ("only if" part) Note that the cube of $(1\ 2)(4\ 5\ 6)$ equals $(1\ 2)$. Therefore $(1\ 2)(4\ 5\ 6)$ -symmetric spatial embeddings of $K_{3,3}$ are $(1\ 2)$ -symmetric. Then we have the result by (7) ("only if" part).

Proof of Theorem 1.11. Let G be K_5 or $K_{3,3}$. For a spatial embedding f of G and a self-homeomorphism φ of \mathbb{S}^3 we have $\mathcal{L}(\varphi \circ f) = \mathcal{L}(f)$ if φ is orientation-preserving and $\mathcal{L}(\varphi \circ f) = -\mathcal{L}(f)$ if φ is orientation-reversing. By combining this fact with Lemma 3.2 and Remark 3.3, we have the results.

Remark 4.3. (1) All symmetries shown by various examples in this paper are realized by periodic self-homeomorphisms of \mathbb{S}^3 . Hence all symmetric spatial graphs in this paper are rigidly symmetric.

(2) There are alternative proofs of the "only if" parts of Theorems 1.9 (1) and 1.10 (1), based on Theorem 4.1 and the Smith conjecture [8], just as the proofs of Theorems 1.9 (5) and (6) and 1.10 (7) and (8) given above are based on Theorems 4.1 and 4.2. However, as we said before, the latter theorems are consequences of deep results of 3-manifold topology. The proof of the "only if" parts of Theorems 1.9 (1) and 1.10 (1) given above are elementary. To find elementary proofs of the "only if" parts of Theorems 1.9 (5) and (6) and 1.10 (7) and (8) is an open problem.

References

- [1] E. Flapan, Symmetries of Möbius ladders, Math. Ann. 283 (1989), 271–289.
- [2] E. Flapan, When topology meets chemistry. A topological look at molecular chirality. Outlooks. Cambridge University Press, Cambridge; Mathematical Association of America, Washington, DC, 2000.
- [3] M. E. Kidwell, Two types of amphichiral links, J. Knot Theory Ramifications 15 (2006), 651–658.
- [4] P. Kirk and C. Livingston, Vassiliev invariants of two component links and the Casson-Walker invariant, *Topology* 36 (1997), 1333–1353.

- [5] D. Kuck and A. Schuster, Centrohexaindan, the first hydrocarbon with topologically nonplanar molecular structure, Angew. Chem. Int. Ed. Engl. 27 (1988), 1192–1194.
- [6] C. Kuratowski, Sur le probléme des courbes gauches en topologie, (French) Fund. Math. 15 (1930), 271–283.
- [7] C. Livingston, Enhanced linking numbers, Amer. Math. Monthly 110 (2003), 361–385.
- [8] J. W. Morgan and H. Bass, ed., The Smith conjecture, Papers presented at the symposium held at Columbia University, New York, 1979. Edited by John W. Morgan and Hyman Bass. Pure and Applied Mathematics, **112**. Academic Press, Inc., Orlando, FL, 1984.
- [9] T. Motohashi and K. Taniyama, Delta unknotting operation and vertex homotopy of graphs in R³, KNOTS '96 (Tokyo), 185–200, World Sci. Publ., River Edge, NJ, 1997.
- [10] H. Murakami and Y. Nakanishi, On a certain move generating link-homology, Math. Ann. 284 (1989), 75–89.
- [11] R. Nikkuni, Achirality of spatial graphs and the Simon invariant, Intelligence of Low Dimensional Topology 2006 (Hiroshima), 239-243, Ser. Knots Everything, 40, World Sci. Publ., Hackensack, NJ, 2007.
- [12] L. A. Paquette and M. Vazeux, Threefold transannular epoxide cyclization. Synthesis of a heterocyclic 17-hexaquinane, *Tetrahedron Lett.* 22 (1981), 291–294.
- [13] K. S. Sarkaria, A one-dimensional Whitney trick and Kuratowski's graph planarity criterion, Israel J. Math. 73 (1991), 79–89.
- [14] H. E. Simmons and J. E. Maggio, Synthesis of the first topologically non-planar molecule, *Tetrahedron Lett.* 22 (1981), 287–290.
- [15] J. Simon, Topological chirality of certain molecules, Topology 25 (1986), 229–235.
- [16] J. Simon, A topological approach to the stereochemistry of nonrigid molecules, Graph theory and topology in chemistry (Athens, Ga., 1987), 43–75, Stud. Phys. Theoret. Chem., 51, Elsevier, Amsterdam, 1987.
- [17] R. Shinjo and K. Taniyama, Homology classification of spatial graphs by linking numbers and Simon invariants, *Topology Appl.* **134** (2003), 53–67.
- [18] P. A. Smith, Transformations of finite period. II, Ann. of Math. (2) 40 (1939), 690-711.
- [19] T. Soma, Symmetries of minimum graphs in the 3-sphere, J. Knot Theory Ramifications 7 (1998), 651–657.
- [20] K. Taniyama, Link homotopy invariants of graphs in R³, Rev. Mat. Univ. Complut. Madrid 7 (1994), 129–144.
- [21] K. Taniyama, Cobordism, homotopy and homology of graphs in R³, Topology 33 (1994), 509–523.
- [22] K. Taniyama, Homology classification of spatial embeddings of a graph, *Topology Appl.* 65 (1995), 205–228.
- [23] D. M. Walba, Stereochemical topology, Chemical applications of topology and graph theory (Athens, Ga., 1983), 17–32, Stud. Phys. Theoret. Chem., 28, Elsevier, Amsterdam, 1983.
- [24] D. M. Walba, R. M. Richards and R. C. Haltiwanger, Total synthesis of the first molecular Möbius strip, J. Am. Chem. Soc. 104 (1982), 3219–3221.
- [25] W. T. Wu, On the isotopy of a complex in a Euclidean space. I, Sci. Sinica 9 (1960), 21-46.

DEPARTMENT OF MATHEMATICS, SCHOOL OF ARTS AND SCIENCES, TOKYO WOMAN'S CHRISTIAN UNIVERSITY, 2-6-1 ZEMPUKUJI, SUGINAMI-KU, TOKYO 167-8585, JAPAN

E-mail address: nick@lab.twcu.ac.jp

DEPARTMENT OF MATHEMATICS, SCHOOL OF EDUCATION, WASEDA UNIVERSITY, NISHI-WASEDA 1-6-1, SHINJUKU-KU, TOKYO, 169-8050, JAPAN

E-mail address: taniyama@waseda.jp

16