Vacuum driven accelerated expansion*

Bogusław Broda[†], Piotr Bronowski[‡], Marcin Ostrowski[§] and Michał Szanecki[¶]

Department of Theoretical Physics, University of Łódź

Pomorska 149/153, 90-236 Łódź, Poland

Key Words Accelerated expansion, cosmological constant, quantum vacuum energy, Casimir effect, dark energy.

PACS 95.36.+x, 04.62.+v, 98.80.Es

It has been shown that an improved estimation of quantum vacuum energy can yield not only theoretically acceptable but also experimentally realistic results. Our idea consists in a straightforward extraction of gravitationally interacting part of the full quantum vacuum energy by means of gauge transformations. The implementation of the idea has been performed in the euclidean version of the formalism of effective action, in the language of Schwinger's proper time and the Seeley–DeWitt heat kernel expansion, in the background of the Friedmann–Robertson–Walker–Lemaître geometry.

1 Introduction

There are three famous problems in modern physics and cosmology, which can, in principle, be treated as independent ones or, just the opposite, (all or any two of them) as mutually related:

- accelerated expansion of the Universe [1, 2] (proven by astrophysicists);
- cosmological constant or dark energy (very small, though non-vanishing) [3, 4];
- quantum vacuum energy density (theoretically very huge, experimentally very small) [5, 6].

^{*}Devoted to the memory of professor Ryszard Rączka on the occasion of the 12-th anniversary of his death.

[†]bobroda@uni.lodz.pl

[‡]bronowski@gmail.com

[§]m.ostrowski@merlin.phys.uni.lodz.pl

[¶]michalszanecki@wp.pl

The accelerated expansion of the Universe is by now rather a well-established by astronomical observations fact, in particular, Supernovae Ia data [1]. That mysterious phenomenon still awaits an explanation. There are dozens of candidates for the solution of the problem (a number of approaches is reviewed in [2]). One of the possible solutions and, in principle, the simplest one is the introduction of the cosmological constant Λ . Another solution (or the same, it depends on the point of view) is quantum vacuum energy [5]. That solution is, in a sense, traditional because it seems to be theoretically the most natural and simple one, and it was proposed long ago. Its "only drawback" is the fact that, as it seems, it does not work well.

The cosmological constant problem [3] troubles physicists from nearly the very beginning of the existence of general relativity. There are also dozens of candidates for the solution of this problem (they are even catalogued in [7]). Unfortunately, explanation of the accelerated expansion by the vanishingly small value of the cosmological constant shifts only the problem rather than solves it.

Traditional approach to the issue of the cosmological constant Λ uses quantum vacuum energy as a source of the origin of that quantity. But still the mechanism, being very appealing, does not work, as it seems, properly. It appears that the traditionally calculated, Casimir-like value of quantum vacuum energy density is definitely too big than accepted, and two orders of orders too big than required! Entirely independently of the problem of the accelerated expansion and of the problem of the cosmological constant such a drastically huge value of the vacuum energy density is a serious problem in itself. That means that one should solve the quantum vacuum energy problem independently whether it could or should be later related to the accelerated expansion and the cosmological constant or not. There is quite a numerous collection of potential explanations of the above issues in literature. They are mainly given in the context of the dark energy [2]. It is not our intention to list or review them but it seems to be useful for our further purposes to mention at least some.

One of early standard ideas was to lower the ultraviolet cutoff scale Λ_{UV} using supersymmetry arguments. It helps a bit, but only a little bit, if we want to be in accordance with current experimental facts (roughly, it cuts the order of discrepancy by two [3]).

As another, rather a radical solution one should mention the idea assuming that quantum vacuum energy does not, for one or another reason, influence on gravity. For example, the authors of [8], referring to the Casimir effect, rule out the possibility that the observed cosmological constant arises from the zero-point energy which is made finite by a suitable cut-off. In [6] it is claimed that the huge contribution of the zero-point motion of the quantum fields to the vacuum energy is exactly cancelled by the higher-energy degrees of freedom of the quantum vacuum (automatic compensation of zero-point energy). Or simply, "zero-point energy does not gravitate in vacuum". And finally, the author of the paper [9] provocatively asks: "...Why the vacuum does not gravitate ..."?

It appears, and the aim of our paper is to show it, that, in principle, it is possible to reasonably estimate the value of quantum vacuum energy obtaining an experimentally acceptable result. Moreover, the result is not only realistic in itself (the quantum vacuum energy is not huge) but experimentally expected as well. Our approach does not appeal to any more or less clever, arbitrary or exotic assumption, and this is, in our opinion, its main advantage. Just the opposite, our idea is supposed to adhere to standard quantum

field theory formalism as closely as possible. In fact, our proposal only works provided standard philosophy of quantum field theory is carefully taken into account.

2 Quantum vacuum energy

The well-known, standard (but, regrettably, not properly working) approach to estimation of the quantum vacuum energy density ϱ_{vac} calculates the Casimir-like energy density for the whole Universe. The result of such a calculation for a single bosonic scalar mode (in mass units) is [3]

$$\varrho_{\text{vac}} = \frac{1}{2} \int_{0}^{\Lambda_{\text{uv}}} \frac{4\pi}{(2\pi\hbar)^3 c} \sqrt{(mc)^2 + k^2} \, k^2 dk, \tag{1}$$

where m is the mass of the mode. For a large ultraviolet (UV) momentum cutoff $\Lambda_{\rm UV}$

$$\varrho_{\rm vac} \approx \frac{1}{(4\pi)^2} \frac{\Lambda_{\rm uv}^4}{\hbar^3 c}.$$
 (2)

Setting $\Lambda_{UV} = \Lambda_P$, where Λ_P is the Planck momentum,

$$\Lambda_{\rm P} = \sqrt{\frac{\hbar c^3}{G}} \approx 6.5 \,\mathrm{kg} \,\mathrm{m/s},$$
 (3)

here G is the newtonian gravitational constant, we obtain

$$\varrho_{\text{vac}}^{\text{P}} \approx \frac{c^5}{(4\pi)^2 \hbar G^2} \approx 3.4 \times 10^{94} \,\text{kg/m}^3,$$
(4)

an enormously huge value, whereas the experimentally estimated value is of the order of, the so-called, critical density of the Universe $\varrho_{\rm crit}=3H_0^2/8\pi G~(\approx 10^{-26}\,{\rm kg/m^3})$, where H_0 is the present day Hubble expansion rate, i.e. more than 120 orders less! Lowering $\Lambda_{\rm uv}$ to, say, the supersymmetry scale $\Lambda_{\rm susy}\approx 1\,{\rm TeV/c}$, only slightly improves the situation, namely, $\varrho_{\rm vac}^{\rm susy}\approx 1.5\times 10^{30}\,{\rm kg/m^3}$, but it does not change the general impression that the whole calculation is principally erroneous. Therefore, as a desperate response to this dramatic situation, the earlier mentioned idea has emerged that gravitational field is insensitive to quantum vacuum fluctuations, yielding $\varrho_{\rm vac}^0=0$.

The both described, extreme approaches, the ordinary, purely Casimir-like calculation and the insensitiveness idea actually yield entirely, from experimental point of view, incorrect results. That fact should become obvious also from theoretical point of view for the following reasons. First of all, we observe that the ordinary, purely Casimir-like calculation of quantum vacuum energy should not give any measurable contribution to gravitational (or any other) field "by construction". Actually, any classically analysed process (interaction) is an approximation of a quantum one. Therefore, we are allowed to reason in the language of Feynman diagrams. Let us observe that the ordinary, purely Casimir-like calculus gives rise to contributions coming from closed loops (see, Fig. 1)

without any external lines. They do not influence gravitational field because this possibility has not been taken into account by virtue of the construction, i.e. there are no "classical" external lines establishing contact of the internal "matter" loops with the outer gravitational field.




Fig.1: A single closed loop representing the ordinary, purely Casimir-like contribution of a free matter field to the effective action.

But one can easily correct this trivial result performing improved calculations. Namely, one should consider contributions coming from closed "matter" loops with classical external gravitational lines attached. Such an approach is not only in full accordance with paradigms of standard quantum field theory, without any additional assumptions, but also, moreover, it would bring us to realistic results.

Summarizing, roughly speaking, we should calculate all quantum vacuum fluctuations of a matter field in an external classical gravitational field, retaining the most divergent parts (as it will become clear in the next section), and next we should discard the term without gravitational field.

In this section, we have sketched our idea of a proposed estimation of the quantum vacuum energy. A concrete implementation of the idea will be introduced in the following section. We intentionally stress the difference between the idea and the implementation because we believe that the proposed implementation of the estimation is not final nor unique.

3 Implementation of the estimation

Full quantum contribution coming from a single (non-self-interacting) mode is included in the effective action of the form [10]

$$S_{\text{eff}} = \pm \frac{\hbar}{2} \log \det \mathcal{D},\tag{5}$$

where \mathcal{D} is a non-negative, second-order differential operator, in general, with classical external fields, and the upper (plus) sign corresponds to bosonic statistics whereas the lower (minus) one corresponds to fermionic statistics, respectively. For simplicity, we work in the euclidean framework throughout. Since (5) is UV divergent we should regularize it, analogously to (1). The most convenient and systematic way to control infinities in Eq. (5) is to use Schwinger's proper-time method. In this approach [10, 11] we can formally rewrite (5) as

$$S_{\text{eff}} = \pm \frac{\hbar}{2} \text{Tr} \log \mathcal{D} = \mp \frac{\hbar}{2} \int_{0}^{\infty} \frac{\mathrm{d}s}{s} \, \text{Tr} \, e^{-s\mathcal{D}}. \tag{6}$$

Now, the UV regularized version of (6) is

$$S_{\text{eff}}^{\varepsilon} = \mp \frac{\hbar}{2} \int_{s}^{\infty} \frac{\mathrm{d}s}{s} \operatorname{Tr} e^{-s\mathcal{D}} \equiv \mp \frac{\hbar}{2} \int_{s}^{\infty} \frac{\mathrm{d}s}{s} T_{\mathcal{D}}(s), \tag{7}$$

where ε is an UV cutoff in the units: length to the power two. Next, we apply the following Seeley–DeWitt heat kernel expansion [10, 12]:

$$T_{\mathcal{D}}(s) \equiv \int t(s; x) \sqrt{g} \, \mathrm{d}^4 x \,,$$
 (8)

where, in four dimensions without boundary,

$$t(s;x) = \frac{1}{(4\pi s)^2} \sum_{n=0}^{\infty} a_n(x) s^n.$$
 (9)

A physical motivation to use the external lines approach is as follows. A generally accepted and also our main assumption is the statement that quantum vacuum fluctuations of "matter" (in fact, all non-gravitational) fields somehow yield non-zero (effective) cosmological constant (or dark energy). In other words, the vacuum quantum fluctuations influence on gravitational field. In technical terms, the influence, in the language of quantum field theory, is established by lines of Feynman diagrams. Since we are working in the domain of classical gravity, graviton lines are supposed to be classical external lines, whereas (quantum) "matter" fields are confined to single loops.

The full expansion (7) corresponds to all one-loop Feynman diagrams for a matter field, those depicted in Fig. 2 and also that in Fig. 1. But the previously discussed, ordinary, purely Casimir-like vacuum diagram in Fig. 1 should be discarded as a trivial one because, according to our earlier discussion, by construction, it does not contain any coupling to an external gravitational field. That trivial vacuum diagram appears in the first coefficient of the Seeley-DeWitt expansion (9), $a_0(x)$. It is obvious because only $a_0(x)$ survives the vanishing external field limit. For any external gravitational (and not only gravitational) non-vanishing or vanishing field, we have $a_0(x) = 1$.




Fig.2: Closed matter loops influencing gravitational field via attached classical external lines.

Other $a_n(x)$'s (for n > 0) contain various powers and derivatives of curvature with dimensionality governed by n. In particular, $a_1(x) = \frac{1}{6}R$ for an ordinary massless scalar mode, where R is the scalar curvature, and it finitely renormalizes or induces [13] (dependently on the point of view) the classical Hilbert-Einstein action of gravity. For that

mode, the philosophy of the induced gravity yields, by virtue of (7)–(9),

$$S_{\text{ind}} = -\frac{\hbar}{2} \frac{1}{\varepsilon} \frac{1}{(4\pi)^2} \int \frac{1}{6} R \sqrt{g} \, d^4 x$$

$$= -\frac{\hbar}{12} \frac{1}{L_P^2} \frac{1}{(4\pi)^2} \int R \sqrt{g} \, d^4 x$$

$$= -\frac{1}{12\pi} \frac{c^3}{16\pi G} \int R \sqrt{g} \, d^4 x,$$
(10)

where the planckian UV cutoff has been assumed, i.e.

$$\varepsilon = L_{\rm P}^2 = \frac{\hbar G}{c^3}.\tag{11}$$

Hence, amazingly, the induced coupling constant for a single mode is approximately 12π times less than the standard classical value! The next term, $a_2(x)$, and also further terms, yields genuine quantum corrections to the classical theory, and thus is uninteresting for us.

As is commonly expected, the (effective) cosmological constant Λ or dark energy can be induced by the zeroth term, a_0 , and therefore we will concentrate on that term henceforth. The zeroth term a_0 yields, according to (7)–(9) and (11), Casimir-like contribution of the form

$$S_{\text{Cas}} = \mp \frac{\hbar}{2} \frac{1}{2\varepsilon^2} \frac{1}{(4\pi)^2} \int \sqrt{g} \, d^4 x$$

$$= \mp \frac{\hbar}{4} \frac{1}{L_{\text{P}}^4} \frac{1}{(4\pi)^2} \int \sqrt{g} \, d^4 x$$

$$= \mp \frac{1}{4} \frac{c^6}{(4\pi)^2 \hbar G^2} \int \sqrt{g} \, d^4 x.$$
(12)

In the flat space limit it corresponds (in mass density units) to the value exactly 4 times less than that calculated earlier in (4). This unessential difference is coming from different regularization procedures. Besides, we should remember that hamiltonian and lagrangian are different objects. Since, by assumption, we only perform estimations we may not care of that difference, and concentrate only on orders of values. In a sense, we have derived a covariant version of the result (4). According to our strategy we have to extract from Eq. (12) only the part corresponding to gravitational field.

For technical simplicity, but in accordance with experimental realm, we can take the metric of the spatially flat Friedmann–Robertson–Walker–Lemaître (FRWL) form with the scale factor a(t),

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0\\ 0 & a^2(t) & 0 & 0\\ 0 & 0 & a^2(t) & 0\\ 0 & 0 & 0 & a^2(t) \end{pmatrix}. \tag{13}$$

For calculational purposes, let us assume that the coordinate time t=0 corresponds to the present moment, and normalize the coordinates according to the equality

$$a(0) = 1. (14)$$

Now, power-series expanding around t = 0, we have

$$a(t) = a(0) + \dot{a}(0)t + \ddot{a}(0)t^{2} + O(t^{3})$$

$$= 1 + H_{0}t - \frac{1}{2}q_{0}H_{0}^{2}t^{2} + O(t^{3}),$$
(15)

where the present day Hubble expansion rate,

$$H_0 \equiv \frac{\dot{a}(0)}{a(0)} = \dot{a}(0),$$

and q_0 is the present day deceleration parameter,

$$q_0 \equiv -H_0^{-2} a^{-1}(0) \ddot{a}(0) = -H_0^{-2} \ddot{a}(0).$$

Hence

$$\sqrt{g} = \left[a^2(t)\right]^{\frac{3}{2}} = \left[1 + 2H_0t + (1 - q_0)H_0^2t^2 + O(t^3)\right]^{\frac{3}{2}}.$$
 (16)

It is of vital importance for our further considerations to show that the second term in (16), linear in t, can be discarded by virtue of gauge symmetry. A general, model-independent discussion of this fact is given in [14]. Physically, such a potential possibility corresponds to the obvious fact that not any perturbation of flat metric represents a genuine gravitational field but only that which is gauge nontrivial. The direct proof goes as follows. Infinitesimal gauge transformations around flat metric are given by

$$\delta g_{\mu\nu} = \partial_{\mu} \xi_{\nu} + \partial_{\nu} \xi_{\mu},\tag{17}$$

where $\xi_{\mu}=(\xi\equiv\xi_0,\ \xi_i)$ are gauge parameters. Explicitly, the first equation is

$$\delta g_{00} = 2\dot{\xi} = 0,\tag{18}$$

because $g_{00} = 1$ should be left undisturbed. A general solution of Eq. (18) is then $\xi = \xi(\mathbf{x})$. The second equation is of the form

$$\delta g_{0i} \equiv \delta g_{i0} = \dot{\xi}_i + \partial_i \xi = 0, \tag{19}$$

because also $g_{0i} = 0$ should be left intact. Hence $\dot{\xi}_i = -\partial_i \xi(\mathbf{x})$, and consequently

$$\xi_i = -t \,\partial_i \xi(\mathbf{x}) + \eta_i(\mathbf{x}). \tag{20}$$

For purely spatial indices we have

$$\delta g_{ij} = \partial_i \xi_j + \partial_j \xi_i
= -2t \, \partial_{ij} \xi(\mathbf{x}) + \partial_i \eta_i(\mathbf{x}) + \partial_j \eta_i(\mathbf{x}). \tag{21}$$

Now, we put

$$\delta g_{ij} = \begin{cases} 0, & \text{for } i \neq j \\ f(t), & \text{for } i = j. \end{cases}$$
 (22)

From Eq. (21) it immediately follows that the most general function f(t) which can be gauged away is linear in t. As a final, sufficiently general solution of our equation we assume the particular form

$$\xi(\mathbf{x}) = \frac{1}{2} \sum_{i,j=1}^{3} \xi_{ij} x^{i} x^{j}, \quad \eta(\mathbf{x}) = 0,$$
(23)

where the constant matrix ξ_{ij} , in view of Eq. (22), should be diagonal (even scalar), i.e. $\xi_{ij} = \frac{1}{2}H_0\delta_{ij}$, after appropriate normalization. Then, the solution of our problem is of the form

$$\xi_{\mu} = \left(\frac{1}{4}H_0\mathbf{x}^2, -\frac{1}{2}H_0tx^i\right). \tag{24}$$

Gauging away the term linear in t in Eq. (16), using (24), we have for small t

$$\sqrt{g} = 1 + \frac{3}{2}(1 - q_0)H_0^2 t^2 + O(t^3). \tag{25}$$

Since the integrand in Eq. (12) is only t-dependent we can painless divide it by the spatial volume $\int d^3x$. Instead, dividing by the time coordinate, at least for small t, is nothing but an averaging procedure with respect to t. As our analysis is perturbative in the time t, the longer the time the smaller the reliability of our analysis. The shortest possible time, in the realm of quantum field theory, is $t = T_P$ (the Planck time). Therefore, time averaging $\langle \cdot \rangle_t$ around present moment (t = 0) is given by the formula

$$\langle \cdot \rangle_t \equiv \lim_{T \to T_{\rm P}} \frac{1}{T} \int_0^T \mathrm{d}t \; (\cdot).$$
 (26)

Thus, an estimated density is according to Eq. (12) of the order

$$\varrho = \mp \frac{1}{4} \frac{c^5}{(4\pi)^2 \hbar G^2} \lim_{T \to T_P} \frac{1}{T} \int_0^T dt \, (\sqrt{g} - 1)$$

$$\approx \mp \frac{1}{4} \frac{c^5}{(4\pi)^2 \hbar G^2} \frac{1}{2} (1 - q_0) H_0^2 T_P^2,$$
(27)

where the substraction in the parentheses in the first term corresponds to discarding gravitationally non-interacting part of the effective action. Equivalently, in Eq. (5) one should consider $\det \mathcal{D}/\det \mathcal{D}_0$ instead of $\det \mathcal{D}$, where \mathcal{D}_0 is a flat space version of \mathcal{D} . Or, in another language, the subtraction corresponds to normalization of the functional measure. Therefore, the subtraction is by no means, as it could seem, an *ad hoc* procedure. Since $T_P^2 = \hbar G/c^5$, and $H_0^2 = \frac{8}{3}\pi G \varrho_{\rm crit}$, we finally obtain

$$\varrho \approx \mp \frac{1}{48\pi} (1 - q_0) \varrho_{\text{crit}}.$$
 (28)

Amazingly, Eq. (28) predicts a highly reasonable result. Inserting $q_0 \approx -0.7$, which is phenomenologically a realistic assumption [15], yields the following numerical result:

$$\varrho \approx \mp 0.01 \ \varrho_{\rm crit}.$$
 (29)

The experimental value is roughly $\varrho_{\rm exp} \approx 0.76 \ \varrho_{\rm crit}$, therefore we finally have

$$\varrho \approx \mp 0.01 \varrho_{\rm exp} \tag{30}$$

per a single mode, which is a very good estimation in our opinion. Taking into account the remarks directly following Eq. (12), ϱ could be just as well 0.04 $\varrho_{\rm exp}$. We should bear in mind that our analysis is an estimation, and we are only interested in the orders of values.

4 Conclusions

Using coordinate gauge freedom we have managed to extract from the full quantum vacuum term induced in an external classical gravitational background by a fluctuating mode of a matter field the fraction corresponding to interaction with gravitational field. An explicit calculus has been performed in the framework of the spatially flat FRWL geometry. The value of the contribution coming from a single mode, which appears to be of the order of one hundredth of the experimentally expected value, seems to be very promising. Thus, the old, primary expectation that quantum vacuum fluctuations could be the source of the cosmological constant Λ does not seem to be unjustified.

Although, our idea, in its definite form (as well as the derivation), is novel, actually it emerges in many papers as more or less explicitly expressed thoughts. For example, the author of [9] claims that zero-point energy gravitates in some environments and not in vacuum. The author of [16] is even closer to our point of view claiming that we should consider the fluctuations in the vacuum energy density. Interestingly, Fig. 1(a) of [17] suits our point of view excellently.

Finally, we would like to comment on the celebrated Casimir effect and its possible relation to our approach. In fact, it happens that the Casimir effect is often referred to by many authors (including us) in the context of quantum vacuum. In particular, it is usually claimed that reality of quantum vacuum energy is proved by experimental confirmation of the Casimir effect itself. Rather surprisingly, Casimir forces are correctly described by a crude count in the spirit of the derivation of Eq. (1), without any external lines present. Therefore, we have an apparent paradox. On the one hand the ordinary Casimir-like calculation yields reasonable results in the case of the very Casimir effect. On the other hand it yields useless results in the context of gravity. The resolution is simple. The ordinary Casimir-like calculation is a simplified approach that (perhaps accidentally) somehow works. It appears that a deeper approach to the Casimir effect [18] is in spirit similar to our considerations. In particular, it uses external lines and a subtraction of the ordinary Casimir-like vacuum diagram (compare Fig. 3 in [18] and our Fig. 2).

We would like to stress that we do not claim that we have found a final solution of the problem of the accelerated expansion. We are aware that there are a lot of more or less sensible competing proposals in that area. All of them, including ours, have some drawbacks. Our approach should be just interpreted in this context, as a voice in the discussion, indicating a direction for a possible further study.

Acknowledgments

This work was supported in part by the Polish Ministry of Science and Higher Education Grant PBZ/MIN/008/P03/2003 and by the University of Łódź grant.

References

- [1] A. G. Riess et al., Observational evidence from supernovae for an accelerating universe and a cosmological constant, Astronom. J. **116**, 1009–1038 (1998).
 - S. Perlmutter et al., Measurements of Ω and Λ from 42 high-redshift Supernovae, Astrophys. J. **517**, 565–586 (1999).
- [2] E. J. Copeland, M. Sami, and S. Tsujikawa, Dynamics of dark energy, Int. J. Mod. Phys. D 15, 1753–1936 (2006); hep-th/0603057.
- [3] S. Weinberg, The cosmological constant problem, Rev. Mod. Phys. 61, 1–23 (1989).
- [4] S. M. Carroll, The cosmological constant, Liv. Rev. Rel. 4 (2001); astro-ph/0004075.
 T. Padmanabhan, Cosmological constant—the weight of the vacuum, Phys. Rep. 380, 235–320 (2003); hep-th/0212290.
 - T. Padmanabhan, Dark energy: mystery of the millennium, AIP Conf. Proc. 861, 179–196 (2006), Albert Einstein Century International Conference; astro-ph/0603114.
- [5] Y. B. Zel'dovich, Cosmological constant and elementary particles, JETP Lett. **6**, 316–317 (1967), translated from Zh. Eksp. Teor. Fiz., Pis'ma Redaktsiyu, **6** 883–884 (1967).
- [6] G. E. Volovik, Vacuum energy: myths and reality, Int. J. Mod. Phys. D 15, 1987–2010 (2006); gr-qc/0604062.
 - G. E. Volovik, Cosmological constant and vacuum energy, Ann. Phys. (Leipzig) **14**, 165–176 (2005); gr-qc/0405012.
- [7] S. Nobbenhuis, Categorizing different approaches to the cosmological constant problem, Found. Phys. **36**, 613–680 (2006); gr-qc/0411093.
- [8] G. Mahajan, S. Sarkar, and T. Padmanabhan, Casimir effect confronts cosmological constant, Phys. Lett. B **641**, 6–10 (2006); astro-ph/0604265.
- [9] J. Polchinski, The cosmological constant and the string landscape, hep-th/0603249.
- [10] B. S. DeWitt, Quantum field theory in curved spacetime, Phys. Rep. 19, 295–357 (1975).
 - B. DeWitt, The Global Approach to Quantum Field Theory, (Clarendon Press, 2003).
- [11] K. Huang, Quarks, Leptons and Gauge Fields, (World Scientific, 1992), Chapt. 10.4.
 P. Ramond, Field Theory: A Modern Primer, (Addison-Wesley, 2001), Chapt. 3.
 See also:

- T. P. Cheng and L. F. Li, Gauge Theory of Elementary Particle Physics, (Oxford University Press, 2000), Chapt. 6.4.
- S. Weinberg, The Quantum Theory of Fields Volume 2: Modern Applications, (Cambridge University Press, 1996), Chapt. 16.2.
- [12] R. D. Ball, Chiral gauge theory, Phys. Rep. **182**, 1–186 (1989).
- [13] M. Visser, Sakharov's induced gravity: a modern perspective, Mod. Phys. Lett. A 17, 977–992 (2002); gr-qc/0204062.
- [14] I. L. Shapiro, Effective action of vacuum: semiclassical approach, 0801.0216 [gr-qc], Chapt. 3.1.
- [15] J. M. Virey, P. Taxil, A. Tilquin, A. Ealet, C. Tao, and D. Fouchez, On the determination of the deceleration parameter from Supernovae data, Phys. Rev. D **72**, 061302 (2005); astro-ph/0502163.
- [16] T. Padmanabhan, Dark energy: the cosmological challenge of the millennium, Curr. Sci. 88, 1057–1068 (2005); astro-ph/0411044.
- [17] R. Bousso, TASI lectures on the cosmological constant, 0708.4231 [hep-th].
- [18] R. L. Jaffe, The Casimir effect and the quantum vacuum, Phys. Rev. D **72**, 021301 (2005); hep-th/0503158.